

# QuSpin: a Python Package for Dynamics and Exact Diagonalisation of Quantum Many Body Systems. Part II: bosons, fermions and higher spins

Phillip Weinberg\* and Marin Bukov

Department of Physics, Boston University,  
590 Commonwealth Ave., Boston, MA 02215, USA

\* weinbe58@bu.edu

May 26, 2017

## Abstract

We present a major update to QuSpin, *SciPostPhys.2.1.003*, – an open-source Python package for exact diagonalization and quantum dynamics of arbitrary boson, fermion and spin many-body systems, supporting the use of various symmetries in one dimension and (imaginary) time evolution following a user-specified driving protocol. We explain how to use the new features of QuSpin using six detailed examples of various complexity: (i) the transverse-field Ising model and the Jordan-Wigner transformation, (ii) free particle systems: the SSH model, (iii) the many-body localised Fermi-Hubbard model, (iv) the Bose-Hubbard model in a ladder geometry, (v) nonlinear (imaginary) time evolution and the Gross-Pitaevskii equation, and (vi) integrability breaking and thermalising dynamics in the spin-1 transverse-field Ising model. This easily accessible package can serve various purposes, including educational and cutting-edge experimental and theoretical research.

## Contents

<b>1 Which Problems can I Study with QuSpin?</b>	<b>2</b>
<b>2 How do I use the New Features of QuSpin?</b>	<b>3</b>
2.1 The Spectrum of the Transverse Field Ising Model and the Jordan-Wigner Transformation	3
2.2 Free Particle Systems: the Fermionic SSH Chain	8
2.3 Many-Body Localization in the Fermi-Hubbard Model	13
2.4 The Bose-Hubbard Model on Translationally Invariant Ladder	18
2.5 The Gross-Pitaevskii Equation and Nonlinear Time Evolution	22
2.6 Integrability Breaking and Thermalising Dynamics in the Spin-1 Transverse-Field Ising Model	27
<b>3 New Horizons for QuSpin</b>	<b>30</b>
<b>A Installation Guide in a Few Steps</b>	<b>30</b>

A.1 Mac OS X/Linux	30
A.2 Windows	31
<b>B Basic Use of Command Line to Run Python</b>	<b>32</b>
B.1 Mac OS X/Linux	32
B.2 Windows	33
B.3 Execute Python Script (any operating system)	33
<b>C Package Documentation</b>	<b>34</b>
<b>D Complete Example Codes</b>	<b>35</b>
<b>References</b>	<b>44</b>

---

## 1 Which Problems can I Study with QuSpin?

- re-label numbering of examples to match order in text.

Understanding the physics of many-body quantum condensed matter systems often involves a great deal of numerical simulations, be it to gain intuition about the complicated problem of interest, or because the latter does not admit an analytical solution which can be expressed in a closed form. This motivates the development of open-source packages [CITE], the purpose of which is to facilitate the study of condensed matter systems, without the need to understand and implement complicated numerical methods which required years to understand and fully develop. Here, we report on a major upgrade to QuSpin [1] – a Python library for exact diagonalisation (ED) and simulation of the dynamics in arbitrary quantum many-body systems.

Although ED methods are vastly outperformed by more sophisticated numerical techniques in the study of equilibrium problems [CITE], as of present date ED remains essential for most dynamical non-equilibrium problems. The reason for this often times relies on the fact that the underlying physics of these problems cannot be explained without taking into consideration the contribution from high-energy states excited during the evolution. Some prominent examples of such problems include the study of many-body localisation (MBL) [CITE], the Eigenstate Thermalisation hypothesis [CITE], quantum quench dynamics [CITE], periodically-driven systems [CITE], adiabatic and counter-diabatic state preparation, applications of Machine Learning to non-equilibrium physics [CITE], optimal control [CITE], and many more **did I forget smth important?**.

It is, thus, arguably useful to have a toolbox available at our disposal which allows one to quickly simulate and study these and related nonequilibrium problems. As such, QuSpin offers easy access to performing numerical simulations, which can facilitate the development and inspiration of new ideas and the discovery of novel phenomena, eliminating the cost of spending time to develop a reliable code. Besides theorists, the new version of QuSpin will hopefully even prove valuable to experimentalists working on problems containing dynamical setups, as it

can help students and researchers focus on perfecting the experiment, rather than worry about writing the supporting simulation code. Last but not least, with the computational processing power growing higher than ever before, the role played by simulations for theoretical research becomes increasingly more important too. It can, therefore, be expected that in the near future quantum simulations become an integral part of the standard physics university curriculum, and having easily accessible toolboxes, such as QuSpin, is one of the required prerequisites for this anticipated change.

## 2 How do I use the New Features of QuSpin?

New in QuSpin 2.0, we have added the following features and toolboxes:

- ...

*Installing QuSpin is quick and efficient; just follow the steps outlined in App. A.*

Before we carry on, we refer the interested reader to examples (i)-(iv) from the original QuSpin paper [1]. The examples below focus predominantly on the newly introduced features, and are thus to be considered complementary. We emphasize that, since they serve the purpose of explaining how to use QuSpin, for the sake of brevity we shall not discuss the interesting physics related to the interpretation of the results.

### 2.1 The Spectrum of the Transverse Field Ising Model and the Jordan-Wigner Transformation

This example shows how to

- construct fermionic hopping,  $p$ -wave pairing and on-site potential terms, and spin-1/2 interactions and transverse fields,
- implement periodic and anti-periodic boundary conditions,
- use particle conservation modulo 2, spin inversion, reflection, and translation symmetries,
- handle the default built-in particle conservation and symmetry checks,
- obtain the spectrum of a QuSpin Hamiltonian.

*Physics Setup*—The transverse field Ising (TFI) chain is paradigmatic in our understanding of quantum phase transitions, since it represents an exactly solvable model[CITE Sachdev]. The Hamiltonian is given by

$$H = \sum_{j=0}^{L-1} -J\sigma_{j+1}^z\sigma_j^z - h\sigma_j^x, \quad (1)$$

where the nearest-neighbour (nn) spin interaction is  $J$ ,  $h$  denotes the transverse field, and  $\sigma_j^\alpha$  are the Pauli spin-1/2 matrices. We use periodic boundary conditions and label the  $L$  lattice sites  $0, \dots, L-1$  to conform with Python's convention. This model has gapped, fermionic elementary excitations, and exhibits a phase transition from an antiferromagnet to a paramagnet at  $(h/J)_c = 1$  [CHECK!](#). The Hamiltonian possesses the following symmetries: magnetisation

conservation, parity (reflection w.r.t. the centre of the chain), spin inversion, and (many-body) momentum conservation.

In one dimension, the TFI Hamiltonian can be mapped to spinless  $p$ -wave superconducting fermions via the Jordan-Wigner (JW) transformation[CITE Sachdev, other paper]:

$$c_i = \prod_{j<i} \sigma_j^z \sigma_i^-, \quad c_i^\dagger = \prod_{j<i} \sigma_j^z \sigma_i^+, \quad (2)$$

where the fermionic operators satisfy  $\{c_i, c_j^\dagger\} = \delta_{ij}$ . The Hamiltonian is readily shown to take the form

$$H = \sum_{j=0}^{L-1} J \left( -c_j^\dagger c_{j+1} + c_j c_{j+1}^\dagger \right) + J \left( -c_j^\dagger c_{j+1}^\dagger + c_j c_{j+1} \right) + 2h \left( n_j - \frac{1}{2} \right). \quad (3)$$

In the fermionic representation, the spin  $zz$ -interaction maps to nn hopping and a  $p$ -wave pairing term with coupling constant  $J$ , while the transverse field translates to an on-site potential shift of magnitude  $h$ . In view of the implementation of the model using QuSpin, we have ordered the terms such that the site index is growing to the right, which comes at the cost of a few negative signs due to the fermion statistics. We emphasize that this ordering is not dictated by QuSpin, but it is merely our choice to adopt it here for the sake of consistency. The fermion Hamiltonian posses the symmetries: particle conservation modulo 2, parity and (many-body) “momentum” conservation.

Here, we are interested in studying the spectrum of the TFI model in both the spin and fermion representations. However, if one naïvely carries out the JW transformation, and computes the spectra of Eqs. (1) and (3), one might be surprised that they do not match exactly. The reason lies in the form of the boundary condition required to make the JW mapping exact – a subtle issue often left aside in favour of discussing the interesting physics of the TFI model itself.

We recall that the starting is the periodic boundary condition imposed on the spin Hamiltonian in Eq. (1). Due to the symmetries of the spin Hamiltonian (1), we can define the JW transformation on every symmetry sector separately. To make the JW mapping exact, we supplement Eq. (2) with the following boundary conditions: (i) the negative spin-inversion symmetry sector maps to the fermion Hamiltonian (3) with *periodic* boundary conditions (PBC) and *odd* total number of fermions; (ii) the positive spin-inversion symmetry sector maps to the fermion Hamiltonian (3) with *anti-periodic* boundary conditions (APBC) and *even* total number of fermions. Anti-periodic boundary conditions differ from PBC by a negative sign attached to all coupling constants that cross a single, fixed lattice bond (the bond itself is arbitrary as all bonds are equal for PBC). APBC and PBC are special cases of the more general twisted boundary conditions where, instead of a negative sign, one attaches an arbitrary phase factor.

In the following, we show how to compute the spectra of the Hamiltonians in Eqs. (1) and (3) with the correct boundary conditions using QuSpin. Figure 1 shows that they match exactly in both the PBC and APBC cases discussed above.

*Code Analysis*—We begin by loading the QuSpin operator and basis constructors, as well as some standard Python libraries.

```
1 from quspin.operators import hamiltonian # Hamiltonians and operators
2 from quspin.basis import spin_basis_1d, fermion_basis_1d # Hilbert space spin basis
```




Figure 1: Comparison of the spectra of the spin and fermion representation of the transverse field Ising Hamiltonian in the spin (1) and fermion (3) representations. The degeneracy in the spectrum is due to the remaining parity and momentum conservations which could but are not taken into account (see text). The parameters are  $J = 1.0$ ,  $h = \sqrt{2}$ , and  $L = 8$ .

```
3 import numpy as np # generic math functions
4 import matplotlib.pyplot as plt # figure/plot library
```

First, we define the models parameters.

```
6 ##### define model parameters #####
7 L=8 # system size
8 J=1.0 # spin zz interaction
9 h=np.sqrt(2) # z magnetic field strength
```

We have to consider two cases when computing the spectrum, as discussed in the theory section above. In one case, the fermionic system has PBC and the particle number sector is odd, while the spins are constrained to the negative spin inversion symmetry sector, while in the second – the fermion model has APBC with even particle number sector, and the spin model is considered in the positive spin inversion sector. To this end, we introduce the variables  $\mathbf{zblock} \in \{\pm 1\}$  and  $\mathbf{PBC} \in \{\pm 1\}$ , where  $\mathbf{PBC} = -1$  denotes APBC. Note that the only meaningful combinations are  $(\mathbf{zblock}, \mathbf{PBC}) = (-1, 1), (1, -1)$ , which we loop over:

```
11 # loop over spin inversion symmetry block variable and boundary conditions
12 for zblock,PBC in zip([-1,1],[1,-1]):
```

Within this loop, the code is divided in two independent parts: first, we compute the spectrum of the TFI system, and then – that of the equivalent fermionic model. Let us discuss the spins.

```
14 ##### define spin model
```

In QuSpin, operators are stored as sparse lists. These lists contain two parts: (i) the lattice sites on which the operator acts together with the coupling strength, which we call a *site-coupling* list, and (ii) the types of the operators involved, i.e. the *operator-string*. For example, the operator  $\mathcal{O} = g \sum_{j=0}^{L-1} \sigma_j^\mu$  can be uniquely represented by the site-coupling list  $[[\mathbf{g}, \emptyset], [\mathbf{g}, 1], \dots, [\mathbf{g}, L-1]]$ , and the information that it is the Pauli matrix  $\mu$ . The components lists are nothing but the tuples of the field strength and the site index  $[\mathbf{g}, j]$ . It is straightforward to generalise this to non-uniform fields  $\mathbf{g} \rightarrow \mathbf{g}[j]$ . Similarly, any two-body operator  $\mathcal{O} = J_{zz} \sum_{j=0}^{L-1} \sigma_j^\mu \sigma_{j+1}^\nu$  can be fully represented by the two sites it acts on, and its coupling strength:  $[\mathbf{J}, j, j+1]$ . We then stack up these elementary lists together into the site-coupling list:  $[[\mathbf{J}, \emptyset, 1], [\mathbf{J}, 1, 2], \dots, [\mathbf{J}, L-2, L-1], [\mathbf{J}, L-1, \emptyset]]$ .

```

15 # site-coupling lists (PBC for both spin inversion sectors)
16 h_field=[[-h,i] for i in range(L)]
17 J_zz=[[-J,i,(i+1)%L] for i in range(L)] # PBC

```

Notice the way we defined the periodic boundary condition for the spin-spin interaction using the modulo operator `%`, which effectively puts a coupling between sites  $L-1$  and  $0$ . We mention in passing that the above procedure generalises so one can define any multi-body local or nonlocal operator using QuSpin.

In order to specify the types of the on-site single-particle operators, we use operator strings. For instance, the transverse field operator  $\mathcal{O} = g \sum_{j=0}^{L-1} \sigma_j^x$  becomes `['x',h_field]`, while the two-body interaction is `['zz',J_zz]`. It is important to notice that the order of the letters in the operator string corresponds to the order the operators are listed in the site-coupling lists. Putting everything into one final list yields the `static` list for the spin model:

```

18 # define spin static and dynamic lists
19 static_spin=[["zz",J_zz],["x",h_field]] # static part of H

```

In QuSpin, the user can define both static and dynamic operators. Since this example does not contain any time evolution, we postpone the explanation of how to use dynamic lists to Sec. 2.5, and use an empty list instead.

```

20 dynamic_spin=[] # time-dependent part of H

```

The last step before we can construct the Hamiltonian is to build the basis for it. This is done using the basis constructors. For spin systems, we use `spin_basis_1d` which allows to use the operator strings `'z'`, `'+'`, `'-'`, `'I'`, and for spin-1/2 additionally `'x'`, `'y'`. The first and required argument is the number of sites  $L$ . Optional arguments are used to parse symmetry sectors. For instance, if we want to construct an operator in the spin-inversion block with quantum number  $+1$ , we can conveniently do this using the flag `zblock=1`.

```

21 # construct spin basis in pos/neg spin inversion sector depending on APBC/PBC
22 basis_spin = spin_basis_1d(L=L,zblock=zblock)

```

Having specified the static and dynamic lists, as well as the basis, building up the Hamiltonian is a one-liner, using the `hamiltonian` constructor. The required arguments in order of appearance are the `static` and `dynamic` lists, respectively. Optional arguments include the `basis`, and the precision or data type `dtype`. If no basis is passed, the constructor uses `spin_basis_1d` by default. The default data type is `np.complex128`.

```

23 # build spin Hamiltonians
24 H_spin=hamiltonian(static_spin,dynamic_spin,basis=basis_spin,dtype=np.float64)

```

The Hamiltonian is stored as a Scipy sparse matrix for efficiency. It can be cast to a full array for a more convenient inspection using the attribute `H.toarray()`. To calculate its spectrum, we use the attribute `H.eigvalsh()`, which returns all eigenvalues. Other attributes for diagonalisation were discussed in [Example 0](#), c.f. Ref. [1].

```
25 # calculate spin energy levels
26 E_spin=H_spin.eigvalsh()
```

Let us now move to the second part of the loop which defines the fermionic  $p$ -wave superconductor. We start by defining the site-coupling list for the local potential

```
28 ##### define fermion model
29 # define site-coupling lists for external field
30 h_pot=[[2.0*h,i] for i in range(L)]
```

Let us focus on the case of periodic boundary conditions `PBC=1` first.

```
31 if PBC==1: # periodic BC: odd particle number subspace only
```

In the fermion model, we have two types of two-body terms: hopping terms  $c_i^\dagger c_{i+1} - c_i c_{i+1}^\dagger$ , and pairing terms  $c_i^\dagger c_{i+1}^\dagger - c_i c_{i+1}$ . While QuSpin allows any ordering of the operators, for the sake of consistency we set a convention: the site indices grow to the right. To take into account the opposite signs resulting from the fermion statistics, we have to code the site-coupling lists for all four terms separately. This is analogous to the spin-spin interaction above:

```
32 # define site-coupling lists (including boundary couplings)
33 J_pm=[[-J,i,(i+1)%L] for i in range(L)] # PBC
34 J_mp=[[+J,i,(i+1)%L] for i in range(L)] # PBC
35 J_pp=[[-J,i,(i+1)%L] for i in range(L)] # PBC
36 J_mm=[[+J,i,(i+1)%L] for i in range(L)] # PBC
```

To construct a fermionic operator, we make use of the basis constructor `fermion_basis_1d`. Once again, we pass the number of sites `L`. As we explained in the analysis above, we need to consider all odd particle number sectors in the case of PBC. This is done by specifying the particle number sector `Nf`.

```
37 # construct fermion basis in the odd particle number subsector
38 basis_fermion = fermion_basis_1d(L=L,Nf=range(1,L+1,2))
```

In the case of APBC, we first construct all two-body site-coupling lists as if the boundaries were open, and supplement the APBC links with negative coupling strength in the end:

```
39 elif PBC==-1: # anti-periodic BC: even particle number subspace only
40 # define bulk site coupling lists
41 J_pm=[[-J,i,i+1] for i in range(L-1)]
42 J_mp=[[+J,i,i+1] for i in range(L-1)]
43 J_pp=[[-J,i,i+1] for i in range(L-1)]
44 J_mm=[[+J,i,i+1] for i in range(L-1)]
45 # add boundary coupling between sites (L-1,0)
46 J_pm.append([+J,L-1,0]) # APBC
47 J_mp.append([-J,L-1,0]) # APBC
48 J_pp.append([+J,L-1,0]) # APBC
49 J_mm.append([-J,L-1,0]) # APBC
```

The construction of the basis is the same, except that this time we need all even particle number sectors:



```

50 # construct fermion basis in the even particle number subsector
51 basis_fermion = fermion_basis_1d(L=L,Nf=range(0,L+1,2))

```

As before, we need to specify the type of operators that go in the fermion Hamiltonian using operator string lists. The `fermion_basis_1d` class accepts the following strings `'+', '-', 'n', 'I'`, and additionally the particle-hole symmetrised density operator `'z' = n - 1/2`. The static and dynamic lists read as

```

52 # define fermionic static and dynamic lists
53 static_fermion = [['+-', J_pm], ['-+', J_mp], ['++', J_pp], ['--', J_mm], ['z', h_pot]]
54 dynamic_fermion=[]

```

Constructing and diagonalising the fermion Hamiltonian is the same as for the spin-1/2 system. Note that one can disable the automatic built-in checks for particle conservation `check_pcon=False` and all other symmetries `check_symm=False` if one wishes to suppress the checks.

```

55 # build fermionic Hamiltonian
56 H_fermion=hamiltonian(static_fermion,dynamic_fermion,basis=basis_fermion,
57                      dtype=np.float64,check_pcon=False,check_symm=False)
58 # calculate fermionic energy levels
59 E_fermion=H_fermion.eigvalsh()

```

The complete code including the lines that produce Fig. 1 is available in Example Code 1.

## 2.2 Free Particle Systems: the Fermionic SSH Chain

This example shows how to

- construct free-particle Hamiltonians in real space,
- implement translation invariance with a two-site unit cell and construct the single-particle Hamiltonian in momentum space in block-diagonal form,
- compute non-equal time correlation functions,
- time evolve quantum states simultaneously

*Physics Setup*—The Su-Schrieffer-Heeger (SSH) model of a free-particle on a dimerised chain is widely used to introduce the concept of edge states, topology, Berry phase, etc., in one spatial dimension. The Hamiltonian is given by

$$H = \sum_{j=0}^{L-1} -(J + (-1)^j \delta J) (c_j c_{j+1}^\dagger - c_j^\dagger c_{j+1}) + \Delta (-1)^j n_j, \quad (4)$$

where  $\{c_i, c_j^\dagger\} = \delta_{ij}$  obey fermionic commutation relations. The uniform part of the hopping matrix element is  $J$ ,  $\delta J$  defines the bond dimerisation, and  $\Delta$  is the staggered potential. We assume periodic boundary conditions.

Below, we show how one can use QuSpin to study the physics of free fermions in the SSH chain. One way of doing this would be to work in the many-body (Fock space) basis, see Sec. ????. However, whenever the particles are non-interacting, the exponential scaling of the Hilbert space dimension with the number of lattice sites imposes an artificial limitation on the system sizes one can do. Luckily, with no interactions present, the many-body wave functions



factorise in a product of single-particle states. Hence, it is possible to study the behaviour of many free bosons and fermions by simulating the physics of a single particle.

Making use of translation invariance, a straightforward Fourier transformation to momentum space,  $a_k = \sqrt{2/L} \sum_{j \text{ even}}^{L-1} e^{-ikj} c_j$  and  $b_k = \sqrt{2/L} \sum_{j \text{ odd}}^{L-1} e^{-ikj} c_j$ , casts the SSH Hamiltonian in the following form

$$H = \sum_{k \in \text{BZ}'} (a_k^\dagger, b_k^\dagger) \begin{pmatrix} \Delta & -(J + \delta J)e^{-ik} - (J - \delta J)e^{+ik} \\ -(J + \delta J)e^{+ik} - (J - \delta J)e^{-ik} & -\Delta \end{pmatrix} \begin{pmatrix} a_k \\ b_k \end{pmatrix}, \quad (5)$$

where the reduced Brillouin zone is defined as  $\text{BZ}' = [-\pi/2, \pi/2)$ . We thus see that the Hamiltonian reduces further to a set of independent  $2 \times 2$  matrices. The spectrum of the SSH model is gapped, see Fig. ???.

Since we are dealing with free fermions, the ground state is the Fermi sea,  $|\text{FS}\rangle$ , defined by filling up the lowest band completely. We are interested in measuring the real-space non-equal time correlation function

$$C_{ij}(t) = \langle \text{FS} | n_i(t) n_j(0) | \text{FS} \rangle = \langle \text{FS}(t) | n_i(0) \underbrace{U(t, 0) n_j(0) | \text{FS} \rangle}_{|\text{FS}(t)\rangle}. \quad (6)$$

To evaluate the correlator numerically, we shall use the right-hand side of this equation.

As we are studying free particles, it is enough to work with the single-particle states. For instance the Fermi sea is given by  $|\text{FS}\rangle = \prod_{k \leq k_F} c_k^\dagger |0\rangle$ . Denoting the density operator in momentum space by  $\hat{n}_i$ , one can cast the correlator in momentum space in the following form:

$$C_{ij}(t) = \sum_{k \leq k_F} \langle k | \hat{n}_i(t) \hat{n}_j(0) | k \rangle. \quad (7)$$

If we want to consider finite-temperature  $\beta^{-1}$ , the above formula generalises to

$$C_{ij}(t, \beta) = \sum_{k \leq k_F} n_{\text{FD}}(k, \beta) \langle k | \hat{n}_i(t) \hat{n}_j(0) | k \rangle, \quad (8)$$

where  $n_{\text{FD}}(k, \beta) = 1/(\exp(\beta E) + 1)$  is the Fermi-Dirac distribution at temperature  $\beta^{-1}$ . Figure ??? shows the time evolution of  $C_{ij}(t, \beta)$  for two sites, separated by the maximal distance on the ring:  $L/2$ .

*Code Analysis.*—Let us explain how one can do all this quickly and efficiently using QuSpin. As always, we start by loading the required packages and libraries.

```
1 from quspin.operators import hamiltonian, exp_op # Hamiltonians and operators
2 from quspin.basis import fermion_basis_1d # Hilbert space fermion basis
3 from quspin.tools.block_tools import block_diag_hamiltonian # block diagonalisation
4 import numpy as np # generic math functions
5 import matplotlib.pyplot as plt # plotting library
6 try: # import python 3 zip function in python 2 and pass if using python 3
7     import itertools.izip as zip
8 except ImportError:
9     pass
```

After that, we define the model parameters

```

1 ##### define model parameters #####
2 L=100 # system size
3 J=1.0 # uniform hopping contribution
4 deltaJ=0.1 # bond dimerisation
5 Delta=0.5 # staggered potential
6 beta=100.0 # set inverse temperature for Fermi-Dirac distribution

```

In the following, we construct the fermionic SSH Hamiltonian first in real space. We then show how one can also construct it in momentum space where, provided we use periodic boundary conditions, it appear block-diagonal. Let us define the fermionic site-coupling lists. Once again, we mention that fermion systems require special care in defining the hopping terms: Eq. (4) is conveniently cast in the form where all site indices on the operators grow to the right, and all signs due to the fermion statistics are explicitly spelt out.

```

1 ##### construct single-particle Hamiltonian #####
2 # define site-coupling lists
3 hop_pm=[[-J-deltaJ*(-1)**i,i,(i+1)%L] for i in range(L)] # PBC
4 hop_mp=[[J+deltaJ*(-1)**i,i,(i+1)%L] for i in range(L)] # PBC
5 stagg_pot=[[Delta*(-1)**i,i] for i in range(L)]

```

Defining the static list assigns the specific SSH operator structure. Since our problem does not possess any explicit time dependence, we leave the **dynamic** list empty.

```

1 # define static and dynamic lists
2 static=[["+-",hop_pm],["-+",hop_mp],['n',stagg_pot]]
3 dynamic=[]

```

Defining the fermion basis with the help of the constructor **fermion\_basis\_1d** proceeds as smoothly as in Sec. 2.1. Notice a cheap trick: by specifying a total of **Nf=1** fermion in the lattice, QuSpin actually allows to define single-particle models, as a special case of the more general many-body Hamiltonians. Unlike many body models, however, due to the exponentially reduced Hilbert space size, this allows us to scale up the system size **L**.

```

1 # define basis
2 basis=fermion_basis_1d(L,Nf=1)

```

We then build the real-valued SSH Hamiltonian in real space by passing the static and dynamic lists, as well as the basis and the data type, and diagonalise it.

```

1 # build real-space Hamiltonian
2 H=hamiltonian(static,dynamic,basis=basis,dtype=np.float64)
3 # diagonalise real-space Hamiltonian
4 E,V=H.eigh()

```

For translation invariant single-particle models, however, the user might prefer to use momentum space, where the Hamiltonian becomes block diagonal, as we showed above. This can be achieved using QuSpin's **block\_tools**. The idea behind it is simple: the main purpose is to create the full Hamiltonian in block-diagonal form, where the blocks correspond to pre-defined quantum numbers. In our case, we would like to use momentum or **kblock**'s. Note that the unit cell in the SSH model contains two sites, which we encode using the variable **a=2**. Thus, we can create a list of dictionaries **blocks**, each element of which defines a single symmetry block. If we combine all blocks, we exhaust the full Hilbert space. All other basis arguments, such as the system size, we store in the variable **basis\_args**. We mention in passing that this procedure is independent of the symmetry, and can be done using all symmetries supported by QuSpin, not only translation.

```

1 ##### compute Fourier transform and momentum-space Hamiltonian #####
2 # define basis blocks and arguments
3 blocks=[dict(Nf=1,kblock=i,a=2) for i in range(L//2)] # only L//2 distinct momenta
4 basis_args = (L,)

```

To create the block-diagonal Hamiltonian, we invoke the `block_diag_hamiltonian` method. It takes both required and optional arguments, and returns the transformation which block-diagonalises the Hamiltonian (in our case the Fourier transform) and the block-diagonal Hamiltonian object. Required arguments, in order of appearance, are the `blocks`, the `static` and `dynamic` lists, the basis constructor, `basis_args`, and the data type. Since we expect the Hamiltonian to contain the Fourier factors  $\exp(-ik)$ , we know to choose a complex data type. `block_diag_hamiltonian` also accepts some optional arguments, such as the flags for disabling the automatic built-in symmetry checks. More about this function can be found in the documentation, cf. App. C.

```

1 # construct block-diagonal Hamiltonian
2 FT,Hblock = block_diag_hamiltonian(blocks,static,dynamic,fermion_basis_1d,
3                                     basis_args,np.complex128,get_proj_kwargs=dict(pcon=True))

```

We can then use all functions and methods of the `hamiltonian` class, for instance the diagonalisation routines:

```

1 # diagonalise momentum-space Hamiltonian
2 Eblock,Vblock=Hblock.eigh()

```

We now proceed to calculate the correlation function from Eq. (6). To this end, we shall split the correlator according to the RHS of Eq. (6). Thus, the strategy is to evolve both the Fermi sea  $|\text{FS}(t)\rangle$  and the auxiliary state  $|\text{nFS}(t)\rangle$  in time, and compute the expectation value of the time-independent operator  $n_i(0)$  in between the two states as a function of time. Last, keep in mind that we do not need to construct the Fermi sea as a many-body state explicitly, so we rather work with single-particle states.

The first step is to collect all momentum eigenstates into the array `psi`. We then build the operators  $n_{j=0}$  and  $n_{j=L/2}$  in real space.

```

1 ##### prepare the density observables and initial states #####
2 # grab single-particle states and treat them as initial states
3 psi0=Vblock
4 # construct operator n_1 = $n_{j=0}$
5 n_1_static=[[ 'n', [[1.0,0]]]]
6 n_1=hamiltonian(n_1_static,[],basis=basis,dtype=np.float64,
7                 check_herm=False,check_pcon=False)
8 # construct operator n_2 = $n_{j=L/2}$
9 n_2_static=[[ 'n', [[1.0,L//2]]]]
10 n_2=hamiltonian(n_2_static,[],basis=basis,dtype=np.float64,
11                 check_herm=False,check_pcon=False)

```

We then transform these two operators to momentum space using the method `rotate_by()`. Setting the flag `generator=False` treats the Fourier transform `FT` as a unitary, rather than a generator of a unitary.

```

1 # transform n_j operators to momentum space
2 n_1=n_1.rotate_by(FT,generator=False)
3 n_2=n_2.rotate_by(FT,generator=False)

```

Let us proceed with the time-evolution. We first define the time vector `t` and the state `n_psi0`.

to report a bug pls visit <https://github.com/weinbe58/QuSpin/issues>

```

1 ##### evaluate nonequal time correlator <FS|n_2(t) n_1(0)|FS> #####
2 # define time vector
3 t=np.linspace(0.0,90.0,901)
4 # calculate state acted on by n_1
5 n_psi0=n_1.dot(psi0)

```

We can perform the time evolution in two ways: (i) we calculate the time-evolution operator  $U$  using the `exp_op` class, and apply it to the momentum states `psi0` and `n_psi0`. The `exp_op` class calculates the matrix exponential  $\exp(aH)$  of an operator  $H$  multiplied by a complex number  $a$ . One can also conveniently compute a series of matrix exponentials  $\exp(aHt)$  for every time  $t$  by specifying the starting point `start`, endpoint `stop` and the number of elements `num` which define the time array  $t$  via `t=np.linspace(start,stop,num)`. Last, by parsing the flag `iterate=True` we create a generator – an object evaluated only at the time it is called, i.e. not pre-computed, which can save both time and memory.

```

1 # construct time-evolution operator using exp_op class (sometimes faster)
2 U = exp_op(Hblock,a=-1j,start=t.min(),stop=t.max(),num=len(t),iterate=True)
3 # evolve states
4 psi_t=U.dot(psi0)
5 n_psi_t = U.dot(n_psi0)

```

Another way of doing the time evolution, (ii), is to use `evolve()` method of the Hamiltonian class. The idea here is that every Hamiltonian defines a generator of time translations. This method solves Schrödinger equation using SciPy's ODE integration routines, see App. C for more details. The required arguments, in order of appearance, are the initial state, the initial time, and the time vector. The `evolve()` method also supports the option to create the output as a generator using the flag `iterate=True`. Both ways (i) and (ii) time-evolve all momentum states `psi` at once.

```

1 # alternative method for time evolution using Hamiltonian class
2 #psi_t=Hblock.evolve(psi0,0.0,t,iterate=True)
3 #n_psi_t=Hblock.evolve(n_psi0,0.0,t,iterate=True)

```

To evaluate the correlator, we first preallocate memory by defining the empty array `correlators`, which contains the correlator in every single-particle momentum mode  $|k\rangle$ . Using generators then allows us to loop only once over the time-evolved states `psi_t` and `n_psi_t`. In doing so, we evaluate the expectation value  $\langle \text{FS}(t) | n_i(0) | n \text{FS}(t) \rangle$  using the `matrix_ele()` method of the `hamiltonian` class. The flag `diagonal=True` makes sure only the diagonal matrix elements are calculated<sup>1</sup> and returned as an one-dimensional array.

```

1 # preallocate variable
2 correlators=np.zeros(t.shape+psi0.shape[1:])
3 # loop over the time-evolved states
4 for i, (psi,n_psi) in enumerate( zip(psi_t,n_psi_t) ):
5     correlators[i,:]=n_2.matrix_ele(psi,n_psi,diagonal=True).real

```

Finally, we weigh all single-state correlators by the Fermi-Dirac distribution to obtain the finite-temperature non-equal time correlation function  $C_{0,L/2}(t, \beta)$ .

```

1 # evaluate correlator at finite temperature
2 n_FD=1.0/(np.exp(beta*E)+1.0)
3 correlator = (n_FD*correlators).sum(axis=-1)

```

<sup>1</sup>Recall that `psi_t` and `n_psi_t` contain many time-evolved states, and if one uses the default `diagonal=False`, all off-diagonal matrix elements will be computed as well, so the result will be an array

The complete code including the lines that produce Fig. ?? is available in Example Code 2.

### 2.3 Many-Body Localization in the Fermi-Hubbard Model

This example shows how to:

- construct Hamiltonians for spinful fermions using the `tensor_basis` class.
- how to use `ops_dict()` class to construct Hamiltonians with varying parameters.
- use new basis functionality to construct simple product states
- use `obs_vs_time` functionality to measure observables as a function of time

A class of exciting new problems in the field of non-equilibrium physics is that of Many-body localised (MBL) models. The MBL transition is a dynamical phase transition in the eigenstates of a many-body Hamiltonian. Driven primarily by quenched disorder, the transition is distinguished by the system having ergodic eigenstates in the weak disorder limit and non-ergodic eigenstates in the strong disorder limit. The MBL phase is reminiscent of integrable systems as one can construct quasi-local integrals of motion, but these integrals of motion are much more robust in the sense that they are not sensitive to small perturbations as is the case in many classes of integrable systems[CITE MBL misc].

Motivated by recent experiments in cold atomic gasses [CITE Bloch MBL I think there are a few others too] we explore MBL in the context of fermions using QuSpin. The model we will consider is the Fermi-Hubbard model with quenched random disorder which has the following Hamiltonian:

$$H = -J \sum_{i=0,\sigma}^{L-2} \left( c_{i\sigma}^\dagger c_{i+1,\sigma} - c_{i\sigma} c_{i+1,\sigma}^\dagger \right) + U \sum_{i=0}^{L-1} n_{i\uparrow} n_{i\downarrow} + \sum_{i=0,\sigma}^{L-1} V_i n_{i\sigma} \quad (9)$$

where  $c_{i\sigma}$  and  $c_{i\sigma}^\dagger$  are the fermionic creation and annihilation operators on site  $i$  for spin  $\sigma$ , respectively. We will work in the sector of 1/4 filling for both up and down spins. Preparing an initial configuration of fermions of alternating spin on every other site, we will then measure the sublattice imbalance [CITE exp]:

$$I = (N_A - N_B)/N_{\text{tot}} \quad (10)$$

where  $A$  and  $B$  refer to the different sublattices of the chain and  $N$  is the particle number. Evolving the initial state under the Hamiltonian (9) we will calculate the time dependence of the imbalance which will give us information about the ergodicity of the Hamiltonian (or the lack thereof). If the Hamiltonian is ergodic then this quantity will decay to 0 in the limit  $t \rightarrow \infty$  as one would expect due to thermalising dynamics, while if the Hamiltonian is MBL then some memory of its initial state will be retained, and therefore the imbalance will remain non-zero even at infinite times.

Because the Hilbert space dimension grows so quickly for this Hamiltonian with the lattice size, we will only consider the dynamics after a finite amount of time, and even with this it is a fairly long calculation to do, due to, (say which parts take longest time)

*Code Analysis*— Once again we start out by loading a set of libraries which we need to proceed with the simulation of MBL fermions.




Figure 2: Sublattice imbalance  $I$  as a function of time, averaged over 100 disorder realizations for different disorder strengths. This data was taken on a chain of length  $L = 8$ .

```

1 from quspin.operators import hamiltonian,exp_op,ops_dict # operators
2 from quspin.basis import tensor_basis,fermion_basis_1d # Hilbert spaces
3 from quspin.tools.measurements import obs_vs_time # calculating dynamics
4 import numpy as np # general math functions
5 from numpy.random import uniform,choice # tools for doing random sampling
6 from time import time # tool for calculating computation time

```

While we already encountered most of these libraries and functions, in this example we introduce the `ops_dict()` class which defines an operator, parametrized by multiple parameters, as opposed to the Hamiltonian which is only parametrized by time. Also, since this example requires us to do many different disorder realisations, we use NumPy's random number library to do random sampling. We load `uniform` to generate the uniformly distributed random potential as well as `choice` which we use to estimate the uncertainties of the disorder averages using a bootstrap re-sampling procedure which we explain below [below, or in App.?](#). In order to time how long each realization takes, we use the `time` function from python's `time` library.

After importing all the required libraries and functions we set up the parameters for the simulation including the number of realizations `n_real`, and the physical couplings  $J$ ,  $U$ , the number of up and down fermions, etc.

```

10 ##### setting parameters for simulation
11 # simulation parameters
12 n_real = 100 # number of realizations
13 n_boot = 100 # number of bootstrap samples to calculate error
14 # physical parameters
15 L = 8 # system size
16 N = L//2 # number of particles
17 N_up = N//2 + N % 2 # number of fermions with spin up
18 N_down = N//2 # number of fermions with spin down
19 w_list = [1.0,4.0,10.0] # disorder strength
20 J = 1.0 # hopping strength
21 U = 5.0 # interaction strength
22 # range in time to evolve system
23 start,stop,num=0.0,35.0,101
24 t = np.linspace(start,stop,num=num,endpoint=True)

```



Next we set up the basis, introducing here the `tensor_basis` constructor class. In its full-fledged generality, `tensor_basis` takes  $n$  basis objects which it then uses to construct the matrix elements in the tensor product space:

$$\mathcal{H} = \mathcal{H}_1 \otimes \mathcal{H}_2 \otimes \cdots \otimes \mathcal{H}_n. \quad (11)$$

Here we consider the case of two Hilbert spaces,  $\mathcal{H} = \mathcal{H}_1 \otimes \mathcal{H}_2$ , one for fermions with spin up and one for spin-down fermions<sup>2</sup>. One disadvantage of using the tensor basis is that it does not allow for the use of symmetries, beyond particle-conservation (magnetisation in the case of spin systems). [we'll have to change this later when we have the ladder classes](#).

```

26 ##### create the basis
27 # build the two bases to tensor together to spinful fermions
28 basis_up = fermion_basis_1d(L,Nf=N_up) # up basis
29 basis_down = fermion_basis_1d(L,Nf=N_down) # down basis
30 basis = tensor_basis(basis_up,basis_down) # spinful fermions

```

The next step in the procedure is to set up the site-coupling and operator lists:

```

32 ##### create model
33 # define site-coupling lists
34 hop_right = [[-J,i,i+1] for i in range(L-1)] # hopping to the right OBC
35 hop_left = [[J,i,i+1] for i in range(L-1)] # hopping to the left OBC
36 int_list = [[U,i,i] for i in range(L)] # onsite interaction
37 # site-coupling list to create the sublattice imbalance observable
38 sublat_list = [[(-1.0)**i/N,i] for i in range(0,L)]
39 # create static lists
40 operator_list_0 = [
41     ["+|" , hop_left], # up hop left
42     ["-|" , hop_right], # up hop right
43     ["|+-" , hop_left], # down hop left
44     ["|-+" , hop_right], # down hop right
45     ["n|n" , int_list], # onsite interaction
46 ]
47 imbalance_list = [["n|",sublat_list],["|n",sublat_list]]

```

Notice here that the `"|"` character is used to separate the operators which belong to the up (left side of tensor product) and down (right side of tensor product) Hilbert spaces in `basis`. If no string is present the operator is assumed to be the identity `'I'`. The site-coupling list, on the other hand, does not require separating the two sides of the tensor product, as it is assumed that the operator string lines up with the correct site index when the `'|'` character is removed.

In the last couple of lines defining model (see below), we create a python `dictionary` object in which we add static operator lists as `values`, indexed by a particular string known as the corresponding `key`. This dictionary, which we refer to as `operator_dict`, is then passed into the `ops_dict()` class which, for each `key` in `operator_dict`, constructs the operator listed in the site-coupling list for that `key`. When one wants to evaluate this operator for a particular set of parameters, one uses a second dictionary (see `params_dict` below [give line number](#)) with the same keys as `operator_dict`: the value corresponding to each key in the `params_dict` multiplies the operator stored at that same key in `operator_dict`. This allows one to parametrize many-body operators in more complicated and general ways. In the present

<sup>2</sup>to construct a `tensor_basis` object in general: `t_basis = tensor_basis(basis_1,basis_2,...,basis_n)`



example, we define a key for the Fermi-Hubbard Hamiltonian, and then, as we need to change the disorder strength in between realisations, we also define keys for the local density operator for both up and down spins on each site. By doing this, we can then construct any disordered Hamiltonian by specifying the disorder at each site, and changing its value from one realisation to another.

```

48 # create operator dictionary for ops_dict class
49 # add key for Hubbard hamiltonian
50 operator_dict=dict(H0=operator_list_0)
51 # add keys for local potential in each site
52 for i in range(L):
53     # add to dictionary keys h0,h1,h2,...,hL with local potential operator
54     operator_dict["n"+str(i)] = [{"n|",[[1.0,i]]},["|n",[[1.0,i]]]]

```

The `ops_dict()` class constructs operators in almost an identical manner as a `hamiltonian()` class with the exception that there is no dynamic list. Next, we construct our initial state with the fermions dispersed over the lattice on every other site. To get the index of the `basis` state which this initial state corresponds to, one can use the `index` function of the `tensor_basis` class. This function takes a string or integer representing the product state for each of the Hilbert spaces and then searches to find the full product state, returning the corresponding index. We can then create an empty array `psi_0` of dimension the total Hilbert space size, and insert unity at the index corresponding to the initial state. This allows us to easily define the many-body product state used in the rest of the simulation.

```

56 ##### setting up operators
57 # set up hamiltonian dictionary and observable (imbalance I)
58 no_checks = dict(check_pcon=False,check_symm=False,check_herm=False)
59 H_dict = ops_dict(operator_dict,basis=basis,**no_checks)
60 I = hamiltonian(imbalance_list,[],basis=basis,**no_checks)
61 # strings which represent the initial state
62 s_up = "".join("1000" for i in range(N_up))
63 s_down = "".join("0010" for i in range(N_down))
64 # basis.index accepts strings and returns the index
65 # which corresponds to that state in the basis list
66 i_0 = basis.index(s_up,s_down) # find index of product state
67 psi_0 = np.zeros(basis.Ns) # allocate space for state
68 psi_0[i_0] = 1.0 # set MB state to be the given product state
69 print("H-space size: {:d}, initial state: |{:s}>(x)|{:s}>".format(basis.Ns,s_up,
    s_down))

```

Now that the operators are all set up, we can proceed with the simulation of the dynamics. First, we define a function which, given a disorder realization of the local potential, calculates the time evolution of  $\mathcal{I}(t)$ . We shall guide the reader through this function step by step. The syntax for this begins as follows:

```

71 # define function to do dynamics for different disorder realizations.
72 def real(H_dict,I,psi_0,w,t,i):
73     # body of function goes below
74     ti = time() # start timing function for duration of reach realisation

```

The first step is to construct the Hamiltonian from the `disorder` list which is as simple as

```

75 # create a parameter list which specifies the onsite potential with disorder
76 params_dict=dict(H0=1)

```

```

77     for i in range(L):
78         params_dict["n"+str(i)] = uniform(-w,w)
79     # using the parameters dictionary construct a hamiltonian object with those
80     # parameters defined in the list
81     H = H_dict.tohamiltonian(params_dict)

```

using the `tohamiltonian()` method of `H_dict` class, which accepts as argument the dictionary `params_dict`, which shares the same keys as `operator_dict`, but whose values are determined by the `disorder` list which changes from one realisation to another.

Once the Hamiltonian has been constructed we want to time-evolve the initial state with it. To this end, we use the fact that, for time-independent Hamiltonians, the time-evolution operator coincides with the matrix exponential  $\exp(-itH)$ . In QuSpin, a convenient way to define matrix exponentials is offered by the `exp_op` class. Given an operator  $A$ , it calculates  $\exp(atA)$  for any complex-valued number  $a$ . The time grid for  $t$  is specified using the optional arguments `start`, `stop` and `num`. If the latter are omitted, default is  $t = 1$ . The `exp_op` objects are either a list of arrays, the elements of which correspond to the operator  $\exp(atA)$  at the times defined or, if `iterate=True` a generator for this list is returned. Here, we use `exp_op` to create a generator list containing the time-evolved states as follows

```

82     # use exp_op to get the evolution operator
83     U = exp_op(H,a=-1j,start=t.min(),stop=t.max(),num=len(t),iterate=True)
84     psi_t = U.dot(psi_0) # get generator psi_t for time evolved state

```

To calculate the expectation value of the imbalance operator in time, we use the `obs_vs_time` function. Since the usage of this function was extensively discussed in Example2 and Example3 of Ref. [1], here we only mention that it accepts a (collection of) state(s) [or a generator] `psi_t`, a time vector `t`, and a dictionary `dict(I=I)`, whose values are the observables to calculate the expectation value of. The function returns a dictionary, the keys of which correspond to the keys every observable was defined under.

```

85     # use obs_vs_time to evaluate the dynamics
86     t = U.grid # extract time grid stored in U, and defined in exp_op
87     obs_t = obs_vs_time(psi_t,t,dict(I=I))

```

The function ends by printing the time of executing and returning the value for  $I$  as a function of time for this realization

```

88     # print reporting the computation time for realization
89     print("realization {}/{ } completed in {:.2f} s".format(i+1,n_real,time()-ti))
90     # return observable values
91     return obs_t["I"]

```

Now we are all set to run the disorder realizations for the different disorder strengths which in principle can be spilt up over multiple simulations, e.g. using `joblib` [c.f. Example2 from Ref. [1]] but for completeness we do all of the calculations in one script.

```

93     ##### looping over differnt disorder strengths
94     for w in w_list:
95         I_data = np.vstack([real(H_dict,I,psi_0,w,t,i) for i in range(n_real)])

```

Last, we calculate the average and its error bars using bootstrap re-sampling (See Appendix??)

```

96     ##### averaging and error estimation
97     I_avg = I_data.mean(axis=0) # get mean value of I for all time points
98     # generate bootstrap samples

```

```

99     bootstrap_gen = (I_data[choice(n_real,size=n_real)].mean(axis=0) for i in range(
        n_boot))
100     # generate the fluctuations about the mean of I
101     sq_fluc_gen = ((bootstrap-I_avg)**2 for bootstrap in bootstrap_gen)
102     I_error = np.sqrt(sum(sq_fluc_gen)/n_boot)

```

The complete code including the lines that produce Fig. 2 is available in Example Code 3.

## 2.4 The Bose-Hubbard Model on Translationally Invariant Ladder

This example shows how to:

- construct interacting Hamiltonians for bosonic systems.
- construct a Hamiltonian on a ladder geometry.
- use the `block_ops` class to evolve over several symmetry sectors at once.
- measure the entanglement entropy of a state on the ladder.

*Physics Setup*— In this example we use QuSpin to simulate the dynamics of the Bose-Hubbard model (BHM) on a ladder geometry. The BHM is a minimal model for interacting lattice bosons which is most often experimentally realizable in cold atom experiments [CITE]. The Hamiltonian is given by

$$H_{\text{BHM}} = -J \sum_{\langle ij \rangle} (a_i^\dagger a_j + \text{h.c.}) + \frac{U}{2} \sum_i n_i (n_i - 1) \quad (12)$$

where  $a_i$  and  $a_i^\dagger$  are bosonic creation and annihilation operators on site  $i$  respectively and the sum  $\langle ij \rangle$  is a sum over nearest neighbors on the ladder. We consider a half filled ladder of length  $L$  with  $N = 2L$  sites and cylindrical boundary condition along the ladder leg direction. Since we are interested in the limit  $U/J \gg 1$ , we restrict the local Hilbert space to allow at most two particles per site, effectively using a total of three states per site.

The system is initialised in a random product state, and then evolved with the Hamiltonian (12). Since the BHM is non-integrable, we expect that the system eventually thermalizes so that the occupation is roughly uniform over the entire system. Besides the local density, we also measure the entanglement entropy between the two legs of the ladder.

As we consider a translational invariant ladder, the Hilbert space factorizes into subspaces corresponding to the different many-body momentum blocks. This is similar to what was discussed in Sec. 2.2 for the SSH model, but slightly different as we consider translations of the *many-body* Fock states as opposed to the single particle states [2]. In certain cases, it happens that, even though the Hamiltonian features symmetries, the initial state does not obey them, as is the case in the present example, due to the initial state being a random product state. Thus, in this section we project the wavefunction to the different symmetry sectors and evolve each of the projections separately under the Hamiltonian for that symmetry sector. After the evolution, each of these block wavefunctions is transformed back to the local Fock space basis, and summed up to recover the properly evolved the initial state. We can then measure quantities such as the on-site density and the entanglement entropy. Figure 3 shows...




Figure 3: Quantities measured in the BHM: (a) the leg-to-leg entanglement entropy density and (b) the local density on each site as a function of time. The red dots on the entanglement plot shows the time points at which the snapshots of the local density are taken. The parameters are  $J = 1$ ,  $U = 10$ , and  $L = 6$ .

*Code Analysis*—Let us now show how one can simulate this system using QuSpin. Following the code structure of the previous examples, we begin by loading the modules needed for the computation:

```
1 from __future__ import print_function, division #import python 3 functions
2 from quspin.operators import hamiltonian # Hamiltonians and operators
3 from quspin.basis import boson_basis_1d # bosonic Hilbert space
4 from quspin.tools.block_tools import block_ops # dynamics in symmetry blocks
5 import numpy as np # general math functions
```

Next, we set up the model parameters defining the length of the ladder  $L$ , the total number of sites in the chain  $N=2*L$ , as well as the filling factor for the bosons  $nb$ , and the maximum number of states per site (i.e. the local on-site Hilbert space dimension)  $sps$ . The hopping matrix elements  $J_{\perp}$ ,  $J_{\parallel,1}$ , and  $J_{\parallel,2}$  (see Fig. ??create figure showing ladder and couplings etc...) correspond to the python script variables  $J\_perp$ ,  $J\_par\_1$ , and  $J\_par\_2$ , respectively. The on-site interaction is denoted by  $U$ .

```
1         ["+",hop_list_hc], # hopping h.c.
2         ["nn",int_list_2], # U n_i^2
3         ["n",int_list_1] # -U n_i
4     ]
5 dynamic = [] # no dynamic operators
6 # create block_ops object
7 blocks=[dict(kblock=kblock) for kblock in range(L)] # blocks to project on to
```

```

8 basis_args = (N,) # boson_basis_1d mandatory arguments
9 basis_kwargs = dict(nb=nb,sps=sps,a=2) # boson_basis_1d optional args

```

Next we set up the times at which we would like to solve the Schrödinger equation. Due to the same restrictions on the exponential solver discussed Sec. 2.3 we will only consider time points which are linearly spaced defined by the variables `start`, `stop`, and `num`

```

1 get_proj_kwargs = dict(pcon=True) # set projection to full particle basis
2 U_block = block_ops(blocks,static,dynamic,boson_basis_1d,basis_args,np.complex128,
3                 basis_kwargs=basis_kwargs,get_proj_kwargs=get_proj_kwargs)

```

For bosonic systems we have '+', '-', and 'n' as possible operators to use. In order to set up the Hubbard local interaction we must define two coupling lists for the  $U \sum_i n_i^2$  and the  $-U \sum_i n_i$ :

```

1 # setting up basis for local fock basis
2 basis = boson_basis_1d(N,nb=nb,sps=sps)
3 # setting up observables

```

We also define the hopping lists

```

1 no_checks = dict(check_herm=False,check_symm=False,check_pcon=False)
2 n_list = [hamiltonian(["n",[[1.0,i]]],[],basis=basis,dtype=np.float64,**no_checks)
3         for i in range(N)]
4 # set up initial state
5 i0 = np.random.randint(basis.Ns) # pick random state from basis set
6 psi = np.zeros(basis.Ns,dtype=np.float64)

```

where we use the `list_1.extend(list_2)` method to concatenate two lists together<sup>3</sup>. The way the ladder is defined, the even sites correspond to the bottom side while the odds sites are the top part of the ladder, therefore the hopping on the bottom/top rung is defined by `[J_par,...,i,(i+2)%N]`, while hopping from top to bottom is defined by `[J_perp,i,(i+1)%N]`. Finally we can define the static and dynamic lists which we will not use to construct a `hamiltonian` object but instead we will use the `block_ops` class.

```

1 psi[i0] = 1.0
2 # print info about setup
3 state_str = "".join(str(int((basis[i0]//basis.sps*i)%basis.sps)) for i in range(N))
4 print("total H-space size: {}, initial state: |{}>".format(basis.Ns,state_str))
5 ##### do time evolution
6 # calculating the evolved states
7 n_jobs = 1 # paralelisation: increase to see if calculation runs faster!
8 psi_t = U_block.expm(psi,start=start,stop=stop,num=num,block_diag=False,n_jobs=
    n_jobs)

```

The purpose of `block_ops` is to provide a simple interface for solving the Schrödinger equation when an initial state may not obey the symmetries of the Hamiltonian which is generating the dynamics. We have seen an example of this in Sec. 2.2 when trying to measure non-equal space-time correlation functions of local operators in a translationally invariant system, while in this section we explicitly start out with a state which does not obey translational invariance. To construct the `block_ops` object we use the follow set of code

<sup>3</sup>Note that the `extend` function is done inplace so if you try to do `new_list=list_1.extend(list_2)`, `new_list` will be `None` and `list_1` will have all of the elements of `list_2` appended to it.

```

1 # calculating entanglement entropy
2 sub_sys_A = range(0,N,2) # bottom side of ladder
3 gen = (basis.ent_entropy(psi,sub_sys_A=sub_sys_A)["Sent_A"])/L for psi in psi_t.T[:]
4 ent_t = np.fromiter(gen,dtype=np.float64,count=num)
5 # calculating the local densities as a function of time
6 expt_n_t = np.vstack([n.expt_value(psi_t).real for n in n_list]).T
7 # reshape data for plotting

```

Firstly, **blocks** is a list of dictionaries which define the different symmetry sectors to evolve the initial state over<sup>4</sup>. Secondly **basis\_args** and **basis\_kwargs** define optional arguments which apply to every symmetry sector. Finally **get\_proj\_kwargs** contains the optional arguments to construct the projectors<sup>5</sup>. For more information about this class we refer the user to the Documentation. Finally we define the initial state and the local density operators using the **boson\_basis\_1d** class.

```

1 n_t = np.zeros((num,2,L))
2 n_t[:,0,:] = expt_n_t[:,0::2]
3 n_t[:,1,:] = expt_n_t[:,1::2]
4 # plotting static figures
5 """
6 fig, ax = plt.subplots(nrows=5,ncols=1)
7 im=[]
8 im_ind = []
9 for i,t in enumerate(np.logspace(-1,np.log10(stop-1),5,base=10)):
10     j = times.searchsorted(t)
11     im_ind.append(j)
12     im.append(ax[i].imshow(n_t[j],cmap="hot",vmax=n_t.max(),vmin=0))
13     ax[i].tick_params(labelbottom=False,labelleft=False)

```

Next we can calculate the time dependent states by using the **expm** function of the **block\_ops** class

```

1 cax = fig.add_axes([0.85, 0.1, 0.03, 0.8])
2 fig.colorbar(im[2],cax)
3 plt.savefig("boson_density.pdf")

```

We define this function to have almost identical arguments as that of the **exp\_op** class, but with some major exceptions. For one, because the hamiltonian factorizes the evolution over each block can be done separately (e.g. just trivially loop through them sequentially), but in some cases where there are a lot of small blocks like in a single particle Hamiltonian, then it actually makes sense to calculate the matrix exponential in block diagonal form which is why we added the optional argument **block\_diag** which toggles between these two options. Another option we have put in allows the user to spawn multiply python processes to do split up the work of doing the calculations for different blocks simultaneously, by setting **n\_jobs** to be greater than 1. On most systems these processes will be distributed over multiple CPUs which can speed up the calculations if the resources are there. This also works on conjunction with the **block\_diag** where each process creates its own block diagonal matrix for the calculation. Once all the calculations for each block are completed the results are then combined and projected to the local Fock basis. Finally, with the time dependent states calculated we can calculate the expectation values and entanglement entropy.

<sup>4</sup>**block\_ops** will not evolve a particular symmetry sector if the projection is 0.

<sup>5</sup>In this case setting **pcon=True** means that the projector takes the state from the symmetry reduced basis to the fixed particle number basis

```

1 plt.figure()
2 plt.plot(times,ent_t,lw=2)
3 plt.plot(times[im_ind],ent_t[im_ind],marker="o",linestyle="",color="red")
4 plt.xlabel("$t/J$", fontsize=20)
5 plt.ylabel("$s_{\mathrm{ent}}(t)$", fontsize=20)

```

In the newer versions of QuSpin we have moved the entanglement entropy calculations to the basis classes themselves (keeping of course backwards compatible functions from older versions), the rational being that this calculation is highly dependent on the type of system one is studying. Here we show how to use the basis function to calculate the entanglement entropy but this function can also calculate the reduced density matrix and its eigenvalues for pure and mixed states. Once again we refer the reader to the Documentation to learn more about how to use this function.

## 2.5 The Gross-Pitaevskii Equation and Nonlinear Time Evolution

This example shows how to

- simulate time-dependent nonlinear equations of motion
- use imaginary time dynamics to find a lowest energy configuration

*Physics Setup*—The Gross-Pitaevskii wave equation (GPE) has been shown to govern the physics of weakly-interacting bosonic systems. It constitutes the starting point for studying Bose-Einstein condensates, but can also appear in non-linear optics, and represents the natural description of Hamiltonian mechanics in the wave picture. One of its interesting features is that it can exhibit chaotic classical dynamics, a physical manifestation of the presence of a cubic non-linear term.

Here, we study the time-dependent GPE on a one-dimensional lattice:

$$\begin{aligned}
 i\partial_t \psi_j(t) &= -J(\psi_{j-1}(t) + \psi_{j+1}(t)) + \frac{1}{2}\kappa_{\text{trap}}(t)(j - j_0)^2 \psi_j(t) + U|\psi_j(t)|^2 \psi_j(t), \\
 \kappa_{\text{trap}}(t) &= (\kappa_f - \kappa_i)t/t_{\text{ramp}} + \kappa_i
 \end{aligned} \tag{13}$$

where  $J$  is the hopping matrix element,  $\kappa_{\text{trap}}(t)$  – the harmonic trap strength which varies slowly in time over a scale  $t_{\text{ramp}}$ , and  $U$  – the mean-field interaction strength. The lattice sites are labelled by  $j = 0, \dots, L - 1$ , and  $j_0$  is the centre of the 1d chain. We set the lattice constant to unity, and use open boundary conditions.

Whenever  $U = 0$ , the system is non-interacting and the GPE reduces to the Heisenberg EOM for the bosonic field operator  $\hat{\psi}_j(t)$ . Thus, for the purposes of using QuSpin to simulate the GPE, it is instructive to cast Eq. (13) in the following generic form

$$i\partial_t \vec{\psi}(t) = H_{\text{sp}}(t)\vec{\psi}(t) + U\vec{\psi}^*(t) \circ \vec{\psi}(t) \circ \vec{\psi}(t), \tag{14}$$

where  $[\vec{\psi}(t)]_j = \psi_j(t)$ , and  $\circ$  represents the element-wise multiplication

$$\vec{\psi}(t) \circ \vec{\psi}(t) = \left( \psi_0(t)\phi_0(t), \psi_1(t)\phi_1(t), \dots, \psi_{L-1}(t)\phi_{L-1}(t) \right)^t.$$

The time-dependent single-particle Hamiltonian in real space is represented as an  $L \times L$  matrix,  $H_{\text{sp}}(t)$ , which comprises the hopping term, and the harmonic trap.



We want to initiate the time-evolution of the system at  $t = 0$  in its lowest energy state. To this end, we can define a ‘ground state’ for the GPE equation, in terms of the configuration which minimises the energy of the system:

$$\begin{aligned}\vec{\psi}_{\text{GS}} &= \inf_{\vec{\psi}} \left( \vec{\psi}^\dagger H_{\text{sp}}(0) \vec{\psi} + \frac{U}{2} \sum_{j=0}^{L-1} |\psi_j|^4 \right), \\ &= \inf_{\psi_j} \left( -J \sum_{j=0}^{L-2} (\psi_{j+1}^* \psi_j + \text{c.c.}) + \frac{1}{2} \kappa_{\text{trap}}(0) \sum_{j=0}^{L-1} (j - j_0)^2 |\psi_j|^2 + \frac{U}{2} \sum_{j=0}^{L-1} |\psi_j|^4 \right). \quad (15)\end{aligned}$$

One way to find the configuration  $\vec{\psi}_{\text{GS}}$ , is to solve the GPE in imaginary time ( $it \rightarrow \tau$ ), which induces exponential decay in all modes of the system, which singles out the lowest-energy state in the longer run. In doing so, we keep the norm of the solution fixed:

$$\begin{aligned}\partial_\tau \vec{\varphi}(\tau) &= - \left[ H_{\text{sp}}(0) \vec{\varphi}(\tau) + U \vec{\varphi}^*(\tau) \circ \vec{\varphi}(\tau) \circ \vec{\varphi}(\tau) \right], \quad ||\vec{\varphi}(\tau)|| = \text{const.}, \\ \vec{\psi}_{\text{GS}} &= \lim_{\tau \rightarrow \infty} \vec{\varphi}(\tau)\end{aligned} \quad (16)$$

Once we have the initial state  $\vec{\psi}_{\text{GS}}$ , we evolve it according to the time-dependent GPE, Eq. (13), and track down the time evolution of the condensate density  $\rho_j(t) = |\psi_j(t)|^2$ . Fig. ??? shows the result.

*Code Analysis*—In the following, we demonstrate how one can code the above physics using QuSpin. As usual, we begin by loading the necessary packages:

```
1 from quspin.operators import hamiltonian # Hamiltonians and operators
2 from quspin.basis import boson_basis_1d # Hilbert space boson basis
3 from quspin.tools.measurements import evolve # nonlinear evolution
4 import numpy as np # generic math functions
5 import matplotlib.pyplot as plt # plot library
```

Next, we define the model parameters. We distinguish between static parameters and dynamic parameters – those involved in the time-dependent trap widening.

```
7 ##### define model parameters #####
8 L=300 # system size
9 # calculate centre of chain
10 if L%2==0:
11     j0 = L//2-0.5 # centre of chain
12 else:
13     j0 = L//2 # centre of chain
14 sites=np.arange(L)-j0
15 # static parameters
16 J=1.0 # hopping
17 U=1.0 # Bose-Hubbard interaction strength
18 # dynamic parameters
19 kappa_trap_i=0.001 # initial chemical potential
20 kappa_trap_f=0.0001 # final chemical potential
21 t_ramp=40.0/J # set total ramp time
```

In order to do time evolution, we code up the trap widening protocol from Eq. (13) in the function `ramp`. Since we want to make use of QuSpin’s time-dependent operator features, the

first argument must be the time  $t$ , followed by all protocol parameters. These same parameters are then explicitly stored in the variable `ramp_args`.

```

22 # ramp protocol
23 def ramp(t,kappa_trap_i,kappa_trap_f,t_ramp):
24     return (kappa_trap_f - kappa_trap_i)*t/t_ramp + kappa_trap_i
25 # ramp protocol parameters
26 ramp_args=[kappa_trap_i,kappa_trap_f,t_ramp]

```

With this, we are ready to construct the single-particle Hamiltonian  $H_{\text{sp}}(t)$ . The first step is to define the site-coupling lists, and the static and dynamic lists. Note that the dynamic list, which defines the harmonic potential in the single-particle Hamiltonian, contains four elements: apart from the operator string and the corresponding site-coupling list, the third and fourth elements are the time-dependent function `ramp` and its argument list `ramp_args`, and this order is crucial.

```

28 ##### construct single-particle Hamiltonian #####
29 # define site-coupling lists
30 hopping=[[-J,i,(i+1)%L] for i in range(L-1)]
31 trap=[[0.5*(i-j0)**2,i] for i in range(L)]
32 # define static and dynamic lists
33 static=[["+-",hopping],["-+",hopping]]
34 dynamic=[['n',trap,ramp,ramp_args]]

```

To create the single-particle Hamiltonian, we choose to use the bosonic basis constructor `boson_basis_1d` specifying the number sector to `Nb=1` boson for the entire lattice, and a local Hilbert space of `sps=2` states per site (empty and filled).

```

35 # define basis
36 basis = boson_basis_1d(L,Nb=1,sps=2)

```

Then we call the `hamiltonian` constructor to build the single-particle matrix. We can obtain the single-particle ground state without fully diagonalising this matrix, by using the sparse diagonalisation attribute `Hsp.eigsh()`. The `eigsh()` routine accepts the optional flags `k=1` and `'which'='SA'` which restrict the routine to find the first eigenstate starting from the bottom of the spectrum, i.e. the ground state.

```

37 # build Hamiltonian
38 Hsp=hamiltonian(static,dynamic,basis=basis,dtype=np.float64)
39 E,V=Hsp.eigsh(time=0.0,k=1,which='SA')

```

Having set up the `Hsmiltonian`, the next step in the simulation is to compute the ground state of the GPE using imaginary time evolution from Eq. (16). To this end, we first define the function `GPE_imag_time` which evaluate the RHS. It is required that the first argument for this function is (imaginary) time `tau`, followed by the state `phi`. All other arguments, such as the single-particle Hamiltonian and the interaction strength are listed last. Note that we evaluate the time-dependent Hamiltonian at `time=0`, since we are interested in finding the GPE GS for the initial trap configuration  $\kappa_i$ . Similar to before, we store these optional arguments in a list which we call `GPE_params`.

```

41 ##### imaginary-time evolution to compute GS of GPE #####
42 def GPE_imag_time(tau,phi,Hsp,U):
43     """
44     This function solves the real-valued GPE in imaginary time:
45     $$ -\dot{\phi}(\tau) = Hsp(t=0)\phi(\tau) + U |\phi(\tau)|^2 \phi(\tau) $$

```

to report a bug pls visit <https://github.com/weinbe58/QuSpin/issues>

```

46     """
47     return -( Hsp.dot(phi,time=0) + U*np.abs(phi)**2*phi )
48 # define ODE parameters
49 GPE_params = (Hsp,U)

```

Any initial value problem requires us to pick an initial state. In the case of imaginary evolution, this state can often be arbitrary, but needs to possess the same symmetries as the true GPE ground state. Here, we choose the ground state of the single-particle Hamiltonian for an initial state, and normalise it to one particle per site. We also define the imaginary time vector `tau`. This array has to contain sufficiently long times so that we make sure we end up in the long imaginary time limit  $\tau \rightarrow \infty$ , as required by Eq. (16). Since imaginary time evolution is not unitary, QuSpin will be normalising the vector every  $\tau$ -step. Thus, one also needs to make sure these steps are small enough to avoid convergence problems with the ODE solver.

```

50 # define initial state to flow to GS from
51 phi0=V[:,0]*np.sqrt(L) # initial state normalised to 1 particle per site
52 # define imaginary time vector
53 tau=np.linspace(0.0,35.0,71)

```

Performing imaginary time evolution is done using the `evolve()` method of the `measurements` tool. This function accepts an initial state `phi0`, initial time `tau[0]`, and a time vector `tau` and solves any user-defined ODE, here `GPE_imag_time`. The parameters of the ODE are passed using the keyword argument `f_params=GPE_params`. To ensure the normalisation of the state at each  $\tau$ -step we use the flag `imag_time=True`. Real-valued output can be specified by `real=True`. Last, we request `evolve()` to create a generator object using the keyword argument `iterate=True`. Many of the keyword arguments of `evolve()` are the same as in the `H.evolve()` method of the `hamiltonian` class: for instance, one can choose a specific SciPy solver and its arguments, or the solver's absolute and relative tolerance. We refer the interested reader to the documentation, cf. App. C.

```

54 # evolve state in imaginary time
55 psi_tau = evolve(phi0,tau[0],tau,GPE_imag_time,f_params=GPE_params,
56                 imag_time=True,real=True,iterate=True)

```

Looping over the generator `psi_tau` we have access to the solution, which we display in a form of a movie:

```

75 # display state evolution
76 for i,psi0 in enumerate(psi_tau):
77     # compute energy
78     E_GS=(Hsp.matrix_ele(psi0,psi0,time=0) + 0.5*U*np.sum(np.abs(psi0)**4) ).real
79     # plot wave function
80     plt.plot(sites, abs(phi0)**2, color='r',marker='s',alpha=0.2,
81             label='$|\\phi_j(0)|^2$')
82     plt.plot(sites, abs(psi0)**2, color='b',marker='o',
83             label='$|\\phi_j(\\tau)|^2$')
84     plt.xlabel('$\\mathrm{lattice\\ sites}$',fontsize=14)
85     plt.title('$J\\tau=\\tau=0.2f,\\ E\\mathrm{GS}(\\tau)=0.4fJ$'%(tau[i],E_GS)
86             ,fontsize=14)
87     plt.ylim([-0.01,max(abs(phi0)**2)+0.01])
88     plt.legend(fontsize=14)
89     plt.draw() # draw frame
90     plt.pause(0.005) # pause frame

```

```

91 plt.clf() # clear figure
92 plt.close()

```

Last, we use our GPE ground state, to time-evolve it in real time according to the trap widening protocol **ramp**, hard-coded into the single-particle Hamiltonian. We proceed analogously – first we define the real-time GPE and the time vector. In defining the GPE function, we split the ODE into a time-independent static part and a time-dependent dynamic part. The single-particle Hamiltonian for the former is accessed using the **hamiltonian** attribute **Hsp.static** which returns a SciPy sparse array. We can then manually add the non-linear cubic mean-field interaction term. In order to access the time-dependent part of the Hamiltonian, and evaluate it, we loop over the dynamic list **Hsp.dynamic**, which is a list consisting of the corresponding operator **Hd** together with the time-dependent function **f** which multiplies it, and its arguments **f\_args**. In the very end, we multiply the final output vector by the Schrödinger  $-i$ , which ensures the unitarity of real-time evolution.

```

77 ##### real-time evolution of GPE #####
78 def GPE(time,psi):
79     """
80     This function solves the complex-valued time-dependent GPE:
81     $$ i\dot{\psi}(t) = Hsp(t)\psi(t) + U |\psi(t)|^2 \psi(t) $$
82     """
83     # solve static part of GPE
84     psi_dot = Hsp.static.dot(psi) + U*np.abs(psi)**2*psi
85     # solve dynamic part of GPE
86     for Hd,f,f_args in Hsp.dynamic:
87         psi_dot += f(time,*f_args)*Hd.dot(psi)
88     return -1j*psi_dot
89 # define real time vector
90 t=np.linspace(0.0,t_ramp,101)

```

To perform the real-time evolution explicitly we once again use the **evolve()** function. This time, however, since the solution of the GPE is anticipated to be complex-valued, and because we do not do imaginary time, we do not need to pass the flags **real** and **imag\_time**. Instead, we decided to show the flags for the relative and absolute tolerance of the solver.

```

91 # time-evolve state according to GPE
92 psi_t = evolve(psi_0,t[0],t,GPE,iterate=True,atol=1E-12,rtol=1E-12)

```

Finally, we can enjoy the movie displaying real-time evolution

```

94 # display state evolution
95 for i,psi in enumerate(psi_t):
96     # compute energy
97     E=(Hsp.matrix_ele(psi,psi,time=t[i]) + 0.5*U*np.sum(np.abs(psi)**4)).real
98     # compute trap
99     kappa_trap=ramp(t[i],kappa_trap_i,kappa_trap_f,t_ramp)*(sites)**2
100    # plot wave function
101    plt.plot(sites, abs(psi)**2, color='r',marker='s',alpha=0.2
102             ,label='$|\psi_{\mathrm{GS},j}|^2$')
103    plt.plot(sites, abs(psi)**2, color='b',marker='o',label='$|\psi_j(t)|^2$')
104    plt.plot(sites, kappa_trap,'--',color='g',label='$\mathrm{trap}$')
105    plt.ylim([-0.01,max(abs(psi_0)**2)+0.01])
106    plt.xlabel('$\mathrm{lattice\ sites}$',fontsize=14)
107    plt.title('$Jt=0.2f, E(t)-E_{\mathrm{GS}}=0.4f$'%(t[i],E-E_GS),fontsize=14)

```

```

108 plt.legend(loc='upper right', fontsize=14)
109 plt.draw() # draw frame
110 plt.pause(0.00005) # pause frame
111 plt.clf() # clear figure
112 plt.close()

```

The complete code including the lines that produce Fig. ?? is available in Example Code 4.

## 2.6 Integrability Breaking and Thermalising Dynamics in the Spin-1 Transverse-Field Ising Model

This example shows how to:

- construct Hamiltonians for Higher spin operators.
- find ground state of a Hamiltonian
- use `obs_vs_time` function with costume user defined generator to calculate the expectation value of operators as a function of time.
- use the new functionality of the basis class to calculate the entanglement entropy for higher spin.

*Physics Setup*— In the previous section we introduced the TFI model and showed how one can solve the problem using the Jordan-Wigner transformation. This transformation allows one to get an exact analytic solution to the Hamiltonian (when the system obeys translational invariance). The fact that this solution exists is deeply connected to the notion of Integrability which has implications of how the system responds to a periodic modulation [CITE]. For non-integrable system when periodic driving, energy is no longer conserved and so generically one would expect that the system will heat up to infinite temperature, while in an integrable system, even though energy is not conserved, there are an extensive number of other static conserved quantities which may be conserved under the drive. If this is the case, then the system will not heat up at long times, but instead reach some steady state. By simply taking the transverse field ising model and promoting the spin-1/2 operators to spin-1, there is no longer a simple mapping to a quadratic Hamiltonian and therefore the model is no longer integrable. Here we will show this explicitly by driving the two different systems and checking if they heat or not. To do this we will define two Hamiltonians

$$H_{zz} = - \sum_{i=0}^{L-1} S_i^z S_{i+1}^z, \quad H_x = - \sum_{i=0}^{L-1} S_i^x \quad (17)$$

and evolve the ferromagnetic ground state of  $H_{zz}$  with the following piecewise periodic Hamiltonian:

$$H(t) = H_{zz} - \Omega \operatorname{sgn}(\cos(\Omega t)) H_x \quad (18)$$

where  $\Omega$  is the driving frequency and  $T$  is the period. As the Hamiltonian obeys translation, parity and spin-inversion symmetries we will use this to speed up the evolution by working in the symmetry sector which contains the ground state.

In order to measure the difference in heating between spin-1 and spin-1/2 we measure the expectation value of  $H_{zz}$  as a function of time. This operator has a symmetric spectrum and




Figure 4: Comparing the dynamics of  $Q(t)$  (a) and  $s_{\text{ent}}(t)$  (b) for  $S = 1$  (orange) and  $S = 1/2$  (blue) at stroboscopic times ( $t = nT$ ). For  $S = 1$  and  $S = 1/2$  we take  $L = 11$  and  $18$  respectively as to make sure the many-body Hilbert spaces have roughly the same number of state.  $s_{\text{ent}}$  is normalized by the Page entropy per site[CITE Page]. Note that for both systems  $\Omega = 4$ .

so following ref.[CITE heading papers] we define  $Q$ :

$$Q(t) = \left\langle \psi(t) \left| \frac{2(H_{zz} - E_{\min})}{E_{\max} - E_{\min}} \right| \psi(t) \right\rangle = \left\langle \psi(t) \left| \frac{H_{zz} - E_{\min}}{-E_{\min}} \right| \psi(t) \right\rangle \quad (19)$$

where the last equality comes from the symmetry of the spectrum:  $E_{\max} = -E_{\min}$ . This quantity is defined such that an infinite temperature state has  $Q = 1$ . Another measure of heating we will use is the entanglement entropy density

$$s_{\text{ent}}(t) = -\frac{1}{|A|} \text{tr}_A [\rho_A(t) \log \rho_A(t)], \quad \rho_A(t) = \text{tr}_{A^c} |\psi(t)\rangle \langle \psi(t)| \quad (20)$$

of subsystem A, defined to contain the left half of the chain and  $|A| = L/2$ . We denoted the reduced density matrix of subsystem A by  $\rho_A$ , and  $A^c$  is the complement of A.

*Code Analysis*—...

```
1 from __future__ import print_function, division
2
3 import numpy as np
4 from quspin.operators import ops_dict, hamiltonian, exp_op
5 from quspin.basis import spin_basis_1d
6 from quspin.tools.measurements import obs_vs_time
7 import matplotlib.pyplot as plt
8 import sys, os
9 # user defined generator
```

```

10 # generates stroboscopic dynamics
11 def evolve_gen(psi0,nT,*U_list):
12     yield psi0
13     for i in range(nT): # loop over number of periods
14         for U in U_list: # loop over unitaries
15             psi0 = U.dot(psi0)
16         yield psi0
17 # frequency and period for driving.
18 omega = 2
19 T = 2*np.pi/omega
20 nT = 200 # number of periods to evolve to.
21 times = np.arange(0,nT+1,1)*T
22 L_1 = 18 # length of chain for spin 1/2
23 L_2 = 11 # length of chain for spin 1
24 ##### setting up basis #####
25 basis_1 = spin_basis_1d(L_1,S="1/2",kblock=0,pblock=1,zblock=1) # spin 1/2 basis
26 basis_2 = spin_basis_1d(L_2,S="1",kblock=0,pblock=1,zblock=1) # spin 1 basis
27 # print information about the basis
28 print("S = {S:3s}, L = {L:2d}, Size of H-space: {Ns:d}".format(S="1/2",L=L_1,Ns=
    basis_1.Ns))
29 print("S = {S:3s}, L = {L:2d}, Size of H-space: {Ns:d}".format(S="1",L=L_2,Ns=
    basis_2.Ns))
30 # setting up coupling lists
31 Jzz_1 = [[-1.0,i,(i+1)%L_1] for i in range(L_1)]
32 hx_1 = [[-1.0,i] for i in range(L_1)]
33 Jzz_2 = [[-1.0,i,(i+1)%L_2] for i in range(L_2)]
34 hx_2 = [[-1.0,i] for i in range(L_2)]
35 # dictioanry to turn off checks
36 no_checks = dict(check_symm=False,check_herm=False)
37 # setting up hamiltonians
38 Hzz_1 = hamiltonian([["zz",Jzz_1]],[],basis=basis_1,dtype=np.float64)
39 Hx_1 = hamiltonian([["+","hx_1],[-","hx_1]],[],basis=basis_1,dtype=np.float64)
40 Hzz_2 = hamiltonian([["zz",Jzz_2]],[],basis=basis_2,dtype=np.float64,**no_checks)
41 Hx_2 = hamiltonian([["+","hx_2],[-","hx_2]],[],basis=basis_2,dtype=np.float64,**
    no_checks)
42 # calculating bandwidth for non-driven hamiltonian
43 [E_1_min,psi_1] = Hzz_1.eigsh(k=1,which="SA")
44 [E_2_min,psi_2] = Hzz_2.eigsh(k=1,which="SA")
45 # setting up initial states
46 psi0_1 = psi_1.ravel()
47 psi0_2 = psi_2.ravel()
48 # creating generators of time evolution
49 U1_1 = exp_op(Hzz_1+omega*Hx_1,a=-1j*T/4)
50 U2_1 = exp_op(Hzz_1-omega*Hx_1,a=-1j*T/2)
51 U1_2 = exp_op(Hzz_2+omega*Hx_2,a=-1j*T/4)
52 U2_2 = exp_op(Hzz_2-omega*Hx_2,a=-1j*T/2)
53 # get generator objects to get time dependent states
54 psi_1_t = evolve_gen(psi0_1,nT,U1_1,U2_1,U1_1)
55 psi_2_t = evolve_gen(psi0_2,nT,U1_2,U2_2,U1_2)

```

The complete code including the lines that produce Fig. ?? is available in Example Code ??.



### 3 New Horizons for QuSpin

- 2D lattices
- single-particle Hamiltonian class
- Liouville dynamics

We would much appreciate it if the users could report bugs using the [issues](#) forum in the QuSpin online repository.

### Acknowledgements

We would like to thank L. Pollet, M. Kolodrubetz, S. Capponi ... for various stimulating discussions and for providing comments on the draft. The authors are pleased to acknowledge that the computational work reported on in this paper was performed on the Shared Computing Cluster which is administered by [Boston University's Research Computing Services](#). The authors also acknowledge the Research Computing Services group for providing consulting support which has contributed to the results reported within this paper. We would also like to thank [Github](#) for providing the online resources to help develop and maintain this project.

**Funding information** This work was supported by ???

### A Installation Guide in a Few Steps

QuSpin is currently only being supported for Python 2.7 and Python 3.5 and so one must make sure to install this version of Python. We recommend the use of the free package manager [Anaconda](#) which installs Python and manages its packages. For a lighter installation, one can use [miniconda](#).

#### A.1 Mac OS X/Linux

To install Anaconda/miniconda all one has to do is execute the installation script with administrative privilege. To do this, open up the terminal and go to the folder containing the downloaded installation file and execute the following command:

```
$ sudo bash <installation_file>
```

You will be prompted to enter your password. Follow the prompts of the installation. We recommend that you allow the installer to prepend the installation directory to your PATH variable which will make sure this installation of Python will be called when executing a Python script in the terminal. If this is not done then you will have to do this manually in your bash profile file:

```
$ export PATH="path\_to/anaconda/bin:$PATH"
```

**Installing via Anaconda.**—Once you have Anaconda/miniconda installed, all you have to do to install QuSpin is to execute the following command into the terminal:

```
$ conda install -c weinbe58 quspin
```

If asked to install new packages just say ‘yes’. To keep the code up-to-date, just run this command regularly.

**Installing Manually.**—Installing the package manually is not recommended unless the above method failed. Note that you must have the Python packages NumPy, SciPy, and Joblib installed before installing QuSpin. Once all the prerequisite packages are installed, one can download the source code from [github](#) and then extract the code to whichever directory one desires. Open the terminal and go to the top level directory of the source code and execute:

```
$ python setup.py install --record install_file.txt
```

This will compile the source code and copy it to the installation directory of Python recording the installation location to `install_file.txt`. To update the code, you must first completely remove the current version installed and then install the new code. The `install_file.txt` can be used to remove the package by running:

```
$ cat install_file.txt | xargs rm -rf.
```

## A.2 Windows

To install Anaconda/miniconda on Windows, download the installer and execute it to install the program. Once Anaconda/miniconda is installed open the conda terminal and do one of the following to install the package:

**Installing via Anaconda.**—Once you have Anaconda/miniconda installed all you have to do to install QuSpin is to execute the following command into the terminal:

```
> conda install -c weinbe58 quspin
```

If asked to install new packages just say ‘yes’. To update the code just run this command regularly.

**Installing Manually.**—Installing the package manually is not recommended unless the above method failed. Note that you must have NumPy, SciPy, and Joblib installed before installing QuSpin. Once all the prerequisite packages are installed, one can download the source code from [github](#) and then extract the code to whichever directory one desires. Open the terminal and go to the top level directory of the source code and then execute:

```
> python setup.py install --record install_file.txt
```

This will compile the source code and copy it to the installation directory of Python and record the installation location to `install_file.txt`. To update the code you must first completely remove the current version installed and then install the new code.

## B Basic Use of Command Line to Run Python

In this appendix we will review how to use the command line for Windows and OS X/Linux to navigate your computer's folders/directories and run the Python scripts.

### B.1 Mac OS X/Linux

Some basic commands:

- change directory:

```
$ cd < path_to_directory >
```

- list files in current directory:

```
$ ls
```

list files in another directory:

```
$ ls < path_to_directory >
```

- make new directory:

```
$ mkdir <path>/< directory_name >
```

- copy file:

```
$ cp < path >/< file_name > < new_path >/< new_file_name >
```

- move file or change file name:

```
$ mv < path >/< file_name > < new_path >/< new_file_name >
```

- remove file:

```
$ rm < path_to_file >/< file_name >
```

Unix also has an auto complete feature if one hits the TAB key. It will complete a word or stop when it matches more than one file/folder name. The current directory is denoted by "." and the directory above is "..". Now, to execute a Python script all one has to do is open your terminal and navigate to the directory which contains the python script. To execute the script just use the following command:

```
$ python script.py
```

It's that simple!

## B.2 Windows

Some basic commands:

- change directory:

```
> cd < path_to_directory >
```

- list files in current directory:

```
> dir
```

list files in another directory:

```
> dir < path_to_directory >
```

- make new directory:

```
> mkdir <path>\< directory_name >
```

- copy file:

```
> copy < path >\< file_name > < new_path >\< new_file_name >
```

- move file or change file name:

```
> move < path >\< file_name > < new_path >\< new_file_name >
```

- remove file:

```
> erase < path >\< file_name >
```

Windows also has a auto complete feature using the TAB key but instead of stopping when there multiple files/folders with the same name, it will complete it with the first file alphabetically. The current directory is denoted by "." and the directory above is "..".

## B.3 Execute Python Script (any operating system)

To execute a Python script all one has to do is open up a terminal and navigate to the directory which contains the Python script. Python can be recognised by the extension `.py`. To execute the script just use the following command:

```
python script.py
```

It's that simple!

## C Package Documentation

In QuSpin quantum many-body operators are represented as matrices. The computation of these matrices are done through custom code written in Cython. Cython is an optimizing static compiler which takes code written in a syntax similar to Python, and compiles it into a highly efficient C/C++ shared library. These libraries are then easily interfaced with Python, but can run orders of magnitude faster than pure Python code [3]. The matrices are stored in a sparse matrix format using the sparse matrix library of SciPy [4]. This allows QuSpin to easily interface with mature Python packages, such as NumPy, SciPy, any many others. These packages provide reliable state-of-the-art tools for scientific computation as well as support from the Python community to regularly improve and update them [5, 6, 7, 4]. Moreover, we have included specific functionality in QuSpin which uses NumPy and SciPy to do many desired calculations common to ED studies, while making sure the user only has to call a few NumPy or SciPy functions directly. The complete up-to-date documentation for the package is available online under:

<https://github.com/weinbe58/QuSpin/#quspin>

*to report a bug pls visit <https://github.com/weinbe58/QuSpin/issues>*

## D Complete Example Codes

In this appendix, we give the complete python scripts for the dix examples discussed in Sec. 2. In case the reader has trouble with the TAB spaces when copying from the code environments below, the scripts can be downloaded from github at:

<https://github.com/weinbe58/QuSpin/tree/master/examples>

QuSpin *Example Code 1*: The Spectrum of the Transverse Field Ising Model and the Jordan-Wigner Transformation

```

1 from quspin.operators import hamiltonian # Hamiltonians and operators
2 from quspin.basis import spin_basis_1d, fermion_basis_1d # Hilbert space spin basis
3 import numpy as np # generic math functions
4 import matplotlib.pyplot as plt # figure/plot library
5 #
6 ##### define model parameters #####
7 L=8 # system size
8 J=1.0 # spin zz interaction
9 h=np.sqrt(2) # z magnetic field strength
10 #
11 # loop over spin inversion symmetry block variable and boundary conditions
12 for zblock,PBC in zip([-1,1],[1,-1]):
13     #
14     ##### define spin model
15     # site-coupling lists (PBC for both spin inversion sectors)
16     h_field=[[-h,i] for i in range(L)]
17     J_zz=[[-J,i,(i+1)%L] for i in range(L)] # PBC
18     # define spin static and dynamic lists
19     static_spin=[["zz",J_zz],["x",h_field]] # static part of H
20     dynamic_spin=[] # time-dependent part of H
21     # construct spin basis in pos/neg spin inversion sector depending on APBC/PBC
22     basis_spin = spin_basis_1d(L=L,zblock=zblock)
23     # build spin Hamiltonians
24     H_spin=hamiltonian(static_spin,dynamic_spin,basis=basis_spin,dtype=np.float64)
25     # calculate spin energy levels
26     E_spin=H_spin.eigvalsh()
27     #
28     ##### define fermion model
29     # define site-coupling lists for external field
30     h_pot=[[2.0*h,i] for i in range(L)]
31     if PBC==1: # periodic BC: odd particle number subspace only
32         # define site-coupling lists (including boudary couplings)
33         J_pm=[[-J,i,(i+1)%L] for i in range(L)] # PBC
34         J_mp=[[+J,i,(i+1)%L] for i in range(L)] # PBC
35         J_pp=[[-J,i,(i+1)%L] for i in range(L)] # PBC
36         J_mm=[[+J,i,(i+1)%L] for i in range(L)] # PBC
37         # construct fermion basis in the odd particle number subsector
38         basis_fermion = fermion_basis_1d(L=L,Nf=range(1,L+1,2))
39     elif PBC==-1: # anti-periodic BC: even particle number subspace only
40         # define bulk site coupling lists
41         J_pm=[[-J,i,i+1] for i in range(L-1)]
42         J_mp=[[+J,i,i+1] for i in range(L-1)]

```

to report a bug pls visit <https://github.com/weinbe58/QuSpin/issues>

```

43     J_pp=[[-J,i,i+1] for i in range(L-1)]
44     J_mm=[[+J,i,i+1] for i in range(L-1)]
45     # add boundary coupling between sites (L-1,0)
46     J_pm.append([+J,L-1,0]) # APBC
47     J_mp.append([-J,L-1,0]) # APBC
48     J_pp.append([+J,L-1,0]) # APBC
49     J_mm.append([-J,L-1,0]) # APBC
50     # construct fermion basis in the even particle number subsector
51     basis_fermion = fermion_basis_1d(L=L,Nf=range(0,L+1,2))
52     # define fermionic static and dynamic lists
53     static_fermion = [[ "+-", J_pm], [ "-+", J_mp], [ "++", J_pp], [ "--", J_mm], [ 'z', h_pot]]
54     dynamic_fermion=[]
55     # build fermionic Hamiltonian
56     H_fermion=hamiltonian(static_fermion,dynamic_fermion,basis=basis_fermion,
57                           dtype=np.float64,check_pcon=False,check_symm=False)
58     # calculate fermionic energy levels
59     E_fermion=H_fermion.eigvalsh()
60     #
61     ##### plot spectra
62     plt.plot(np.arange(H_fermion.Ns),E_fermion/L,marker='o'
63              ,color='b',label='fermion')
64     plt.plot(np.arange(H_spin.Ns),E_spin/L,marker='x'
65              ,color='r',markersize=2,label='spin')
66     plt.xlabel('state number',fontsize=16)
67     plt.ylabel('energy',fontsize=16)
68     plt.xticks(fontsize=16)
69     plt.yticks(fontsize=16)
70     plt.legend(fontsize=16)
71     plt.grid()
72     plt.tight_layout()
73     plt.show()

```



QuSpin *Example Code 2*: Free Particle Systems: the Fermionic SSH Chain

```

1 from quspin.operators import hamiltonian, exp_op # Hamiltonians and operators
2 from quspin.basis import fermion_basis_1d # Hilbert space fermion basis
3 from quspin.tools.block_tools import block_diag_hamiltonian # block diagonalisation
4 import numpy as np # generic math functions
5 import matplotlib.pyplot as plt # plotting library
6 try: # import python 3 zip function in python 2 and pass if using python 3
7     import itertools.izip as zip
8 except ImportError:
9     pass
10 ##### define model parameters #####
11 L=100 # system size
12 J=1.0 # uniform hopping contribution
13 deltaJ=0.1 # bond dimerisation
14 Delta=0.5 # staggered potential
15 beta=100.0 # set inverse temperature for Fermi-Dirac distribution
16 ##### construct single-particle Hamiltonian #####
17 # define site-coupling lists
18 hop_pm=[[-J-deltaJ*(-1)**i,i,(i+1)%L] for i in range(L)] # PBC
19 hop_mp=[[+J+deltaJ*(-1)**i,i,(i+1)%L] for i in range(L)] # PBC
20 stagg_pot=[[Delta*(-1)**i,i] for i in range(L)]
21 # define static and dynamic lists
22 static=[["+-",hop_pm],["-+",hop_mp],['n',stagg_pot]]
23 dynamic=[]
24 # define basis
25 basis=fermion_basis_1d(L,Nf=1)
26 # build real-space Hamiltonian
27 H=hamiltonian(static,dynamic,basis=basis,dtype=np.float64)
28 # diagonalise real-space Hamiltonian
29 E,V=H.eigh()
30 ##### compute Fourier transform and momentum-space Hamiltonian #####
31 # define basis blocks and arguments
32 blocks=[dict(Nf=1,kblock=i,a=2) for i in range(L//2)] # only L//2 distinct momenta
33 basis_args = (L,)
34 # construct block-diagonal Hamiltonian
35 FT,Hblock = block_diag_hamiltonian(blocks,static,dynamic,fermion_basis_1d,
36                                     basis_args,np.complex128,get_proj_kwargs=dict(pcon=True))
37 # diagonalise momentum-space Hamiltonian
38 Eblock,Vblock=Hblock.eigh()
39 ##### prepare the density observables and initial states #####
40 # grab single-particle states and treat them as initial states
41 psi0=Vblock
42 # construct operator n_1 = $n_{j=0}$
43 n_1_static=[['n',[[1.0,0]]]]
44 n_1=hamiltonian(n_1_static,[],basis=basis,dtype=np.float64,
45                 check_herm=False,check_pcon=False)
46 # construct operator n_2 = $n_{j=L/2}$
47 n_2_static=[['n',[[1.0,L//2]]]]
48 n_2=hamiltonian(n_2_static,[],basis=basis,dtype=np.float64,
49                 check_herm=False,check_pcon=False)
50 # transform n_j operators to momentum space
51 n_1=n_1.rotate_by(FT,generator=False)

```

to report a bug pls visit <https://github.com/weinbe58/QuSpin/issues>

```

52 n_2=n_2.rotate_by(FT,generator=False)
53 ##### evaluate nonequal time correlator <FS|n_2(t) n_1(0)|FS> #####
54 # define time vector
55 t=np.linspace(0.0,90.0,901)
56 # calcualte state acted an by n_1
57 n_psi0=n_1.dot(psi0)
58 # construct time-evolution operator using exp_op class (sometimes faster)
59 U = exp_op(Hblock,a=-1j,start=t.min(),stop=t.max(),num=len(t),iterate=True)
60 # evolve states
61 psi_t=U.dot(psi0)
62 n_psi_t = U.dot(n_psi0)
63 # alternative method for time evolution using Hamiltonian class
64 #psi_t=Hblock.evolve(psi0,0.0,t,iterate=True)
65 #n_psi_t=Hblock.evolve(n_psi0,0.0,t,iterate=True)
66 # preallocate variable
67 correlators=np.zeros(t.shape+psi0.shape[1:])
68 # loop over the time-evolved states
69 for i, (psi,n_psi) in enumerate( zip(psi_t,n_psi_t) ):
70     correlators[i,:]=n_2.matrix_ele(psi,n_psi,diagonal=True).real
71 # evaluate correlator at finite temperature
72 n_FD=1.0/(np.exp(beta*E)+1.0)
73 correlator = (n_FD*correlators).sum(axis=-1)
74 ##### plot spectra
75 plt.plot(np.arange(H.Ns),E/L,
76          marker='o',color='b',label='real space')
77 plt.plot(np.arange(Hblock.Ns),Eblock/L,
78          marker='x',color='r',markersize=2,label='momentum space')
79 plt.xlabel('state number',fontsize=16)
80 plt.ylabel('energy',fontsize=16)
81 plt.xticks(fontsize=16)
82 plt.yticks(fontsize=16)
83 plt.legend(fontsize=16)
84 plt.grid()
85 plt.tight_layout()
86 plt.show()
87 ##### plot correlator
88 plt.plot(t,correlator,linewidth=2)
89 plt.xlabel('$t$',fontsize=16)
90 plt.ylabel('$C_{0,L/2}(t,\\beta)$',fontsize=16)
91 plt.xticks(fontsize=16)
92 plt.yticks(fontsize=16)
93 plt.grid()
94 plt.tight_layout()
95 plt.show()

```

QuSpin *Example Code 3*: Many-Body Localisation in the Fermi-Hubbard Model

```

1 from __future__ import print_function, division
2 from quspin.operators import hamiltonian, exp_op, ops_dict # operators
3 from quspin.basis import tensor_basis, fermion_basis_1d # Hilbert spaces
4 from quspin.tools.measurements import obs_vs_time # calculating dynamics
5 import numpy as np # general math functions
6 from numpy.random import uniform, choice # tools for doing random sampling
7 from time import time # tool for calculating computation time
8 import matplotlib.pyplot as plt # plotting
9 #
10 ##### setting parameters for simulation
11 # simulation parameters
12 n_real = 100 # number of realizations
13 n_boot = 100 # number of bootstrap samples to calculate error
14 # physical parameters
15 L = 8 # system size
16 N = L//2 # number of particles
17 N_up = N//2 + N % 2 # number of fermions with spin up
18 N_down = N//2 # number of fermions with spin down
19 w_list = [1.0, 4.0, 10.0] # disorder strength
20 J = 1.0 # hopping strength
21 U = 5.0 # interaction strength
22 # range in time to evolve system
23 start, stop, num = 0.0, 35.0, 101
24 t = np.linspace(start, stop, num=num, endpoint=True)
25 #
26 ##### create the basis
27 # build the two bases to tensor together to spinful fermions
28 basis_up = fermion_basis_1d(L, Nf=N_up) # up basis
29 basis_down = fermion_basis_1d(L, Nf=N_down) # down basis
30 basis = tensor_basis(basis_up, basis_down) # spinful fermions
31 #
32 ##### create model
33 # define site-coupling lists
34 hop_right = [[-J, i, i+1] for i in range(L-1)] # hopping to the right OBC
35 hop_left = [[J, i, i+1] for i in range(L-1)] # hopping to the left OBC
36 int_list = [[U, i, i] for i in range(L)] # onsite interaction
37 # site-coupling list to create the sublattice imbalance observable
38 sublat_list = [[(-1.0)**i/N, i] for i in range(0, L)]
39 # create static lists
40 operator_list_0 = [
41     ["+-|", hop_left], # up hop left
42     ["-+|", hop_right], # up hop right
43     ["|+-", hop_left], # down hop left
44     ["|-+", hop_right], # down hop right
45     ["n|n", int_list], # onsite interaction
46 ]
47 imbalance_list = [["n|", sublat_list], ["|n", sublat_list]]
48 # create operator dictionary for ops_dict class
49 # add key for Hubbard hamiltonian
50 operator_dict = dict(H0=operator_list_0)
51 # add keys for local potential in each site

```

to report a bug pls visit <https://github.com/weinbe58/QuSpin/issues>

```

52 for i in range(L):
53     # add to dictioanry keys h0,h1,h2,...,hL with local potential operator
54     operator_dict["n"+str(i)] = [["n|",[[1.0,i]]],["|n",[[1.0,i]]]]
55 #
56 ##### setting up operators
57 # set up hamiltonian dictionary and observable (imbalance I)
58 no_checks = dict(check_pcon=False,check_symm=False,check_herm=False)
59 H_dict = ops_dict(operator_dict,basis=basis,**no_checks)
60 I = hamiltonian(imbalance_list,[],basis=basis,**no_checks)
61 # strings which represent the initial state
62 s_up = "".join("1000" for i in range(N_up))
63 s_down = "".join("0010" for i in range(N_down))
64 # basis.index accepts strings and returns the index
65 # which corresponds to that state in the basis list
66 i_0 = basis.index(s_up,s_down) # find index of product state
67 psi_0 = np.zeros(basis.Ns) # allocate space for state
68 psi_0[i_0] = 1.0 # set MB state to be the given product state
69 print("H-space size: {:d}, initial state: |{:s}>(x)|{:s}>".format(basis.Ns,s_up,
    s_down))
70 #
71 # define function to do dynamics for different disorder realizations.
72 def real(H_dict,I,psi_0,w,t,i):
73     # body of function goes below
74     ti = time() # start timing function for duration of reach realisation
75     # create a parameter list which specifies the onsite potential with disorder
76     params_dict=dict(H0=1)
77     for i in range(L):
78         params_dict["n"+str(i)] = uniform(-w,w)
79     # using the parameters dictionary construct a hamiltonian object with those
80     # parameters defined in the list
81     H = H_dict.tohamiltonian(params_dict)
82     # use exp_op to get the evolution operator
83     U = exp_op(H,a=-1j,start=t.min(),stop=t.max(),num=len(t),iterate=True)
84     psi_t = U.dot(psi_0) # get generator psi_t for time evolved state
85     # use obs_vs_time to evaluate the dynamics
86     t = U.grid # extract time grid stored in U, and defined in exp_op
87     obs_t = obs_vs_time(psi_t,t,dict(I=I))
88     # print reporting the computation time for realization
89     print("realization {}/{} completed in {:.2f} s".format(i+1,n_real,time()-ti))
90     # return observable values
91     return obs_t["I"]
92 #
93 ##### looping over differnt disorder strengths
94 for w in w_list:
95     I_data = np.vstack([real(H_dict,I,psi_0,w,t,i) for i in range(n_real)])
96     ##### averaging and error estimation
97     I_avg = I_data.mean(axis=0) # get mean value of I for all time points
98     # generate bootstrap samples
99     bootstrap_gen = (I_data[choice(n_real,size=n_real)].mean(axis=0) for i in range(
    n_boot))
100     # generate the fluctuations about the mean of I
101     sq_fluc_gen = ((bootstrap-I_avg)**2 for bootstrap in bootstrap_gen)
102     I_error = np.sqrt(sum(sq_fluc_gen)/n_boot)

```

```
103     ##### plotting results
104     plt.errorbar(t,I_avg,I_error,marker=".",label="w={:.2f}".format(w))
105 # configuring plots
106 plt.xlabel("$t/J$",fontsize=18)
107 plt.ylabel("$\mathcal{I}$",fontsize=18)
108 plt.grid(True)
109 plt.tick_params(labelsize=16)
110 plt.legend(loc=0)
111 plt.savefig('fermion_MBL.pdf', bbox_inches='tight')
112 plt.show()
```

*to report a bug pls visit <https://github.com/weinbe58/QuSpin/issues>*

QuSpin *Example Code 4*: The Gross-Pitaevskii Equation and Nonlinear Time Evolution

```

1 from quspin.operators import hamiltonian # Hamiltonians and operators
2 from quspin.basis import boson_basis_1d # Hilbert space boson basis
3 from quspin.tools.measurements import evolve # nonlinear evolution
4 import numpy as np # generic math functions
5 import matplotlib.pyplot as plt # plot library
6 #
7 ##### define model parameters #####
8 L=300 # system size
9 # calculate centre of chain
10 if L%2==0:
11     j0 = L//2-0.5 # centre of chain
12 else:
13     j0 = L//2 # centre of chain
14 sites=np.arange(L)-j0
15 # static parameters
16 J=1.0 # hopping
17 U=1.0 # Bose-Hubbard interaction strength
18 # dynamic parameters
19 kappa_trap_i=0.001 # initial chemical potential
20 kappa_trap_f=0.0001 # final chemical potential
21 t_ramp=40.0/J # set total ramp time
22 # ramp protocol
23 def ramp(t,kappa_trap_i,kappa_trap_f,t_ramp):
24     return (kappa_trap_f - kappa_trap_i)*t/t_ramp + kappa_trap_i
25 # ramp protocol parameters
26 ramp_args=[kappa_trap_i,kappa_trap_f,t_ramp]
27 #
28 ##### construct single-particle Hamiltonian #####
29 # define site-coupling lists
30 hopping=[[-J,i,(i+1)%L] for i in range(L-1)]
31 trap=[[0.5*(i-j0)**2,i] for i in range(L)]
32 # define static and dynamic lists
33 static=[["+-",hopping],["-+",hopping]]
34 dynamic=[['n',trap,ramp,ramp_args]]
35 # define basis
36 basis = boson_basis_1d(L,Nb=1,sps=2)
37 # build Hamiltonian
38 Hsp=hamiltonian(static,dynamic,basis=basis,dtype=np.float64)
39 E,V=Hsp.eigsh(time=0.0,k=1,which='SA')
40 #
41 ##### imaginary-time evolution to compute GS of GPE #####
42 def GPE_imag_time(tau,phi,Hsp,U):
43     """
44     This function solves the real-valued GPE in imaginary time:
45     $$ -\dot{\phi}(\tau) = Hsp(t=0)\phi(\tau) + U |\phi(\tau)|^2 \phi(\tau) $$
46     """
47     return -( Hsp.dot(phi,time=0) + U*np.abs(phi)**2*phi )
48 # define ODE parameters
49 GPE_params = (Hsp,U)
50 # define initial state to flow to GS from
51 phi0=V[:,0]*np.sqrt(L) # initial state normalised to 1 particle per site

```

to report a bug pls visit <https://github.com/weinbe58/QuSpin/issues>

```

52 # define imaginary time vector
53 tau=np.linspace(0.0,35.0,71)
54 # evolve state in imaginary time
55 psi_tau = evolve(phi0,tau[0],tau,GPE_imag_time,f_params=GPE_params,
56                 imag_time=True,real=True,iterate=True)
57 #
58 # display state evolution
59 for i,psi0 in enumerate(psi_tau):
60     # compute energy
61     E_GS=(Hsp.matrix_ele(psi0,psi0,time=0) + 0.5*U*np.sum(np.abs(psi0)**4)).real
62     # plot wave function
63     plt.plot(sites, abs(phi0)**2, color='r',marker='s',alpha=0.2,
64             label='$|\\phi_j(0)|^2$')
65     plt.plot(sites, abs(psi0)**2, color='b',marker='o',
66             label='$|\\phi_j(\\tau)|^2$')
67     plt.xlabel('$\\mathrm{lattice\\ sites}$',fontsize=14)
68     plt.title('$J\\tau=0.2f,\\ E\\mathrm{\\{GS\\}}(\\tau)=0.4f$'%(tau[i],E_GS)
69             ,fontsize=14)
70     plt.ylim([-0.01,max(abs(phi0)**2)+0.01])
71     plt.legend(fontsize=14)
72     plt.draw() # draw frame
73     plt.pause(0.005) # pause frame
74     plt.clf() # clear figure
75 plt.close()
76 #
77 ##### real-time evolution of GPE #####
78 def GPE(time,psi):
79     """
80     This function solves the complex-valued time-dependent GPE:
81     $$ i\\dot{\\psi}(t) = Hsp(t)\\psi(t) + U |\\psi(t)|^2 \\psi(t) $$
82     """
83     # solve static part of GPE
84     psi_dot = Hsp.static.dot(psi) + U*np.abs(psi)**2*psi
85     # solve dynamic part of GPE
86     for Hd,f,f_args in Hsp.dynamic:
87         psi_dot += f(time,*f_args)*Hd.dot(psi)
88     return -1j*psi_dot
89 # define real time vector
90 t=np.linspace(0.0,t_ramp,101)
91 # time-evolve state according to GPE
92 psi_t = evolve(psi0,t[0],t,GPE,iterate=True,atol=1E-12,rtol=1E-12)
93 #
94 # display state evolution
95 for i,psi in enumerate(psi_t):
96     # compute energy
97     E=(Hsp.matrix_ele(psi,psi,time=t[i]) + 0.5*U*np.sum(np.abs(psi)**4)).real
98     # compute trap
99     kappa_trap=ramp(t[i],kappa_trap_i,kappa_trap_f,t_ramp)*(sites)**2
100    # plot wave function
101    plt.plot(sites, abs(psi0)**2, color='r',marker='s',alpha=0.2
102            ,label='$|\\psi_{\\mathrm{GS},j}|^2$')
103    plt.plot(sites, abs(psi)**2, color='b',marker='o',label='$|\\psi_j(t)|^2$')
104    plt.plot(sites, kappa_trap, '--',color='g',label='$\\mathrm{\\{trap\\}}$')

```



```

105 plt.ylim([-0.01,max(abs(psi0)**2)+0.01])
106 plt.xlabel('$\mathrm{lattice}$ sites$', fontsize=14)
107 plt.title('$Jt=0.2f, E(t)-E_\mathrm{GS}=0.4fJ$'%(t[i],E-E_GS), fontsize=14)
108 plt.legend(loc='upper right', fontsize=14)
109 plt.draw() # draw frame
110 plt.pause(0.00005) # pause frame
111 plt.clf() # clear figure
112 plt.close()

```

## References

- [1] P. Weinberg and M. Bukov, *QuSpin: a Python Package for Dynamics and Exact Diagonalisation of Quantum Many Body Systems part I: spin chains*, SciPost Phys. **2**, 003 (2017), doi:[10.21468/SciPostPhys.2.1.003](https://doi.org/10.21468/SciPostPhys.2.1.003).
- [2] A. W. Sandvik, *Computational studies of quantum spin systems*, AIP Conference Proceedings **1297**(1), 135 (2010), doi:<http://dx.doi.org/10.1063/1.3518900>.
- [3] S. Behnel, R. Bradshaw, C. Citro, L. Dalcin, D. S. Seljebotn and K. Smith, *Cython: The best of both worlds*, Computing in Science & Engineering **13**(2), 31 (2011), doi:<http://dx.doi.org/10.1109/MCSE.2010.118>.
- [4] E. Jones, T. Oliphant, P. Peterson *et al.*, *SciPy: Open source scientific tools for Python* (2001–).
- [5] S. v. d. Walt, S. C. Colbert and G. Varoquaux, *The numpy array: A structure for efficient numerical computation*, Computing in Science & Engineering **13**(2), 22 (2011), doi:<http://dx.doi.org/10.1109/MCSE.2011.37>.
- [6] T. E. Oliphant, *Python for scientific computing*, Computing in Science & Engineering **9**(3), 10 (2007), doi:<http://dx.doi.org/10.1109/MCSE.2007.58>.
- [7] K. J. Millman and M. Aivazis, *Python for scientists and engineers*, Computing in Science & Engineering **13**(2), 9 (2011), doi:<http://dx.doi.org/10.1109/MCSE.2011.36>.