



For numbered affiliations see end of article.

Correspondence to

Dr Heng-Cheng Hsu;
b101092037@gmail.com

Received 4 April 2024
Accepted 11 July 2024
Published Online First
7 August 2024

Cervical cancer: a new era

Giuseppe Caruso ^{1,2,3} Matthew K Wagar ⁴ Heng-Cheng Hsu ^{5,6,7} Jorge Hoegl ⁸
Guido Martin Rey Valzacchi ⁹ Andreina Fernandes ¹⁰ Giuseppe Cucinella ¹¹
Seda Sahin Aker ^{12,13} Aarthi S Jayraj ¹⁴ Jessica Mauro¹⁵ Rene Pareja ¹⁶
Pedro T Ramirez ¹⁷

ABSTRACT

Cervical cancer is a major global health issue, ranking as the fourth most common cancer in women worldwide. Depending on stage, histology, and patient factors, the standard management of cervical cancer is a combination of treatment approaches, including (fertility- or non-fertility-sparing) surgery, radiotherapy, platinum-based chemotherapy, and novel systemic therapies such as bevacizumab, immune checkpoint inhibitors, and antibody-drug conjugates. While ambitious global initiatives seek to eliminate cervical cancer as a public health problem, the management of cervical cancer continues to evolve with major advances in imaging modalities, surgical approaches, identification of histopathological risk factors, radiotherapy techniques, and biomarker-driven personalized therapies. In particular, the introduction of immune checkpoint inhibitors has dramatically altered the treatment of cervical cancer, leading to significant survival benefits in both locally advanced and metastatic/recurrent settings. As the landscape of cervical cancer therapies continues to evolve, the aim of the present review is to provide a comprehensive discussion of the current state and the latest practice-changing updates in cervical cancer.

INTRODUCTION

Cervical cancer is the fourth most common cancer in women worldwide and the most common gynecological cancer globally.¹ Despite the effectiveness of the human papillomavirus (HPV) vaccination and screening,^{2,3} cervical cancer remains a major global health issue, with a particularly high burden in low- and middle-income countries due to wide disparities in HPV vaccination coverage, access to preventive interventions, and socioeconomic factors.^{4,5} The 5-year overall survival rate for all stages combined is 67%: 91% for early stage, 60% for locally advanced disease, and 19% for metastatic disease.⁶ Almost all cases of cervical cancer (~95%) are caused by persistent infection with high-risk HPV. The most common histologies are squamous cell carcinoma (~80%) and adenocarcinoma (~20%), while other histologic subtypes remain rare.^{7,8}

Cervical cancer is generally curable if diagnosed and treated at an early stage. The recommended treatment for early-stage cervical cancer (International Federation of Gynecology and Obstetrics (FIGO) 2018 stages IA–IB2, and IIA1) is surgery followed by

observation or adjuvant therapy (radiotherapy±concurrent chemotherapy) based on histopathologic risk factors on definitive pathology. Conservative and fertility-sparing approaches may be considered in carefully selected patients. Standard of care for locally advanced cervical cancer (FIGO 2018 stages IB3, IIA2–IVA) is definitive concurrent chemoradiotherapy.^{9–11} However, recent findings have the potential to improve the standard of care of patients with locally advanced cervical cancer. Based on results of the KEYNOTE-A18 trial, the US Food and Drug Administration (FDA) recently approved pembrolizumab in addition to concurrent chemoradiotherapy for patients with high-risk locally advanced cervical cancer.¹² Moreover, results from the INTERLACE trial presented at the 2023 European Society for Medical Oncology (ESMO) congress demonstrated that induction chemotherapy before standard concurrent chemoradiotherapy provides a significant improvement in overall survival.¹³ Patients with metastatic (FIGO 2018 stage IVB), persistent, or recurrent cervical cancer should be treated at tertiary referral centers in a multidisciplinary setting. The management of these patients poses a major clinical challenge and relies on combined treatment modalities, including platinum-based chemotherapy±bevacizumab±pembrolizumab, locoregional treatments (surgery, radiotherapy), enrollment in clinical trials, and/or palliative care.^{9–11}

The WHO Global Initiative was launched to accelerate the elimination of cervical cancer as a public health problem by using three main pillars: HPV vaccination, effective screening (Pap test, HPV test, colposcopy), and treatment of precancerous lesions. Meanwhile, there have been continuous and significant improvements in imaging modalities, surgical approaches (tailored parametrectomy, less radical surgery, sentinel lymph node (SLN) biopsy), pathological diagnosis, radiotherapy techniques (image-guided, intensity-modulated radiation), and novel treatment options with targeted agents, including immune checkpoint inhibitors and antibody-drug conjugates.

Given recent practice-changing advances in the management of cervical cancer, this narrative review aimed to summarize the contemporary state and latest updates in the management of cervical cancer.



© IGCS and ESGO 2024. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Caruso G, Wagar MK, Hsu H-C, et al. *Int J Gynecol Cancer* 2024;**34**:1946–1970.

METHODS

A formal systematic review was not performed, but relevant literature in cervical cancer was identified by searching PubMed for English-language articles from inception to March 2024. Priority was given to the selection of randomized controlled trials, systematic reviews, meta-analyses, and practice guidelines. The references of identified articles were also reviewed. Selected articles were mutually agreed by the authors. Details of the study design and results of the pivotal studies underpinning treatment recommendations in cervical cancer are reported in Table 1.

Epidemiology

Cervical cancer is the fourth leading cause of cancer incidence and mortality in women worldwide, with more than 85% of cases occurring in low-resource settings.¹⁴ In 2022, there were an estimated 661 021 cervical cancer cases and 348 189 deaths globally.¹ The cumulative lifetime risk of developing and dying from cervical cancer are 1.50% and 0.79%, respectively.¹ The burden of cervical cancer incidence is not equal worldwide, with a clear socioeconomic gradient. In countries with a low human development index, cervical cancer incidence and mortality rates are three and six times higher, respectively, than in countries with a high human development index.¹⁵ Specific regions of the world with particularly high incidence and mortality rates include East and Southeast Asia (57.3% of global cases, 57.6% of deaths), sub-Saharan Africa (16.6% of cases, 19.3% of deaths), and Latin America (9.2% of cases, 8.5% of deaths).¹⁵

The incidence of cervical cancer by stage varies internationally as well, with patients in low human development index countries more likely to be diagnosed at a later stage. Worldwide, around 40–50% of cervical cancers are diagnosed at an early stage, 35–40% at locally advanced stage, and 10–25% at metastatic stage.¹⁶ Geographic and socioeconomic disparities in the incidence and global burden of cervical cancer reflect inequalities in the availability of the HPV vaccine and accessibility to adequate screening and treatment of precancerous lesions. The main risk factor for cervical cancer is persistent HPV infection, with HPV 16 and 18 being the most common high-risk genotypes.¹⁷ More than 70% of cervical cancer diagnoses are attributed to HPV 16/18 genotypes, while other known risk factors include smoking, higher number of sexual partners, and HIV infection, highlighting the importance of risk reduction strategies aimed at smoking cessation and HIV prevention.¹⁸

Pathophysiology

High-risk HPV is a sexually transmitted virus capable of establishing persistent infections leading to precancerous lesions and cervical cancer. HPV is a non-enveloped, double-stranded DNA virus of 8 kilobase pairs containing eight open reading frames, including two structural genes (*L1*, *L2*) and six early genes involved in viral replication (*E1*, *E2*, *E4*, *E5*, *E6*, *E7*).^{19,20} After HPV infection, distinct mRNA splicing patterns and multiple promoters control the expression of viral genes, allowing expression of gene products at different stages of the viral life cycle.

E6 and *E7* have an essential function during the viral life cycle, increasing viral fitness and virion production.¹⁹ The *E6* protein is able to directly interact with p53, a tumor suppressor protein, and induce ubiquitin-dependent proteolysis. The *E7* protein binds to pRb

suppressor protein to facilitate the release of transcription factors such as E2F and induce gene transcription, resulting in disruption of cell cycle regulatory pathways, resistance to apoptosis, and dysregulated cell proliferation.^{21,22} Persistent HPV infection leads to integration of the viral genome into the cellular genome, promoting sustained expression of *E6* and *E7*, and loss of the *E2* gene, a repressor of the expression of these oncoproteins (Figure 1). Additionally, HPV integration induces genetic alterations, amplification of oncogenes, chromosomal rearrangements, and instability.²² Most cervical cancer cases demonstrate integrated HPV genomes. In cases where the genome is in an episomal form, dysregulation occurs by epigenetic modifications such as DNA methylation, histone modifications, non-coding RNA regulation and chromatin regulation.²³

Prevention and Screening

In 2020, WHO launched a global strategy to eliminate cervical cancer as a public health problem by 2130, with the goal of a global incidence of less than 4 per 100 000 women per year. The strategy focuses on three key pillars: (1) prevention through HPV vaccination; (2) screening and treatment of precancerous lesions; and (3) adequate treatment of invasive cervical cancer. WHO countries are expected to achieve the following three main targets by 2030 to reduce the median cervical cancer incidence by 10%: (1) 90% of girls fully vaccinated against HPV by 15 years of age; (2) 70% of women screened with high-performance tests for HPV detection by 35 years of age and again by 45 years of age; and (3) 90% of women adequately treated for precancerous lesions.²⁴

Prophylactic HPV vaccination has been shown to decrease the burden of HPV-associated disease. Currently, there are five commercial prophylactic HPV vaccines: two bivalent vaccines (HPV 16 and 18), two tetravalent vaccines (HPV 6, 11, 16, and 18), and one nonavalent vaccine (HPV 6, 11, 16, 18, 31, 33, 45, 52, and 58). In addition, there is emerging evidence on the safety and immunogenicity of a new nonavalent vaccine produced in *Escherichia coli*, a simple and cost-effective system for the production of L1 virus-like particles, with potential to alleviate supply constraints and affordability of HPV vaccine.^{20,25,26} WHO recommends either a single or two-dose schedule in women under 20 years of age to improve vaccination coverage.^{27,28} The primary target of vaccination is girls aged 9–14 years, prior to the start of sexual activity. Vaccination of secondary targets, such as males and older females, is also recommended where feasible and affordable.

Primary screening strategies are a critical component of cervical cancer prevention and include cervical cytology (Pap smear), HPV co-testing, and colposcopy (with biopsy). The main objective of screening programs is to detect precancerous lesions to provide prompt treatment and appropriate follow-up before cancer develops.²⁰ About 50% of cervical cancers occur in women with inadequate screening.²⁹ In a recent estimate of screening coverage across 202 countries, 1.6 billion (67%) of 2.3 billion women aged 20–70 years had never been screened for cervical cancer. According to Bruni et al, 69% of countries surveyed have official recommendations for primary cervical cancer screening, with only 29% sending individual reminders to women.³⁰

The use of polymerase chain reaction (PCR)-based tests for HPV genome detection has proven to be cost-effective while increasing sensitivity for detecting high-grade lesions.^{20,31,32} Several countries

Table 1 Pivotal prospective studies underpinning treatment recommendations in cervical cancer

Study	Design	Setting	N	Treatment	Results	Conclusion
Early-stage disease and LACC						
SHAPE ⁶⁷	Phase III, multicenter, non-inferiority RCT	FIGO 2009 stage IA2–IB1 squamous cell carcinoma, adenocarcinoma, or adenosquamous carcinoma, with stromal invasion <10 mm on LEEP/cone biopsy or <50% invasion on preoperative MRI. LVSI+ allowed	700	Arm 1 (n=350): simple hysterectomy with LND Arm 2 (n=350): radical hysterectomy with LND	Incidence of pelvic recurrence at 3 years was 2.17% in the radical hysterectomy group and 2.52% in the simple hysterectomy group (an absolute difference of 0.35 percentage points; 90% CI –1.62 to 2.32). The incidence of urinary incontinence was lower in the simple hysterectomy group than in the radical hysterectomy group within 4 weeks after surgery (2.4% vs 5.5%; p=0.048) and beyond 4 weeks (4.7% vs 11.0%; p=0.003). The incidence of urinary retention in the simple hysterectomy group was also lower than that in the radical hysterectomy group within 4 weeks after surgery (0.6% vs 11.0%; p<0.001) and beyond 4 weeks (0.6% vs 9.9%; p<0.001)	In patients with low-risk cervical cancer, simple hysterectomy is not inferior to radical hysterectomy with respect to the 3-year incidence of pelvic recurrence and is associated with a lower risk of urinary incontinence or retention
SENTIX/CEEGOG-cx01/ENGOT-cx2 ⁶⁹	Multicenter, prospective cohort study	FIGO 2009 stage IA1 LVSI+ to IB2 (<4 cm) squamous cell carcinoma, adenocarcinoma, or adenosquamous carcinoma, with no suspicious lymph nodes on imaging	647	All patients underwent SLN biopsy with ultrastaging	The 2-year recurrence rate was 6%. SLN biopsy showed a bilateral detection rate of 92.3% and was able to detect lymph node involvement in >56% of cases by frozen section and >90% after ultrastaging	The study validated the hypothesis that the 2-year recurrence rate post-SLN biopsy was non-inferior to the reference recurrence rate in historical cohorts following pelvic lymph node dissection
LACC ⁶⁰	Phase III, multicenter RCT	FIGO 2009 stages IA1 LVSI+ to IB1 squamous cell carcinoma, adenocarcinoma, or adenosquamous carcinoma	631	Arm 1 (n=319): MIS radical hysterectomy Arm 2 (n=312): open radical hysterectomy	Disease-free survival at 4.5 years was 86.0% with MIS and 96.5% with open surgery (95% CI –16.4 to –4.7). MIS was associated with a lower rate of disease-free survival than open (3-year rate 91.2% vs 97.1%; HR 3.74; 95% CI 1.63 to 8.58). MIS was also associated with a lower rate of 3-year overall survival (93.8% vs 99.0%; HR 6.00; 95% CI 1.77 to 20.30)	MIS radical hysterectomy is associated with lower rates of disease-free survival and overall survival than open radical hysterectomy in early-stage cervical cancer

Continued

Table 1 Continued

Study	Design	Setting	N	Treatment	Results	Conclusion
ConCerv ⁶⁶	Multicenter, prospective, single-arm study	FIGO 2009 stage IA2–IB1 squamous cell carcinoma (any grade), or adenocarcinoma (grade 1 or 2), with tumor size <2 cm, no LVSI, depth of stromal invasion <10 mm, negative imaging for metastatic disease, and negative conization margins	100	Patients underwent conservative surgery with either conization (if fertility desired) and LND (n=44), conization followed by simple hysterectomy and LND (n=40), or inadvertent simple hysterectomy followed by LND (n=16). One additional conization was allowed	Five patients (5%) were found to have positive lymph nodes. In one out of 40 patients, residual disease was observed in the post-conization hysterectomy specimen, indicating an immediate failure rate of 2.5%. The median follow-up period was 36.3 months, ranging from 0.0 to 68.3 months. Within 2 years post-surgery, three patients experienced recurrent disease, corresponding to a cumulative incidence of 3.5% (95% CI 0.9% to 9.0%) There was a term delivery rate of 92.9% among the 14 pregnancies of 11 patients who underwent cervical conization for fertility preservation and remained in the study	Patients diagnosed with early-stage low-risk cervical carcinoma may be considered as candidates for conservative surgical interventions
GOG 278 ⁶⁸	Phase I/II, multicenter, prospective study	FIGO 2009 stage IA1 (LVSI+) and IA2–IB1 (≤2 cm); squamous cell carcinoma, adenocarcinoma, or adenosquamous carcinoma	224	Cone biopsy (n=72) or simple hysterectomy (n=152) plus pelvic LND	Of the 201 patients eligible for survival analysis, only three experienced a recurrence after a median follow-up of 37 months. Among the 68 patients who underwent cone biopsy and were eligible for analysis, 31 patients attempted to conceive and 16 pregnancies (15 patients) were obtained, including nine full-term pregnancies	Non-radical surgery for early-stage cervical cancer appears safe. Patients treated by cone biopsy and pelvic LND can achieve successful fertility outcomes
GOG 92 (Sedlis criteria) ⁹⁶	Multicenter RCT	Stage IB squamous cell carcinoma, adenocarcinoma, or adenosquamous carcinoma, with LVSI+, >1/3 stromal invasion and large tumor diameter following radical hysterectomy with LND	277	Arm 1 (n=137): pelvic RT Arm 2 (n=140): no further treatment (NFT)	47% reduction in risk of recurrence (RR 0.53; p=0.008, one-tail) among the RT group, with recurrence-free rates at 2 years of 88% vs 79% for the RT and NFT groups, respectively 6% grade 3–4 AEs in RT vs 2.1% in the NFT group	Adjuvant pelvic RT following radical surgery reduces the number of recurrences in patients meeting Sedlis criteria There was no statistically significant difference in overall survival at a median follow-up of 10 years
GOG 109/SWOG 8797/RTOG 9112 (Peters criteria) ⁹⁴	Multicenter RCT	Stage IA2, IB, and IIA cervical cancer, treated with radical hysterectomy and pelvic LND, and with positive pelvic lymph nodes, positive margins, and/or microscopic involvement of the parametrium	243	Arm 1 (n=127): RT + cisplatin/5-FU chemotherapy Arm 2 (n=116): RT alone	HRs for progression-free survival and overall survival in the RT only arm vs the CRT arm are 2.01 (p=0.003) and 1.96 (p=0.007)	Concurrent cisplatin-based CRT significantly improves progression-free survival and overall survival for patients with high-risk, early-stage disease who undergo radical hysterectomy and pelvic LND

Continued

Table 1 Continued

Study	Design	Setting	N	Treatment	Results	Conclusion
NRG oncology/ RTOG 0724/GOG- 0724 ⁹⁵	Phase III, multicenter RCT	Stage IA2, IB, or IIA cervical carcinoma with at least one of the high- risk features after radical hysterectomy: positive pelvic nodes, positive parametrium, and/or positive para-aortic nodes	212	Arm 1 (n=109): CRT Arm 2 (n=103): CRT followed by four cycles of adjuvant chemotherapy with paclitaxel and carboplatin (CRT + chemo)	4 year disease-free survival estimates of 76% (90% CI 69% to 83%) and 77% (90% CI 70% to 84%) for the CRT and CRT+chemo arms, respectively (HR 1.05; 90% CI 0.65 to 1.68; p=0.56) 4-years overall survival is 87% (90% CI 82% to 93%) and 89% (90% CI 84% to 94%) for the CRT and CRT+chemo arms, respectively (HR 0.91; 90% CI 0.49 to 1.69; one- sided log-rank p=0.40)	The addition of systemic chemotherapy following CRT did not improve disease-free or overall survival in patients with cervical cancer and high- risk factors after radical hysterectomy
EMBRACE I ¹⁶¹	Multicenter, prospective cohort study	FIGO 2009 stage IB–IVB squamous cell carcinoma or adenocarcinoma undergoing CRT with curative intent	1341	CRT (weekly cisplatin + external beam RT) followed by MRI-based IGABT	Actuarial overall 5- year local control was 92% (95% CI 90% to 93%). Actuarial cumulative 5- year incidence of grade 3–5 morbidity was 6.8% (95% CI 5.4% to 8.6%) for genitourinary events, 8.5% (95% CI 6.9% to 10.6%) for gastrointestinal events, 5.7% (95% CI 4.3% to 7.6%) for vaginal events, and 3.2% (95% CI 2.2% to 4.5%) for fistulae	CRT and MRI-based IGABT result in effective and stable long-term local control across all stages of locally advanced cervical cancer, with a limited severe morbidity per organ
STARS ¹⁶²	RCT	Stage IB to IIA, with adverse pathological factors following radical hysterectomy.	1048	Arm1 (n=350): RT Arm 2 (n=345): CCRT with weekly cisplatin Arm 3 (n=353): SCRT with cisplatin+paclitaxel in a 21-day cycle, given two cycles before and two cycles after RT	SCRT was associated with a higher rate of 3-year disease-free survival than RT (90.0% vs 82.0%; HR 0.52; 95% CI 0.35 to 0.76) and CCRT (90.0% vs 85.0%; HR 0.65; 95% CI 0.44 to 0.96). Treatment with SCRT decreased cancer death risk compared with RT (5- year rate 92.0% vs 88.0%; HR 0.58; 95% CI 0.35 to 0.95) after adjustment for lymph node involvement. However, neither disease- free survival nor cancer death risk was different among patients treated with CCRT or RT	SCRT, rather than CCRT, resulted in a higher disease-free survival and lower risk of cancer death than RT among women with early-stage cervical cancer
Landoni et al ¹⁶³	RCT	Stage IB and IIA squamous cell carcinoma or adenocarcinoma	343	Arm 1 (n=172): radical surgery Arm 2 (n=171): RT	5- year overall survival and disease-free survival were identical in the surgery and RT groups (83% and 74%, respectively, for both groups). 86 women developed recurrent disease: 42 (25%) in the surgery group and 44 (26%) in the RT group	There is no treatment of choice for early-stage cervical carcinoma in terms of overall survival or disease-free survival. The combination of surgery and RT has the worst morbidity, especially urological complications. The optimum therapy for each patient should take account of clinical factors

Continued

Table 1 Continued

Study	Design	Setting	N	Treatment	Results	Conclusion
GOG 123 ¹⁶⁴	Phase III RCT	Stage IB bulky cervical tumor (>4 cm)	369	Arm 1 (n=186): RT followed by hysterectomy Arm 2 (n=183): RT + weekly cisplatin (CRT) followed by hysterectomy	RRs of disease progression and death in CRT compared with RT alone were 0.51 (95% CI 0.34 to 0.75) and 0.54 (95% CI 0.34 to 0.86), respectively. The rates of progression-free survival (p<0.001) and overall survival (p=0.008) were significantly higher in the CRT group at 4 years. Pathologically, no residual disease in the uterus at hysterectomy in 41% of RT group compared with 52% of CRT group	Adding weekly infusions of cisplatin to pelvic RT followed by hysterectomy significantly reduced the risk of disease recurrence and death in women with stage IB bulky cervical cancers. The value of hysterectomy in this setting is unclear
RTOG 9001 ¹⁰⁶	Phase III RCT	Stage IIB, III, IVA, and stage IB-IIA with tumor >5 cm and positive lymph nodes	386	Arm 1 (n=193): cisplatin + 5-FU weekly with radiotherapy (CRT) Arm 2 (n=193): RT (with extended para-aortic field)	Cumulative rates of 5-year overall survival were 73% among patients treated with CRT and 58% among patients treated with RT alone (p=0.004). Cumulative rates of 5-year disease-free survival were 67% among patients in the CRT group and 40% in the RT group (p<0.001). The rates of both distant metastases (p<0.001) and locoregional recurrences (p<0.001) were significantly higher among patients treated with RT alone	The addition of chemotherapy with cisplatin and 5-FU to treatment with external-beam and intracavitary RT significantly improved survival among women with locally advanced cervical cancer
GOG 165 ¹⁶⁵	Phase III, open-label RCT	Stage IIB, IIIB, and IVA cervical cancer with clinically negative aortic nodes	316	Arm 1 (n=159): weekly cisplatin + RT (cisplatin/RT) Arm 2 (n=157): 5 - FU + RT (5-FU/RT)	Closed prematurely when a planned interim futility analysis indicated that 5-FU/RT had a higher treatment failure rate (35% higher) and most likely would not result in an improvement in progression-free survival compared with weekly cisplatin/RT. The 5-FU/RT arm showed a higher risk of treatment failure (unadjusted RR 1.29) and a higher mortality rate (unadjusted RR 1.37). There was no difference in pelvic treatment failure between regimens, but there was an increase in the failure rate at distant sites in the 5-FU arm	5-FU showed no improvement over weekly cisplatin

Continued

Table 1 Continued

Study	Design	Setting	N	Treatment	Results	Conclusion
EORTC 55994 ¹²⁵	Phase III RCT	FIGO 2009 stage IB2–IIB cervical cancer	626	Arm 1 (n=314): NACT followed by surgery (NACT-S) Arm 2 (n=312): CCRT	The 5-year overall survival was not significantly different between NACT-S (72%; 95% CI 66% to 77%) and CCRT (76%; 95% CI 70% to 80%). Short-term AEs ≥grade 3 occurred more frequently with NACT-S (41% vs 23%) and long-term AEs ≥grade 3 more often with CCRT (21% vs 15%). Additional RT after completed NACT-S was given to 107 patients (48%), and additional surgery to 20 patients (8%) after completed CCRT	NACT-S is not superior to CCRT. Additional RT was used for a substantial proportion of patients with stage IB2–IIB cervical cancer who were assigned to NACT-S. CCRT remains the standard of care in this setting
Gupta et al ¹²⁴	Phase III, single-center RCT	FIGO 1994 stage IB2, IIA, or IIB cervical cancer.	633	Arm 1 (n=316): NACT-S Arm 2 (n=317): RT	The 5-year disease-free survival rate was 69.3% in the NACT-S cohort and 76.7% in the CCRT group (HR, 1.38; 95% CI 1.02 to 1.87; p=0.038)	Cisplatin-based CCRT showed superior disease-free survival compared with NACT-S for patients with locally advanced cervical cancer
Uterus-11 ¹¹⁴	RCT	FIGO 2009 stage IIB–IVA squamous cell carcinoma, adenocarcinoma, adenosquamous carcinoma	255	Arm 1 (n=130): surgical staging Arm 2 (n=125): clinical staging	Laparoscopic staging led to upstaging in 39 of 120 (33%) patients. However, after a median follow-up of 90 months (range 1–123) in both arms, there was no difference in disease-free survival between the groups (p=0.084)	No difference in disease-free survival between surgical and clinical staging in patients with locally advanced cervical cancer
CALLA ¹²⁸	Phase III, double blind RCT	FIGO 2009 stage IB2–IIB lymph node positive, or stage ≥III any lymph node status; squamous cell carcinoma, adenocarcinoma, adenosquamous carcinoma	770	Arm 1 (n=385): CRT + placebo Arm 2 (n=385): CRT + durvalumab up to 24 cycles	12-month progression-free survival was 76.0% (95% CI 71.3% to 80.0%) with durvalumab and 73.3% (95% CI 68.4% to 77.5%) with placebo. Serious AEs occurred for 106 (28%) patients who received durvalumab and 89 (23%) patients who received placebo	Durvalumab concurrent with CRT was well tolerated in participants with locally advanced cervical cancer; however, it did not significantly improve progression-free survival in a biomarker unselected, all comers population
ENGOT-cx11/ GOG-3047/ KEYNOTE-A18 ¹²	Phase III, double-blind, placebo-controlled RCT	High-risk locally advanced cervical cancer: FIGO 2014 stage IB2–IIB with node-positive disease or stage III–IVA regardless of nodal status; squamous cell carcinoma, adenocarcinoma, or adenosquamous carcinoma	1060	Arm 1 (n=529): pembrolizumab-CRT Arm 2 (n=531): placebo-CRT	Median 24-month progression-free survival was 68% for pembrolizumab-CRT and 57% for placebo-CRT. The HR for disease progression or death was 0.70 (95% CI 0.55 to 0.89; p=0.0020). At 24 months, overall survival was 87% and 81%, respectively (information fraction 42.9%). The HR for death was 0.73 (95% CI 0.49 to 1.07), not reaching statistical significance	The combination of pembrolizumab and CRT demonstrated statistically significant enhancement in progression-free survival among individuals diagnosed with high-risk, locally advanced cervical cancer

Continued

Table 1 Continued

Study	Design	Setting	N	Treatment	Results	Conclusion
OUTBACK ¹³¹	Phase III, open-label RCT	FIGO 2009 stage IB1 disease with nodal involvement, or stage IB2, II, IIIB, or IVA squamous cell carcinoma, adenocarcinoma, or adenosquamous carcinoma	926	Arm 1 (n=461): CRT Arm 2 (n=465): CRT + adjuvant chemotherapy (four cycles of carboplatin/paclitaxel after completion of CRT)	5-year overall survival was 72% (95% CI 67% to 76%) in the adjuvant chemotherapy group (105 deaths) and 71% (95% CI 66% to 75%) in the CRT only group (116 deaths; difference 1% (95% CI -6 to 7%); HR 0.90 (95% CI 0.70 to 1.17; p=0.81). Serious AEs occurred in 107 (30%) in the adjuvant chemotherapy group vs 98 (22%) in the CRT only group, most commonly due to infectious complications	Adjuvant carboplatin/paclitaxel chemotherapy given after standard cisplatin-based CRT for unselected locally advanced cervical cancer increased short-term toxicity and did not improve overall survival; therefore, it should not be given in this setting
INTERLACE ¹³	Phase III RCT	FIGO 2009 stage IB1 node positive, IB2, II, IIIB, IVA squamous cell carcinoma, adenocarcinoma, adenosquamous carcinoma	500	Arm 1 (n=250): CRT Arm 2 (n=250): weekly carboplatin (AUC 2)/paclitaxel (80 mg/m ²) for 6 weeks followed by CRT	5-year progression-free survival rate was 73% with induction chemotherapy and CRT and 64% with CRT alone (HR 0.65; 95% CI 0.46 to 0.91; p=0.013). The corresponding 5-year overall survival rates are 80% and 72% (HR 0.61; 95% CI 0.40 to 0.91; p=0.04)	Induction chemotherapy followed by CRT significantly improves progression-free and overall survival in locally advanced cervical cancer
Metastatic/recurrent disease						
First-line systemic therapy						
GOG 169 ¹⁶⁶	Phase III RCT	Metastatic, recurrent or persistent squamous cell carcinoma of the cervix with measurable disease	264	Arm 1 (n=134): cisplatin 50 mg/m ² Arm 2 (n=130): cisplatin 50 mg/m ² + paclitaxel 135 mg/m ² (C+P) every 3 weeks for six cycles	The majority of all patients had prior RT (cisplatin 92%; C+P 91%). ORR was 19% (6% complete plus 13% partial) of patients receiving cisplatin vs 36% (15% complete plus 21% partial) receiving C+P (p=0.002). The median progression-free survival was 2.8 months and 4.8 months, respectively for cisplatin vs C+P (p<0.001). There was no difference in median survival (8.8 months vs 9.7 months). Grade 3–4 anemia and neutropenia were more common in the combination arm	C+P is superior to cisplatin alone with respect to ORR and progression-free survival
GOG 204 ¹⁶⁷	Multicenter, phase III RCT	Metastatic, recurrent, or persistent squamous cell carcinoma, adenocarcinoma, or adenosquamous carcinoma with measurable disease	434	Arm 1 (n=103): paclitaxel 135 mg/m ² over 24 hours plus cisplatin 50 mg/m ² day 2 every 3 weeks (PC, reference arm) Arm 2 (n=108): vinorelbine 30 mg/m ² days 1 and 8 plus cisplatin 50 mg/m ² day 1 every 3 weeks (VC) Arm 3 (n=112): gemcitabine 1000 mg/m ² day 1 and 8 plus cisplatin 50 mg/m ² day 1 every 3 weeks (GC) Arm 4 (n=111): topotecan 0.75 mg/m ² days 1, 2, and 3 plus cisplatin 50 mg/m ² day 1 every 3 weeks (TC)	The experimental-to-PC HR of death were 1.15 (95% CI 0.79 to 1.67) for VC, 1.32 (95% CI 0.91 to 1.92) for GC, and 1.26 (95% CI 0.86 to 1.82) for TC. The HRs for progression-free survival were 1.36 (95% CI 0.97 to 1.90) for VC, 1.39 (95% CI 0.99 to 1.96) for GC, and 1.27 (95% CI 0.90 to 1.78) for TC. Response rates for PC, VC, GC, and TC were 29.1%, 25.9%, 22.3%, and 23.4%, respectively	VC, GC, and TC are not superior to PC in terms of overall survival. However, the trend in response rates, progression-free survival, and overall survival favors PC

Continued

Table 1 Continued

Study	Design	Setting	N	Treatment	Results	Conclusion
JCOG0505 ¹⁶⁸	Multicenter, open-label, phase III RCT	Metastatic, recurrent or persistent squamous cell carcinoma, adenocarcinoma, or adenosquamous carcinoma. Allowed up to one prior platinum-based regimen	253	Arm 1 (n=127): cisplatin/paclitaxel (TP) Arm 2 (n=126): carboplatin/paclitaxel (TC)	The HR of overall survival was 0.994 (90% CI 0.79 to 1.25; non-inferiority p=0.032 by stratified Cox regression). Median overall survival was 18.3 months with TP vs 17.5 months with TC. Among patients who had not received prior cisplatin, overall survival was shorter with TC (13.0 vs 23.2 months; HR 1.571; 95% CI 1.06 to 2.32)	TC was non-inferior to TP and should be a standard treatment option for metastatic or recurrent cervical cancer. However, cisplatin remains a key drug for patients who have not received platinum agents
GOG 240 ¹³²	Multicenter, phase III RCT	Metastatic, recurrent, or persistent with measurable disease	452	Arm 1 (n=114): cisplatin/paclitaxel Arm 2 (n=115): cisplatin/paclitaxel + bevacizumab Arm 3 (n=111): topotecan/paclitaxel Arm 4 (n=112): topotecan / paclitaxel + bevacizumab	Topotecan-paclitaxel was not superior to cisplatin-paclitaxel (HR for death 1.20). With the data for the two chemotherapy regimens combined, the addition of bevacizumab to chemotherapy was associated with increased overall survival (17.0 vs 13.3 months; HR 0.71; 98% CI 0.54 to 0.95; p=0.004 in a one-sided test) and higher response rates (48% vs 36%; p=0.008)	The addition of bevacizumab to combined chemotherapy in patients with recurrent, persistent, or metastatic cervical cancer was associated with an improvement of 3.7 months in median overall survival
KEYNOTE-826 ¹³³	Multicenter, double-blind, phase III RCT	Metastatic, recurrent or persistent squamous cell carcinoma, adenocarcinoma, or adenosquamous carcinoma	617	Arm 1 (n=309): paclitaxel + cisplatin (or carboplatin) with or without bevacizumab + placebo Arm 2 (n=308): paclitaxel + cisplatin (or carboplatin) with or without bevacizumab + pembrolizumab 200 mg every 3 weeks	In patients with a PD-L1 combined positive score $\geq$ 1, median progression-free survival was 10.4 months in the pembrolizumab group vs 8.2 months in the placebo group (HR 0.62; 95% CI 0.50 to 0.77; p<0.001). In the intention-to-treat population, median progression-free survival was 10.4 months vs 8.2 months, respectively (HR 0.65; 95% CI 0.53 to 0.79; p<0.001). In patients with a PD-L1 combined positive score $\geq$ 1, overall survival at 24 months was 53.0% in the pembrolizumab group vs 41.7% in the placebo group (HR 0.64; 95% CI 0.50 to 0.81; p<0.001). In the intention-to-treat population, overall survival at 24 months was 50.4% vs 40.4% (HR 0.67; 95% CI 0.54 to 0.84; p<0.001)	Progression-free and overall survival were significantly longer with pembrolizumab than with placebo among patients with persistent, recurrent, or metastatic cervical cancer who were also receiving chemotherapy with or without bevacizumab
BEATcc/ ENGOT-cx10 ¹³⁵	International, open-label, phase III RCT	Metastatic, persistent, or recurrent squamous cell carcinoma, adenocarcinoma, adenosquamous carcinoma	410	Arm 1 (n=204): cisplatin (or carboplatin) + paclitaxel + bevacizumab every 3 weeks Arm 2 (n=206): cisplatin (or carboplatin) + paclitaxel + bevacizumab + atezolizumab 1200 mg every 3 weeks	Median progression-free survival was 13.7 months (95% CI 12.3 to 16.6) with atezolizumab and 10.4 months (95% CI 9.7 to 11.7) with standard therapy (HR 0.62; 95% CI 0.49 to 0.78; p<0.0001). At the interim overall survival analysis, median overall survival was 32.1 months (95% CI 25.3 to 36.8) vs 22.8 months (95% CI 20.3 to 28.0), respectively (HR 0.68; 95% CI 0.52 to 0.88; p=0.0046)	Adding atezolizumab to a standard bevacizumab plus platinum regimen for metastatic, persistent, or recurrent cervical cancer significantly improves progression-free and overall survival

Continued

Table 1 Continued

Study	Design	Setting	N	Treatment	Results	Conclusion
Subsequent line of systemic therapy						
EMPOWER-Cervical 1/GOG-3016/ENGOT-cx9 ¹³⁷	Multicenter, open-label, phase III RCT	Recurrent or metastatic cervical cancer progressing after platinum-containing chemotherapy. Patients must have received taxane and bevacizumab. It was required to have measurable disease	608	Arm 1 (n=304): investigator's choice single agent chemotherapy Arm 2 (n=304): cemiplimab (350 mg every 3 weeks)	Median overall survival was longer in the cemiplimab group than in the chemotherapy group (12.0 vs 8.5 months; HR for death 0.69; 95% CI 0.56 to 0.84; two-sided p<0.001). In the overall population, ORR occurred in 16.4% (95% CI 12.5 to 21.1) of the patients in the cemiplimab group compared with 6.3% (95% CI 3.8 to 9.6) in the chemotherapy group. ORR occurred in 18% (95% CI 11 to 28) of the cemiplimab-treated patients with PD-L1 expression ≥1% and in 11% (95% CI 4 to 25) of those with PD-L1 expression <1%	Survival was significantly longer with cemiplimab than with single-agent chemotherapy among patients with recurrent cervical cancer after first-line platinum-containing chemotherapy
KEYNOTE-158 ¹³⁸	Phase II, open-label basket trial investigating the antitumor activity and safety of pembrolizumab in multiple cancer types	Previously treated advanced cervical cancer	98	Pembrolizumab 200 mg every 3 weeks for 2 years or until progression, intolerable toxicity, or physician or patient decision	82 patients (83.7%) had PD-L1-positive tumors (combined positive score ≥1). ORR was 12.2%, with 3 complete and 9 partial responses. All 12 responses were in patients with PD-L1-positive tumors, for an ORR of 14.6%. Treatment-related grade 3 to 4 AEs occurred in 12.2% of patients	Pembrolizumab monotherapy demonstrated durable antitumor activity and manageable safety in patients with advanced PD-L1-positive cervical cancer who experienced progression during or after chemotherapy
GOG 324/ENGOT-cx8/innovaTV 205 ¹⁴⁴	Phase Ib/2, open-label, multicenter study	Recurrent or metastatic cervical cancer	142	Dose-escalation arms (n=41): tisotumab vedotin-tftv (TV) 2 mg/kg plus bevacizumab 15 mg/kg every 3 weeks (arm A), pembrolizumab 200 mg every 3 weeks (arm B), or carboplatin AUC 5 every 3 weeks (arm C) Arm D (n=33): 1L TV + carboplatin Arm E (n=33): 1L TV + pembrolizumab Arm F (n=35): 2L/3L TV + pembrolizumab	In dose escalation (n=41), no drug-limiting toxicities were observed. In dose expansion, the ORR was 54.5% (n/N, 18/33; 95% CI 36.4% to 71.9%) with first-line TV+carboplatin, 40.6% (n/N, 13/32; 95% CI 23.7% to 59.4%) with first-line TV+pembrolizumab, and 35.3% (n/N, 12/34; 95% CI 19.7% to 53.5%) with second-line/third-line TV+pembrolizumab. The median DOR was 8.6 months, not reached, and 14.1 months in arms 1, 2, and 3, respectively. Grade ≥3 AEs (≥15%) were anemia, diarrhea, nausea, and thrombocytopenia in arm 1 and anemia in arm 3 (none ≥15%, arm 2)	TV in combination with bevacizumab, carboplatin, or pembrolizumab demonstrated manageable safety and encouraging anti-tumor activity in treatment-naïve and previously treated recurrent or metastatic cervical cancer
GOG 3023/ENGOT-cx6/innovaTV 204 ¹⁴²	Multicenter, open-label, single-arm, phase II study	Recurrent or metastatic squamous cell carcinoma, adenocarcinoma, or adenosquamous carcinoma previously treated with platinum doublet chemotherapy with bevacizumab (two or fewer systemic regimens), with measurable disease	102	TV 2 mg/kg every 3 weeks until disease progression or unacceptable toxicity	The confirmed ORR was 24% (95% CI 16% to 33%), with 7 (7%) complete responses and 17 (17%) partial responses. Serious treatment-related AEs occurred in 13 (13%) patients, the most common of which included peripheral sensorimotor neuropathy (2%) and fever (2%)	TV showed clinically meaningful and durable anti-tumour activity with a manageable and tolerable safety profile in women with previously treated recurrent or metastatic cervical cancer

Continued

Table 1 Continued

Study	Design	Setting	N	Treatment	Results	Conclusion
GOG 3057/ ENGOT-cx12/ innoVA TV 301 ¹⁴³	International, open-label, phase III RCT	Recurrent or metastatic disease following platinum doublet therapy with or without bevacizumab and with or without anti-PD-L1 therapy. Measurable disease required	502	Arm 1 (n=253): TV Arm 2 (n=249): investigator's choice single agent chemotherapy	TV arm had a 30% reduction in risk of death vs chemotherapy (HR 0.70; 95% CI 0.54 to 0.89; p=0.004), with significantly longer median overall survival (11.5 vs 9.5 months). Progression-free survival was superior in tTV vs chemotherapy arm (median 4.2 vs 2.9 months; HR 0.67; 95% CI 0.54 to 0.82; p<0.001). Confirmed ORR was 17.8% and 5.2% in the TV and chemotherapy arms, respectively (OR 4.0; 95% CI 2.1 to 7.6; p<0.001)	TV showed a statistically significant and clinically meaningful improvement in overall survival, progression-free survival, and ORR compared with chemotherapy, with a manageable and tolerable safety profile in previously treated patients with recurrent or metastatic cervical cancer
DESTINY- PanTumor02 ¹⁴⁵	Open-label, phase II study	HER2-expressing (IHC 2+/3+) locally advanced or metastatic disease after ≥1 systemic treatment or without alternative treatments	267	Trastuzumab deruxtecan 5.4 mg/kg every 3 weeks	In all patients, the ORR was 37.1% (n=99; 95% CI 31.3% to 43.2%), with responses in all cohorts. Median DOR was 11.3 months (95% CI 9.6 to 17.8); the median progression-free survival was 6.9 months (95% CI 5.6 to 8.0); the median overall survival was 13.4 months (95% CI 11.9 to 15.5) In patients with central HER2 IHC 3+ expression (n=75), ORR was 61.3% (95% CI 49.4% to 72.4%). Median DOR was 22.1 months (95% CI 9.6 to not reached); the median progression-free survival was 11.9 months (95% CI 8.2 to 13.0); the median overall survival was 21.1 months (95% CI 15.3 to 29.6)	HER2-expressing solid tumors demonstrate durable clinical benefit and meaningful survival outcomes with trastuzumab deruxtecan with the greatest benefit observed in the IHC 3+ population

AE, adverse event; AUC, area under the curve; CCRT, concurrent chemoradiotherapy; CRT, chemoradiotherapy; dMMR, mismatch repair deficient; DOR, duration of response; FIGO, International Federation of Gynecology and Obstetrics; 5-FU, 5-fluorouracil; IGABT, imaging guided adaptive brachytherapy; IHC, immunohistochemistry; LACC, Laparoscopic Approach to Cervical Cancer; LEEP, loop electrosurgical excision procedure; LND, lymphadenectomy; LVSI, lymphovascular space invasion; MIS, minimally invasive surgery; MRI, magnetic resonance imaging; MSI-H, microsatellite instability-high; NACT, neoadjuvant chemotherapy; NFT, no further treatment; ORR, objective response rate; PD-L1, programmed cell death ligand 1; RCT, randomized clinical trial; RR, relative risk; RT, radiotherapy; SCRT, sequential chemoradiotherapy; SLN, sentinel lymph node; TV, tisotumab vedotin-ttfv.

have implemented primary cervical cancer screening with HPV testing,^{33 34} coupled with triage testing (reflex cytology, HPV genotyping, p16 immunostaining, and DNA methylation) to decrease the follow-up of transient infections.²⁰ However, there are many barriers to universal adoption of HPV-based screening, which has made primary screening coverage challenging, particularly in low-income and middle-income countries.³⁵ Approximately 35% of countries have implemented HPV testing-based programs and at least 21 countries are in transition from cytology to HPV testing.³⁰ The utilization of HPV testing in screening has the potential to improve the identification of adenocarcinoma.³⁶ In resource-limited countries, screening still relies on direct visual inspection using acetic acid. Interestingly, the FDA recently approved the first self-test collection kit for HPV, a move that could expand access to screening and reduce barriers.³⁷

Clinical Presentation

The most frequent clinical presentation of cervical cancer is vaginal discharge/bleeding, postcoital bleeding, or intermittent spotting.

Early-stage cervical cancer is often asymptomatic or associated with vague and nonspecific symptoms that are typically identified during screening programs.^{38 39} Cervical cancer spreads locally into the vagina, parametrium, uterine corpus, and nearby organs, as well as through lymphatic channels to regional and distant lymph nodes. Metastasis can also occur via hematogenous routes to the lungs, liver, and skeleton.⁴⁰ Advanced stages may present with a wide range of signs and symptoms, including hematuria, malodorous vaginal discharge, dyspareunia, and pelvic or back pain. Pelvic sidewall involvement may lead to a triad of lower limb edema, flank pain, and sciatica. Rarely, vesicovaginal or rectovaginal fistulas are encountered during initial presentations.⁴¹

Diagnosis

Workup for patients with suspicious symptoms includes history, pelvic examination, cervical biopsy/conization biopsy, laboratory tests, and imaging. If a clear lesion is visible or the cervix appears irregular or enlarged during speculum examination, direct biopsy is




Figure 1 Pathophysiology of the HPV infection. HPV, human papillomavirus; p53, tumor protein P53; pRB, retinoblastoma protein.

sufficient for diagnosis. Colposcopy with biopsy or conization are performed in cases where a lesion is not macroscopically visible. Cervical conization should not be performed in the setting of a grossly visible lesion.

Staging

Patients with cervical cancer should be staged according to the TNM classification and the revised FIGO 2018 staging system.⁴² Imaging techniques that may be considered for clinical staging include pelvic ultrasound (performed by a trained sonographer), CT chest-abdomen-pelvis, abdominal/pelvic MRI, 18-fluorodeoxyglucose (¹⁸FDG) positron emission tomography (PET)/CT, and/or chest radiograph, as indicated and depending on resource availability. Cystoscopy and proctoscopy are recommended only when bladder or rectal extension is suspected ($\geq$ FIGO 2018 stage IB3).^{9–11}

CT has high sensitivity in detecting lymph node and distant metastases and ¹⁸FDG-PET/CT may add value if CT results are indeterminate.⁴³ MRI has better sensitivity than CT for assessing local tumor extent⁴⁴; pelvic ultrasound can be a valid alternative if performed by highly experienced sonographers.⁴⁵ All CT/MRI should be performed with contrast unless contraindicated.^{46–48} Ongoing research is exploring innovative imaging techniques such as diffusion-weighted imaging MRI,^{49–50} dynamic contrast-enhanced MRI,^{51–52} contrast-enhanced ultrasound, and hybrid imaging with PET-MRI.^{53–54} With the progressive integration of artificial intelligence, these techniques may have the potential to assist clinicians in diagnostic and prognostic assessment at diagnosis, refining risk stratification, guiding decision making and personalized treatments.⁵⁵

Treatment

Treatment planning should be made in specialized referral centers within a multidisciplinary setting and be tailored to each patient. Patients should be carefully selected and counseled on the standard

treatment and potential alternatives, aligning expectations. Treatment depends on several factors: (a) disease-related factors (stage at diagnosis, histology, tumor size, parametrial invasion, lymph node involvement); (b) patient-related factors (age, performance status, desire for fertility, medical comorbidities); (c) resources available at the treating facility (multidisciplinary team, experience, financial resources). Treatment options may vary across countries depending on national guidelines, resources, and reimbursement policies.^{10–56} Squamous cell carcinomas and usual-type (HPV-associated) adenocarcinomas are typically treated in a similar manner.^{57–59}

Early-stage Cervical Cancer

The standard primary treatment of early-stage (FIGO 2018 stages IA–IB2, IIA1) cervical cancer is surgery followed by observation or adjuvant therapy based on histopathological risk factors on final pathology (Figure 2). Conservative surgical approaches should be considered after careful patient selection. Fertility-sparing approaches should be reserved for patients with a desire for childbearing and low-risk disease (Figure 3). Conservative and fertility-sparing approaches should be considered only within a multidisciplinary setting with expert gynecological oncologists, pathologists, and radiologists to identify appropriate candidates. Definitive radiotherapy ($\pm$ concurrent platinum-containing chemotherapy) may be a valid alternative to surgery in patients deemed to have inoperable cancer due to poor performance status.^{9–11}

Surgery

Three major concepts in the treatment of early-stage cervical cancer have been examined and have evolved in recent years aiming to optimize oncologic outcomes and tailor less radical surgery to specific patients: (1) surgical approach; (2) criteria




Figure 2 Primary non-fertility-sparing treatment algorithm for early-stage cervical cancer. All staging is based on updated FIGO 2018 staging. Treatment planning should be multidisciplinary and individualized. Treatment options may vary across countries depending on national guidelines and socioeconomic factors. *Negative (at least 1 mm) for invasive disease or histologic high-grade squamous intraepithelial lesion (HSIL). If positive margins, repeat cone biopsy or perform simple hysterectomy. Cold-knife conization is preferable to LEEP or laser conization. Endocervical curettage should be performed as clinically indicated. †Ovarian preservation may be considered in premenopausal women (<45 years of age) with SCC or HPV-associated ADC/ADS. *Risk factors may not be limited to the traditional Sedlis criteria. ADC, adenocarcinoma; ADS, adenosquamous carcinoma; BRT, brachytherapy; CCRT, concurrent chemoradiotherapy; CHT, chemotherapy; EBRT, external beam radiotherapy; FIGO, International Federation of Gynecology and Obstetrics; HPV, human papillomavirus; LEEP, loop electrosurgical excision procedure; LVSI, lymphovascular space invasion; PALND, para-aortic lymph node dissection; PLND, pelvic lymph node dissection; SCC, squamous cell carcinoma; SLN, sentinel lymph node.

for conservative surgical approaches; and (3) the role of pelvic SLN biopsy (Figure 2).

Surgical Approach

In 2018, a phase III, multicenter, randomized trial (Laparoscopic Approach to Cervical Cancer—LACC) reported higher recurrence rate and worse overall survival with minimally invasive surgery for early-stage cervical cancer, establishing the open approach as the standard of care (Table 1).⁶⁰ The rate of disease-free survival at 4.5 years was 86.0% with minimally invasive surgery and 96.5% with open surgery, a difference of −10.6 percentage points (95% CI −16.4 to −4.7). The final overall survival analysis has recently been published and showed that the overall survival rate at 4.5 years was 90.6% versus 96.2% for the minimally invasive and open surgery groups, respectively (HR for death of any cause 2.71; 95% CI 1.32 to 5.59; $p=0.007$).⁶¹

Potential theories for these findings have been proposed, including: (1) the impact of CO₂ pneumoperitoneum; (2) the lack

of use of peritoneal protective maneuvers to avoid tumor exposure; and (3) the use of a uterine manipulator.^{62 63} To further clarify this topic, two ongoing randomized controlled trials, RACC⁶⁴ and ROCC (NCT04831580), are investigating the oncological safety of a robotic versus an open approach. These studies do not allow the use of an intrauterine manipulator and require documentation of standardized surgical techniques for tumor containment.

Conservative Approaches

The type of hysterectomy and extent of parametrectomy (type A-C2 according to the 2017 revised Querleu-Morrow classification) have traditionally been based on preoperative prognostic risk factors (tumor size, maximum stromal invasion, and lymphovascular space invasion), which are used to categorize tumors at low, intermediate, and high risk of recurrence.⁶⁵ Recent advances in the surgical management of cervical cancer have identified selected patients who may benefit from less radical surgery for early-stage cervical cancer.




Figure 3 Primary fertility-sparing management of early-stage cervical cancer. All staging is based on updated FIGO 2018 staging. Treatment planning should be multidisciplinary and individualized. Treatment options may vary across countries depending on national guidelines and socioeconomic factors. *Negative (at least 1 mm) for invasive disease or histologic high-grade squamous intraepithelial lesion (HSIL). If positive margins, repeat cone biopsy or perform simple trachelectomy. Loop electrosurgical excision procedure (LEEP) or laser conization is preferable to cold-knife conization in women wanting to preserve fertility. Endocervical curettage should be performed as clinically indicated. †Radical trachelectomy should be considered if conservative surgery criteria are not met (stromal invasion ≥10 mm on LEEP/cone or ≥50% on MRI). ADC, adenocarcinoma; ADS, adenosquamous carcinoma; FIGO, International Federation of Gynecology and Obstetrics; HPV, human papillomavirus; LVSI, lymphovascular space invasion; NACT, neoadjuvant chemotherapy; PALND, para-aortic lymph node dissection; PLND, pelvic lymph node dissection; SCC, squamous cell carcinoma; SLN, sentinel lymph node; US, ultrasound.

The ConCerv trial, a prospective, single-arm, multicenter study, examined the feasibility of conservative surgery to reduce surgical complications and morbidity in low-risk, early-stage cervical cancer (Table 1).⁶⁶ Eligibility criteria included: (1) FIGO 2009 stage IA2–IB1 cervical cancer; (2) squamous cell (any grade) or adenocarcinoma (grade 1 or 2 only) histology; (3) tumor size <2 cm; (4) no lymphovascular space invasion; (5) depth of invasion <10 mm; (6) negative imaging for metastatic disease; and (7) negative conization margins. Cervical conization was performed to determine eligibility, with one repeat cone permitted. Patients who met these inclusion criteria underwent a second surgery with pelvic lymph node assessment (if fertility preservation was desired) or a simple hysterectomy with lymph node assessment. The 2-year cumulative recurrence rate in the study population was 3.5% (95% CI 0.9% to 9.0%).⁶⁶

The randomized phase III SHAPE trial recently evaluated simple hysterectomy compared with radical hysterectomy in a non-inferiority design (Table 1).⁶⁷ This study included patients with squamous cell carcinoma, adenocarcinoma, or adenosquamous carcinoma of any histologic grade, with or without lymphovascular space involvement, FIGO 2009 stage IA2 or IB1 with tumors measuring no more than 2 cm, and with limited depth of cervical stromal invasion (invasion of <50% of stromal tissue or to a depth of <10 mm, or both). Notably, 80.2% of patients had undergone diagnostic loop electrosurgical excision procedure or conization before surgery. Findings demonstrated simple hysterectomy was not inferior to radical hysterectomy with respect to the 3-year incidence of

pelvic recurrence (absolute difference of 0.35 percentage points; 90% CI –1.62 to 2.32).⁶⁷

Results from the GOG 278 trial were presented at the 2024 Society of Gynecologic Oncology (SGO) annual meeting (Table 1).⁶⁸ This phase I/II multicenter prospective trial aimed to evaluate the impact of non-radical surgical treatment, either simple hysterectomy or cone biopsy (for fertility preservation) and pelvic lymphadenectomy, on functional outcomes of bladder, bowel, and sexual function and on the incidence and severity of lymphedema in patients with FIGO 2009 stage IA1 (lymphovascular space invasion present) and IA2–IB1 (≤ 2 cm) cervical cancer with depth of stromal invasion ≤10 mm and no evidence of metastasis on radiologic staging. Of the 201 patients eligible for survival analysis, only three experienced a recurrence, suggesting that non-radical surgery for early-stage cancer appears safe and is associated with low perioperative morbidity.

Sentinel Lymph Node Biopsy

While there is no doubt that lymph node assessment is a crucial part of surgical staging in cervical cancer, there is no consensus on the appropriate modality: pelvic (±para-aortic) systematic lymphadenectomy and/or SLN biopsy. Systematic lymphadenectomy has traditionally been considered the standard of care. However, SLN biopsy without systematic pelvic lymphadenectomy has recently emerged as a reasonable strategy in early-stage cervical tumors <4 cm that is associated with less morbidity than pelvic lymphadenectomy. The SENTIX/ENGOT-cx2 trial was

a single-arm, multicenter, prospective cohort study to evaluate whether SLN biopsy is non-inferior to systematic pelvic lymphadenectomy in terms of recurrence rate in early-stage cervical cancer (Table 1).⁶⁹ Eligible patients had FIGO 2009 stage IA1 with lymphovascular space invasion, IA2, and IB1 (<2 cm for fertility sparing) with negative lymph nodes on preoperative imaging. In the SENTIX trial, the recurrence rate after SLN biopsy was 6%, which is non-inferior to the reference recurrence rate of 7% after pelvic lymphadenectomy in historical cohorts. SLN biopsy showed a bilateral detection rate of 92.3% and was able to detect lymph node involvement in more than 56% of cases by frozen section and over 90% after ultrastaging. These recent findings are consistent with meta-analyses of pooled data from SLN mapping studies reporting SLN detection rates of 89–92%, sensitivity of ~90%, and negative predictive value >98%.^{70–72} All suspicious or grossly enlarged nodes should be removed at the time of surgical staging regardless of SLN mapping. An ancillary analysis of two prospective multicentric trials (SENTICOL I and II) on SLN biopsy for cervical cancer also suggested that omitting systematic pelvic lymphadenectomy for patients with bilateral negative SLN was not associated with an increased risk of recurrence.⁷³ The results of the ongoing SENTICOL III validation study of SLN biopsy in early cervical cancer are eagerly awaited.⁷⁴

Another area of controversy is whether to routinely perform SLN frozen section evaluation. While the European Society of Gynecological Oncology (ESGO) 2023 guidelines¹¹ recommend intraoperative assessment of lymph node status by frozen section, other guidelines such as the National Comprehensive Cancer Network (NCCN),⁹ the Japanese Society of Gynecologic Oncology,⁷⁵ and the Korean Society of Gynecologic Oncology (KSGO)⁷⁶ do not comment on intraoperative lymph node assessment. A recent meta-analysis showed the pooled sensitivity of intraoperative frozen section analysis was 65% for macrometastases, micrometastases, and isolated tumor cells, and 72% when patients with isolated tumor cells were excluded.⁷⁷ The rationale for SLN frozen section is that it may prevent unnecessary surgery in patients with positive SLN on frozen section who should be referred to definitive concurrent chemoradiation, avoiding the increased toxicity associated with combined modalities.^{78,79}

If no SLNs are detected on either side, systematic dissection should be performed on the ipsilateral no-mapping side. All sentinel nodes should be processed with ultrastaging. Para-aortic lymphadenectomy (at least to the inferior mesenteric artery) may be considered in patients with FIGO 2018 stage $\geq$ IB1, even though there is limited and controversial evidence to suggest either a therapeutic benefit or a diagnostic role due to the low risk of isolated para-aortic metastasis.⁸⁰

Fertility-sparing Approaches

Cervical cancer is frequently diagnosed in women of reproductive age, with one third of patients younger than 45 years, and 40% nulliparous or with only one child at diagnosis.⁸¹ Thus, there is a growing need for fertility-sparing strategies for women with cervical cancer who have not yet completed childbearing. Fertility-sparing approaches may be considered in highly selected patients who have been thoroughly counseled regarding oncologic and obstetric risks (Figure 3). Consultation at a center with experience in oncofertility and discussion of each case in a multidisciplinary

team are essential before offering fertility-sparing treatment. The specific goal of fertility-sparing surgery is complete resection of the tumor with adequate free margins and preservation of the upper cervix.

Fertility-sparing surgical options include cervical conization, simple trachelectomy, vaginal radical trachelectomy, and radical abdominal trachelectomy. Neoadjuvant chemotherapy followed by conservative surgery is considered an alternative approach, although without any prospective evidence to confirm its oncologic safety. Negative lymph node status is required for fertility-sparing approaches and preoperative imaging has a crucial role to rule out nodal involvement. Surgical lymph node staging follows the same principles of non-fertility-sparing management and can be performed through a minimally invasive approach. Surgical lymph node staging can be performed using different approaches based on national guidelines and availability of resources: (1) SLN biopsy (with or without frozen section) and fertility-sparing procedure in the same surgery; (2) systematic pelvic lymphadenectomy and fertility-sparing procedure in the same surgery; (3) two-step approach first performing lymph node assessment (either SLN or systematic pelvic lymphadenectomy) and then fertility-sparing surgery only in patients with negative lymph nodes on definitive pathology.^{9–11} Intraoperative placement of permanent cerclage should be performed after trachelectomy. Routine hysterectomy after completion of childbearing is not mandatory.

Vaginal radical trachelectomy has traditionally been the most widely used fertility-sparing procedure since its development in the early 1990s,⁸² with recurrence, pregnancy, and live birth rates for tumors <2 cm of 4.7%, 58.7%, and 71%, respectively.⁸³ Patients with favorable tumor features (tumor size $\leq$ 2 cm, absence of lymphovascular space invasion, depth of invasion <10 mm, and negative lymph nodes) have a risk of parametrial involvement <1%^{84,85} and can be excellent candidates for more conservative procedures (conization or simple trachelectomy).^{86,87} The ConCerv⁶⁶ and GOG 278 trials⁶⁸ reported on fertility outcomes of patients with cervical cancer treated with conization and pelvic node assessment and suggested that this conservative strategy is safe and can lead to successful fertility outcomes in selected patients with low-risk early-stage cervical cancer (Table 1). In the ConCerv trial, 14 pregnancies have been reported among 11 of 40 women (27.5%) who underwent fertility-sparing surgery and remained in the study. Of these 14 pregnancies, 13 (92.9%) delivered at term.⁶⁶ Among the 68 patients who underwent fertility-sparing surgery and were eligible for analysis in the GOG 278 trial, 31 patients attempted to conceive and 16 pregnancies (15 patients) were obtained, including nine full-term pregnancies.⁶⁸

Fertility-sparing approaches are contraindicated in tumors >2 cm and should be considered experimental for selected FIGO 2018 stages IB2–IB3.⁸⁸ Low recurrence rates have been observed after radical trachelectomy using an abdominal laparotomic approach (4.8%); however, laparotomy has been associated with poor fertility outcomes (pregnancy rate: 36%; live birth rate: 66.6%).⁸³ Platinum-based neoadjuvant chemotherapy before fertility-sparing surgery has been explored to shrink tumors that would not be amenable to upfront fertility-sparing approaches.^{89,90} This multimodal approach is associated with better obstetric outcomes, with a pregnancy rate of 74.5% and a live birth rate of 78.7%⁸³; however, oncologic outcomes remain controversial (recurrence rates between

0 and 22.2%).^{91 92} Although there are no clear recommendations on the safety and standardization of neoadjuvant chemotherapy schemes, an ongoing phase II trial will clarify the oncologic and fertility outcomes in women with FIGO 2018 stage IB2 cervical cancer treated with three cycles of neoadjuvant chemotherapy and conservative surgery (CoNteSSA trial).⁹³ The primary outcome is to evaluate the rate of functional uterus defined as successful fertility-sparing surgery and no need for postoperative adjuvant therapy.

Adjuvant Therapy

Adjuvant management after surgery for early-stage cervical cancer is still largely debated and should be individualized based on histopathological risk factors on definitive pathology (Figure 2). Patients with at least one of three high-risk GOG-109 criteria (Peters criteria), such as positive margins, lymph nodes, or parametria, have a high risk of recurrence (25–30%) and adjuvant platinum-based chemoradiotherapy with or without vaginal brachytherapy boost is accepted as standard of care (Table 1).⁹⁴ Notably, recent results of the NRG oncology/RTOG 0724/GOG-0724 study presented at the 2024 American Society of Clinical Oncology (ASCO) Annual Meeting showed no survival benefit for the addition of systemic chemotherapy following chemoradiotherapy in patients with cervical cancer and high-risk factors after radical hysterectomy (Table 1).⁹⁵ In the absence of high-risk criteria, patients meeting at least two of intermediate-risk GOG-92 criteria (Sedlis criteria), such as lymph vascular space invasion, >1/3 stromal invasion, and bulky tumor (largest dimension ≥4 cm), have historically been considered at intermediate risk of recurrence (15–20%) and receive adjuvant external pelvic radiotherapy±concurrent chemotherapy±vaginal brachytherapy boost (Table 1).^{96–98} Patients who do not meet either criterion have generally been considered at low risk of recurrence.

However, the Sedlis criteria were established 25 years ago, and since then there have been tremendous advances in imaging modalities, surgical approaches, radiotherapy techniques, and diagnostic and molecular pathology. The population analyzed by Sedlis et al in 1999 is likely to be different from patients with cervical cancer undergoing hysterectomy today. There is currently wide heterogeneity in the definition of intermediate-risk and low-risk early-stage cervical cancer across centers. Several alternative criteria, such as modified Sedlis criteria or the GOG score, have been developed over time to guide postoperative decision making in clinical practice.^{99 100} The best management for patients meeting only one of the Sedlis criteria and/or having other potentially important risk factors (high grade, age<50 years, non-squamous histology, Silva pattern of stromal invasion, pathological tumor-free distance, close surgical margins) remains controversial.

Postoperative management in the intermediate-risk population is still largely debated and heterogeneous. The benefit of adding chemotherapy to radiotherapy seems to be minimal.¹⁰¹ Moreover, the need for adjuvant therapy has recently been questioned. A recent sub-analysis of the SCCAN study evaluating 692 patients (274 observed after surgery and 418 who received radiotherapy or chemoradiotherapy) suggested that patients with intermediate-risk early-stage cervical cancer who are treated with adequate radical surgery alone achieve similar survival rates compared with those who receive adjuvant therapy.¹⁰² The 5-year disease-free survival (83.2% vs 80.3%; $p=0.365$) and overall survival (88.7% vs 89.0%; $p=0.281$) were not significantly different between the two groups. A

recent meta-analysis by Gómez-Hidalgo et al also reported similar recurrence and mortality rates for patients undergoing observation versus adjuvant treatment.¹⁰³

Reaching an evidence-based consensus for the definition and management of intermediate-risk early-stage cervical cancer has important practical implications. Avoiding unnecessary combined regimens with surgery and radiotherapy whenever possible should be part of the therapeutic goals to preserve quality of life and sexual function. Several ongoing trials (CERVANTES/ENGOT-cx16 (NCT04989647),¹⁰⁴ ACCEPT (NCT05277688), GOG 263 (NCT01101451, NCT04723875)) are currently investigating the role of different types of adjuvant therapy after surgery for early-stage cervical cancer.

Locally Advanced Cervical Cancer

Standard Treatment

Concurrent cisplatin-containing chemoradiation plus brachytherapy is the standard of care for patients with locally advanced cervical cancer or patients who are not candidates for surgery. In 1999, five clinical trials demonstrated that concurrent chemoradiation reduced the risk of death by 30–50% compared with radiotherapy alone.^{94 105–108} A subsequent Cochrane meta-analysis reported a 19% relative reduction in the risk of death with chemoradiotherapy compared with radiotherapy (HR 0.81; 95% CI 0.71 to 0.91; $p<0.001$), which translates into an absolute survival benefit of 6% (from 60% to 66%) at 5 years.¹⁰⁹ Figure 4 shows the primary treatment algorithm for locally advanced cervical cancer (FIGO 2018 stages IB3, IIA2–IVA).

Fertility preservation options such as oocyte, embryo or ovarian tissue preservation, and oocyte donation should be discussed with patients. Ovarian transposition may be considered before pelvic radiotherapy to preserve endocrine function for selected patients <45 years of age with squamous cell cancer or usual-type (HPV-associated) adenocarcinoma, except in the case of positive nodes.^{110–113}

Para-aortic Lymph Node Staging

Historically, it has been widely debated whether para-aortic lymphadenectomy provides a therapeutic benefit or has a diagnostic role to avoid extended-field radiotherapy in patients with clinically negative para-aortic lymph nodes.⁸⁰ Routine adoption of this approach into clinical practice is highly heterogeneous among institutions worldwide.

However, the role of lymphadenectomy has been strongly challenged by recent evidence. The prospective multicenter randomized Uterus-11 trial compared surgical (pelvic and para-aortic lymphadenectomy) versus imaging (abdominal CT and/or MRI and chest radiograph, CT or PET-CT) staging prior to primary chemoradiation for locally advanced cervical cancer and demonstrated no statistically significant difference in disease-free survival between the two groups after a median follow-up of 90 months (Table 1).¹¹⁴ Therefore, systematic para-aortic lymphadenectomy should not be recommended. Extended-field para-aortic radiation can be considered based on the presence and number of positive pelvic nodes (more than two level I nodes or at least one level II (common iliac) nodes) or tumor size (>2 cm).^{115–117}

Routine preoperative PET/CT imaging as an advanced diagnostic tool has the potential to further improve the reliability of preoperative




Figure 4 Primary treatment for locally advanced cervical cancer. All staging is based on updated FIGO 2018 staging. Treatment planning should be multidisciplinary and individualized. Treatment options may vary across countries depending on national guidelines and socioeconomic factors. *Pelvic exenteration is generally indicated as the last therapeutic option for patients with advanced, persistent, or recurrent cervical cancer and should be considered an individualized approach after careful patient selection and counseling by a multidisciplinary team in specialized centers. †Ovarian preservation may be considered in premenopausal women (<45 years of age) with squamous cell carcinoma or HPV-associated adenocarcinoma. BRT, brachytherapy; CCRT, concurrent chemoradiotherapy; CHT, chemotherapy; EBRT, external beam radiotherapy; FIGO, International Federation of Gynecology and Obstetrics; HPV, human papillomavirus; LACC, locally advanced cervical cancer; NACT, neoadjuvant chemotherapy; PALND, para-aortic lymph node dissection; PLND, pelvic lymph node dissection.

radiologic staging, with a sensitivity of 75% and specificity of 92% for detecting lymph node metastases in the para-aortic region.¹¹⁸ Patients in the Uterus-11 trial did not routinely undergo preoperative PET-CT. The PAROLA trial is assessing whether para-aortic surgical staging followed by tailored chemoradiation improves disease-free survival while avoiding unnecessary extended-field chemoradiation compared with radiologic PET/CT staging alone in patients with stage IIIC1 cervical cancer, the subset at higher risk of positive para-aortic lymph nodes.¹¹⁹ However, it should be noted that requiring preoperative PET/CT staging for every patient may limit applicability to the real-world setting due to disparities in the availability of diagnostic resources. PET/CT imaging is not available in 95% of low-income countries and 92% of lower-middle-income countries; therefore, treatment standards may vary across countries depending on available resources.¹²⁰

Radical Surgery versus Chemoradiotherapy

Radical surgery (with or without neoadjuvant chemotherapy) as a potential alternative to standard concurrent chemoradiotherapy for locally advanced cervical cancer remains controversial.^{121–123} Two recent randomized clinical trials (Gupta et al¹²⁴ and EORTC 55994¹²⁴) failed to demonstrate superiority of the neoadjuvant chemotherapy-surgery approach compared with concomitant chemoradiotherapy; therefore, concomitant chemoradiotherapy remains the standard of care for locally advanced cervical cancer (Table 1).

Nonetheless, radical surgery after neoadjuvant chemotherapy may yield non-inferior overall survival outcomes with acceptable toxicity and health-related quality of life in selected patients with FIGO 2018 stage IB3/IIA2 cervical cancer.^{124 125} In particular, radical

surgery may be considered an alternative approach to radiotherapy for patients with FIGO 2018 stage IB3/IIA2 disease in the following scenarios: (1) young patients who decline radiotherapy for concerns about possible radiotherapy-induced toxicity and want to preserve overall quality of life and sexual function; (2) poor availability and/or accessibility to adequate radiotherapy facilities, especially in low-income countries; (3) contraindications to radiotherapy (active inflammatory bowel disease, inability to maintain patient positioning, prior radiotherapy).^{10 126 127} Patients should be discussed by a multidisciplinary team and counseled about the advantages and disadvantages of both treatment options (surgery vs radiotherapy). To avoid the combination of radical surgery and postoperative radiotherapy, decision making regarding adjuvant management after radical surgery should not be rigorously based on traditional selection criteria for adjuvant treatment, but should be individualized.¹¹

Novel Approaches

Immunotherapy has demonstrated activity not only in the management of metastatic, persistent, and recurrent cervical cancer but more recently in locally advanced cases. While the addition of durvalumab (anti-programmed cell death ligand 1 (PD-L1) immune checkpoint inhibitor) to chemoradiation therapy for locally advanced cervical cancer did not provide a survival benefit in the randomized phase III CALLA trial,¹²⁸ the results of the randomized phase III KEYNOTE-A18/ENGOT-cx11 trial demonstrated the efficacy and safety of pembrolizumab (anti-PD-1 immune checkpoint inhibitor) versus placebo in combination with concurrent chemoradiotherapy (Table 1).¹² Key differences between the CALLA and KEYNOTE-A18

trials include the different type of immune checkpoint inhibitor (anti-PD-L1 vs anti-PD-1, respectively) and patient risk factors. The CALLA study enrolled lower risk patients with approximately 10% positive para-aortic lymph nodes versus 22% in the KEYNOTE-A18.

The KEYNOTE-A18 study included patients with high-risk locally advanced cervical cancer: FIGO 2014 stage IB2–IIB with node-positive disease or FIGO 2014 stage III–IVA disease.¹² Patients in the experimental arm received pembrolizumab 200 mg via intravenous infusion every 3 weeks for five cycles in combination with chemoradiotherapy (with single-agent cisplatin), followed by 400 mg of pembrolizumab intravenously every 6 weeks until disease progression, unacceptable toxicity, or for up to 24 months (further 15 cycles). The primary endpoints were progression-free survival and overall survival. Compared with chemoradiation alone, the addition of pembrolizumab demonstrated a statistically significant improvement in 24-month progression-free survival in the overall population (68% vs 57%; HR 0.70; 95% CI 0.55 to 0.89; $p=0.002$). Exploratory subgroup analyses showed that the benefit was primarily attributed to patients with FIGO 2014 stage III–IVA disease. As a result, the FDA approved pembrolizumab with chemoradiotherapy only for patients with FIGO 2014 stage III–IVA cervical cancer. In a recent press release, it was anticipated that a pre-specified interim analysis also demonstrated a statistically significant improvement in overall survival.¹²⁹

Immunotherapy may have a promising role earlier in the treatment paradigm of cervical cancer. The combination of neoadjuvant chemo-immunotherapy with surgery holds potential as a novel therapeutic approach for locally advanced cervical cancer. The single-arm, phase 2 NACI study showed promising antitumour activity (objective response rate: 98%) and a manageable toxicity profile for the combination of camrelizumab (anti-PD-1) with neoadjuvant chemotherapy in patients with FIGO 2018 stage IB3, IIA2, or IIB/IIIC1r with a tumor diameter ≥ 4 cm.¹³⁰ The ongoing single-arm, phase 2 MITO CERV 3 trial (NCT04238988) is evaluating the role of pembrolizumab combined with three cycles of platinum-based chemotherapy as neoadjuvant treatment before radical surgery for patients with FIGO 2009 stage IB2–IIB disease.

Furthermore, while the multicenter randomized OUTBACK trial failed to demonstrate a survival benefit for patients who received adjuvant carboplatin/paclitaxel-based chemotherapy after concurrent chemoradiotherapy at the expense of a higher toxicity,¹³¹ results from the INTERLACE trial presented at the 2023 ESMO congress might be practice changing (Table 1).¹³ This randomized multicenter study involving 500 patients evaluated short-course weekly induction with six cycles of carboplatin AUC2 plus paclitaxel 80 mg/m² administered before concurrent chemoradiotherapy. Even though methodological concerns have been raised about the type of radiation therapy and the enrollment timeline, induction chemotherapy was shown to significantly improve 5-year progression-free (73% vs 64%; HR 0.65; 95% CI 0.46 to 0.91; $p=0.013$) and overall survival (80% vs 72%; HR 0.61; 95% CI 0.40 to 0.91; $p=0.04$) and was feasible in diverse healthcare settings and countries, which adds value considering that most cases are diagnosed in low-income countries with limited resources.¹³

Metastatic, Persistent, or Recurrent Cervical Cancer

The management of metastatic, persistent or recurrent cervical cancer poses a significant clinical challenge and depends on

multidisciplinary assessment of clinical factors and patient goals, including age, performance status, location and extent of disease, prior treatment modalities, treatment-free interval, and patient preferences. The standard of care is based primarily on systemic therapies and enrollment in clinical trials is strongly encouraged. For recurrent cervical cancer, repeat biopsy could be considered to perform molecular/genetic analysis and identify targetable biomarkers, including PD-L1, MMR, HER2, tumor mutational burden, *NTRK* and *RET* gene fusion testing. Early involvement of a palliative care specialist is advised. Best supportive care should be provided in patients unfit to receive further treatments.^{9,11}

Disease Not Amenable to Local Treatment

The management of metastatic, persistent, or distant recurrent cervical cancer not amenable to local treatment is outlined in Figure 5. Based on the practice-changing results from the GOG 240¹³² and KEYNOTE-826 trials,¹³³ platinum-based chemotherapy $\pm$ bevacizumab $\pm$ (anti-PD-1) pembrolizumab (in patients with PD-L1 positive tumors, assessed as combined positive score ≥ 1) is the current first-line treatment of choice (Table 1). The phase III, randomized GOG 240 trial demonstrated that the addition of bevacizumab (15 mg/m² every 3 weeks until disease progression or unacceptable toxicity) to combination chemotherapy (cisplatin 50 mg/m² plus paclitaxel 135–175 mg/m²) in patients with recurrent, persistent, or metastatic cervical cancer was associated with an improvement of 3.7 months in median overall survival.¹³² It should be noted that bevacizumab has been associated with an increased risk of hypertension, thromboembolic events, gastrointestinal fistulas and perforations, and delayed wound healing; therefore, caution should be exercised to exclude any individual contraindications. In the phase III, randomized KEYNOTE-826 trial, progression-free and overall survival were significantly longer with pembrolizumab (200 mg every 3 weeks until disease progression, unacceptable toxicity, or for up to 35 cycles) than with placebo among patients with persistent, recurrent, or metastatic cervical cancer who were also receiving chemotherapy with or without bevacizumab.¹³³ Since no benefit was found for tumors not expressing PD-L1 (combined positive score <1), the FDA approved the addition of pembrolizumab for tumors with PD-L1 combined positive score ≥ 1 . Locoregional therapies may be combined with systemic treatment, such as (1) pelvic radiotherapy in the case of complete response of distant metastases with response or stability of pelvic disease; or (2) surgical resection or radiotherapy for residual extrapelvic oligometastatic disease.¹³⁴

The BEATcc trial showed that the anti-PD-L1 immune checkpoint inhibitor atezolizumab (1200 mg every 3 weeks until disease progression or unacceptable toxicity) significantly improved progression-free and overall survival when combined with standard platinum-based chemotherapy plus bevacizumab, paving the way for another practice-changing strategy in this setting (Table 1).¹³⁵ Moreover, chemotherapy-free immunotherapy with nivolumab alone or in combination with ipilimumab provided encouraging durable responses with manageable toxicity in recurrent or metastatic HPV-associated cervical cancer, regardless of tumor PD-L1 expression, according to updated findings from the phase I/II CheckMate-358 trial.¹³⁶

Second-line treatment should include immunotherapy if not previously administered. The phase III, randomized EMPOWER-Cervical




Figure 5 Management of metastatic, persistent, or distant recurrent cervical cancer. Treatment planning should be multidisciplinary and individualized. Treatment options may vary across countries depending on national guidelines and socioeconomic factors. CHT, chemotherapy; CR, complete response; PD-L1 CPS, programmed cell death ligand 1 combined positive score; PR, partial response; RT, radiotherapy.

1/GOG-3016/ENGOT-cx9 trial demonstrated that cemiplimab (anti-PD-1 immune checkpoint inhibitor; 350 mg every 3 weeks until disease progression, unacceptable toxicity, or for up to 96 weeks) significantly improves overall survival compared with single-agent chemotherapy in patients with recurrent cervical cancer after first-line platinum-containing chemotherapy, regardless of PD-L1 status (Table 1).¹³⁷ In addition, the phase II KEYNOTE-158 basket study showed that pembrolizumab monotherapy (200 mg every 3 weeks until progression, unacceptable toxicity, or for up to 2 years) provides durable anti-tumor activity and manageable safety in patients with advanced PD-L1-positive cervical cancer who experienced progression during or after chemotherapy (Table 1).¹³⁸ Estimates of PD-L1 expression in cervical cancer are 71% in squamous cell carcinoma and 33% in adenocarcinoma.¹³⁷ Pembrolizumab has additional FDA-approved biomarkers, such as microsatellite instability-high (MSI-H)/mismatch repair deficiency (MMRd) and tumor mutational burden-high (≥ 10 mutations/megabase), which have prevalence rates of 2.6–11.3% and 21.3%, respectively, in cervical cancer.^{139–141}

Standard chemotherapy can be considered in patients who had already received immunotherapy.^{9–11} In addition, based on the

results from the phase II ENGOT-cx6/InnovaTV 204 trial¹⁴² and the phase III ENGOT-cx12/InnovaTV 301 trial,¹⁴³ tisotumab vedotin-tftv has recently been approved for patients with disease progression on or after standard chemotherapy±bevacizumab±anti-PD-(L)1 therapy. Tisotumab vedotin-tftv is an antibody-drug conjugate and acts by combining the targeting properties of tissue factor directed monoclonal antibodies with the anti-cancer activity of a microtubule inhibitor payload. The target is the tissue factor, which is highly expressed on many solid tumors, but it is not used as a biomarker for patient selection. In the phase III innovaTV 301 study, tisotumab vedotin-tftv showed a statistically significant and clinically meaningful improvement in overall survival, progression-free survival, and objective response rate versus chemotherapy, with a manageable and tolerable safety profile, although particular attention should be paid to potential ocular toxicity (Table 1).¹⁴³ Interestingly, the phase Ib/II ENGOT-cx8/InnovaTV205 trial suggested that tisotumab vedotin-tftv may have a promising role when combined with carboplatin or pembrolizumab in the first-line setting and with pembrolizumab in the recurrent setting.¹⁴⁴

The pipeline of novel biomarker-driven targeted treatments is constantly evolving, and more treatment options are expected




Figure 6 Treatment algorithm for locoregional and oligometastatic recurrent cervical cancer. Treatment planning should be multidisciplinary and individualized. Treatment options may vary across countries depending on national guidelines and socioeconomic factors. BRT, brachytherapy; CHT, chemotherapy; EBRT, external beam radiotherapy; IORT, intraoperative radiotherapy; LEER, laterally extended endopelvic resection; RT, radiotherapy.

to be available in the near future. Immune checkpoint inhibitors and antibody-drug conjugates are the cornerstone of this new era of precision oncology in cervical cancer. The FDA recently granted accelerated approval to fam-trastuzumab deruxtecan-nxki (antibody-drug conjugate targeting HER2) for adult patients with unresectable or metastatic HER2-expressing (immunohistochemistry 3+) solid tumors who have received prior systemic treatment and have no satisfactory alternative treatment options. Positive results from the primary analysis of the phase II DESTINY-PanTumor02 trial showed that fam-trastuzumab deruxtecan-nxki provided durable responses and clinically meaningful survival benefit in multiple HER2-expressing advanced solid tumors, including cervical cancer (Table 1).¹⁴⁵ Single-agent larotrectinib and entrectinib are indicated for *NTRK* gene fusion tumors, with cervical cancer showing a 1.5% *NTRK* gene fusion positivity rate.¹⁴⁶ Selpercatinib is designated for *RET* gene fusion tumors, and cervical cancer has a 1.52% occurrence of altered *RET*.¹⁴⁷

Disease Amenable to Local Treatment

The treatment algorithm for recurrent cervical cancer amenable to local therapy, including locoregional and oligometastatic disease, is outlined in Figure 6. Multifocal local relapse or multiple/distant metastases (non-oligometastatic disease) not amenable to local therapies should be treated like metastatic disease. Locoregional pelvic recurrences are classified into central (if the tumor is confined to the vagina, bladder, rectum, and/or parametrium) or non-central/lateral (if the tumor involves the muscles and vasculature of the lateral pelvic wall). Oligometastatic disease refers to cases with one to five metastatic or recurrent lesions.^{9 148 149} In the case of suspected tumor persistence or recurrence, histological verification by biopsy is strongly recommended if feasible before treatment planning.

The treatment for locoregional relapse depends on whether the patient has had previous radiotherapy. Previously irradiated patients should be considered for surgery. Pelvic exenteration is generally indicated as the last therapeutic option for patients with advanced, persistent, or recurrent cervical cancer and should be considered an individualized approach after careful patient selection and counseling by a multidisciplinary team in specialized centers.^{9-11 150} It is generally recommended for central pelvic recurrence (<2 cm) in the absence of pelvic sidewall involvement, extrapelvic nodes, or peritoneal disease. Extended pelvic surgery (laterally extended endopelvic resection)±intraoperative radiotherapy can also be considered in carefully selected cases with pelvic sidewall involvement. In the era of targeted therapies, systemic therapy could be considered before and/or after surgery for locoregional recurrence. For patients who wish to avoid pelvic exenteration, systemic treatment alone remains a viable option, especially in the bevacizumab/immunotherapy-naïve setting. Individualized re-irradiation with external beam radiotherapy±chemotherapy can be also considered an alternative to surgical resection for locoregional relapse after primary radiotherapy.

Radiotherapy-naïve patients and those with outside-field relapses should be treated with external beam radiotherapy±chemotherapy±brachytherapy. Surgical resection±intraoperative chemotherapy is another option. Brachytherapy alone might be considered for small superficial vaginal relapses (<5 mm thickness). For highly selected patients with isolated distant metastases (nodal, lung, liver, bone), either at initial diagnosis or at relapse, amenable to local therapy, combined locoregional treatment modalities including surgery, external beam radiotherapy±concurrent chemotherapy, and/or local ablative therapies might be considered, possibly followed by adjuvant systemic therapy.⁹⁻¹¹

- its clinical impact in cervical cancer. *Int J Radiat Oncol Biol Phys* 2010;77:502–8.
- 52 Lund KV, Simonsen TG, Hompland T, *et al*. Short-term pretreatment DCE-MRI in prediction of outcome in locally advanced cervical cancer. *Radiother Oncol* 2015;115:379–85.
 - 53 Sarabhai T, Schaarschmidt BM, Wetter A, *et al*. Comparison of (18)F-FDG PET/MRI and MRI for pre-therapeutic tumor staging of patients with primary cancer of the uterine cervix. *Eur J Nucl Med Mol Imaging* 2018;45:67–76.
 - 54 Gruenewald J, Schaarschmidt BM, Heubner M, *et al*. Integrated PET/MRI for whole-body staging of patients with primary cervical cancer: preliminary results. *Eur J Nucl Med Mol Imaging* 2015;42:1814–24.
 - 55 Ou Z, Mao W, Tan L, *et al*. Prediction of postoperative pathologic risk factors in cervical cancer patients treated with radical hysterectomy by machine learning. *Curr Oncol* 2022;29:9613–29.
 - 56 Pujade-Lauraine E, Tan DSP, Leary A, *et al*. Comparison of global treatment guidelines for locally advanced cervical cancer to optimize best care practices: a systematic and scoping review. *Gynecol Oncol* 2022;167:360–72.
 - 57 Baalbergen A, Veenstra Y, Stalpers LL, *et al*. Primary surgery versus primary radiation therapy with or without chemotherapy for early adenocarcinoma of the uterine cervix. *Cochrane Database Syst Rev* 2013;2013.
 - 58 Gien LT, Beauchemin MC, Thomas G. Adenocarcinoma: a unique cervical cancer. *Gynecol Oncol* 2010;116:140–6.
 - 59 Park J-Y, Kim D-Y, Kim J-H, *et al*. Outcomes after radical hysterectomy in patients with early-stage adenocarcinoma of uterine cervix. *Br J Cancer* 2010;102:1692–8.
 - 60 Ramirez PT, Frumovitz M, Pareja R, *et al*. Minimally invasive versus abdominal radical hysterectomy for cervical cancer. *N Engl J Med* 2018;379:1895–904.
 - 61 Ramirez PT, Robledo KP, Frumovitz M, *et al*. LACC trial: final analysis on overall survival comparing open versus minimally invasive radical hysterectomy for early-stage cervical cancer. *J Clin Oncol* 2024.
 - 62 Bogani G, Di Donato V, Muzii L, *et al*. Assessing the role of minimally invasive radical hysterectomy for early-stage cervical cancer. *Eur J Obstet Gynecol Reprod Biol* 2022;275:64–9.
 - 63 Touhami O, Plante M. Minimally invasive surgery for cervical cancer in light of the LACC trial: what have we learned? *Curr Oncol* 2022;29:1093–106.
 - 64 Falconer H, Palsdottir K, Stalberg K, *et al*. Robot-assisted approach to cervical cancer (RACC): an international multi-center, open-label randomized controlled trial. *Int J Gynecol Cancer* 2019;29:1072–6.
 - 65 Querleu D, Cibula D, Abu-Rustum NR. 2017 update on the querleu-morrow classification of radical hysterectomy. *Ann Surg Oncol* 2017;24:3406–12.
 - 66 Schmeler KM, Pareja R, Lopez Blanco A, *et al*. ConCerv: a prospective trial of conservative surgery for low-risk early-stage cervical cancer. *Int J Gynecol Cancer* 2021;31:1317–25.
 - 67 Plante M, Kwon JS, Ferguson S, *et al*. Simple versus radical hysterectomy in women with low-risk cervical cancer. *N Engl J Med* 2024;390:819–29.
 - 68 Evaluation of physical function and quality of life before and after non-radical surgical therapy (extra fascial hysterectomy or cone biopsy with pelvic lymphadenectomy) for stage IA1 (LVSI+) and IA2-IB1 cervical cancer (GOG-278). SGO 2024 Annual Meeting on Women's Cancer; San Diego (CA, USA), 2024:16–8.
 - 69 Cibula D, Dusek J, Jarkovsky J, *et al*. A prospective multicenter trial on sentinel lymph node biopsy in patients with early-stage cervical cancer (SENTIX). *Int J Gynecol Cancer* 2019;29:212–5.
 - 70 Favre G, Guani B, Balaya V, *et al*. Sentinel lymph-node biopsy in early-stage cervical cancer: the 4-year follow-up results of the senticol 2 trial. *Front Oncol* 2020;10:621518.
 - 71 Kadkhodayan S, Hasanizadeh M, Treglia G, *et al*. Sentinel node biopsy for lymph nodal staging of uterine cervix cancer: a systematic review and meta-analysis of the pertinent literature. *Eur J Surg Oncol* 2015;41:1–20.
 - 72 Wu Y, Li Z, Wu H, *et al*. Sentinel lymph node biopsy in cervical cancer: a meta-analysis. *Mol Clin Oncol* 2013;1:1025–30.
 - 73 Balaya V, Guani B, Morice P, *et al*. Long-term oncological safety of sentinel lymph node biopsy in early-stage cervical cancer: a post-hoc analysis of SENTICOL I and SENTICOL II cohorts. *Gynecol Oncol* 2022;164:53–61.
 - 74 Lecuru FR, McCormack M, Hillemanns P, *et al*. SENTICOL III: an international validation study of sentinel node biopsy in early cervical cancer. A GINECO, ENGOT, GCIG and multicenter study. *Int J Gynecol Cancer* 2019;29:829–34.
 - 75 Seino M, Nagase S, Tokunaga H, *et al*. Japan society of gynecologic oncology 2022 guidelines for uterine cervical neoplasm treatment. *J Gynecol Oncol* 2024;35:e15.
 - 76 Yoo JG, Lee SJ, Nam EJ, *et al*. Clinical practice guidelines for cervical cancer: the Korean society of gynecologic oncology guidelines. *J Gynecol Oncol* 2024;35:e44.
 - 77 Agustí N, Viveros-Carreño D, Mora-Soto N, *et al*. Diagnostic accuracy of sentinel lymph node frozen section analysis in patients with early-stage cervical cancer: a systematic review and meta-analysis. *Gynecol Oncol* 2023;177:157–64.
 - 78 Sandadi S, Tanner EJ, Khoury-Collado F, *et al*. Radical surgery with individualized postoperative radiation for stage IB cervical cancer: oncologic outcomes and severe complications. *Int J Gynecol Cancer* 2013;23:553–8.
 - 79 van den Akker MJE, Horeweg N, Beltman JJ, *et al*. Efficacy and toxicity of postoperative external beam radiotherapy or chemoradiation for early-stage cervical cancer. *Int J Gynecol Cancer* 2020;30:1878–86.
 - 80 Gold MA, Tian C, Whitney CW, *et al*. Surgical versus radiographic determination of para-aortic lymph node metastases before chemoradiation for locally advanced cervical carcinoma: a gynecologic oncology group study. *Cancer* 2008;112:1954–63.
 - 81 Benard VB, Watson M, Castle PE, *et al*. Cervical carcinoma rates among young females in the United States. *Obstet Gynecol* 2012;120:1117–23.
 - 82 Dargent D, Martin X, Sacchetoni A, *et al*. Laparoscopic vaginal radical trachelectomy: a treatment to preserve the fertility of cervical carcinoma patients. *Cancer* 2000;88:1877–82.
 - 83 Morice P, Maulard A, Scherier S, *et al*. Oncologic results of fertility sparing surgery of cervical cancer: an updated systematic review. *Gynecol Oncol* 2022;165:169–83.
 - 84 Wright JD, Grigsby PW, Brooks R, *et al*. Utility of parametrectomy for early stage cervical cancer treated with radical hysterectomy. *Cancer* 2007;110:1281–6.
 - 85 Frumovitz M, Sun CC, Schmeler KM, *et al*. Parametrial involvement in radical hysterectomy specimens for women with early-stage cervical cancer. *Obstet Gynecol* 2009;114:93–9.
 - 86 Ramirez PT, Pareja R, Rendón GJ, *et al*. Management of low-risk early-stage cervical cancer: should conization, simple trachelectomy, or simple hysterectomy replace radical surgery as the new standard of care? *Gynecol Oncol* 2014;132:254–9.
 - 87 Batman SH, Schmeler KM. Fertility-sparing and less radical surgery for cervical cancer. *Curr Oncol Rep* 2022;24:1541–8.
 - 88 Di Donato V, Caruso G, Sassu CM, *et al*. Fertility-sparing surgery for women with stage I cervical cancer of 4 cm or larger: a systematic review. *J Gynecol Oncol* 2021;32:e83.
 - 89 Buda A, Borghese M, Puppo A, *et al*. Neoadjuvant chemotherapy prior fertility-sparing surgery in women with FIGO 2018 stage IB2 cervical cancer: a systematic review. *Cancers (Basel)* 2022;14:797.
 - 90 Pareja R, Rendón GJ, Vasquez M, *et al*. Immediate radical trachelectomy versus neoadjuvant chemotherapy followed by conservative surgery for patients with stage IB1 cervical cancer with tumors 2cm or larger: a literature review and analysis of oncological and obstetrical outcomes. *Gynecol Oncol* 2015;137:574–80.
 - 91 Burbano J, Heredia F, Sanabria D, *et al*. Neoadjuvant chemotherapy prior to fertility-sparing surgery in cervical tumors larger than 2 cm: a systematic review on fertility and oncologic outcomes. *Int J Gynecol Cancer* 2021;31:387–98.
 - 92 Ronsini C, Solazzo MC, Bizzarri N, *et al*. Fertility-sparing treatment for early-stage cervical cancer ≥2cm: a problem with a thousand nuances—a systematic review of oncological outcomes. *Ann Surg Oncol* 2022;29:8346–58.
 - 93 Plante M, van Trommel N, Lheureux S, *et al*. FIGO 2018 stage IB2 (2–4 cm) cervical cancer treated with neo-adjuvant chemotherapy followed by fertility sparing surgery (CONTESSA); neo-adjuvant chemotherapy and conservative surgery in cervical cancer to preserve fertility (NEOCON-F). A PMHC, DGO, GCIG/CCRN and multicenter study. *Int J Gynecol Cancer* 2019;29:969–75.
 - 94 Peters WA III, Liu PY, Barrett RJ II, *et al*. Concurrent chemotherapy and pelvic radiation therapy compared with pelvic radiation therapy alone as adjuvant therapy after radical surgery in high-risk early-stage cancer of the cervix. *JCO* 2000;18:1606–13.
 - 95 Adjuvant chemotherapy following concurrent chemoradiation (CRT) in patients with high-risk early-stage cervical carcinoma following radical hysterectomy: results of NRG oncology/RTOG 0724/ GOG0724. ASCO Annual Meeting; 2024
 - 96 Sedlis A, Bundy BN, Rotman MZ, *et al*. A randomized trial of pelvic radiation therapy versus no further therapy in selected patients with stage IB carcinoma of the cervix after radical hysterectomy and

- pelvic lymphadenectomy: a gynecologic oncology group study. *Gynecol Oncol* 1999;73:177–83.
- 97 Rotman M, Sedlis A, Piedmonte MR, *et al*. A phase III randomized trial of postoperative pelvic irradiation in stage IB cervical carcinoma with poor prognostic features: follow-up of a gynecologic oncology group study. *Int J Radiat Oncol Biol Phys* 2006;65:169–76.
 - 98 Rogers L, Siu SSN, Luesley D, *et al*. Radiotherapy and chemoradiation after surgery for early cervical cancer. *Cochrane Database Syst Rev* 2012;2012.
 - 99 Levinson K, Beavis AL, Purdy C, *et al*. Beyond sedlis—a novel histology-specific nomogram for predicting cervical cancer recurrence risk: an NRG/GOG ancillary analysis. *Gynecol Oncol* 2021;162:532–8.
 - 100 Kinjo Y, Matsuura Y, Ohguri T, *et al*. Retrospective analysis of prognosis using the gynecology oncology group score of stage IB-IIA node negative uterine cervical cancer after radical hysterectomy and trachelectomy. *Mol Clin Oncol* 2022;16:105.
 - 101 Rodriguez J, Viveros-Carreño D, Pareja R. Adjuvant treatment after radical surgery for cervical cancer with intermediate risk factors: is it time for an update? *Int J Gynecol Cancer* 2022;32:1219–26.
 - 102 Cibula D, Akilli H, Jarkovsky J, *et al*. Role of adjuvant therapy in intermediate-risk cervical cancer patients - subanalyses of the SCCAN study. *Gynecol Oncol* 2023;170:195–202.
 - 103 Gómez-Hidalgo NR, Acosta Ú, Rodríguez TG, *et al*. Adjuvant therapy in early-stage cervical cancer after radical hysterectomy: are we overtreating our patients? A meta-analysis. *Clin Transl Oncol* 2022;24:1605–14.
 - 104 Cibula D, Borčinová M, Kocian R, *et al*. CERVANTES: an international randomized trial of radical surgery followed by adjuvant (chemo) radiation versus no further treatment in patients with early-stage, intermediate-risk cervical cancer (CEEEOG-CX-05; ENGOT-CX16). *Int J Gynecol Cancer* 2022;32:1327–31.
 - 105 Keys HM, Bundy BN, Stehman FB, *et al*. Cisplatin, radiation, and adjuvant hysterectomy compared with radiation and adjuvant hysterectomy for bulky stage IB cervical carcinoma. *N Engl J Med* 1999;340:1154–61.
 - 106 Morris M, Eifel PJ, Lu J, *et al*. Pelvic radiation with concurrent chemotherapy compared with pelvic and para-aortic radiation for high-risk cervical cancer. *N Engl J Med* 1999;340:1137–43.
 - 107 Rose PG, Bundy BN, Watkins EB, *et al*. Concurrent cisplatin-based radiotherapy and chemotherapy for locally advanced cervical cancer. *N Engl J Med* 1999;340:1144–53.
 - 108 Whitney CW, Sause W, Bundy BN, *et al*. Randomized comparison of fluorouracil plus cisplatin versus hydroxyurea as an adjunct to radiation therapy in stage IIB-IVA carcinoma of the cervix with negative para-aortic lymph nodes: a gynecologic oncology group and southwest oncology group study. *J Clin Oncol* 1999;17:1339–48.
 - 109 Chemoradiotherapy for Cervical Cancer Meta-analysis Collaboration (CCCMAC). Reducing uncertainties about the effects of chemoradiotherapy for cervical cancer: individual patient data meta-analysis. *Cochrane Database Syst Rev* 2010;2010.
 - 110 Hoekman EJ, Broeders E, Louwe LA, *et al*. Ovarian function after ovarian transposition and additional pelvic radiotherapy: a systematic review. *Eur J Surg Oncol* 2019;45:1328–40.
 - 111 Morice P, Juncker L, Rey A, *et al*. Ovarian transposition for patients with cervical carcinoma treated by radiosurgical combination. *Fertil Steril* 2000;74:743–8.
 - 112 Pahisa J, Martínez-Román S, Martínez-Zamora MA, *et al*. Laparoscopic ovarian transposition in patients with early cervical cancer. *Int J Gynecol Cancer* 2008;18:584–9.
 - 113 Wo JY, Viswanathan AN. Impact of radiotherapy on fertility, pregnancy, and neonatal outcomes in female cancer patients. *Int J Radiat Oncol Biol Phys* 2009;73:1304–12.
 - 114 Marnitz S, Tsunoda AT, Martus P, *et al*. Surgical versus clinical staging prior to primary chemoradiation in patients with cervical cancer FIGO stages IIB-IVA: oncologic results of a prospective randomized international multicenter (Uterus-11) intergroup study. *Int J Gynecol Cancer* 2020;30:1855–61.
 - 115 Gouy S, Morice P, Narducci F, *et al*. Prospective multicenter study evaluating the survival of patients with locally advanced cervical cancer undergoing laparoscopic para-aortic lymphadenectomy before chemoradiotherapy in the era of positron emission tomography imaging. *J Clin Oncol* 2013;31:3026–33.
 - 116 Huang H, Liu J, Li Y, *et al*. Metastasis to deep obturator and para-aortic lymph nodes in 649 patients with cervical carcinoma. *Eur J Surg Oncol* 2011;37:978–83.
 - 117 Sakuragi N, Satoh C, Takeda N, *et al*. Incidence and distribution pattern of pelvic and paraaortic lymph node metastasis in patients with stages IB, IIA, and IIB cervical carcinoma treated with radical hysterectomy. *Cancer* 1999;85:1547–54.
 - 118 Rose PG, Adler LP, Rodriguez M, *et al*. Positron emission tomography for evaluating para-aortic nodal metastasis in locally advanced cervical cancer before surgical staging: a surgicalpathologic study. *J Clin Oncol* 1999;17:41–5.
 - 119 Martinez A, Lecuru F, Bizzarri N, *et al*. PARa-aortic lymphadenectomy in locally advanced cervical cancer (PAROLA trial): a GINECO, ENGOT, and GCIG study. *Int J Gynecol Cancer* 2023;33:293–8.
 - 120 Verduzco-Aguirre HC, Lopes G, Soto-Perez-De-Celis E. Implementation of diagnostic resources for cancer in developing countries: a focus on PET/CT. *Ecancermedicalscience* 2019;13:ed87.
 - 121 Benedetti-Panici P, Greggi S, Colombo A, *et al*. Neoadjuvant chemotherapy and radical surgery versus exclusive radiotherapy in locally advanced squamous cell cervical cancer: results from the Italian multicenter randomized study. *J Clin Oncol* 2002;20:179–88.
 - 122 Friedlander M, Kaye SB, Sullivan A, *et al*. Cervical carcinoma: a drug-responsive tumor—experience with combined cisplatin, vinblastine, and bleomycin therapy. *Gynecol Oncol* 1983;16:275–81.
 - 123 Kirsten F, Atkinson KH, Coppleson JVM, *et al*. Combination chemotherapy followed by surgery or radiotherapy in patients with locally advanced cervical cancer. *BJOG* 1987;94:583–8.
 - 124 Gupta S, Maheshwari A, Parab P, *et al*. Neoadjuvant chemotherapy followed by radical surgery versus concomitant chemotherapy and radiotherapy in patients with stage IB2, IIA, or IIB squamous cervical cancer: a randomized controlled trial. *J Clin Oncol* 2018;36:1548–55.
 - 125 Kenter GG, Greggi S, Vergote I, *et al*. Randomized phase III study comparing neoadjuvant chemotherapy followed by surgery versus chemoradiation in stage IB2-IIB cervical cancer: EORTC-55994. *J Clin Oncol* 2023;41:5035–43.
 - 126 Caruso G, Bruni S, Lapresa M, *et al*. Dose-dense neoadjuvant chemotherapy before radical surgery in cervical cancer: a retrospective cohort study and systematic literature review. *Int J Gynecol Cancer* 2024;34:47–57.
 - 127 Caruso G, Palaia I, Di Donato V, *et al*. Radical surgery after neoadjuvant chemotherapy for locally advanced neuroendocrine cancer of the cervix. *Anticancer Res* 2021;41:4431–8.
 - 128 Monk BJ, Toita T, Wu X, *et al*. Durvalumab versus placebo with chemoradiotherapy for locally advanced cervical cancer (CALLA): a randomised, double-blind, phase 3 trial. *Lancet Oncol* 2023;24:1334–48.
 - 129 Merck's Keytruda (pembrolizumab) plus chemoradiotherapy (CRT) significantly improved overall survival (OS) versus CRT alone in patients with newly diagnosed high-risk locally advanced cervical cancer. *N R Merck* 2024. Available: <https://tinyurl.com/4n4mfs9a>
 - 130 Li K, Chen J, Hu Y, *et al*. Neoadjuvant chemotherapy plus camrelizumab for locally advanced cervical cancer (NACI study): a multicentre, single-arm, phase 2 trial. *Lancet Oncol* 2024;25:76–85.
 - 131 Mileskin LR, Moore KN, Barnes EH, *et al*. Adjuvant chemotherapy following chemoradiotherapy as primary treatment for locally advanced cervical cancer versus chemoradiotherapy alone (OUTBACK): an international, open-label, randomised, phase 3 trial. *Lancet Oncol* 2023;24:468–82.
 - 132 Tewari KS, Sill MW, Long HJ III, *et al*. Improved survival with bevacizumab in advanced cervical cancer. *N Engl J Med* 2014;370:734–43.
 - 133 Colombo N, Dubot C, Lorusso D, *et al*. Pembrolizumab for persistent, recurrent, or metastatic cervical cancer. *N Engl J Med* 2021;385:1856–67.
 - 134 Macchia G, Nardangeli A, Laliscia C, *et al*. Stereotactic body radiotherapy in oligometastatic cervical cancer (MITO-RT2/RAD study): a collaboration of MITO, AIRO GYN, and MaNGO groups. *Int J Gynecol Cancer* 2022;32:732–9.
 - 135 Oaknin A, Gladieff L, Martínez-García J, *et al*. Atezolizumab plus bevacizumab and chemotherapy for metastatic, persistent, or recurrent cervical cancer (BEATcc): a randomised, open-label, phase 3 trial. *The Lancet* 2024;403:31–43.
 - 136 Oaknin A, Moore K, Meyer T, *et al*. Nivolumab with or without ipilimumab in patients with recurrent or metastatic cervical cancer (CheckMate 358): a phase 1-2, open-label, multicohort trial. *Lancet Oncol* 2024;25:588–602.
 - 137 Tewari KS, Monk BJ, Vergote I, *et al*. Survival with cemiplimab in recurrent cervical cancer. *N Engl J Med* 2022;386:544–55.
 - 138 Chung HC, Ros W, Delord J-P, *et al*. Efficacy and safety of pembrolizumab in previously treated advanced cervical cancer: results from the phase II KEYNOTE-158 study. *J Clin Oncol* 2019;37:1470–8.

- 139 Bonneville R, Paruchuri A, Wing MR, *et al.* Characterization of clonal evolution in microsatellite unstable metastatic cancers through multiregional tumor sequencing. *Mol Cancer Res* 2021;19:465–74.
- 140 Marabelle A, Fakih M, Lopez J, *et al.* Association of tumour mutational burden with outcomes in patients with advanced solid tumours treated with pembrolizumab: prospective biomarker analysis of the multicohort, open-label, phase 2 KEYNOTE-158 study. *Lancet Oncol* 2020;21:1353–65.
- 141 Noh JJ, Kim MK, Choi MC, *et al.* Frequency of mismatch repair deficiency/high microsatellite instability and its role as a predictive biomarker of response to immune checkpoint inhibitors in gynecologic cancers. *Cancer Res Treat* 2022;54:1200–8.
- 142 Coleman RL, Lorusso D, Gennigens C, *et al.* Efficacy and safety of tisotumab vedotin in previously treated recurrent or metastatic cervical cancer (innovaTV 204/GOG-3023/ENGOT-cx6): a multicentre, open-label, single-arm, phase 2 study. *Lancet Oncol* 2021;22:609–19.
- 143 Vergote I, González-Martín A, Fujiwara K, *et al.* Tisotumab Vedotin as Second- or Third-Line Therapy for Recurrent Cervical Cancer. *N E J Med* 2024;391:44–55.
- 144 Vergote I, Van Nieuwenhuysen E, O’Cearbhaill RE, *et al.* Tisotumab vedotin in combination with carboplatin, pembrolizumab, or bevacizumab in recurrent or metastatic cervical cancer: results from the innovaTV 205/GOG-3024/ENGOT-cx8 study. *J Clin Oncol* 2023;41:5536–49.
- 145 Meric-Bernstam F, Makker V, Oaknin A, *et al.* Efficacy and safety of trastuzumab deruxtecan in patients with HER2-expressing solid tumors: primary results from the DESTINY-PanTumor02 phase II trial. *J Clin Oncol* 2024;42:47–58.
- 146 Gatalica Z, Xiu J, Swensen J, *et al.* Molecular characterization of cancers with NTRK gene fusions. *Mod Pathol* 2019;32:147–53.
- 147 Consortium APG. AACR project GENIE: powering precision medicine through an international consortium. *Cancer Discov* 2017;7:818–31.
- 148 Hong J-H, Tsai C-S, Lai C-H, *et al.* Recurrent squamous cell carcinoma of cervix after definitive radiotherapy. *Int J Radiat Oncol Biol Phys* 2004;60:249–57.
- 149 Kobayashi K, Murakami N, Takahashi K, *et al.* Local radiotherapy or chemotherapy for oligo-recurrent cervical cancer in patients with prior pelvic irradiation. *In Vivo* 2019;33:1659–65.
- 150 Sardain H, Lavoue V, Redpath M, *et al.* Curative pelvic exenteration for recurrent cervical carcinoma in the era of concurrent chemotherapy and radiation therapy. a systematic review. *Eur J Surg Oncol* 2015;41:975–85.
- 151 Zola P, Macchi C, Cibula D, *et al.* Follow-up in gynecological malignancies: a state of art. *Int J Gynecol Cancer* 2015;25:1151–64.
- 152 Brooks RA, Rader JS, Dehdashti F, *et al.* Surveillance FDG-PET detection of asymptomatic recurrences in patients with cervical cancer. *Gynecol Oncol* 2009;112:104–9.
- 153 Elit L, Fyles AW, Oliver TK, *et al.* Follow-up for women after treatment for cervical cancer. *Curr Oncol* 2010;17:65–9.
- 154 Salani R, Khanna N, Frimer M, *et al.* An update on post-treatment surveillance and diagnosis of recurrence in women with gynecologic malignancies: society of gynecologic oncology (SGO) recommendations. *Gynecol Oncol* 2017;146:3–10.
- 155 Sartori E, Pasinetti B, Carrara L, *et al.* Pattern of failure and value of follow-up procedures in endometrial and cervical cancer patients. *Gynecol Oncol* 2007;107:S241–7.
- 156 Brown AJ, Shah JS, Fleming ND, *et al.* Role of cervical cytology in surveillance after radical trachelectomy for cervical cancer. *Gynecol Oncol* 2016;142:283–5.
- 157 Orr JM, Barnett JC, Leath CA. Incidence of subsequent abnormal cytology in cervical cancer patients completing five-years of post treatment surveillance without evidence of recurrence. *Gynecol Oncol* 2011;122:501–4.
- 158 Cibula D, Dostálek L, Jarkovsky J, *et al.* The annual recurrence risk model for tailored surveillance strategy inpatients with cervical cancer. *Eur J Cancer* 2021;158:111–22.
- 159 Suzuki Y, Huang Y, Ferris J, *et al.* Prescription of hormone replacement therapy among cervical cancer patients with treatment-induced premature menopause. *Int J Gynecol Cancer* 2023;33:26–34.
- 160 Hickey M, Basu P, Sassarini J, *et al.* Managing menopause after cancer. *Lancet* 2024;403:984–96.
- 161 Pötter R, Tanderup K, Schmid MP, *et al.* MRI-guided adaptive brachytherapy in locally advanced cervical cancer (EMBRACE-I): a multicentre prospective cohort study. *Lancet Oncol* 2021;22:538–47.
- 162 Huang H, Feng Y-L, Wan T, *et al.* Effectiveness of sequential chemoradiation vs concurrent chemoradiation or radiation alone in adjuvant treatment after hysterectomy for cervical cancer: the STARS phase 3 randomized clinical trial. *JAMA Oncol* 2021;7:361–9.
- 163 Landoni F, Maneo A, Colombo A, *et al.* Randomised study of radical surgery versus radiotherapy for stage Ib-IIa cervical cancer. *Lancet* 1997;350:535–40.
- 164 Stehman FB, Ali S, Keys HM, *et al.* Radiation therapy with or without weekly cisplatin for bulky stage 1B cervical carcinoma: follow-up of a gynecologic oncology group trial. *Am J Obstet Gynecol* 2007;197:503.
- 165 Lanciano R, Calkins A, Bundy BN, *et al.* Randomized comparison of weekly cisplatin or protracted venous infusion of fluorouracil in combination with pelvic radiation in advanced cervix cancer: a gynecologic oncology group study. *J Clin Oncol* 2005;23:8289–95.
- 166 Moore DH, Blessing JA, McQuellon RP, *et al.* Phase III study of cisplatin with or without paclitaxel in stage IVB, recurrent, or persistent squamous cell carcinoma of the cervix: a gynecologic oncology group study. *J Clin Oncol* 2004;22:3113–9.
- 167 Monk BJ, Sill MW, McMeekin DS, *et al.* Phase III trial of four cisplatin-containing doublet combinations in stage IVB, recurrent, or persistent cervical carcinoma: a gynecologic oncology group study. *J Clin Oncol* 2009;27:4649–55.
- 168 Kitagawa R, Katsumata N, Shibata T, *et al.* Paclitaxel plus carboplatin versus paclitaxel plus cisplatin in metastatic or recurrent cervical cancer: the open-label randomized phase III trial JCOG0505. *J Clin Oncol* 2015;33:2129–35.