

What Actually Causes Sleep Paralysis

Written by Sumbal Shiraz

Sleep paralysis is one of the most disorienting experiences the human body can produce. A person wakes up, or is about to fall asleep, and finds they cannot move, speak, or sometimes even open their eyes fully, while remaining completely conscious. For many, this alone is frightening. For others, it comes paired with a sensed presence in the room, pressure on the chest, or vivid, threatening hallucinations. Cultures across the world have given this experience different names, from the “night hag” in Western folklore to spiritual explanations rooted in regional belief systems. But behind the fear and the folklore is a well-documented neurological event with a clear physiological cause.

The Root Cause: A Timing Error in REM Sleep

During REM (rapid eye movement) sleep, the stage in which most vivid dreaming occurs, the brain deliberately paralyzes the body's skeletal muscles. This is not a malfunction. It is a protective mechanism. Without it, people would physically act out their dreams, which can lead to injury. The brainstem region primarily responsible for triggering this temporary paralysis, known as the subcoeruleus nucleus, sends inhibitory signals through the spinal cord that switch off voluntary muscle movement for the duration of REM sleep.

Sleep paralysis happens when consciousness resumes before this paralysis has fully switched off. In other words, the mind wakes up, but the body has not yet been given the signal to move again. The result is a brief but very real state in which a person is fully aware of their surroundings while their body remains locked in the muscular shutdown normally reserved for dreaming sleep.

This same REM-related mechanism explains the hallucinations that often accompany sleep paralysis. Because the brain is still partially in a dream-generating state even as the person becomes consciously aware, dream imagery and sensory distortions can bleed directly into waking perception. This is why so many sleep paralysis experiences involve a sensed intruder in the room or a feeling of pressure on the chest. The brain is essentially producing dream content while the eyes are open.

Who Is More Likely to Experience It

Sleep paralysis is not rare. Research estimates that roughly 7.6 percent of the general population experiences at least one episode in their lifetime, and the number rises sharply in specific groups. Among college and university students, studies place lifetime prevalence as high as 28.3 percent, a gap researchers attribute to the irregular sleep schedules, elevated stress, and disrupted sleep hygiene common in student life.

The clearest and most consistently replicated risk factors are:

- **Sleep deprivation and irregular sleep-wake schedules.** Shift workers and people who sleep at inconsistent times show measurably higher rates of sleep paralysis.
- **Anxiety and generalized worry.** Studies using standardized anxiety measures have found that people with higher trait anxiety experience sleep paralysis more frequently.
- **PTSD.** Multiple studies, including research on firefighters and nurses, found a significant association between PTSD symptoms and sleep paralysis frequency, with one study on firefighters finding that a probable PTSD diagnosis nearly doubled the odds of experiencing an episode.
- **Substance use.** Caffeine, alcohol, and nicotine use have each been linked to higher rates of sleep paralysis, likely through their disruptive effects on normal REM cycling.
- **Sleeping on the back.** While less rigorously established than the factors above, supine sleep position is frequently reported alongside sleep paralysis episodes in self-report studies.

Notably, research has found no meaningful difference in sleep paralysis prevalence by gender or age, which rules out some of the more common assumptions people make about who is “prone” to it.

Why It Feels So Threatening

The fear response during sleep paralysis is not incidental, it is built into the biology of the event itself. Because the amygdala, the brain's threat-detection center, remains active during REM sleep, a person regaining consciousness mid-paralysis is neurologically primed for fear before any hallucination even occurs. The combination of genuine physical immobility, an already-activated fear circuit, and intruding dream imagery creates a psychological experience that feels, in the moment, indistinguishable from real danger, even though the person is in no physical danger at all.

What Actually Helps

Sleep paralysis is classified as a parasomnia and, in most cases, is not itself a serious medical condition. It is typically treated by addressing the underlying contributors rather than the episodes directly. The most evidence-supported approaches include stabilizing sleep schedules, reducing chronic stress, limiting alcohol and caffeine intake, and treating any underlying anxiety or trauma-related condition. In cases where episodes are frequent, distressing, or linked to narcolepsy, a sleep specialist can assess whether further evaluation, such as a sleep study, is warranted.

Understanding sleep paralysis as a real, explainable neurological event, rather than something mystical or dangerous, is often the first and most effective step in reducing both the frequency of episodes and the fear that surrounds them.

Sources

Akhtar, Z.B. "Voices in the Night: Sleep Paralysis and the Intersection of Brain, Trauma, and Traditions." *ProBiologists*, 2025.

Denis, D., French, C.C., & Gregory, A.M. "A systematic review of variables associated with sleep paralysis." *Sleep Medicine Reviews*, 2018.

Jachowicz, A. et al. "Risk factors of sleep paralysis in a population of Polish students." *BMC Psychiatry*, 2022.

Jachowicz, A. et al. "Sleep Paralysis among Professional Firefighters and a Possible Association with PTSD." *International Journal of Environmental Research and Public Health*, 2021.

Jachowicz, A. et al. "Characteristics of Sleep Paralysis and Its Association with Anxiety Symptoms, Perceived Stress, PTSD, and Other Variables Related to Lifestyle in Selected High Stress Exposed Professions." *International Journal of Environmental Research and Public Health*, 2022.

Sharpless, B.A., & Barber, J.P. "Lifetime prevalence rates of sleep paralysis: A systematic review." *Sleep Medicine Reviews*, 2011.

StatPearls. "Sleep Paralysis." *StatPearls Publishing*, 2025.