

Affine 0-Schur algebras and affine double flag varieties.

Tom Crawley

October 22, 2020

Contents

1	Introduction	3
2	Background: The double flag variety approach to q-Schur algebras	4
3	Geometric approach to affine q-Schur algebras	5
3.1	Affine q -Schur algebras via Hecke algebras	5
3.2	The cyclic flags realisation of affine q -Schur algebras	5
3.2.1	Cyclic flags	6
3.2.2	A product of orbits	10
3.2.3	Triple products	10
3.2.4	Convolution algebras	11
3.2.5	Affine q -Schur algebras	12
3.3	Affine zero-Schur algebras	12
4	Presenting affine q-Schur algebras	13
4.1	The distinguished basis	13
4.1.1	Elementary basis elements	13
4.1.2	Transpose involution	14
4.1.3	Fundamental multiplication rules	15
4.1.4	The hook order	16
4.1.5	Shifting	19
4.2	Quivers and relations	21
4.2.1	Relations in affine q -Schur algebras	22
4.2.2	A quiver algebra	25
4.2.3	Mapping to the q -Schur algebra	26
4.2.4	Change of rings	27
4.3	Relations for the $n=2$ case	30
5	A generic affine algebra	32
5.1	Introduction	32
5.2	Grassmannians and related varieties	33
5.3	Geometry of affine flag varieties	35
5.3.1	Action through an algebraic group	36
5.3.2	Incidence in affine flag varieties	38
5.4	Geometry of orbits	39
5.5	Geometry of orbit products	40
5.6	Degenerations of orbits and the combinatorial partial order	41
5.7	Associativity of the generic product	42

5.8	The generic affine algebra	46
5.8.1	A categorical perspective	47
6	A realisation of affine zero Schur algebras	48
6.1	Preliminary results on the generic affine algebra	48
6.1.1	Elementary basis elements	48
6.1.2	Transpose involution	50
6.1.3	Shifting and periodicity	51
6.2	Multiplicative bases in affine zero Schur algebras: motivating example	53
6.3	Aperiodicity in the generic affine algebra	54
6.4	Quiver presentation of the generic affine algebra.	57
6.4.1	Standard paths	62
6.5	The isomorphism result	63
6.6	The period 2 case	65
7	Further directions	67
7.1	Further results on affine zero Schur algebras	67
7.2	Deformed group algebras of symmetric groups	67
7.3	back matter	67

Chapter 1

Introduction

Chapter 2

**Background: The double flag variety
approach to q -Schur algebras**

Chapter 3

Geometric approach to affine q-Schur algebras

3.1 Affine q-Schur algebras via Hecke algebras

Describe the original construction/ definition of affine q-Schur algebras via Hecke endomorphism algebras.

3.2 The cyclic flags realisation of affine q-Schur algebras

Fix integers $n, r \geq 1$.

Definition 3.2.1. A *composition of r into n parts* is an n -tuple $\lambda = (\lambda_1, \dots, \lambda_n) \in \mathbb{Z}^n$ of non-negative integers whose sum equals r . Denote the set of compositions of r into n parts by Λ_0 .

A composition $\lambda \in \Lambda_0$ is said to be *sincere* if $\lambda_i > 0$ for each $i \in \{1, \dots, n\}$ and otherwise λ is said to be *insincere*.

For each $i \in \{1, \dots, n\}$, let

$$\alpha_i = \varepsilon_i - \varepsilon_{i+1},$$

where $\varepsilon_{n+1} = \varepsilon_1$.

Definition 3.2.2. Let Λ_1 be the set of matrices $A = (a_{i,j})_{i,j \in \mathbb{Z}}$ with integer entries $a_{i,j}$ satisfying the following conditions:

- i. $a_{i,j} \geq 0$ for each $i, j \in \mathbb{Z}$;
- ii. each row or column has only finitely many non-zero entries;
- iii. the sum of the entries in any n consecutive rows or columns equals r ;
- iv. $a_{i-n,j-n} = a_{i,j}$ for each $i, j \in \mathbb{Z}$.

These matrices are referred to as *infinite periodic matrices*.

Definition 3.2.3. Given $A \in \Lambda_1$, let $\text{ro}(A)$ and $\text{co}(A)$ be the compositions of r into n parts given by

$$\text{ro}(A) = \left(\sum_{j \in \mathbb{Z}} a_{1,j}, \dots, \sum_{j \in \mathbb{Z}} a_{n,j} \right)$$

and

$$\text{co}(A) = \left(\sum_{i \in \mathbb{Z}} a_{i,1}, \dots, \sum_{i \in \mathbb{Z}} a_{i,n} \right).$$

The *source* of A is $\text{co}(A)$ and the *target* of A is $\text{ro}(A)$.

The row and column sums are finite since each row and column of A contains only finitely many nonzero entries, according to the definition of Λ_1 .

For each $i, j \in \mathbb{Z}$, let $\mathcal{E}_{i,j}$ be the $\mathbb{Z} \times \mathbb{Z}$ ‘elementary periodic matrix’ with entries given by

$$(\mathcal{E}_{i,j})_{s,t} = 1$$

if $(s, t) = (i + cn, j + cn)$ for some $c \in \mathbb{Z}$ and $(\mathcal{E}_{i,j})_{s,t} = 0$ otherwise. Clearly $\mathcal{E}_{i,j} = \mathcal{E}_{i+n,j+n}$ for each $i, j \in \mathbb{Z}$.

Given $\lambda \in \Lambda_0$, let $D_\lambda \in \Lambda_1$ be the diagonal matrix with source and target λ , which is given by

$$D_\lambda = \lambda_1 \mathcal{E}_{1,1} + \dots + \lambda_n \mathcal{E}_{n,n}. \quad (3.2.1)$$

3.2.1 Cyclic flags

Fix $n, r \in \mathbb{N}$ and let \mathbf{k} be a field. Let \mathcal{S} be the \mathbf{k} -algebra $\mathbf{k}[\varepsilon, \varepsilon^{-1}]$ and let \mathcal{R} be the subalgebra generated by ε , so $\mathcal{R} = \mathbf{k}[\varepsilon]$. Let V be a free \mathcal{S} -module of rank r . Let G be the automorphism group of the \mathcal{S} -module V , so G is isomorphic to $\text{GL}_r(\mathcal{S})$.

Definition 3.2.4. A *lattice* in V is a \mathcal{R} -submodule L of V with

$$\mathcal{S} \otimes_{\mathcal{R}} L = V.$$

In particular, a lattice is an \mathcal{R} -submodule of V which is a free \mathcal{R} -module of rank r .

Lemma 3.2.5. Let L be a lattice in V . $L/\varepsilon L$ is a torsion \mathcal{R} -module, where ε acts as zero and $L/\varepsilon L$ is an r -dimensional \mathbf{k} -vector space.

Proof. L is a free \mathcal{R} -module of rank r , with $L \subset V$. Given an \mathcal{R} -basis $\{x_1, \dots, x_r\}$ of L , $\{\varepsilon x_1, \dots, \varepsilon x_r\}$ is an \mathcal{R} -basis of εL . Finally, the cosets $\{x_1 + \varepsilon L, \dots, x_r + \varepsilon L\}$ give a basis for $L/\varepsilon L$ over $\mathcal{R}/\langle \varepsilon \rangle \cong \mathbf{k}$. \square

Observe that if L and L' are lattices in V , then both the sum $L + L'$ and the intersection $L \cap L'$ are lattices in V . Moreover,

$$\begin{aligned} g(L + L') &= g(L) + g(L') \\ g(L \cap L') &= g(L) \cap g(L') \end{aligned}$$

for each $g \in G$.

Definition 3.2.6. A *cyclic flag* in V is a collection $(L_i)_{i \in \mathbb{Z}}$ of lattices in V with $L_i \subset L_{i+1}$ and $\varepsilon L_i = L_{i-n}$ for each $i \in \mathbb{Z}$. The space of cyclic flags in V is denoted by $\mathcal{F} = \mathcal{F}(n, r)$.

The group $G = \text{Aut}(V)$ acts on \mathcal{F} by $(g \cdot L)_i = g(L_i)$ for each $i \in \mathbb{Z}$, $g \in G$ and $L \in \mathcal{F}$. The G -orbits in \mathcal{F} are indexed by the set Λ_0 of compositions of r into n parts. In particular, the G -orbit in \mathcal{F} corresponding to $\lambda \in \Lambda_0$ is

$$\mathcal{F}_\lambda = \left\{ L \in \mathcal{F} : \dim \left(\frac{L_i}{L_{i-1}} \right) = \lambda_i \text{ for each } i \in \mathbb{Z} \right\}.$$

Consider the space $\mathcal{F} \times \mathcal{F}$ of pairs of flags with the diagonal action of G , given by

$$g \cdot (L, L') = (gL, gL')$$

for $g \in G$ and $(L, L') \in \mathcal{F} \times \mathcal{F}$. Denote the G -orbit of (L, L') by $[L, L']$. The set of G -orbits in $\mathcal{F} \times \mathcal{F}$ admits a combinatorial description as will be described below.

Definition 3.2.7. The periodic characteristic matrix of a pair of cyclic flags $(L, L') \in \mathcal{F} \times \mathcal{F}$ is the matrix $A(L, L') = (a_{i,j})_{i,j \in \mathbb{Z}}$ with entries

$$a_{i,j} = \dim \left(\frac{L_i \cap L'_j}{L_i \cap L'_{j-1} + L_{i-1} \cap L'_j} \right)$$

for each $i, j \in \mathbb{Z}$.

The next lemma shows that the characteristic matrix of a pair of flags depends only on the G -orbit of the pair.

Lemma 3.2.8. Given $(L, L') \in \mathcal{F} \times \mathcal{F}$ and $g \in G$,

$$A(gL, gL') = A(L, L').$$

Proof. Write $A = A(L, L')$ and $B = A(gL, gL')$. For each $i, j \in \mathbb{Z}$, g induces a linear isomorphism

$$\frac{L_i \cap L'_j}{L_{i-1} \cap L'_j + L_i \cap L'_{j-1}} \rightarrow \frac{g(L_i \cap L'_j)}{g(L_{i-1} \cap L'_j + L_i \cap L'_{j-1})},$$

so

$$\begin{aligned} b_{i,j} &= \dim \left(\frac{gL_i \cap gL'_j}{gL_{i-1} \cap gL'_j + gL_i \cap gL'_{j-1}} \right) \\ &= \dim \left(\frac{g(L_i \cap L'_j)}{g(L_{i-1} \cap L'_j + L_i \cap L'_{j-1})} \right) \\ &= \dim \left(\frac{L_i \cap L'_j}{L_{i-1} \cap L'_j + L_i \cap L'_{j-1}} \right) \\ &= a_{i,j}, \end{aligned}$$

since the action of g commutes with sums and intersections of lattices. Therefore $A = B$ as claimed. \square

The following result gives another useful set of expressions for the characteristic matrix.

Lemma 3.2.9. For each $i, j \in \mathbb{Z}$,

$$a_{i,j} = \dim \left(\frac{L_{i-1} + L_i \cap L'_j}{L_{i-1} + L_i \cap L'_{j-1}} \right).$$

Proof. Set $U = L_i \cap L'_j$ and $U' = L_{i-1} + L_i \cap L'_{j-1}$. Then $U + U' = L_{i-1} + L_i \cap L'_j$ and $U \cap U' = L_i \cap L'_j \cap L_{i-1} + L_i \cap L'_{j-1}$. Applying the isomorphism theorems, $U + U'/U'$ is naturally isomorphic to $U/U \cap U'$ as a vector space. In particular,

$$\frac{L_{i-1} + L_i \cap L'_j}{L_{i-1} + L_i \cap L'_{j-1}} = \frac{L_i \cap L'_j}{L_{i-1} \cap L'_j + L_i \cap L'_{j-1}}$$

and thus the dimensions of these spaces are both equal to $a_{i,j}$. \square

Lemma 3.2.10. For each $(L, L') \in \mathcal{F} \times \mathcal{F}$, $A(L, L') \in \Lambda_1$.

Proof. Let $A = A(L, L')$, fix $\lambda \in \Lambda_0$ such that $L \in \mathcal{F}_\lambda$ and let $i \in \{1, \dots, n\}$. **COMPLETE THIS PROOF** □

Finish giving a complete and rigorous description of the orbit-matrix correspondence. It is fairly elementary but used extensively in the rest of the thesis, so it is worth giving a full account.

Lemma 3.2.11. Given a pair of flags $(L, L') \in \mathcal{F}^2$, the matrices $A(L, L')$ and $A(L', L)$ are related by the transpose. In particular, $A(L, L')_{i,j} = A(L', L)_{j,i}$ for each $i, j \in \mathbb{Z}$.

Proof. By swapping the roles of i and j and swapping L and L' it is clear that $A(L, L')_{i,j}$ and $A(L', L)_{j,i}$ are both equal to the dimension of the \mathbf{k} -vector space

$$\frac{L_i \cap L'_j}{L_{i-1} \cap L'_j + L_i \cap L'_{j-1}},$$

for each $i, j \in \mathbb{Z}$. □

Lemma 3.2.12. Given $A \in \Lambda_1$ and $(L, L') \in \mathcal{O}_A$,

$$\dim_{\mathbf{k}} \left(\frac{L_i}{L_i \cap L'_j} \right) = \sum_{s \leq i, t > j} a_{s,t}$$

and

$$\dim_{\mathbf{k}} \left(\frac{L'_j}{L_i \cap L'_j} \right) = \sum_{s > i, t \leq j} a_{s,t},$$

for each $i, j \in \mathbb{Z}$.

Proof. For each $s, t \in \mathbb{Z}$ define a lattice

$$M_{s,t} = L_i \cap L'_j + L_{s-1} + L_s \cap L'_t.$$

Observe that $L_i \cap L'_j$ is a sublattice of each $M_{s,t}$ and when $s \leq i$, L_s is contained in L_i , so $M_{s,t}$ is a sublattice of L_i . The collection of lattices $(M_{s,t})_{s \leq i, t > j}$ are totally ordered by subset inclusion, as will be shown below, so give a chain of lattices each containing $L_i \cap L'_j$ and contained in L_i . This chain of lattices induces a filtration of $L_i/L_i \cap L'_j$ and it will be shown that the dimensions of the quotients are precisely $a_{s,t}$ for $s \leq i$ and $t > j$.

Let $s \leq i$ and $t \geq j$.

$$M_{s,t} \subset M_{s,t+1}$$

and

$$\begin{aligned} M_{s,j} &= L_i \cap L'_j + L_{s-1} + L_s \cap L'_j \\ &= L_i \cap L'_j + L_{s-1}. \end{aligned}$$

If t is sufficiently large, $L_s \subset L'_t$ and then

$$\begin{aligned} M_{s,t} &= L_i \cap L'_j + L_s \\ &= M_{s+1,j}. \end{aligned}$$

It follows that the collection of lattices is totally ordered, with $M_{s,t} \leq M_{u,v}$ if and only if $s < u$ or $s = u$ and $t \leq v$. Thus $L_i/L_i \cap L'_j$ has a filtration given by the spaces $M_{s,t}/L_i \cap L'_j$ for all $s \leq i$ and $t > j$.

$$\begin{aligned} M_{i,t} &= L_i \cap L'_j + L_{i-1} + L_i \cap L'_t \\ &= L_{i-1} + L_i \cap L'_t \end{aligned}$$

and if t is sufficiently large that $L_i \subset L'_t$, then $M_{i,t} = L_i$.

If s is sufficiently small that $L_s \subset L_i \cap L'_j$ then

$$\begin{aligned} M_{s,t} &= L_i \cap L'_j + L_{s-1} + L_s \cap L'_t \\ &= L_i \cap L'_j. \end{aligned}$$

$$\begin{aligned} \frac{M_{s,t}}{L_i \cap L'_j} &= \frac{L_i \cap L'_j + L_{s-1} + L_s \cap L'_t}{L_i \cap L'_j} \\ &= \frac{L_{s-1} + L_s \cap L'_t}{L_i \cap L'_j \cap (L_{s-1} + L_s \cap L'_t)} \\ &= \frac{L_{s-1} + L_s \cap L'_t}{L_s \cap L'_j} \end{aligned}$$

Then for each $s \leq i$ and $t > j$

$$\begin{aligned} \dim \left(\frac{M_{s,t}}{M_{s,t-1}} \right) &= \dim \left(\frac{L_{s-1} + L_s \cap L'_t}{L_{s-1} + L_s \cap L'_{t-1}} \right) \\ &= a_{s,t}, \end{aligned}$$

so

$$\dim \left(\frac{L_i}{L_i \cap L'_j} \right) = \sum_{s \leq i, t > j} a_{s,t}.$$

To deduce the second formula observe that $(L', L) \in \mathcal{O}_{A^\top}$, by Lemma 3.2.11, so

$$\begin{aligned} \dim \left(\frac{L'_j}{L_i \cap L'_j} \right) &= \sum_{t \leq j, s > i} a_{t,s}^\top \\ &= \sum_{s > i, t \leq j} a_{s,t}. \end{aligned}$$

□

Lemma 3.2.13. *Given $(L, L') \in \mathcal{F}^2$, $L' \subset L$ if and only if $A(L, L')_{i,j} = 0$ for $i, j \in \mathbb{Z}$ with $i > j$.*

Proof. Suppose $L, L' \in \mathcal{F}$ with $L' \subset L$, meaning $L'_j \subset L_j$ for each $j \in \mathbb{Z}$. Then for $i > j$, $L_i \cap L'_j = L'_j$, $L_{i-1} \cap L'_j = L'_j$ and $L_i \cap L'_{j-1}$, which shows

$$A(L, L')_{i,j} = \dim \left(\frac{L'_j}{L'_{j-1} + L'_j} \right) = 0$$

as required. Conversely, suppose $A(L, L')$ is upper triangular, meaning $A(L, L')_{i,j} = 0$ when $i > j$. Using Lemma 3.2.12,

$$\dim \left(\frac{L'_i}{L'_i \cap L_i} \right) = \sum_{s>i, t \leq i} a_{s,t} = 0,$$

so $L_i \cap L'_i = L'_i$ and thus $L'_i \subset L_i$ for each $i \in \mathbb{Z}$, as required. \square

Corollary 3.2.14. *Given $L, L' \in \mathcal{F}$, $L = L'$ if and only if $A(L, L')_{i,j} = 0$ whenever $i \neq j$. In particular,*

$$\mathcal{O}_{D_\lambda} = \{(L, L) \in \mathcal{F}^2 : L \in \mathcal{F}_\lambda\},$$

for each $\lambda \in \Lambda_0$.

3.2.2 A product of orbits

Given $A, B \in \Lambda_1$ with $\text{co}(A) = \text{ro}(B)$, define

$$Y_{A,B} = \{(L, L', L'') \in \mathcal{F}^3 : (L, L') \in \mathcal{O}_A \text{ and } (L', L'') \in \mathcal{O}_B\},$$

$$X_{A,B} = \{(L, L'') \in \mathcal{F}^2 : \exists L' \in \mathcal{F} \text{ with } (L, L') \in \mathcal{O}_A \text{ and } (L', L'') \in \mathcal{O}_B\}.$$

If also $L \in \mathcal{F}_{\text{ro}(A)}$, define the L -slices of $Y_{A,B}$ and $X_{A,B}$ respectively as

$$Y_{A,B}^L = \{(L', L'') \in \mathcal{F}^2 : (L, L', L'') \in Y_{A,B}\},$$

$$X_{A,B}^L = \{L'' \in \mathcal{F} : (L, L'') \in X_{A,B}\}.$$

Remark 3.2.15. There are only finitely many G -orbits in $X_{A,B}$.

Lemma 3.2.16. *Given $A \in \Lambda_1$, $X_{D_\lambda, A} = \mathcal{O}_A$ if $\lambda = \text{ro}(A)$ and $X_{A, D_\lambda} = \mathcal{O}_A$ if $\lambda = \text{co}(A)$.*

Proof. Let $A \in \Lambda_1$ and set $\lambda = \text{ro}(A)$. $Y_{D_\lambda, A}$ is the set of triples $(L, L', L'') \in \mathcal{F}^3$ with $(L, L') \in \mathcal{O}_{D_\lambda}$, thus $L = L'$ by Corollary 3.2.14, and $(L', L'') \in \mathcal{O}_A$. $X_{D_\lambda, A}$ is the projection of $Y_{D_\lambda, A}$, which equals \mathcal{O}_A .

Similarly, if $\lambda = \text{co}(A)$, Y_{A, D_λ} is the set of triples $(L, L', L'') \in \mathcal{F}^3$ with $(L, L') \in \mathcal{O}_A$ and $L'' = L'$, so X_{A, D_λ} is exactly the orbit \mathcal{O}_B . \square

3.2.3 Triple products

Given $A, B, C \in \Lambda_1$ with $\text{co}(A) = \text{ro}(B)$ and $\text{co}(B) = \text{ro}(C)$ and $L \in \mathcal{F}_{\text{ro}(A)}$, there are spaces $X_{A,B,C}$, $Y_{A,B,C}$ and their respective L -slices, defined as follows:

$$Y_{A,B,C} = \{(L, L', L'', L''') \in \mathcal{F}^4 : (L, L') \in \mathcal{O}_A, (L', L'') \in \mathcal{O}_B \text{ and } (L'', L''') \in \mathcal{O}_C\},$$

$$X_{A,B,C} = \{(L, L''') \in \mathcal{F}^2 : \exists (L', L'') \in \mathcal{O}_B \text{ with } (L, L') \in \mathcal{O}_A \text{ and } (L'', L''') \in \mathcal{O}_C\},$$

$$Y_{A,B,C}^L = \{(L', L'', L''') \in \mathcal{F}^3 : (L, L', L'', L''') \in Y_{A,B,C}\},$$

$$X_{A,B,C}^L = \{L''' \in \mathcal{F} : (L, L''') \in X_{A,B,C}\}.$$

3.2.4 Convolution algebras

Suppose \mathbf{k} is a finite field and let q denote the number of elements of \mathbf{k} . Consider the set S of G -invariant functions $\mathcal{F} \times \mathcal{F} \rightarrow \mathbb{Z}$ with constructible support. S is a free \mathbb{Z} -module with a basis consisting of the indicator functions of the G -orbits in $\mathcal{F} \times \mathcal{F}$. Define an operation \star on S as follows: for each $f, g \in S$, $f \star g \in S$ is given by

$$(f \star g)(L, L'') = \sum_{L' \in \mathcal{F}} f(L, L')g(L', L''),$$

for $(L, L'') \in \mathcal{F} \times \mathcal{F}$.

$f \star g$ is well defined since the supports of f and g consist of finitely many G -orbits, so there are only finitely many $L' \in \mathcal{F}$ such that $f(L, L')g(L', L'') \neq 0$, given $(L, L'') \in \mathcal{F} \times \mathcal{F}$. $f \star g$ is constant on G -orbits and is supported on finitely many G -orbits, so $f \star g \in S$.

Lemma 3.2.17. *The set S together with the operation \star is an associative \mathbb{Z} -algebra with identity element ι given by $\iota(L, L) = 1$ and $\iota(L, L') = 0$ for $L' \neq L$.*

Proof. Given $f, g, h \in S$ and $(L, L''') \in \mathcal{F} \times \mathcal{F}$,

$$\begin{aligned} ((f \star g) \star h)(L, L''') &= \sum_{L''} (f \star g)(L, L'')h(L'', L''') \\ &= \sum_{L''} \sum_{L'} f(L, L')g(L', L'')h(L'', L''') \\ &= (f \star (g \star h))(L, L'''), \end{aligned}$$

thus \star is associative. ι is the multiplicative identity since

$$(\iota \star f)(L, L'') = \sum_{L'} \iota(L, L')f(L', L'') = f(L, L'')$$

and

$$(f \star \iota)(L, L'') = \sum_{L'} f(L, L')\iota(L', L'') = f(L, L''),$$

for each $f \in S$ and $(L, L'') \in \mathcal{F} \times \mathcal{F}$. □

Given $A \in \Lambda_1$, let $e_A \in S$ denote the indicator function of the orbit \mathcal{O}_A . S is a free \mathbb{Z} -module with basis $\{e_A : A \in \Lambda_1\}$. There exist $\gamma_{A,B,C;q} \in \mathbb{Z}$ for $A, B, C \in \Lambda_1$ such that

$$e_A \star e_B = \sum_{C \in \Lambda_1} \gamma_{A,B,C;q} e_C$$

for each $A, B \in \Lambda_1$. Then

$$\begin{aligned} \gamma_{A,B,C;q} &= (e_A \star e_B)(L, L'') \\ &= \sum_{L'} e_A(L, L')e_B(L', L'') \\ &= \#\{L' : (L, L') \in \mathcal{O}_A \text{ and } (L', L'') \in \mathcal{O}_B\}, \end{aligned}$$

for any $(L, L'') \in \mathcal{O}_C$.

3.2.5 Affine q-Schur algebras

There exist polynomials $\gamma_{A,B,C} \in \mathbb{Z}[q]$ for $A, B, C \in \Lambda_1$ such that $\gamma_{A,B,C}(q) = \gamma_{A,B,C;q}$ for any prime power q , following [28, section 4]. The affine q -Schur algebra $\hat{S}_q(n, r)$ is a $\mathbb{Z}[q]$ -algebra which is a free $\mathbb{Z}[q]$ -module with basis $\{e_A : A \in \Lambda_1\}$ and with multiplication given by

$$e_A e_B = \sum_C \gamma_{A,B,C} e_C.$$

Given the existence of these ‘universal polynomials’ $\gamma_{A,B,C} \in \mathbb{Z}[q]$, it follows from Lemma 3.2.17 that $\hat{S}_q(n, r)$ is an associative $\mathbb{Z}[q]$ -algebra with multiplicative identity given by

$$1 = \sum_{\lambda \in \Lambda_0} e_{D_\lambda}.$$

3.3 Affine zero-Schur algebras

Fix integers $n, r \geq 1$.

Definition 3.3.1. The affine zero Schur algebra $\hat{S}_0(n, r)$ is the \mathbb{Z} -algebra

$$\hat{S}_0(n, r) = \mathbb{Z}[q]/(q) \otimes_{\mathbb{Z}[q]} \hat{S}_q(n, r).$$

Observe that $\hat{S}_0(n, r)$ is a free \mathbb{Z} -module, since $\hat{S}_q(n, r)$ is a free $\mathbb{Z}[q]$ -module. Moreover, $\hat{S}_0(n, r)$ has a \mathbb{Z} -basis $\{e_A : A \in \Lambda_1\}$ with multiplication given by

$$e_A \cdot e_B = \sum_{C \in \Lambda_1} \gamma_{A,B,C}(0) e_C,$$

for each $A, B \in \Lambda_1$.

Chapter 4

Presenting affine q-Schur algebras

4.1 The distinguished basis

4.1.1 Elementary basis elements

Recall that $\mathcal{E}_{i,j}$, for $i, j \in \mathbb{Z}$ is the $\mathbb{Z} \times \mathbb{Z}$ elementary periodic matrix, given by

$$(\mathcal{E}_{i,j})_{s,t} = 1$$

if $(s, t) = (i + cn, j + cn)$ for some $c \in \mathbb{Z}$ and $(\mathcal{E}_{i,j})_{s,t} = 0$ otherwise.

Recall that the diagonal matrix with source and target λ is

$$D_\lambda = \lambda_1 \mathcal{E}_{1,1} + \cdots + \lambda_n \mathcal{E}_{n,n},$$

as in Equation 3.2.1.

The corresponding basis elements e_{D_λ} , for $\lambda \in \Lambda_0$, are pairwise orthogonal idempotents in $\hat{S}_q(n, r)$ with

$$\sum_{\lambda \in \Lambda_0} e_{D_\lambda} = 1,$$

as a result of Lemma 3.2.16.

For each $i \in \{1, \dots, n\}$ and $\lambda \in \Lambda_0$ with $\lambda_{i+1} > 0$, define

$$E_{i,\lambda} = e_{D_\lambda + \mathcal{E}_{i,i+1} - \mathcal{E}_{i+1,i+1}}$$

and define

$$E_i = \sum_{\lambda \in \Lambda_0: \lambda_{i+1} > 0} E_{i,\lambda}.$$

Also define, for each $i \in \{1, \dots, n\}$ and $\lambda \in \Lambda_0$ with $\lambda_i > 0$,

$$F_{i,\lambda} = e_{D_\lambda + \mathcal{E}_{i+1,i} - \mathcal{E}_{i,i}}$$

and define

$$F_i = \sum_{\lambda \in \Lambda_0: \lambda_i > 0} F_{i,\lambda}.$$

For each $i \in \{1, \dots, n\}$, let $\alpha_i = \varepsilon_i - \varepsilon_{i+1}$. Then

$$\text{co}(E_{i,\lambda}) = \text{co}(F_{i,\lambda}) = \lambda,$$

$$\text{ro}(E_{i,\lambda}) = \lambda + \alpha_i$$

and

$$\text{ro}(F_{i,\lambda}) = \lambda - \alpha_i.$$

4.1.2 Transpose involution

Let S be the $\mathbb{Z}[q]$ -module automorphism of $\hat{S}_q(n, r)$ given by

$$S(e_A) = e_{A^\top},$$

for each $A \in \Lambda_1$.

Lemma 4.1.1. *The map S is a $\mathbb{Z}[q]$ -algebra antihomomorphism of order 2. In particular,*

$$S(e_A e_B) = S(e_B) S(e_A)$$

for each $A, B \in \Lambda_1$.

Proof. Let $A, B, C \in \Lambda_1$ and let \mathbf{k} be a finite field with $q = \#\mathbf{k}$ elements. If $(L, L'') \in \mathcal{O}_C$ then $(L'', L) \in \mathcal{O}_{C^\top}$ and

$$\begin{aligned} \gamma_{A,B,C;q} &= \#\{L' : (L, L') \in \mathcal{O}_A \text{ and } (L', L'') \in \mathcal{O}_B\} \\ &= \#\{L' : (L'', L') \in \mathcal{O}_{B^\top} \text{ and } (L', L) \in \mathcal{O}_{A^\top}\} \\ &= \gamma_{B^\top, A^\top, C^\top;q} \end{aligned}$$

It follows that

$$S(e_A e_B) = S(e_B) S(e_A),$$

for each $A, B \in \Lambda_1$ and therefore S is a $\mathbb{Z}[q]$ -algebra antihomomorphism. Moreover, $S \circ S$ is the identity map on $\hat{S}_q(n, r)$ since $(A^\top)^\top = A$. \square

The action of S on E_i , F_i and 1_λ is as follows:

$$S(1_\lambda) = 1_\lambda$$

for each $\lambda \in \Lambda_0$,

$$S(E_{i,\lambda}) = F_{i,\lambda+\alpha_i}$$

for each $i \in \{1, \dots, n\}$ and $\lambda \in \Lambda_0$ with $\lambda_{i+1} > 0$, and

$$S(F_{i,\lambda}) = E_{i,\lambda-\alpha_i}$$

for each $i \in \{1, \dots, n\}$ and $\lambda \in \Lambda_0$ with $\lambda_i > 0$.

In particular,

$$\begin{aligned} S(E_i) &= F_i, \\ S(F_i) &= E_i, \\ S(1_\lambda) &= 1_\lambda \end{aligned}$$

for $i \in \{1, \dots, n\}$ and $\lambda \in \Lambda_0$.

4.1.3 Fundamental multiplication rules

For each $m \in \mathbb{N}$, define the q -integer $[[m]] \in \mathbb{Z}[q]$ by

$$[[m]] = \frac{1 - q^m}{1 - q},$$

so that

$$\begin{aligned} [[0]] &= 0 \\ [[1]] &= 1 \\ [[2]] &= 1 + q \\ [[3]] &= 1 + q + q^2 \end{aligned}$$

and

$$[[m]] = 1 + q + \dots + q^{m-1}$$

for $m \geq 1$.

Lemma 4.1.2. *Given $A \in \Lambda_1$ and $i \in \{1, \dots, n\}$ with $\text{ro}(A)_{i+1} > 0$,*

$$E_i e_A = \sum_{p \in \mathbb{Z}: a_{i+1,p} > 0} q^{\sum_{j>p} a_{i,j}} [[a_{i,p} + 1]] e_{A+\mathcal{E}_{i,p}-\mathcal{E}_{i+1,p}}.$$

Given $A \in \Lambda_1$ and $i \in \{1, \dots, n\}$ with $\text{ro}(A)_i > 0$,

$$F_i e_A = \sum_{p \in \mathbb{Z}: a_{i,p} > 0} q^{\sum_{j<p} a_{i+1,j}} [[a_{i+1,p} + 1]] e_{A+\mathcal{E}_{i+1,p}-\mathcal{E}_{i,p}}.$$

Note that these formulas are still valid in the cases $E_i e_A = 0$ and $F_i e_A = 0$. If the convention that $e_B = 0$ whenever B is not in Λ_1 is used, then the conditions on p in the above sums may be ignored.

Corollary 4.1.3. *Given $A \in \Lambda_1$ and $j \in \{1, \dots, n\}$ with $\text{co}(A)_{j+1} > 0$,*

$$e_A F_j = \sum_{p \in \mathbb{Z}: a_{p,j+1} > 0} q^{\sum_{i>p} a_{i,j}} [[a_{p,j} + 1]] e_{A+\mathcal{E}_{p,j}-\mathcal{E}_{p,j+1}}.$$

Given $A \in \Lambda_1$ and $j \in \{1, \dots, n\}$ with $\text{co}(A)_j > 0$,

$$e_A E_j = \sum_{p \in \mathbb{Z}: a_{p,j} > 0} q^{\sum_{i<p} a_{i,j+1}} [[a_{p,j+1} + 1]] e_{A+\mathcal{E}_{p,j+1}-\mathcal{E}_{p,j}}.$$

Proof.

$$\begin{aligned} e_A F_j &= S(E_j e_{A^\top}) \\ &= S \left(\sum_{p \in \mathbb{Z}: a_{p,j+1} > 0} q^{\sum_{i>p} a_{i,j}} [[a_{p,j} + 1]] e_{A^\top + \mathcal{E}_{j,p} - \mathcal{E}_{j+1,p}} \right) \\ &= \sum_{p \in \mathbb{Z}: a_{p,j+1} > 0} q^{\sum_{i>p} a_{i,j}} [[a_{p,j} + 1]] e_{A+\mathcal{E}_{p,j}-\mathcal{E}_{p,j+1}}, \end{aligned}$$

where the second equality comes from Lemma 4.1.2. Similarly,

$$\begin{aligned}
e_A E_j &= S(F_j e_{A^\top}) \\
&= S\left(\sum_{p \in \mathbb{Z}: a_{p,j} > 0} q^{\sum_{i < p} a_{i,j+1}} [[a_{p,j+1} + 1]] e_{A^\top + \mathcal{E}_{j+1,p} - \mathcal{E}_{j,p}}\right) \\
&= \sum_{p \in \mathbb{Z}: a_{p,j} > 0} q^{\sum_{i < p} a_{i,j+1}} [[a_{p,j+1} + 1]] e_{A + \mathcal{E}_{p,j+1} - \mathcal{E}_{p,j}}.
\end{aligned}$$

□

4.1.4 The hook order

For each $i, j \in \mathbb{Z}$, let $d_{i,j}$ and $\bar{d}_{i,j}$ be the maps from Λ_1 to \mathbb{Z} given by

$$d_{i,j}(A) = \sum_{s \leq i, t > j} a_{s,t}$$

and

$$\bar{d}_{i,j}(A) = \sum_{s > i, t \leq j} a_{s,t}$$

for each $A \in \Lambda_1$.

Lemma 4.1.4. *For each $A \in \Lambda_1$ and $i, j \in \mathbb{Z}$, the following equations hold:*

$$\begin{aligned}
d_{i,j}(A) - d_{i-1,j}(A) &= \sum_{t > j} a_{i,t} \\
d_{i,j}(A) - d_{i,j-1}(A) &= -\sum_{s \leq i} a_{s,j}
\end{aligned}$$

and

$$\begin{aligned}
\bar{d}_{i,j}(A) - \bar{d}_{i,j-1}(A) &= -\sum_{t \leq j} a_{i,t} \\
\bar{d}_{i,j}(A) - \bar{d}_{i-1,j}(A) &= \sum_{s > i} a_{s,j}
\end{aligned}$$

Proof. Let $i, j \in \mathbb{Z}$ and $A \in \Lambda_1$. Then

$$d_{i,j}(A) - d_{i-1,j}(A) = \sum_{s \leq i, t > j} a_{s,t} - \sum_{s \leq i-1, t > j} a_{s,t} = \sum_{t > j} a_{i,t}$$

and

$$d_{i,j}(A) - d_{i,j-1}(A) = \sum_{s \leq i, t > j} a_{s,t} - \sum_{s \leq i, t > j-1} a_{s,t} = -\sum_{s \leq i} a_{s,j}.$$

Similarly,

$$\bar{d}_{i,j}(A) - \bar{d}_{i-1,j}(A) = \sum_{s > i, t \leq j} a_{s,t} - \sum_{s > i-1, t \leq j} a_{s,t} = -\sum_{t \leq j} a_{i,t}$$

and

$$\bar{d}_{i,j}(A) - \bar{d}_{i,j-1}(A) = \sum_{s > i, t \leq j} a_{s,t} - \sum_{s > i, t \leq j-1} a_{s,t} = \sum_{s > i} a_{s,j}.$$

□

Lemma 4.1.5. For each $A \in \Lambda_1$ and $i, j \in \mathbb{Z}$,

$$a_{i,j} = d_{i,j-1}(A) - d_{i-1,j-1}(A) - d_{i,j}(A) + d_{i-1,j}(A)$$

and

$$a_{i,j} = \bar{d}_{i,j-1}(A) - \bar{d}_{i-1,j-1}(A) - \bar{d}_{i,j}(A) + \bar{d}_{i-1,j}(A).$$

Proof. As a result of Lemma 4.1.4,

$$\begin{aligned} d_{i,j-1}(A) - d_{i-1,j-1}(A) - d_{i,j}(A) + d_{i-1,j}(A) &= \sum_{t>j-1} a_{i,t} - \sum_{t>j} a_{i,t} \\ &= a_{i,j} \end{aligned}$$

and

$$\begin{aligned} \bar{d}_{i,j-1}(A) - \bar{d}_{i-1,j-1}(A) - \bar{d}_{i,j}(A) + \bar{d}_{i-1,j}(A) &= - \sum_{t \leq j-1} a_{i,t} + \sum_{t \leq j} a_{i,t} \\ &= a_{i,j}. \end{aligned}$$

□

Define a relation \leq on Λ_1 by $A \leq B$ if and only if the following conditions are satisfied:

- $\text{ro}(A) = \text{ro}(B)$ and $\text{co}(A) = \text{co}(B)$.
- For each $i, j \in \mathbb{Z}$, $d_{i,j}(A) \leq d_{i,j}(B)$.
- For each $i, j \in \mathbb{Z}$, $\bar{d}_{i,j}(A) \leq \bar{d}_{i,j}(B)$.

Lemma 4.1.6. The relation \leq defines a partial order on Λ_1 .

Proof. It is clear that \leq is reflexive and transitive.

Suppose $A, B \in \Lambda_1$ with $A \leq B$ and $B \leq A$. Then $d_{i,j}(A) = d_{i,j}(B)$ for each $i, j \in \mathbb{Z}$ with $i \leq j$, which shows $a_{s,t} = b_{s,t}$ whenever $s < t$, as a result of Lemma 4.1.5. Similarly, $\bar{d}_{i,j}(A) = \bar{d}_{i,j}(B)$ for each $i, j \in \mathbb{Z}$ with $i \geq j$, so $a_{s,t} = b_{s,t}$ whenever $s > t$. Moreover, $a_{i,i} = b_{i,i}$ for each $i \in \mathbb{Z}$, since $\text{co}(A) = \text{co}(B)$. Thus $A = B$, which shows \leq is antisymmetric and therefore \leq is a partial order on Λ_1 . □

This partial order is sometimes called the hook order. The following lemma will be invoked later in induction arguments.

Lemma 4.1.7. For any $A \in \Lambda_1$, the set $\{B \in \Lambda_1 : B \leq A\}$ is finite.

Proof. Let $B \in \Lambda_1$. Only finitely many of the $d_{i,j}(B)$ and $\bar{d}_{i,j}(B)$ are sufficient to determine B and $B \leq A$ if and only if

$$0 \leq d_{i,j}(B) \leq d_{i,j}(A)$$

and

$$0 \leq \bar{d}_{i,j}(B) \leq \bar{d}_{i,j}(A)$$

for each $i, j \in \mathbb{Z}$, thus there are only finitely many possible values of $d_{i,j}(B)$ and $\bar{d}_{i,j}(B)$ provided $B \leq A$. Therefore there are only finitely many $B \in \Lambda_1$ such that $B \leq A$. □

Lemma 4.1.8. *The transpose operation on Λ_1 is order preserving. In particular, $B \leq A$ if and only if $B^\top \leq A^\top$.*

Proof. Suppose $A, B \in \Lambda_1$ with $B \leq A$. The condition $\text{co}(A) = \text{co}(B)$ and $\text{ro}(A) = \text{ro}(B)$ is preserved by the transpose operation.

For each $i, j \in \mathbb{Z}$,

$$d_{i,j}(A^\top) = \sum_{s \leq i, t > j} a_{t,s} = \bar{d}_{j,i}(A)$$

and

$$\bar{d}_{i,j}(A^\top) = \sum_{s > i, t \leq j} a_{t,s} = d_{j,i}(A).$$

It follows that $B^\top \leq A^\top$ and therefore the transpose is order preserving. \square

Lemma 4.1.9. *Suppose $A, B \in \Lambda_1$ with*

$$B = A + \mathcal{E}_{i,j} - \mathcal{E}_{s,j} + \mathcal{E}_{s,t} - \mathcal{E}_{i,t}$$

for some $i, j, s, t \in \mathbb{Z}$ with $i < s$ and $j < t$. Then $B < A$.

Proof. Let $p, q \in \mathbb{Z}$. Then

$$d_{p,q}(B) = \begin{cases} d_{p,q}(A) - 1 & : i \leq p < s \text{ and } j \leq q < t, \\ d_{p,q}(A) & : \text{otherwise,} \end{cases}$$

and

$$\bar{d}_{p,q}(B) = \begin{cases} \bar{d}_{p,q}(A) - 1 & : i \leq p < s \text{ and } j \leq q < t, \\ \bar{d}_{p,q}(A) & : \text{otherwise,} \end{cases}$$

which proves that $B < A$. \square

Let $A \in \Lambda_1$ and $i \in \{1, \dots, n\}$ with $\text{ro}(A)_{i+1} > 0$. Using the fundamental multiplication rules 4.1.2 and Lemma 4.1.9,

$$E_i e_A = \sum_{s=1}^m q^{\sum_{t>j_s} a_{i,t}} [[a_{i,j_s} + 1]] e_{A + \mathcal{E}_{i,j_s} - \mathcal{E}_{i+1,j_s}}$$

where $j_1, \dots, j_m \in \mathbb{Z}$ with $j_1 < j_2 < \dots < j_m$ and

$$\{j_1, \dots, j_m\} = \{j \in \mathbb{Z} : a_{i+1,j} > 0\}.$$

The basis elements appearing in the above expression are totally ordered, with

$$A + \mathcal{E}_{i,j_s} - \mathcal{E}_{i+1,j_s} < A + \mathcal{E}_{i,j_{s+1}} - \mathcal{E}_{i+1,j_{s+1}}$$

for $s = 1, \dots, m-1$. Thus the term with $s = m$ is the maximum.

The partial order on Λ_1 induces a partial order on the set of G -orbits in $\mathcal{F} \times \mathcal{F}$, such that $\mathcal{O}_A \leq \mathcal{O}_B$ if and only if $A \leq B$. The following is a restatement of Lemma 3.2.12 and gives some geometric significance to the hook order on Λ_1 .

Lemma 4.1.10. *Let $A \in \Lambda_1$ and $(L, L') \in \mathcal{O}_A$. Then*

$$\dim \left(\frac{L_i}{L_i \cap L'_j} \right) = d_{i,j}(A)$$

and

$$\dim \left(\frac{L'_j}{L_i \cap L'_j} \right) = \bar{d}_{i,j}(A),$$

for each $i, j \in \mathbb{Z}$.

4.1.5 Shifting

In this subsection it is shown that the operations on Λ_1 given by shifting up by one row or to the right by one column may be described by the action, on the left or right respectively, of an invertible element R of $\hat{S}_q(n, r)$.

For each $A \in \Lambda_1$ and $m \in \mathbb{Z}$, the row shift of A by m is the element $[m]A$ of Λ_1 given by

$$([m]A)_{i,j} = a_{i+m,j},$$

for each $i, j \in \mathbb{Z}$.

The column shift of A by m is the element $A[m]$ given by

$$(A[m])_{i,j} = a_{i,j+m},$$

for each $i, j \in \mathbb{Z}$.

For $\lambda \in \Lambda_0$ and $m \in \mathbb{Z}$, the translation of λ by m is the element $\lambda[m]$ of Λ_0 given by

$$(\lambda[m])_i = \lambda_{i+m},$$

for each $i \in \mathbb{Z}$, where the indices of λ are taken modulo n .

Example 4.1.11. Let $\lambda = (2, 1, 3)$. Then $\lambda[1] = (1, 3, 2)$, $\lambda[2] = (3, 2, 1)$ and $\lambda[3] = \lambda$.

For each $\lambda \in \Lambda_0$, define

$$\begin{aligned} R_\lambda &= e_{[1]D_\lambda} \\ &= e_{\lambda_1 \mathcal{E}_{0,1} + \dots + \lambda_n \mathcal{E}_{n-1,n}} \end{aligned}$$

and let

$$R = \sum_{\lambda \in \Lambda_0} R_\lambda.$$

Recall that

$$\mathcal{O}_{D_\lambda} = \{(L, L) : L \in \mathcal{F}_\lambda\},$$

so

$$\mathcal{O}_{[m]D_\lambda} = \{([m]L, L) : L \in \mathcal{F}_\lambda\}$$

and

$$\mathcal{O}_{D_\lambda[m]} = \{(L, [m]L) : L \in \mathcal{F}_\lambda\}.$$

This leads to a simple rule for multiplication by R in terms of these shifts on matrices.

Lemma 4.1.12. *If $A \in \Lambda_1$ then*

$$Re_A = e_{[1]A}$$

and

$$e_AR = e_{A[-1]}.$$

Proof. Let $\mu = \text{ro}(A)$ and $\lambda = \text{co}(A)[-1]$. Observe that

$$\{(L, L', L'') \in \mathcal{F}^3 : (L, L') \in \mathcal{O}_{[1]D_\mu}, (L', L'') \in \mathcal{O}_A\} = \{(L'[1], L', L'') : (L', L'') \in \mathcal{O}_A\},$$

and the image under the projection onto the first and last components is

$$\{(L'[1], L'') : (L', L'') \in \mathcal{O}_A\} = \mathcal{O}_{[1]A}.$$

The coefficient of $\mathcal{O}_{[1]A}$ in the product Re_A is 1 since, for any $(N, N'') \in \mathcal{O}_{[1]A}$,

$$\{N' \in \mathcal{F} : (N, N') \in \mathcal{O}_{[1]D_\mu}, (N', N'') \in \mathcal{O}_A\} = \{N[-1]\},$$

so it follows that $Re_A = e_{[1]A}$.

To compute the product e_AR , consider

$$\{(L, L', L'') \in \mathcal{F}^3 : (L, L') \in \mathcal{O}_A, (L', L'') \in \mathcal{O}_{[1]D_\lambda}\} = \{(L, L', L'[-1]) : (L, L') \in \mathcal{O}_A\}.$$

The image under the projection onto the first and last components is

$$\{(L, L'[-1]) : (L, L') \in \mathcal{O}_A\} = \mathcal{O}_{A[-1]}$$

and, for any $(N, N'') \in \mathcal{O}_{A[-1]}$,

$$\{N' \in \mathcal{F} : (N, N') \in \mathcal{O}_A, (N', N'') \in \mathcal{O}_{[1]D_\lambda}\} = \{N''[1]\}.$$

Therefore $e_AR = e_{A[-1]}$. □

Lemma 4.1.13. *The element R is invertible and*

$$RS(R) = S(R)R = 1.$$

In particular,

$$R^{-1} = \sum_{\lambda \in \Lambda_0} e_{[-1]D_\lambda}.$$

Proof. Recall that

$$S(R) = \sum_{\lambda \in \Lambda_0} e_{[-1]D_\lambda}.$$

Then it follows from Lemma 4.1.12 that

$$RS(R) = \sum_{\lambda \in \Lambda_0} e_{[1][-1]D_\lambda} = 1$$

and

$$\begin{aligned} S(R)R &= \sum_{\lambda \in \Lambda_0} e_{[-1]D_\lambda[-1]} \\ &= \sum_{\lambda \in \Lambda_0} e_{D(\lambda[-1])} \\ &= 1. \end{aligned}$$

□

As a visual cue, acting on a basis element e_A on the left by R corresponds to moving the matrix A up by one row, while acting on the right by R corresponds to moving the matrix to the right by one column. Then conjugating by R corresponds to the composition of a shift to the left by one and a shift up by one, which is a shift by one along the diagonal, so conjugating by R^n leaves e_A invariant. Thus conjugation by R gives a $\mathbb{Z}[q]$ -algebra automorphism of $\hat{S}_q(n, r)$ which has order n .

Multiplication on the left by $S(R)$ sends e_A to $e_{[-1]A}$, while multiplication on the right by $S(R)$ sends e_A to $e_{A[1]}$.

Lemma 4.1.14. *For each $\lambda \in \Lambda_0$,*

$$R1_\lambda S(R) = 1_{[1]\lambda}$$

and, for each $i \in \{1, \dots, n\}$,

$$RE_i S(R) = E_{i-1}$$

and

$$RF_i S(R) = F_{i-1}.$$

Proof. It follows from Lemma 4.1.12 and Lemma 4.1.13 that

$$Re_A S(R) = e_{[1]A[1]},$$

for each $A \in \Lambda_1$. In particular,

$$R1_\lambda S(R) = 1_{\lambda[1]}$$

for each $\lambda \in \Lambda_0$,

$$RE_{i,\lambda} S(R) = E_{i-1,\lambda[1]}$$

for each $(\lambda, i) \in \Lambda_0 \times \mathbb{Z}$ with $\lambda_{i+1} > 0$, and

$$RF_{i,\lambda} S(R) = F_{i-1,\lambda[1]}$$

for each $(\lambda, i) \in \Lambda_0 \times \mathbb{Z}$ with $\lambda_i > 0$.

It now follows that

$$RE_i S(R) = E_{i-1}$$

and

$$RF_i S(R) = F_{i-1}$$

as claimed. □

[THIS IS DEFINITELY TRUE – PROVE IT.]

Although I can't be sure, I suspect that conjugation by R gives a realisation of the Auslander-Reiten translation on the nilpotent representations of a cyclic quiver determined by the upper triangular matrices in Λ_1 . This is at least plausible since the A.R translation τ sends the simple representation at vertex i to the simple representation at vertex $i - 1$, which is consistent with the conjugation by R , which sends E_i to E_{i-1} .

4.2 Quivers and relations

Assume n and r are integers with $n \geq 3$ and $r \geq 1$.

4.2.1 Relations in affine q -Schur algebras

Lemma 4.2.1. *If $i, j \in \{1, \dots, n\}$ and $i \neq j$, then*

$$E_i F_j - F_j E_i = 0.$$

For each $i \in \{1, \dots, n\}$,

$$E_i F_i - F_i E_i = \sum_{\lambda \in \Lambda_0} ([[\lambda_i]] - [[\lambda_{i+1}]]) 1_\lambda.$$

Proof. Denote e_A by $[A]$. Fix $i, j \in \{1, \dots, n\}$ with $i \neq j$. Then

$$\begin{aligned} E_i F_j &= \sum_{\lambda \in \Lambda_0} E_i [D_\lambda + \mathcal{E}_{j+1,j} - \mathcal{E}_{j,j}] \\ &= \sum_{\lambda \in \Lambda_0} [D_\lambda + \mathcal{E}_{j+1,j} - \mathcal{E}_{j,j} + \mathcal{E}_{i,i+1} - \mathcal{E}_{i+1,i+1}]. \end{aligned}$$

Observe that the nonzero terms in the above sum are those for which $\lambda_j > 0$ and $\lambda_{i+1} > 0$. Similarly,

$$\begin{aligned} F_j E_i &= \sum_{\lambda \in \Lambda_0} F_j [D_\lambda + \mathcal{E}_{i,i+1} - \mathcal{E}_{i+1,i+1}] \\ &= \sum_{\lambda \in \Lambda_0} [D_\lambda + \mathcal{E}_{i,i+1} - \mathcal{E}_{i+1,i+1} + \mathcal{E}_{j+1,j} - \mathcal{E}_{j,j}], \end{aligned}$$

where the sum is taken over those λ such that $\lambda_{i+1} > 0$ and $\lambda_j > 0$. Therefore

$$E_i F_j - F_j E_i = 0.$$

Again using Lemma 4.1.2,

$$\begin{aligned} E_i F_i &= \sum_{\lambda \in \Lambda_0} E_i [D_\lambda + \mathcal{E}_{i+1,i} - \mathcal{E}_{i,i}] \\ &= \sum_{\lambda \in \Lambda_0} [D_\lambda + \mathcal{E}_{i+1,i} - \mathcal{E}_{i,i} + \mathcal{E}_{i,i+1} - \mathcal{E}_{i+1,i+1}] + [[\lambda_i]] [D_\lambda] \end{aligned}$$

and

$$\begin{aligned} F_i E_i &= \sum_{\lambda \in \Lambda_0} F_i [D_\lambda + \mathcal{E}_{i,i+1} - \mathcal{E}_{i+1,i+1}] \\ &= \sum_{\lambda \in \Lambda_0} [D_\lambda + \mathcal{E}_{i,i+1} - \mathcal{E}_{i+1,i+1} + \mathcal{E}_{i+1,i} - \mathcal{E}_{i,i}] + [[\lambda_{i+1}]] [D_\lambda]. \end{aligned}$$

Therefore

$$E_i F_i - F_i E_i = \sum_{\lambda \in \Lambda_0} ([[\lambda_i]] - [[\lambda_{i+1}]]) 1_\lambda,$$

as required. \square

An explicit version of these relations will be given after defining some terminology. Given $\lambda \in \Lambda_0$ and $i \in \{1, \dots, n\}$, say that λ is internal with respect to i if $\lambda - \alpha_i, \lambda + \alpha_i \in \Lambda_0$. Say that λ is initial with respect to i if $\lambda - \alpha_i \notin \Lambda_0$ and that λ is final with respect to i if $\lambda + \alpha_i \notin \Lambda_0$.

Then the expression for the commutator $[E_i, F_i]$ in Lemma 4.2.1 gives the following relations in $\hat{S}_q(n, r)$:

- If λ is internal with respect to i then

$$E_{i,\lambda-\alpha_i}F_{i,\lambda} - F_{i,\lambda+\alpha_i}E_{i,\lambda} = 0.$$

- If λ is initial with respect to i then

$$F_{i,\lambda+\alpha_i}E_{i,\lambda} - 1_\lambda = 0.$$

- If λ is final with respect to i then

$$E_{i,\lambda-\alpha_i}F_{i,\lambda} - 1_\lambda = 0.$$

Lemma 4.2.2. *The following relations hold in $\hat{S}_q(n, r)$, when $n \geq 3$:*

$$E_i E_j - E_j E_i = 0$$

and

$$F_i F_j - F_j F_i = 0$$

for $i, j \in \{1, \dots, n\}$ such that $j \geq i + 2$,

$$\begin{aligned} E_i E_{i+1}^2 - (1+q)E_{i+1}E_i E_{i+1} + qE_{i+1}^2 E_i &= 0 \\ E_i^2 E_{i+1} - (1+q)E_i E_{i+1} E_i + qE_{i+1} E_i^2 &= 0 \end{aligned}$$

and

$$\begin{aligned} F_{i+1} F_i^2 - (1+q)F_i F_{i+1} F_i + qF_i^2 F_{i+1} &= 0 \\ F_{i+1}^2 F_i - (1+q)F_{i+1} F_i F_{i+1} + qF_i F_{i+1}^2 &= 0. \end{aligned}$$

for $i \in \{1, \dots, n\}$.

Proof. Denote e_A by $[A]$.

$$\begin{aligned} E_i E_{i+1}^2 &= \sum_{\lambda \in \Lambda_0} [[2]] [D_\lambda + \mathcal{E}_{i,i+2} + \mathcal{E}_{i+1,i+2} - 2\mathcal{E}_{i+2,i+2}] \\ &\quad + [[2]] [D_\lambda + \mathcal{E}_{i,i+1} - \mathcal{E}_{i+1,i+1} + 2\mathcal{E}_{i+1,i+2} - 2\mathcal{E}_{i+2,i+2}] \end{aligned}$$

$$\begin{aligned} E_{i+1} E_i E_{i+1} &= \sum_{\lambda \in \Lambda_0} [D_\lambda + \mathcal{E}_{i,i+2} + \mathcal{E}_{i+1,i+2} - 2\mathcal{E}_{i+2,i+2}] \\ &\quad + [[2]] [D_\lambda + \mathcal{E}_{i,i+1} - \mathcal{E}_{i+1,i+1} + 2\mathcal{E}_{i+1,i+2} - 2\mathcal{E}_{i+2,i+2}] \end{aligned}$$

$$E_{i+1}^2 E_i = \sum_{\lambda \in \Lambda_0} [[2]] [D_\lambda + \mathcal{E}_{i,i+1} - \mathcal{E}_{i+1,i+1} + 2\mathcal{E}_{i+1,i+2} - 2\mathcal{E}_{i+2,i+2}]$$

$$E_i E_{i+1}^2 - (1+q)E_{i+1}E_i E_{i+1} + qE_{i+1}^2 E_i = \sum_{\lambda \in \Lambda_0} ([2] - (1+q)) [X_\lambda] + ([2] - (1+q)[2] + q[[2]]) [Y_\lambda]$$

where

$$X_\lambda = D_\lambda + \mathcal{E}_{i,i+2} + \mathcal{E}_{i+1,i+2} - 2\mathcal{E}_{i+2,i+2}$$

and

$$Y_\lambda = D_\lambda + \mathcal{E}_{i,i+1} - \mathcal{E}_{i+1,i+1} + 2\mathcal{E}_{i+1,i+2} - 2\mathcal{E}_{i+2,i+2}.$$

It follows

$$E_i E_{i+1}^2 - (1+q)E_{i+1}E_iE_{i+1} + qE_{i+1}^2E_i = 0$$

and so

$$F_{i+1}^2F_i - (1+q)F_{i+1}F_iF_{i+1} + qF_iF_{i+1}^2 = 0,$$

by applying the transpose involution to the first relation.

$$\begin{aligned} E_i^2 E_{i+1} &= \sum_{\lambda \in \Lambda_0} [[2]] [D_\lambda + \mathcal{E}_{i,i+1} - \mathcal{E}_{i+1,i+1} + \mathcal{E}_{i,i+2} - \mathcal{E}_{i+2,i+2}] \\ &\quad + [[2]] [D_\lambda + 2\mathcal{E}_{i,i+1} - 2\mathcal{E}_{i+1,i+1} + \mathcal{E}_{i+1,i+2} - \mathcal{E}_{i+2,i+2}] \end{aligned}$$

$$\begin{aligned} E_i E_{i+1} E_i &= \sum_{\lambda \in \Lambda_0} [D_\lambda + \mathcal{E}_{i,i+1} - \mathcal{E}_{i+1,i+1} + \mathcal{E}_{i,i+2} - \mathcal{E}_{i+2,i+2}] \\ &\quad + [[2]] [D_\lambda + 2\mathcal{E}_{i,i+1} - 2\mathcal{E}_{i+1,i+1} + \mathcal{E}_{i+1,i+2} - \mathcal{E}_{i+2,i+2}] \end{aligned}$$

$$E_{i+1} E_i^2 = \sum_{\lambda \in \Lambda_0} [[2]] [D_\lambda + 2\mathcal{E}_{i,i+1} - 2\mathcal{E}_{i+1,i+1} + \mathcal{E}_{i+1,i+2} - \mathcal{E}_{i+2,i+2}]$$

So

$$E_i^2 E_{i+1} - (1+q)E_i E_{i+1} E_i + qE_{i+1} E_i = \sum_{\lambda \in \Lambda_0} ([2] - (1+q))A_\lambda + ([2] - (1+q)[2] + q[[2]])B_\lambda,$$

where

$$A_\lambda = D_\lambda + \mathcal{E}_{i,i+1} - \mathcal{E}_{i+1,i+1} + \mathcal{E}_{i,i+2} - \mathcal{E}_{i+2,i+2}$$

and

$$B_\lambda = D_\lambda + 2\mathcal{E}_{i,i+1} - 2\mathcal{E}_{i+1,i+1} + \mathcal{E}_{i+1,i+2} - \mathcal{E}_{i+2,i+2}.$$

Therefore

$$E_i^2 E_{i+1} - (1+q)E_i E_{i+1} E_i + qE_{i+1} E_i = 0$$

and

$$F_{i+1} F_i^2 - (1+q)F_i F_{i+1} F_i + qF_i^2 F_{i+1} = 0,$$

where the second relation follows from the first by applying the transpose involution. \square

Recall the result of Lemma 4.1.14, which gives relations involving R :

$$\begin{aligned} R1_\lambda R^{-1} &= 1_{\lambda[1]} \\ RE_{i,\lambda} R^{-1} &= E_{i-1,\lambda[1]} \\ RF_{i,\lambda} R^{-1} &= F_{i-1,\lambda[1]}. \end{aligned}$$

4.2.2 A quiver algebra

Define a quiver $\Gamma = \Gamma(n, r)$ associated with the affine q -Schur algebra $\hat{S}_q(n, r)$ as follows:

- The set of vertices is $\Gamma_0 = \Lambda_0$.
- The set of edges is Γ_1 , consisting of edges

$$e_{i,\lambda}: \lambda \rightarrow \lambda + \alpha_i$$

for $i \in \{1, \dots, n\}$ and $\lambda \in \Lambda_0$ such that $\lambda_{i+1} > 0$ and

$$f_{i,\lambda}: \lambda \rightarrow \lambda - \alpha_i$$

for $i \in \{1, \dots, n\}$ and $\lambda \in \Lambda_0$ such that $\lambda_i > 0$.

The path $\mathbb{Z}[q]$ -algebra of Γ is an associative $\mathbb{Z}[q]$ -algebra with a unit, which has a $\mathbb{Z}[q]$ -basis consisting of the paths in Γ , where the multiplication is defined by concatenation of paths. That is, if p and q are paths in Γ , then the product pq is the path ‘ q followed by p ’ if the target of q equals the source of p , or equals zero otherwise.

For each $\lambda \in \Lambda_0$, denote the constant path at λ by k_λ . These elements form a set of pairwise orthogonal idempotents and the multiplicative identity in $\mathbb{Z}[q]\Gamma$ is

$$\sum_{\lambda \in \Lambda_0} k_\lambda.$$

For each $i \in \{1, \dots, n\}$, define

$$e_i = \sum_{\lambda \in \Lambda_0: \lambda_{i+1} > 0} e_{i,\lambda}$$

and

$$f_i = \sum_{\lambda \in \Lambda_0: \lambda_i > 0} f_{i,\lambda}.$$

To avoid long subscripts, define $e_{i,\lambda} = 0$ if $\lambda_{i+1} = 0$ and define $f_{i,\lambda} = 0$ if $\lambda_i = 0$.

Let $I = I(n, r)$ be the ideal in $\mathbb{Z}[q]\Gamma$ generated by the following expressions:

$$\begin{aligned} e_i e_j - e_j e_i \\ f_i f_j - f_j f_i \end{aligned}$$

for $i, j \in \{1, \dots, n\}$ such that $j \neq i \pm 1$,

$$\begin{aligned} e_i e_{i+1}^2 - [[2]] e_{i+1} e_i e_{i+1} + q e_{i+1}^2 e_i \\ e_i^2 e_{i+1} - [[2]] e_i e_{i+1} e_i + q e_{i+1} e_i^2 \\ f_{i+1}^2 f_i - [[2]] f_{i+1} f_i f_{i+1} + q f_i f_{i+1}^2 \\ f_{i+1} f_i^2 - [[2]] f_i f_{i+1} f_i + q f_i^2 f_{i+1} \end{aligned}$$

for $i \in \{1, \dots, n\}$,

$$e_i f_j - f_j e_i$$

for $i, j \in \{1, \dots, n\}$ such that $i \neq j$,

$$e_i f_i - f_i e_i - \sum_{\lambda \in \Lambda_0} ([[\lambda_i]] - [[\lambda_{i+1}]]) k_\lambda$$

for $i \in \{1, \dots, n\}$.

4.2.3 Mapping to the q-Schur algebra

Lemma 4.2.3. *There is a $\mathbb{Z}[q]$ -algebra homomorphism*

$$\phi: \mathbb{Z}[q]\Gamma/I \rightarrow \hat{S}_q(n, r)$$

defined by

$$\begin{aligned}\phi(e_{i,\lambda} + I) &= E_{i,\lambda}, \\ \phi(f_{i,\lambda} + I) &= F_{i,\lambda}, \\ \phi(k_\lambda + I) &= 1_\lambda,\end{aligned}$$

for $i \in \{1, \dots, n\}$ and $\lambda \in \Lambda_0$.

Proof. Lemma 4.2.1 and Lemma 4.2.2 shows that each equation defining the ideal I corresponds to a zero relation in $\hat{S}_q(n, r)$, so there is a unique homomorphism of $\mathbb{Z}[q]$ -algebras given by

$$\begin{aligned}\phi(e_{i,\lambda} + I) &= E_{i,\lambda}, \\ \phi(f_{i,\lambda} + I) &= F_{i,\lambda}, \\ \phi(k_\lambda + I) &= 1_\lambda,\end{aligned}$$

for each $i \in \{1, \dots, n\}$ and $\lambda \in \Lambda_0$. □

In fact, ϕ is determined by its values on e_1, \dots, e_n , f_1, \dots, f_n and k_λ for $\lambda \in \Lambda_0$. In order to describe the image of this map we introduce the notion of standard paths in Γ .

Definition 4.2.4. A standard path in Γ is a path of the following form:

$$p = k_\mu p_1 \cdots p_l \cdot p'_1 \cdots p'_l$$

for some $l \in \mathbb{N}$, where

$$p_s = e_{i_0+s-1}^{\alpha_{i_0,s}} e_{i_0+s-2}^{\alpha_{i_0-1,s}} \cdots e_{i_0+s-n}^{\alpha_{i_0-n+1,s}}$$

and

$$p'_s = f_{i_1-s+1}^{\beta_{i_1,s}} f_{i_1-s+2}^{\beta_{i_1+1,s}} \cdots f_{i_1-s+n}^{\beta_{i_1+n-1,s}}$$

for $s \in \{1, \dots, l\}$, determined by two collections of non-negative integers $(\alpha_{i,s})$ and $(\beta_{i,s})$, for $i \in \{1, \dots, n\}$ and $s \in \{1, \dots, l\}$, satisfying the following conditions:

$$\alpha_{i,1} + \beta_{i-1,1} \leq \mu_i,$$

$$\alpha_{i,1} \geq \alpha_{i,2} \geq \cdots \geq \alpha_{i,l} \geq 0$$

and

$$\beta_{i,1} \geq \beta_{i,2} \geq \cdots \geq \beta_{i,l} \geq 0,$$

for $i \in \{1, \dots, n\}$; where

$$i_0 = \max\{i : i \in \{1, \dots, n\} \text{ with } \alpha_{i+1,s} = 0 \text{ for } s \in \{1, \dots, l\}\}$$

and

$$i_1 = \min\{i : i \in \{1, \dots, n\} \text{ with } \beta_{i,s} = 0 \text{ for } s \in \{1, \dots, l\}\}.$$

Observe that the definition of standard paths includes the constant paths k_λ for $\lambda \in \Lambda_0$. Suppose p is a standard path and $A \in \Lambda_1$ with $\rho(p + \mathcal{J}) = e_A$, then there are the following linear relations between the exponents in the standard expression for p and the entries of A :

$$\begin{aligned}\alpha_{i,s} &= \sum_{t \geq i+s} a_{i,t} \\ \beta_{i,s} &= \sum_{t \leq i-s+1} a_{i+1,t},\end{aligned}$$

for $i \in \{1, \dots, n\}$ and $s \in \mathbb{N}$; and

$$\begin{aligned}a_{i,i+s} &= \alpha_{i,s} - \alpha_{i,s+1} \\ a_{i,i-s} &= \beta_{i-1,s} - \beta_{i-1,s+1} \\ a_{i,i} &= \mu_i - \alpha_{i,1} - \beta_{i-1,1}.\end{aligned}$$

Thus there is a bijection between the set of standard paths in Γ and the standard basis of the affine q -Schur algebra indexed by Λ_1 .

Lemma 4.2.5. *Let $A \in \Lambda_1$ and let p be the standard path associated to A . Then*

$$\phi(p + I) = \left(\prod_{i \in \{1, \dots, n\}, s \geq 1} [[\alpha_{i,s}]]! [[\beta_{i,s}]]! \right) e_A + \sum_{B \in \Lambda_1: B < A} g_B e_B$$

Proof. **THIS IS A REALLY FUNDAMENTAL LEMMA – PROVE IT!** □

4.2.4 Change of rings

The following is based on the change of rings for quiver presentations result, Lemma 5.2 in [25]. Let R and S be commutative rings and suppose $f: R \rightarrow S$ is a ring homomorphism with $f(1) = 1$. Let Σ be a quiver and let $I \subset R\Sigma$ be an ideal of relations in Σ . The homomorphism f defines an R -algebra structure on S with $r \cdot s = f(r)s$ for all $r \in R$ and $s \in S$. Let $\bar{f}: R\Sigma \rightarrow S\Sigma$ be the R -algebra homomorphism induced by f , which is given by

$$\bar{f}(rp) = f(r)p$$

for each path p in Σ and $r \in R$.

Applying the right exact functor $S \otimes_R -$ to the short exact sequence

$$0 \rightarrow I \xrightarrow{i} R\Sigma \rightarrow R\Sigma/I \rightarrow 0$$

of R -modules gives the exact sequence

$$S \otimes_R I \xrightarrow{1 \otimes i} S \otimes_R R\Sigma \rightarrow S \otimes_R R\Sigma/I \rightarrow 0$$

of S -modules.

Let $m: S \otimes_R R\Sigma \rightarrow S\Sigma$ be the S -algebra homomorphism given by

$$m(s \otimes (rp + I)) = sf(r)p + S\bar{f}(I),$$

for all $r \in R$, $s \in S$ and paths p in Γ . The S -algebra homomorphism $S\Sigma \rightarrow S \otimes_R R\Sigma$ given by sending sp to $s \otimes p$ is inverse to m , so m is an isomorphism of S -algebras. Observe that m is also

R -linear, so is an isomorphism of R -algebras. The image of $m \circ (1 \otimes i): S \otimes_R I \rightarrow S\Sigma$ is $S\bar{f}(I)$ since the image is spanned by elements of the form

$$\begin{aligned} m((1 \otimes i)(s \otimes x)) &= m(s \otimes x) \\ &= s\bar{f}(x), \end{aligned}$$

for $s \in S$ and $x \in I$.

Thus we have a commuting diagram of S -modules with exact rows and columns:

$$\begin{array}{ccccccc} & & 0 & & \ker \bar{m} & & \\ & & \downarrow & & \downarrow & & \\ S \otimes_R I & \xrightarrow{1 \otimes i} & S \otimes_R R\Sigma & \rightarrow & S \otimes_R R\Sigma/I & \rightarrow & 0 \\ \downarrow & & \downarrow m & & \downarrow \bar{m} & & \\ 0 \rightarrow & S\bar{f}(I) & \rightarrow & S\Sigma & \rightarrow & S\Sigma/S\bar{f}(I) & \rightarrow 0 \\ \downarrow & & \downarrow & & \downarrow & & \\ & 0 & & 0 & & \text{coker } \bar{m} & \end{array}$$

The morphism \bar{m} is given by the universal property of cokernels and can be computed explicitly using the commuting diagram, with

$$\begin{aligned} \bar{m}(s \otimes (rp + I)) &= m(s \otimes rp) + S\bar{f}(I) \\ &= s\bar{f}(r)p + S\bar{f}(I), \end{aligned}$$

for all $r \in R$, $s \in S$ and paths p in Σ .

Lemma 4.2.6. [25] *The morphism*

$$\bar{m}: S \otimes_R R\Sigma/I \rightarrow S\Sigma/S\bar{f}(I)$$

is both an isomorphism of R -algebras and an isomorphism of S -algebras.

Proof. Using the snake lemma on the above commuting diagram gives an exact sequence of S -modules

$$0 \rightarrow \ker \bar{m} \rightarrow 0 \rightarrow 0 \rightarrow \text{coker } \bar{m} \rightarrow 0,$$

so $\ker \bar{m}$ and $\text{coker } \bar{m}$ are both zero and therefore \bar{m} is an isomorphism of S -algebras. Moreover, \bar{m} is R -linear, so is also an isomorphism of R -algebras. \square

Recall that the q -integers are given by

$$[[0]] = 0$$

and

$$[[m]] = 1 + q + \cdots + q^{m-1} = 1 + q[[m-1]]$$

for $m \in \mathbb{Z}$ with $m \geq 1$. For $m \in \mathbb{N}$, define the q -factorial

$$[[m]]! = \prod_{a=1}^m [[a]].$$

Given integers a and b with $0 < a < b$,

$$[[b]] - [[a]] = q^a [[b-a]]$$

and the product $[[a]][[b]]$ can be computed recursively as follows:

$$\begin{aligned} [[a]][[b]] &= (1 + q[[a-1]])(1 + q[[b-1]]) \\ &= 1 + q([a-1] + [b-1] + q[a-1][b-1]). \end{aligned}$$

The set of q -integers is not multiplicatively closed since, for example $[[2]]^2 = 1 + 2q + q^2$, but the set $1 + q\mathbb{Z}[q]$ is multiplicatively closed and contains the q -integers. Let \mathcal{Q} be the localisation of $\mathbb{Z}[q]$ at the set of elements of the form $1 + qf$ for $f \in \mathbb{Z}[q]$, so \mathcal{Q} is the subring of $\mathbb{Q}(q)$ given by

$$\mathcal{Q} = \left\{ \frac{f}{1 + qg} : f, g \in \mathbb{Z}[q] \right\}.$$

Observe that $\mathbb{Z}[q]$ is a subring of \mathcal{Q} , so \mathcal{Q} is a $\mathbb{Z}[q]$ -algebra. The \mathcal{Q} -form of the affine q -Schur algebra $\hat{S}_q(n, r)$ is defined to be the \mathcal{Q} -algebra

$$\hat{S}_{\mathcal{Q}}(n, r) = \mathcal{Q} \otimes_{\mathbb{Z}[q]} \hat{S}_q(n, r).$$

Lemma 4.2.7. *The \mathcal{Q} -algebra homomorphism*

$$\bar{m}: \mathcal{Q} \otimes_{\mathbb{Z}[q]} \mathbb{Z}[q]\Gamma/I \rightarrow \mathcal{Q}\Gamma/\mathcal{Q}I$$

given by

$$\bar{m}(s \otimes (hp + I)) = shp + \mathcal{Q}I,$$

for all $s \in \mathcal{Q}$, $h \in \mathbb{Z}[q]$ and paths p in Γ , is an isomorphism.

Proof. Applying Lemma 4.2.6 for the inclusion $\mathbb{Z}[q] \hookrightarrow \mathcal{Q}$ proves that \bar{m} is an isomorphism of \mathcal{Q} -algebras and an isomorphism of $\mathbb{Z}[q]$ -algebras. \square

Let $\phi_{\mathcal{Q}}$ be the \mathcal{Q} -algebra homomorphism

$$\phi_{\mathcal{Q}} = (1 \otimes \phi) \circ \bar{m}^{-1}: \mathcal{Q}\Gamma/\mathcal{Q}I \rightarrow \hat{S}_{\mathcal{Q}}(n, r),$$

which is given by

$$\begin{aligned} \phi_{\mathcal{Q}}(e_{i,\lambda}) &= E_{i,\lambda} \\ \phi_{\mathcal{Q}}(f_{i,\lambda}) &= F_{i,\lambda} \\ \phi_{\mathcal{Q}}(k_{\lambda}) &= 1_{\lambda}, \end{aligned}$$

for $i \in \{1, \dots, n\}$ and $\lambda \in \Lambda_0$.

Proposition 4.2.8. *If $r < n$ then $\phi_{\mathcal{Q}}$ is surjective.*

Proof. Fix $A \in \Lambda_1$ and let p be the standard path in Γ corresponding to A . Then

$$\phi_{\mathcal{Q}}(p + \mathcal{Q}I) = \sum_{B: B \leq A} g_B e_B$$

for some $g_B \in \mathbb{Z}[q]$, where

$$g_A = \prod_{i \in \{1, \dots, n\}, s \geq 1} [[\alpha_{i,s}]]! [[\beta_{i,s}]]!,$$

by Lemma 4.2.5. The coefficient of the leading term g_A is a unit in \mathcal{Q} , so

$$e_A = \phi_{\mathcal{Q}}(g_A^{-1}p + \mathcal{Q}I) - \sum_{B: B < A} g_B g_A^{-1} e_B.$$

There are only finitely many $B \in \Lambda_1$ with $B < A$ and for each such B , e_B admits a similar expression, which shows that e_A can be expressed as the image of a \mathcal{Q} -linear combination of the standard paths corresponding to the matrices B with $B \leq A$ and therefore $\phi_{\mathcal{Q}}$ is surjective. \square

Conjecture 4.2.9. If $r < n$, the quiver with relations (Γ, I) gives a presentation of $\hat{S}_{\mathcal{Q}}(n, r)$ over \mathcal{Q} .

Ideas for proof. The only thing that remains to be shown is that the map from the quiver algebra is injective, since Proposition 4.2.8 shows that this map is surjective.

I hope to deduce this from the presentation of the affine generic algebra by tensoring the surjective map between \mathcal{Q} -forms of the path algebra and q -Schur algebra with the \mathcal{Q} -algebra $\mathcal{Q}/(q)$ and observing this map is an isomorphism of \mathbb{Z} -algebras. \square

4.3 Relations for the $n=2$ case

In this section we give relations in $\hat{S}_q(2, r)$. Compare with the relations given in [\[REFERENCE - a double hall algebra approach to affine quantum schur weyl theory p.13\]](#) in the presentation of quantum affine \mathfrak{sl}_n .

Lemma 4.3.1. *The following equations hold in $\hat{S}_q(2, r)$:*

$$\begin{aligned} qE_1E_2^3 - [[3]]E_2E_1E_2^2 + [[3]]E_2^2E_1E_2 - qE_2^3E_1 &= 0 \\ qE_1^3E_2 - [[3]]E_1^2E_2E_1 + [[3]]E_1E_2E_1^2 - qE_2E_1^3 &= 0 \\ qF_2F_1^3 - [[3]]F_1F_2F_1^2 + [[3]]F_1^2F_2F_1 - qF_1^3F_2 &= 0 \\ qF_2^3F_1 - [[3]]F_2^2F_1F_2 + [[3]]F_2F_1F_2^2 - qF_1F_2^3 &= 0. \end{aligned}$$

Proof. It suffices to prove the first of these relations holds, since the second relation is obtained by applying the shifting automorphism of $\hat{S}_q(n, r)$ given by conjugation by R , which sends E_1 to E_2 and E_2 to E_1 , and then the last two relations are obtained by applying the transpose operator S on $\hat{S}_q(n, r)$, which sends E_i to F_i (for $i = 1, 2$) and reverses the order of multiplication.

Next, the first relation will be established by an explicit computation using the fundamental multiplication rules 4.1.3.

Write

$$\begin{aligned}
W &= D_\lambda + \mathcal{E}_{1,2} - \mathcal{E}_{1,1} + 3\mathcal{E}_{2,3} - 3\mathcal{E}_{2,2} \\
X &= D_\lambda + 2\mathcal{E}_{2,3} + \mathcal{E}_{2,4} - 3\mathcal{E}_{2,2} \\
Y &= D_\lambda + \mathcal{E}_{1,3} - \mathcal{E}_{1,1} + 2\mathcal{E}_{2,3} - 2\mathcal{E}_{2,2} \\
Z &= D_\lambda + \mathcal{E}_{2,3} + \mathcal{E}_{2,5} - 2\mathcal{E}_{2,2}.
\end{aligned}$$

$$E_1 E_2^3 = \sum_{\lambda \in \Lambda_0} [[2]][[3]]e_W + [[2]][[3]]e_Y$$

$$E_2 E_1 E_2^2 = \sum_{\lambda \in \Lambda_0} [[2]][[3]]e_W + [[2]]e_X + [[2]]^2 e_Y + [[2]]e_Z$$

$$E_2^2 E_1 E_2 = \sum_{\lambda \in \Lambda_0} [[2]][[3]]e_W + [[2]]^2 e_X + [[2]]e_Y + [[2]]e_Z$$

$$E_2^3 E_1 = \sum_{\lambda \in \Lambda_0} [[2]]3e_W + [[2]][[3]]e_X$$

Thus

$$\begin{aligned}
qE_1 E_2^3 - [[3]]E_2 E_1 E_2^2 + [[3]]E_2^2 E_1 E_2 - qE_2^3 E_1 &= [[2]][[3]](q - [[3]] + [[3]] - q)e_W \\
&\quad + [[2]][[3]](-1 + [[2]] - q)e_X \\
&\quad + [[2]][[3]](q - [[2]] + 1)e_Y \\
&\quad + ([[2]][[3]] - [[2]][[3]])e_Z,
\end{aligned}$$

which proves that the first relation holds and hence all the relations hold. \square

Chapter 5

A generic affine algebra

5.1 Introduction

Assume $\mathbf{k} = \mathbb{C}$ and fix $n, r \geq 1$. Let \mathcal{S} be the \mathbf{k} -algebra $\mathbf{k}[\varepsilon, \varepsilon^{-1}]$ and let \mathcal{R} be the subalgebra generated by ε , namely $\mathcal{R} = \mathbf{k}[\varepsilon]$. Let V be a free \mathcal{S} -module of rank r and let $\mathcal{F} = \mathcal{F}_{\mathbf{k}}(n, r)$ be the set of n -periodic cyclic flags in V ; so \mathcal{F} consists of collections $L = (L_i)_{i \in \mathbb{Z}}$ of \mathcal{R} -lattices in V with $L_i \subset L_{i+1}$ for $i \in \mathbb{Z}$ and $\varepsilon L_i = L_{i-n}$ for $i \in \mathbb{Z}$.

Let G be the group of \mathcal{S} -module automorphisms of V . Thus G is isomorphic to $\mathrm{GL}_r(\mathcal{S})$. G acts on \mathcal{F} with orbits $\{\mathcal{F}_\lambda : \lambda \in \Lambda_0\}$, where Λ_0 is the set of compositions of r into n parts, as in Definition 3.2.1.

The diagonal action of G on $\mathcal{F} \times \mathcal{F}$ has orbits $\{\mathcal{O}_A : A \in \Lambda_1\}$, where \mathcal{O}_A consists of those pairs of flags with periodic characteristic matrix equal to A . Definitions of the periodic characteristic matrix and the set Λ_1 are given in Definition 3.2.7 and Definition 3.2.2 respectively.

Recall that the periodic characteristic matrix of a pair $(L, L') \in \mathcal{F} \times \mathcal{F}$ is the $\mathbb{Z} \times \mathbb{Z}$ matrix $A = (a_{i,j})_{i,j \in \mathbb{Z}}$, with

$$a_{i,j} = \dim \left(\frac{L_i \cap L'_j}{L_{i-1} \cap L'_j + L_i \cap L'_{j-1}} \right)$$

for each $i, j \in \mathbb{Z}$.

Recall that ro and co are the maps $\Lambda_1 \rightarrow \Lambda_0$ given by

$$\mathrm{ro}(A) = \left(\sum_{j \in \mathbb{Z}} a_{1,j}, \dots, \sum_{j \in \mathbb{Z}} a_{n,j} \right)$$

and

$$\mathrm{co}(A) = \left(\sum_{i \in \mathbb{Z}} a_{i,1}, \dots, \sum_{i \in \mathbb{Z}} a_{i,n} \right)$$

for each $A \in \Lambda_1$. Given $A \in \Lambda_1$, write $A: \mathrm{co}(A) \rightarrow \mathrm{ro}(A)$.

The purpose of this chapter is to define an associative \mathbb{Z} -algebra with a multiplicative basis by defining a modified form of the product in the affine q -Schur algebra. In particular, given $A, B \in \Lambda_1$, the orbit product

$$X_{A,B} = \{(L, L'') \in \mathcal{F} \times \mathcal{F} : \exists L' \in \mathcal{F} \text{ with } (L, L') \in \mathcal{O}_A, (L', L'') \in \mathcal{O}_B\}$$

consists of finitely many G -orbits and it will be shown that there is a unique ‘generic’ orbit in $X_{A,B}$, denoted \mathcal{O}_{A*B} , with the property that

$$\dim \left(\frac{L_i}{L_i \cap L_j''} \right) \leq \dim \left(\frac{N_i}{N_i \cap N_j''} \right)$$

and

$$\dim \left(\frac{L_j''}{L_i \cap L_j''} \right) \leq \dim \left(\frac{N_j''}{N_i \cap N_j''} \right)$$

for all $i, j \in \mathbb{Z}$, $(N, N'') \in \mathcal{O}_{A*B}$ and $(L, L'') \in X_{A,B}$. It will be shown that the above ‘generic product’ of orbits is associative, so the free \mathbb{Z} -module on the set of G -orbits in $\mathcal{F} \times \mathcal{F}$ with \mathbb{Z} -bilinear multiplication given by

$$\mathcal{O}_A * \mathcal{O}_B = \mathcal{O}_{A*B},$$

for each $A, B \in \Lambda_1$ with $\text{co}(A) = \text{ro}(B)$, and

$$\mathcal{O}_A * \mathcal{O}_B = 0$$

for $A, B \in \Lambda_1$ with $\text{co}(A) \neq \text{ro}(B)$, is an associative \mathbb{Z} -algebra with multiplicative identity given by

$$\sum_{\lambda \in \Lambda_0} \mathcal{O}_{D_\lambda},$$

where D_λ is the diagonal matrix with $\text{co}(D_\lambda) = \lambda$. The resulting \mathbb{Z} -algebra is called the *generic affine algebra* (of rank r and period n), denoted $\hat{G}(n, r)$.

5.2 Grassmannians and related varieties

Here we collect a few elementary results on Grassmannians and some related varieties. In this section, let V be an n -dimensional \mathbf{k} -vector space and let $0 \leq d \leq n$ be an integer. There is a linear map

$$\phi^{(d)} : \Lambda^d(V) \rightarrow \text{Hom}(V, \Lambda^{d+1}(V))$$

given by

$$\phi^{(d)}(\alpha)(v) = \alpha \wedge v$$

for $\alpha \in \Lambda^d(V)$ and $v \in V$. The kernel of $\phi^{(d)}(\alpha)$ is the space of divisors of α ,

$$D_\alpha = \{v \in V : \alpha \wedge v = 0\}.$$

An element $\alpha \in \Lambda^d(V)$ is said to be totally decomposable if $\alpha = \alpha_1 \wedge \cdots \wedge \alpha_d$, where $\alpha_1, \dots, \alpha_d \in V$ are linearly independent. The dimension of D_α is at most d and $\dim(D_\alpha) = d$ precisely when α is totally decomposable. Consequently, the rank of $\phi^{(d)}(\alpha)$ is at least $n - d$ and α is totally decomposable if and only if $\text{rank } \phi^{(d)}(\alpha) \leq n - d$, which holds if and only if the $(n - d + 1) \times (n - d + 1)$ -minors of a matrix of $\phi^{(d)}(\alpha)$ are all zero.

Lemma 5.2.1. $\{(U_1, U_2) \in \text{Gr}_{d_1}(V) \times \text{Gr}_{d_2}(V) : \dim(U_1 \cap U_2) \geq a\}$ is a projective variety, for each $d_1, d_2, a \in \mathbb{N}$ with $d_1, d_2, a \leq n$.

Proof. As above, there is a linear map $\Psi: \Lambda^{d_1}V \oplus \Lambda^{d_2}V \rightarrow \text{Hom}(V, \Lambda^{d_1+1}(V) \oplus \Lambda^{d_2+1}(V))$ given by $\Psi(\alpha, \beta)(v) = (\alpha \wedge v, \beta \wedge v)$. Given $\alpha \in \Lambda^{d_1}(V)$ and $\beta \in \Lambda^{d_2}(V)$, the kernel of $\Psi(\alpha, \beta)$ is $D_\alpha \cap D_\beta$ and so the rank of $\Psi(\alpha, \beta)$ is $n - \dim(D_\alpha \cap D_\beta)$.

Let $U_i \in \text{Gr}_{d_i}(V)$ and suppose $p_i(U_i) = [\alpha_i]$, where p_i is the Plücker embedding of $\text{Gr}_{d_i}(V)$ in $\mathbb{P}(\Lambda^{d_i}(V))$, so $U_i = D_{\alpha_i} = \ker \phi^{(d_i)}(\alpha)$. Therefore the kernel of $\Psi(\alpha_1, \alpha_2)$ is $U_1 \cap U_2$, so the condition that $\dim(U_1 \cap U_2) \geq a$ is equivalent to the condition that $\Psi(\alpha_1, \alpha_2)$ has rank at most $n - a$. After fixing a basis of V , this condition is given by the vanishing of the $(n - a + 1) \times (n - a + 1)$ minors of the matrix of $\Psi(\alpha_1, \alpha_2)$ with respect to this basis. Therefore $\{(U_1, U_2) \in \text{Gr}_{d_1}(V) \times \text{Gr}_{d_2}(V) : \dim(U_1 \cap U_2) \geq a\}$ is a closed subset of the product of Grassmannians $\text{Gr}_{d_1}(V) \times \text{Gr}_{d_2}(V)$, so is a projective variety.

More precisely, the entries of a matrix of $\Psi(\alpha_1, \alpha_2)$ are homogeneous polynomials of degree 1 in the Plücker coordinates on $\text{Gr}_{d_1}(V) \times \text{Gr}_{d_2}(V)$ since Ψ is linear and so the minors of $\Psi(\alpha_1, \alpha_2)$ are also homogeneous polynomials in the Plücker coordinates. \square

Lemma 5.2.2. *Let V be an n -dimensional vector space over \mathbf{k} and let $d_1, d_2, a \in \mathbb{N}$ with $d_1, d_2, a \leq n$. The following hold:*

1. $\{(U_1, U_2) \in \text{Gr}_{d_1}(V) \times \text{Gr}_{d_2}(V) : \dim(U_1 \cap U_2) = a\}$ is a quasiprojective variety;
2. $\{(U_1, U_2) \in \text{Gr}_{d_1}(V) \times \text{Gr}_{d_2}(V) : U_1 \subset U_2\}$ is a projective variety;
3. Given $U_2 \in \text{Gr}_{d_2}(V)$, $\{U_1 \in \text{Gr}_{d_1}(V) : \dim(U_1 \cap U_2) \geq a\}$ is a projective variety;
4. Given $U_2 \in \text{Gr}_{d_2}(V)$, $\{U_1 \in \text{Gr}_{d_1}(V) : \dim(U_1 \cap U_2) = a\}$ is a quasiprojective variety;
5. Given $U_2 \in \text{Gr}_{d_2}(V)$, $\{U_1 \in \text{Gr}_{d_1}(V) : U_1 \subset U_2\}$ is a projective variety;
6. Given $U_2 \in \text{Gr}_{d_2}(V)$, $\{U_1 \in \text{Gr}_{d_1}(V) : U_2 \subset U_1\}$ is a projective variety.

Proof. Let X_i denote the space in statement i of the lemma. To emphasise the dependence of X_i on a , write $X_{i,a}$.

X_1 is a quasiprojective variety since it is equal to the intersection of the projective variety $\{(U_1, U_2) \in \text{Gr}_{d_1}(V) \times \text{Gr}_{d_2}(V) : \dim(U_1 \cap U_2) \geq a\}$ with the open set $\{(U_1, U_2) \in \text{Gr}_{d_1}(V) \times \text{Gr}_{d_2}(V) : \dim(U_1 \cap U_2) \leq a\}$.

Given $(U_1, U_2) \in \text{Gr}_{d_1}(V) \times \text{Gr}_{d_2}(V)$, $U_1 \subset U_2$ if and only if $\dim(U_1 \cap U_2) \geq d_1$, so Lemma 5.2.1 shows X_2 is a projective variety.

Let $\pi_i: \text{Gr}_{d_1}(V) \times \text{Gr}_{d_2}(V) \rightarrow \text{Gr}_{d_i}(V)$ be the projection map onto the i -th factor, for $i = 1, 2$. The completeness property of projective varieties ensures that π_i is a closed morphism. Observe that

$$\begin{aligned} X_3 &= \{U_1 \in \text{Gr}_{d_1}(V) : \dim(U_1 \cap U_2) \geq a\} \\ &= \pi_1(\{(U_1, W) \in \text{Gr}_{d_1}(V) \times \text{Gr}_{d_2}(V) : \dim(U_1 \cap W) \geq a\} \cap \pi_2^{-1}\{U_2\}). \end{aligned}$$

The fibre of π_2 over U_2 is closed, so the intersection of the fibre with the variety from Lemma 5.2.1 is closed and then the image of this intersection under π_1 is closed. This shows X_3 is a projective variety.

X_4 is a quasiprojective variety since it is the complement of the subvariety $X_{3,a+1}$ in $X_{3,a}$. Finally, 5-6 follow as special cases of 3 since $X_5 = X_{3,d_1}$ and $X_6 = X_{3,d_2}$. \square

5.3 Geometry of affine flag varieties

Given $L \in \mathcal{F}$, $N, a \in \mathbb{N}$ and $\lambda \in \Lambda_0$ define

$$\Pi_{N,\lambda}(L) = \{L' \in \mathcal{F}_\lambda : \varepsilon^N L_0 \subset L'_0 \subset \varepsilon^{-N} L_0\}.$$

and

$$\Pi_{N,\lambda}^a(L) = \left\{ L' \in \mathcal{F}_\lambda : \varepsilon^N L_0 \subset L'_0 \subset \varepsilon^{-N} L_0, \dim \left(\frac{\varepsilon^{-N} L_0}{L'_0} \right) = a \right\}.$$

Lemma 5.3.1. *Given $L \in \mathcal{F}$, $N \in \mathbb{N}$ and $\lambda \in \Lambda_0$,*

$$\Pi_{N,\lambda}(L) = \bigcup_{a: 0 \leq a \leq 2Nr} \Pi_{N,\lambda}^a(L).$$

Proof. If $L' \in \Pi_{N,\lambda}(L)$ then $\varepsilon^N L_0 \subset L'_0 \subset \varepsilon^{-N} L_0$ and the $\mathbf{k}[\varepsilon]$ -module $\varepsilon^{-N} L_0 / L'_0$ is naturally isomorphic to $(\varepsilon^{-N} L_0 / \varepsilon^N L_0) / (L'_0 / \varepsilon^N L_0)$, so

$$\dim \left(\frac{\varepsilon^{-N} L_0}{L'_0} \right) \leq \dim \left(\frac{\varepsilon^{-N} L_0}{\varepsilon^N L_0} \right) = 2Nr.$$

□

Lemma 5.3.2. *Given $L \in \mathcal{F}$, $N \in \mathbb{N}$, $\lambda \in \Lambda_0$ and $a \in \mathbb{N}$ with $0 \leq a \leq 2Nr$, $\Pi_{N,\lambda}^a(L)$ is a projective algebraic variety.*

Proof. Let W be the $\mathbf{k}[\varepsilon]$ -module $\varepsilon^{-1-N} L_0 / \varepsilon^N L_0$, which has dimension $(2N+1)r$ over \mathbf{k} . Let $d_i = 2Nr - a + \lambda_1 + \dots + \lambda_i$ for each $i = 1, \dots, n$. The correspondence between submodules of $\varepsilon^{-1-N} L_0$ which contain $\varepsilon^N L_0$ and submodules of $\varepsilon^{-1-N} L_0 / \varepsilon^N L_0$ determines a map

$$\rho: \Pi_{N,\lambda}^a(L) \rightarrow \mathrm{Gr}_{d_1}(W) \times \dots \times \mathrm{Gr}_{d_n}(W),$$

with $\rho(L') = (L'_1 / \varepsilon^N L_0, \dots, L'_n / \varepsilon^N L_0)$.

Let \mathcal{X} be the space of $(U_1, \dots, U_n) \in \mathrm{Gr}_{d_1}(W) \times \dots \times \mathrm{Gr}_{d_n}(W)$ with $U_i \subset U_{i+1}$ for $i = 1, \dots, n-1$ and $\varepsilon U_n \subset U_1$. Lemma 5.2.2 shows that each of these conditions is closed, so \mathcal{X} is a closed subset of $\mathrm{Gr}_{d_1}(W) \times \dots \times \mathrm{Gr}_{d_n}(W)$, therefore \mathcal{X} is a projective algebraic variety.

The image of ρ is contained in \mathcal{X} since

$$\varepsilon L'_n / \varepsilon^N L_0 = L'_0 / \varepsilon^N L_0 \subset L'_1 / \varepsilon^N L_0 \subset \dots \subset L'_n / \varepsilon^N L_0.$$

Suppose $(U_1, \dots, U_n) \in \mathcal{X}$. Then U_i is a $\mathbf{k}[\varepsilon]$ -module, since $\varepsilon U_i \subset \varepsilon U_n \subset U_1 \subset U_i$, for each $i = 1, \dots, n$, so U_i lifts uniquely to a $\mathbf{k}[\varepsilon]$ -module L'_i with $\varepsilon^N L_0 \subset L'_i \subset \varepsilon^{-1-N} L_0$. Therefore L'_1, \dots, L'_n are $\mathbf{k}[\varepsilon]$ -lattices with $L_i \subset L_{i+1}$ for $i = 1, \dots, n-1$ and $\varepsilon L'_n \subset L'_1$, with

$$\dim(\varepsilon^{-1-N} L_0 / L'_n) = \dim(W / W_n) = (2N+1)r - d_n = a$$

and

$$\dim(L'_i / L'_{i-1}) = \dim(W_i / W_{i-1}) = d_i - d_{i-1} = \lambda_i,$$

for each $i = 2, \dots, n$. Therefore there is a unique $L' \in \Pi_{N,\lambda}^a(L)$ such that $\rho(L') = (W_1, \dots, W_n)$, where L' is given by $L'_{i+cn} = \varepsilon^{-c} L'_i$ for $i = 1, \dots, n$ and $c \in \mathbb{Z}$. It follows ρ is injective and $\mathrm{im} \rho = \mathcal{X}$, which is a projective variety, so $\Pi_{N,\lambda}^a(L)$ is a projective variety. □

Lemma 5.3.3. *Given $L \in \mathcal{F}$, $N \in \mathbb{N}$, $\lambda \in \Lambda_0$ and $a \in \mathbb{N}$ with $0 \leq a \leq 2Nr$, $\Pi_{N,\lambda}^a(L)$ is closed in $\Pi_{N+1,\lambda}^{a+r}(L)$.*

Proof. If $L' \in \Pi_{N,\lambda}^a(L)$, then $\varepsilon^{N+1}L_0 \subset \varepsilon^N L_0 \subset L'_0 \subset \varepsilon^{-N}L_0 \subset \varepsilon^{-(N+1)}L_0$ and

$$\dim \left(\frac{\varepsilon^{-(1+n)}L_0}{L'_0} \right) = \dim \left(\frac{L_0}{\varepsilon L_0} \right) + \dim \left(\frac{\varepsilon^{-N}L_0}{L'_0} \right) = r + a,$$

which shows that $\Pi_{N,\lambda}^a(L) \subset \Pi_{N+1,\lambda}^{a+r}(L)$. For $L' \in \Pi_{N+1,\lambda}^{a+r}(L)$, if additionally $\varepsilon^N L_0 \subset L'_0 \subset \varepsilon^{-N}L_0$, then

$$\dim \left(\frac{\varepsilon^{-(N+1)}L_0}{L'_0} \right) = r + \dim \left(\frac{\varepsilon^{-N}L_0}{L'_0} \right),$$

which shows $L' \in \Pi_{N,\lambda}^a(L)$. Therefore $\Pi_{N,\lambda}^a(L)$ is the subspace of $\Pi_{N+1,\lambda}^{a+r}(L)$ defined by the two closed conditions $\varepsilon^N L_0 \subset L'_0$ and $L'_0 \subset \varepsilon^{-N}L_0$, using Lemma 5.2.2. \square

Lemma 5.3.4. *Let $\lambda \in \Lambda_0$, $M, N \in \mathbb{N}$, $L, \tilde{L} \in \mathcal{F}$, $0 \leq a \leq 2Nr$, $0 \leq b \leq 2Mr$. $\Pi_{N,\lambda}^a(L) \cap \Pi_{M,\lambda}^b(\tilde{L})$ is a closed set in $\Pi_{N,\lambda}^a(L)$. In particular, if the intersection is nonempty it is a projective algebraic variety.*

Proof. Observe that $\Pi_{N,\lambda}^a(L) \cap \Pi_{M,\lambda}^b(\tilde{L})$ is the subset of $\Pi_{N,\lambda}^a(L)$ defined by the additional conditions that $\varepsilon^M \tilde{L}_0 \subset L'_0$ and $L'_0 \subset \varepsilon^{-M} \tilde{L}_0$, so is a closed subset of $\Pi_{N,\lambda}^a(L)$, using 5.2.2. \square

Lemma 5.3.5. *Suppose $L \in \mathcal{F}$, $N, a \in \mathbb{N}$ and $\lambda \in \Lambda_0$ with $a \leq 2Nr$. For each $g \in G$, the natural map (restriction of the action map) $\Pi_{N,\lambda}^a(L) \rightarrow \Pi_{N,\lambda}^a(gL)$ is an isomorphism of projective varieties.*

Proof. If $L' \in \Pi_{N,\lambda}^a(L)$, then $\varepsilon^N L_0 \subset L'_0 \subset \varepsilon^{-N}L_0$ and so $\varepsilon^N g(L_0) \subset g(L'_0) \subset \varepsilon^{-N}g(L_0)$, so $gL' \in \Pi_{N,\lambda}^a(gL)$. Thus g and g^{-1} induce mutually inverse morphisms of varieties $g: \Pi_{N,\lambda}^a(L) \rightarrow \Pi_{N,\lambda}^a(gL)$ and $g^{-1}: \Pi_{N,\lambda}^a(gL) \rightarrow \Pi_{N,\lambda}^a(L)$. \square

5.3.1 Action through an algebraic group

Let W be the $\mathbb{C}[\varepsilon]$ -module $\varepsilon^{-(1+N)}L_0/\varepsilon^N L_0$. ε^{2N+1} acts as zero on W and $\mathbb{C}[\varepsilon]/\langle \varepsilon^{2N+1} \rangle \otimes_{\mathbb{C}[\varepsilon]} W$ is a free $\mathbb{C}[\varepsilon]/\langle \varepsilon^{2N+1} \rangle$ -module of rank r . In particular, W is a complex vector space of dimension $(2N+1)r$.

Each element $g \in G_L$ determines an endomorphism \bar{g} of W , given by

$$\bar{g}(x + \varepsilon^N L_0) = g(x) + \varepsilon^N L_0,$$

for each $x \in \varepsilon^{-(1+N)}L_0$. Given $g, h \in G_L$, $\overline{gh} = \bar{g}\bar{h}$ and so \bar{g} is an automorphism of W with $\bar{g}^{-1} = \overline{g^{-1}}$. Therefore the map $\bar{\cdot}: G_L \mapsto \text{GL}(W)$ given by $g \mapsto \bar{g}$ is a group homomorphism with kernel

$$H_{N,L} := \{g \in G_L : \bar{g} = 1\},$$

which consists of those $g \in G_L$ such that

$$g(x) - x \in \varepsilon^N L_0$$

for each $x \in \varepsilon^{-(1+N)}L_0$. Thus $G_L/H_{N,L}$ may be identified with a subgroup of $\text{GL}(W)$.

Lemma 5.3.6. $G_L/H_{N,L}$ is a connected algebraic group.

Proof. As a result of the first isomorphism theorem, $G_L/H_{N,L}$ is isomorphic to the image of G_L in $\text{GL}(W)$, which will be described explicitly by equations in the coordinate functions on $\text{GL}(W)$, with respect to a fixed basis of W .

Let $\{\tilde{x}_1, \dots, \tilde{x}_r\}$ be a basis of L_n/L_0 over \mathbb{C} which is adapted to the flag

$$L_1/L_0 \subset \dots \subset L_{n-1}/L_0 \subset L_n/L_0,$$

so that

$$L_i/L_0 = \langle \tilde{x}_1, \dots, \tilde{x}_{\lambda_1 + \dots + \lambda_i} \rangle$$

for each $i \in \{1, \dots, n\}$. Fix $x_1, \dots, x_r \in L_n$ such that $\tilde{x}_i = x_i + L_0$ for each $i = 1, \dots, r$, then

$$L_i = L_0 + \langle x_1, \dots, x_{\lambda_1 + \dots + \lambda_i} \rangle$$

for $i = 1, \dots, r$.

Then W has a \mathbb{C} -basis $\{y_j : 1 \leq j \leq (2N+1)r\}$ given by

$$y_{i+cr} = \varepsilon^{-c+N} x_i$$

for each $i \in \{1, \dots, r\}$ and $c \in \{0, \dots, 2N\}$. Observe that $\varepsilon y_i = 0$ for $i \in \{1, \dots, r\}$ and $\varepsilon y_i = y_{i-r}$ for $r < i \leq (2N+1)r$.

The coordinate functions on $\text{GL}(W)$ with respect to this choice of basis are the maps

$$\gamma_{i,j} : \text{GL}(W) \rightarrow \mathbb{C}$$

for $i, j \in \mathbb{Z}$ with $1 \leq i, j \leq (2N+1)r$, given by

$$g(y_j) = \sum_i \gamma_{ij}(g) y_i,$$

for each $j = 1, \dots, (2N+1)r$.

The image of G_L in $\text{GL}(W)$ is the subgroup defined by the conditions

$$\gamma_{i,j} = \gamma_{i-r,j-r}$$

for each $i, j \in \{r+1, \dots, (2N+1)r\}$ and

$$\gamma_{i,j} = 0$$

for each $i, j \in \{1, \dots, (2N+1)r\}$ with $i > \lambda_1 + \dots + \lambda_s$ and $j \leq \lambda_1 + \dots + \lambda_s$ for some $s \in \{1, \dots, r\}$. This shows that the image of G_L in $\text{GL}(W)$ is a connected algebraic group and therefore $G_L/H_{N,L}$ is a connected algebraic group.

With respect to the basis $\{y_i : i \in \{1, \dots, (2N+1)r\}\}$, the image of G_L in $\text{GL}(W)$ consists of matrices of the form

$$\begin{pmatrix} A_0 & A_1 & A_2 & \cdots & A_{2N} \\ 0 & A_0 & A_1 & \cdots & A_{2N-1} \\ 0 & 0 & A_0 & \cdots & A_{2N-2} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & A_0 \end{pmatrix}$$

where $A_0 \in \mathcal{P}_\lambda$ and $A_1, \dots, A_{2N} \in M_r(\mathbb{C})$, where \mathcal{P}_λ is the parabolic subgroup of $\text{GL}_r(\mathbb{C})$ which is the stabiliser of the flag

$$L_1/L_0 \subset \dots \subset L_{n-1}/L_0 \subset L_n/L_0.$$

□

Given $g \in G$, the map $G_L \rightarrow G_{gL}$ sending h to ghg^{-1} is a group isomorphism which descends to an isomorphism of algebraic groups $G_L/H_{N,L} \rightarrow G_{gL}/H_{N,gL}$. Thus we have a commuting diagram of morphisms of varieties, where the vertical arrows are isomorphisms:

$$\begin{array}{ccc} G_L/H_{N,L} \times \Pi_{N,\lambda}^a(L) & \longrightarrow & \Pi_{N,\lambda}^a(L) \\ \downarrow & & \downarrow \\ G_{gL}/H_{N,gL} \times \Pi_{N,\lambda}^a(gL) & \longrightarrow & \Pi_{N,\lambda}^a(gL) \end{array}$$

5.3.2 Incidence in affine flag varieties

Lemma 5.3.7. *Given $N, a, b, c \in \mathbb{N}$, $\lambda, \mu \in \Lambda_0$, $L \in \mathcal{F}$ and $i, j \in \mathbb{Z}$,*

$$\left\{ (L', L'') \in \Pi_{N,\lambda}^a(L) \times \Pi_{N,\mu}^b(L) : \dim \left(\frac{L'_i}{L'_i \cap L''_j} \right) \leq c \right\}$$

is a closed set in the projective variety $\Pi_{N,\lambda}^a(L) \times \Pi_{N,\mu}^b(L)$.

Proof. There is $M \geq N$ so that $\varepsilon^M L_0 \subset L'_i \subset \varepsilon^{-M} L_0$ and $\varepsilon^M L_0 \subset L''_j \subset \varepsilon^{-M} L_0$. Let $a' = a + (M - N)r$ and $b' = b + (M - N)r$. Lemma 5.3.3 shows that $\Pi_{N,\lambda}^a(L)$ is a subvariety of $\Pi_{M,\lambda}^{a'}(L)$, so $\Pi_{N,\lambda}^a(L) \times \Pi_{N,\mu}^b(L)$ is a subvariety of $\Pi_{M,\lambda}^{a'}(L) \times \Pi_{M,\mu}^{b'}(L)$.

The fact that

$$\dim \left(\frac{L'_i}{L'_i \cap L''_j} \right) = \dim \left(\frac{L'_i/\varepsilon^M L_0}{L'_i/\varepsilon^M L_0 \cap L''_j/\varepsilon^M L_0} \right),$$

together with Lemma 5.3.2 and Lemma 5.2.1, shows that

$$\left\{ (L', L'') \in \Pi_{M,\lambda}^{a'}(L) \times \Pi_{M,\mu}^{b'}(L) : \dim \left(\frac{L'_i}{L'_i \cap L''_j} \right) \leq c \right\}$$

is closed, so the intersection with $\Pi_{N,\lambda}^a(L) \times \Pi_{N,\mu}^b(L)$ is closed. □

Lemma 5.3.8. *Given $N, a, c \in \mathbb{N}$, $\lambda \in \Lambda_0$, $L \in \mathcal{F}$ and $i, j \in \mathbb{Z}$,*

$$\left\{ L' \in \Pi_{N,\lambda}^a(L) : \dim \left(\frac{L_i}{L_i \cap L'_j} \right) \leq c \right\}$$

and

$$\left\{ L' \in \Pi_{N,\lambda}^a(L) : \dim \left(\frac{L'_j}{L_i \cap L'_j} \right) \leq c \right\}$$

are closed sets in $\Pi_{N,\lambda}^a(L)$.

Proof. This is a result of Lemma 5.2.2, since

$$\dim \left(\frac{L_i}{L_i \cap L'_j} \right) = \dim \left(\frac{L_i/\varepsilon^M L_0}{L_i/\varepsilon^M L_0 \cap L'_j/\varepsilon^M L_0} \right),$$

where $M \geq N$ is chosen so that $\varepsilon^M L_0 \subset L_i \subset \varepsilon^{-M} L_0$ and $\varepsilon^M L_0 \subset L'_j \subset \varepsilon^{-M} L_0$ for each $L' \in \Pi_{N,\lambda}^a(L)$. □

5.4 Geometry of orbits

Let $A \in \Lambda_1$ and $L \in \mathcal{F}_{\text{ro}(A)}$ and write $\lambda = \text{co}(A)$. Recall that

$$X_A^L = \{L' \in \mathcal{F}_\lambda : (L, L') \in \mathcal{O}_A\}.$$

Lemma 5.4.1. *There is $N \in \mathbb{N}$ such that $X_A^L \subset \Pi_{N,\lambda}^a(L)$, where $a = d_{nN,0}A$.*

Proof. There is $N \in \mathbb{N}$ so that $a_{i,j} = 0$ whenever $|j - i| > nN$. If $(L, L') \in \mathcal{O}_A$ then

$$\dim \left(\frac{L'_0}{L'_0 \cap \varepsilon^{-N} L_0} \right) = \dim \left(\frac{L'_0}{L'_0 \cap L_{nN}} \right) = \sum_{s > nN, t \leq 0} a_{s,t} = 0,$$

so it follows $L'_0 \subset \varepsilon^{-N} L_0$. Similarly,

$$\dim \left(\frac{\varepsilon^N L_0}{\varepsilon^N L_0 \cap L'_0} \right) = \dim \left(\frac{L_{-nN}}{L_{-nN} \cap L'_0} \right) = \sum_{s \leq -nN, t > 0} a_{s,t} = 0,$$

which shows $\varepsilon^N L_0 \subset L'_0$. Moreover,

$$\dim \left(\frac{\varepsilon^{-N} L_0}{L'_0} \right) = \dim \left(\frac{\varepsilon^{-N} L_0}{\varepsilon^{-N} L_0 \cap L'_0} \right) = \sum_{s \leq nN, t > 0} a_{s,t} = d_{nN,0}(A),$$

as a result of Lemma 3.2.12. \square

Assume $N \in \mathbb{N}$ is chosen so that $X_A^L \subset \Pi_{N,\lambda}^a(L)$, where $a = d_{nN,0}A$, as in Lemma 5.4.1.

Lemma 5.4.2. *X_A^L is a locally closed subset of $\Pi_{N,\lambda}^a(L)$. In particular, X_A^L is a quasiprojective variety.*

Proof. If $L' \in \Pi_{N,\lambda}^a(L)$ then

$$L_{-Nn} = \varepsilon^N L_0 \subset L'_0 \subset L'_1 \subset L'_n \subset \varepsilon^{-1-N} L_0 = L_{(N+1)n}.$$

Therefore X_A^L is the subset of $\Pi_{N,\lambda}^a(L)$ defined by the conditions $\dim(L_i/L_i \cap L'_j) = d_{i,j}A$ for $i : -Nn \leq i < j$ and $\dim(L'_j/L_i \cap L'_j) = \bar{d}_{i,j}A$ for $i : j < i \leq (N+1)n$, for $j = 1, \dots, n$.

The set of $L' \in \Pi_{N,\lambda}^a(L)$ with $\dim(L_i/L_i \cap L'_j) \leq d_{i,j}A$ for $j = 1, \dots, n$ and $i : -Nn \leq i < j$ and $\dim(L'_j/L_i \cap L'_j) \leq \bar{d}_{i,j}A$ for $j = 1, \dots, n$ and $i : j < i \leq (N+1)n$ is a closed subset of $\Pi_{N,\lambda}^a(L)$, as a result of Lemma 5.3.8.

On the other hand, the set of $L' \in \Pi_{N,\lambda}^a(L)$ satisfying the conditions $\dim(L_i/L_i \cap L'_j) \geq d_{i,j}A$ (for $i < j$) and $\dim(L'_j/L_i \cap L'_j) \geq \bar{d}_{i,j}A$ (for $i > j$) is open in $\Pi_{N,\lambda}^a(L)$ since the complement is closed, as a result of Lemma 5.3.8.

Therefore X_A^L is the intersection of an open set and a closed set in $\Pi_{N,\lambda}^a(L)$, so X_A^L is locally closed. It follows that X_A^L is an open subset of the projective variety $\overline{X_A^L}$, so is a quasiprojective variety as claimed. \square

Lemma 5.4.3. *X_A^L is irreducible.*

Proof. For any $L' \in X_A^L$, $X_A^L = G_L/H_{N,L} \cdot L'$. Lemma 5.3.6 shows that $G_L/H_{N,L}$ is a connected algebraic group which acts algebraically on $\Pi_{N,\lambda}^a(L)$. The image of $G_L/H_{N,L}$ under the morphism $g \mapsto gL'$ equals X_A^L , which shows X_A^L is irreducible since $G_L/H_{N,L}$ is irreducible. \square

Consequently, $\overline{X_A^L}$ is an irreducible projective variety and the action of $G_L/H_{N,L}$ on $\Pi_{N,\lambda}^a(L)$ restricts to an algebraic group action on $\overline{X_A^L}$ for which there are finitely many orbits. In particular, $\overline{X_A^L} \setminus X_A^L$ is a union of finitely many orbits which are so-called degenerations of the orbit X_A^L .

5.5 Geometry of orbit products

Let $A, B \in \Lambda_1$ with $\text{co}(A) = \text{ro}(B)$ and write $\lambda = \text{co}(A)$ and $\mu = \text{co}(B)$. Fix $L \in \mathcal{F}_{\text{ro}(A)}$. Recall

$$Y_{A,B}^L = \{(L', L'') \in \mathcal{F}_\lambda \times \mathcal{F}_\mu : L' \in X_A^L, L'' \in X_B^{L'}\}$$

and

$$X_{A,B}^L = \{L'' \in \mathcal{F}_\mu : \exists L' \in X_A^L \text{ with } L'' \in X_B^{L'}\}$$

Lemma 5.5.1. *There is $N \in \mathbb{N}$ such that*

$$Y_{A,B}^L \subset \Pi_{N,\lambda}^a(L) \times \Pi_{2N,\mu}^{a+b}(L),$$

where $a = d_{nN,0}(A)$ and $b = d_{nN,0}(B)$.

Proof. There is $N \in \mathbb{N}$ such that $\varepsilon^N L_0 \subset L'_0 \subset \varepsilon^{-N} L_0$ and $\varepsilon^N L'_0 \subset L''_0 \subset \varepsilon^{-N} L'_0$ for each $(L', L'') \in Y_{A,B}^L$, using Lemma 5.4.1. Set $a = d_{nN,0}(A)$ and $b = d_{nN,0}(B)$.

Then for any $(L', L'') \in Y_{A,B}^L$,

$$\varepsilon^{2N} L_0 \subset \varepsilon^N L'_0 \subset L''_0 \subset \varepsilon^{-N} L'_0 \subset \varepsilon^{-2N} L_0$$

and

$$\begin{aligned} \dim \left(\frac{\varepsilon^{-2N} L_0}{L''_0} \right) &= \dim \left(\frac{\varepsilon^{-N} L'_0}{L''_0} \right) + \dim \left(\frac{\varepsilon^{-2N} L_0}{\varepsilon^{-N} L'_0} \right) \\ &= \dim \left(\frac{\varepsilon^{-N} L'_0}{L''_0} \right) + \dim \left(\frac{\varepsilon^{-N} L_0}{L'_0} \right) \\ &= a + b, \end{aligned}$$

as a result of Lemma 3.2.12, so $(L', L'') \in \Pi_{N,\lambda}^a(L) \times \Pi_{2N,\mu}^{a+b}(L)$ as required. \square

Now assume $N \in \mathbb{N}$ is chosen so that $Y_{A,B}^L \subset \Pi_{N,\lambda}^a(L) \times \Pi_{2N,\mu}^{a+b}(L)$, where $a = d_{nN,0}(A)$ and $b = d_{nN,0}(B)$, using Lemma 5.5.1.

Lemma 5.5.2. *$Y_{A,B}^L$ is a locally closed subset of $\Pi_{N,\lambda}^a(L) \times \Pi_{2N,\mu}^{a+b}(L)$. In particular, $Y_{A,B}^L$ is a quasiprojective variety.*

Proof. $Y_{A,B}^L$ is the subset of $\Pi_{N,\lambda}^a(L) \times \Pi_{2N,\mu}^{a+b}(L)$ consisting of those (L', L'') satisfying the following conditions: $\dim(L_i/L_i \cap L'_j) = d_{i,j}(A)$ for $i < j$, $\dim(L'_j/L_i \cap L'_j) = \bar{d}_{i,j}(A)$ for $i > j$, $\dim(L'_i/L'_i \cap L''_j) = d_{i,j}(B)$ for $i < j$ and $\dim(L''_j/L'_i \cap L''_j) = \bar{d}_{i,j}(B)$. Only finitely many conditions are required to define $Y_{A,B}^L$ since there are only finitely many nonzero entries in A and B modulo the (n, n) -periodicity.

The conditions $\dim(L_i/L_i \cap L'_j) \leq d_{i,j}(A)$, $\dim(L'_j/L'_i \cap L''_j) \leq d_{i,j}(B)$, $\dim(L'_j/L_i \cap L'_j) \leq \bar{d}_{i,j}(A)$ and $\dim(L''_j/L'_i \cap L''_j) \leq \bar{d}_{i,j}(B)$ define closed subsets of $\Pi_{N,\lambda}^a(L) \times \Pi_{2N,\mu}^{a+b}(L)$ for each $i, j \in \mathbb{Z}$, as a result of Lemma 5.3.7 and Lemma 5.3.8.

On the other hand, the conditions $\dim(L_i/L_i \cap L'_j) \geq d_{i,j}(A)$, $\dim(L'_j/L'_i \cap L''_j) \geq d_{i,j}(B)$, $\dim(L'_j/L_i \cap L'_j) \geq \bar{d}_{i,j}(A)$ and $\dim(L''_j/L'_i \cap L''_j) \geq \bar{d}_{i,j}(B)$ define open subsets of $\Pi_{N,\lambda}^a(L) \times \Pi_{2N,\mu}^{a+b}(L)$ for each $i, j \in \mathbb{Z}$, using Lemma 5.3.7 and Lemma 5.3.8.

Therefore $Y_{A,B}^L$ is the intersection of finitely many open sets and finitely many closed sets in $\Pi_{N,\lambda}^a(L) \times \Pi_{2N,\mu}^{a+b}(L)$, so $Y_{A,B}^L$ is locally closed. In particular, $Y_{A,B}^L$ is a quasiprojective variety. \square

Lemma 5.5.3. *For any $L' \in X_A^L$, $Y_{A,B}^L = G_L \cdot (\{L'\} \times X_B^{L'})$.*

Proof. Let $L' \in X_A^L$, then $\{L'\} \times X_B^{L'}$ is contained in $Y_{A,B}^L$ and G_L acts on $Y_{A,B}^L$, so $G_L \cdot (\{L'\} \times X_B^{L'})$ is contained in $Y_{A,B}^L$. If $(N', N'') \in Y_{A,B}^L$, then $N' = \sigma L'$ for some $\sigma \in G_L$, since $N' \in X_A^L$. Then $(N', N'') = \sigma(L', \sigma^{-1}N'')$ and $\sigma^{-1}N'' \in X_B^{\sigma^{-1}N'} = X_B^{L'}$, so $(N', N'') \in \sigma \cdot (\{L'\} \times X_B^{L'})$. Therefore $Y_{A,B}^L = G_L \cdot (\{L'\} \times X_B^{L'})$ as claimed. \square

Proposition 5.5.4. *$Y_{A,B}^L$ is irreducible.*

Proof. Let $L' \in X_A^L$. $G_L/H_{2N,L}$ is a connected algebraic group acting algebraically on $\Pi_{N,\lambda}^a(L) \times \Pi_{2N,\mu}^{a+b}(L)$ by Lemma 5.3.6. $X_B^{L'}$ is an irreducible locally closed subset of $\Pi_{2N,\mu}^{a+b}(L)$, so $\{L'\} \times X_B^{L'}$ is an irreducible locally closed set in $\Pi_{N,\lambda}^a(L) \times \Pi_{2N,\mu}^{a+b}(L)$. $Y_{A,B}^L = G_L \cdot (\{L'\} \times X_B^{L'}) = G_L/H_{2N,L} \cdot (\{L'\} \times X_B^{L'})$, by Lemma 5.5.3, so it follows that $Y_{A,B}^L$ is irreducible. \square

Let p_2 be the projection onto the second factor $\Pi_{N,\lambda}^a(L) \times \Pi_{2N,\mu}^{a+b}(L) \rightarrow \Pi_{2N,\mu}^{a+b}(L)$. p_2 is a closed morphism since $\Pi_{N,\lambda}^a(L)$ is a projective variety and therefore complete, by Lemma 5.3.2. Therefore $p_2(\overline{Y_{A,B}^L}) = \overline{X_{A,B}^L}$, since $p_2(Y_{A,B}^L) = X_{A,B}^L$.

Lemma 5.5.5. *$X_{A,B}^L$ is irreducible and constructible.*

Proof. Proposition 5.5.4 shows that $Y_{A,B}^L$ is irreducible and locally closed, so it follows $X_{A,B}^L$ is irreducible and constructible, since $X_{A,B}^L = p_2(Y_{A,B}^L)$. \square

Proposition 5.5.6. *There is a unique open G_L -orbit in $X_{A,B}^L$.*

Proof. $X_{A,B}^L$ consists of finitely many G_L -orbits and is an irreducible topological space, by Lemma 5.5.5. Consequently, X_C^L is dense in $X_{A,B}^L$ for some $C \in \Lambda_1^{A,B}$. Lemma 5.4.2 shows that X_C^L is locally closed in $X_{A,B}^L$, so X_C^L is open in $\overline{X_C^L} = X_{A,B}^L$. Irreducibility of $X_{A,B}^L$ shows that there is a unique open G_L -orbit, since two nonempty open sets in $X_{A,B}^L$ intersect nontrivially, thus any two open G_L orbits in $X_{A,B}^L$ coincide. \square

Let $A * B \in \Lambda_1$ be the matrix corresponding to the dense open G_L -orbit in $X_{A,B}^L$, so $\overline{X_{A*B}^L} = \overline{X_{A,B}^L}$.

5.6 Degenerations of orbits and the combinatorial partial order

Theorem 5.6.1. *Let $A, B \in \Lambda_1$ with $\text{ro}(A) = \text{ro}(B)$ and $\text{co}(A) = \text{co}(B)$, then $B \leq A$ if and only if $X_B^L \subset \overline{X_A^L}$ for any $L \in \mathcal{F}_{\text{ro}(A)}$.*

Proof. Let $\lambda = \text{co}(A)$, $\mu = \text{ro}(A)$ and fix $L \in \mathcal{F}_\mu$. Assume $N \in \mathbb{N}$ is chosen so that $X_A^L \subset \Pi_{N,\lambda}^a(L)$ and $X_B^L \subset \Pi_{N,\lambda}^b(L)$, where $a = d_{nN,0}(A)$ and $b = d_{nN,0}(B)$. Then X_A^L is an open subset of the projective variety consisting of those $L' \in \Pi_{N,\lambda}^a(L)$ such that

$$\dim \left(\frac{L_i}{L_i \cap L'_j} \right) \leq d_{i,j}(A)$$

and

$$\dim \left(\frac{L'_j}{L_i \cap L'_j} \right) \leq \bar{d}_{i,j}(A),$$

for all $i, j \in \mathbb{Z}$.

Assume $X_B^L \subset \overline{X_A^L}$, then

$$d_{i,j}(B) = \dim \left(\frac{L_i}{L_i \cap L'_j} \right) \leq d_{i,j}(A)$$

and

$$\bar{d}_{i,j}(B) = \dim \left(\frac{L'_j}{L_i \cap L'_j} \right) \leq \bar{d}_{i,j}(A),$$

for each $i, j \in \mathbb{Z}$, for any $L' \in X_B^L$. So $B \leq A$ if $X_B^L \leq \overline{X_A^L}$.

Conversely, suppose $A \leq B$.

□

Corollary 5.6.2. *The maximum in $\Lambda_1^{A,B}$ is $A * B$.*

5.7 Associativity of the generic product

Let $A, B, C \in \Lambda_1$ with $\text{co}(A) = \text{ro}(B)$ and $\text{co}(B) = \text{ro}(C)$ and fix $L \in \mathcal{F}_{\text{ro}(A)}$. Write $\lambda = \text{co}(A)$, $\mu = \text{co}(B)$ and $\nu = \text{co}(C)$. Define

$$Y_{A,B,C}^L = \left\{ (L', L'', L''') \in \mathcal{F}^3 : L' \in X_A^L, L'' \in X_B^L, L''' \in X_C^L \right\}$$

and

$$X_{A,B,C}^L = \left\{ L''' \in \mathcal{F} : \exists (L', L'') \in \mathcal{F}^2 \text{ with } (L', L'', L''') \in Y_{A,B,C}^L \right\}.$$

Lemma 5.7.1. *There is $N \in \mathbb{N}$ such that $Y_{A,B,C}^L$ is contained in $\Pi_{N,\lambda}^a(L) \times \Pi_{2N,\mu}^{a+b}(L) \times \Pi_{3N,\nu}^{a+b+c}(L)$, where $a = d_{nN,0}(A)$, $b = d_{nN,0}(B)$ and $c = d_{nN,0}(C)$.*

Proof. Lemma 5.4.1 shows that there is $N \in \mathbb{N}$ such that $\varepsilon^N L_0 \subset L'_0 \subset \varepsilon^{-N} L_0$, $\varepsilon^N L'_0 \subset L''_0 \subset \varepsilon^{-N} L'_0$ and $\varepsilon^N L''_0 \subset L'''_0 \subset \varepsilon^{-N} L''_0$ for each $(L', L'', L''') \in Y_{A,B,C}^L$. Using the proof of Lemma 5.5.1, it follows $L'' \in \Pi_{2N,\mu}^{a+b}(L)$ and $L''' \in \Pi_{2N,\nu}^{b+c}(L') \subset \Pi_{3N,\nu}^{a+b+c}(L)$. □

Assume $N \in \mathbb{N}$ is chosen so that $Y_{A,B,C}^L \subset \Pi_{N,\lambda}^a(L) \times \Pi_{2N,\mu}^{a+b}(L) \times \Pi_{3N,\nu}^{a+b+c}(L)$, where $a = d_{nN,0}(A)$, $b = d_{nN,0}(B)$ and $c = d_{nN,0}(C)$, as in Lemma 5.7.1.

Lemma 5.7.2. *$Y_{A,B,C}^L$ is a locally closed subset of $\Pi_{N,\lambda}^a(L) \times \Pi_{2N,\mu}^{a+b}(L) \times \Pi_{3N,\nu}^{a+b+c}(L)$. In particular, $Y_{A,B,C}^L$ is a quasiprojective variety.*

Proof. Write $\Pi = \Pi_{N,\lambda}^a(L) \times \Pi_{2N,\mu}^{a+b}(L) \times \Pi_{3N,\nu}(L)$. Then $Y_{A,B,C}^L$ consists of those $(L', L'', L''') \in \Pi$ satisfying the following conditions:

$$\dim \left(\frac{L_i}{L_i \cap L'_j} \right) = d_{i,j}(A), \quad (5.7.1)$$

$$\dim \left(\frac{L'_i}{L'_i \cap L''_j} \right) = d_{i,j}(B), \quad (5.7.2)$$

$$\dim \left(\frac{L''_i}{L''_i \cap L'''_j} \right) = d_{i,j}(C), \quad (5.7.3)$$

for $(i, j) \in \{1, \dots, n\} \times \mathbb{Z}$ with $i < j < (N+1)n$, and

$$\dim \left(\frac{L'_j}{L_i \cap L'_j} \right) = \bar{d}_{i,j}(A), \quad (5.7.4)$$

$$\dim \left(\frac{L''_j}{L'_i \cap L''_j} \right) = \bar{d}_{i,j}(B), \quad (5.7.5)$$

$$\dim \left(\frac{L'''_j}{L''_i \cap L'''_j} \right) = \bar{d}_{i,j}(C), \quad (5.7.6)$$

for $(i, j) \in \{1, \dots, n\} \times \mathbb{Z}$ with $-Nn < j < i$.

For $i < j$, the conditions

$$\begin{aligned} \dim(L_i/L_i \cap L'_j) &\leq d_{i,j}(A), \\ \dim(L'_i/L'_i \cap L''_j) &\leq d_{i,j}(B) \end{aligned}$$

and

$$\dim(L''_i/L''_i \cap L'''_j) \leq d_{i,j}(C)$$

define closed subsets of Π , by Lemma 5.3.7. For $i > j$, the conditions

$$\begin{aligned} \dim(L'_j/L_i \cap L'_j) &\leq \bar{d}_{i,j}(A), \\ \dim(L''_j/L'_i \cap L''_j) &\leq \bar{d}_{i,j}(B) \end{aligned}$$

and

$$\dim(L'''_j/L''_i \cap L'''_j) \leq \bar{d}_{i,j}(C)$$

also define closed subsets of Π .

On the other hand, the conditions $\dim(L_i/L_i \cap L'_j) \geq d_{i,j}(A)$, $\dim(L'_i/L'_i \cap L''_j) \geq d_{i,j}(B)$ and $\dim(L''_i/L''_i \cap L'''_j) \geq d_{i,j}(C)$ for $i < j$ define open subsets of Π . Similarly, the conditions $\dim(L'_j/L_i \cap L'_j) \geq \bar{d}_{i,j}(A)$, $\dim(L''_j/L'_i \cap L''_j) \geq \bar{d}_{i,j}(B)$ and $\dim(L'''_j/L''_i \cap L'''_j) \geq \bar{d}_{i,j}(C)$ for $i > j$ define open subsets of Π .

Therefore $Y_{A,B,C}^L$ is the intersection of finitely many closed sets in Π with finitely many open subsets of Π , so $Y_{A,B,C}^L$ is locally closed. In particular, $Y_{A,B,C}^L$ is a quasiprojective variety. \square

Lemma 5.7.3. For any $(L', L'', L''') \in Y_{A,B,C}^L$,

$$Y_{A,B,C}^L = \{ \alpha \cdot (L', \beta L'', \beta \gamma L''') : \alpha \in G_L, \beta \in G_{L'}, \gamma \in G_{L''} \}.$$

In particular,

$$Y_{A,B,C}^L = G_L \cdot (\{L'\} \times Y_{B,C}^{L'})$$

for each $L' \in X_A^L$.

Proof. Let $(L', L'', L''') \in Y_{A,B,C}^L$. Given $\alpha \in G_L$, $\beta \in G_{L'}$ and $\gamma \in G_{L''}$, $(\alpha L', \alpha \beta L'', \alpha \beta \gamma L''')$ is in $Y_{A,B,C}^L$ since

$$\begin{aligned} (L, \alpha L') &= \alpha(L, L') \in \mathcal{O}_A \\ (\alpha L', \alpha \beta L'') &= \alpha \beta (L', L'') \in \mathcal{O}_B \\ (\alpha \beta L'', \alpha \beta \gamma L''') &= \alpha \beta \gamma (L'', L''') \in \mathcal{O}_C \end{aligned}$$

For each $(N', N'', N''') \in Y_{A,B,C}^L$ there exist $\sigma_1, \sigma_2, \sigma_3 \in G$ with

$$\begin{aligned} (L, N') &= \sigma_1(L, L') \\ (N', N'') &= \sigma_2(L', L'') \\ (N'', N''') &= \sigma_3(L'', L'''). \end{aligned}$$

Let $\alpha = \sigma_1$, $\beta = \sigma_1^{-1} \sigma_2$ and $\gamma = \sigma_2^{-1} \sigma_3$, so $\sigma_2 = \alpha \beta$ and $\sigma_3 = \alpha \beta \gamma$. It follows that

$$(N', N'', N''') = (\alpha L', \alpha \beta L'', \alpha \beta \gamma L'''),$$

which proves the first claim. The second claim follows from the first since $(L'', L''') \in Y_{B,C}^{L'}$ and therefore

$$Y_{B,C}^{L'} = \{ (\beta L'', \beta \gamma L''') : \beta \in G_{L'}, \gamma \in G_{L''} \},$$

as required. \square

Proposition 5.7.4. $Y_{A,B,C}^L$ is irreducible.

Proof. Write

$$\Pi = \Pi_{N,\lambda}^a(L) \times \Pi_{2N,\mu}^{a+b}(L) \times \Pi_{3N,\nu}^{a+b+c}(L).$$

Lemma 5.3.2 shows that Π is a projective algebraic variety and Lemma 5.3.6 shows that $G_L/H_{3N,L}$ is a connected algebraic group acting algebraically on Π by the diagonal action.

Let $L' \in X_A^L$. As a result of Lemma 5.7.3

$$\begin{aligned} Y_{A,B,C}^L &= G_L \cdot (\{L'\} \times Y_{B,C}^{L'}) \\ &= G_L/H_{3N,L} \cdot (\{L'\} \times Y_{B,C}^{L'}). \end{aligned}$$

Proposition 5.5.4 shows that $Y_{B,C}^{L'}$ is irreducible, so $\{L'\} \times Y_{B,C}^{L'}$ is irreducible. The image of $\{L'\} \times Y_{B,C}^{L'}$ under the action of $G_L/H_{3N,L}$ is irreducible, since $G_L/H_{3N,L}$ is connected and therefore irreducible. Therefore $Y_{A,B,C}^L$ is irreducible. \square

Let p_3 be the projection of $\Pi_{N,\lambda}^a(L) \times \Pi_{2N,\mu}^{a+b}(L) \times \Pi_{3N,\nu}^{a+b+c}(L)$ onto the third factor. By the completeness property of projective varieties, p_3 is a closed morphism. The image of $Y_{A,B,C}^L$ under p_3 is $X_{A,B,C}^L$, so $p_3(\overline{Y_{A,B,C}^L}) = \overline{X_{A,B,C}^L}$.

Lemma 5.7.5. $X_{A,B,C}^L$ is irreducible and constructible.

Proof. Lemma 5.7.2 and Proposition 5.7.4 show that $Y_{A,B,C}^L$ is locally closed and irreducible. It follows $X_{A,B}^L$ is irreducible and constructible, since $X_{A,B,C}^L$ is the image of $Y_{A,B,C}^L$ under the morphism p_3 . \square

Lemma 5.7.6. There is a unique open and dense G_L -orbit in $X_{A,B,C}^L$.

Proof. There are only finitely many G_L -orbits in $X_{A,B,C}^L$. In particular,

$$X_{A,B,C}^L = \bigcup_{D \in \Lambda_1^{A,B}} X_{D,C}^L = \bigcup_{D \in \Lambda_1^{A,B}} \bigcup_{D' \in \Lambda_1^{D,C}} X_{D'}^L$$

and

$$\overline{X_{A,B,C}^L} = \bigcup_{D \in \Lambda_1^{A,B}} \bigcup_{D' \in \Lambda_1^{D,C}} \overline{X_{D'}^L}.$$

There is $D \in \Lambda_1$ such that $\overline{X_D^L} = \overline{X_{A,B,C}^L}$, since $X_{A,B,C}^L$ is irreducible, by Lemma 5.7.5. By Lemma 5.4.2, X_D^L is open in $\overline{X_D^L} = \overline{X_{A,B,C}^L}$, so X_D^L is open in $X_{A,B,C}^L$.

If X_D^L and $X_{D'}^L$ are open in $X_{A,B,C}^L$, then X_D^L and $X_{D'}^L$ have nonempty intersection since $X_{A,B,C}^L$ is irreducible, then $X_D^L = X_{D'}^L$. \square

Lemma 5.7.7. $p_3^{-1}(X_{A*B,C}^L)$ is open in $\overline{Y_{A,B,C}^L}$.

Proof. Projection onto the second component is a closed morphism of varieties $p_2: \overline{Y_{A,B,C}^L} \rightarrow \overline{X_{A,B}^L}$ with $p_2(Y_{A,B,C}^L) = X_{A,B}^L$. It follows that $p_3^{-1}(X_{A*B,C}^L)$ is open in $\overline{Y_{A,B,C}^L}$ since $p_3^{-1}(X_{A*B,C}^L) = p_2^{-1}(X_{A*B}^L)$ and X_{A*B}^L is open in $\overline{X_{A,B}^L}$. \square

Lemma 5.7.8. $p_3^{-1}(X_{A,B*C}^L)$ is open in $\overline{Y_{A,B,C}^L}$.

Proof. $p_3^{-1}(X_{A,B*C}^L)$ consists of those $(L', L'', L''') \in \overline{Y_{A,B,C}^L}$ such that $\dim(L'_i/L'_i \cap L'''_j) \geq d_{i,j}(B * C)$ for $i < j$ and $\dim(L'''_j/L'_i \cap L'''_j) \geq \bar{d}_{i,j}(B * C)$ for $i > j$. Each of these conditions defines an open subset of $\overline{Y_{A,B,C}^L}$ as a result of Lemma 5.3.7 and only finitely many conditions are required to determine $p_3^{-1}(X_{A,B*C}^L)$, as before. Therefore $p_3^{-1}(X_{A,B*C}^L)$ is the intersection of finitely many open sets in $\overline{Y_{A,B,C}^L}$, so is open as claimed. \square

Proposition 5.7.9. $X_{A*(B*C)}^L = X_{(A*B)*C}^L$.

Proof. The unique open G_L -orbit in $X_{A*B,C}^L$ is $X_{(A*B)*C}^L$, so $p_3^{-1}(X_{(A*B)*C}^L)$ is open in $p_3^{-1}(X_{A*B,C}^L)$. Lemma 5.7.7 shows that $p_3^{-1}(X_{A*B,C}^L)$ is open in $\overline{Y_{A,B,C}^L}$, so $p_3^{-1}(X_{(A*B)*C}^L)$ is open in $\overline{Y_{A,B,C}^L}$.

Similarly, $X_{A*(B*C)}^L$ is open in $X_{A,B*C}^L$, so $p_3^{-1}(X_{A*(B*C)}^L)$ is open in $p_3^{-1}(X_{A,B*C}^L)$. Lemma 5.7.8 shows that $p_3^{-1}(X_{A,B*C}^L)$ is open in $\overline{Y_{A,B,C}^L}$, so it follows $p_3^{-1}(X_{A*(B*C)}^L)$ is open in $\overline{Y_{A,B,C}^L}$.

Therefore $f^{-1}(X_{A*(B*C)}^L)$ has nonempty intersection with $f^{-1}(X_{(A*B)*C}^L)$, since $Y_{A,B,C}^L$ is irreducible by Proposition 5.7.4. It follows that the G_L -orbits $X_{A*(B*C)}^L$ and $X_{(A*B)*C}^L$ have nonempty intersection and therefore $X_{A*(B*C)}^L$ equals $X_{(A*B)*C}^L$. \square

5.8 The generic affine algebra

The generic affine algebra of rank r and period n , denoted by $\hat{G}(n, r)$, is a free \mathbb{Z} -module with basis $\{e_A : A \in \Lambda_1\}$ and \mathbb{Z} -bilinear multiplication given by

$$e_A * e_B = e_{A*B}$$

for $A, B \in \Lambda_1$ with $\text{co}(A) = \text{ro}(B)$, and

$$e_A * e_B = 0$$

for $A, B \in \Lambda_1$ with $\text{co}(A) \neq \text{ro}(B)$.

Proposition 5.8.1. *The generic algebra $\hat{G}(n, r)$ is an associative \mathbb{Z} -algebra with 1, with*

$$1 = \sum_{\lambda \in \Lambda_0} 1_\lambda$$

where

$$1_\lambda = e_{D_\lambda},$$

for each $\lambda \in \Lambda_0$.

Proof. Let $A, B, C \in \Lambda_1$. If $\text{co}(A) \neq \text{ro}(B)$ or $\text{co}(B) \neq \text{ro}(C)$, then

$$(e_A * e_B) * e_C = 0 = e_A * (e_B * e_C),$$

so we may now suppose $\text{co}(A) = \text{ro}(B)$ and $\text{co}(B) = \text{ro}(C)$.

As a result of Proposition 5.7.9,

$$\begin{aligned} (e_A * e_B) * e_C &= e_{(A*B)*C} \\ &= e_{A*(B*C)} \\ &= e_A * (e_B * e_C), \end{aligned}$$

so it follows $\hat{G}(n, r)$ is an associative \mathbb{Z} -algebra.

The expression for the multiplicative identity follows from Lemma 3.2.16, since

$$e_A * \left(\sum_{\lambda \in \Lambda_0} 1_\lambda \right) = e_A * 1_{\text{co}(A)} = e_A$$

and

$$\left(\sum_{\lambda \in \Lambda_0} 1_\lambda \right) * e_A = 1_{\text{ro}(A)} * e_A = e_A,$$

for each $A \in \Lambda_1$. □

5.8.1 A categorical perspective

Proposition 5.8.2. *The following constitutes a small category: the set of objects is Λ_0 and the set of morphisms is Λ_1 . Given compositions $\lambda, \mu \in \Lambda_0$, the morphisms with source μ and target λ are those matrices $A \in \Lambda_1$ with $\text{co}(A) = \mu$ and $\text{ro}(A) = \lambda$. Given $\lambda, \mu, \nu \in \Lambda_0$ and $A, B \in \Lambda_1$ with $\text{co}(B) = \nu$, $\text{ro}(B) = \mu = \text{co}(A)$ and $\text{ro}(A) = \lambda$, their composition is $A * B$, with source $\text{co}(A * B) = \text{co}(B) = \nu$ and target $\text{ro}(A * B) = \text{ro}(A) = \lambda$.*

Proof. Proposition 5.7.9 shows that the generic product $*$ is associative. For each object $\lambda \in \Lambda_0$, the identity morphism $\lambda \rightarrow \lambda$ is the diagonal matrix D_λ . \square

Then the generic affine algebra $\hat{G}(n, r)$ may be realised as the \mathbb{Z} -algebra of this category. Observe that there are only finitely many objects in this category and distinct objects are non-isomorphic, so the isomorphism classes in this category are in one to one correspondence with Λ_0 . The \mathbb{Z} -algebra of this category is the free \mathbb{Z} -module on Λ_1 with \mathbb{Z} -bilinear multiplication given by the generic product $*$.

Chapter 6

A realisation of affine zero Schur algebras

The purpose of this chapter is to study the link between the generic affine algebra $\hat{G}(n, r)$ and the affine 0-Schur algebra $\hat{S}_0(n, r)$.

The main result is the construction of an isomorphism of \mathbb{Z} -algebras from $\hat{G}(n, r)$ to $\hat{S}_0(n, r)$ such that $E_i \mapsto E_i$, $F_i \mapsto F_i$ and $1_\lambda \mapsto 1_\lambda$, in the case that $n, r \geq 1$ with $r < n$.

6.1 Preliminary results on the generic affine algebra

Recall that the generic affine algebra $\hat{G}(n, r)$ is an associative \mathbb{Z} -algebra with a multiplicative basis $\{e_A : A \in \Lambda_1\}$ over \mathbb{Z} , where

$$e_A * e_B = e_{A*B}$$

for $A, B \in \Lambda_1$ with $\text{co}(A) = \text{ro}(B)$, and

$$e_A * e_B = 0$$

for $A, B \in \Lambda_1$ with $\text{co}(A) \neq \text{ro}(B)$.

6.1.1 Elementary basis elements

For $i \in \{1, \dots, n\}$ and $\lambda \in \Lambda_0$ such that $\lambda_{i+1} > 0$, define

$$E_{i,\lambda} = e_{D_\lambda + \mathcal{E}_{i,i+1} - \mathcal{E}_{i+1,i+1}}$$

and let

$$E_i = \sum_{\lambda \in \Lambda_0 : \lambda_{i+1} > 0} E_{i,\lambda}$$

for each $i \in \{1, \dots, n\}$

For $i \in \{1, \dots, n\}$ and $\lambda \in \Lambda_0$ such that $\lambda_i > 0$, define

$$F_{i,\lambda} = e_{D_\lambda + \mathcal{E}_{i+1,i} - \mathcal{E}_{i,i}}$$

and let

$$F_i = \sum_{\lambda \in \Lambda_0 : \lambda_i > 0} F_{i,\lambda}$$

for each $i \in \{1, \dots, n\}$.

Lemma 6.1.1. *Let $i \in \{1, \dots, n\}$ and $A \in \Lambda_1$ and write $\mu = \text{ro}(A)$.*

*If $\mu_{i+1} = 0$ then $E_i * e_A = 0$. If $\mu_{i+1} > 0$, then*

$$E_i * e_A = e_{A+\mathcal{E}_{i,p}-\mathcal{E}_{i+1,p}},$$

where

$$p = \max\{j \in \mathbb{Z} : a_{i+1,j} > 0\}.$$

*If $\mu_i = 0$ then $F_i * e_A = 0$. If $\mu_i > 0$ then*

$$F_i * e_A = e_{A+\mathcal{E}_{i+1,q}-\mathcal{E}_{i,q}},$$

where

$$q = \min\{j \in \mathbb{Z} : a_{i,j} > 0\}.$$

Proof. Suppose $\mu_{i+1} > 0$. Recall that the corresponding product in the affine q -Schur algebra $\hat{S}_q(n, r)$ is

$$E_i \cdot e_A = \sum_{j \in \mathbb{Z} : a_{i+1,j} > 0} q^{\sum_{t > j} a_{i,t}} [[a_{i,j} + 1]] e_{A+\mathcal{E}_{i,j}-\mathcal{E}_{i+1,j}},$$

by Lemma 4.1.2.

Suppose $B \in \Lambda_1$ with $B = A + \mathcal{E}_{i,j} - \mathcal{E}_{i+1,j}$ for some $j \in \mathbb{Z}$. For $s \in \{1, \dots, n\}$ and $t \in \mathbb{Z}$,

$$d_{s,t}(B) = \begin{cases} d_{s,t}(A) + 1 & : s = i \text{ and } t < j, \\ d_{s,t}(A) & : \text{otherwise,} \end{cases}$$

and

$$\bar{d}_{s,t}(B) = \begin{cases} \bar{d}_{s,t}(A) - 1 & : s = i \text{ and } t \geq j, \\ \bar{d}_{s,t}(A) & : \text{otherwise.} \end{cases}$$

It follows that if $j' < j$, then

$$A + \mathcal{E}_{i,j'} - \mathcal{E}_{i+1,j'} < A + \mathcal{E}_{i,j} - \mathcal{E}_{i+1,j}.$$

Therefore, the product in $\hat{G}(n, r)$ is given by

$$E_i * e_A = e_{A+\mathcal{E}_{i,p}-\mathcal{E}_{i+1,p}},$$

where

$$p = \max\{j \in \mathbb{Z} : a_{i+1,j} > 0\}.$$

The argument for the action of F_i is similar, but there is a pleasing symmetry in the two proofs.

Now suppose $\mu_i > 0$. Using Lemma 4.1.2,

$$F_i \cdot e_A = \sum_{j \in \mathbb{Z} : a_{i,j} > 0} q^{\sum_{t < j} a_{i+1,t}} [[a_{i+1,j} + 1]] e_{A+\mathcal{E}_{i+1,j}-\mathcal{E}_{i,j}},$$

in $\hat{S}_q(n, r)$.

Suppose $B \in \Lambda_1$ with $B = A + \mathcal{E}_{i+1,j} - \mathcal{E}_{i,j}$, for some $j \in \mathbb{Z}$. Then for $i \in \{1, \dots, n\}$ and $j \in \mathbb{Z}$,

$$d_{s,t}(B) = \begin{cases} d_{s,t}(A) - 1 & : s = i \text{ and } t < j, \\ d_{s,t}(A) & : \text{otherwise,} \end{cases}$$

and

$$\bar{d}_{s,t}(B) = \begin{cases} \bar{d}_{s,t}(A) + 1 & : s = i \text{ and } t \geq j, \\ \bar{d}_{s,t}(A) & : \text{otherwise.} \end{cases}$$

Then if $j' < j$ it follows

$$A + \mathcal{E}_{i+1,j'} - \mathcal{E}_{i,j'} > A + \mathcal{E}_{i+1,j} - \mathcal{E}_{i,j},$$

so the terms with nonzero coefficients in the product $F_i \cdot e_A$ are totally ordered and the maximum is

$$F_i * e_A = e_{A + \mathcal{E}_{i+1,q} - \mathcal{E}_{i,q}},$$

where $q = \min\{j \in \mathbb{Z} : a_{i,j} > 0\}$. □

6.1.2 Transpose involution

Let S be the \mathbb{Z} -module automorphism of $\hat{G}(n, r)$ given by

$$S(e_A) = e_{A^\top}$$

for each $A \in \Lambda_1$.

Lemma 6.1.2. *The map S is a \mathbb{Z} -algebra antihomomorphism. In particular,*

$$e_{A^\top} * e_{B^\top} = e_B * e_A,$$

for each $A, B \in \Lambda_1$.

Proof. Lemma 4.1.8 show that the transpose preserves the partial order on Λ_1 and so

$$(B * A)^\top = A^\top * B^\top,$$

using Lemma 4.1.1. □

For any $A \in \Lambda_1$,

$$S(S(e_A)) = e_{(A^\top)^\top} = e_A,$$

so $S \circ S$ is the identity map on $\hat{S}_q(n, r)$.

For each $i \in \{1, \dots, n\}$ and $\lambda \in \Lambda_0$ with $\lambda_{i+1} > 0$,

$$S(E_{i,\lambda}) = F_{i,\lambda + \alpha_i},$$

for each $i \in \{1, \dots, n\}$ and $\lambda \in \Lambda_0$ with $\lambda_i > 0$,

$$S(F_{i,\lambda}) = E_{i,\lambda - \alpha_i}, \text{ and}$$

and

$$S(1_\lambda) = 1_\lambda,$$

for each $\lambda \in \Lambda_0$.

Lemma 6.1.3. *Let $i \in \{1, \dots, n\}$ and $A \in \Lambda_1$ and write $\lambda = \text{co}(A)$.*

*If $\lambda_j = 0$ then $e_A * E_j = 0$. If $\lambda_j > 0$ then*

$$e_A * E_j = e_{A+\varepsilon_{p,j+1}-\varepsilon_{p,j}},$$

where

$$p = \min\{i \in \mathbb{Z} : a_{i,j} > 0\}.$$

*If $\lambda_{j+1} = 0$ then $e_A * F_j = 0$. If $\lambda_{j+1} > 0$ then*

$$e_A * F_j = e_{A+\varepsilon_{q,j}-\varepsilon_{q,j+1}},$$

where

$$q = \max\{i \in \mathbb{Z} : a_{i,j+1} > 0\}.$$

Proof. This follows immediately on applying the transpose involution to the formulas for the action of E_i and F_i on the left given in Lemma 6.1.1.

Equally, this result can be proven directly using the formulas for the action of E_i and F_i on the right in Lemma 4.1.3, as in the proof of Lemma 6.1.1. \square

6.1.3 Shifting and periodicity

For each $\lambda \in \Lambda_0$, define

$$R_\lambda = e_{[1]D_\lambda} = e_{\lambda_1\varepsilon_{0,1}+\dots+\lambda_n\varepsilon_{n-1,n}}$$

and set

$$R = \sum_{\lambda \in \Lambda_0} R_\lambda.$$

Lemma 6.1.4. *For each $A \in \Lambda_1$,*

$$R * e_A = e_{[1]A}$$

and

$$e_A * R = e_{A[-1]}.$$

Proof. Lemma 4.1.12 shows that the same formulas hold in $\hat{S}_q(n, r)$, then the result follows for the generic multiplication $*$, since each product $R * e_A$ and $e_A * R$ is supported on one orbit, so the generic multiplication and the product on $\hat{S}_q(n, r)$ are the same in this instance. \square

Observe that

$$\begin{aligned} S(R_\lambda) &= e_{\lambda_1\varepsilon_{1,0}+\dots+\lambda_n\varepsilon_{n,n-1}} \\ &= e_{[-1]D_{[1]\lambda}} \end{aligned}$$

so

$$S(R) = \sum_{\lambda \in \Lambda_0} e_{[-1]D_\lambda}.$$

Lemma 6.1.5. *The element R of $\hat{G}(n, r)$ is invertible, with*

$$R * S(R) = 1 = S(R) * R.$$

Proof. Lemma 6.1.4 shows that

$$\begin{aligned} R * S(R)1_\lambda &= Re_{[-1]D_{[1]}\lambda} \\ &= e_{D_{[1]}\lambda} \\ &= 1_{[1]\lambda} \end{aligned}$$

for each $\lambda \in \Lambda_0$, so

$$R * S(R) = 1.$$

Similarly,

$$\begin{aligned} S(R) * R &= \sum_{\lambda \in \Lambda_0} e_{D_\lambda[1]} * R \\ &= \sum_{\lambda \in \Lambda_0} e_{D_\lambda} \\ &= 1. \end{aligned}$$

□

Let τ be the \mathbb{Z} -algebra automorphism of $\hat{G}(n, r)$ given by conjugation by R , so

$$\begin{aligned} \tau(e_A) &= R * e_A * S(R) \\ &= R * e_A * R^{-1}, \end{aligned}$$

for each $A \in \Lambda_1$.

Observe that τ has order n , by the (n, n) -periodicity condition on Λ_1 .

As in Lemma 4.1.14, it follows from Lemma 6.1.4 that

$$\tau(E_{i,\lambda}) = E_{i-1,[1]\lambda}$$

for $i \in \{1, \dots, r\}$ and $\lambda \in \Lambda_0$ with $\lambda_{i+1} > 0$,

$$\tau(F_{i,\lambda}) = F_{i-1,[1]\lambda}$$

for $i \in \{1, \dots, n\}$ and $\lambda \in \Lambda_0$ with $\lambda_i > 0$, and

$$\tau(1_\lambda) = 1_{[1]\lambda}$$

for $\lambda \in \Lambda_0$.

In particular,

$$\begin{aligned} \tau(E_i) &= E_{i-1} \\ \tau(F_i) &= F_{i-1} \end{aligned}$$

for $i \in \{1, \dots, r\}$.

As earlier, I can not be sure but I think this map τ is related to the Auslander-Reiten translation on the isomorphism classes of nilpotent representations of the cyclic quiver on n vertices. The result that $\tau(E_i) = E_{i-1}$ is consistent with the fact the A.R translation sends the simple representation at vertex i to the simple representation at vertex $i - 1$.

6.2 Multiplicative bases in affine zero Schur algebras: motivating example

Recall that the affine 0-Schur algebra $\hat{S}_0(n, r)$ is defined to be the associative \mathbb{Z} -algebra

$$\hat{S}_0(n, r) = \mathbb{Z}[q]/(q) \otimes_{\mathbb{Z}[q]} \hat{S}_q(n, r).$$

In particular, $\hat{S}_0(n, r)$ has a \mathbb{Z} -basis

$$\{e_A : A \in \Lambda_1\}$$

with \mathbb{Z} -bilinear product given by

$$e_A e_B = \sum_{C \in \Lambda_1} \gamma_{A,B,C}(0) e_C$$

for $A, B, C \in \Lambda_1$; where $\gamma_{A,B,C} \in \mathbb{Z}[q]$ are the structure polynomials of the affine q -Schur algebra $\hat{S}_q(n, r)$ with respect to this distinguished basis.

The multiplicative identity in $\hat{S}_0(n, r)$ is

$$\sum_{\lambda \in \Lambda_0} 1_\lambda.$$

The result of the shifting lemma, Lemma 4.1.12, also holds in $\hat{S}_0(n, r)$. In particular,

$$Re_A = e_{[1]A}$$

and

$$e_A R = e_{A[-1]},$$

for each $A \in \Lambda_1$.

Now assume $r = 1$, so

$$\Lambda_1(n, 1) = \{\mathcal{E}_{i,j} : (i, j) \in \mathbb{Z} \times \{1, \dots, n\}\}$$

and

$$\Lambda_0(n, 1) = \{\varepsilon_n, \dots, \varepsilon_1\}.$$

Lemma 6.2.1. *The distinguished basis $\{e_A : A \in \Lambda_1(n, 1)\}$ is a multiplicative basis of $\hat{S}_0(n, 1)$. More precisely,*

$$e_{\mathcal{E}_{i,j}} e_{\mathcal{E}_{j,k}} = e_{\mathcal{E}_{i,k}}$$

for $i, j, k \in \mathbb{Z}$, and

$$e_{\mathcal{E}_{i,j}} e_{\mathcal{E}_{k,l}} = 0$$

for $i, j, k, l \in \mathbb{Z}$ with $j \neq k$ modulo n .

Proof. Let $i, j \in \mathbb{Z}$. Lemma 4.1.12 shows that

$$e_{\mathcal{E}_{i,j}} = R^{j-i} 1_{\varepsilon_j},$$

where the subscript of ε_j is taken modulo n .

If $i, j, k, l \in \mathbb{Z}$ with $j \neq k$ modulo n , then

$$\text{co}(\mathcal{E}_{i,j}) = \varepsilon_j \neq \varepsilon_k = \text{ro}(\mathcal{E}_{k,l}),$$

so

$$e_{\mathcal{E}_{i,j}} e_{\mathcal{E}_{k,l}} = 0.$$

Finally, let $i, j, k \in \mathbb{Z}$. Then

$$\begin{aligned} e_{\mathcal{E}_{i,j}} e_{\mathcal{E}_{j,k}} &= R^{j-i} 1_{\varepsilon_j} R^{k-j} 1_{\varepsilon_k} \\ &= R^{j-i} R^{k-j} 1_{\varepsilon_k} \\ &= R^{k-i} 1_{\varepsilon_k} \\ &= e_{\mathcal{E}_{i,k}}. \end{aligned}$$

This proves that the basis $\{e_A : A \in \Lambda_1(n, 1)\}$ of $\hat{S}_0(n, 1)$ is a multiplicative basis. \square

This result also shows that the product in $\hat{S}_0(n, 1)$ is the same as the generic product, since

$$e_A e_B = e_{A*B}$$

if $\text{co}(A) = \text{ro}(B)$, and

$$e_A e_B = 0$$

if $\text{co}(A) \neq \text{ro}(B)$, for $A, B \in \Lambda_1(n, 1)$.

Corollary 6.2.2. *For each integer $n \geq 1$,*

$$\hat{S}_0(n, 1) = \hat{G}(n, 1).$$

Proof. This is a consequence of Lemma 6.2.1 and the comment which follows the proof. \square

6.3 Aperiodicity in the generic affine algebra

Definition 6.3.1. An element $A \in \Lambda_1$ is aperiodic if for each $l \in \mathbb{Z} \setminus \{0\}$ there exists $i \in \mathbb{Z}$ such that $a_{i,i+l} = 0$.

An element of $\hat{G}(n, r)$ is said to be aperiodic if it is a \mathbb{Z} -linear combination of basis elements e_A corresponding to the aperiodic elements in Λ_1 .

For example, the diagonal matrix D_λ is aperiodic so 1_λ is aperiodic, for any $\lambda \in \Lambda_0$. The elementary basis elements $E_{i,\lambda}$ and $F_{i,\lambda}$ introduced earlier are also aperiodic.

When $r < n$, any element $A \in \Lambda_1$ is aperiodic since $\text{co}(A)$ is insincere and therefore A has a zero column.

Lemma 6.3.2. *Suppose $A \in \Lambda_1$ is aperiodic and write $\mu = \text{ro}(A)$. If $\mu_{i+1} > 0$, then $E_i * e_A$ is aperiodic. If $\mu_i > 0$, then $F_i * e_A$ is aperiodic.*

Proof. Let $A \in \Lambda_1$ be aperiodic and let $\mu = \text{ro}(A)$.

Suppose $\mu_{i+1} > 0$. There is $p \in \mathbb{Z}$ such that $a_{i+1,p} > 0$ and $a_{i+1,p'} = 0$ whenever $p' > p$. Lemma 4.1.2 shows that $E_i * e_A = e_B$, where $B = A + \mathcal{E}_{i,p} - \mathcal{E}_{i+1,p}$. Let $l \in \mathbb{Z} \setminus \{0\}$. If $l \notin \{p-i, p-i-1\}$, then $b_{s,s+l} = a_{s,s+l}$ for each $s \in \mathbb{Z}$, so there is $s \in \mathbb{Z}$ such that $b_{s,s+l} = a_{s,s+l} = 0$, since A is aperiodic. If $l = p-i$, then $b_{i+1,i+1+l} = b_{i+1,p+1} = a_{i+1,p+1} = 0$, by maximality of p . If

$l = p - i - 1$, there is $s \neq i + 1$ such that $a_{s,s+l} = 0$, since A is aperiodic and $a_{i+1,i+1+l} = a_{i+1,p} > 0$, so $b_{s,s+l} = a_{s,s+l} = 0$. Therefore, $B = A + \mathcal{E}_{i,p} - \mathcal{E}_{i+1,p}$ is aperiodic.

Suppose $\mu_i > 0$. Lemma 4.1.2 shows that $F_i * e_A = e_C$ where $C = A + \mathcal{E}_{i+1,p} - \mathcal{E}_{i,p}$ and $p = \min\{p' \in \mathbb{Z} : a_{i,p'} > 0\}$. Let $l \in \mathbb{Z} \setminus \{0\}$. If $l \notin \{p - i, p - i - 1\}$ then $c_{s,s+l} = a_{s,s+l}$ for each $s \in \mathbb{Z}$, so there is $s \in \mathbb{Z}$ such that $c_{s,s+p} = a_{s,s+p} = 0$, by aperiodicity of A . If $l = p - i$, then $a_{i,i+l} = a_{i,p} > 0$, so there is $s \neq i$ such that $a_{s,s+l} = 0$. Then $c_{s,s+l} = a_{s,s+l} = 0$. Finally, if $l = p - i - 1$, then $c_{i,i+l} = a_{i,p-1} = 0$ by minimality of p . Thus C is aperiodic as required. \square

Suppose $\lambda \in \Lambda_0$ and

$$\omega = \omega_1 \cdots \omega_m,$$

where

$$\omega_1, \dots, \omega_m \in \{E_1, \dots, E_n\} \cup \{F_1, \dots, F_n\}.$$

Either $\omega * 1_\lambda = 0$ or $\omega * 1_\lambda = e_A$ for some $A \in \Lambda_1$, where A is aperiodic, as a result of Lemma 6.3.2.

The next step is to prove a converse of this result. It will be shown that each of the aperiodic basis elements e_A in $\hat{G}(n, r)$ can be expressed in the form $\omega 1_\lambda$, where ω is a word in E_1, \dots, E_n and F_1, \dots, F_n and $\lambda = \text{co}(A)$. This will be proven by induction on the ‘weight’ of a matrix by showing how any aperiodic basis element can be written as the product of some E_i or F_i with an aperiodic basis element of strictly smaller weight.

Definition 6.3.3. For each $A \in \Lambda_1$, define the weight of A to be the non negative integer $\text{wt}(A)$ given by

$$\text{wt}(A) = \sum_{i \in \{1, \dots, n\}, j \in \mathbb{Z}} |j - i| a_{i,j}.$$

Observe that

$$\text{wt}(A) = \sum_{[i,j]: i < j} (j - i) a_{i,j} + \sum_{[i,j]: i > j} (i - j) a_{i,j}.$$

Also write $\text{wt}(e_A) = \text{wt}(A)$. Then 1_λ has weight 0, and $E_{i,\lambda}$ and $F_{i,\lambda}$ have weight 1. In fact, the converses also hold in that $\text{wt}(e_A) = 0$ implies $e_A = 1_\lambda$ for some λ , and $\text{wt}(e_A) = 1$ implies e_A is $E_{i,\lambda}$ or $F_{i,\lambda}$ for some i and λ .

Lemma 6.3.4. Let $A \in \Lambda_1$ and write $\mu = \text{ro}(A)$. Suppose $\mu_{i+1} > 0$ and set

$$p = \max\{p' \in \mathbb{Z} : a_{i+1,p'} > 0\}.$$

If $p > i$ then

$$\text{wt}(E_i * e_A) = 1 + \text{wt}(e_A)$$

and if $p \leq i$ then

$$\text{wt}(E_i * e_A) = -1 + \text{wt}(e_A).$$

Proof. Lemma 6.1.1 shows that

$$E_i * e_A = e_{A + \mathcal{E}_{i,p} - \mathcal{E}_{i+1,p}}$$

so

$$\text{wt}(E_i * e_A) - \text{wt}(e_A) = |p - i| - |p - i - 1|,$$

which equals 1 if $p > i$ and equals -1 if $p \leq i$. \square

Lemma 6.3.5. Let $A \in \Lambda_1$ and $\mu = \text{ro}(A)$. Suppose $i \in \{1, \dots, n\}$ is such that $\mu_i > 0$ and let

$$q = \min\{q' \in \mathbb{Z} : a_{i,q'} > 0\}.$$

If $q \leq i$ then

$$\text{wt}(F_i * e_A) = \text{wt}(e_A) + 1$$

and if $q > i$ then

$$\text{wt}(F_i * e_A) = \text{wt}(e_A) - 1.$$

Proof. Again using Lemma 6.1.1,

$$F_i * e_A = e_{A + \mathcal{E}_{i+1,q} - \mathcal{E}_{i,q}},$$

so

$$\text{wt}(F_i * e_A) - \text{wt}(e_A) = |q - i - 1| - |q - i|,$$

which equals -1 if $q > i$ and equals 1 if $q \leq i$. □

Lemma 6.3.6. If $A \in \Lambda_1$ is aperiodic, then

$$e_A = \omega_1 \cdots \omega_m 1_\lambda$$

for some

$$\omega_1, \dots, \omega_m \in \{E_1, \dots, E_n\} \cup \{F_1, \dots, F_n\},$$

where $\lambda = \text{co}(A)$.

Proof. The proof uses induction on the weight of A .

If $\text{wt}(A) = 0$ then $A = D_\lambda$, where $\lambda = \text{co}(A)$, so

$$e_A = 1_\lambda.$$

Assume $\text{wt}(A) > 0$. Then A has at least one nonzero entry which is not on the diagonal.

Suppose the upper part of A is nonzero and set

$$h^+ = \max\{j - i : a_{i,j} \neq 0\}.$$

There is $i \in \{1, \dots, n\}$ such that $a_{i,i+h^+} > 0$ and $a_{i+1,i+1+h^+} = 0$, using the aperiodicity property of A . Let p be the smallest integer such that $p > i$, $a_{i,p} > 0$ and $a_{i+1,j} = 0$ for $j > p$.

Then

$$e_A = E_i * e_B$$

where $B = A + \mathcal{E}_{i+1,p} - \mathcal{E}_{i,p}$. Moreover, B is aperiodic and

$$\text{wt}(B) = \text{wt}(A) - 1,$$

using Lemma 6.3.4.

Next suppose the lower part of A is nonzero and set

$$h^- = \max\{i - j : a_{i,j} > 0\}.$$

There is $i \in \{1, \dots, n\}$ such that $a_{i,i-h^-} = 0$ and $a_{i+1,i+1-h^-} > 0$, by the aperiodicity property of A . Let q be the largest integer such that $q < i + 1$, $a_{i+1,q} > 0$ and $a_{i,j} = 0$ for $j < q$. Then $q \geq i - h^-$ and

$$e_A = F_i e_B,$$

where

$$B = A + \mathcal{E}_{i,q} - \mathcal{E}_{i+1,q}.$$

Observe B is aperiodic and

$$\text{wt}(B) = \text{wt}(A) - 1,$$

by Lemma 6.3.5.

Therefore, if $\text{wt}(A) > 0$ there exists an aperiodic element $B \in \Lambda_1$ with

$$\text{wt}(B) = \text{wt}(A) - 1$$

and such that

$$e_A = \omega e_B$$

for some $\omega \in \{E_1, \dots, E_n\} \cup \{F_1, \dots, F_n\}$.

It follows that any aperiodic basis element e_A is the product of a word of length $\text{wt}(A)$ in E_1, \dots, E_n and F_1, \dots, F_n with the idempotent 1_λ , where $\lambda = \text{co}(A)$. \square

Proposition 6.3.7. *The subalgebra of $\hat{G}(n, r)$ generated by E_i and F_i for $i \in \{1, \dots, n\}$ and 1_λ for $\lambda \in \Lambda_0$ has \mathbb{Z} -basis*

$$\{e_A : A \in \Lambda_1 \text{ is aperiodic}\}.$$

Proof. By definition, this subalgebra is spanned by the nonzero products in E_i and F_i for $i \in \{1, \dots, n\}$ and 1_λ for $\lambda \in \Lambda_0$, which are exactly the aperiodic basis elements, by Lemma 6.3.2 and Lemma 6.3.6. \square

Lemma 6.3.8. *In the case $r < n$, $\hat{G}(n, r)$ is generated by E_i and F_i for $i \in \{1, \dots, n\}$ and 1_λ for $\lambda \in \Lambda_0$.*

Proof. When $r < n$, any $A \in \Lambda_1$ is aperiodic since $\text{co}(A)$ has a zero entry, so A has a column of zero entries. Therefore each of the basis elements e_A in $\hat{G}(n, r)$ may be written as a product of the E_i , F_i and 1_λ , using Proposition 6.3.7. \square

6.4 Quiver presentation of the generic affine algebra.

Let n and r be integers with $n \geq 3$ and $r \geq 1$. Let $\Gamma = \Gamma(n, r)$ be the quiver associated to the affine q -Schur algebra $\hat{S}_q(n, r)$, as defined in Section 4.2.2.

Recall that Γ is the quiver with set of vertices $\Gamma_0 = \Lambda_0$ and set of arrows $\Gamma_1 = \Gamma_1^+ \cup \Gamma_1^-$, where Γ_1^+ consists of the arrows

$$e_{i,\lambda}: \lambda \rightarrow \lambda + \alpha_i \text{ for } (i, \lambda) \in \{1, \dots, n\} \times \Lambda_0 \text{ with } \lambda_{i+1} > 0,$$

and Γ_1^- consists of the arrows

$$f_{i,\lambda}: \lambda \rightarrow \lambda - \alpha_i \text{ for } (i, \lambda) \in \{1, \dots, n\} \times \Lambda_0 \text{ with } \lambda_i > 0.$$

Recall that the path \mathbb{Z} -algebra of Γ is an associative \mathbb{Z} -algebra with a \mathbb{Z} -basis consisting of the paths in Γ and with multiplication defined by concatenation of paths. If p and q are paths in Γ then the product pq is the path q followed by p if the target of q equals the source of p , otherwise pq equals zero.

For each $i \in \{1, \dots, n\}$, define

$$e_i = \sum_{\lambda \in \Lambda_0: \lambda_{i+1} > 0} e_{i,\lambda}$$

and

$$f_i = \sum_{\lambda \in \Lambda_0: \lambda_i > 0} f_{i,\lambda}.$$

Let \mathcal{J} be the ideal in $\mathbb{Z}\Gamma$ generated by the following expressions, which are obtained from the relations in the q -Schur algebra by setting q equal to 0:

$$\begin{aligned} e_i e_j - e_j e_i, \\ f_i f_j - f_j f_i \end{aligned}$$

for $i, j \in \{1, \dots, n\}$ with $j > i + 1$;

$$\begin{aligned} e_i e_{i+1}^2 - e_{i+1} e_i e_{i+1}, \\ e_i^2 e_{i+1} - e_i e_{i+1} e_i, \\ f_{i+1}^2 f_i - f_{i+1} f_i f_{i+1}, \\ f_{i+1} f_i^2 - f_i f_{i+1} f_i \end{aligned}$$

for $i \in \{1, \dots, n\}$;

$$e_i f_j - f_j e_i$$

for $i, j \in \{1, \dots, n\}$ with $i < j$;

$$e_i f_i - f_i e_i - \sum_{\lambda \in \Lambda_0} c_{i,\lambda} k_\lambda$$

for $i \in \{1, \dots, n\}$, where

$$c_{i,\lambda} = \begin{cases} 1 & : \text{if } \lambda_{i+1} = 0, \lambda_i > 0 \\ 0 & : \text{if } \lambda_i > 0, \lambda_{i+1} > 0 \\ -1 & : \text{if } \lambda_i = 0, \lambda_{i+1} > 0. \end{cases}$$

Multiplying each expression above with the idempotents k_λ for $\lambda \in \Lambda_0$ gives a relation involving paths with common source and target vertices, thus \mathcal{J} is an ideal of \mathbb{Z} -linear relations in Γ .

The ideal \mathcal{J} in $\mathbb{Z}\Gamma$ is generated by the following set of relations:

$$\begin{aligned} e_{i,\lambda+\alpha_j} e_{j,\lambda} - e_{j,\lambda+\alpha_i} e_{i,\lambda}, \\ f_{i,\lambda-\alpha_j} f_{j,\lambda} - f_{j,\lambda-\alpha_i} f_{i,\lambda}, \end{aligned}$$

for $i, j \in \{1, \dots, n\}$ with $j > i + 1$;

$$\begin{aligned} e_{i,\lambda+2\alpha_{i+1}} e_{i+1,\lambda+\alpha_{i+1}} e_{i+1,\lambda} - e_{i+1,\lambda+\alpha_i+\alpha_{i+1}} e_{i,\lambda+\alpha_{i+1}} e_{i+1,\lambda}, \\ e_{i,\lambda+\alpha_i+\alpha_{i+1}} e_{i,\lambda+\alpha_{i+1}} e_{i+1,\lambda} - e_{i,\lambda+\alpha_i+\alpha_{i+1}} e_{i+1,\lambda+\alpha_i} e_{i,\lambda}, \\ f_{i+1,\lambda-\alpha_i-\alpha_{i+1}} f_{i+1,\lambda-\alpha_i} f_{i,\lambda} - f_{i+1,\lambda-\alpha_i-\alpha_{i+1}} f_{i,\lambda-\alpha_{i+1}} f_{i+1,\lambda}, \\ f_{i+1,\lambda-2\alpha_i} f_{i,\lambda-\alpha_i} f_{i,\lambda} - f_{i,\lambda-\alpha_i-\alpha_{i+1}} f_{i+1,\lambda-\alpha_i} f_{i,\lambda}, \end{aligned}$$

for $i \in \{1, \dots, n\}$;

$$e_{i,\lambda-\alpha_j} f_{j,\lambda} - f_{j,\lambda+\alpha_i} e_{i,\lambda}$$

for $i, j \in \{1, \dots, n\}$ with $i \neq j$;

$$e_{i,\lambda-\alpha_i} f_{i,\lambda} - f_{i,\lambda+\alpha_i} e_{i,\lambda}$$

for $i \in \{1, \dots, n\}$ and $\lambda \in \Lambda_0$ with $\lambda_i > 0$ and $\lambda_{i+1} > 0$;

$$e_{i,\lambda-\alpha_i} f_{i,\lambda} - k_\lambda$$

for $i \in \{1, \dots, n\}$ and $\lambda \in \Lambda_0$ such that $\lambda_i > 0$ and $\lambda_{i+1} = 0$;

$$f_{i, \lambda + \alpha_i} e_{i, \lambda} - k_\lambda$$

for $i \in \{1, \dots, n\}$ and $\lambda \in \Lambda_0$ with $\lambda_i = 0$ and $\lambda_{i+1} > 0$.

Lemma 6.4.1. *The following equations hold in the generic affine algebra $\hat{G}(n, r)$:*

$$E_i E_j = E_j E_i$$

$$F_i F_j = F_j F_i$$

for $i, j \in \{1, \dots, n\}$ with $|j - i| \neq 1$;

$$E_i E_{i+1}^2 = E_{i+1} E_i E_{i+1}$$

$$E_i^2 E_{i+1} = E_i E_{i+1} E_i$$

$$F_{i+1}^2 F_i = F_{i+1} F_i F_{i+1}$$

$$F_{i+1} F_i^2 = F_i F_{i+1} F_i$$

for $i \in \{1, \dots, n\}$;

$$E_i F_j = F_j E_i$$

for $i, j \in \{1, \dots, n\}$ with $i \neq j$;

$$E_i F_i - F_i E_i = \sum_{\lambda \in \Lambda_0} c_{i, \lambda} 1_\lambda$$

for $i \in \{1, \dots, n\}$.

Proof. Suppose $i, j \in \{1, \dots, n\}$ with $j > i + 1$, so $\{i, i + 1\}$ and $\{j, j + 1\}$ are disjoint, then

$$\begin{aligned} E_i E_j &= \sum_{\lambda \in \Lambda_0} E_i [D_\lambda + \mathcal{E}_{j, j+1} - \mathcal{E}_{j+1, j+1}] \\ &= \sum_{\lambda \in \Lambda_0} [D_\lambda + \mathcal{E}_{i, i+1} - \mathcal{E}_{i+1, i+1} + \mathcal{E}_{j, j+1} - \mathcal{E}_{j+1, j+1}] \\ &= E_j E_i \end{aligned}$$

Then applying the transpose involution yields the second equation:

$$F_i F_j - F_j F_i = -S([E_i, E_j]) = 0.$$

Using the fundamental multiplication rules 6.1.1 and 6.1.3, for each $i \in \{1, \dots, n\}$,

$$\begin{aligned} E_i E_{i+1}^2 &= \sum_{\lambda \in \Lambda_0} E_i [D_\lambda + 2\mathcal{E}_{i+1, i+2} - 2\mathcal{E}_{i+2, i+2}] \\ &= \sum_{\lambda \in \Lambda_0} [D_\lambda + 2\mathcal{E}_{i+1, i+2} - 2\mathcal{E}_{i+2, i+2} + \mathcal{E}_{i, i+2} - \mathcal{E}_{i+1, i+2}] \\ &= \sum_{\lambda \in \Lambda_0} [D_\lambda + \mathcal{E}_{i, i+2} + \mathcal{E}_{i+1, i+2} - 2\mathcal{E}_{i+2, i+2}] \end{aligned}$$

and

$$\begin{aligned} E_{i+1}E_iE_{i+1} &= \sum_{\lambda \in \Lambda_0} E_{i+1} [D_\lambda + \mathcal{E}_{i,i+2} - \mathcal{E}_{i+2,i+2}] \\ &= \sum_{\lambda \in \Lambda_0} [D_\lambda + \mathcal{E}_{i,i+2} + \mathcal{E}_{i+1,i+2} - 2\mathcal{E}_{i+2,i+2}], \end{aligned}$$

so $E_iE_{i+1}^2 = E_{i+1}E_iE_{i+1}$.

$$\begin{aligned} E_i^2E_{i+1} &= \sum_{\mu \in \Lambda_0} [D_\mu + 2\mathcal{E}_{i,i+1} - 2\mathcal{E}_{i,i}] E_{i+1} \\ &= \sum_{\mu \in \Lambda_0} [D_\mu + \mathcal{E}_{i,i+2} + \mathcal{E}_{i,i+1} - 2\mathcal{E}_{i,i}] \end{aligned}$$

and

$$\begin{aligned} E_iE_{i+1}E_i &= \sum_{\mu \in \Lambda_0} [D_\mu + \mathcal{E}_{i,i+2} - \mathcal{E}_{i,i}] E_i \\ &= \sum_{\mu \in \Lambda_0} [D_\mu + \mathcal{E}_{i,i+2} + \mathcal{E}_{i,i+1} - 2\mathcal{E}_{i,i}], \end{aligned}$$

so $E_i^2E_{i+1} = E_iE_{i+1}E_i$.

The relations between F_i and F_{i+1} may be deduced using the transpose involution as follows:

$$F_{i+1}^2F_i = S(E_iE_{i+1}^2) = S(E_{i+1}E_iE_{i+1}) = F_{i+1}F_iF_{i+1}$$

and

$$F_{i+1}F_i^2 = S(E_i^2E_{i+1}) = S(E_iE_{i+1}E_i) = F_iF_{i+1}F_i.$$

Suppose $i, j \in \{1, \dots, n\}$ with $i \neq j$. Then

$$\begin{aligned} E_iF_j &= \sum_{\lambda \in \Lambda_0} E_i [D_\lambda + \mathcal{E}_{j+1,j} - \mathcal{E}_{j,j}] \\ &= \sum_{\lambda \in \Lambda_0} [D_\lambda + \mathcal{E}_{j+1,j} - \mathcal{E}_{j,j} + \mathcal{E}_{i,i+1} - \mathcal{E}_{i+1,i+1}] \end{aligned}$$

and

$$\begin{aligned} F_jE_i &= \sum_{\lambda \in \Lambda_0} F_j [D_\lambda + \mathcal{E}_{i,i+1} - \mathcal{E}_{i+1,i+1}] \\ &= \sum_{\lambda \in \Lambda_0} [D_\lambda + \mathcal{E}_{i,i+1} - \mathcal{E}_{i+1,i+1} + \mathcal{E}_{j+1,j} - \mathcal{E}_{j,j}], \end{aligned}$$

so $E_iF_j = F_jE_i$.

Finally, for $i \in \{1, \dots, n\}$,

$$E_iF_i = \sum_{\lambda: \lambda_i > 0, \lambda_{i+1} = 0} 1_\lambda + \sum_{\lambda: \lambda_i > 0, \lambda_{i+1} > 0} [D_\lambda + \mathcal{E}_{i,i+1} - \mathcal{E}_{i+1,i+1} + \mathcal{E}_{i+1,i} - \mathcal{E}_{i,i}]$$

and

$$F_i E_i = \sum_{\lambda: \lambda_i=0, \lambda_{i+1}>0} 1_\lambda + \sum_{\lambda: \lambda_i>0, \lambda_{i+1}>0} [D_\lambda + \mathcal{E}_{i,i+1} - \mathcal{E}_{i+1,i+1} + \mathcal{E}_{i+1,i} - \mathcal{E}_{i,i}],$$

so

$$\begin{aligned} E_i F_i - F_i E_i &= \sum_{\lambda: \lambda_i>0, \lambda_{i+1}=0} 1_\lambda - \sum_{\lambda: \lambda_i=0, \lambda_{i+1}>0} 1_\lambda \\ &= \sum_{\lambda \in \Lambda_0} c_{i,\lambda} 1_\lambda. \end{aligned}$$

□

Lemma 6.4.1 shows that there is a homomorphism of \mathbb{Z} -algebras

$$\rho: \mathbb{Z}\Gamma/\mathcal{J} \rightarrow \hat{G}(n, r)$$

defined by

$$\begin{aligned} \rho(k_\lambda + \mathcal{J}) &= 1_\lambda \\ \rho(e_{i,\lambda} + \mathcal{J}) &= E_{i,\lambda} \\ \rho(f_{i,\lambda} + \mathcal{J}) &= F_{i,\lambda}, \end{aligned}$$

for all $i \in \{1, \dots, n\}$ and $\lambda \in \Lambda_0$.

Proposition 6.4.2. *The image of ρ is spanned by the aperiodic basis elements. If $r < n$ then ρ is surjective.*

Proof. The image of ρ is the subalgebra of $\hat{G}(n, r)$ generated by E_i and F_i for $i \in \{1, \dots, n\}$ and 1_λ for $\lambda \in \Lambda_0$, which has \mathbb{Z} -basis

$$\{e_A : A \in \Lambda_1, A \text{ is aperiodic.}\},$$

using Proposition 6.3.7. If $r < n$ then every $A \in \Lambda_1$ is aperiodic, since A must contain a zero row or column. Therefore ρ is surjective when $r < n$. □

The following lemma gives a few relations which may be derived from the relations given in Lemma 6.4.1. This lemma will be used later to show that certain paths are congruent modulo \mathcal{J} . Observe that the relations given below also correspond to relations in the bound quiver algebra, since they are derived only from the braid type relations in Lemma 6.4.1 and the corresponding relations in \mathcal{J} , respectively.

Lemma 6.4.3. *For each $a \geq 1$ and $i \in \{1, \dots, n\}$ the following relations hold:*

$$E_i^2 E_{i+1}^a - E_i E_{i+1}^a E_i = 0 \tag{6.4.1}$$

$$E_{i+1} E_i^a E_{i+1} - E_i^a E_{i+1}^2 = 0 \tag{6.4.2}$$

$$F_{i+1}^a F_i^2 - F_i F_{i+1}^a F_i = 0 \tag{6.4.3}$$

$$F_{i+1}^2 F_i^a - F_{i+1} F_i^a F_{i+1} = 0. \tag{6.4.4}$$

Proof. Proceed by induction on a to first establish the relations including E_i and E_{i+1} . If $a = 1$ the relations follows from Lemma 6.4.1. Now suppose $a \geq 2$ and the statement of the lemma holds for all smaller values. Then

$$\begin{aligned}
E_i E_{i+1}^a E_i &= E_{i+1} E_i E_{i+1}^{a-1} E_i \\
&= E_{i+1} E_i^2 E_{i+1}^{a-1} \text{ (inductive hypothesis.)} \\
&= E_{i+1} E_i E_{i+1} E_i E_{i+1}^{a-2} \\
&= E_i E_{i+1}^2 E_i E_{i+1}^{a-2} \\
&= E_i E_{i+1} E_i E_{i+1}^{a-1} \\
&= E_i^2 E_{i+1}^a.
\end{aligned}$$

Similarly,

$$\begin{aligned}
E_{i+1} E_i^a E_{i+1} &= E_{i+1} E_i^{a-1} E_{i+1} E_i \\
&= E_i^{a-1} E_{i+1}^2 E_i \text{ (inductive hypothesis.)} \\
&= E_i^{a-2} E_{i+1} E_i E_{i+1} E_i \\
&= E_i^{a-2} E_{i+1} E_i^2 E_{i+1} \\
&= E_i^{a-1} E_{i+1} E_i E_{i+1} \\
&= E_i^a E_{i+1}^2.
\end{aligned}$$

Thus the first two relations hold for each $a \in \mathbb{N}$. Finally, applying the transpose involution to these relations yields the relations involving F_i and F_{i+1} :

$$F_{i+1}^a F_i^2 - F_i F_{i+1}^a F_i = \top(E_i E_{i+1}^a E_i - E_i^2 E_{i+1}^a) = 0$$

and

$$F_{i+1}^2 F_i^a - F_{i+1} F_i^a F_{i+1} = \top(E_{i+1} E_i^a E_{i+1} - E_i^a E_{i+1}^2) = 0.$$

□

6.4.1 Standard paths

Recall the definition of standard paths in Γ , from Definition 4.2.4. As noted earlier, there is a bijection between the set of standard paths in Γ and the standard monomial basis in $\hat{G}(n, r)$ indexed by Λ_1 .

Lemma 6.4.4. *Let p be a standard path in Γ . If q is a path in Γ with $q = pe_i$ or $q = pf_i$ for some $i \in \{1, \dots, n\}$, then q is congruent to a standard path modulo \mathcal{J} .*

Proof. **IMPORTANT: need to prove this beyond doubt.**

□

Proposition 6.4.5. *When $r < n$, any path in Γ is congruent to a standard path modulo \mathcal{J} .*

Proof. Let p be a path in Γ and proceed by induction on the length of p . If p has length zero then $p = k_\mu$ for some $\mu \in \Lambda_0$, so p is a standard path. If p has length one then $p = k_\mu e_i$ or $p = k_\mu f_i$ for some $\mu \in \Lambda_0$ and $i \in \{1, \dots, n\}$.

Suppose p has length at least two and that any strictly shorter path is congruent to a standard path. Pulling out the first arrow, write $p = p'e_i$ or $p = p'f_i$ for some $i \in \{1, \dots, n\}$. Using the

inductive hypothesis we may assume p' is a standard path, so it follows from Lemma 6.4.4 that p is congruent to a standard path.

Note: The lemma on extending standard paths still needs to be proven for this proof to be complete. \square

Theorem 6.4.6. *If $r < n$ then ρ is a \mathbb{Z} -algebra isomorphism. Thus $\hat{G}(n, r)$ admits a presentation by the quiver Γ and the ideal of relations \mathcal{J} in $\mathbb{Z}\Gamma$.*

Proof. Under the assumption $r < n$, ρ is a surjective homomorphism of \mathbb{Z} -algebras, by Proposition 6.4.2.

Suppose p and p' are paths in Γ and $A \in \Lambda_1$ with

$$\rho(p + \mathcal{J}) = \rho(p' + \mathcal{J}) = e_A.$$

Proposition 6.4.5 shows that p and p' are both congruent modulo \mathcal{J} to the standard path corresponding to A , so p and p' are congruent modulo \mathcal{J} . Therefore ρ is injective, so is an isomorphism of \mathbb{Z} -algebras as claimed. \square

Conjecture 6.4.7. Even when $r \geq n$, ρ gives a \mathbb{Z} -algebra isomorphism

$$\mathbb{Z}\Gamma/\mathcal{J} \rightarrow \hat{G}(n, r)^{ap},$$

where $\hat{G}(n, r)^{ap}$ is the subalgebra of $\hat{G}(n, r)$ spanned by the aperiodic basis elements.

Rough. There will have to be an extended or new notion of standard path for the case of aperiodic basis elements which do not have a row or column of zero entries. \square

6.5 The isomorphism result

This section gives a realisation of the affine 0-Schur algebra by the generic affine algebra in the case that $r < n$. Recall that the affine 0-Schur algebra $\hat{S}_0(n, r)$ is defined to be the \mathbb{Z} -algebra

$$\hat{S}_0(n, r) = \mathbb{Z}[q]/(q) \otimes_{\mathbb{Z}} [q]\hat{S}_q(n, r).$$

The inclusion of $\mathbb{Z}[q]$ into \mathcal{Q} sending f to $f/1$ gives an isomorphism of \mathbb{Z} algebras

$$\mathbb{Z}[q]/q\mathbb{Z}[q] \rightarrow \mathcal{Q}/q\mathcal{Q} : a + q\mathbb{Z}[q] \mapsto a + q\mathcal{Q},$$

and both are isomorphic to \mathbb{Z} itself. Therefore

$$\hat{S}_0(n, r) = \mathcal{Q}/q\mathcal{Q} \otimes_{\mathcal{Q}} \hat{S}_{\mathcal{Q}}(n, r)$$

Let $n, r \geq 1$ with $r < n$. Recall

$$\phi: \mathbb{Z}[q]\Gamma/I \rightarrow \hat{S}_q(n, r)$$

is the homomorphism of $\mathbb{Z}[q]$ -algebras defined by

$$\begin{aligned} \phi(e_{i,\lambda}) &= E_{i,\lambda} \\ \phi(f_{i,\lambda}) &= F_{i,\lambda} \\ \phi(k_\lambda) &= 1_\lambda, \end{aligned}$$

for $i \in \{1, \dots, n\}$ and $\lambda \in \Lambda_0$. Let ϕ_0 be the \mathbb{Z} -algebra homomorphism

$$\phi_0 = \mathcal{Q}/(q) \otimes_{\mathcal{Q}} \phi_{\mathcal{Q}}: \mathbb{Z}\Gamma/\mathcal{J} \rightarrow \hat{S}_0(n, r).$$

Let ρ be the \mathbb{Z} -algebra isomorphism

$$\rho: \mathbb{Z}\Gamma/\mathcal{J} \rightarrow \hat{G}(n, r)$$

from Theorem 6.4.6.

Let Ψ be the \mathbb{Z} -algebra homomorphism

$$\Psi = \phi_0 \circ \rho^{-1}: \hat{G}(n, r) \rightarrow \hat{S}_0(n, r) \quad (6.5.1)$$

with

$$\begin{aligned} \Psi(E_{i,\lambda}) &= E_{i,\lambda} \\ \Psi(F_{i,\lambda}) &= F_{i,\lambda} \\ \Psi(1_\lambda) &= 1_\lambda, \end{aligned}$$

for $i \in \{1, \dots, n\}$ and $\lambda \in \Lambda_0$.

Proposition 6.5.1. *The map Ψ is surjective.*

Proof. Proposition 4.2.8 shows that

$$\phi_{\mathcal{Q}}: \mathcal{Q}\Gamma/\mathcal{Q}I \rightarrow \hat{S}_{\mathcal{Q}}(n, r)$$

is a surjective \mathcal{Q} -algebra homomorphism, so

$$\phi_0 = \mathcal{Q}/(q) \otimes_{\mathcal{Q}} \phi_{\mathcal{Q}}: \mathbb{Z}\Gamma/\mathcal{J} \rightarrow \hat{S}_0(n, r)$$

is a surjective \mathbb{Z} -algebra homomorphism, using right exactness of tensor products. It follows that Ψ is surjective since $\Psi \circ \rho = \phi_0$ and ρ is an isomorphism. \square

Lemma 6.5.2. *For each $A \in \Lambda_1$,*

$$\Psi(e_A) = e_A + \sum_{B: B < A} c_B e_B$$

for some $c_B \in \mathbb{N}$.

Proof. Given $A \in \Lambda_1$, let p be the standard path in Γ corresponding to A , so

$$\rho(p + \mathcal{J}) = e_A.$$

Then

$$\begin{aligned} \Psi(e_A) &= \phi_0(p + \mathcal{J}) \\ &= \sum_{B \in \Lambda_1: B \leq A} g_B(0) e_B, \end{aligned}$$

for some $g_B \in \mathbb{Z}[q]$, where

$$\begin{aligned} g_A(0) &= \left(\prod_{i \in \{1, \dots, n\}, s \geq 1} [[\alpha_{i,s}]]! [[\beta_{i,s}]]! \right)_{q=0} \\ &= 1, \end{aligned}$$

by Lemma 4.2.5. \square

Theorem 6.5.3. *In the case $r < n$, the map*

$$\Psi: \hat{G}(n, r) \rightarrow \hat{S}_0(n, r)$$

is an isomorphism of \mathbb{Z} -algebras.

Proof. Proposition 6.5.1 shows that Ψ is a surjective \mathbb{Z} -algebra homomorphism.

To prove that Ψ is injective, suppose x is a nonzero element of $\hat{G}(n, r)$ and write

$$x = \sum_{A \in \Lambda_1} c_A e_A$$

and let $\Xi = \{A \in \Lambda_1 : c_A \neq 0\}$. Fix a maximal element D in Ξ , which exists since Ξ is nonempty and finite. Then

$$\begin{aligned} \Psi(x) &= \sum_{A \in \Xi} c_A \Psi(e_A) \\ &= \sum_{A \in \Xi} \left(c_A e_A + \sum_{B \in \Xi: B < A} c_{A,B} e_B \right) \\ &= c_D e_D + \sum_{A \in \Xi: A \neq D} c'_A e_A \end{aligned}$$

by Lemma 6.5.2, so $\Psi(x) \neq 0$, which proves that Ψ is injective and therefore Ψ is an isomorphism of \mathbb{Z} -algebras. \square

6.6 The period 2 case

In the case $n = 2$ the quiver $\Gamma = \Gamma(2, r)$ associated to $\hat{G}(2, r)$ consists of $r + 1$ vertices (totally ordered) with two pairs of edges between adjacent vertices, (e_1, f_1) and (e_2, f_2) .

The following equations are a $q = 0$ form of the q -Serre relations in Lemma 4.3.1:

Lemma 6.6.1. *The following equations hold in $\hat{G}(2, r)$, for $i \in \mathbb{Z}/2\mathbb{Z}$:*

$$\begin{aligned} E_i E_{i+1} E_i^2 &= E_i^2 E_{i+1} E_i \\ F_i F_{i+1} F_i^2 &= F_i^2 F_{i+1} F_i. \end{aligned}$$

Proof.

$$\begin{aligned} E_1 E_2 E_1^2 &= \sum_{\mu \in \Lambda_0} [D_\mu + \mathcal{E}_{1,3} - \mathcal{E}_{1,1}] E_1^2 \\ &= \sum_{\mu \in \Lambda_0} [D_\mu + \mathcal{E}_{1,4} - \mathcal{E}_{1,1}] E_1 \\ &= \sum_{\mu \in \Lambda_0} [D_\mu + \mathcal{E}_{1,4} + \mathcal{E}_{1,2} - 2\mathcal{E}_{1,1}] \end{aligned}$$

and

$$\begin{aligned}
E_1^2 E_2 E_1 &= \sum_{\mu \in \Lambda_0} [D_\mu + 2\mathcal{E}_{1,2} - 2\mathcal{E}_{1,1}] E_2 E_1 \\
&= \sum_{\mu \in \Lambda_0} [D_\mu + \mathcal{E}_{1,3} + \mathcal{E}_{1,2} - 2\mathcal{E}_{1,1}] E_1 \\
&= \sum_{\mu \in \Lambda_0} [D_\mu + \mathcal{E}_{1,4} + \mathcal{E}_{1,2} - 2\mathcal{E}_{1,1}],
\end{aligned}$$

so $E_1 E_2 E_1^2 = E_1^2 E_2 E_1$.

Recall that conjugation by R defines an automorphism τ of $\hat{G}(n, r)$ of degree 2, with $\tau(E_1) = E_2$ and $\tau(E_2) = E_1$, so

$$E_2 E_1 E_2^2 - E_2^2 E_1 E_2 = \tau(E_1 E_2 E_1^2 - E_1^2 E_2 E_1) = 0.$$

Finally, the equations involving F_i and F_{i+1} follow by applying the transpose involution:

$$F_i F_{i+1} F_i^2 - F_i^2 F_{i+1} F_i = S(E_i^2 E_{i+1} E_i - E_i E_{i+1} E_i^2) = 0,$$

for $i \in \{1, 2\}$. □

Chapter 7

Further directions

7.1 Further results on affine zero Schur algebras

[1] Investigate link between this generic product and the generic extension of representations. Shifting to the non-negative subalgebra to do computations purely in terms of generic extensions of quiver representations.

7.2 Deformed group algebras of symmetric groups

[2] Degenerate group algebras of symmetric groups: write down a presentation of the degenerate group algebras, with generators given by the transpositions, or 2-cycles. Type up the computations done for degenerate group algebras for S_3 and S_4 . Formulate propositions for the general case: the transpositions generate the degenerate group algebra; lemma: ‘these’ relations hold; these generators and relations give a presentation of the degenerate group algebras.

Terminology: deformed group algebra.

7.3 back matter

[1] $Y_{A,B}^L$ is the image of $G_L \times G_{L'}$ under the action map $(\alpha, \beta) \mapsto \alpha\beta \cdot (L', L'')$, for any $(L', L'') \in Y_{A,B}^L$. Lemma 5.3.6 shows that $G_L/H_{N,L}$ is a connected algebraic group. Moreover, $G_{L'}/H_{2N,L}$ is an irreducible affine variety, so $G_L/H_{N,L} \times G_{L'}/H_{2N,L}$ is an irreducible affine variety. It follows that $Y_{A,B}^L$ is irreducible and constructible.

Bibliography

- [1] A. A. Beilinson, G. Lusztig, R. MacPherson, et al. “A geometric setting for the quantum deformation of GL_n .” In: *Duke Mathematical Journal* 61.2 (1990), pp. 655–677.
- [2] K. Bongartz. “On degenerations and extensions of finite dimensional modules”. In: *Advances in Mathematics* 121.2 (1996), pp. 245–287.
- [3] T. Bridgeland. “Quantum groups via Hall algebras of complexes”. In: *Annals of Mathematics* (2013), pp. 739–759.
- [4] W. Crawley-Boevey and J. Sauter. “On quiver Grassmannians and orbit closures for representation-finite algebras”. In: *Mathematische Zeitschrift* 285.1-2 (2017), pp. 367–395.
- [5] B. Deng, J. Du, and Q. Fu. *A double Hall algebra approach to affine quantum Schur–Weyl theory*. Vol. 401. Cambridge University Press, 2012.
- [6] B. Deng, J. Du, and A. Mah. “Generic extensions and composition monoids of cyclic quivers”. In: *Contem. Math* 602 (2013), pp. 99–114.
- [7] B. Deng, J. Du, and B. Parshall. *Finite dimensional algebras and quantum groups*. 150. American Mathematical Soc., 2008.
- [8] B. Deng and G. Yang. “On 0-Schur algebras”. In: *Journal of Pure and Applied Algebra* 216.6 (2012), pp. 1253–1267.
- [9] Bangming Deng and Shiquan Ruan. *Hall polynomials for tame type*. 2015. eprint: [arXiv:1512.03504](#).
- [10] R. Dipper and G. James. “q-tensor space and q-Weyl modules”. In: *Transactions of the American Mathematical Society* 327.1 (1991), pp. 251–282.
- [11] R. Dipper and G. James. “The q-Schur Algebra”. In: *Proceedings of the London Mathematical Society* 3.1 (1989), pp. 23–50.
- [12] S. Doty and A. Giaquinto. “Presenting Schur algebras”. In: *International Mathematics Research Notices* 2002.36 (2002), pp. 1907–1944.
- [13] S. R. Doty and R. M. Green. “Presenting affine q-Schur algebras”. In: *Mathematische Zeitschrift* 256.2 (2007), pp. 311–345.
- [14] R. Dou, Yong Jiang, and Jie Xiao. *Hall algebra approach to Drinfeld’s presentation of quantum loop algebras*. 2010. eprint: [arXiv:1002.1316](#).
- [15] Zhaobing Fan et al. *Affine flag varieties and quantum symmetric pairs*. 2016. [arXiv:1602.04383 \[math.RT\]](#).
- [16] V. Ginzburg and E. Vasserot. “Langlands reciprocity for affine quantum groups of type A_n ”. In: *International Mathematics Research Notices* 1993.3 (1993), pp. 67–85.

- [17] R. M. Green. “q-Schur algebras as quotients of quantized enveloping algebras”. In: *Journal of algebra* 185.3 (1996), pp. 660–687.
- [18] Joe Harris. *Algebraic geometry: a first course*. Vol. 133. Springer Science & Business Media, 2013.
- [19] Andrew Hubery. *Hall polynomials for affine quivers*. 2007. eprint: [arXiv:math/0703178](https://arxiv.org/abs/math/0703178).
- [20] Andrew Hubery. “The composition algebra of an affine quiver”. In: *arXiv preprint math/0403206* (2004).
- [21] D. A. Hudec. “The Grassmanian as a Projective Variety”. In: (2007).
- [22] J. Humphreys. *Linear Algebraic Groups*. Springer-Verlag, 1981.
- [23] B. T. Jensen and X. Su. “A geometric realisation of 0-Schur and 0-Hecke algebras”. In: *Journal of Pure and Applied Algebra* 219.2 (2015), pp. 277–307.
- [24] B. T. Jensen, X. Su, and G. Yang. “Degenerate 0-Schur algebras and Nil-Temperley-Lieb algebras”. In: *arXiv preprint arXiv:1705.06084* (2017).
- [25] Bernt Tore Jensen and Xiuping Su. *A geometric realisation of 0-Schur and 0-Hecke algebras*. 2012. eprint: [arXiv:1207.6769](https://arxiv.org/abs/1207.6769).
- [26] Bernt Tore Jensen, Xiuping Su, and Guiyu Yang. *Projective modules of 0-Schur algebras*. 2013. eprint: [arXiv:1312.5487](https://arxiv.org/abs/1312.5487).
- [27] G. Lusztig. “Introduction to quantized enveloping algebras”. In: *New developments in Lie theory and their applications*. Springer, 1992, pp. 49–65.
- [28] George Lusztig. “Aperiodicity in quantum affine gl n ”. In: *Asian Journal of Mathematics* 3.1 (1999), pp. 147–178.
- [29] Patrick J Morandi. “Algebraic Groups, Grassmannians, and Flag Varieties”. In: (1998).
- [30] M. Reineke. “Generic extensions and multiplicative bases of quantum groups at $q = 0$ ”. In: *Represent. Theory* 5 (2001), pp. 147–163.
- [31] M. Reineke. “The monoid of families of quiver representations”. In: *Proceedings of the London Mathematical Society* 84.3 (2002), pp. 663–685.
- [32] M. Reineke. “The quantic monoid and degenerate quantized enveloping algebras”. In: *arXiv preprint math/0206095* (2002).
- [33] M. Reineke. “The use of geometric and quantum group techniques for wild quivers”. In: *Representations of finite dimensional algebras and related topics in Lie theory and geometry* 40 (2004), pp. 365–390.
- [34] C. M. Ringel. “Hall algebras”. In: *Banach Center Publications* 26.1 (1990), pp. 433–447.
- [35] C. M. Ringel. *The Hall algebra approach to quantum groups*. Sonderforschungsbereich 343, 1993.
- [36] X. Su. “A generic multiplication in quantized Schur algebras”. In: *Quarterly journal of mathematics* 61.4 (2010), pp. 497–510.