



Netherlands Food and Consumer
Product Safety Authority
*Ministry of Agriculture,
Nature and Food Quality*



HPAI in the Netherlands



Timeline HPAI outbreaks

20 Feb Uithuizermeeden

NL-HPAI(P)-2026-00010

Laying hen farm with approx. 39.900 animals kept in one stable. Started with an increase of the number of dead animals. Clinical signs consisted of blue combs and petechiae in belly and heart fat. There is 1 farm in the 1 km zone and 1 in the 3 km zone and 16 in the 10 km zone.

23 Feb Ede

NL-HPAI(P)-2026-00011

HPAI found on a free-range egg-laying poultry farm with two stables with approx. 23.500 chickens. An increase in the number of dead animals was the first sign of the disease. Corresponding clinical symptoms were observed.

There are 2 other poultry farms in the 1 km zone. One of which is located just 35 meters from the infected farm and belongs to the same owner. In the 3 km zone are 10 other poultry farms and in the 10 km 142.

In addition to the usual measures, the adjacent farm of the same owner was culled preemptively

2 March Lunteren

NL-HPAI(P)-2026-00012

Laying hen farm with around 30.500 animals distributed across 2 stables. In increase in dead animals and sick animals with severe clinical signs were found in one stable. Also, a reduction in food and water intake was seen. There are 4 poultry farms in the 1 km zone and 44 in the 3 km zone and 205 in the 10 km zone.

All outbreaks, the measures according 2016/429 and 2020/68 have been taken





Timeline HPAI outbreaks

4 March Neede

NL-HPAI(P)-2026-00013

Rearing farm for broiler chicks with approx. 95.000 chickens spread across 5 stables. In one of the stables with 9.000 animals an increased mortality rate was observed.

There are no poultry farms in the 1 km and 44 in the 3 km zone and 25 in the 10 km zone.

7 March Putten

NL-HPAI(P)-2026-00014

A laying hen farm with approx. 22.500 hens divide across 5 stables was infected with HPAI. An increase in mortality was found. In one of the stable the hens were lethargic and had loose stools. In another stable, the onset of the same symptoms was also observed. There are no poultry farms in the 1 km zone. In the 3 km zone are two poultry related companies and in the 10 km zone there are 79 poultry related companies, like farms and egg packing stations.

In all outbreaks, the measures according 2016/429 and 2020/68 have been taken.





Timeline HPAI outbreaks

20 March Oudemolen

NL-HPAI(P)-2026-00015

HPAI found in a broiler farm with approx. 13.500 broiler chickens in one stable. An increased mortality rate was observed. There are no poultry farms in the 1 km and 2 in the 3 km zone and 10 in the 10 km zone.

21 March Geesburg

NL-HPAI(P)-2026-00016


A laying hen farm with approx. 24.000 hens in one stable was infected with HPAI. An increase in mortality was found in one compartment of that stable. There are no poultry farms in the 1 km zone and a bird society in the 3 km zone. In the 10 km zone there are 30 79 poultry related companies, like farms and one egg packing station.

In all outbreaks, the measures according 2016/429 and 2020/68 have been taken.





HPAI (H5N1) in the wild





Thank you for your
attention!

Questions?