Upper GI Endoscopy

Date: 01 Jan 2026

Patient	FICTIONAL PATIENT	**Identifier**	DEMO-001

Indication: Persistent dyspepsia (fictional demonstration).

Premedication / Sedation: Topical pharyngeal anaesthesia; no intravenous sedation in this fictional example.

Findings

Oesophagus

Upper oesophagus: Mucosa appears unremarkable.

Middle oesophagus: No visible mucosal lesion.

Lower oesophagus: No visible erosion or luminal narrowing.

Gastro-oesophageal junction: Z-line at 40 cm from the incisors; no visible hiatus hernia.

Stomach

Cardia: Unremarkable on retroflexion.

Fundus: No visible mucosal lesion.

Body: Mucosa appears unremarkable.

Antrum: Mild patchy erythema; no visible ulcer.

Pylorus: Patent; traversed without difficulty.

Duodenum

Bulb: No visible ulcer or deformity.

Second part: Mucosa appears unremarkable to the examined extent.

Biopsy taken: Yes

Biopsy site: Antrum; body

Number of biopsies: Four; labelled by sampling site.

Assessment / Impression: Mild antral erythema. Gastric biopsies obtained; histology pending.

Recommendation: Review the histology with the referring clinician. This fictional example is not a management recommendation.

Clinician: DEMONSTRATION ONLY - NOT SIGNED **Signature:** _____