

Linear Algebra & Linear Dynamical Systems

AA 203 Recitation #1

April 10th, 2020

- ① Linear Algebra
- ② Linear Dynamical Systems (LDS)
 - a. Discrete time LDS.
 - b. Continuous time LDS.

Overview (cont.)

1 Linear Algebra

- a. Vector Spaces.
- b. Matrices and linear functions.
- d. Matrix Multiplication and Matrix Inverse.
- e. Singular Value Decomposition.
- f. Eigenvalue Decomposition.

Overview (cont.)

① Linear Algebra

- a. Vector Spaces.
- b. Matrices and linear functions.
- d. Matrix Multiplication and Matrix Inverse.
- e. Singular Value Decomposition.
- f. Eigenvalue Decomposition.

② Discrete time Linear Dynamical Systems

- a. Stabilization and tracking.
- b. Constraint structure.
- c. Operator norm & stability via linear feedback.

Overview (cont.)

① Linear Algebra

- a. Vector Spaces.
- b. Matrices and linear functions.
- d. Matrix Multiplication and Matrix Inverse.
- e. Singular Value Decomposition.
- f. Eigenvalue Decomposition.

② Discrete time Linear Dynamical Systems

- a. Stabilization and tracking.
- b. Constraint structure.
- c. Operator norm & stability via linear feedback.

③ Continuous time Linear Dynamical Systems

- a. Stabilization and tracking.
- b. State evolution.
- c. Diagonalization & stability via linear feedback.

1. Linear Algebra

1. Linear Algebra - Vector Spaces

We will use Euclidean vector spaces in this course.

- \mathbb{R}^n is the set of n -vectors.

1. Linear Algebra - Vector Spaces

We will use Euclidean vector spaces in this course.

- \mathbb{R}^n is the set of n -vectors.
- \mathbb{R} is the set of scalars.

1. Linear Algebra - Vector Spaces

We will use Euclidean vector spaces in this course.

- \mathbb{R}^n is the set of n -vectors.
- \mathbb{R} is the set of scalars.
- Vector addition is **entrywise**: $x + y = z$ means $x_i + y_i = z_i$ for all $1 \leq i \leq n$. **Example**:

$$\begin{bmatrix} 2 \\ 5 \end{bmatrix} + \begin{bmatrix} 9 \\ 2 \end{bmatrix} = \begin{bmatrix} 11 \\ 7 \end{bmatrix}$$

1. Linear Algebra - Vector Spaces

We will use Euclidean vector spaces in this course.

- \mathbb{R}^n is the set of n -vectors.
- \mathbb{R} is the set of scalars.
- Vector addition is **entrywise**: $x + y = z$ means $x_i + y_i = z_i$ for all $1 \leq i \leq n$. **Example**:

$$\begin{bmatrix} 2 \\ 5 \end{bmatrix} + \begin{bmatrix} 9 \\ 2 \end{bmatrix} = \begin{bmatrix} 11 \\ 7 \end{bmatrix}$$

- Scalar multiplication is **entrywise**: $cx = y$ means $cx_i = y_i$ for all $1 \leq i \leq n$. **Example**:

$$2 \begin{bmatrix} 4 \\ 3 \end{bmatrix} = \begin{bmatrix} 8 \\ 6 \end{bmatrix}$$

1. Linear Algebra - Linear Combinations

Definition (Linear Combination)

A linear combination of the vectors x_1, x_2, \dots, x_m is any vector of the form

$$\sum_{i=1}^m c_i x_i \text{ where } c_1, \dots, c_m \in \mathbb{R}.$$

1. Linear Algebra - Linear Combinations

Definition (Linear Combination)

A linear combination of the vectors x_1, x_2, \dots, x_m is any vector of the form

$$\sum_{i=1}^m c_i x_i \text{ where } c_1, \dots, c_m \in \mathbb{R}.$$

Definition (Span)

The span of a collection of vectors x_1, \dots, x_m , denoted by $\text{span}(x_1, \dots, x_m)$ is the collection of all possible linear combinations.

$$\text{span}(x_1, \dots, x_m) := \left\{ y : y = \sum_{i=1}^m c_i x_i \text{ for some } c_1, \dots, c_m \in \mathbb{R} \right\}$$

1. Linear Algebra - Linear Independence

Definition (Linear Independence)

A collection of vectors x_1, \dots, x_m is linearly independent if no member can be written as a linear combination of the other members.

Equivalently, the vectors x_1, \dots, x_m are independent if

$$\sum_{i=1}^m c_i x_i = 0 \implies c_i = 0 \text{ for all } 1 \leq i \leq m.$$

1. Linear Algebra - Linear Subspaces

Definition (Linear Subspaces)

A set $V \subset \mathbb{R}^n$ is a linear subspace if it is

- 1 Closed under scalar multiplication: $v \in V \implies cv \in V$ for all $c \in \mathbb{R}$.
- 2 Closed under vector addition: $u, v \in V \implies u + v \in V$.

1. Linear Algebra - Linear Subspaces

Definition (Linear Subspaces)

A set $V \subset \mathbb{R}^n$ is a linear subspace if it is

- ① Closed under scalar multiplication: $v \in V \implies cv \in V$ for all $c \in \mathbb{R}$.
- ② Closed under vector addition: $u, v \in V \implies u + v \in V$.

Example 1: \mathbb{R}^n itself is a linear subspace!

1. Linear Algebra - Linear Subspaces

Definition (Linear Subspaces)

A set $V \subset \mathbb{R}^n$ is a linear subspace if it is

- ① Closed under scalar multiplication: $v \in V \implies cv \in V$ for all $c \in \mathbb{R}$.
- ② Closed under vector addition: $u, v \in V \implies u + v \in V$.

Example 1: \mathbb{R}^n itself is a linear subspace!

Example 2: For any set of vectors V_0 , $\text{span}(V_0)$ is a linear subspace.

1. Linear Algebra - Linear Subspaces

Definition (Linear Subspaces)

A set $V \subset \mathbb{R}^n$ is a linear subspace if it is

- ① Closed under scalar multiplication: $v \in V \implies cv \in V$ for all $c \in \mathbb{R}$.
- ② Closed under vector addition: $u, v \in V \implies u + v \in V$.

Example 1: \mathbb{R}^n itself is a linear subspace!

Example 2: For any set of vectors V_0 , $\text{span}(V_0)$ is a linear subspace.

Example 3: $\{x \in \mathbb{R}^n : x_1 = 0\}$ is a linear subspace.

1. Linear Algebra - Spanning sets, Basis, Dimension

Definition (Spanning Set)

Given a linear subspace V , a collection of vectors x_1, \dots, x_m is a spanning set if $\text{span}(x_1, \dots, x_m) = V$.

1. Linear Algebra - Spanning sets, Basis, Dimension

Definition (Spanning Set)

Given a linear subspace V , a collection of vectors x_1, \dots, x_m is a spanning set if $\text{span}(x_1, \dots, x_m) = V$.

Definition (Basis)

A collection of vectors x_1, \dots, x_m is a basis for a subspace V if a) it is a spanning set of V and b) it is linearly independent.

1. Linear Algebra - Spanning sets, Basis, Dimension

Definition (Spanning Set)

Given a linear subspace V , a collection of vectors x_1, \dots, x_m is a spanning set if $\text{span}(x_1, \dots, x_m) = V$.

Definition (Basis)

A collection of vectors x_1, \dots, x_m is a basis for a subspace V if a) it is a spanning set of V and b) it is linearly independent.

Definition (Dimension)

Every basis of a subspace V contains the same number of vectors. This number is the dimension of V .

1. Linear Algebra - Vector Spaces

Definition

Given $x, y \in \mathbb{R}^n$, their dot product $\langle x, y \rangle$ is defined as:

$$\langle x, y \rangle := \sum_{i=1}^n x_i y_i.$$

1. Linear Algebra - Matrices

A matrix $A \in \mathbb{R}^{m \times n}$ is an array of numbers with m rows and n columns. A_{ij} is the number in the i th row and j th column.

1. Linear Algebra - Matrices

A matrix $A \in \mathbb{R}^{m \times n}$ is an array of numbers with m rows and n columns. A_{ij} is the number in the i th row and j th column.

Example: Below is a 4×3 matrix, A . Note that $A_{1,3} = 5$.

$$A = \begin{bmatrix} 4 & 1 & 5 \\ 2 & 2 & 1 \\ 3 & 0 & 9 \\ 0 & 0 & 7 \end{bmatrix}$$

1. Linear Algebra - Matrices

A matrix $A \in \mathbb{R}^{m \times n}$ is an array of numbers with m rows and n columns. A_{ij} is the number in the i th row and j th column.

Example: Below is a 4×3 matrix, A . Note that $A_{1,3} = 5$.

$$A = \begin{bmatrix} 4 & 1 & 5 \\ 2 & 2 & 1 \\ 3 & 0 & 9 \\ 0 & 0 & 7 \end{bmatrix}$$

Vectors are also matrices! A vector $x \in \mathbb{R}^n$ is a $n \times 1$ matrix.

1. Linear Algebra - Matrix Transpose

If $A \in \mathbb{R}^{m \times n}$, then its transpose is $A' \in \mathbb{R}^{n \times m}$ and

$$A_{ij} = A'_{ji}.$$

1. Linear Algebra - Matrix Transpose

If $A \in \mathbb{R}^{m \times n}$, then its transpose is $A' \in \mathbb{R}^{n \times m}$ and

$$A_{ij} = A'_{ji}.$$

Example:

$$A = \begin{bmatrix} 1 & 6 \\ 0 & 2 \\ 5 & 3 \end{bmatrix}, A' = \begin{bmatrix} 1 & 0 & 5 \\ 6 & 2 & 3 \end{bmatrix}$$

1. Linear Algebra - Matrix Multiplication

Matrix Multiplication: For $A \in \mathbb{R}^{m \times n}$, $B \in \mathbb{R}^{p \times q}$, AB exists if and only if $n = p$. In this case, AB is a $m \times q$ matrix.

1. Linear Algebra - Matrix Multiplication

Matrix Multiplication: For $A \in \mathbb{R}^{m \times n}$, $B \in \mathbb{R}^{p \times q}$, AB exists if and only if $n = p$. In this case, AB is a $m \times q$ matrix.

Furthermore, $(AB)_{ij} = \langle A_i, B_j \rangle$.

A_i = i th row of A

B_j is the j th column of B .

1. Linear Algebra - Matrix Multiplication

Matrix Multiplication: For $A \in \mathbb{R}^{m \times n}$, $B \in \mathbb{R}^{p \times q}$, AB exists if and only if $n = p$. In this case, AB is a $m \times q$ matrix.

Furthermore, $(AB)_{ij} = \langle A_i, B_j \rangle$.

A_i = i th row of A

B_j is the j th column of B .

Dot product as matrix multiplication: For $x, y \in \mathbb{R}^n$,

$$\langle x, y \rangle = x'y = y'x.$$

1. Linear Algebra - Linear functions

Definition (Linear functions)

A function $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is linear if for any $x, y \in \mathbb{R}^n$ and any $c_1, c_2 \in \mathbb{R}$,

$$f(c_1x + c_2y) = c_1f(x) + c_2f(y).$$

1. Linear Algebra - Linear functions

Definition (Linear functions)

A function $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is linear if for any $x, y \in \mathbb{R}^n$ and any $c_1, c_2 \in \mathbb{R}$,

$$f(c_1x + c_2y) = c_1f(x) + c_2f(y).$$

Example: Any matrix $A \in \mathbb{R}^{m \times n}$ defines a linear function $f(x) := Ax$.

$$\begin{aligned} f(c_1x + c_2y) &= A(c_1x + c_2y) = A(c_1x) + A(c_2y) = c_1Ax + c_2Ay \\ &= c_1f(x) + c_2f(y). \end{aligned}$$

1. Linear Algebra - Linear Representation

All linear functions are matrices!

1. Linear Algebra - Linear Representation

All linear functions are matrices!

Theorem

For every linear $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$, there exists a matrix A so that $f(x) = Ax$ for every $x \in \mathbb{R}^n$.

1. Linear Algebra - Linear Representation

All linear functions are matrices!

Theorem

For every linear $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$, there exists a matrix A so that $f(x) = Ax$ for every $x \in \mathbb{R}^n$.

Idea: Let e_1, e_2, \dots, e_n be a basis of \mathbb{R}^n . By linearity,

$$f(x) = f\left(\sum_{i=1}^n x_i e_i\right) = \sum_{i=1}^n x_i f(e_i).$$

1. Linear Algebra - Linear Representation

All linear functions are matrices!

Theorem

For every linear $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$, there exists a matrix A so that $f(x) = Ax$ for every $x \in \mathbb{R}^n$.

Idea: Let e_1, e_2, \dots, e_n be a basis of \mathbb{R}^n . By linearity,

$$f(x) = f\left(\sum_{i=1}^n x_i e_i\right) = \sum_{i=1}^n x_i f(e_i).$$

$\implies f$ is entirely determined by its behavior on a basis.

1. Linear Algebra - Linear Representation

All linear functions are matrices!

Theorem

For every linear $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$, there exists a matrix A so that $f(x) = Ax$ for every $x \in \mathbb{R}^n$.

Idea: Let e_1, e_2, \dots, e_n be a basis of \mathbb{R}^n . By linearity,

$$f(x) = f\left(\sum_{i=1}^n x_i e_i\right) = \sum_{i=1}^n x_i f(e_i).$$

$\implies f$ is entirely determined by its behavior on a basis.

A matrix A specifies the image of the standard basis vectors.

1. Linear Algebra - Linear Representation

All linear functions are matrices!

Theorem

For every linear $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$, there exists a matrix A so that $f(x) = Ax$ for every $x \in \mathbb{R}^n$.

Idea: Let e_1, e_2, \dots, e_n be a basis of \mathbb{R}^n . By linearity,

$$f(x) = f\left(\sum_{i=1}^n x_i e_i\right) = \sum_{i=1}^n x_i f(e_i).$$

$\implies f$ is entirely determined by its behavior on a basis.

A matrix A specifies the image of the standard basis vectors.

e_i is mapped to the i th column of A .

1. Linear Algebra - Operator Subspaces

For $A \in \mathbb{R}^{m \times n}$, let

- ① $R \subset \mathbb{R}^n$ be the set of rows of A .
- ② $C \subset \mathbb{R}^m$ be the set of columns of A .

1. Linear Algebra - Operator Subspaces

For $A \in \mathbb{R}^{m \times n}$, let

- 1 $R \subset \mathbb{R}^n$ be the set of rows of A .
- 2 $C \subset \mathbb{R}^m$ be the set of columns of A .

Definition (Row space)

The row space of A is denoted $\text{row}(A) := \text{span}(R)$.

1. Linear Algebra - Operator Subspaces

For $A \in \mathbb{R}^{m \times n}$, let

- 1 $R \subset \mathbb{R}^n$ be the set of rows of A .
- 2 $C \subset \mathbb{R}^m$ be the set of columns of A .

Definition (Row space)

The row space of A is denoted $\text{row}(A) := \text{span}(R)$.

Definition (Column space)

The column space of A is denoted $\text{col}(A) := \text{span}(C)$.

1. Linear Algebra - Operator Subspaces

For $A \in \mathbb{R}^{m \times n}$, let

- 1 $R \subset \mathbb{R}^n$ be the set of rows of A .
- 2 $C \subset \mathbb{R}^m$ be the set of columns of A .

Definition (Row space)

The row space of A is denoted $\text{row}(A) := \text{span}(R)$.

Definition (Column space)

The column space of A is denoted $\text{col}(A) := \text{span}(C)$.

Definition

The kernel of A is denoted $\text{ker}(A) := \{x \in \mathbb{R}^n : Ax = 0\}$.

1. Linear Algebra - Operator Subspaces

$\text{row}(A)$, $\text{col}(A)$, $\text{ker}(A)$ are all subspaces.

1. Linear Algebra - Operator Subspaces

$\text{row}(A), \text{col}(A), \ker(A)$ are all subspaces.

$\text{row}(A), \text{col}(A)$ have the same dimension. This dimension is also the rank of A .

1. Linear Algebra - Operator Subspaces

$\text{row}(A)$, $\text{col}(A)$, $\text{ker}(A)$ are all subspaces.

$\text{row}(A)$, $\text{col}(A)$ have the same dimension. This dimension is also the rank of A .

A is full rank if its rank is $\min(n, m)$.

1. Linear Algebra - Operator Subspaces

$\text{row}(A)$, $\text{col}(A)$, $\text{ker}(A)$ are all subspaces.

$\text{row}(A)$, $\text{col}(A)$ have the same dimension. This dimension is also the rank of A .

A is full rank if its rank is $\min(n, m)$.

Rank-Nullity: $\text{rank}(A) + \dim(\text{ker}(A)) = n$.

1. Linear Algebra - Identity Matrix

$I_n \in \mathbb{R}^{n \times n}$ is the identity matrix:

$$(I_n)_{ij} = \begin{cases} 1 & \text{if } i = j \\ 0 & \text{otherwise.} \end{cases}$$

I_n is a **multiplicative identity**: $AI_n = A$ and $I_n B = B$ whenever A, B have compatible dimensions to multiply with I_n .

1. Linear Algebra - Matrix Inverse & Orthogonal Matrices

Definition (Matrix Inverse)

The inverse of a matrix $A \in \mathbb{R}^{n \times n}$ is a matrix A^{-1} such that $AA^{-1} = A^{-1}A = I_n$.

Remark: Not all matrices have inverses. A matrix has an inverse if and only if it is full rank.

1. Linear Algebra - Matrix Inverse & Orthogonal Matrices

Definition (Matrix Inverse)

The inverse of a matrix $A \in \mathbb{R}^{n \times n}$ is a matrix A^{-1} such that $AA^{-1} = A^{-1}A = I_n$.

Remark: Not all matrices have inverses. A matrix has an inverse if and only if it is full rank.

Definition (Orthogonal Matrix)

A matrix $A \in \mathbb{R}^{n \times n}$ is orthogonal if and only if $A' = A^{-1}$.

Orthogonal Matrices are Isometries

Theorem (Orthogonal Matrices preserve length)

If $U \in \mathbb{R}^{n \times n}$ is orthogonal, then for every $x \in \mathbb{R}^n$,

$$\|Ux\|_2 = \|x\|_2.$$

Orthogonal Matrices are Isometries

Theorem (Orthogonal Matrices preserve length)

If $U \in \mathbb{R}^{n \times n}$ is orthogonal, then for every $x \in \mathbb{R}^n$,

$$\|Ux\|_2 = \|x\|_2.$$

Proof:

$$\|x\|_2^2 = x'x = x'I_nx = x'U'Ux = (Ux)'(Ux) = \|Ux\|_2^2.$$

Take square roots of both sides.

1. Linear Algebra - Singular Value Decomposition

For any matrix $A \in \mathbb{R}^{m \times n}$, we can factorize

$$A = U\Sigma V'$$

where $U \in \mathbb{R}^{m \times m}$, $V \in \mathbb{R}^{n \times n}$ are orthogonal matrices,

$\Sigma \in \mathbb{R}^{m \times n}$ is diagonal:

- $\Sigma_{ij} = 0$ if $i \neq j$.
- $\Sigma_{ii} \geq 0$.

1. Linear Algebra - Eigenvalue Decomposition

For any symmetric matrix $A \in \mathbb{R}^{n \times n}$, we can factorize

$$A = U\Lambda U'$$

where $U \in \mathbb{R}^{n \times n}$ is orthogonal and $\Lambda \in \mathbb{R}^{n \times n}$ is a diagonal matrix.

2. Discrete Linear Dynamical Systems

2. Discrete Linear Dynamical Systems

Dynamics:

$$x_{k+1} = Ax_k + Bu_k$$

- Time is discrete $k \in \{1, 2, 3, \dots\}$.

2. Discrete Linear Dynamical Systems

Dynamics:

$$x_{k+1} = Ax_k + Bu_k$$

- Time is discrete $k \in \{1, 2, 3, \dots\}$.
- $x_k \in \mathbb{R}^n$ describes the system state at timestep k .

2. Discrete Linear Dynamical Systems

Dynamics:

$$x_{k+1} = Ax_k + Bu_k$$

- Time is discrete $k \in \{1, 2, 3, \dots\}$.
- $x_k \in \mathbb{R}^n$ describes the system state at timestep k .
- $u_k \in \mathbb{R}^m$ is the control/action that is applied at timestep k .

2. Discrete Linear Dynamical Systems

Dynamics:

$$x_{k+1} = Ax_k + Bu_k$$

- Time is discrete $k \in \{1, 2, 3, \dots\}$.
- $x_k \in \mathbb{R}^n$ describes the system state at timestep k .
- $u_k \in \mathbb{R}^m$ is the control/action that is applied at timestep k .
- $A \in \mathbb{R}^{n \times n}, B \in \mathbb{R}^{n \times m}$.

2. Discrete Linear Dynamical Systems - Stabilizing

Stabilization Objective: Try to get x_k to converge to the origin as fast as possible.

$$\begin{aligned} & \underset{\{u_k\}_{k=1}^{T-1}}{\text{minimize}} && \sum_{k=1}^T \|x_k\|_2^2 \\ & \text{s.t.} && x_{k+1} = Ax_k + Bu_k \text{ for } 1 \leq k \leq T-1 \end{aligned}$$

2. Discrete Linear Dynamical Systems - Stabilizing

Stabilization Objective:

- We can penalize deviations in different coordinates by different amounts.

2. Discrete Linear Dynamical Systems - Stabilizing

Stabilization Objective:

- We can penalize deviations in different coordinates by different amounts.
- We can add penalty for using large control u_k .

2. Discrete Linear Dynamical Systems - Stabilizing

Stabilization Objective:

- We can penalize deviations in different coordinates by different amounts.
- We can add penalty for using large control u_k .

$$\begin{aligned} \underset{\{u_k\}_{k=1}^{T-1}}{\text{minimize}} \quad & \sum_{k=1}^T x_k' R x_k + u_k' Q u_k \\ \text{s.t.} \quad & x_{k+1} = A x_k + B u_k \text{ for } 1 \leq k \leq T-1 \end{aligned}$$

where $R, Q \succeq 0$.

2. Discrete Linear Dynamical Systems - Optimization

The relation between x_{k+1} , x_k , u_k is **linear**. We can incorporate the dynamics constraints as one large matrix equation: $\mathcal{A}z = 0$.

2. Discrete Linear Dynamical Systems - Optimization

The relation between x_{k+1}, x_k, u_k is **linear**. We can incorporate the dynamics constraints as one large matrix equation: $Az = 0$.

$$\underbrace{\begin{bmatrix} A & B & -I_n & 0 & 0 & \dots & 0 & 0 & 0 \\ 0 & 0 & A & B & -I_n & \dots & 0 & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \dots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & 0 & 0 & \dots & A & B & -I_n \end{bmatrix}}_A \underbrace{\begin{bmatrix} x_1 \\ u_1 \\ x_2 \\ u_2 \\ x_3 \\ \vdots \\ x_{T-1} \\ u_{T-1} \\ x_T \end{bmatrix}}_z = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 0 \end{bmatrix}$$

2. Discrete Linear Dynamical Systems - Optimization

Thus the stabilization and tracking problems can be solved by optimizing an objective function subject to linear equality constraints:

$$\begin{aligned} & \underset{z}{\text{minimize}} \ J(z) \\ & \text{s.t.} \ \mathcal{A}z = 0. \end{aligned}$$

2. Discrete Linear Dynamical Systems - Control Synthesis

How do we choose u so that $\lim_{k \rightarrow \infty} x_k = 0$?

2. Discrete Linear Dynamical Systems - Control Synthesis

How do we choose u so that $\lim_{k \rightarrow \infty} x_k = 0$?

Idea: Let's try linear state feedback, where $u_k = -Kx_k$ for some $K \in \mathbb{R}^{m \times n}$.

2. Discrete Linear Dynamical Systems - Control Synthesis

How do we choose u so that $\lim_{k \rightarrow \infty} x_k = 0$?

Idea: Let's try linear state feedback, where $u_k = -Kx_k$ for some $K \in \mathbb{R}^{m \times n}$. Then the dynamics become

$$\begin{aligned}x_{k+1} &= Ax_k + Bu_k \\&= Ax_k - BKx_k = (A - BK)x_k\end{aligned}$$

2. Discrete Linear Dynamical Systems - Control Synthesis

How do we choose u so that $\lim_{k \rightarrow \infty} x_k = 0$?

Idea: Let's try linear state feedback, where $u_k = -Kx_k$ for some $K \in \mathbb{R}^{m \times n}$. Then the dynamics become

$$\begin{aligned}x_{k+1} &= Ax_k + Bu_k \\ &= Ax_k - BKx_k = (A - BK)x_k\end{aligned}$$

If $(A - BK)$ is symmetric, then we can factor it $(A - BK) = U\Lambda U'$

2. Discrete Linear Dynamical Systems - Control Synthesis

How do we choose u so that $\lim_{k \rightarrow \infty} x_k = 0$?

Idea: Let's try linear state feedback, where $u_k = -Kx_k$ for some $K \in \mathbb{R}^{m \times n}$. Then the dynamics become

$$\begin{aligned}x_{k+1} &= Ax_k + Bu_k \\&= Ax_k - BKx_k = (A - BK)x_k\end{aligned}$$

If $(A - BK)$ is symmetric, then we can factor it $(A - BK) = U\Lambda U'$ Thus

$$\begin{aligned}x_{k+1} &= (A - BK)x_k \\ \iff x_{k+1} &= U\Lambda U'x_k \\ \implies U'x_{k+1} &= \underbrace{U'U}_{I_n}\Lambda U'x_k\end{aligned}$$

2. Discrete Linear Dynamical Systems - Control Synthesis

How do we choose u so that $\lim_{k \rightarrow \infty} x_k = 0$?

Idea: Let's try linear state feedback, where $u_k = -Kx_k$ for some $K \in \mathbb{R}^{m \times n}$. Then the dynamics become

$$\begin{aligned}x_{k+1} &= Ax_k + Bu_k \\&= Ax_k - BKx_k = (A - BK)x_k\end{aligned}$$

If $(A - BK)$ is symmetric, then we can factor it $(A - BK) = U\Lambda U'$ Thus

$$\begin{aligned}x_{k+1} &= (A - BK)x_k \\ \implies U'x_{k+1} &= \underbrace{U'U}_{I_n} \Lambda U'x_k\end{aligned}$$

2. Discrete Linear Dynamical Systems - Control Synthesis

How do we choose u so that $\lim_{k \rightarrow \infty} x_k = 0$?

Idea: Let's try linear state feedback, where $u_k = -Kx_k$ for some $K \in \mathbb{R}^{m \times n}$. Then the dynamics become

$$\begin{aligned}x_{k+1} &= Ax_k + Bu_k \\&= Ax_k - BKx_k = (A - BK)x_k\end{aligned}$$

If $(A - BK)$ is symmetric, then we can factor it $(A - BK) = U\Lambda U'$ Thus

$$\begin{aligned}x_{k+1} &= (A - BK)x_k \\&\iff x_{k+1} = U\Lambda U'x_k \\&\implies U'x_{k+1} = \underbrace{U'U}_{I_n}\Lambda U'x_k\end{aligned}$$

2. Discrete Linear Dynamical Systems - Control Synthesis

Defining $y_k := U'x_k$ gives:

2. Discrete Linear Dynamical Systems - Control Synthesis

Defining $y_k := U'x_k$ gives:

$$U'x_{k+1} = \Lambda U'x_k \iff y_{k+1} = \Lambda y_k$$

2. Discrete Linear Dynamical Systems - Control Synthesis

Defining $y_k := U'x_k$ gives:

$$U'x_{k+1} = \Lambda U'x_k \iff y_{k+1} = \Lambda y_k$$

So $y_{k+1,i} = \Lambda_{ii}y_{k,i}$ for each coordinate.

2. Discrete Linear Dynamical Systems - Control Synthesis

Defining $y_k := U'x_k$ gives:

$$U'x_{k+1} = \Lambda U'x_k \iff y_{k+1} = \Lambda y_k$$

So $y_{k+1,i} = \Lambda_{ii}y_{k,i}$ for each coordinate.

Iterate: $y_{k,i} = (\Lambda_{ii})^k y_{0,i}$.

2. Discrete Linear Dynamical Systems - Control Synthesis

$$U \text{ orthogonal} \implies \|x_k\|_2 = \|U'x_k\|_2 = \|y_k\|_2.$$

2. Discrete Linear Dynamical Systems - Control Synthesis

$$U \text{ orthogonal} \implies \|x_k\|_2 = \|U'x_k\|_2 = \|y_k\|_2.$$

$$\text{So } x_k \rightarrow 0 \iff y_k \rightarrow 0.$$

2. Discrete Linear Dynamical Systems - Control Synthesis

$$U \text{ orthogonal} \implies \|x_k\|_2 = \|U'x_k\|_2 = \|y_k\|_2.$$

$$\text{So } x_k \rightarrow 0 \iff y_k \rightarrow 0.$$

$$\text{Next, } y_k \rightarrow 0 \iff y_{k,i} \rightarrow 0 \text{ for all } i.$$

2. Discrete Linear Dynamical Systems - Control Synthesis

$$U \text{ orthogonal} \implies \|x_k\|_2 = \|U'x_k\|_2 = \|y_k\|_2.$$

$$\text{So } x_k \rightarrow 0 \iff y_k \rightarrow 0.$$

$$\text{Next, } y_k \rightarrow 0 \iff y_{k,i} \rightarrow 0 \text{ for all } i.$$

$$y_{k,i} = (\Lambda_{ii})^k y_{0,i} \rightarrow 0 \iff |\Lambda_{ii}| < 1.$$

2. Discrete Linear Dynamical Systems - Control Synthesis

$$U \text{ orthogonal} \implies \|x_k\|_2 = \|U'x_k\|_2 = \|y_k\|_2.$$

$$\text{So } x_k \rightarrow 0 \iff y_k \rightarrow 0.$$

$$\text{Next, } y_k \rightarrow 0 \iff y_{k,i} \rightarrow 0 \text{ for all } i.$$

$$y_{k,i} = (\Lambda_{ii})^k y_{0,i} \rightarrow 0 \iff |\Lambda_{ii}| < 1.$$

So $x_k \rightarrow 0$ if all eigenvalues of $A - BK$ have magnitude less than 1.

2. Discrete Linear Dynamical Systems - Control Synthesis

$$U \text{ orthogonal} \implies \|x_k\|_2 = \|U'x_k\|_2 = \|y_k\|_2.$$

$$\text{So } x_k \rightarrow 0 \iff y_k \rightarrow 0.$$

$$\text{Next, } y_k \rightarrow 0 \iff y_{k,i} \rightarrow 0 \text{ for all } i.$$

$$y_{k,i} = (\Lambda_{ii})^k y_{0,i} \rightarrow 0 \iff |\Lambda_{ii}| < 1.$$

So $x_k \rightarrow 0$ if all eigenvalues of $A - BK$ have magnitude less than 1.

If we can choose K so that all eigenvalues of $A - BK$ have magnitude less than 1, then $u(t) = -Kx(t)$ will be a stabilizing controller!

3. Continuous Linear Dynamical Systems

3. Continuous Linear Dynamical Systems

Dynamics:

$$\frac{d}{dt}x(t) = Ax(t) + Bu(t)$$

(shorthand ver.) $\dot{x} = Ax + Bu$

- Time is continuous $t \in \mathbb{R}$.

3. Continuous Linear Dynamical Systems

Dynamics:

$$\frac{d}{dt}x(t) = Ax(t) + Bu(t)$$

(shorthand ver.) $\dot{x} = Ax + Bu$

- Time is continuous $t \in \mathbb{R}$.
- $x(t) \in \mathbb{R}^n$ describes the system state at time t .

3. Continuous Linear Dynamical Systems

Dynamics:

$$\frac{d}{dt}x(t) = Ax(t) + Bu(t)$$

(shorthand ver.) $\dot{x} = Ax + Bu$

- Time is continuous $t \in \mathbb{R}$.
- $x(t) \in \mathbb{R}^n$ describes the system state at time t .
- $u(t) \in \mathbb{R}^m$ is the control/action that is applied at time t .

3. Continuous Linear Dynamical Systems

Dynamics:

$$\frac{d}{dt}x(t) = Ax(t) + Bu(t)$$

(shorthand ver.) $\dot{x} = Ax + Bu$

- Time is continuous $t \in \mathbb{R}$.
- $x(t) \in \mathbb{R}^n$ describes the system state at time t .
- $u(t) \in \mathbb{R}^m$ is the control/action that is applied at time t .
- $A \in \mathbb{R}^{n \times n}$, $B \in \mathbb{R}^{n \times m}$.

3. Continuous Linear Dynamical Systems - Stabilizing

Stabilizing Objective: Guide $x(t)$ to zero.

$$\begin{aligned} & \underset{u: [0, T] \rightarrow \mathbb{R}^m}{\text{minimize}} && \int_0^T x'(t) R x(t) + u'(t) Q u(t) dt \\ & \text{s.t.} && \dot{x}(t) = A x(t) + B u(t). \end{aligned}$$

where $R, Q \succeq 0$.

3. Continuous Linear Dynamical Systems - Control Synthesis

Objective: How do we choose u so that $\lim_{t \rightarrow \infty} x(t) \rightarrow 0$?

3. Continuous Linear Dynamical Systems - Control Synthesis

Objective: How do we choose u so that $\lim_{t \rightarrow \infty} x(t) \rightarrow 0$?

Idea: Let's try linear state feedback, where $u(t) = -Kx(t)$ for some $K \in \mathbb{R}^{m \times n}$.

3. Continuous Linear Dynamical Systems - Control Synthesis

Objective: How do we choose u so that $\lim_{t \rightarrow \infty} x(t) \rightarrow 0$?

Idea: Let's try linear state feedback, where $u(t) = -Kx(t)$ for some $K \in \mathbb{R}^{m \times n}$. The dynamics become

$$\begin{aligned}\dot{x}(t) &= Ax(t) + Bu(t) \\ &= Ax(t) - BKx(t) = (A - BK)x(t).\end{aligned}$$

3. Continuous Linear Dynamical Systems - Control Synthesis

Objective: How do we choose u so that $\lim_{t \rightarrow \infty} x(t) \rightarrow 0$?

Idea: Let's try linear state feedback, where $u(t) = -Kx(t)$ for some $K \in \mathbb{R}^{m \times n}$. The dynamics become

$$\begin{aligned}\dot{x}(t) &= Ax(t) + Bu(t) \\ &= Ax(t) - BKx(t) = (A - BK)x(t).\end{aligned}$$

If $A - BK$ is symmetric, then $A - BK = U\Lambda U'$ for some diagonal Λ and orthogonal U . Thus

$$\begin{aligned}\dot{x}(t) &= U\Lambda U'x(t) \\ \implies U'\dot{x}(t) &= \Lambda U'x(t).\end{aligned}$$

3. Continuous Linear Dynamical Systems - Control Synthesis

Define $y(t) := U'x(t)$.

3. Continuous Linear Dynamical Systems - Control Synthesis

Define $y(t) := U'x(t)$. Then

$$U'\dot{x}(t) = \Lambda U'x(t) \iff \dot{y}(t) = \Lambda y(t).$$

3. Continuous Linear Dynamical Systems - Control Synthesis

Define $y(t) := U'x(t)$. Then

$$U'\dot{x}(t) = \Lambda U'x(t) \iff \dot{y}(t) = \Lambda y(t).$$

Since Λ is diagonal, $\dot{y}_i(t) = \Lambda_{ii}y_i(t)$ for all i .

3. Continuous Linear Dynamical Systems - Control Synthesis

Define $y(t) := U'x(t)$. Then

$$U'\dot{x}(t) = \Lambda U'x(t) \iff \dot{y}(t) = \Lambda y(t).$$

Since Λ is diagonal, $\dot{y}_i(t) = \Lambda_{ii}y_i(t)$ for all i . Solve this ODE!

$$\begin{aligned}\frac{\dot{y}_i(t)}{y_i(t)} &= \Lambda_{ii} \\ \implies \ln(y_i(t)) &= \Lambda_{ii}t + C \\ \implies y_i(t) &= e^C e^{\Lambda_{ii}t} = y_i(0)e^{\Lambda_{ii}t}.\end{aligned}$$

3. Continuous Linear Dynamical Systems - Control Synthesis

$$U \text{ orthogonal} \implies \|x(t)\|_2 = \|U'x(t)\|_2 = \|y(t)\|_2.$$

3. Continuous Linear Dynamical Systems - Control Synthesis

$$U \text{ orthogonal} \implies \|x(t)\|_2 = \|U'x(t)\|_2 = \|y(t)\|_2.$$

$$\text{So } x(t) \rightarrow 0 \iff y(t) \rightarrow 0.$$

3. Continuous Linear Dynamical Systems - Control Synthesis

$$U \text{ orthogonal} \implies \|x(t)\|_2 = \|U'x(t)\|_2 = \|y(t)\|_2.$$

$$\text{So } x(t) \rightarrow 0 \iff y(t) \rightarrow 0.$$

$$\text{Next, } y(t) \rightarrow 0 \iff y_i(t) \rightarrow 0 \text{ for all } i.$$

3. Continuous Linear Dynamical Systems - Control Synthesis

$$U \text{ orthogonal} \implies \|x(t)\|_2 = \|U'x(t)\|_2 = \|y(t)\|_2.$$

$$\text{So } x(t) \rightarrow 0 \iff y(t) \rightarrow 0.$$

$$\text{Next, } y(t) \rightarrow 0 \iff y_i(t) \rightarrow 0 \text{ for all } i.$$

$$y_i(t) = y_i(0)e^{\Lambda_{ii}t} \rightarrow 0 \iff \Lambda_{ii} < 0.$$

3. Continuous Linear Dynamical Systems - Control Synthesis

$$U \text{ orthogonal} \implies \|x(t)\|_2 = \|U'x(t)\|_2 = \|y(t)\|_2.$$

$$\text{So } x(t) \rightarrow 0 \iff y(t) \rightarrow 0.$$

$$\text{Next, } y(t) \rightarrow 0 \iff y_i(t) \rightarrow 0 \text{ for all } i.$$

$$y_i(t) = y_i(0)e^{\Lambda_{ii}t} \rightarrow 0 \iff \Lambda_{ii} < 0.$$

$$\text{So } x(t) \rightarrow 0 \text{ if all eigenvalues of } A - BK \text{ are negative.}$$

3. Continuous Linear Dynamical Systems - Control Synthesis

$$U \text{ orthogonal} \implies \|x(t)\|_2 = \|U'x(t)\|_2 = \|y(t)\|_2.$$

$$\text{So } x(t) \rightarrow 0 \iff y(t) \rightarrow 0.$$

$$\text{Next, } y(t) \rightarrow 0 \iff y_i(t) \rightarrow 0 \text{ for all } i.$$

$$y_i(t) = y_i(0)e^{\Lambda_{ii}t} \rightarrow 0 \iff \Lambda_{ii} < 0.$$

So $x(t) \rightarrow 0$ if all eigenvalues of $A - BK$ are negative.

If we can choose K so that $A - BK \prec 0$, then $u(t) = -Kx(t)$ will be a stabilizing controller!