

1 Introduction

2 Throughout the paper, every time we speak about matrices we mean binary
3 matrices (also called 01-matrices) and we omit the word binary. If we speak
4 about a *pattern*, we again mean a binary matrix and we use the word in order to
5 distinguish among more matrices as well as to indicate relationship.

6 When dealing with matrices, we always index rows and column starting with
7 one and when we speak about a row r , we simply mean a row with index r . A
8 *line* is a common word for both a row and a column. When we order a set of
9 lines, we first put all rows and then all columns. For $M \in \{0, 1\}^{m \times n}$, $[m]$ is a set
10 of all rows and $[m + n]$ is a set of all lines, where m -th element is the last row.
11 This goes with the usual notation.

12 **Notation 1.** For $n \in \mathbb{N}$ let $[n] := \{1, 2, \dots, n\}$ and for $m \in \mathbb{N}$, where $n \leq m$ let
13 $[n, m] := \{n, n + 1, \dots, m\}$.

14 **Notation 2.** For a matrix $M \in \{0, 1\}^{m \times n}$ and $L \subseteq [m + n]$ let $M[L]$ denote a
15 submatrix of M induced by lines in L .

16 **Notation 3.** For a matrix $M \in \{0, 1\}^{m \times n}$, $R \subseteq [m]$ and $C \subseteq [n]$ let $M[R, C]$
17 denote a submatrix of M induced by rows in R and columns in C . Furthermore,
18 for $r \in [m]$ and $c \in [n]$ let $M[r, c] := M[\{r\}, \{c\}] = M[\{r, c + m\}]$.

19 **Definition 1.** We say a matrix $M \in \{0, 1\}^{m \times n}$ *contains* a pattern $P \in \{0, 1\}^{k \times l}$
20 *as a submatrix* and denote it by $P \leq M$ if there are $R \in [m]$ and $C \in [n]$
21 such that $|R| = k$, $|C| = l$ and for every $r \in R$ and $c \in C$ if $P[r, c] = 1$, then
22 $M[R, C][r, c] = 1$.

23 This does not necessarily mean $P = M[R, C]$ as $M[R, C]$ can have more
24 one-entries than P does.

25 **Notation 4.** For a matrix $M \in \{0, 1\}^{m \times n}$ and $L \subseteq [m + n]$ let $M_{\preceq}[L]$ denote a
26 matrix acquired from M by applying following operation for each $l \in L$:

- 27 • If l is the first row in L then we replace the first l rows by one row that is
28 a bitwise OR of replaced rows.
- 29 • If l is the first column in L then we replace the first $l - m$ columns by one
30 column that is a bitwise OR of replaced columns.
- 31 • Otherwise, we take l 's predecessor $l' \in L$ in the standard ordering and
32 replace lines $[l' + 1, l]$ by one line that is a bitwise OR of replaced lines.

33 **Notation 5.** For a matrix $M \in \{0, 1\}^{m \times n}$, $R \subseteq [m]$ and $C \subseteq [n]$ let $M_{\preceq}[R, C] :=$
34 $M_{\preceq}[R \cup \{c + m | c \in C\}]$.

35 **Definition 2.** We say a matrix $M \in \{0, 1\}^{m \times n}$ *contains* a pattern $P \in \{0, 1\}^{k \times l}$
36 *as an interval minor* and denote it by $P \preceq M$ if there are $R \in [m]$ and $C \in [n]$
37 such that $|R| = k$, $|C| = l$ and for every $r \in R$ and $c \in C$ if $P[r, c] = 1$, then
38 $M_{\preceq}[R, C][r, c] = 1$.

39 **Observation 1.** For all matrices M and P , $P \leq M \Rightarrow P \preceq M$.

40 **Observation 2.** *For all matrices M and P , if P is a permutation matrix, then*
41 *$P \leq M \Leftrightarrow P \preceq M$.*

42 *Proof.* If we have $P \preceq M$, then there is a partitioning of M into rectangles and for
43 each one-entry of P there is at least one one-entry in the corresponding rectangle
44 of M . Since P is a permutation matrix, it is sufficient to take rows and columns
45 having at least one one-entry in the right rectangle and we can always do so.

46 Together with Observation 1 this gives us the statement. \square

0.1 Characterizations

Observation 3. Let $P \in \{0, 1\}^{k \times l}$ and $P' \in \{0, 1\}^{k \times l+1}$ such that $P' = P \oplus_h 0^{k \times 1}$, similarly let $M \in \{0, 1\}^{m \times n}$ and $M' \in \{0, 1\}^{m \times n+1}$ such that $M' = M \oplus_h 0^{m \times 1}$, then $P \preceq M \Leftrightarrow P' \preceq M'$.

Proof. \Rightarrow Clearly we can map the last column of P' to the last column of M' and then map (using OR) $P'[[k], [l]]$ to $M'[[m], [n]]$ the same way P is mapped to M .

\Leftarrow If $P' \preceq M$ we are done. Otherwise, the last column of P' needs to be mapped to the last column of M' and by deleting both from their matrix we get $P'[[k], [l]] \preceq M'[[m], [n]]$ which is the same as $P \preceq M$.

□

The same proof can be also used for adding an empty column as the first column or an empty row as the first or the last row. Using induction we can easily show that a pattern P' is avoided by a matrix M' if and only if P is avoided by M where P is derived from P' by excluding all empty beginning or ending rows and columns and M is derived from M' by excluding the same number of beginning or ending rows and columns. Therefore, when characterizing matrices avoiding a forbidden pattern, we do not need to consider patterns having empty rows or columns on their boundary.

Definition 3. A *walk* in a matrix M is a sequence of some of its entries, beginning in the top left corner and ending in the bottom right one. If an entry $M[i, j]$ is in the sequence, the next one is either $M[i + 1, j]$ or $M[i, j + 1]$.

Definition 4. We call a binary matrix M a *walking matrix* if there is a walk in M such that all one-entries of M are contained on the walk.

Definition 5. An *extended walk of size $k \times l$* in a matrix M is a subset of some of its entries, beginning in the top left corner and ending in the bottom right one. If an entry $M[i, j]$ is in the subset there is also either $M[i + 1, j]$ or $M[i, j + 1]$. The size describes that no more than k entries directly above each other are in the subset and no more than l entries directly next to each other are in the subset. We say that an extended walk of size $k \times l$ in M starts with a walk w , if the extended walk is a subset of entries of M that

- lie on w or below w and
- lie on w shifted by $k - 1$ down and by $l - 1$ to the left or above it.

Definition 6. For $M \in \{0, 1\}^{m \times n}$ and $r \in [m], c \in [n]$ we say $M[r, c]$ is

- *top-left empty* if $M[[r - 1], [c - 1]]$ is an empty matrix,
- *top-right empty* if $M[[r - 1], [c + 1, n]]$ is empty,
- *bottom-left empty* if $M[[r - 1], [c + 1, n]]$ is empty,
- *bottom-right empty* if $M[[r - 1], [c + 1, n]]$ is empty.

0.1.1 Patterns of size 2×2 and their generalization

Theorem 4. Let $P = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$, then for all M : $P \not\leq M \Leftrightarrow M$ is a walking matrix.

Proof. Since P is a permutation matrix, $P \not\leq M \Leftrightarrow P \not\leq M$ and it is easy to see $P \not\leq M \Leftrightarrow M$ is a walking matrix. \square

Now consider a generalization of the pattern from above:

Theorem 5. Let $P \in \{0, 1\}^{k \times l}$ be a matrix having only two one-entries – $P[1, n]$ and $P[m, 1]$, then for all M : $P \not\leq M \Leftrightarrow M$ has an extended walk of size $k-1 \times l-1$ containing all one-entries.

Proof. \Rightarrow Let $P \not\leq M$ and consider the left-most top-right empty elements of M . They necessarily form a walk w . For contradiction, assume there is a one-entry e below the extended walk of size $k-1 \times l-1$ starting with w . Since e is below the extended walk, there is an element e' – the right-most element of M that is neither below e nor to the right from e and at the same time still below the extended walk (it is possible $e = e'$). Let $e = M[r, c]$ and notice $M[r-k, c-l]$ is part of walk w and because of the choice of e' neither $M[r-k-1, c-l]$ nor $M[r-k, c-l-1]$ are on the walk w and $M[r-k, c-l]$ must be a one-entry; therefore, together with e it forms the forbidden pattern in M , which is a contradiction.

\Leftarrow Let $M[r, c]$ be any one-entry of M , which then necessarily lie in the extended walk. Because the size of the walk is $k-1 \times l-1$, $M[r-k+1, c-l+1]$ is top-left empty and $M[r+k-1, c+l-1]$ is bottom-right empty; therefore e cannot be a part of a mapping of P . \square

Theorem 6. Let $P = \begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix}$, then for all $M \in \{0, 1\}^{m \times n}$: $P \not\leq M \Leftrightarrow$ there exist a row r and a column c such that (see Figure 1)

- $M[[r-1], [c-1]]$ is empty,
- $M[[r-1], [c+1, n]]$ is empty,
- $M[[r+1, m], [c-1]]$ is empty and
- $M[[r, m], [c, n]]$ is a walking matrix.

Proof. \Rightarrow If $\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \not\leq M$ then M is a walking matrix and we set $r = c = 1$. Otherwise, there are one-entries $M[r, c']$ and $M[r', c]$ such that $r' < r$ and $c' < c$. If there is a one-entry in regions $M[[r-1], [c-1]]$, $M[[r-1], [c+1, n]]$ or $M[[r+1, m], [c-1]]$ then $P \leq M$. If $M[[r, m], [c, n]]$ is not a walking matrix then it contains $\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ and we again get a contradiction.

\Leftarrow For contradiction, assume that M described in Figure 1 contains P as an interval minor. It means that there is a partition of the matrix into four quadrants such that there is at least one one-entry in each quadrant besides the bottom right one. If the matrix is partitioned above the r -th row, then there is only one column containing one-entries and it is not possible for both

			c
	0		0
r	0		M'

Figure 1: Characterization of a matrix avoiding $\begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix}$ as an interval minor. Matrix M' is a walking matrix

top quadrants to have a one-entry. Similarly, if the matrix is partitioned to the left of the c -th column, there is only one row containing one-entries and there is no one-entry in either top-left or bottom-left quadrant. Therefore, the partitioning lies below the r -th row and to the right of the c -th column, but if the quadrants contain one-entries, there is a $\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ interval minor in M' , which is a contradiction with it being a walking matrix. \square

Theorem 7. Let $P \in \{0, 1\}^{k \times l}$ be a matrix having only three one-entries – $P[1, 1]$, $P[1, n]$ and $P[m, 1]$, then for all M : $P \not\leq M \Leftrightarrow$ there exist a row r and a column c such that (see Figure 1 and imagine rows and columns being extended)

- $M[[r - 1], [c - 1]]$ is empty,
- $M[[r - 1], [c + l, n]]$ is empty,
- $M[[r + k, m], [c - 1]]$ is empty and
- $M[[r, m], [c, n]]$ has an extended walk of size $k - 1 \times l - 1$ containing all one-entries.

Proof. Let $P' = P$ and set $P'[m, 1] = 0$ (P' is a generalization of $\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$).

\Rightarrow If $P' \not\leq M$ then M is a matrix having an extended walk of size $k - 1 \times l - 1$ containing all one-entries and we set $r = c = 1$. Otherwise, there are one-entries $M[r_1, c_1]$ and $M[r_2, c_2]$ such that $r_2 < r_1$ and $c_1 < c_2$. We now choose $M[r_3, c_3]$ to be the bottom-most one-entry that still forms P' with $M[r_2, c_2]$. We choose $M[r_4, c_4]$ to be the left-most one-entry that forms P' with $M[r_3, c_3]$ and set $r = r_3 - k + 1$ and $c = c_4 - l + 1$. If there is a one-entry in regions $M[[r - 1], [c - 1]]$, $M[[r - 1], [c + l, n]]$ or $M[[r + k, m], [c - 1]]$ then $P \leq M$. If $M[[r, m], [c, n]]$ is not a walking matrix then it contains P' and we again get a contradiction.

\Leftarrow Because of the sizes of areas with no one-entries and the condition for $M[[r, m], [c, n]]$, there cannot be P' anywhere but in $M[[r + k - 1], [c + l - 1]]$. Since $M[[r - 1], [c - 1]]$ is empty, there is no one-entry to map $P[1, 1]$ to; therefore, $P \not\leq M$.

154

□

155 **Lemma 8.** Let $P = \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix}$ and let $M \in \{0, 1\}^{m \times n}$ avoid P as an interval minor,
 156 then there exists a row r and a column c such that $M[r, c]$ is either

157 1. both top-left empty and bottom-right empty and $[r, c] \notin \{[1, n], [m, 1]\}$ or

158 2. both top-right empty and bottom-left empty and $[r, c] \notin \{[1, 1], [m, n]\}$.

159 *Proof.*

□

160 **Theorem 9.** Let $P = \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix}$, then for all M : $P \not\preceq M \Leftrightarrow M$ looks like one of the
 161 matrices in Figure 2, where $\begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix} \not\preceq M_1$, $\begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix} \not\preceq M_2$, $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \not\preceq M_3$ and $\begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix} \not\preceq M_4$.




Figure 2: Characterization of a matrix avoiding $\begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix}$ as an interval minor.

162 *Proof.*

163 \Rightarrow We proceed by induction by the size of M .

164 If $M \in \{0, 1\}^{2 \times 2}$ then it either avoids $\begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix}$ or $\begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix}$ and we are done.

165 For bigger M there is, from Lemma 8, “the element”. Assume the first case
 166 (top-right and bottom-left empty (will change this when I have some notation)).

167 If M_1 is non-empty, then $\begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix} \not\preceq M_2$; otherwise, $P \preceq M$. Similarly, $\begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix} \not\preceq M_1$ if
 168 M_2 is non-empty. If one of them is empty, the other is a smaller matrix avoiding
 169 P as an interval minor and by induction hypothesis, it can be partitioned. Adding
 170 empty rows and columns does not break any condition and we get a partitioning
 171 of the whole M .

172 \Leftarrow Without loss of generality, let us assume M looks like the left matrix in Figure 2.

173 For contradiction, assume $P \preceq M$. In that case, we can partition M into four
 174 quadrants such that there is at least one one-entry in each of them. It does not
 175 matter where we partition it, every time we either get $\begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix} \preceq M_1$ or $\begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix} \preceq M_2$,
 176 which is a contradiction. □

177 **Theorem 10.** Let $P \in \{0, 1\}^{k \times l}$ be a matrix having only four one-entries –
 178 $P[1, 1]$, $P[1, n]$, $P[m, 1]$ and $P[m, n]$, then for all M : $P \not\preceq M \Leftrightarrow M$ looks like
 179 one of the matrices in Figure 2, where generalized $\begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix} \not\preceq M_1$, $\begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix} \not\preceq M_2$,
 180 $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \not\preceq M_3$ and $\begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix} \not\preceq M_4$.

181 0.1.2 Matrices of size 2×3

182 **Theorem 11.** *Let $P = \begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}$, then for all M : $P \not\preceq M \Leftrightarrow M = M_1 \oplus_h M_2$ where*
 183 *$\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \not\preceq M_1$ and $\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \not\preceq M_2$.*

184 *Proof.* \Rightarrow Let $e = [r, c]$ be the top-most one-entry of M . If $\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \preceq M[[m], [c - 1]]$, together with e it forms P . If $\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \not\preceq M[[m], [c, n]]$ then we are done.
 185
 186 Let us assume it is not the case and let $e_{0,0}, e_{1,1}$ be any two one-entries forming the forbidden pattern. Symmetrically, let $\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \preceq M[[m], [c]]$ and
 187
 188 let $e_{0,1}, e_{1,0}$ be any two one-entries forming the forbidden pattern. Now if we take $e_{0,0}, e_{0,1}$ and $e_{1,0}$ or $e_{1,1}$ with bigger row, we get the forbidden
 189
 190 pattern P as an interval minor of M .

191 \Leftarrow For contradiction, let us assume $P \preceq M$ and $M = M_1 \oplus_h M_2$. If $P \preceq M$,
 192 look at the one-entry of M where the bottom one-entry of P is mapped. If
 193 it is in M_1 then $P \not\preceq M$ because $\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \not\preceq M_1$. Otherwise, $P \not\preceq M$ because
 194 $\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \not\preceq M_2$.
 195 □

196 **Lemma 12.** *Let $P = \begin{pmatrix} 1 & 1 & 1 \\ 0 & 1 & 0 \end{pmatrix}$, then for all M : $P \not\preceq M \Rightarrow M = M_1 \oplus_h M_2$ where*

- 197 1. $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \not\preceq M_1$ and $\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \not\preceq M_2$ or
 198 2. $\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \not\preceq M_1$ and $\begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix} \not\preceq M_2$.

199 *Proof.* Let $e = [r, c]$ be the top-most one-entry of M . If $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \preceq M[[m], [c - 1]]$,
 200 together with e it would be the whole P . Similarly, $\begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix} \not\preceq M[[m], [c + 1, n]]$.
 201 For contradiction with the statement, let $\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \preceq M[[m], [c]]$ and $e_{0,0}, e_{1,1}$ (none
 202 of them equal to e , since e lies in the top-right corner) be any two one-entries
 203 forming the pattern. Symmetrically, let $\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \preceq M[[m], [c, n]]$ and $e_{0,1}, e_{1,0}$ be
 204 any two one-entries forming the pattern. In that case $e_{0,0}, e, e_{0,1}$ and $e_{1,0}$ or $e_{1,1}$
 205 with bigger row give us the forbidden pattern P as an interval minor of M . □

206 **Theorem 13.** *Let $P = \begin{pmatrix} 1 & 1 & 1 \\ 0 & 1 & 0 \end{pmatrix}$, then for all M : $P \not\preceq M \Leftrightarrow M$ looks like the*
 207 *matrix in Figure 3 and $\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \not\preceq M_1$ and $\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \not\preceq M_2$.*




Figure 3: Characterization of a matrix avoiding $\begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix}$ as an interval minor.

208 *Proof.* \Rightarrow From Lemma 12 we know $M = M'_1 \oplus_h M'_2$ where $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \not\leq M'_1$ and
209 $\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \not\leq M'_2$. The second case would be dealt with symmetrically. From
210 Theorem 6 we have that M'_1 can be characterized exactly like $M[[m], [c_2 - 1]$
211 and $M[[m], [c_2, n]]$ forms a walking matrix. The only problem with our claim
212 would be if there were two different columns having a one-entry above the
213 r -th row. In that case, those two one-entries together with a one-entry in
214 the r -th row between the columns c_1 and c_2 and a one-entry in the c_1 -th
215 column above the r -th row form P as an interval minor.

216 \Leftarrow The bottom-middle one-entry of P can not be mapped anywhere but to the
217 r -th row, but in that case there are at most two columns having one-entries
218 above it.
219 □

0.2 Extremal function

Notation 6. Let M be a matrix. We denote $|M|$ the weight of M , the number of one-entries in M .

Usually $|M|$ stands for a determinant of matrix M . However, in this paper we do not work with determinants at all so the notation should not lead to misunderstanding.

Definition 7. For a matrix P we define $Ex(P, m, n) := \max\{|M| \mid M \in \{0, 1\}^{m \times n}, P \not\preceq M\}$. We denote $Ex(P, n) := Ex(P, n, n)$.

Definition 8. For a matrix P we define $Ex_{\preceq}(P, m, n) := \max\{|M| \mid M \in \{0, 1\}^{m \times n}, P \not\preceq M\}$. We denote $Ex_{\preceq}(P, n) := Ex_{\preceq}(P, n, n)$.

Observation 14. For all P, m, n ; $Ex_{\preceq}(P, m, n) \leq Ex(P, m, n)$.

Observation 15. If $P \in \{0, 1\}^{k \times l}$ has a one-entry at position $[a, b]$, then

$$Ex(P, m, n) \geq \begin{cases} m \cdot n & k > m \vee l > m \\ (k-1)n + (l-1)m - (k-1)(l-1) & \text{otherwise.} \end{cases}$$

Observation 16. The same holds for $Ex_{\preceq}(P, m, n)$.

Definition 9. $P \in \{0, 1\}^{k \times l}$ is (strongly) minimalist if

$$Ex(P, m, n) = \begin{cases} m \cdot n & k > m \vee l > m \\ (k-1)n + (l-1)m - (k-1)(l-1) & \text{otherwise.} \end{cases}$$

Definition 10. $P \in \{0, 1\}^{k \times l}$ is weakly minimalist if

$$Ex_{\preceq}(P, m, n) = \begin{cases} m \cdot n & k > m \vee l > m \\ (k-1)n + (l-1)m - (k-1)(l-1) & \text{otherwise.} \end{cases}$$

Observation 17. If P is strongly minimalist, then P is weakly minimalist.

0.2.1 Known results

Fact 18. 1. $\begin{pmatrix} 1 \end{pmatrix}$ is strongly minimalist.

2. If $P \in \{0, 1\}^{k \times l}$ is strongly minimalist and there is a one-entry in the last row in the c -th column, then $P' \in \{0, 1\}^{k+1 \times l}$, which is created from P by adding a new row having a one-entry only in the c -th column, is strongly minimalist.

3. If P is strongly minimalist, then after changing a one-entry into a zero-entry it is still strongly minimalist.

Fact 19. Let $P = \begin{pmatrix} 1 & \dots & 1 \\ 1 & \dots & 1 \end{pmatrix}$ have l columns, then P is weakly minimalist.

Proof. Let $M \in \{0, 1\}^{m \times n}$ be a matrix avoiding $P = \{1\}^{2 \times l}$ as an interval minor and $A_i = \{j \in [n] \mid \text{weight of } M[[i, \{j\}] > 0 \wedge \text{weight of } M[[i + 1, m], \{j\}] > 0\}$. Clearly $|A_i| \leq l - 1$, otherwise $P \preceq M$. Let b_j denote the number of one-entries in the j -th column. Each column j of M appears in at least $b_j - 1$ of sets A_i , $0 \leq i \leq m - 2$. It follows that

$$\text{weight of } M = \sum_{j=0}^n b_j = \sum_{j=0}^n (b_j - 1) + n \leq \sum_{i=0}^{m-2} |A_i| + n \leq (l - 1)(m - 1) + n$$

242

□

243 This result is indeed very important because it shows that there are matrices
244 like $\begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix}$, which are weakly minimalist, although it is known they are not strongly
245 minimalist.

246 **Fact 20.** Let $P = \begin{pmatrix} 1 & \dots & 1 \\ 1 & \dots & 1 \end{pmatrix}$ have l columns, then P is weakly minimalist.

Proof. Let $M \in \{0, 1\}^{m \times n}$ be a matrix avoiding $P = \{1\}^{3 \times l}$ as an interval minor and $A_i = \{j \in [n] \mid \text{weight of } M[[i - 1, \{j\}] > 0 \wedge \text{weight of } M[[i + 1, m], \{j\}] > 0 \wedge M[i, j] \text{ one-entry}\}$. Clearly $|A_i| \leq l - 1$, otherwise $P \preceq M$. Let b_j denote the number of one-entries in the j -th column. Each column j of M (for which $b_j \geq 2$) appears in exactly $b_j - 2$ of sets A_i , $1 \leq i \leq m - 1$. It follows that

$$\text{weight of } M = \sum_{j=0}^n b_j = \sum_{j=0}^n (b_j - 2) + 2n \leq \sum_{i=1}^{m-2} |A_i| + 2n \leq (l - 1)(m - 2) + 2n$$

247

□