

Sasa Mardi, EE23BTECH11222

Question NM 25: Consider the contour integral $\oint \frac{dz}{z^4 + z^3 - 2z^2}$, along the curve $|z| = 3$ oriented in the counterclockwise direction. If $\text{Res}[f, z_0]$ denotes the residue of $f(z)$ at the point z_0 , then which of the following are TRUE?

- 1) $\text{Res}[f, 0] = -\frac{1}{4}$
- 2) $\text{Res}[f, 1] = \frac{1}{3}$
- 3) $\text{Res}[f, -2] = -\frac{1}{12}$
- 4) $\text{Res}[f, 2] = -1$

(GATE NM 2023)

Solution:

$$\frac{dz}{z^4 + z^3 - 2z^2} = \frac{dz}{z^2(z-1)(z+2)} \quad (1)$$

Poles: $z = 0, 1, -2$

Curve: $|z| = 3$, all poles inside it

Given the function $f(z) = \frac{1}{z^2(z-1)(z+2)}$, with poles at $z = 0$, $z = 1$, and $z = -2$, and considering the curve $|z| = 3$, where all poles are inside it. We want to find the residues of $f(z)$ at these poles.

The general formula for finding the residue at a pole z_0 is:

$$\text{Res}(f, z_0) = \frac{1}{(n-1)!} \lim_{z \rightarrow z_0} \frac{d^{n-1}}{dz^{n-1}} ((z - z_0)^n f(z)) \quad (2)$$

where n denotes how many times the pole is repeated.

- For $z_0 = 0$, where $n = 2$, we have:

$$\text{Res}(f, 0) = \frac{1}{1!} \lim_{z \rightarrow 0} \frac{d}{dz} \left(\frac{1}{(z-1)(z+2)} \right) \quad (3)$$

$$= -\frac{1}{4} \quad (4)$$

- For $z_0 = 1$, where $n = 1$, we have:

$$\text{Res}(f, 1) = \frac{1}{(1-1)!} \lim_{z \rightarrow 1} \frac{z-1}{z^2(z+2)} \quad (5)$$

$$= \frac{1}{3} \quad (6)$$

- For $z_0 = -2$, where $n = 1$, we have:

$$\text{Res}(f, -2) = \frac{1}{(1-1)!} \lim_{z \rightarrow -2} \frac{z-(-2)}{z^2(z-1)} \quad (7)$$

$$= -\frac{1}{12} \quad (8)$$

Therefore, the correct answers are: options A, B, and C.




Fig. 1: Region of Convergence (ROC) for $|z| = 3$