

INVESTIGATING SUPERMASSIVE BLACK HOLES AND
THEIR VARIABILITY BY USING A STRUCTURE FUNCTION

by

Derek Blue

A THESIS SUBMITTED IN PARTIAL FULFILMENT OF
THE REQUIREMENTS FOR THE DEGREE OF

BACHELOR OF SCIENCE

in

Honours Mathematics

(Co-Supervised by Dr. Tina Harriott, Mount Saint Vincent University, Department
of Mathematics and Dr. Luigi Gallo, Saint Mary's University, Department of
Astronomy and Physics and Mount Saint Vincent University, Adjunct Professor)

.....
.....
.....
.....
.....

MOUNT SAINT VINCENT UNIVERSITY

March 10, 2019

© Derek Blue, 2019

ABSTRACT

INVESTIGATING SUPERMASSIVE BLACK HOLES AND THEIR VARIABILITY BY USING A STRUCTURE FUNCTION

Our universe is pockmarked with galaxies as far as we can currently observe from our humble planet. The currently accepted theoretical model is that at the center of these galaxies are super-massive black holes holding them together. This has been proven for a majority of these observable galaxies, including our own, and for some the central black hole is actively accreting matter. We call the galaxies hosting these black holes active galaxies. Active galaxies are of particular interest because they are some of the brightest objects in the night sky and yet their emission spectra are incredibly variable. Unfortunately since the data acquired from these objects is generally unevenly sampled, we cannot use a Fourier analysis as it suffers from windowing and aliasing. For these reasons we turn to the structure function which has a similar effect as the Fourier analysis but with the added benefits of remaining in the time domain and sporting a robustness against windowing and aliasing due to uneven sampling.

by *Derek Blue*

submitted on March 10, 2019:

Contents

Contents	iii
List of Figures	v
List of Tables	vi
1 INTRODUCTION	1
1.1 OVERVIEW	1
1.2 GALAXIES, ACTIVE GALAXIES, AND BLACK HOLES	2
1.2.1 GALAXIES	2
1.2.2 ACTIVE GALAXIES	4
1.2.3 BLACK HOLES	4
1.3 HOW AGN ARE OBSERVED	4
1.3.1 ORBITAL X-RAY TELESCOPES	4
1.3.2 THE UNIFIED MODEL	4
1.4 EXPECTED ACHIEVEMENTS	4
2 THE GENERAL RELATIVISTIC DESCRIPTION	5
2.1 GENERAL RELATIVITY	5
2.1.1 AN INTRODUCTION TO TENSORS	5

2.1.2	THE METRIC TENSOR	5
2.1.3	THE SPACETIME INTERVAL	5
2.2	THE KERR SPINNING BLACK HOLE	5
2.2.1	THE KERR SOLUTION	6
2.2.2	PROPERTIES OF THE KERR SOLUTION	6
3	STATISTICAL ANALYSIS	7
3.1	THE STRUCTURE FUNCTION	7
3.2	THE DATA	7
4	PROGRAMMING SFA	8
4.1	SFA USAGE AND DESCRIPTION	8
4.2	THE SFA ALGORITHM	8
4.2.1	PROCESS	8
4.2.2	COST AND COMPLEXITY	8
4.3	EXPECTED OUTPUT	8
5	RESULTS AND CONCLUSIONS	9
5.1	RESULTS	9
5.2	CONCLUSIONS	9
6	FUTURE WORK	10
A	SFA	11
	Bibliography	12

List of Figures

1.1 Hubble - de Vaucouleurs diagram for galaxy morphology. Image credit:

Antonio Ciccolella / M. De Leo - <https://en.wikipedia.org/wiki/File::Hubble->

Vaucouleurs.png 3

List of Tables

Chapter 1

INTRODUCTION

1.1 OVERVIEW

This chapter will give an overview of the theoretical models used by modern astronomers and physicists to describe black holes. The majority of this thesis is concerned with the analysis of the electromagnetic spectrum emitted by real super-massive black holes at the centers of various galaxies by using a structure function, which will be described in detail.

To begin, this chapter will give a brief overview of galaxies, active galaxies, black holes and their similarities and differences. This chapter will then go into detail on how we can observe these astronomical objects and the model used to explain what we observe from active galaxies. Finally, this chapter will go over what is expected from the work that was conducted and the motivation behind it.

1.2 GALAXIES, ACTIVE GALAXIES, AND BLACK

HOLES

Before we can dive into the work that was conducted for this thesis we must first understand the objects that were observed and how they are classified. Galaxies are the easiest of these objects to classify and explain.

1.2.1 GALAXIES

Galaxies are relatively faint objects in comparison to close stars when being observed with the naked eye. So it comes as no surprise that they were not discovered until telescope technology had advanced to a sufficient degree as to allow a significant enough amount of light in to view very distant objects. In 1845, William Parsons of Ireland built a telescope that would be the largest of its time. In using this telescope he was able to observe groupings of stars and gas and dust distant from our own Milky Way. Initially Parsons believed galaxies to be nebulae but soon concluded that they were "great clouds of stars" Seeds and Backman (2013). These great clouds of stars are now theorized by modern astronomers and physicists to contain supermassive black holes at their centers holding them together. Supermassive black holes, as their name suggests and as this chapter will explain later are extremely massive.

GALAXIES CLASSIFICATION

Since their discovery, many various types of galaxies have been discovered and classified. Galaxies can be categorized by the following classifications.

The diagram illustrates the Hubble sequence of galaxies, showing the progression from Ellipticals to Irregulars. The sequence is organized into several categories, each with representative galaxy images and labels:

- Ellipticals:** A horizontal row of galaxies labeled E₀ through E₇, showing a progression from circular to increasingly elongated shapes.
- Lenticular:** Two galaxies labeled S0A and S0B, positioned between the Ellipticals and the main spiral sequence.
- Spirals:** A curved path of galaxies labeled SABa, SABb, SABc, SABd, SAb, SAa, SAc, SAd, and SAm, showing a progression from tightly wound to more loosely wound and eventually to irregular spiral forms.
- Barred Spirals:** A curved path of galaxies labeled SBa, SBb, SBc, SBd, and SBm, showing a progression from tightly wound to more loosely wound and eventually to irregular spiral forms.
- Irregulars:** Two galaxies labeled Im and dSph, positioned at the end of the sequence.

The labels for the galaxies are color-coded: blue for Ellipticals, brown for Lenticular, orange for Spirals, red for Barred Spirals, and green for Irregulars.

1: ELLIPTICAL GALAXIES

Seeds and Backman (2013) Describes elliptical galaxies as round or elliptical with no visible gas and dust and lacking hot bright stars. These galaxies classified from E0 to E7 based on how elliptical they are.

Spiral galaxies are the most popular of the classifications of galaxies, and also the most pleasing to the eye. They consist of a disk and spiral arms extruding from their center. Spiral galaxies have some extremely unique characteristics, the least of which are their spiral arms from which they derive their name. A sub-class of spiral galaxy known as the *Barred Spiral Galaxy* is called so because of the presence of a uniquely inherited bar at their center from which their spiral arms protrude.

3: IRREGULAR GALAXIES

Not quite belonging to either the spiral nor the elliptical classes of galaxies, *Irregular Galaxies* are collections of gas and dust with no real structure or shape.

1.2.2 ACTIVE GALAXIES

Astronomical description of active galaxies(AGN).

1.2.3 BLACK HOLES

Astronomical description of black holes and super-massive black holes.

1.3 HOW AGN ARE OBSERVED

1.3.1 ORBITAL X-RAY TELESCOPES

A description of the orbital observatories used and the bands of the EM spectrum they can observe

1.3.2 THE UNIFIED MODEL

A description of the unified model and how, if the model holds, we can observe the inner workings.

1.4 EXPECTED ACHIEVEMENTS

What is expected to be achieved from this project

Chapter 2

THE GENERAL RELATIVISTIC DESCRIPTION

2.1 GENERAL RELATIVITY

An introduction to general relativity and its relation to black holes.

2.1.1 AN INTRODUCTION TO TENSORS

An introduction to tensors and their mathematical properties

2.1.2 THE METRIC TENSOR

Role of the metric tensor

2.1.3 THE SPACETIME INTERVAL

Role of the spacetime interval

2.2 THE KERR SPINNING BLACK HOLE

Description of the concept of the Kerr spinning black hole and its importance.

2.2.1 THE KERR SOLUTION

The mathematics of the Kerr solution.

2.2.2 PROPERTIES OF THE KERR SOLUTION

Properties of the Kerr solution.

Chapter 3

STATISTICAL ANALYSIS

3.1 THE STRUCTURE FUNCTION

(Collier and Peterson, 2001) (author1, year1) (?) What the structure function is and how it applies to this data sample. Why it was chosen.

3.2 THE DATA

An overview of the data. This section will contain many figures.

Chapter 4

PROGRAMMING SFA

4.1 SFA USAGE AND DESCRIPTION

A description of what SFA was developed for and its usage.

4.2 THE SFA ALGORITHM

4.2.1 PROCESS

The logical process SFA follows to generate the structure function

4.2.2 COST AND COMPLEXITY

An overview of the general running cost and Big-Oh order of SFA

4.3 EXPECTED OUTPUT

What one should expect for results from running SFA

Chapter 5

RESULTS AND CONCLUSIONS

5.1 RESULTS

A review of the results of running SFA on observational data

5.2 CONCLUSIONS

Theoretical conclusions based on the findings of SFA

Chapter 6

FUTURE WORK

Potential future work and applications of SFA

Appendix A

SFA

If you wanted an appendix, it would go in like this. It would be referenced using Appendix A.

Bibliography

author1. title1. *journal1*, volume1:pages1, year1.

S. Collier and B. M. Peterson. Characteristic ultraviolet/optical timescales in active galactic nuclei. *The Astrophysical Journal*, 555:775, 2001.

Michael A. Seeds and Dana E. Backman. *Stars and Galaxies*. Brooks/Cole, 8th edition, 2013.