

A healthy diet is built on six essential nutrients — **protein, carbohydrates, fats, vitamins, minerals, and water** — each serving critical functions from energy production to tissue repair. The key to optimal health lies not just in consuming these nutrients, but in choosing the **right types**: complex carbohydrates over refined sugars, unsaturated fats over saturated and trans fats, and fiber-rich whole foods over processed options. By following evidence-based guidelines like filling **half your plate with vegetables and fruits, one-quarter with lean protein, and one-quarter with whole grains**, while limiting sodium, added sugars, and empty calories, you can significantly reduce your risk of chronic diseases including heart disease, type 2 diabetes, and certain cancers. ([WHO](#))

Healthy Nutrition: A Comprehensive Guide to Eating Well

1. Understanding the Six Essential Nutrients

The human body requires a diverse array of nutrients to function optimally, sustain energy, repair tissues, and defend against disease. According to the National Institutes of Health (NIH), there are **six major classes of nutrients** essential for human health: carbohydrates, lipids (fats), proteins, vitamins, minerals, and water. ([National Center for Biotechnology Information](#)) These nutrients work synergistically to support every physiological process, from the beating of your heart to the firing of neurons in your brain. Understanding what each nutrient does, where to find it, and how much you need is the foundation of making informed dietary choices that promote long-term health and vitality.

Carbohydrates, proteins, and fats are classified as **macronutrients** because the body requires them in relatively large amounts to provide energy and support structural functions. Vitamins and minerals are **micronutrients**, needed in smaller quantities but equally critical for metabolic processes, immune function, and cellular health. Water, though often overlooked, is perhaps the most essential nutrient of all, serving as the medium for virtually every biochemical reaction in the body and comprising approximately **60% of adult body weight**. ([CDC](#)) Each of these nutrient categories contains both beneficial and harmful subtypes, and learning to distinguish between them is key to constructing a diet that nourishes rather than harms your body.

1.1 What the Body Uses Nutrients For

The six essential nutrients serve four primary physiological purposes that keep the body functioning at its best. These fundamental roles — **energy production, growth and development, tissue repair, and regulation of body processes** — explain why a balanced diet is not merely about calorie counting but about ensuring that every cell in your body receives the specific building blocks it needs to perform its designated function. When any of these nutritional needs are compromised, the body begins to show signs of dysfunction, ranging from fatigue and poor concentration to serious chronic diseases.

Nutrient Function	Description	Key Nutrients Involved
Energy	Powers all physical activity, brain function, and cellular processes	Carbohydrates, fats, proteins (Cleveland Clinic)
Growth	Builds new tissues, especially during childhood, adolescence, and pregnancy	Protein, calcium, vitamin D, vitamin A (Cleveland Clinic)

Table 1 – continued

Nutrient Function	Description	Key Nutrients Involved
Repair	Heals wounds, rebuilds damaged tissues, and maintains muscle mass	Protein, vitamin C, zinc, vitamin K (Cleveland Clinic)
Regulation	Controls body temperature, hormone production, enzyme activity, and immune response	Vitamins, minerals, water, proteins (National Center for Biotechnology Information)

Table 1: The four primary functions of nutrients in the human body and the key nutrients responsible for each function.

1.1.1 Energy Production Energy is the most fundamental requirement for life, and the body obtains it primarily through the metabolism of **carbohydrates, fats, and proteins**. Carbohydrates are the body’s preferred and most efficient source of energy, particularly for the brain and central nervous system, which rely almost exclusively on glucose — the simple sugar derived from carbohydrate digestion. ([Cleveland Clinic](#)) When carbohydrates are consumed, the digestive system breaks them down into glucose, which enters the bloodstream and is transported to cells throughout the body. The hormone insulin facilitates the uptake of glucose into cells, where it undergoes cellular respiration to produce **adenosine triphosphate (ATP)**, the body’s primary energy currency. Excess glucose is stored in the liver and muscles as glycogen for short-term energy needs, and any remaining surplus is converted to fat for long-term storage.

Fats serve as the body’s **most concentrated source of energy**, providing approximately **9 calories per gram** compared to 4 calories per gram from carbohydrates and proteins. ([Harvard Health](#)) While the body preferentially uses carbohydrates for immediate energy needs, fats become increasingly important during prolonged physical activity, fasting, or when carbohydrate intake is insufficient. Dietary fats are broken down into fatty acids, which can enter energy-producing pathways to generate ATP. Beyond their role in energy production, fats are essential for the absorption of fat-soluble vitamins (A, D, E, and K), the formation of cell membranes, and the production of important signaling molecules such as hormones and prostaglandins. Proteins can also be used for energy when carbohydrate and fat stores are depleted, but this is inefficient and can lead to the breakdown of muscle tissue.

1.1.2 Growth and Development Growth and development are nutrient-intensive processes that require a steady supply of specific building blocks, particularly during periods of rapid change such as **childhood, adolescence, pregnancy, and recovery from illness or injury**. Protein is the cornerstone of growth, providing the amino acids necessary to build new tissues, muscles, organs, and enzymes. ([Cleveland Clinic](#)) The body requires **20 different amino acids** to construct the vast array of proteins it needs, nine of which are considered “essential” because the body cannot synthesize them and must obtain them from dietary sources. Foods rich in high-quality protein — such as lean meats, poultry, fish, eggs, dairy products, beans, and nuts — are therefore critical during growth phases.

Beyond protein, several micronutrients play pivotal roles in growth and development. **Calcium and vitamin D** work in partnership to build and maintain strong bones and teeth, with calcium providing the mineral matrix and vitamin D facilitating its absorption from the digestive tract. ([National Center for Biotechnology Information](#)) Vitamin A is essential for cellular differentiation and gene expression,

processes that guide cells to develop into specialized tissues and organs. The B-complex vitamins, particularly **folate (vitamin B9) and vitamin B12**, are crucial for DNA synthesis and cell division, which is why adequate folate intake before and during pregnancy is strongly recommended to prevent neural tube defects in developing fetuses. ([National Center for Biotechnology Information](#)) Without adequate intake of these growth-promoting nutrients, children may experience stunted growth, delayed development, and increased susceptibility to infections.

1.1.3 Tissue Repair The body's ability to repair damaged tissues — whether from a minor cut, a surgical incision, or the microscopic wear and tear of daily life — depends heavily on adequate nutrition. **Protein** is once again the star player in tissue repair, as it provides the raw materials needed to rebuild damaged cells and synthesize collagen, the structural protein that forms the scaffold of skin, blood vessels, and connective tissues. ([Cleveland Clinic](#)) When tissue is damaged, the body enters a state of increased metabolic demand, requiring additional protein and energy to support the healing process. Chronic wounds or slow-healing injuries are often signs of protein deficiency or malnutrition.

Vitamin C is another critical nutrient for tissue repair, as it serves as an essential cofactor in the enzymatic reactions that produce collagen. Without sufficient vitamin C, collagen synthesis is impaired, leading to weakened blood vessels, poor wound healing, and the development of scurvy in severe cases. ([National Center for Biotechnology Information](#)) **Zinc** supports tissue repair by facilitating cell division, protein synthesis, and immune function at the wound site. **Vitamin K** plays a vital role in blood clotting, preventing excessive bleeding when tissues are damaged and allowing the healing process to begin. ([National Center for Biotechnology Information](#)) Together, these nutrients ensure that the body can efficiently repair itself and recover from injuries, infections, and surgical procedures.

1.1.4 Regulation of Body Processes The regulation of body processes encompasses a vast array of physiological functions that maintain **homeostasis** — the stable internal environment necessary for life. Nutrients play essential roles as **cofactors for enzymes, components of hormones, regulators of gene expression, and modulators of immune responses**. Vitamins and minerals are particularly important in this regulatory role. The B-complex vitamins function as coenzymes in hundreds of metabolic reactions involved in energy production, neurotransmitter synthesis, and DNA repair. ([National Center for Biotechnology Information](#)) Vitamin D acts as a hormone, regulating calcium absorption, bone mineralization, and immune function through its interactions with vitamin D receptors throughout the body.

Minerals such as **sodium, potassium, and chloride** serve as electrolytes, maintaining fluid balance, nerve transmission, and muscle contraction. ([National Center for Biotechnology Information](#)) **Magnesium** participates in over **300 enzymatic reactions**, including those involved in glucose metabolism, protein synthesis, and blood pressure regulation. **Iron** is essential for oxygen transport in the blood and for the function of enzymes involved in energy production. Proteins also serve regulatory functions as hormones (insulin, glucagon), enzymes (digestive enzymes, metabolic enzymes), antibodies (immune defense), and transport molecules (hemoglobin, albumin). Without adequate intake of these regulatory nutrients, the body's ability to maintain homeostasis is compromised, leading to metabolic dysfunction, impaired immunity, and increased disease risk.

1.2 Macronutrients vs. Micronutrients

Understanding the distinction between macronutrients and micronutrients is essential for planning a balanced diet that meets all of the body's nutritional needs. **Macronutrients** — carbohydrates, proteins, and fats — are required in gram quantities daily and serve as the primary sources of energy and structural

components for the body. [\(National Center for Biotechnology Information\)](#) **Micronutrients** — vitamins and minerals — are needed in much smaller amounts (milligrams or micrograms) but are equally critical for health, serving as cofactors for enzymes, antioxidants, and regulators of physiological processes. Both categories are essential, and deficiencies in either can lead to significant health problems.

Category	Nutrients	Daily Requirement	Primary Functions	Main Food Sources
Macronutrients	Carbohydrates, Proteins, Fats	Grams (g) quantities: Carbs 45-65% of calories; Protein 10-35%; Fat 20-35% (American Heart Association)	Energy, tissue building, hormone production, cell structure	Grains, meats, dairy, oils, legumes (Cleveland Clinic)
Micronutrients	Vitamins (A, B, C, D, E, K) and Minerals (calcium, iron, zinc, etc.)	Milligrams (mg) or micrograms (mcg) quantities (National Center for Biotechnology Information)	Enzyme cofactors, immune function, bone health, antioxidant defense	Fruits, vegetables, dairy, fortified foods (Cleveland Clinic)
Water	H O	~2.7-3.7 liters per day (including food sources) (CDC)	Temperature regulation, nutrient transport, waste removal, joint lubrication	Water, beverages, fruits, vegetables (CDC)

Table 2: Comparison of macronutrients, micronutrients, and water, including their daily requirements, primary functions, and main food sources.

1.2.1 Carbohydrates: The Body’s Main Fuel Source Carbohydrates are the body’s **primary and preferred source of energy**, particularly for the brain, red blood cells, and muscles during moderate to high-intensity exercise. [\(Cleveland Clinic\)](#) When consumed, carbohydrates are broken down into glucose, which circulates in the bloodstream and is taken up by cells to produce ATP through cellular respiration. The brain alone consumes approximately **120 grams of glucose per day**, accounting for about 20% of the body’s total energy expenditure, despite representing only 2% of body weight. This heavy reliance on glucose explains why low-carbohydrate diets can cause symptoms such as headaches, fatigue, and difficulty concentrating — the brain is being deprived of its primary fuel source.

Not all carbohydrates are created equal, however. **Complex carbohydrates** — found in whole grains, legumes, vegetables, and fruits — contain longer chains of sugar molecules that take longer to digest, resulting in a slower, more sustained release of glucose into the bloodstream. [\(harvard.edu\)](#) They also tend to be rich in fiber, vitamins, and minerals. **Simple carbohydrates** — such as table sugar, honey, and the sugars found in candy and sweetened beverages — are composed of one or two sugar molecules that are rapidly absorbed, causing sharp spikes in blood glucose and insulin levels. [\(harvard.edu\)](#) Repeated consumption of simple carbohydrates has been linked to weight gain, insulin resistance, type 2 diabetes, and cardiovascular disease. The **glycemic index (GI)** was developed to rank carbohydrate-containing foods based on how quickly they raise blood sugar levels, with low-GI foods (whole grains, legumes, most fruits) being preferable for sustained energy and metabolic health. [\(harvard.edu\)](#)

1.2.2 Proteins: Building and Repairing the Body Proteins are often described as the “**building blocks of life**” because they are essential for the construction, repair, and maintenance of virtually every tissue and structure in the body. ([National Institute on Aging](#)) Proteins are large molecules composed of chains of **20 different amino acids**, which are linked together in specific sequences determined by genetic code. The body can synthesize 11 of these amino acids, but the remaining **nine are “essential”** and must be obtained from dietary sources. Foods that contain all nine essential amino acids in adequate proportions are called “complete proteins” and include animal products such as meat, poultry, fish, eggs, and dairy, as well as some plant sources like quinoa and soy.

The functions of proteins extend far beyond tissue building. **Enzymes**, which catalyze virtually every chemical reaction in the body, are proteins. **Hormones** such as insulin and glucagon, which regulate blood sugar levels, are proteins or protein-derived. **Antibodies**, which defend the body against pathogens, are proteins. **Hemoglobin**, the molecule that transports oxygen in the blood, is a protein. Even **neurotransmitters** — the chemical messengers of the nervous system — are synthesized from amino acids. The Recommended Dietary Allowance (RDA) for protein is **0.8 grams per kilogram of body weight per day** for adults, but needs may be higher for athletes, older adults, pregnant women, and individuals recovering from illness or injury. ([National Institute on Aging](#))

1.2.3 Fats: Essential for Health and Energy Storage Dietary fats have been unfairly demonized in popular nutrition culture, but they are **essential for human health** and serve numerous critical functions in the body. Fats are the most energy-dense macronutrient, providing **9 calories per gram** — more than twice the energy of carbohydrates or proteins. ([Harvard Health](#)) This makes them an efficient form of energy storage; the body can store virtually unlimited amounts of fat in adipose tissue, which can be mobilized during periods of fasting or increased energy demand. Beyond energy storage, fats are essential components of cell membranes, where they influence membrane fluidity and the function of membrane-bound proteins. They are also necessary for the absorption of fat-soluble vitamins (A, D, E, and K) and the synthesis of steroid hormones, including estrogen, testosterone, and cortisol.

The type of fat consumed is far more important than the total amount. **Unsaturated fats** — including monounsaturated fats (found in olive oil, avocados, and nuts) and polyunsaturated fats (found in fatty fish, flaxseeds, and walnuts) — are considered “healthy fats” because they can improve blood cholesterol levels, reduce inflammation, and lower the risk of cardiovascular disease. ([Harvard Health](#)) **Omega-3 fatty acids**, a type of polyunsaturated fat found in fatty fish, have particularly potent anti-inflammatory effects and are associated with reduced risk of heart disease, improved brain function, and better mental health. ([National Institute on Aging](#)) In contrast, **saturated fats** (found in butter, fatty meats, and full-fat dairy) and **trans fats** (found in partially hydrogenated oils and many processed foods) can raise LDL (“bad”) cholesterol levels and increase the risk of heart disease and stroke. ([American Heart Association](#))

1.2.4 Vitamins and Minerals: Small but Mighty Vitamins and minerals are **micronutrients** that the body needs in relatively small amounts — milligrams or even micrograms per day — yet their importance to health cannot be overstated. ([National Center for Biotechnology Information](#)) Vitamins are organic compounds that serve as coenzymes, antioxidants, and regulators of gene expression. They are classified as either **water-soluble** (the B-complex vitamins and vitamin C) or **fat-soluble** (vitamins A, D, E, and K), a distinction that affects how they are absorbed, transported, stored, and excreted by the body. Water-soluble vitamins are not stored in significant amounts and must be consumed regularly, while fat-soluble vitamins can be stored in the liver and fatty tissues for longer periods.

Minerals are **inorganic elements** that serve structural, regulatory, and catalytic functions in the body. **Macrominerals** — including calcium, phosphorus, magnesium, sodium, potassium, and chloride — are

required in amounts greater than 100 mg per day. ([National Center for Biotechnology Information](#)) Calcium and phosphorus form the mineral matrix of bones and teeth; sodium, potassium, and chloride maintain fluid balance and nerve transmission; and magnesium participates in over 300 enzymatic reactions. **Microminerals** or **trace minerals** — including iron, zinc, copper, selenium, and iodine — are needed in smaller amounts but are no less essential. Iron is critical for oxygen transport; zinc supports immune function and wound healing; iodine is necessary for thyroid hormone production; and selenium functions as an antioxidant and supports thyroid health. Deficiencies in any of these micronutrients can lead to characteristic deficiency diseases and impaired physiological function.

1.2.5 Water: The Most Essential Nutrient Water is frequently overlooked in discussions of nutrition, yet it is **the most essential nutrient for human survival**. The human body is approximately **60% water by weight**, and every system in the body depends on water to function properly. ([CDC](#)) Water serves as the **solvent for biochemical reactions**, the **medium for nutrient transport**, the **regulator of body temperature**, the **lubricant for joints**, and the **medium for waste removal** through urine, sweat, and bowel movements. Without adequate water intake, the body quickly becomes dehydrated, leading to symptoms ranging from thirst and dry mouth to fatigue, dizziness, confusion, and in severe cases, organ failure and death.

The daily water requirements vary depending on age, sex, body size, activity level, and environmental conditions. The **National Academies of Sciences, Engineering, and Medicine** recommend approximately **3.7 liters per day for men** and **2.7 liters per day for women** from all sources, including beverages and food. ([nih.gov](#)) Physical activity, heat exposure, fever, diarrhea, and vomiting all increase water needs. While the “8 glasses a day” rule is a reasonable general guideline, individual needs may vary considerably. Foods with high water content — such as fruits, vegetables, soups, and dairy products — can contribute significantly to daily fluid intake. Replacing sugary beverages with plain water is one of the simplest and most effective dietary changes for improving health, as it reduces calorie intake while ensuring adequate hydration. ([CDC](#))

2. Good Nutrients: What They Do for Your Body

Nutrient-dense foods — those that provide substantial amounts of vitamins, minerals, fiber, and other beneficial compounds relative to their calorie content — are the cornerstone of a healthy diet. These foods support optimal physiological function, protect against chronic disease, promote healthy aging, and enhance quality of life. Understanding which nutrients are beneficial and how they work in the body empowers you to make food choices that actively promote health rather than merely prevent deficiency.



Figure 1: A healthy balanced meal plate with vegetables, lean protein, and whole grains

Figure 1: A healthy balanced meal following the plate method — half vegetables and fruits, one-quarter lean protein, and one-quarter whole grains, with water as the beverage of choice.

2.1 Fiber: The Digestive Champion

Dietary fiber is a type of carbohydrate that the **body cannot digest or absorb**, yet it plays a remarkably diverse and important role in human health. [\(Mayo Clinic\)](#) Unlike other carbohydrates that are broken down into sugars and absorbed, fiber passes relatively intact through the stomach, small intestine, and colon, exerting beneficial effects along the entire length of the digestive tract. Fiber is found exclusively in plant-based foods, including fruits, vegetables, whole grains, legumes, nuts, and seeds. Despite its well-documented health benefits, fiber remains one of the most underconsumed nutrients in the modern diet, with the **average American consuming only about 15 grams per day** — roughly half the recommended amount. [\(Mayo Clinic\)](#)

There are two main types of dietary fiber, each with distinct properties and health benefits. **Soluble fiber** dissolves in water to form a gel-like substance in the digestive tract. This gel slows gastric emptying and the absorption of glucose and cholesterol, which helps regulate blood sugar levels and lower blood cholesterol. [\(Mayo Clinic\)](#) Good sources of soluble fiber include oats, peas, beans, apples, citrus fruits, carrots, and barley. **Insoluble fiber** does not dissolve in water and adds bulk to stool, promoting regular bowel movements and preventing constipation. [\(Mayo Clinic\)](#) It is found in whole wheat flour, wheat bran, nuts, beans, and vegetables such as cauliflower and green beans. Most plant foods contain both types of fiber, and consuming a wide variety of fiber-rich foods ensures that you receive the full spectrum of benefits.

2.1.1 Health Benefits of Fiber The health benefits of dietary fiber extend far beyond its well-known effect on bowel regularity. A high-fiber diet has been associated with a **reduced risk of numerous chronic diseases**, including heart disease, type 2 diabetes, certain cancers, and obesity. [\(Mayo Clinic\)](#) The mechanisms underlying these protective effects are diverse and include the modulation of gut mi-

crobiota, improvement of glycemic control, reduction of blood cholesterol levels, enhancement of satiety, and reduction of systemic inflammation.

Health Benefit	Mechanism	Key Evidence
Heart disease prevention	Soluble fiber binds to cholesterol in the gut, preventing its absorption and lowering LDL levels (Mayo Clinic)	A meta-analysis of 67 controlled trials found significant reduction in total and LDL cholesterol with increased soluble fiber intake (harvard.edu)
Type 2 diabetes prevention	Fiber slows glucose absorption, reducing blood sugar spikes and improving insulin sensitivity (nih.gov)	High cereal fiber intake is associated with a 20-30% lower risk of developing T2DM (harvard.edu)
Weight management	High-fiber foods increase satiety and reduce overall calorie intake (Mayo Clinic)	The POUNDS Lost study found that fiber intake predicted weight loss and dietary adherence (nih.gov)
Colorectal cancer prevention	Fiber increases stool bulk and reduces transit time, minimizing exposure to carcinogens (Mayo Clinic)	Meta-analyses show a significant association between high-fiber diets and lower colorectal cancer risk (harvard.edu)
Gut health	Fermentable fiber feeds beneficial gut bacteria, producing short-chain fatty acids that nourish colon cells (nih.gov)	Studies show that increased fiber intake improves gut microbial diversity and reduces inflammation (nih.gov)

Table 3: The major health benefits of dietary fiber, the mechanisms by which they occur, and key supporting evidence.

One of the most significant benefits of dietary fiber is its role in **cardiovascular health**. Soluble fiber, in particular, has been shown to lower blood levels of **low-density lipoprotein (LDL) cholesterol** — the “bad” cholesterol that contributes to the formation of atherosclerotic plaques in arteries. (Mayo Clinic) The mechanism involves the binding of bile acids (which contain cholesterol) to soluble fiber in the intestine, preventing their reabsorption and promoting their excretion in feces. To replace the lost bile acids, the liver draws cholesterol from the blood, thereby reducing circulating LDL levels. A meta-analysis of 67 controlled trials found that increased soluble fiber intake was associated with modest but significant reductions in both total and LDL cholesterol. (harvard.edu) Epidemiological studies have consistently found that higher fiber intake is associated with a lower risk of heart disease and cardiovascular mortality.

Fiber also plays a critical role in **glycemic control and the prevention of type 2 diabetes**. Because fiber slows the digestion and absorption of carbohydrates, it helps prevent the rapid spikes in blood glucose that occur after consuming high-glycemic foods. (nih.gov) This is particularly beneficial for individuals with diabetes, as it helps stabilize blood sugar levels and may reduce the need for medication. Large cohort studies have found that diets high in cereal fiber and whole grains are associated with a **20-30% lower risk of developing type 2 diabetes**, independent of other dietary and lifestyle factors. (harvard.edu) The benefits appear to be strongest for insoluble cereal fiber, suggesting that the type and source of fiber matter as much as the total amount consumed.

2.1.2 Best Sources of Fiber Meeting the recommended daily fiber intake of **25 grams for women and 38 grams for men** requires a deliberate focus on whole, plant-based foods. (eatright.org - [Academy of Nutrition and Dietetics](#)) Refined and processed foods are generally low in fiber because the refining process removes the bran and germ from grains, which are the parts richest in fiber and other nutrients. To increase fiber intake, focus on incorporating a variety of the following foods into your daily diet:

- **Whole grains:** Brown rice, oatmeal, quinoa, whole wheat bread and pasta, barley, bulgur, and bran cereals are excellent sources of fiber. When choosing grain products, look for “whole grain” or “whole wheat” as the first ingredient on the label. ([Mayo Clinic](#))
- **Legumes:** Beans, lentils, chickpeas, and split peas are among the most fiber-rich foods available, providing **10-15 grams of fiber per cup** of cooked beans. They are also excellent sources of plant-based protein. ([Mayo Clinic](#))
- **Fruits:** Berries, apples, pears, oranges, and bananas are all good sources of fiber. Eating the skin of fruits (when edible) increases fiber intake further. A medium apple with skin provides about **4.4 grams of fiber**. ([Mayo Clinic](#))
- **Vegetables:** Broccoli, Brussels sprouts, carrots, sweet potatoes, and leafy greens are fiber-rich and also provide a wealth of vitamins, minerals, and antioxidants. ([Mayo Clinic](#))
- **Nuts and seeds:** Almonds, walnuts, chia seeds, and flaxseeds provide fiber along with healthy fats and protein. A quarter-cup of almonds contains about **4.5 grams of fiber**. ([Mayo Clinic](#))

When increasing fiber intake, it is important to do so **gradually** over several weeks and to drink **plenty of water**. A sudden increase in fiber can cause gas, bloating, and cramping as the gut microbiota adjust to the increased substrate. Adding fiber slowly allows the digestive system to adapt and minimizes discomfort.

2.2 Lean Proteins: Building Blocks for Life

Protein is an essential macronutrient that serves as the **primary building material for the body’s tissues and structures**. Every cell in the body contains protein, and it is needed for the growth, repair, and maintenance of muscles, bones, skin, hair, nails, and internal organs. ([National Institute on Aging](#)) Proteins also function as enzymes, hormones, antibodies, and transport molecules, making them indispensable for virtually every physiological process. The quality and source of dietary protein matter significantly for health, with lean and plant-based protein sources being preferable to high-fat animal products.

The **protein food group** encompasses a wide variety of foods, including meat, poultry, seafood, eggs, beans, peas, lentils, nuts, seeds, and soy products. ([National Institute on Aging](#)) The Dietary Guidelines for Americans recommend consuming a variety of protein foods from both animal and plant sources to ensure adequate intake of all essential amino acids and to obtain the diverse array of vitamins, minerals, and other beneficial compounds found in different protein-rich foods. For example, fatty fish like salmon and tuna provide not only high-quality protein but also **omega-3 fatty acids**, which have potent anti-inflammatory and heart-protective effects.

Protein Source	Serving Size	Protein (g)	Key Benefits	Notes
Chicken breast (skinless)	3 oz (85g)	26g	Lean, versatile, high in B vitamins	Choose skinless to reduce saturated fat (National Institute on Aging)

Table 4 – continued

Protein Source	Serving Size	Protein (g)	Key Benefits	Notes
Salmon	3 oz (85g)	22g	Rich in omega-3 fatty acids, heart-healthy	Aim for 2 servings of fatty fish per week (National Institute on Aging)
Lentils (cooked)	1 cup	18g	High in fiber and iron, plant-based, economical	Excellent meat alternative (NHS)
Greek yogurt (non-fat)	1 cup	20g	Probiotics for gut health, high in calcium	Choose plain, unsweetened varieties (NHS)
Eggs	2 large	12g	Complete protein, versatile, affordable	The yolk contains vitamins and minerals (NHS)
Almonds	1/4 cup	8g	Healthy fats, fiber, vitamin E	Calorie-dense; watch portion sizes (NHS)
Tofu	1/2 cup	10g	Plant-based, complete protein, cholesterol-free	Good source of calcium and iron (National Institute on Aging)
Black beans (cooked)	1 cup	15g	High in fiber, folate, and potassium	Versatile in soups, salads, and burritos (NHS)

Table 4: Common protein sources with their protein content, key benefits, and practical notes for healthy eating.

2.2.1 Animal-Based Proteins Animal-based proteins are considered “**complete proteins**” because they contain all nine essential amino acids in proportions that closely match human needs. ([National Institute on Aging](#)) Lean meats such as beef, pork, and lamb provide high-quality protein along with important nutrients such as iron, zinc, and vitamin B12. However, they can also be significant sources of saturated fat, which is why choosing lean cuts and trimming visible fat is recommended. Poultry, particularly **skinless chicken and turkey breast**, is an excellent lean protein source, providing high-quality protein with relatively low fat content.

Seafood is arguably one of the **healthiest protein choices** available. Fish and shellfish are excellent sources of high-quality protein and provide a unique array of nutrients not found in other protein sources. **Fatty fish** such as salmon, mackerel, sardines, trout, and herring are particularly rich in **omega-3 fatty acids (EPA and DHA)**, which have been shown to reduce inflammation, lower triglyceride levels, slow the development of atherosclerotic plaques, and reduce the risk of fatal arrhythmias. ([National Institute on Aging](#)) The American Heart Association recommends eating **at least two servings of fish per week**, particularly fatty fish, to obtain these cardiovascular benefits. Shellfish such as clams, crabs, and lobster are low in fat and provide protein along with minerals such as zinc and selenium.

Eggs are one of the **most nutrient-dense foods** available, providing high-quality complete protein along with vitamins A, D, E, B12, riboflavin, folate, and minerals such as iron and selenium. Although

eggs do contain cholesterol (about 186 mg per large egg), research has shown that dietary cholesterol has a relatively small impact on blood cholesterol levels for most people, and the **American Heart Association now recommends one whole egg per day** as part of a healthy diet. Dairy products such as milk, yogurt, and cheese are also excellent sources of complete protein and provide calcium, vitamin D, and potassium. Choosing low-fat or fat-free dairy products helps reduce saturated fat intake while maintaining protein and calcium content.

2.2.2 Plant-Based Proteins Plant-based proteins offer significant health advantages, including **lower saturated fat content, zero cholesterol, and higher fiber content** compared to animal proteins. ([National Institute on Aging](#)) While most individual plant proteins are “incomplete” (lacking one or more essential amino acids), consuming a variety of plant protein sources throughout the day ensures that all essential amino acid needs are met. Beans, peas, and lentils (collectively known as legumes or pulses) are nutritional powerhouses, providing not only protein but also significant amounts of fiber, iron, folate, potassium, and magnesium. A cup of cooked lentils provides approximately **18 grams of protein and 15 grams of fiber** — a combination that supports both muscle health and digestive health.

Nuts and seeds are concentrated sources of **plant protein, healthy unsaturated fats, fiber, and vitamin E**. ([NHS](#)) Almonds, walnuts, pistachios, chia seeds, flaxseeds, and hemp seeds can be easily incorporated into meals and snacks to boost protein and nutrient intake. Nut butters (without added sugars or oils) are another convenient option. Soy products such as tofu, tempeh, and edamame are unique among plant foods because they contain **all nine essential amino acids**, making them complete proteins comparable to animal sources. Tofu is also an excellent source of calcium and iron, particularly when calcium-set. Processed soy products such as veggie burgers and soy milk can be healthy alternatives to meat and dairy, but it is important to choose products that are low in sodium and added sugars.

2.3 Complex Carbohydrates: Sustained Energy

Complex carbohydrates are the **foundation of a healthy diet**, providing the body with a steady, sustained source of energy along with a wealth of vitamins, minerals, fiber, and phytonutrients. ([Cleveland Clinic](#)) Unlike simple carbohydrates, which cause rapid spikes and crashes in blood sugar, complex carbohydrates are digested and absorbed more slowly, providing a more stable and prolonged release of glucose into the bloodstream. This steady energy supply supports sustained physical and mental performance, reduces hunger and cravings, and promotes better metabolic health.

The key distinction between healthy and unhealthy carbohydrates lies in their **degree of processing**. Whole, unprocessed carbohydrate sources — such as whole grains, legumes, vegetables, and fruits — retain their natural fiber, vitamins, minerals, and phytonutrients. ([American Heart Association](#)) Refined carbohydrate sources — such as white bread, white rice, pastries, and sugary cereals — have been processed to remove the bran and germ, stripping away most of their nutritional value and leaving behind primarily starch and calories. The refining process also increases the glycemic index of these foods, causing more rapid blood sugar elevation.

Carbohydrate Type	Examples	Glycemic Effect	Nutrient Content	Health Impact
Complex carbs (whole grains)	Brown rice, oatmeal, quinoa, whole wheat bread (Cleveland Clinic)	Slow, steady glucose release	High in fiber, B vitamins, minerals	Supports sustained energy, weight management, and metabolic health (American Heart Association)
Complex carbs (starchy vegetables)	Sweet potatoes, corn, peas, butternut squash (Cleveland Clinic)	Moderate glucose release	High in fiber, vitamins A and C, potassium	Nutrient-dense energy source (CDC)
Complex carbs (legumes)	Beans, lentils, chickpeas (Cleveland Clinic)	Very slow glucose release	High in protein, fiber, iron, folate	Best for blood sugar control and satiety (CDC)
Simple carbs (natural)	Fruits, milk, dairy (American Heart Association)	Moderate glucose release	Contains vitamins, minerals, fiber (in fruit)	Healthy in moderation as part of balanced diet (American Heart Association)
Simple carbs (added sugars)	Soda, candy, cookies, cakes (American Heart Association)	Rapid glucose spike	Empty calories, no nutritional value	Linked to obesity, diabetes, heart disease (Harvard Health)
Refined grains	White bread, white rice, pastries (American Heart Association)	Rapid glucose spike	Low in fiber and nutrients (unless enriched)	Associated with weight gain and metabolic syndrome (CDC)

Table 5: Comparison of different carbohydrate types, their glycemic effects, nutrient content, and health impacts.

2.3.1 Whole Grains: The Powerhouse Carbs Whole grains are grains that retain all three parts of the kernel — the **bran, germ, and endosperm** — in their original proportions. ([National Institute on Aging](#)) The bran is the outer layer and is rich in fiber, B vitamins, and antioxidants. The germ is the embryo of the seed and contains healthy fats, vitamin E, and phytochemicals. The endosperm is the starchy middle layer that provides energy. When grains are refined, the bran and germ are removed, leaving only the endosperm. This process gives refined grains a finer texture and longer shelf life but strips away approximately **25% of the protein and 17 key nutrients**, including fiber, iron, and B vitamins.

The health benefits of whole grains are well-documented and extensive. Regular consumption of whole grains has been associated with a **reduced risk of heart disease, type 2 diabetes, obesity, and certain cancers**. ([harvard.edu](#)) The fiber in whole grains helps lower cholesterol, regulate blood sugar, and promote healthy digestion. The vitamins, minerals, and phytochemicals in whole grains provide antioxidant protection and support overall metabolic health. Examples of whole grains include **whole**

wheat, oats, brown rice, quinoa, barley, bulgur, millet, buckwheat, and cornmeal. [\(National Institute on Aging\)](#) When shopping for grain products, look for the word “whole” as the first ingredient on the label (e.g., “whole wheat flour” rather than “enriched wheat flour”).

2.3.2 Fruits and Vegetables: Nature’s Vitamin Packages Fruits and vegetables are among the **most nutrient-dense foods** available, providing a wide array of vitamins, minerals, fiber, and phytonutrients with relatively few calories. [\(WHO\)](#) The World Health Organization (WHO) recommends eating at least **400 grams (about 5 servings) of fruits and vegetables per day** to reduce the risk of noncommunicable diseases. [\(WHO\)](#) People with diets rich in fruits and vegetables have a significantly lower risk of obesity, heart disease, stroke, diabetes, and certain types of cancer.

Different colored fruits and vegetables contain different **phytonutrients** — beneficial plant compounds that give these foods their vibrant colors and provide antioxidant, anti-inflammatory, and other health-promoting effects. For example, orange and yellow fruits and vegetables (carrots, sweet potatoes, oranges) are rich in **beta-carotene**, which the body converts to vitamin A. Dark green leafy vegetables (spinach, kale, collard greens) are excellent sources of **vitamins A, C, and K, folate, iron, and calcium**. Red and purple fruits and vegetables (tomatoes, berries, grapes) contain **lycopene and anthocyanins**, powerful antioxidants that protect against cellular damage. Eating a wide variety of colorful fruits and vegetables ensures that you receive the full spectrum of these beneficial compounds.

2.4 Healthy Fats: Unsaturated Fats and Omega-3s

The era of “fat-free” dieting has given way to a more nuanced understanding of dietary fats. Research has clearly established that the **type of fat consumed is far more important than the total amount**. [\(Harvard Health\)](#) While trans fats and excessive saturated fats are harmful to health, unsaturated fats — particularly monounsaturated and polyunsaturated fats — are beneficial and should be actively included in a healthy diet. These healthy fats support cardiovascular health, reduce inflammation, improve cholesterol profiles, and may even protect against cognitive decline.

Monounsaturated fats are found in high concentrations in olive oil, peanut oil, canola oil, avocados, and most nuts. [\(harvard.edu\)](#) The health benefits of monounsaturated fats were first highlighted by the **Seven Countries Study** in the 1960s, which found that people in Mediterranean countries had low rates of heart disease despite consuming a relatively high-fat diet, because their primary fat source was olive oil. [\(Harvard Health\)](#) **Polyunsaturated fats**, which include omega-3 and omega-6 fatty acids, are found in fatty fish, walnuts, flaxseeds, sunflower oil, and corn oil. [\(harvard.edu\)](#) Because the body cannot synthesize omega-3 and omega-6 fatty acids, they are considered “essential” and must be obtained from dietary sources.

Fat Type	Food Sources	Effect on LDL	Effect on HDL	Health Impact	Recommendation
Monounsaturated	Olive oil, avocados, almonds, peanuts (harvard.edu)	Lowers LDL	Maintains or raises HDL	Reduces heart disease risk, anti-inflammatory	Use as primary cooking oil and fat source (Harvard Health)

Table 6 – continued

Fat Type	Food Sources	Effect on LDL	Effect on HDL	Health Impact	Recommendation
Polyunsaturated (Omega-3)	Salmon, mackerel, sardines, walnuts, flaxseeds (harvard.edu)	Lowers LDL	Raises HDL	Reduces inflammation, supports brain health, lowers triglycerides	Eat fatty fish 2x/week; include plant sources daily (National Institute on Aging)
Polyunsaturated (Omega-6)	Corn oil, soybean oil, sunflower seeds (harvard.edu)	Lowers LDL when replacing saturated fat	Neutral	Essential for health, but balance with omega-3 is important	Use in moderation; maintain omega-6 to omega-3 ratio (Harvard Health)
Saturated	Butter, fatty meat, cheese, coconut oil (American Heart Association)	Raises LDL	Neutral	Increases cardiovascular disease risk	Limit to <6% of total calories (American Heart Association)
Trans fats	Partially hydrogenated oils, fried foods, baked goods (harvard.edu)	Raises LDL	Lowers HDL	Strongly increases heart disease risk	Avoid completely (harvard.edu)

Table 6: Comparison of different types of dietary fats, their effects on cholesterol, health impacts, and dietary recommendations.

2.4.1 Omega-3 Fatty Acids: Heart and Brain Protectors Omega-3 fatty acids are a special class of polyunsaturated fats that have received considerable attention for their potent **anti-inflammatory and cardioprotective effects**. ([National Institute on Aging](#)) The three main types of omega-3 fatty acids are **eicosapentaenoic acid (EPA)**, **docosahexaenoic acid (DHA)**, and **alpha-linolenic acid (ALA)**. EPA and DHA are found primarily in fatty fish and are the most biologically active forms. ALA is found in plant sources such as flaxseeds, chia seeds, walnuts, and canola oil and must be converted to EPA and DHA in the body — a process that is relatively inefficient.

The cardiovascular benefits of omega-3 fatty acids are among the best-documented in nutritional science. EPA and DHA have been shown to **lower triglyceride levels, reduce blood pressure, decrease platelet aggregation (reducing clot risk), reduce inflammation, and stabilize heart rhythm**. ([National Institute on Aging](#)) Higher blood levels of omega-3 fats are associated with a lower risk of premature death from all causes among older adults. ([harvard.edu](#)) Beyond heart health, omega-3 fatty acids are essential for **brain development and function**. DHA is a major structural component of the brain and retina, and adequate intake during pregnancy and early childhood is critical for cognitive and visual development. Some research also suggests that omega-3 fatty acids may have protective effects against depression, anxiety, and age-related cognitive decline.

2.5 Vitamins and Minerals: Essential Micronutrients

Vitamins and minerals are **micronutrients** that the body needs in small amounts to function properly, but their importance to health is immense. [\(Cleveland Clinic\)](#) These nutrients serve as cofactors for enzymes, antioxidants, structural components, and regulators of gene expression. Deficiencies in vitamins and minerals can lead to characteristic diseases and impaired physiological function, while adequate intake supports optimal health, disease prevention, and healthy aging. The best way to obtain these essential micronutrients is through a varied diet rich in whole foods, including fruits, vegetables, whole grains, lean proteins, and dairy products.

Vitamin/Mineral	Key Functions	Best Food Sources	RDA (Adults)	Deficiency Risks
Vitamin A	Vision, immune function, cell growth, reproduction (National Center for Biotechnology Information)	Sweet potatoes, carrots, spinach, liver, fortified milk	900 mcg (M) / 700 mcg (F)	Night blindness, weakened immunity (National Center for Biotechnology Information)
Vitamin C	Collagen synthesis, wound healing, antioxidant, immune support (National Center for Biotechnology Information)	Citrus fruits, strawberries, bell peppers, broccoli	90 mg (M) / 75 mg (F)	Scurvy, poor wound healing (National Center for Biotechnology Information)
Vitamin D	Calcium absorption, bone health, immune function (National Center for Biotechnology Information)	Fatty fish, fortified milk, sunlight exposure	15-20 mcg (600-800 IU)	Rickets, osteoporosis, weakened immunity (National Center for Biotechnology Information)
Vitamin E	Antioxidant, cell membrane protection, immune function (National Center for Biotechnology Information)	Nuts, seeds, vegetable oils, spinach	15 mg	Nerve and muscle damage (National Center for Biotechnology Information)
Calcium	Bone and teeth structure, muscle contraction, nerve transmission (National Center for Biotechnology Information)	Dairy, fortified plant milks, leafy greens, tofu	1,000 mg	Osteoporosis, muscle cramps (National Center for Biotechnology Information)

Table 7 – continued

Vitamin/Mineral	Key Functions	Best Food Sources	RDA (Adults)	Deficiency Risks
Iron	Oxygen transport (hemoglobin), energy metabolism (National Center for Biotechnology Information)	Red meat, spinach, lentils, fortified cereals	8 mg (M) / 18 mg (F)	Anemia, fatigue, weakness (National Center for Biotechnology Information)
Potassium	Fluid balance, nerve transmission, muscle contraction, blood pressure regulation (National Center for Biotechnology Information)	Bananas, potatoes, beans, yogurt, leafy greens	2,600-3,400 mg	Hypertension, muscle weakness (National Center for Biotechnology Information)
Magnesium	Enzyme function, energy production, muscle and nerve function, bone health (National Center for Biotechnology Information)	Nuts, seeds, whole grains, leafy greens, legumes	400-420 mg (M) / 310-320 mg (F)	Muscle cramps, irregular heartbeat, fatigue (National Center for Biotechnology Information)

Table 7: Essential vitamins and minerals, their key functions, best food sources, recommended daily allowances, and risks of deficiency.

2.5.1 Vitamin C, D, and B-Complex **Vitamin C (ascorbic acid)** is a water-soluble vitamin that functions as a powerful antioxidant and is essential for the synthesis of collagen, the protein that provides structure to skin, blood vessels, bones, and connective tissue. ([National Center for Biotechnology Information](#)) Vitamin C also enhances iron absorption from plant sources, supports immune function, and helps regenerate other antioxidants such as vitamin E. The RDA for vitamin C is **90 mg per day for men and 75 mg per day for women**, with smokers requiring an additional 35 mg per day due to increased oxidative stress. ([National Center for Biotechnology Information](#)) Excellent sources include citrus fruits, strawberries, kiwi, bell peppers, broccoli, and tomatoes.

Vitamin D is unique among vitamins because it can be synthesized by the body when skin is exposed to sunlight. ([National Center for Biotechnology Information](#)) However, many people do not get adequate sun exposure, particularly during winter months or in northern latitudes, making dietary sources and supplements important. Vitamin D is essential for calcium absorption and bone health, and it also plays roles in immune function, muscle function, and cell growth regulation. The RDA for adults is **15-20 micrograms (600-800 IU) per day**. ([National Center for Biotechnology Information](#)) Food sources include fatty fish, fortified milk and plant milks, egg yolks, and fortified cereals. Deficiency is associated with rickets in children, osteomalacia in adults, and increased risk of falls and fractures in older adults.

The **B-complex vitamins** (thiamin, riboflavin, niacin, pantothenic acid, vitamin B6, biotin, folate,

and vitamin B12) function primarily as **coenzymes in energy metabolism**, helping the body convert carbohydrates, fats, and proteins into usable energy. ([National Center for Biotechnology Information](#)) They also play critical roles in the synthesis of red blood cells, DNA, and neurotransmitters. Folate (vitamin B9) is particularly important for women of childbearing age, as adequate intake before and during pregnancy prevents neural tube defects in developing fetuses. Vitamin B12 is found naturally only in animal products, so vegans and some vegetarians may need fortified foods or supplements. Good sources of B vitamins include whole grains, meat, poultry, fish, eggs, dairy, legumes, leafy greens, and fortified cereals.

2.5.2 Calcium, Iron, and Potassium **Calcium** is the most abundant mineral in the body and is essential for the structure and strength of bones and teeth. ([National Center for Biotechnology Information](#)) Approximately **99% of the body's calcium is stored in the skeleton**, where it provides the mineral matrix that gives bones their hardness and strength. The remaining 1% circulates in the blood and is essential for muscle contraction, nerve transmission, blood clotting, and hormone secretion. The RDA for calcium is **1,000 mg per day for adults**, increasing to 1,200 mg for women over 50 and men over 70. ([National Center for Biotechnology Information](#)) Dairy products are the most concentrated sources, but calcium is also found in fortified plant milks, leafy greens (such as kale and collard greens), tofu (when calcium-set), almonds, and sardines with bones.

Iron is a critical component of **hemoglobin**, the protein in red blood cells that transports oxygen from the lungs to tissues throughout the body. ([National Center for Biotechnology Information](#)) Iron is also a component of myoglobin (which stores oxygen in muscles) and is involved in energy metabolism, immune function, and cognitive development. The RDA for iron is **8 mg per day for men and 18 mg per day for women** (due to menstrual blood loss), increasing to 27 mg during pregnancy. ([National Center for Biotechnology Information](#)) Iron from animal sources (heme iron) is more readily absorbed than iron from plant sources (non-heme iron). Good sources include red meat, poultry, seafood, lentils, beans, spinach, and fortified cereals. Vitamin C enhances the absorption of non-heme iron, so consuming iron-rich plant foods with vitamin C-rich foods (such as adding lemon juice to spinach) can improve iron uptake.

Potassium is an essential mineral and electrolyte that plays a critical role in **maintaining fluid balance, nerve transmission, muscle contraction, and blood pressure regulation**. ([National Center for Biotechnology Information](#)) Adequate potassium intake helps counteract the blood pressure-raising effects of sodium and is associated with a reduced risk of stroke and kidney stones. The RDA for potassium is **2,600-3,400 mg per day** for adults. Despite its importance, many people do not consume enough potassium. Good sources include bananas, potatoes, sweet potatoes, beans, lentils, yogurt, leafy greens, and tomatoes. The DASH (Dietary Approaches to Stop Hypertension) diet, which emphasizes potassium-rich fruits and vegetables along with low-fat dairy and lean proteins, has been shown to effectively lower blood pressure.

2.6 Water: Hydration and Body Function

Water is the **most abundant substance in the human body** and is essential for life itself. Every cell, tissue, and organ in the body depends on water to function properly. Water serves as the **solvent for biochemical reactions**, the **medium for transporting nutrients and oxygen** to cells, the **regulator of body temperature** through sweating and respiration, the **lubricant for joints**, the **shock absorber for the brain and spinal cord**, and the **vehicle for removing waste products** through urine, feces, and sweat. ([CDC](#)) Without adequate water intake, the body cannot perform even its most basic functions, and dehydration can quickly become life-threatening.

The daily water requirements vary based on individual factors including age, sex, body size, activity level, and climate. The National Academies recommend approximately **3.7 liters per day for men** and **2.7 liters per day for women** from all sources, including beverages and food. ([nih.gov](#)) About 20% of daily water intake typically comes from food, with the remaining 80% from beverages. Physical activity, heat exposure, fever, and gastrointestinal illness all increase water needs. Thirst is a reliable indicator of hydration needs for most people, but older adults may have a diminished thirst response and need to be more conscious of their fluid intake.

Function	How Water Helps	Signs of Inadequate Intake
Temperature regulation	Sweating and evaporative cooling maintain safe body temperature during heat and exercise (nih.gov)	Overheating, heat exhaustion, reduced exercise performance
Nutrient transport	Water is the medium that carries nutrients, oxygen, and hormones through the bloodstream to cells (CDC)	Fatigue, poor concentration, reduced physical performance
Waste removal	Kidneys use water to filter waste products from blood and excrete them in urine (CDC)	Dark urine, kidney stones, urinary tract infections
Joint lubrication	Synovial fluid, which cushions and lubricates joints, is primarily water (CDC)	Joint discomfort, reduced mobility
Digestion	Water is essential for saliva production, stomach acid dilution, and bowel regularity (CDC)	Constipation, digestive discomfort
Cognitive function	Even mild dehydration (1-2% body weight) can impair mood, memory, and concentration (nih.gov)	Headaches, confusion, irritability

Table 8: The critical functions of water in the body, how water supports each function, and the signs that indicate inadequate intake.

3. Bad Nutrients: What Harm They Cause

Just as certain nutrients actively promote health, others can cause significant harm when consumed in excess. The modern food environment makes it easy to overconsume these harmful nutrients, as they are prevalent in processed foods, fast food, sugary beverages, and snack items that are heavily marketed and readily available. Understanding which nutrients are harmful, how they affect the body, and where they hide in the food supply is essential for making informed choices that protect your health.



Figure 2: Comparison of healthy whole foods versus unhealthy processed foods

Figure 2: A visual comparison of nutrient-dense whole foods (left) versus processed, empty-calorie foods (right). Choosing foods from the left side more often supports long-term health.

3.1 Saturated Fats: Heart Disease Risk

Saturated fats have long been recognized as a **major contributor to cardiovascular disease**, the leading cause of death worldwide. Saturated fats are typically **solid at room temperature** and are found primarily in animal-based foods such as fatty cuts of meat, poultry with skin, full-fat dairy products (butter, cheese, cream, whole milk), and eggs, as well as in tropical oils such as coconut oil, palm oil, and palm kernel oil. ([American Heart Association](#)) When consumed in excess, saturated fats raise levels of **low-density lipoprotein (LDL) cholesterol** in the blood, which contributes to the formation of atherosclerotic plaques in artery walls. These plaques narrow arteries, reduce blood flow, and can rupture, causing heart attacks and strokes.

The mechanism by which saturated fats raise LDL cholesterol is well-understood. Saturated fats stimulate the liver to produce more LDL particles and reduce the activity of LDL receptors on cell surfaces, which are responsible for removing LDL from the bloodstream. ([American Heart Association](#)) The net effect is an increase in circulating LDL cholesterol, which penetrates artery walls, becomes oxidized, and triggers an inflammatory response that leads to plaque formation. The **American Heart Association recommends limiting saturated fat intake to less than 6% of total daily calories**. ([American Heart Association](#)) For someone consuming 2,000 calories per day, this equates to no more than **13 grams of saturated fat**.

Food	Serving Size	Saturated Fat (g)	% of Daily Limit (on 2,000 cal diet)
Butter	1 tbsp	7g	54% (American Heart Association)

Table 9 – continued

Food	Serving Size	Saturated Fat (g)	% of Daily Limit (on 2,000 cal diet)
Cheddar cheese	1 oz (28g)	6g	46% (American Heart Association)
Ground beef (85% lean)	3 oz cooked	5g	38% (American Heart Association)
Coconut oil	1 tbsp	12g	92% (American Heart Association)
Whole milk	1 cup	5g	38% (American Heart Association)
Bacon	3 slices	4g	31% (American Heart Association)
Chicken thigh (with skin)	3 oz cooked	3g	23% (American Heart Association)
Ice cream	1/2 cup	4g	31% (American Heart Association)

Table 9: Saturated fat content of common foods, showing how quickly daily limits can be exceeded.

3.1.1 How Saturated Fats Affect Cholesterol The relationship between saturated fat intake and blood cholesterol levels is one of the most well-established findings in nutritional science. When saturated fats are consumed, they increase the production of **apolipoprotein B (apoB)**-containing lipoproteins in the liver, including LDL cholesterol. ([American Heart Association](#)) These particles transport cholesterol and other fats through the bloodstream to tissues throughout the body. High levels of LDL cholesterol in the blood increase the likelihood that cholesterol will be deposited in artery walls, initiating the process of atherosclerosis.

Research has consistently shown that reducing saturated fat intake and replacing saturated fats with **polyunsaturated or monounsaturated fats** can significantly lower LDL cholesterol levels and reduce cardiovascular risk. ([Harvard Health](#)) A meta-analysis of randomized controlled trials found that replacing 5% of calories from saturated fats with polyunsaturated fats reduced the risk of coronary heart disease by **10%**. The type of fat used to replace saturated fat matters: replacement with polyunsaturated fats (found in vegetable oils, nuts, and fatty fish) provides the greatest cardiovascular benefit, while replacement with refined carbohydrates does not provide the same protective effect and may even increase cardiovascular risk in some cases.

3.1.2 Foods High in Saturated Fats to Limit Reducing saturated fat intake does not require eliminating all animal products from the diet, but rather making **smarter choices within food categories**. When it comes to meat, choose **lean cuts** such as skinless poultry, loin cuts of pork and beef, and ground meat that is at least 90% lean. Trim visible fat from meat before cooking, and opt for grilling, baking, or broiling rather than frying. For dairy, choose **low-fat or fat-free options** such as skim milk, low-fat yogurt, and reduced-fat cheese. These products provide the same protein and calcium as their full-fat counterparts but with significantly less saturated fat.

Tropical oils such as **coconut oil and palm oil** deserve special mention because they have been marketed as health foods in recent years. While it is true that coconut oil contains a type of saturated fat (medium-

chain triglycerides) that is metabolized differently than other saturated fats, it still raises LDL cholesterol levels. ([American Heart Association](#)) The American Heart Association does not recommend coconut oil as a healthy fat source. For cooking and food preparation, it is better to use oils high in monounsaturated and polyunsaturated fats, such as **olive oil, canola oil, avocado oil, and peanut oil**.

3.2 Trans Fats: The Worst Type of Fat

If there is one nutrient that health experts universally agree should be **avoided completely**, it is trans fat. Trans fats are a type of unsaturated fat that have been chemically modified through a process called **partial hydrogenation**, which converts liquid vegetable oils into solid fats. ([harvard.edu](#)) This process was originally developed to create fats with longer shelf life and desirable textural properties for use in baked goods, fried foods, and snack products. However, it also creates a fat configuration that the human body is not designed to metabolize, with devastating consequences for cardiovascular health.

Trans fats are uniquely harmful because they **raise LDL (“bad”) cholesterol while simultaneously lowering HDL (“good”) cholesterol** — a double whammy that dramatically increases the risk of heart disease. ([harvard.edu](#)) HDL cholesterol helps remove LDL cholesterol from the bloodstream and transport it back to the liver for disposal, so reducing HDL impairs the body’s ability to clear harmful cholesterol. Trans fats also promote inflammation and endothelial dysfunction (impaired function of the cells lining blood vessels), further increasing cardiovascular risk. The World Health Organization has estimated that eliminating trans fats from the global food supply could prevent **more than 500,000 deaths from cardiovascular disease each year**.

Food Category	Common Sources	Why They Contain Trans Fats
Fried fast food	French fries, fried chicken, donuts (harvard.edu)	Partially hydrogenated oils used in frying
Baked goods	Cookies, cakes, pastries, pie crusts (harvard.edu)	Solid fats improve texture and shelf life
Margarine and shortening	Stick margarine, vegetable shortening (harvard.edu)	Hydrogenation creates spreadable solid fat
Snack foods	Microwave popcorn, crackers, chips (harvard.edu)	Hydrogenated oils enhance crispness and stability
Frozen foods	Frozen pizza, frozen pies, pot pies (harvard.edu)	Solid fats improve texture during reheating

Table 10: Common food sources of trans fats and why manufacturers use them.

3.2.1 Why Trans Fats Are Banned in Many Countries The overwhelming evidence of harm from trans fats has led many countries to take regulatory action. The **U.S. Food and Drug Administration (FDA)** determined in 2015 that partially hydrogenated oils (the primary source of artificial trans fats) were no longer “generally recognized as safe” and gave food manufacturers three years to remove them from their products. As of 2021, the use of partially hydrogenated oils in the U.S. food supply has been largely eliminated, though small amounts of naturally occurring trans fats (found in small quantities in meat and dairy from ruminant animals) still exist.

Denmark was the first country to ban trans fats in 2003, and many other countries have since followed suit, including Austria, Iceland, Hungary, Norway, Switzerland, and several others. The World Health

Organization has launched a global initiative called **REPLACE** to eliminate trans fats from the global food supply by 2023. Despite these efforts, trans fats remain a concern in many parts of the world, particularly in low- and middle-income countries where regulatory oversight may be weaker. Consumers should continue to read food labels and avoid products that list “partially hydrogenated oils” in the ingredient list, even if the product claims “0 grams trans fat” (manufacturers can make this claim if the product contains less than 0.5 grams per serving, which can still add up).

3.3 Added Sugars: Empty Calories and Disease

Added sugars are sugars and syrups that are **added to foods and beverages during processing or preparation**, as opposed to the naturally occurring sugars found in whole fruits, vegetables, and dairy products. ([Harvard Health](#)) The major sources of added sugars in the American diet include sugar-sweetened beverages (sodas, fruit drinks, sports drinks, energy drinks), desserts and sweet snacks (cakes, cookies, candies, ice cream), sweetened dairy products (flavored yogurt, chocolate milk), and sweetened breakfast cereals. On average, U.S. adults consume approximately **17 teaspoons (about 71 grams) of added sugar per day** — far exceeding the recommended limits. ([Harvard Health](#))

The health consequences of excessive added sugar consumption are extensive and well-documented. High sugar intake is strongly associated with **weight gain and obesity**, as sugary foods and beverages are calorie-dense but not satiating, leading to excess calorie consumption. ([Harvard Health](#)) Sugar-sweetened beverages are particularly problematic because liquid calories do not trigger the same satiety signals as solid foods, making it easy to consume large amounts without feeling full. Excess sugar consumption also contributes to the development of **type 2 diabetes** by promoting insulin resistance and overworking the pancreas’s insulin-producing beta cells.

3.3.1 Link Between Sugar and Obesity The relationship between sugar consumption and obesity is one of the most extensively studied topics in nutritional science. Sugar provides **4 calories per gram**, and sugary foods and beverages are often consumed in large quantities, contributing significantly to total daily calorie intake. ([nih.gov](#)) However, the metabolic effects of sugar extend beyond its calorie content. Fructose, which makes up approximately half of table sugar (sucrose) and high-fructose corn syrup, is metabolized primarily in the liver. When consumed in excess, fructose overwhelms the liver’s metabolic capacity and is converted to fat through a process called **de novo lipogenesis**. ([Harvard Health](#)) This contributes to the accumulation of fat in the liver (non-alcoholic fatty liver disease) and promotes the release of triglycerides into the bloodstream.

Sugar also affects hormones that regulate hunger and satiety. High sugar intake can lead to **leptin resistance**, a condition in which the brain does not respond properly to signals from the hormone leptin that indicate fullness. ([nih.gov](#)) At the same time, sugar consumption does not suppress **ghrelin** (the “hunger hormone”) as effectively as other macronutrients, leading to persistent feelings of hunger despite adequate calorie intake. These hormonal disruptions create a vicious cycle of overeating and weight gain. Reducing sugar intake — particularly from sugar-sweetened beverages — is one of the most effective dietary strategies for weight management.

3.3.2 Sugar’s Role in Diabetes and Heart Disease The link between added sugar and **type 2 diabetes** is well-established. Regular consumption of high-glycemic foods and beverages causes repeated spikes in blood glucose and insulin levels. Over time, cells can become **resistant to insulin’s effects** (insulin resistance), meaning that more insulin is required to clear glucose from the bloodstream. The pancreas compensates by producing more insulin, but eventually, the beta cells may become exhausted and unable to keep up with demand, resulting in elevated blood glucose levels and the diagnosis of

type 2 diabetes. [\(nih.gov\)](#) Studies have shown that individuals who consume sugar-sweetened beverages regularly have a **26% greater risk of developing type 2 diabetes** compared to those who consume them rarely or never.

The impact of sugar on **cardiovascular health** has become increasingly recognized. A large study published in *BMC Medicine* in 2023 analyzed data from over 110,000 people and found that higher intake of added sugars was associated with **increased risks of heart disease and stroke**. [\(Harvard Health\)](#) The higher the added sugar intake, the greater the cardiovascular risk. Sugar contributes to heart disease through multiple mechanisms: it raises blood pressure, promotes inflammation, increases triglyceride levels, and contributes to weight gain and fatty liver disease — all of which are risk factors for cardiovascular disease. The 2025-2030 Dietary Guidelines for Americans recommend that no meal should contain more than **10 grams (2.5 teaspoons) of added sugars**. [\(Harvard Health\)](#)

3.4 Excess Sodium: Blood Pressure and Stroke Risk

Sodium is an essential mineral that the body needs in small amounts for **fluid balance, nerve transmission, and muscle contraction**. [\(National Center for Biotechnology Information\)](#) However, the average person consumes far more sodium than necessary — approximately **3,400 mg per day** in the United States, which is nearly **50% higher** than the recommended limit of 2,300 mg per day. [\(American Heart Association\)](#) The majority of this excess sodium comes not from the salt shaker at the table, but from **processed and restaurant foods**, where it is used as a preservative, flavor enhancer, and texture modifier. Bread, processed meats, canned soups, cheese, snack foods, and restaurant meals are among the biggest contributors to sodium intake.

The primary health concern associated with excess sodium consumption is its effect on **blood pressure**. Sodium attracts water, and when sodium levels in the bloodstream are high, more water is retained, increasing blood volume and putting extra pressure on blood vessel walls. [\(American Heart Association\)](#) Over time, this sustained elevation in blood pressure (hypertension) damages blood vessels and forces the heart to work harder, increasing the risk of **heart attack, heart failure, stroke, and kidney disease**. Meta-analyses have consistently found a direct association between habitual salt intake and the risk of stroke and cardiovascular disease, with a **5-gram reduction in daily salt intake** associated with a **23% lower risk of stroke** and a **17% lower risk of cardiovascular disease**. [\(nih.gov\)](#)

Food Category	Average Sodium Content	% of Daily Limit (2,300 mg)
Deli ham (3 oz)	1,100 mg	48% (American Heart Association)
Canned chicken noodle soup (1 cup)	940 mg	41% (American Heart Association)
Frozen pizza (1/4 of 12-inch)	720 mg	31% (American Heart Association)
Bread (2 slices)	300 mg	13% (American Heart Association)
Pickles (1 large)	1,600 mg	70% (American Heart Association)
Soy sauce (1 tbsp)	900 mg	39% (American Heart Association)
Potato chips (1 oz bag)	170 mg	7% (American Heart Association)

Table 11 – continued

Food Category	Average Sodium Content	% of Daily Limit (2,300 mg)
Restaurant cheeseburger	1,300 mg	57% (American Heart Association)

Table 11: Sodium content of common foods, illustrating how quickly the daily limit is exceeded.

3.4.1 Hidden Sources of Salt in Processed Foods One of the biggest challenges in reducing sodium intake is that **much of the sodium in the modern diet is hidden** in foods that do not necessarily taste salty. Bread and rolls are the top source of sodium in the American diet, not because they taste salty, but because salt is an essential ingredient in the bread-making process that controls yeast activity and enhances flavor. ([American Heart Association](#)) A single sandwich with two slices of bread, deli meat, cheese, and condiments can easily contain over 1,500 mg of sodium — nearly two-thirds of the daily limit. Other surprising sources of hidden sodium include breakfast cereals, salad dressings, ketchup, pasta sauces, and baked goods.

Reducing sodium intake requires vigilance in reading food labels and making conscious choices about food preparation. The **Nutrition Facts panel** on packaged foods lists sodium content per serving, and choosing products labeled “low sodium” (140 mg or less per serving), “reduced sodium” (at least 25% less than the regular version), or “no salt added” can help lower intake. Cooking more meals at home using fresh ingredients allows you to control the amount of salt added to your food. When dining out, ask for sauces and dressings on the side, choose grilled rather than fried options, and request that your meal be prepared with less salt. Flavoring foods with herbs, spices, lemon juice, and vinegar can enhance taste without adding sodium.

3.4.2 Daily Sodium Limits The **Dietary Guidelines for Americans recommend limiting sodium intake to less than 2,300 mg per day** for adults — the amount found in approximately **one teaspoon of table salt**. ([American Heart Association](#)) For individuals with hypertension, diabetes, chronic kidney disease, or those over age 51, the recommended limit is even lower at **1,500 mg per day**. The American Heart Association goes further, recommending an ideal limit of **1,500 mg per day** for most adults, as this level has been shown to provide the greatest reduction in blood pressure and cardiovascular risk.

Reducing sodium intake by even modest amounts can have significant health benefits. A meta-analysis of randomized controlled trials found that reducing salt intake by **6 grams per day** (about 2,400 mg of sodium) was associated with reductions in blood pressure of **7/4 mmHg in people with hypertension** and **4/2 mmHg in those with normal blood pressure**. ([nih.gov](#)) At the population level, these reductions could predict an average **24% lower rate of stroke** and **18% lower rate of coronary heart disease**. It is estimated that reducing sodium intake by 30% could prevent approximately **40 million deaths worldwide** over a 25-year period. ([American Heart Association](#))

3.5 Empty Calories: Foods with No Nutritional Value

Empty calories are calories from foods and beverages that provide **energy but little to no nutritional value** in terms of vitamins, minerals, fiber, protein, or healthy fats. ([Harvard Health](#)) These foods are typically high in added sugars, unhealthy fats, and refined carbohydrates, while being devoid of the beneficial nutrients that support health. Consuming a diet high in empty calories displaces nutrient-dense foods from the diet, leading to nutritional imbalances even when total calorie intake is adequate. Over

time, a diet dominated by empty calories contributes to weight gain, nutrient deficiencies, chronic disease, and poor overall health.

Common sources of empty calories include **sugar-sweetened beverages** (soda, energy drinks, sweetened teas, fruit drinks), **desserts and sweet snacks** (cakes, cookies, candy, ice cream), **fried fast food** (French fries, fried chicken, burgers), **alcoholic beverages**, and **ultra-processed snack foods** (chips, crackers, cheese puffs). (hers) These foods are designed to be highly palatable and easy to overconsume, often leading to a cycle of craving and overeating. The problem is compounded by the fact that empty-calorie foods are often cheaper, more convenient, and more heavily marketed than healthier alternatives, making them an easy choice for time-pressed or budget-conscious consumers.

Empty Calorie Food	Calories	Key Nutrients Provided	Healthier Alternative
Soda (12 oz)	140 cal	None (hers)	Water, sparkling water, unsweetened tea
Donut (1 medium)	250 cal	Minimal (hers)	Whole grain toast with nut butter and fruit
French fries (medium)	380 cal	Minimal (hers)	Baked sweet potato wedges
Candy bar (1.5 oz)	220 cal	Minimal (hers)	Dark chocolate (70%+ cacao) with nuts
Potato chips (1 oz)	160 cal	Minimal (hers)	Raw vegetables with hummus
Alcoholic cocktail	150-300 cal	None (hers)	Sparkling water with fresh fruit
Sugary cereal (1 cup)	120-150 cal	Minimal (may be fortified) (hers)	Oatmeal with berries and nuts

Table 12: Common empty-calorie foods, their caloric content, and healthier nutrient-dense alternatives.

3.5.1 Examples of Empty Calorie Foods **Sugar-sweetened beverages** are the single largest source of empty calories in the American diet and are strongly linked to weight gain, type 2 diabetes, and cardiovascular disease. (Harvard Health) A single 20-ounce bottle of soda contains approximately **65 grams of added sugar** — more than the recommended daily limit in a single serving. These beverages provide a large number of calories without providing any satiety, meaning that people do not compensate by eating less food at meals. Over time, the excess calories from sugary drinks contribute significantly to weight gain and metabolic dysfunction.

Fried and fast foods are another major source of empty calories. French fries, fried chicken, onion rings, and other deep-fried foods are high in calories, unhealthy fats (including trans fats), and sodium, while providing minimal vitamins, minerals, or fiber. (hers) A large order of French fries from a fast-food restaurant can contain **500-600 calories**, primarily from fat and refined carbohydrates, with almost no beneficial nutrients. Similarly, pastries, cakes, cookies, and donuts are high in refined flour, added sugars, and unhealthy fats, providing a calorie-dense but nutrient-poor eating experience.

3.5.2 How Empty Calories Contribute to Weight Gain Empty calories promote weight gain through several interconnected mechanisms. First, foods high in empty calories are typically **very palatable and easy to overconsume**. They are engineered to be hyper-palatable, combining sugar, fat, and

salt in ways that trigger the brain's reward pathways and override natural satiety signals. Second, these foods are **low in fiber, protein, and water** — the three components that contribute most to feelings of fullness and satiety. ([hers](#)) A meal of nutrient-dense whole foods (such as grilled chicken, brown rice, and steamed vegetables) provides the same number of calories as a fast-food meal but keeps you feeling full for much longer, reducing the likelihood of snacking between meals.

Third, empty-calorie foods often cause **rapid fluctuations in blood sugar and insulin levels**, leading to energy crashes and renewed hunger shortly after eating. ([hers](#)) This blood sugar rollercoaster can trigger cravings for more sugary or starchy foods, creating a vicious cycle of overeating. Finally, because empty-calorie foods displace nutrient-dense foods from the diet, they can lead to **nutritional deficiencies** that impair metabolism, energy production, and overall health. When the body lacks essential vitamins and minerals, metabolic processes become less efficient, potentially contributing to weight gain even when calorie intake appears adequate.

3.6 Refined Carbohydrates: Blood Sugar Spikes

Refined carbohydrates are grains and grain products that have been **processed to remove the bran and germ**, leaving only the starchy endosperm. ([American Heart Association](#)) This refining process strips away most of the fiber, vitamins, minerals, and phytonutrients, leaving behind a product that is calorie-dense but nutrient-poor. Common examples of refined carbohydrates include **white flour, white bread, white rice, white pasta, pastries, sugary cereals, and most crackers and snack foods**. ([American Heart Association](#)) While some refined grains are “enriched” with a few added B vitamins and iron, they still lack the full spectrum of nutrients found in whole grains, and the added nutrients do not compensate for the loss of fiber and other beneficial compounds.

The primary health concern with refined carbohydrates is their **rapid digestion and absorption**, which causes sharp spikes in blood glucose and insulin levels. ([harvard.edu](#)) Foods with a high glycemic index (such as white bread, white rice, and potatoes) are broken down quickly into glucose, which floods the bloodstream and triggers a large release of insulin from the pancreas. Over time, repeated exposure to these blood sugar spikes can lead to **insulin resistance**, a condition in which cells become less responsive to insulin's signals. Insulin resistance is a key feature of metabolic syndrome and a major risk factor for type 2 diabetes, cardiovascular disease, and fatty liver disease.

3.6.1 White Bread, Pasta, and Pastries vs. Whole Grain Alternatives The difference between refined and whole grain products is stark when examined from a nutritional perspective. A slice of **white bread** provides approximately **1 gram of fiber**, while a slice of **whole wheat bread** provides **3-4 grams**. ([American Heart Association](#)) This difference may seem small, but over the course of a day, choosing whole grains can add 10-15 grams of fiber to your diet — a substantial contribution toward the recommended daily intake. Whole grains also provide significantly more vitamins (particularly B vitamins and vitamin E), minerals (magnesium, zinc, iron), and phytonutrients than their refined counterparts.

The glycemic impact of refined versus whole grain carbohydrates also differs dramatically. White bread has a **glycemic index of approximately 75**, while whole grain bread has a glycemic index of around **50-60**. ([harvard.edu](#)) This means that white bread raises blood sugar levels more rapidly and to a greater extent than whole grain bread. For individuals with diabetes or insulin resistance, this difference can have significant implications for blood sugar management. Even for individuals without these conditions, choosing lower-glycemic carbohydrates helps maintain more stable energy levels, reduces hunger and cravings, and supports long-term metabolic health.

4. Building a Healthy Daily Diet

Constructing a healthy daily diet is not about following rigid rules or depriving yourself of foods you enjoy. Rather, it is about making **informed, balanced choices** that prioritize nutrient-dense foods while allowing for flexibility and moderation. The evidence-based guidelines from reputable health organizations such as the USDA, Harvard School of Public Health, and the World Health Organization all converge on a similar set of principles: **emphasize vegetables, fruits, whole grains, and lean proteins; limit processed foods, added sugars, unhealthy fats, and excess sodium; and pay attention to portion sizes and overall calorie balance.** ([WHO](#)) By following these principles, you can create a sustainable eating pattern that supports long-term health and well-being.

4.1 The Healthy Plate Method

The **Healthy Plate method** (also known as the Plate Method or MyPlate) is a simple, visual tool for building balanced meals without the need for counting calories or measuring portions precisely. ([harvard.edu](#)) It is based on the principle of dividing your plate into sections that correspond to different food groups, ensuring that each meal contains an appropriate balance of nutrients. This method is flexible enough to accommodate different cuisines, dietary preferences, and cultural food traditions, making it a practical approach for people from all walks of life. The visual nature of the plate method makes it particularly useful for teaching children about healthy eating and for anyone who finds traditional calorie counting too burdensome.

The Healthy Plate: Recommended Proportions




Figure 3: The Healthy Plate proportions chart showing 50% vegetables and fruits, 25% whole grains, and 25% lean protein

Figure 3: The Healthy Plate model recommends filling half your plate with vegetables and fruits, one-quarter with whole grains, and one-quarter with lean protein.

4.1.1 Half Plate: Vegetables and Fruits The foundation of the Healthy Plate is **vegetables and fruits**, which should occupy **half of your plate at each meal**. ([harvard.edu](https://www.harvard.edu)) This recommendation reflects the overwhelming evidence that diets rich in vegetables and fruits are associated with reduced risk of virtually all major chronic diseases, including heart disease, stroke, diabetes, and certain cancers. (WHO) Vegetables and fruits are nutrient-dense — meaning they provide a high concentration of vitamins, minerals, fiber, and phytonutrients relative to their calorie content — making them ideal foods for maintaining a healthy weight while meeting nutritional needs.

When filling half your plate with vegetables and fruits, **variety and color are key**. Different colored vegetables and fruits contain different phytonutrients, so eating a rainbow of produce ensures that you receive the full spectrum of these beneficial compounds. ([harvard.edu](https://www.harvard.edu)) Dark leafy greens (spinach, kale, collards) provide vitamins A, C, and K, folate, iron, and calcium. Orange and yellow vegetables (carrots, sweet potatoes, bell peppers) are rich in beta-carotene and vitamin C. Red vegetables and fruits (tomatoes, watermelon, strawberries) contain lycopene and anthocyanins. Blue and purple produce (blueberries,

eggplant, purple cabbage) provide anthocyanins and resveratrol. White vegetables (cauliflower, onions, garlic) contain allicin and other sulfur compounds with immune-boosting properties.

4.1.2 Quarter Plate: Lean Protein Lean protein should occupy **one-quarter of your plate** at each meal. ([harvard.edu](#)) Protein is essential for building and repairing tissues, producing enzymes and hormones, and supporting immune function. Including a source of protein at each meal also helps promote satiety, reducing the likelihood of overeating and snacking between meals. The key is to choose lean, high-quality protein sources that provide these benefits without excessive saturated fat, sodium, or calories.

Healthy protein choices include **skinless poultry, fish and seafood, lean cuts of beef and pork, eggs, beans, lentils, tofu, tempeh, nuts, and seeds.** ([National Institute on Aging](#)) When selecting animal proteins, choose lean cuts and limit processed meats (bacon, sausage, deli meats, hot dogs), which are high in sodium and saturated fat and have been linked to increased risk of colorectal cancer. When including plant-based proteins, remember that beans, lentils, and peas also count as vegetables, providing the dual benefit of protein and fiber. Aim for a variety of protein sources throughout the week to ensure a diverse intake of nutrients.

4.1.3 Quarter Plate: Whole Grains Whole grains should occupy the remaining **one-quarter of your plate.** ([harvard.edu](#)) Whole grains provide complex carbohydrates for sustained energy, along with fiber, B vitamins, iron, magnesium, and other important nutrients. Choosing whole grains over refined grains is one of the most impactful dietary changes you can make for your health, as it improves glycemic control, supports digestive health, reduces cardiovascular risk, and may help with weight management.

Healthy whole grain choices include **brown rice, quinoa, whole wheat pasta, whole wheat bread, oats, barley, bulgur, and corn.** ([National Institute on Aging](#)) When shopping for grain products, read the ingredient list and look for “whole grain” or “whole wheat” as the first ingredient. Be wary of products that are labeled “multigrain” or “made with whole grains” but contain mostly refined grains. A serving of grains is typically **1/2 cup of cooked grains** or **one slice of bread**, so the quarter-plate guideline translates to approximately **1/2 to 1 cup of cooked whole grains** per meal, depending on your calorie needs.

4.2 The Five Food Groups

The concept of food groups provides a framework for understanding the different types of foods that contribute to a balanced diet and for ensuring that you obtain a wide range of nutrients. The **five main food groups** — vegetables, fruits, grains, protein foods, and dairy — each provide a unique set of nutrients that are difficult to obtain from other sources. ([National Institute on Aging](#)) By including foods from all five groups each day, you create a nutritional safety net that helps prevent deficiencies and supports optimal health.

Food Group	Key Nutrients Provided	Daily Recommendations	Best Choices	Foods to Limit
Vegetables	Fiber, vitamins A, C, K, folate, potassium, magnesium (NHS)	2.5-3 cups per day	Dark leafy greens, broccoli, carrots, peppers, tomatoes (NHS)	Fried vegetables, vegetables with creamy sauces

Table 13 – continued

Food Group	Key Nutrients Provided	Daily Recommendations	Best Choices	Foods to Limit
Fruits	Fiber, vitamin C, potassium, antioxidants (NHS)	1.5-2 cups per day	Berries, apples, oranges, bananas, melons (NHS)	Fruit canned in syrup, fruit juices (WHO)
Grains	Carbohydrates, fiber, B vitamins, iron, magnesium (National Institute on Aging)	5-8 oz equivalents per day	Brown rice, oatmeal, whole wheat bread, quinoa (National Institute on Aging)	White bread, pastries, sugary cereals (American Heart Association)
Protein Foods	Protein, iron, zinc, B12, omega-3s (National Institute on Aging)	5-6.5 oz equivalents per day	Fish, poultry, beans, lentils, nuts, tofu (National Institute on Aging)	Processed meats, fatty cuts of meat (NHS)
Dairy	Calcium, vitamin D, protein, potassium (NHS)	3 cups per day	Low-fat milk, yogurt, cheese, fortified soy milk (NHS)	Full-fat dairy, flavored yogurts with added sugar (NHS)

Table 13: The five food groups, their key nutrients, daily recommendations, best choices, and foods to limit.

4.2.1 Vegetables: Dark Green, Red, Orange, and Starchy Vegetables are the **most nutrient-dense food group**, providing an unparalleled combination of vitamins, minerals, fiber, and phytonutrients with very few calories. (NHS) The vegetable group is typically divided into subgroups based on color and nutrient content: dark green vegetables (spinach, kale, broccoli, romaine lettuce), red and orange vegetables (carrots, sweet potatoes, tomatoes, red peppers), starchy vegetables (potatoes, corn, green peas), and other vegetables (onions, mushrooms, cauliflower, celery). Each subgroup provides a different array of nutrients, so it is important to eat a variety of vegetables from all subgroups throughout the week.

Dark green vegetables are particularly important because they are excellent sources of **vitamins A, C, and K, folate, iron, and calcium**. (National Institute on Aging) These nutrients support immune function, bone health, blood clotting, and oxygen transport. **Red and orange vegetables** are rich in **beta-carotene** (which the body converts to vitamin A), **vitamin C**, and **lycopene**, an antioxidant that may protect against prostate cancer and heart disease. **Starchy vegetables** such as potatoes and corn provide complex carbohydrates for energy along with fiber, potassium, and vitamin C. While they are sometimes criticized for their carbohydrate content, starchy vegetables are nutrient-dense foods that can be part of a healthy diet when prepared without excessive fat or salt.

4.2.2 Fruits: Fresh, Frozen, Canned, and Dried Fruits are nature's sweet treat, providing **natural sugars along with fiber, vitamins, minerals, and antioxidants**. (NHS) The natural sugars in whole fruits are packaged with fiber, which slows their absorption and prevents the blood sugar spikes associated with added sugars. Fruits are also hydrating, with many varieties containing **80-90% water**. The fiber in fruits supports digestive health and promotes satiety, making them a satisfying snack or dessert option.

All forms of fruit — fresh, frozen, canned, and dried — can be part of a healthy diet, but some forms are better choices than others. **Fresh and frozen fruits** are the best options, as they retain most of their nutrients and do not contain added sugars. **Canned fruits** can be healthy if they are packed in water or **100% fruit juice** rather than heavy syrup. **Dried fruits** are concentrated sources of nutrients and fiber but are also calorie-dense, so portion control is important. **Fruit juices**, even 100% juices, lack the fiber of whole fruits and can contribute to excess calorie intake, so they should be consumed in moderation (no more than one small glass per day). ([WHO](#))

4.2.3 Grains: Whole Grains vs. Refined Grains The grains food group includes any food made from **wheat, rice, oats, cornmeal, barley, or other cereal grains**. ([National Institute on Aging](#)) This includes bread, pasta, breakfast cereals, tortillas, and even popcorn. The critical distinction within this group is between **whole grains** and **refined grains**. Whole grains contain the entire grain kernel — the bran, germ, and endosperm — and provide fiber, B vitamins, iron, magnesium, and other nutrients. Refined grains have been processed to remove the bran and germ, which removes most of the fiber and nutrients. The Dietary Guidelines recommend that **at least half of all grains consumed should be whole grains**. ([National Institute on Aging](#))

Making the switch from refined to whole grains is one of the simplest ways to improve your diet. Instead of white bread, choose whole wheat bread. Instead of white rice, choose brown rice or quinoa. Instead of regular pasta, choose whole wheat pasta. Instead of sugary cereals, choose oatmeal or whole grain cereals with minimal added sugar. These substitutions provide more fiber, vitamins, and minerals without requiring major changes to your eating habits. When reading food labels, look for the word “whole” as the first ingredient (e.g., “whole wheat flour” rather than “enriched wheat flour”) and check the fiber content — a good whole grain product should provide **at least 3 grams of fiber per serving**.

4.2.4 Protein: Meat, Poultry, Seafood, Beans, Nuts The protein foods group includes a wide variety of animal and plant-based foods: **meat, poultry, seafood, eggs, beans, peas, lentils, nuts, seeds, and soy products**. ([National Institute on Aging](#)) This diversity allows for flexibility in meeting protein needs while accommodating different dietary preferences, cultural food traditions, and budget constraints. The key principles for choosing healthy protein foods are to **vary your protein routine**, choose lean options, limit processed meats, and include plant-based proteins regularly.

Seafood is an especially valuable protein source because it provides **omega-3 fatty acids** in addition to high-quality protein. ([National Institute on Aging](#)) The Dietary Guidelines recommend eating **at least 8 ounces of seafood per week**, with an emphasis on fatty fish such as salmon, trout, sardines, and mackerel. For meat and poultry, choose lean cuts and remove visible fat and skin to reduce saturated fat intake. Processed meats (bacon, sausage, deli meats, hot dogs) should be limited due to their high sodium content and association with increased risk of colorectal cancer. Plant-based proteins (beans, lentils, nuts, tofu) offer the advantages of being low in saturated fat, high in fiber, and rich in vitamins and minerals.

4.2.5 Dairy: Milk, Yogurt, Cheese, and Fortified Alternatives The dairy group includes **milk, yogurt, cheese, and fortified soy beverages** (soy milk). ([NHS](#)) These foods are the primary sources of **calcium and vitamin D** in the American diet, nutrients that are essential for bone health. Dairy products also provide high-quality protein, potassium, phosphorus, and B vitamins. For individuals who are lactose intolerant or follow a vegan diet, fortified soy milk is the closest nutritional equivalent to cow’s milk, providing similar amounts of protein, calcium, and vitamin D when fortified.

When choosing dairy products, **low-fat and fat-free options** are preferable to full-fat versions, as they

provide the same protein and calcium with less saturated fat and fewer calories. (NHS) Flavored yogurts and milk-based desserts (ice cream, pudding) can be high in added sugars, so it is important to read labels and choose plain, unsweetened versions when possible. For cheese, which is naturally high in sodium and saturated fat, portion control is important — a serving is **1.5 ounces** (about the size of three dice). Fortified plant-based milks (almond, oat, pea) can be part of a healthy diet, but it is important to choose unsweetened, calcium-fortified versions, as they vary widely in their nutritional content.

4.3 Daily Calorie Guidelines

Calories are units of energy that the body uses to fuel all of its activities, from basic metabolic functions to physical exercise. The number of calories a person needs each day depends on several factors, including **age, sex, body size, activity level, and metabolic health**. (National Institute on Aging) Understanding your individual calorie needs and how they change with age and activity level can help you maintain a healthy weight and ensure that you are consuming enough energy to support your lifestyle.




Figure 4: Bar chart showing recommended daily calorie intake by gender, age, and activity level

Figure 4: Recommended daily calorie intake varies by gender, age, and physical activity level. Active individuals require significantly more calories than sedentary individuals.

4.3.1 Factors Affecting Calorie Needs: Age, Gender, Activity Level Age is one of the primary determinants of calorie needs. As people age, their **basal metabolic rate** (the number of calories burned at rest) tends to decline due to loss of muscle mass and changes in hormone levels. (National Institute on Aging) This means that older adults generally need fewer calories than younger adults to maintain the same body weight. However, older adults often have **increased needs for certain nutrients** such as protein, calcium, and vitamin D, making nutrient density (nutrients per calorie) an important consideration.

Gender also affects calorie needs, primarily due to differences in **body composition and muscle mass**. Men typically have more muscle mass and less body fat than women, which means they burn more calories at rest and during physical activity. (National Institute on Aging) As a result, men generally require more calories than women of the same age and activity level. For example, a sedentary 30-year-old man

needs approximately **2,400 calories per day**, while a sedentary 30-year-old woman needs approximately **2,000 calories per day**.

Physical activity level has a major impact on calorie needs. The more active you are, the more calories you burn, and the more calories you need to consume to maintain your weight. (National Institute on Aging) Activity levels are generally categorized as **sedentary** (little to no exercise), **moderately active** (light exercise 1-3 days per week), and **active** (moderate to vigorous exercise most days of the week). An active individual may need **500-1,000 more calories per day** than a sedentary individual of the same age and gender. The table below provides estimated daily calorie needs based on these factors.

Gender	Age	Sedentary	Moderately Active	Active
Female	19-30 years	2,000 cal (National Institute on Aging)	2,000-2,200 cal (National Institute on Aging)	2,400 cal (National Institute on Aging)
Female	31-50 years	1,800 cal (National Institute on Aging)	2,000 cal (National Institute on Aging)	2,200 cal (National Institute on Aging)
Female	51+ years	1,600 cal (National Institute on Aging)	1,800 cal (National Institute on Aging)	2,000-2,200 cal (National Institute on Aging)
Male	19-30 years	2,400 cal (National Institute on Aging)	2,600-2,800 cal (National Institute on Aging)	3,000 cal (National Institute on Aging)
Male	31-50 years	2,200 cal (National Institute on Aging)	2,400-2,600 cal (National Institute on Aging)	2,800-3,000 cal (National Institute on Aging)
Male	51+ years	2,000 cal (National Institute on Aging)	2,200-2,400 cal (National Institute on Aging)	2,400-2,800 cal (National Institute on Aging)

Table 14: Estimated daily calorie needs by gender, age, and physical activity level.

4.3.2 Understanding Calorie Balance for Weight Management **Calorie balance** — the relationship between calories consumed and calories burned — is the fundamental principle underlying weight management. (National Institute on Aging) When calorie intake equals calorie expenditure, weight remains stable. When intake exceeds expenditure, the excess calories are stored as fat, leading to weight gain. When expenditure exceeds intake, the body draws on stored fat for energy, leading to weight loss. While the concept of “calories in, calories out” is often oversimplified, it remains a useful framework for understanding weight management.

However, not all calories are created equal. The **source of calories** matters for health, even when total calorie intake is controlled. A diet of 2,000 calories from whole foods (vegetables, fruits, whole grains, lean proteins, healthy fats) will have very different effects on metabolism, satiety, hormone levels, and overall health than a diet of 2,000 calories from processed foods, sugary beverages, and fast food. (hers) Nutrient-dense foods provide vitamins, minerals, fiber, and other beneficial compounds that support metabolic health and help regulate appetite, while empty-calorie foods can disrupt metabolism and promote overeating. For sustainable weight management, focus on both calorie balance **and** food quality.

4.4 Portion Control and Mindful Eating

Portion sizes in the United States have **increased dramatically** over the past several decades, contributing to overeating and weight gain. ([Croí Heart & Stroke Charity](#)) The average restaurant meal today is more than **four times larger** than it was in the 1950s, and packaged food portions have similarly expanded. This “portion distortion” means that many people are consuming far more calories than they realize, even when they are choosing relatively healthy foods. Learning to recognize appropriate portion sizes and practicing mindful eating habits can help you enjoy your food while maintaining a healthy weight.

4.4.1 Recognizing Appropriate Portion Sizes One of the simplest ways to control portions is to use the **Plate Method** described earlier, which naturally limits portion sizes by dividing the plate into sections for different food groups. ([Croí Heart & Stroke Charity](#)) Using a **smaller plate** (9 inches in diameter rather than the standard 12-inch dinner plate) can also help, as it makes appropriate portions look more visually satisfying. Studies have shown that simply switching to smaller plates can result in reduced calorie intake and weight loss over time.

For foods that are not easily portioned on a plate, using your **hand as a guide** can be helpful: a serving of protein (meat, fish, poultry) should be about the **size of your palm**; a serving of grains or starchy vegetables should be about the **size of your fist**; a serving of fats (nuts, oils, butter) should be about the **size of your thumb**; and a serving of vegetables can be **two cupped handfuls**. ([Baker Heart and Diabetes Institute](#)) Another useful strategy is to **measure foods** occasionally to recalibrate your sense of portion sizes — for example, measuring out 1/2 cup of cereal or 3 ounces of meat to see what a standard serving actually looks like.

4.4.2 Tips for Avoiding Overeating **Mindful eating** is a practice that involves paying full attention to the experience of eating — the taste, texture, aroma, and appearance of food, as well as the body’s hunger and fullness cues. ([Croí Heart & Stroke Charity](#)) In contrast, mindless eating (eating while distracted by television, smartphones, or work) often leads to overconsumption because the brain does not register the food intake as fully. To practice mindful eating, try to eat without distractions, chew your food thoroughly, put your fork down between bites, and check in with your body’s hunger and fullness signals throughout the meal.

Other practical strategies for avoiding overeating include: **eating slowly** (it takes about **20 minutes** for the brain to register fullness after eating begins); **drinking water before and during meals** to promote satiety; **serving meals in the kitchen** rather than at the table to discourage seconds; **packaging leftovers** into portion-sized containers immediately after cooking; and **waiting 20 minutes** after finishing a meal before deciding whether you are still hungry. ([Queensland Health](#)) When eating out, consider sharing an entree, ordering an appetizer as your main course, or asking for a to-go box at the beginning of the meal and packing up half of your food before you start eating.

4.5 MyPlate Recommendations

MyPlate is the **USDA’s current nutrition guide**, introduced in 2011 as a replacement for the Food Pyramid. It uses a familiar mealtime visual — a place setting — to illustrate the five food groups and their recommended proportions. ([harvard.edu](#)) The MyPlate icon shows a plate divided into four sections (vegetables, fruits, grains, and protein) with a side of dairy, representing a balanced meal. MyPlate also emphasizes the importance of choosing nutrient-dense foods, limiting added sugars, saturated fats, and sodium, and balancing calories with physical activity.

4.5.1 Avoid Large Portions MyPlate and other dietary guidelines emphasize the importance of **portion control** as a key strategy for maintaining a healthy weight and preventing overconsumption of calories, sodium, and unhealthy fats. ([Croí Heart & Stroke Charity](#)) Even healthy foods can contribute to weight gain when eaten in excessive amounts. The calorie-dense nature of some healthy foods (nuts, oils, avocado, cheese) means that portion control is especially important for these items. Being mindful of portion sizes at every meal and snack helps ensure that calorie intake aligns with calorie needs.

4.5.2 Reduce Salt Intake MyPlate recommends choosing foods that are **low in sodium** and limiting the addition of salt at the table. ([American Heart Association](#)) As discussed earlier, excess sodium intake is a major contributor to hypertension and cardiovascular disease. Most of the sodium in the American diet comes from processed and restaurant foods, so cooking at home with fresh ingredients is one of the most effective ways to reduce sodium intake. When using canned goods (beans, vegetables, tomatoes), choose low-sodium or no-salt-added versions, and rinse canned foods before using them to remove some of the sodium. Flavor foods with herbs, spices, garlic, lemon juice, and vinegar instead of salt.

4.5.3 Replace Sugary Drinks with Water One of the most impactful dietary changes recommended by MyPlate is to **replace sugary beverages with water**. ([CDC](#)) Sugar-sweetened beverages are the largest source of added sugars in the American diet and provide empty calories without any nutritional benefit. Making water your primary beverage is one of the simplest and most effective ways to reduce calorie intake, improve hydration, and support overall health. If plain water seems boring, try adding slices of lemon, lime, cucumber, or fresh herbs (mint, basil) for flavor without calories. Unsweetened tea and coffee are also healthy beverage choices that can contribute to daily fluid intake.

4.5.4 Fill the Plate with Fruits, Vegetables, and Whole Grains The overarching message of MyPlate is to **fill your plate with nutrient-dense foods** from the vegetable, fruit, and whole grain groups, while including moderate amounts of lean protein and dairy. ([harvard.edu](#)) These foods provide the vitamins, minerals, fiber, and phytonutrients that support optimal health and protect against chronic disease. By making vegetables, fruits, and whole grains the foundation of every meal, you naturally crowd out less healthy foods and create a dietary pattern that promotes long-term well-being. Remember that healthy eating is not about perfection — it is about making consistently good choices most of the time, while allowing for flexibility and enjoyment.

5. Key Nutrition Guidelines to Live By

After reviewing the extensive evidence on nutrition and health, several key principles emerge as the foundation of a healthy eating pattern. These guidelines are supported by decades of scientific research and are endorsed by major health organizations worldwide. Following these principles does not require following a restrictive diet or eliminating favorite foods entirely — rather, it is about making informed choices, practicing moderation, and building sustainable habits that support lifelong health.

5.1 Decrease Saturated Fats, Salt, and Empty Calories

The first principle of healthy eating is to **limit the harmful nutrients** that contribute to chronic disease. This means reducing intake of saturated fats (found in fatty meats, full-fat dairy, and tropical oils), sodium (found in processed and restaurant foods), and empty calories (found in sugary beverages, desserts, fried foods, and ultra-processed snacks). ([American Heart Association](#)) These foods provide calories without beneficial nutrients and are strongly linked to obesity, heart disease, type 2 diabetes, stroke, and certain cancers.

Making these reductions does not mean giving up flavor or enjoyment. **Healthy fats** such as olive oil, avocado, and nuts can replace butter and lard in cooking and at the table. **Herbs, spices, citrus, and vinegar** can add abundant flavor to foods without adding sodium. **Fresh fruits, dark chocolate, and homemade baked goods** with reduced sugar can satisfy a sweet tooth without the excess calories and unhealthy fats of commercially produced desserts. By focusing on what you can eat rather than what you cannot, these changes become sustainable and even enjoyable.

5.2 Increase Fiber Intake

The second principle is to **actively increase dietary fiber intake**. Fiber is one of the most beneficial yet underconsumed nutrients in the modern diet, with average intake falling far short of the recommended **25-38 grams per day**. ([Mayo Clinic](#)) Increasing fiber intake is associated with a remarkable array of health benefits, including improved digestive health, lower cholesterol, better blood sugar control, weight management, reduced risk of heart disease and type 2 diabetes, and possibly even reduced risk of certain cancers and depression.

The best way to increase fiber intake is to **emphasize whole, plant-based foods** at every meal. Choose whole grain breads, cereals, and pasta instead of refined versions. Include beans, lentils, or peas in meals several times per week. Eat a variety of fruits and vegetables, including the skins when edible. Snack on nuts and seeds rather than chips or crackers. These simple substitutions can easily add 15-20 grams of fiber to your daily intake, bringing you much closer to the recommended amount.

5.3 Choose Nutrient-Dense Foods Over Processed Options

The third principle is to **prioritize nutrient-dense foods** — those that provide a high concentration of vitamins, minerals, fiber, and other beneficial compounds relative to their calorie content. ([WHO](#)) Vegetables, fruits, whole grains, legumes, nuts, seeds, lean proteins, and low-fat dairy are all nutrient-dense choices that support health and provide satiety. In contrast, processed and ultra-processed foods tend to be calorie-dense but nutrient-poor, providing excess calories, sodium, unhealthy fats, and added sugars without the beneficial nutrients your body needs.

Shopping the **perimeter of the grocery store** (where fresh produce, meats, dairy, and whole grains are typically located) rather than the center aisles (where processed and packaged foods predominate) is a simple strategy for choosing more nutrient-dense foods. Cooking meals at home using fresh ingredients gives you control over what goes into your food and typically results in meals that are lower in sodium, added sugars, and unhealthy fats than restaurant or takeout meals. When you do choose processed foods, read the Nutrition Facts panel and ingredient list, and select products with shorter ingredient lists, less sodium, less added sugar, and more fiber.

5.4 Balance Calorie Intake with Physical Activity

The fourth principle is to **balance the calories you consume with the calories you burn through physical activity**. ([National Institute on Aging](#)) This principle acknowledges that healthy eating is only one part of the equation — regular physical activity is equally important for maintaining a healthy weight, building and preserving muscle mass, supporting cardiovascular health, improving mood and mental health, and reducing the risk of chronic disease. The Physical Activity Guidelines for Americans recommend at least **150 minutes of moderate-intensity aerobic activity** (such as brisk walking) or **75 minutes of vigorous-intensity activity** (such as running) per week, along with muscle-strengthening activities on two or more days per week.

Physical activity not only burns calories but also improves insulin sensitivity, reduces blood pressure, strengthens bones, boosts mood, and enhances overall quality of life. Finding activities you enjoy — whether it is walking, swimming, dancing, cycling, gardening, or playing sports — makes it easier to maintain an active lifestyle. Combining regular physical activity with a nutrient-dense diet creates a powerful synergy that supports optimal health and well-being at any age.




Figure 5: Bar chart comparing good vs bad nutrients and their health impact scores

Figure 5: A comparison of good nutrients (green bars) versus bad nutrients (red bars), showing their relative impact on health. Prioritizing foods from the green side while limiting foods from the red side is the key to a healthy diet.