


# PP Hypothesis Tests

## Hypothesis Tests for CSR

- With a spatial point process, *a single realization from the underlying point process is observed*.
- Hypothesis tests and Monte Carlo simulation procedures allow testing for departures from Complete Spatial Randomness (CSR).
- Devise and implement a Monte Carlo-based test for CSR. First detail your test statistic and implementation, then assess the results with the four data sets from Lecture 21.



## G and F Functions

- One way to describe a spatial point process, is to consider the probability of being a certain distance from a point or similarly, the number of points expected in a distance from a point.

- Define  $Num(\mathbf{s}, d; \mathbf{S})$  as the number of points from  $\mathbf{S}$  in a circle of diameter  $d$  from point  $\mathbf{s}$ .

- Then we can compute the probability that more than one point exists with a distance  $d$  of an observed point.

$$E_{\mathbf{S}} \left( \sum_{\mathbf{s}_i \in \mathbf{S}} 1(Num(\mathbf{s}_i), d; \mathbf{S}) > 0 \right) = \lambda |\mathbf{D}| P(Num_d(\mathbf{s}, d; \mathbf{S}) > 0)$$

- The term  $P(Num_d(\mathbf{s}, d; \mathbf{S}) > 0)$  is referred to as  $G(d)$ .  $G(d)$  increases in  $d$ .

- The  $G(d)$  function is similar to a CDF of the **nearest neighbor** distance

- A similar statistic is the  $F(d)$  function. Whereas  $G(d)$  is centered at the observed  $\mathbf{s}_i$ ,  $F(d)$  is defined at any arbitrary point. Hence this is a CDF for empty space.

- **QUESTION:** Assume a CSR Poisson process with constant intensity  $\lambda$ , calculate  $F(d)$  or  $G(d)$  (they are the same).

- Discuss how to create an empirical estimate of  $\hat{G}(d)$ , given a realization of a point process.

- With bounded area, edge correction procedures are necessary. *Hence, consider*

$$\hat{G}(d) = \frac{\sum_i 1(d_i \leq d < b_i)}{\sum_i 1(d < b_i)},$$

where  $b_i$  is the distance from  $\mathbf{s}_i$  to the edge and  $d_i$  is the distance to the nearest neighbor of  $\mathbf{s}_i$ .

- The empirical estimates of  $G$  or  $H$  can be compared with  $G$  or  $F$  using a QQ-plot.

- **Discuss:** What would be the implications of shorter tails or longer tails than expected under CSR?  
*shorter tail = clustering/attraction, longer tail = inhibition/repulsion*

- *Describe a natural process that might cluster and another that might repel*