

Appendix B. Parameters

LZU-CHINA

October 10, 2023

Abstract

This appendix presents and processes the data of the fluorescence protein intensity F and cell density mentioned in the aforementioned model. In fact, we added the fluorescence protein gene after the oleic acid inducer operator fadO. When FadR binds with the operator fadO, it leads to the activation of the oleic acid inducer, causing the expression of the fluorescence protein gene. Thus, we can detect the corresponding data.

We tested Raw Fluorescence Intensity and OD600 (Optical Density at 600 nm) in three scenarios of oleic acid concentrations: 5%, 10%, and 15%, under both aerobic and anaerobic conditions. For each type, we conducted six sets of data. Subsequently, we organized and preprocessed the respective experimental data to aid in subsequent parameter estimation.

1 Parameters

Parameter description	Term	Value	Units	Reference
Forward sequestration	k_f	612.55	$\mu\text{M}^{-2} \cdot \text{h}^{-1}$	[2]
Reverse sequestration	k_r	900.73	h^{-1}	[2]
PlsB turnover rate	$k_{\text{cat},B}$	192.91	h^{-1}	[2]
FadD turnover rate	$k_{\text{cat},D}$	49	h^{-1}	[2]
FadR'strength to promote itself(NAR)	a_R	0.0131	$\mu\text{M} \cdot \text{h}^{-1}$	[2]
FadR leaky expression (NAR)	b_R	0.0007	$\mu\text{M} \cdot \text{h}^{-1}$	[2]
FadD'strength to promote itself	a_D	0.0517	$\mu\text{M} \cdot \text{h}^{-1}$	[2]
FadD leaky expression	b_D	0.0108	$\mu\text{M} \cdot \text{h}^{-1}$	[2]
FadR affinity to promote itself (NAR)	K_R	4.3222	μM^{-1}	[2]
FadR affinity to fadD	K_D	305.95	μM^{-1}	[2]
Michaelis constant	$K_{m,B}$	45429	μM	[2]
Michaelis constant	$K_{m,D}$	0.0672	μM	[2]
PlsB concentration	B	0.1369	μM	[2]
Cell growth parameter	λ_{max}	0.1818	h^{-1}	[2], E. coli DH1Δ fadE strain.
E_g promoter strength	a_g	λ_{max}	$\mu\text{M} \cdot \text{h}^{-1}$	In order to ensure full express at λ_{max} .
FadR'strength to promote itself(PAR)	a_R	0.0131^\dagger	$\mu\text{M} \cdot \text{h}^{-1}$	Suppose that the parameters of FadR with NAR are same.
FadR leaky expression (PAR)	b_R	0.0007^\dagger	$\mu\text{M} \cdot \text{h}^{-1}$	Suppose that the parameters of FadR with NAR are same.
Decrease paramameter in growth	s_T	0.7	—	Determined by pgi down-regulation.
FadR affinity to own promoter (PAR)	K_R	$4.3222^\dagger \times 7$	μM^{-1}	Suppose that the parameters of FadR with PAR are same. It is same as system with NAR before induction (OA = 0μM) when steady state FadR approx except scaled.
FadR affinity to E_g promoter	K_g	$= K_R = 4.9114$	μM^{-1}	Suppose designed same promoter as fadR with PAR,for FadR activated expression.
Affinity of FadR for prod synthesis enzyme E_p	K_p	$= K_D = 305.95$	$\mu\text{M} \cdot \text{h}^{-1}$	Suppose designed with FadR operator site from fadD promoter.