

Multiple Comparison Procedures To A Control

For AN(C)OVA Models

Statsomat.com

Contributors*

14 Juni 2021

Contents

Basic Information	2
Model Information	3
Descriptive Plots	3
Dependent Variable	3
Dependent Against Categorical Factors	5
References	7

*Denise Welsch, Markus Neuhäuser, Viktoria Daum, Linda Müller, Damian Nink, Simone Schüttler, Daniela Wüller

Basic Information

Automatic statistics for the file:

File
recovery.csv

Your selection for the encoding: UTF-8

Your selection for the decimal character: .

Observations (rows with at least one non-missing value): 41

Variables (columns with at least one non-missing value): 2

Variables considered continuous: 1

Variables considered continuous
minutes

Variables considered categorical: 1

Variables considered categorical
blanket

Model Information


You defined the following linear model: minutes~blanket

You are interested in the factor: blanket


You are interested in pairwise comparisons to the control factor level: b0

Descriptive Plots

Dependent Variable




Boxplot of minutes



Dependent Against Categorical Factors

Boxplot of minutes ~ blanket







Anova Table (Type III tests)

Response: minutes

	Sum Sq	Df	F value	Pr(>F)
(Intercept)	4380.8	1	652.8851	< 2.2e-16 ***
blanket	152.0	3	7.5499	0.0004619 ***
Residuals	248.3	37		

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1



Simultaneous Tests for General Linear Hypotheses

Multiple Comparisons of Means: Dunnett Contrasts

```
Fit: lm(formula = modelfunction, data = df_factorized)
```

Linear Hypotheses:

	Estimate	Std. Error	t value	Pr(<t)
b1 - b0 >= 0	-2.1333	1.6038	-1.330	0.2411
b2 - b0 >= 0	-7.4667	1.6038	-4.656	<0.001 ***
b3 - b0 >= 0	-1.6667	0.8848	-1.884	0.0923 .

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
(Adjusted p values reported -- single-step method)

Simultaneous Confidence Intervals

Multiple Comparisons of Means: Dunnett Contrasts

```
Fit: lm(formula = modelfunction, data = df_factorized)
```

Quantile = 2.1823
95% family-wise confidence level

Linear Hypotheses:

	Estimate	lwr	upr
b1 - b0 >= 0	-2.1333	-Inf	1.3667
b2 - b0 >= 0	-7.4667	-Inf	-3.9667
b3 - b0 >= 0	-1.6667	-Inf	0.2642

References

- Fox, John, and Sanford Weisberg. 2019. *An R Companion to Applied Regression*. Third. Thousand Oaks CA: Sage. <https://socialsciences.mcmaster.ca/jfox/Books/Companion/>.
- Gross, Juergen, and Uwe Ligges. 2015. *Nortest: Tests for Normality*. <https://CRAN.R-project.org/package=nortest>.
- Madsen, Jacob H. 2018. *DDoutlier: Distance & Density-Based Outlier Detection*. <https://CRAN.R-project.org/package=DDoutlier>.
- R Core Team. 2019. *R: A Language and Environment for Statistical Computing*. Vienna, Austria: R Foundation for Statistical Computing. <https://www.R-project.org/>.
- Zeileis, Achim, and Torsten Hothorn. 2002. "Diagnostic Checking in Regression Relationships." *R News* 2 (3): 7–10. <https://CRAN.R-project.org/doc/Rnews/>.