Multiple Comparison Procedures To A Control

For AN(C)OVA Models

Statsomat.com

Contributors*

22 April 2021

Contents

Basic Information	2
References	6

^{*}Denise Welsch, Markus Neuhäuser, Studentin1, Studentin2, Studentin3, Studentin4

Basic Information

Automatic statistics for the file:	
	File
	recovery.csv
Your selection for the encoding: UTF-8 Your selection for the decimal character: . Observations (rows with at least one non-missing value): 41 Variables (columns with at least one non-missing value): 2 Variables considered continuous: 1	
	Variables considered continuous
Variables considered categorical: 1	
	Variables considered categorical
	blanket


Anova Table (Type III tests)


Response: minutes

Sum Sq Df F value Pr(>F)
(Intercept) 4380.8 1 652.8851 < 2.2e-16 ***
blanket 152.0 3 7.5499 0.0004619 ***

Residuals 248.3 37

Signif. codes: 0 '***' 0.001 '**' 0.05 '.' 0.1 ' ' 1





Simultaneous Tests for General Linear Hypotheses

18₀

0

14

0

Linear Hypotheses:

Quantile = 2.1818

Estimate lwr upr b1 - b0 >= 0 -2.1333 -Inf 1.3658 b2 - b0 >= 0 -7.4667 -Inf -3.9676 b3 - b0 >= 0 -1.6667 -Inf 0.2637

95% family-wise confidence level

Simultaneous Tests for General Linear Hypotheses

Multiple Comparisons of Means: Dunnett Contrasts

Fit: lm(formula = modelfunction, data = df_factorized)

Linear Hypotheses:

Estimate Std. Error t value Pr(<t)b1 - b0 >= 0 -2.1333 1.6038 -1.330 0.0958 . b2 - b0 >= 0 -7.4667 1.6038 -4.656 6.2e-05 *** b3 - b0 >= 0 -1.6667 0.8848 -1.884 0.0640 .

```
Signif. codes: 0 '***' 0.001 '**' 0.05 '.' 0.1 ' ' 1
(Adjusted p values reported -- free method)
    Simultaneous Tests for General Linear Hypotheses
Multiple Comparisons of Means: Dunnett Contrasts
Fit: lm(formula = modelfunction, data = df_factorized)
Linear Hypotheses:
           Estimate Std. Error t value Pr(<t)
b2 - b0 >= 0 -7.4667 1.1095 -6.730 <0.001 ***
b3 - b0 >= 0 -1.6667 0.8642 -1.929 0.0847.
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
(Adjusted p values reported -- single-step method)
    Simultaneous Confidence Intervals
Multiple Comparisons of Means: Dunnett Contrasts
Fit: lm(formula = modelfunction, data = df_factorized)
Quantile = 2.1829
95% family-wise confidence level
Linear Hypotheses:
           Estimate lwr
                           upr
b1 - b0 >= 0 -2.1333 -Inf 1.6531
b2 - b0 >= 0 -7.4667 -Inf -5.0448
b3 - b0 >= 0 -1.6667 -Inf 0.2198
    Simultaneous Tests for General Linear Hypotheses
```

Multiple Comparisons of Means: Dunnett Contrasts

Fit: lm(formula = modelfunction, data = df_factorized)

Linear Hypotheses:

References

Fox, John, and Sanford Weisberg. 2019. An R Companion to Applied Regression. Third. Thousand Oaks CA: Sage. https://socialsciences.mcmaster.ca/jfox/Books/Companion/.

Gross, Juergen, and Uwe Ligges. 2015. Nortest: Tests for Normality. https://CRAN.R-project.org/package=nortest.

Madsen, Jacob H. 2018. DDoutlier: Distance & Density-Based Outlier Detection. https://CRAN.R-project.org/package=DDoutlier.

R Core Team. 2019. R: A Language and Environment for Statistical Computing. Vienna, Austria: R Foundation for Statistical Computing. https://www.R-project.org/.

Zeileis, Achim, and Torsten Hothorn. 2002. "Diagnostic Checking in Regression Relationships." R News 2 (3): 7-10. https://CRAN.R-project.org/doc/Rnews/.