

Multiple Comparison Procedures To A Control

For AN(C)OVA Models

Statsomat.com

Contributors*

22 Juli 2021

Contents

Basic Information	2
Model Information	3
Descriptive Plots	3
Dependent Variable	3
Dependent Against Categorical Factors	5
Analysis of variance	6
Effect of the separate expressions of the given variables (Parameter Estimates)	6
Total influence of factors (ANOVA Type III)	6
Multiple Comparisons of Means to a Control	8
Dunnet	8
References	8

*Denise Welsch, Viktoria Daum, Linda Müller, Damian Nink, Simone Schüttler, Daniela Wüller

Basic Information

Automatic statistics for the file:

File
recovery.csv

Your selection for the encoding: UTF-8

Your selection for the decimal character: .

Observations (rows with at least one non-missing value): 41

Variables (columns with at least one non-missing value): 2

Variables considered continuous: 1

Variables considered continuous
minutes

Variables considered categorical: 1

Variables considered categorical
blanket

Model Information


You defined the following linear model: minutes~blanket

You are interested in the factor: blanket


You are interested in pairwise comparisons to the reference level: b0

Descriptive Plots

Dependent Variable




Boxplot of minutes



Dependent Against Categorical Factors

Boxplot of minutes ~ blanket



Analysis of variance





Effect of the separate expressions of the given variables (Parameter Estimates)

Variable	Value	Std.Error	T.value	P.value	sign. level ¹	Significance at 5 percent error
(Intercept)	11.98	0.57	20.91	<0.001	***	Intercept Significant.
blanket1	2.82	0.70	4.00	<0.001	***	Significant. A Difference between the effect of blanket1 and its reference.
blanket2	0.68	1.20	0.57	0.573		Not Significant. No difference between the effect of blanket2 and its reference.
blanket3	-4.65	1.20	-3.87	<0.001	***	Significant. A Difference between the effect of blanket3 and its reference.

¹ '***': sign. to 0.1% error. '**': sign. to 1% error. '*': sign. to 5% error. '.' : sign. to 10% error. ' ': not sign. ' - ': no statement.

Total influence of factors (ANOVA Type III)

Variable	Sum.Sq	Df	F.value	P.value	Interpretation (5% error)
(Intercept)	2933.11	1	437.13	<0.001	Intercept significantly different from zero.
blanket	151.98	3	7.55	<0.001	There exist significant differences between the levels of factor 2.
Residuals	248.27	37			



Multiple Comparisons of Means to a Control

Theoretical background: Testing multiple hypotheses simultaneously and each at the same pre-specified significance level, increases the probability of false positive effects. The probability to commit at least one false positive decision increases with the number of hypotheses. A solution to overcome this problem is given by multiple comparisons procedures. Here, we do not control the per-hypothesis Type I error but the probability of committing at least one Type I error over all hypotheses. Using p-values adjusted for multiplicity, individual hypotheses can be finally compared with the pre-specified significance level.

Dunnet

Multiple Comparison: Dunnet Contrasts

Test whether the factor level b0 of the factor blanket is different from the other levels. The Null Hypothesis is for example $b1 - b0 = 0$.

Null Hypothesis	Value	Std.Error	T.value	adjusted P.value	Sign. level ¹	Significance at 5 percent Type I error
$b1 - b0 = 0$	-2.13	1.60	-1.33	0.456	***	Not Significant. Level b0 of factor blanket is not different than b1 ²
$b2 - b0 = 0$	-7.47	1.60	-4.66	<0.001		Significant. Level b2 of factor blanket is significantly different than b0 ³
$b3 - b0 = 0$	-1.67	0.88	-1.88	0.182		Not Significant. Level b0 of factor blanket is not different than b3 ²

¹ '***': sign. to 0.1% error. '**': sign. to 1% error. '*': sign. to 5% error. '.' : sign. to 10% error. ' ': not sign. '- ': no statement.

² H1 does not hold significantly.

³ H1 holds significantly.

Simoultaneous Confidence Intervals: Dunnet Contrasts

Simultaneous Confidence Intervals which includes the true value of the difference between the reference level b0 and the other levels of blanket

Null Hypothesis	Value	Lower bound	Upper bound	Interpretation
$b1 - b0 = 0$	-2.13	-6.12	1.86	The interval (-6.12, 1.86) traps the true difference b1-b0 with probability 95 percent. ²
$b2 - b0 = 0$	-7.47	-11.46	-3.48	The interval (-11.46, -3.48) traps the true difference b2-b0 with probability 95 percent. ¹
$b3 - b0 = 0$	-1.67	-3.87	0.53	The interval (-3.87, 0.53) traps the true difference b3-b0 with probability 95 percent. ²

¹ Remark: Zero is not in the confidence interval.

² Remark: Zero is in the confidence interval.

References

- Bretz, Frank, and Peter Westfall Torsten Hothorn. 2010. *Multiple Comparisons Using r*. 1st Edition. Chapman; Hall/CRC. <https://doi.org/10.1201/9781420010909>.
- Fox, John, and Sanford Weisberg. 2019. *An R Companion to Applied Regression*. Third. Thousand Oaks CA: Sage. <https://socialsciences.mcmaster.ca/jfox/Books/Companion/>.
- Gross, Juergen, and Uwe Ligges. 2015. *Nortest: Tests for Normality*. <https://CRAN.R-project.org/package=nortest>.

- R Core Team. 2019. *R: A Language and Environment for Statistical Computing*. Vienna, Austria: R Foundation for Statistical Computing. <https://www.R-project.org/>.
- Zeileis, Achim, and Torsten Hothorn. 2002. “Diagnostic Checking in Regression Relationships.” *R News* 2 (3): 7–10. <https://CRAN.R-project.org/doc/Rnews/>.