Dokładna redukcja potrójnego overlapu Yukawy w TGP

Schwinger, Feynman, PDE, geometria simpleksu, asympotyki i nieelementarność $f_\triangle(t)$

Marzec 2026

Abstract

Prezentujemy kompletne wyprowadzenie i analizę trzycentrowego overlapu Yukawy $I_Y = \int d^3x\, G_m(|x-x_1|)\, G_m(|x-x_2|)\, G_m(|x-x_3|)$ występującego w teorii TGP jako wkład trzyciałowy z członu Φ^4. Główne wyniki:

(i) Dwie równoważne dokładne reprezentacje 2D: Schwingera (5) i Feynmana (7).

(ii) Układ równań PDE Helmholtza (8)–(10) na przestrzeni kształtów trójkąta.

(iii) Dokładna zamknięta formuła gęstości $p(\Delta)$ miary uniformowej na 2-simpleksie (16).

(iv) Redukcja 2D→1D: dokładna jednowymiarowa całka dla $f_\triangle(t)$ (17).

(v) Nieoczekiwana tożsamość (18): $\int_{\Delta_2} \Delta^{-3/2}\, d\alpha = 2\pi$.

(vi) Dokładne asympotyki: Yukawa $t \to \infty$ (19) i Coulomb $t \to 0$ (20).

(vii) Dowód nieelementarności $f_\triangle(t)$ (21).

Contents

1 Definicja i kontekst TGP **4**
- 1.1 Overlap Yukawy . . . 4
- 1.2 Miejsce w TGP . . . 5

2 Reprezentacja Schwingera (przestrzeń fizyczna) **5**

3 Reprezentacja Feynmana (przestrzeń pędów) **5**

4 Układ równań PDE **6**

5 Redukcja do rodzin jednoparametrowych **6**
- 5.1 Trójkąt równoboczny . . . 6
- 5.2 Symetryczna rodzina koliniarna . . . 6

6 Geometria simpleksu: dokładna gęstość $p(\Delta)$ **6**
- 6.1 Struktura poziomic $\Delta = \text{const}$ . . . 6
- 6.2 Punkt krytyczny . . . 6
- 6.3 Pole czapek i gęstość . . . 7

7 Redukcja 2D→1D i kluczowa tożsamość **7**
- 7.1 Dokładna reprezentacja 1D . . . 7
- 7.2 Kluczowa tożsamość: $C_1 = 2\pi$ . . . 7

8 Asympotyki dokładne **8**
- 8.1 Granica Yukawa ($t \to \infty$): metoda siodłowa . . . 8
- 8.2 Granica Coulomba ($t \to 0$): rozw-inięcie logarytmiczne . . . 8

9 Nieelementarność $f_\triangle(t)$ **9**

10 Podsumowanie wyników **9**
10.1 Strategia dla TGP runtime . . . 9
10.2 Wkład trzyciałowy w TGP . . . 10

11 Non-stationary TGP equations of motion **10**
11.1 Starting point: the time-dependent field equation . . . 10
11.2 Linearisation around the vacuum . . . 10
11.3 Retarded solution: the radiation field . . . 10
11.4 Non-stationary pairwise interaction . . . 11
11.5 Radiation: power loss . . . 11
11.6 Non-stationary N-body equations of motion . . . 11
11.7 Hierarchy of approximations . . . 12
11.8 Retardation time scale . . . 12
11.9 Non-stationary 3-body effects: summary . . . 12

12 Konkretne przewidywania TGP dla układów wielociałowych **13**
12.1 Kiedy efekty trójciałowe dominują? Liczba krytyczna N_{crit} . . . 13
12.2 Promieniowanie skalarne TGP: odcięcie częstotliwości . . . 14
12.3 Podsumowanie: trzy nowe przewidywania TGP . . . 14

13 Topological defect structure of TGP bodies **15**
13.1 Motivation: what is a "body" in TGP? . . . 15
13.2 Static spherically symmetric ODE . . . 15
13.3 Linearised far-field: oscillatory, not Yukawa . . . 15
13.4 Implication: Path B as external-source formulation . . . 16
13.5 Physical analogies . . . 16
13.6 Rotation-curve diagnostic (ex14) . . . 16
13.7 Summary of constraints on m_{sp} . . . 16

14 False vacuum structure and spontaneous nucleation in TGP **17**
14.1 The inverted potential: $V(g = 1)$ is a maximum . . . 17
14.2 Linearised field equation and oscillatory tails . . . 17
14.3 Defect energy landscape . . . 17
14.4 Metastability and the nucleation barrier . . . 18
14.5 Cosmological scenario: matter nucleation from the vacuum . . . 18
14.6 Energy scale of nucleated bodies . . . 19
14.7 Summary of vacuum structure . . . 19

15 Exact three-body N-body observability: Feynman integral vs saddle-point **19**
15.1 Setup . . . 19
15.2 Numerical results . . . 19
15.2.1 $\mathcal{R}$ as a function of $t = m_{\text{sp}}d$ . . . 19
15.2.2 Origin of the saddle-point error . . . 20
15.3 Critical coupling C_{crit} for observability . . . 20
15.4 Physical systems . . . 20
15.5 Implication: saddle-point systematically underestimates 3-body effects . . . 21

16 Three-body TGP scattering dynamics for Planck-scale objects **21**
16.1 Setup . . . 21
16.2 Static energetics . . . 21
16.3 Collision dynamics: observer displacement . . . 22
16.4 Triangle stability: three-body compactness . . . 22

16.5 Summary . . . 22

17 Three-body induced binding in TGP **23**
17.1 Static potential landscape . . . 23
17.2 Geometry: triangle vs linear configuration . . . 23
17.3 N-body binding: super-extensive deepening . . . 23
17.4 Virial theorem: exact three-body deepening (ex21) . . . 24
17.5 Summary . . . 25

18 Efimov-like three-body bound states in TGP **25**
18.1 The question . . . 25
18.2 Critical couplings . . . 25
18.3 The 3B-induced window at $m_{\rm sp} = 0.1$ . . . 26
18.4 Connection to Efimov physics . . . 26
18.5 Physical parameters of the 3B-induced state . . . 26
18.6 WKB level count: finite tower (ex24) . . . 26
18.7 Falsifiability . . . 27

19 Optimal three-body geometry **27**
19.1 Question . . . 27
19.2 Setup . . . 28
19.3 Results at $m_{\rm sp} = 0.1$, $C = 0.155$ . . . 28
19.4 Equilateral vs linear: 3B-only windows . . . 28
19.5 Isosceles deformation . . . 29
19.6 Four-body square ($N = 4$) . . . 29
19.7 Summary . . . 29

20 Quantum Efimov window: exact finite-difference result **29**
20.1 Motivation: WKB validity . . . 29
20.2 Finite-difference Hamiltonian . . . 30
20.3 Results: classical vs quantum windows . . . 30
20.4 Planck coupling is inside the quantum window . . . 31
20.5 Wavefunction at $C = 0.207$ . . . 31
20.6 Correction to ex24 WKB . . . 31
20.7 Summary . . . 31

21 Quantum Efimov window versus scalar mass $m_{\rm sp}$ **31**
21.1 Question . . . 31
21.2 Method . . . 32
21.3 Results: window versus $m_{\rm sp}$ . . . 32
21.4 Critical scalar mass range for Planck objects . . . 32
21.5 Significance: non-trivial intersection . . . 33
21.6 Comparison: classical versus quantum window membership . . . 33
21.7 Summary . . . 33

22 Polygon binding: N=3, 4, 5 bodies **34**
22.1 Setup . . . 34
22.2 Binding energies at $C = 0.155$ . . . 34
22.3 Critical coupling thresholds . . . 35
22.4 Pentagon Feynman integrals . . . 35
22.5 Comparison at $C = C_{\rm Pl}$ . . . 35
22.6 Summary . . . 36

23 Ground-state wavefunction at $C = C_{\mathrm{Pl}}$ **36**
23.1 Setup . . . 36
23.2 Ground-state energy vs m_{sp} . . . 36
23.3 Detailed wavefunction at $m_{\mathrm{sp}} = 0.1$ . . . 36
23.4 Second bound state . . . 37
23.5 Wavefunction evolution near window boundary . . . 38

24 Physical interpretation: window bounds and classical/quantum non-overlap **38**
24.1 Why does the quantum Efimov window exist for $\lambda_{\mathrm{sp}} \in (5.1,\, 13.2)\, l_{\mathrm{Pl}}$? . . . 38
24.2 Why are the classical and quantum windows non-overlapping? . . . 39
24.2.1 The role of zero-point energy . . . 39
24.2.2 Why the shift causes non-overlap . . . 39
24.2.3 Schematic . . . 40
24.2.4 Key insight: WKB invalidity as a diagnostic . . . 40
24.3 Quantum state characterisation at $C = C_{\mathrm{Pl}}$ . . . 40
24.4 Summary . . . 41

25 Triangle shape-space stability and the equilateral approximation **41**
25.1 Question . . . 41
25.2 Parameterisation of shape deformations . . . 41
25.3 Numerical results at $C = C_{\mathrm{Pl}}$, $m_{\mathrm{sp}} = 0.1$ . . . 42
25.4 Physical interpretation . . . 42
25.5 Consequence for quantum Efimov window . . . 42
25.6 Normal mode frequencies . . . 43
25.7 Shape stability vs C at quantum mean position . . . 43
25.8 Summary . . . 43

26 Open theoretical questions **44**
26.1 Q1: Yukawa source mechanism . . . 44
26.2 Q2: Dark matter and rotation curves . . . 44
26.3 Q3: Path C — quantum defects and the Yukawa derivation . . . 45
26.4 Q4: Relation to general relativity . . . 46
26.5 Summary of open questions . . . 47

27 Galactic rotation curves: TGP vs CDM and MOND **47**
27.1 Setup . . . 47
27.2 Results . . . 47
27.3 Analysis . . . 47
27.4 Conclusion . . . 48

1 Definicja i kontekst TGP

1.1 Overlap Yukawy

Potencjał Yukawy (ekranowany):

$$G_m(r) \equiv \frac{e^{-mr}}{r}, \qquad (\nabla^2 - m^2)\, G_m(|x|) = -4\pi\, \delta^{(3)}(x). \tag{1}$$

Masa ekranowania TGP (warunek próżni $\beta = \gamma$, wypr. z symetrii $\mathbb{Z}_2$):

$$m_{\mathrm{sp}} = \sqrt{3\gamma - 2\beta}\big|_{\beta=\gamma} = \sqrt{\beta}. \tag{2}$$

Dokładny trójcentrowy overlap:

$$I_Y(x_1, x_2, x_3;\, m) = \int_{\mathbb{R}^3} d^3x\; G_m(|x - x_1|)\, G_m(|x - x_2|)\, G_m(|x - x_3|). \tag{3}$$

Z niezmienniczości przesunięcia/obrotu: $I_Y = I_Y(d_{12}, d_{13}, d_{23};\, m)$.

1.2 Miejsce w TGP

Z członu Φ^4 (współczynnik multinomialny $4!/(1!1!1!1!) = 24$, po symetryzacji wkład na trip-let):

$$V_3(1, 2, 3) = -6\gamma\, C_1 C_2 C_3\, I_Y(d_{12}, d_{13}, d_{23};\, m_{\mathrm{sp}}). \tag{4}$$

2 Reprezentacja Schwingera (przestrzeń fizyczna)

Używamy reprezentacji Schwingera: $\frac{e^{-mr}}{r} = \frac{1}{\sqrt{\pi}} \int_0^\infty s^{-1/2} e^{-sr^2 - m^2/(4s)}\, ds$.

Po podstawieniu dla trzech czynników, całce gaussowskiej po x (z tożsamością kwadratową $\sum_a s_a |x - x_a|^2 = S|x - \bar{x}|^2 + \frac{\sum_{i<j} s_i s_j d_{ij}^2}{S}$, $S = \sum s_a$) i zamianie $s_i = u\alpha_i$ ($\alpha_i > 0$, $\sum \alpha_i = 1$, $u > 0$):

$$\boxed{I_Y = 2\pi \int_{\Delta_2} \frac{d\alpha_1\, d\alpha_2}{\sqrt{\alpha_1 \alpha_2 \alpha_3}}\, K_0\!\left(m\sqrt{A(\boldsymbol{\alpha}) \cdot B(\boldsymbol{\alpha})}\right)} \tag{5}$$

gdzie $A(\boldsymbol{\alpha}) = \alpha_1\alpha_2 d_{12}^2 + \alpha_1\alpha_3 d_{13}^2 + \alpha_2\alpha_3 d_{23}^2$, $B(\boldsymbol{\alpha}) = \frac{m^2}{4}\left(\frac{1}{\alpha_1} + \frac{1}{\alpha_2} + \frac{1}{\alpha_3}\right)$, z użyciem $\int_0^\infty u^{-1} e^{-uA - B/u}\, du = 2K_0(2\sqrt{AB})$.

3 Reprezentacja Feynmana (przestrzeń pędów)

Wstawiamy transformację Fouriera $G_m(r) = \int \frac{d^3k}{(2\pi)^3} \frac{4\pi}{k^2 + m^2} e^{ik\cdot r}$ dla każdego czynnika. Całka po x daje $\delta^{(3)}(k_1 + k_2 + k_3)$. Po eliminacji $k_3 = -(k_1 + k_2)$ otrzymujemy diagram *sunset* z trzema propagatorami.

Forma kwadratowa pędów:

$$M = \begin{pmatrix} \alpha_1 + \alpha_3 & \alpha_3 \\ \alpha_3 & \alpha_2 + \alpha_3 \end{pmatrix}, \qquad \Delta \equiv \det M = \alpha_1\alpha_2 + \alpha_1\alpha_3 + \alpha_2\alpha_3. \tag{6}$$

Całka gaussowska po sześciu składowych pędu, $r_{12} \cdot r_{13} = (d_{12}^2 + d_{13}^2 - d_{23}^2)/2$ i uproszczenie prefiksu $\frac{(4\pi)^3}{(2\pi)^6} \cdot \pi^3 = 1$:

$$\boxed{I_Y(d_{12}, d_{13}, d_{23};\, m) = 2 \int_{\Delta_2} d\alpha\; \Delta^{-3/2}\, K_0\!\left(m\sqrt{\frac{\alpha_2 d_{12}^2 + \alpha_1 d_{13}^2 + \alpha_3 d_{23}^2}{\Delta}}\right)} \tag{7}$$

gdzie $\Delta = \alpha_1\alpha_2 + \alpha_1\alpha_3 + \alpha_2\alpha_3$, $\alpha_3 = 1 - \alpha_1 - \alpha_2$.

Weryfikacja. Obie reprezentacje są równoważne (ta sama całka, inna zmiana zmiennych na simpleksie). W punkcie symetrycznym $\alpha_i = 1/3$: argument K_0 wynosi $md/\sqrt{1/3} = md\sqrt{3}$ w obu przypadkach dla trójkąta równobocznego.

4 Układ równań PDE

Działanie operatorem Helmholtza pod całką (3) i skorzystanie z $(\nabla^2 - m^2)G_m = -4\pi\delta^{(3)}$ daje układ:

$$(\nabla^2_{x_1} - m^2)\, I_Y = -4\pi\, G_m(d_{12})\, G_m(d_{13}), \tag{8}$$
$$(\nabla^2_{x_2} - m^2)\, I_Y = -4\pi\, G_m(d_{12})\, G_m(d_{23}), \tag{9}$$
$$(\nabla^2_{x_3} - m^2)\, I_Y = -4\pi\, G_m(d_{13})\, G_m(d_{23}). \tag{10}$$

W zmiennych względnych $\rho = m|x_1 - x_3|$, $\sigma = m|x_2 - x_3|$, $\mu = \cos\theta$, ze skalowaniem $\mathcal{I} = m\,\mathcal{J}(\rho, \sigma, \mu)$:

$$\Big[\partial^2_\rho + \tfrac{2}{\rho}\partial_\rho + \tfrac{1}{\rho^2}\Big((1-\mu^2)\partial^2_\mu - 2\mu\partial_\mu\Big) - 1\Big]\,\mathcal{J} = -4\pi\,\frac{e^{-\rho}}{\rho}\,\frac{e^{-\delta}}{\delta}, \qquad \delta = \sqrt{\rho^2 + \sigma^2 - 2\rho\sigma\mu}. \tag{11}$$

Rodziny jednoparametrowe. Dla trójkąta równobocznego ($\rho = \sigma = t$, $\mu = 1/2$): PDE się *nie* domyka do ODE dla $f_\triangle(t)$, ponieważ pochodne poprzeczne $\partial_\mu \mathcal{J}$ oceniane na lokusu równobocznym są nieznane. Ist-nienie $f_\triangle$ jako funkcji 1D wynika z symetrii, nie z ODE.

5 Redukcja do rodzin jednoparametrowych

5.1 Trójkąt równoboczny

Dla $d_{12} = d_{13} = d_{23} = d$, $t \equiv md$: $\alpha_2 d^2 + \alpha_1 d^2 + \alpha_3 d^2 = d^2$, zatem:

$$\boxed{f_\triangle(t) \equiv I_\triangle(d, m) = 2\int_{\Delta_2} d\alpha\; \Delta^{-3/2}\, K_0\!\left(\frac{t}{\sqrt{\Delta}}\right)} \tag{12}$$

Argument K_0 zależy od $\Delta(\boldsymbol{\alpha})$. To klucz do redukcji 2D→1D.

5.2 Symetryczna rodzina koliniarna

Dla $(d_{12}, d_{13}, d_{23}) = (d, 2d, d)$, $t = md$ (α_1 to centrum):

$$f_{\mathrm{col}}(t) = 2\int_{\Delta_2} d\alpha\; \Delta^{-3/2}\, K_0\!\left(t\sqrt{\frac{\alpha_2 + \alpha_3 + 4\alpha_1}{\Delta}}\right). \tag{13}$$

6 Geometria simpleksu: dokładna gęstość $p(\Delta)$

6.1 Struktura poziomic $\Delta = \mathrm{const}$

Wprowadzamy $\varepsilon_k = \alpha_k - 1/3$ (Jacobi, $\Sigma\varepsilon_k = 0$). W współrzędnych $(\varepsilon_1, \varepsilon_2)$ z $\varepsilon_3 = -\varepsilon_1 - \varepsilon_2$:

$$\Sigma\alpha_k^2 = \tfrac{1}{3} + 2r^2, \quad r^2 \equiv \varepsilon_1^2 + \varepsilon_1\varepsilon_2 + \varepsilon_2^2, \tag{14}$$
$$\Delta = \tfrac{1}{2}(1 - \Sigma\alpha_k^2) = \tfrac{1}{3} - r^2. \tag{15}$$

Forma kwadratowa r^2 ma macierz $\left[\begin{smallmatrix} 1 & 1/2 \\ 1/2 & 1 \end{smallmatrix}\right]$ z $\det = 3/4$, stąd pole elipsy $r^2 \le R^2$ w mierze $d\varepsilon_1 d\varepsilon_2$ wynosi $2\pi R^2/\sqrt{3}$.

6.2 Punkt krytyczny

Punkt śródkowy krawędzi simpleksu (np. $\alpha_3 = 0$, $\alpha_1 = \alpha_2 = 1/2$): $r^2 = 1/12$, czyli $\Delta = 1/4$. Dla $r^2 \le 1/12$ (*tj.* $\Delta \ge 1/4$) cała elipsa leży wewnątrz simpleksu. Symetria S_3 gwarantuje, że wszystkie trzy boki są równoodległe w metryce r^2.

6.3 Pole czapek i gęstość

Lemat 1 (Gęstość miary simpleksowej)**.** *Dla miary uniformowej $d\alpha_1\, d\alpha_2$ na 2-simpleksie, gęstość $p(\Delta_0) \equiv -d/d\Delta_0[Area\{\Delta \geq \Delta_0\}]$ wynosi:*

$$\boxed{p(\Delta) = \begin{cases} \dfrac{2\pi}{\sqrt{3}} & \Delta \in \left[\dfrac{1}{4}, \dfrac{1}{3}\right], \\ \dfrac{2}{\sqrt{3}}\left[3\arcsin\left(\dfrac{1}{2\sqrt{1-3\Delta}}\right) - \dfrac{\pi}{2}\right] & \Delta \in \left[0, \dfrac{1}{4}\right]. \end{cases}} \tag{16}$$

Proof. Dla $\Delta \geq 1/4$: pełna elipsa $r^2 \leq 1/3 - \Delta$ leży wewnątrz simpleksu. Area $= 2\pi(1/3-\Delta)/\sqrt{3}$, stąd $p = 2\pi/\sqrt{3}$.

Dla $\Delta < 1/4$: elipsa jest obcięta przez trzy boki (symetria S_3; każdy bok obcina równe czapki). Wzmiast $w = \varepsilon_1 + \varepsilon_2$, $z = \varepsilon_1 - \varepsilon_2$, pól-osie elipsy w (w, z) wynoszą $a_w = 2R/\sqrt{3}$, $a_z = 2R$. Pole czapki $w \geq 1/3$ (odpowiadającej bokowi $\alpha_3 = 0$):

$$\text{Pole}_{\text{czap}} = \tfrac{2R^2}{\sqrt{3}}\left[\tfrac{\pi}{2} - \arcsin\xi - \xi\sqrt{1-\xi^2}\right], \qquad \xi = \tfrac{1}{2R\sqrt{3}}.$$

Pole obszaru $\Delta \geq \Delta_0$: $\text{Area}(R^2) = \frac{2R^2}{\sqrt{3}}\left[3\arcsin\xi + 3\xi\sqrt{1-\xi^2} - \frac{\pi}{2}\right]$. Różniczkując po R^2 (z uwzględnieniem $\xi = 1/(2R\sqrt{3})$, $d\xi/d(R^2) = -6\xi^3$, $d(3\arcsin\xi + \ldots)/d\xi = 6\sqrt{1-\xi^2}$):

$$p(\Delta) = \frac{d\,\text{Area}}{d(R^2)} = \frac{2}{\sqrt{3}}\left[3\arcsin\xi - \frac{\pi}{2}\right], \quad \xi = \frac{1}{2\sqrt{1-3\Delta}}. \qquad \square$$

Weryfikacja numeryczna. Monte Carlo (5×10^6 próbek) potwierdza formułę z błędem względnym $< 1\%$; całkowanie numeryczne daje $\int_0^{1/3} p(\Delta)\, d\Delta = 1/2$ (pole simpleksu) z precyzją $< 10^{-9}$.

7 Redukcja 2D→1D i kluczowa tożsamość

7.1 Dokładna reprezentacja 1D

Ponieważ integrand $f_\triangle$ zależy wyłącznie od Δ:

$$\boxed{f_\triangle(t) = 2\int_0^{1/3} p(\Delta)\, \Delta^{-3/2}\, K_0\left(\frac{t}{\sqrt{\Delta}}\right) d\Delta} \tag{17}$$

z dokładnym $p(\Delta)$ z Lematu 1. Jest to dokładna całka 1D (zamiast 2D).

7.2 Kluczowa tożsamość: $C_1 = 2\pi$

Twierdzenie 2 (Tożsamość simpleksowa)**.**

$$\boxed{\int_{\Delta_2} (\alpha_1\alpha_2 + \alpha_1\alpha_3 + \alpha_2\alpha_3)^{-3/2}\, d\alpha_1\, d\alpha_2 = 2\pi.} \tag{18}$$

Proof. Rozbijamy $C_1 = \int_0^{1/3} p(\Delta)\Delta^{-3/2}\, d\Delta$ na dwa przedziały.

Część wysoka $[\frac{1}{4}, \frac{1}{3}]$: z $p = 2\pi/\sqrt{3}$:

$$\frac{2\pi}{\sqrt{3}}\left[-2\Delta^{-1/2}\right]_{1/4}^{1/3} = \frac{2\pi}{\sqrt{3}}(4 - 2\sqrt{3}) = 6\pi - \frac{8\pi}{\sqrt{3}}.$$

Część niska $[0, \frac{1}{4}]$: podstawiamy $u = \cos\psi$ ($u \in [0, \frac{\sqrt{3}}{2}]$ odpowiada $\psi \in [0, \frac{\pi}{3}]$), korzystając z $3\arcsin u - \pi/2 = \pi - 3\psi$ i $4u^2 - 1 = 1 + 2\cos 2\psi$:

$$I_{\text{low}} = 8\int_0^{\pi/3} \frac{(\pi - 3\psi)\sin\psi}{(3 - 4\sin^2\psi)^{3/2}} d\psi.$$

Całkowanie przez części z $g(\psi) = \frac{\cos\psi}{\sqrt{1+2\cos 2\psi}}$ (weryfikacja: $dg/d\psi = -\sin\psi/(3-4\sin^2\psi)^{3/2}$):

$$I_{\text{low}} = 8\left[-\pi/\sqrt{3} + 3\int_0^{\pi/3} \frac{\cos\psi}{\sqrt{1+2\cos 2\psi}}\, d\psi\right].$$

Całka rezydualna: podstawiamy $t = \sin\psi$, $dt = \cos\psi\, d\psi$: $\int_0^{\sqrt{3}/2} \frac{dt}{\sqrt{3-4t^2}}$. Dalej $t = \frac{\sqrt{3}}{2}\sin\phi$:

$$\int_0^{\pi/2} d\phi = \frac{\pi}{2}, \qquad \text{zatem} \int_0^{\pi/3} \frac{\cos\psi\, d\psi}{\sqrt{1+2\cos 2\psi}} = \frac{\pi}{4}.$$

Stąd $I_{\text{low}} = 8(-\pi/\sqrt{3} + 3\pi/4) = 6\pi - 8\pi/\sqrt{3}$.

Suma: $C_1 = (6\pi - 8\pi/\sqrt{3}) + (8\pi/\sqrt{3} - 4\pi) = 2\pi$. □

Weryfikacja. Quadratura numeryczna daje $C_1 = 6.28318531$ z błędem $< 10^{-9}$, zgodna z $2\pi = 6.28318530\ldots$

8 Asympotyki dokładne

8.1 Granica Yukawa ($t \to \infty$): metoda siodłowa

Twierdzenie 3 (Asymptotyka Yukawa)**.**

$$\boxed{f_\triangle(t) \underset{t\to\infty}{\sim} \frac{8\pi\sqrt{\pi}}{\sqrt{2}\cdot 3^{3/4}} \frac{e^{-\sqrt{3}\,t}}{t^{3/2}}.} \tag{19}$$

Proof. Argument $K_0(t/\sqrt{\Delta})$ jest minimalizowany przy $\Delta = \Delta_{\max} = 1/3$ (centrum sympleksu, trójkąt równoboczny — głębokość siodła $t\sqrt{3}$). Używamy $K_0(z) \approx \sqrt{\pi/(2z)}\, e^{-z}$ dla dużych z.

Rozwinięcie wokół centrum: $\Delta \approx 1/3 - r^2$, $t/\sqrt{\Delta} \approx t\sqrt{3}(1+3r^2/2)$. Całkowanie gaussowskie po r z miarą $(2/\sqrt{3})\, 2\pi r\, dr$ (pełna elipsa spełnia warunek $r^2 \le 1/12$, ale siodło dominuje dla $t \gg 1$):

$$\int_0^\infty r\, e^{-3\sqrt{3}\,t\, r^2/2}\, dr = \frac{1}{3\sqrt{3}\,t}.$$

Zbierając: $f_\triangle(t) \approx 2\cdot 3^{3/2}\cdot\sqrt{\pi/(2\sqrt{3}\,t)}\cdot e^{-\sqrt{3}\,t}\cdot\frac{4\pi}{\sqrt{3}}\cdot\frac{1}{3\sqrt{3}\,t}$, co daje (19) po uproszczeniu. □

Interpretacja. Efektywna masa $m_{\text{eff}} = m\sqrt{3}$ pochodzi z faktoru $1/\sqrt{\Delta_{\max}} = \sqrt{3}$. Układ "widzi" całkę przez punkt równoboczny.

8.2 Granica Coulomba ($t \to 0$): rozw-inięcie logarytmiczne

Twierdzenie 4 (Asymptotyka Coulomba)**.**

$$\boxed{f_\triangle(t) \underset{t\to 0}{\sim} -4\pi\ln t + C_0,} \tag{20}$$

gdzie C_0 jest skończoną stałą (obliczalną numerycznie).

Proof. Dla $z \to 0$: $K_0(z) = -\ln(z/2) - \gamma_E + O(z^2\ln z)$, zatem:

$$f_\triangle(t) \approx -2\ln(t)\underbrace{\int_0^{1/3} p(\Delta)\Delta^{-3/2}\, d\Delta}_{C_1=2\pi} + 2\int_0^{1/3} p(\Delta)\Delta^{-3/2}(\ln 2 + \tfrac{1}{2}\ln\Delta - \gamma_E)\, d\Delta.$$

Całka główna daje $-4\pi\ln t$. Zbieżność: blisko wierzchołków $p(\Delta) \sim 3\Delta$, stąd $p(\Delta)\Delta^{-3/2} \sim 3\Delta^{-1/2}$ — zbieżne. □

Fizyczna interpretacja. Czysta całka Coulomba ($m = 0$): $\int d^3x/(r_1 r_2 r_3)$ jest *podczerwie-nie rozbieżna* (logarytmicznie: $\int_R^\infty r^2/r^3\, dr = \ln(\infty/R)$). Masa Yukawa m reguluje podczerwoność. Spójność z $C_1 = 2\pi$ jest dokładna.

9 Nieelementarność $f_\triangle(t)$

Twierdzenie 5 (Nieelementarność)**.** *$f_\triangle(t)$ nie może być zapisana jako skończona kombinacja $K_\nu(a_k t)$ z żadnymi stałymi a_k, ν.*

Proof. Transformata Mellina:

$$\mathcal{M}[f_\triangle](s) = \sqrt{\pi}\,\Gamma\left(\frac{s}{2}\right)^2 \underbrace{\int_{\Delta_2} \Delta^{s/2-3/2}\, d\alpha}_{\mathcal{B}(s)}. \tag{21}$$

Gdyby $f_\triangle = \sum_k c_k K_\nu(a_k t)$, to $\mathcal{M}[f_\triangle]$ byłoby iloczynem funkcji Gamma bez czynnika $\mathcal{B}(s)$. Tymczasem $\mathcal{B}(s) = \int_0^{1/3} p(\Delta)\Delta^{s/2-3/2}\, d\Delta$ jest inna funkcją s niż dowolny iloczyn $\Gamma((s \pm \nu)/2)$: posiada logarytmiczne osobliwości od krawędzi simpleksu, które nie redukują się do biegunów Gamma. □

$f_\triangle(t)$ jest nową, nieelementarną **funkcją specjalną TGP**: dokładnie określoną przez (17), a nie redukującą się do standardowych funkcji Bessela.

10 Podsumowanie wyników

Wynik	Formuła	Status
Repr. 3D $\to$ 2D Schwinger	(5)	Dokładny
Repr. 3D $\to$ 2D Feynman	(7)	Dokładny
Układ PDE Helmholtza	(8)–(10)	Dokładny
Gęstość $p(\Delta)$	(16)	Dokładny, zamknięty
Redukcja 2D $\to$ 1D	(17)	Dokładny
Tożsamość $C_1 = 2\pi$	(18)	Udowodniony
Asymptotyka $t \to \infty$	(19)	Dokładny
Asymptotyka $t \to 0$	(20)	Dokładny
Forma zamknięta $f_\triangle$	—	Nie istnieje (21)

10.1 Strategia dla TGP runtime

1. **Definicja analityczna**: $f_\triangle(t) = 2\int_0^{1/3} p(\Delta)\Delta^{-3/2} K_0(t/\sqrt{\Delta})\, d\Delta$.
2. **Reżym Yukawa** ($md \gg 1$, typ-owe jednostki kodu): $f_\triangle \approx C_\infty\, t^{-3/2} e^{-\sqrt{3}\,t}$.
3. **Reżym Coulomba** ($md \ll 1$, fizyczny $\gamma \sim 10^{-52}$): $f_\triangle \approx -4\pi \ln(md) + C_0$.
4. **Tabulacja numeryczna**: grid $t \in [t_{\min}, t_{\max}]$, interpolacja Chebysheva lub splajnami — precyzja maszynowa przy koszcie jednej ewaluacji zamiast 2D całki.

10.2 Wkład trzyciałowy w TGP

Na mocy (4) i powyższych wyników:

- Dla równobocznego układu równowagi przy d_{well}: $|V_3/V_2| \approx 0.086\%$ (dla $\beta/C = 5$, reżym Yukawa).
- Wkład trzyciałowy jest zawsze perturbatywny względem dwuciałowego.
- Korekta do równowagi: $\delta d \approx -6\gamma C^3 f'_{\triangle}(t_-)/[m_{\text{sp}}(V_2)''(d_-)]$.

11 Non-stationary TGP equations of motion

11.1 Starting point: the time-dependent field equation

The TGP field $\Phi(x,t)$ obeys the Euler–Lagrange equation derived from the action $S = \int dt\, E[\Phi]$ extended with a kinetic term. Including the standard minimal kinetic term for the scalar field gives the field equation

$$\Box\,\Phi + V'(\Phi;\Phi_0) = \sum_{i=1}^{N} C_i\,\delta^{(3)}(x - x_i(t)), \tag{22}$$

where $\Box = \partial_t^2 - \nabla^2$ is the d'Alembertian and the source $C_i\,\delta^{(3)}$ models each body as a point source of field strength C_i. The self-interaction potential is

$$V(\Phi;\Phi_0) = \frac{\beta}{3\Phi_0}\,\Phi^3 - \frac{\gamma}{4\Phi_0^2}\,\Phi^4. \tag{23}$$

Vacuum condition. The field sits in the non-trivial vacuum $\Phi_0 > 0$ with $\beta = \gamma$ (exact TGP vacuum condition), giving a screening mass

$$m_{\text{sp}}^2 = V''(\Phi_0;\Phi_0)\big|_{\beta=\gamma} = \beta. \tag{24}$$

11.2 Linearisation around the vacuum

Write $\Phi(x,t) = \Phi_0\big(1 + \delta(x,t)\big)$ with $\delta \ll 1$. Expanding V' to linear order in δ:

$$V'(\Phi) \approx V'(\Phi_0) + V''(\Phi_0)\,\Phi_0\,\delta = m_{\text{sp}}^2\,\Phi_0\,\delta, \tag{25}$$

where we used $V'(\Phi_0) = 0$ (vacuum equation) and $V''(\Phi_0) = m_{\text{sp}}^2$. Hence

$$\boxed{(\Box - m_{\text{sp}}^2)\delta(x,t) = \frac{1}{\Phi_0}\sum_{i=1}^{N} C_i\,\delta^{(3)}(x - x_i(t)).} \tag{26}$$

This is the *Klein–Gordon equation with time-dependent point sources.*

11.3 Retarded solution: the radiation field

The causal solution of (26) is given by convolution with the retarded Green function G_{ret}:

$$(\Box - m_{\text{sp}}^2)G_{\text{ret}}(x,t;x',t') = \delta^{(3)}(x - x')\,\delta(t - t'), \tag{27}$$

whose explicit form in 3+1 dimensions is (Yukawa propagator):

$$G_{\text{ret}}(x,t;x',t') = -\frac{1}{4\pi|x - x'|}\left[\delta\big(t - t' - |x - x'|\big) - m_{\text{sp}}\,\frac{J_1\big(m_{\text{sp}}\sqrt{(t-t')^2 - |x-x'|^2}\big)}{\sqrt{(t-t')^2 - |x-x'|^2}}\,\theta\big(t - t' - |x - x'|\big)\right]. \tag{28}$$

Therefore the exact linearised field is

$$\delta(x,t) = \frac{1}{\Phi_0}\sum_{i=1}^{N} C_i \int_{-\infty}^{t} dt'\; G_{\text{ret}}\big(x,t;\,x_i(t'),\,t'\big). \tag{29}$$

Physical interpretation. The key difference from the quasi-static case is that:

- each body emits a *wave* that propagates at speed $c = 1$ (in natural units);
- the wave decays exponentially on length scale $m_{\rm sp}^{-1}$ *in the rest frame*;
- the field at body j at time t depends on the *retarded position* $x_i(t - |x_j - x_i|)$, not the current position.

11.4 Non-stationary pairwise interaction

For two slowly moving bodies (velocities $v \ll 1$), the retarded position can be expanded:

$$x_i(t - d_{ij}/c) = x_i(t) - \dot{x}_i(t)\, d_{ij}/c + \tfrac{1}{2}\ddot{x}_i(t)\,(d_{ij}/c)^2 + \cdots \tag{30}$$

This gives corrections to the static Yukawa potential of order v/c.

The leading-order retardation correction to the pairwise energy is (in the $m_{\rm sp} d \ll 1$ limit):

$$\delta V_2^{(v/c)} = \frac{4\pi C_i C_j}{d_{ij}} \cdot v_i \cdot v_j / c^2 + O(v^2/c^2), \tag{31}$$

analogous to the Darwin term in electrodynamics.

11.5 Radiation: power loss

For an accelerating source (e.g. body i with acceleration $\ddot{x}_i$), the scalar field radiates energy. In the far field ($r \to \infty$, $mr \ll 1$), the radiated power is (Larmor formula for massive scalar field):

$$P_{\rm rad,i} = \frac{C_i^2}{6\pi}\,|\ddot{x}_i|^2 \cdot \begin{cases} 1 & m_{\rm sp} = 0 \text{ (massless)}, \\ 1 - \dfrac{3m_{\rm sp}^2}{4\omega^2} + \cdots & \omega \gg m_{\rm sp}, \end{cases} \tag{32}$$

where ω is the orbital frequency. For TGP screening masses $m_{\rm sp} = \sqrt{\beta}$ and typical orbital frequencies, this quantifies the rate of energy loss to scalar radiation.

11.6 Non-stationary N-body equations of motion

Including retardation and radiation reaction, the equation of motion for body i becomes:

$$\boxed{C_i\,\ddot{x}_i(t) = \underbrace{F_i^{(2,\rm quasi)}(x_i(t), \{x_j(t)\})}_{\text{quasi-static pairwise}} + \underbrace{F_i^{(3,\rm quasi)}(\{x_j(t)\})}_{\text{quasi-static 3-body}} + \underbrace{\delta F_i^{(v/c)}(\{x_j(t), \dot{x}_j(t)\})}_{\text{velocity corrections}} + \underbrace{\delta F_i^{(\rm RR)}(t)}_{\text{radiation reaction}} + \cdots} \tag{33}$$

The terms are:

1. *Quasi-static pairwise force* (exact, closed form):

$$F_i^{(2,\rm quasi)} = -\nabla_{x_i} \sum_{j \neq i} V_2(d_{ij}), \quad V_2(d) = -\frac{4\pi C_i C_j}{d} + \frac{8\pi\beta C_i C_j}{d^2} - \frac{12\pi\gamma C_i C_j (C_i + C_j)}{d^3}. \tag{34}$$

2. *Quasi-static 3-body force* (exact via Feynman integral):

$$F_i^{(3,\rm quasi)} = 6\gamma \sum_{\substack{j<k \\ j\neq i, k\neq i}} C_i C_j C_k \left[\frac{\partial I_Y}{\partial d_{ij}}\,\hat{e}_{ij} + \frac{\partial I_Y}{\partial d_{ik}}\,\hat{e}_{ik} \right], \tag{35}$$

where $\partial I_Y / \partial d_{ij}$ is computed exactly via

$$\frac{\partial I_Y}{\partial d_{12}} = -2m\, d_{12} \int_{\Delta_2} \frac{\alpha_2}{\Delta^2 \sqrt{Q}}\, K_1\!\left(m\sqrt{Q/\Delta}\right) d\alpha, \tag{36}$$

with $Q = \alpha_2 d_{12}^2 + \alpha_1 d_{13}^2 + \alpha_3 d_{23}^2$ and $\Delta = \alpha_1\alpha_2 + \alpha_1\alpha_3 + \alpha_2\alpha_3$ (implemented in `three_body_force_exact.py`)

3. *Velocity corrections* (v/c order): retardation shifts, Darwin term, Römer delay in the effective potential. Suppressed by $(v/c)^2 \sim (v_{\rm orb}/1)^2$.

4. *Radiation reaction* (Abraham–Lorentz–Dirac term for a scalar):

$$\delta F_i^{(\rm RR)} = \frac{C_i^2}{6\pi}\, \dddot{x}_i(t). \tag{37}$$

Suppressed by C_i^2 (source strength squared) and relevant only when the orbital period is comparable to the radiation timescale.

11.7 Hierarchy of approximations

Level	Approximation kept	Error scale
0	Full field equation (22) (nonlinear, retarded)	exact
1	Linearised + retarded (26)	$O(\delta^2)$
2	Quasi-static: drop $\partial_t^2\delta$	$O(v^2/c^2)$
3	Yukawa superposition: $\delta = \sum \delta_i$	$O(\delta^2)$
4	Truncate at 3-body: drop $V_4, V_5, \ldots$	$O(C^4)$
5	Exact 3-body force via (36)	0 (within level 4)

What `dynamics_v2.py` implements. Currently levels 0–4 are applied and, for the 3-body term, an additional saddle-point approximation $I_Y \approx (2\pi)^{3/2} m^{-2} e^{-ms}/s$ is used (roughly 5–30% error in the 3-body force).

What `three_body_force_exact.py` provides. The exact 3-body force at level 5, replacing the saddle-point approximation with the exact Feynman 2D integral, at the cost of $O(n_{\rm quad}^2)$ function evaluations per triplet per time step.

11.8 Retardation time scale

The quasi-static approximation (level 2) fails when the retardation delay $\tau = d_{ij}/c$ is comparable to the orbital period $T_{\rm orb}$. The condition for validity is

$$\frac{d_{ij}}{c\,T_{\rm orb}} = \frac{v_{\rm orb}}{c} \ll 1. \tag{38}$$

For TGP bodies moving at non-relativistic speeds ($v_{\rm orb}/c \sim 10^{-3}$–$10^{-6}$ in astrophysical contexts), the quasi-static limit is excellent. Post-Newtonian corrections enter at relative order $v^2/c^2 \sim 10^{-6}$–10^{-12} and are negligible for most TGP applications.

11.9 Non-stationary 3-body effects: summary

The genuine novelty of the non-stationary TGP equations (compared to Newtonian gravity or standard Yukawa) lies in two effects:

1. *Non-trivial 3-body interaction* (already present in the quasi-static limit): the term $V_3 \propto \gamma\, C_1 C_2 C_3\, I_Y$ has no Newtonian analogue. It is attractive, breaks the equivalence with gravity ($C_i \neq$ mass), and is dominant at intermediate separations $m_{\rm sp} d \sim 1$.

2. *Modified radiation*: the massive scalar ($m_{\rm sp} > 0$) radiates less efficiently than a massless field. The radiation spectrum has a low-frequency cutoff at $\omega = m_{\rm sp}$, unlike electromagnetic or gravitational radiation. This means short-period orbits are more stable in TGP than in GR.

The leading observable signatures of non-stationary effects in TGP would therefore be:

- orbital decay rate suppressed below the gravitational-wave analogue by a factor of $(m_{\rm sp}\, d_{\rm orb})^2$ relative to GR,
- three-body secular effects (Kozai–Lidov-type resonances modified by the V_3 coupling) measurable at the $C^3/(C^2) = C$ level compared to pairwise dynamics.

12 Konkretne przewidywania TGP dla układów wielociałowych

Niniejsza sekcja zawiera ilościowe wyniki uzyskane za pomocą dokładnej całki Feynmana (moduł `three_body_force_exact.py`). Wszystkie poprzednie obliczenia trójciałowe używały przybliżenia siodłowego, które ma błąd 160–770% — wyniki poniżej są pierwszymi *dokładnymi* przewidywaniami TGP dla tych zagadnień.

12.1 Kiedy efekty trójciałowe dominują? Liczba krytyczna $N_{\rm crit}$

Schemat. Rozważ N identycznych ciał o sile źródłowej C, przy średniej separacji d (bezwymiarowo $t = m_{\rm sp}\, d$). Energie wielociałowe skalują się jak:

$$V_2^{\rm total} = \binom{N}{2} V_2(d) \sim N^2\, V_2, \tag{39}$$

$$V_3^{\rm total} = \binom{N}{3} V_3(d,d,d) \sim N^3\, V_3, \tag{40}$$

zatem stosunek

$$\frac{|V_3^{\rm total}|}{|V_2^{\rm total}|} = \frac{N-2}{3} \cdot \frac{|V_3^{(1\ \rm trojka)}|}{|V_2^{(1\ \rm para)}|} \xrightarrow{N\to\infty} \infty. \tag{41}$$

Liczba krytyczna, przy której energia trójciałowa zrównuje się z parową:

$$\boxed{N_{\rm crit} = \frac{3}{\left|\, V_3^{(1)}/V_2^{(1)} \,\right|} + 2,} \tag{42}$$

gdzie $V_3^{(1)}/V_2^{(1)}$ to stosunek dla jednej trójki/pary, obliczony dokładnie z całki Feynmana.

Wyniki numeryczne. Tabela 1 zestawia $|V_3/V_2|$ na jedną trójkę (trójkąt równoboczny) i odpowiadające $N_{\rm crit}$:

Table 1: Stosunek $|V_3/V_2|$ (jedna trójka, trójkąt równoboczny) i liczba krytyczna $N_{\rm crit}$. Parametry: $\beta = \gamma = 1$.

$t = m_{\rm sp}\, d$	$C = 0.05$	$C = 0.10$	$C = 0.15$	$C = 0.20$
1.0	1.0% ($N_{\rm crit} = 298$)	3.5% (86)	21% (**16**)	14% (23)
1.5	1.6% (191)	9.5% (33)	14% (23)	6.3% (49)
2.0	1.8% (166)	1.8% (166)	1.8% (166)	1.8% (166)
2.5	0.23% (1295)	0.39% (773)	0.50% (599)	0.59% (513)
3.0	0.07% (4581)	0.12% (2499)	0.17% (1805)	0.21% (1458)

Główny wniosek.

> Dla $C = 0.15$, $t = m_{\rm sp}\, d = 1.0$ (typowy reżim Yukawa): $N_{\rm crit} = \mathbf{16}$. Skupisko zaledwie 16 ciał TGP wykazuje energię trójciałową porównywalną z parową.

Jest to przewidywanie *jakościowo różne* od grawitacji Newtonowskiej (gdzie nie ma trójciałowego członu) i od ogólnej teorii względności (gdzie trójciałowe poprawki post-Newtonowskie są rzędu v^4/c^4, a nie C jak w TGP).

12.2 Promieniowanie skalarne TGP: odcięcie częstotliwości

Mechanizm. Pole TGP jest masywne: $m_{\rm sp} = \sqrt{\beta} > 0$. Dispersion relation dla fal skalarnych:

$$\omega^2 = k^2 + m_{\rm sp}^2 \geq m_{\rm sp}^2. \tag{43}$$

Źródło oscylujące z częstotliwością $\omega < m_{\rm sp}$ nie może wzbudzić propagujących fal — emituje tylko ewanescentne pole bliskie. Promieniowanie (eksport energii do nieskończoności) jest możliwe tylko dla $\omega > m_{\rm sp}$.

Moc promieniowania. Dla ciała o przyspieszeniu $a(t) = A\omega^2 \cos(\omega t)$ moc promieniowania skalarnego wynosi:

$$P_{\rm TGP}(\omega) = \frac{C^2}{6\pi} A^2 \omega^4 \sqrt{1 - \frac{m_{\rm sp}^2}{\omega^2}}\, \theta(\omega - m_{\rm sp}). \tag{44}$$

Porównanie z grawitacją.

- Grawitacja Newtonowska: $P_{\rm GW} \sim G m^2 A^2 \omega^4$ (zawsze).
- TGP: $P_{\rm TGP} = 0$ dla $\omega < m_{\rm sp}$ (orbity długookresowe *dokładnie* stabilne).
- Dla $\omega \gg m_{\rm sp}$: $P_{\rm TGP} \to P_{\rm masless}$ (jak bezmasowe pole skalarne).

Table 2: Moc promieniowania TGP (44) vs bezmasowe pole, dla $C = 1$, $A = 1$, $m_{\rm sp} = 1$.

ω	$P_{\rm TGP}$	$P_{\rm bezm.}$	supresja
$0.5 < m_{\rm sp}$	0	3.4×10^{-3}	**0%**
$0.8 < m_{\rm sp}$	0	2.1×10^{-2}	**0%**
$1.0 = m_{\rm sp}$	4.4×10^{-3}	5.4×10^{-2}	8%
1.1	3.3×10^{-2}	7.8×10^{-2}	42%
1.5	2.0×10^{-1}	2.7×10^{-1}	75%
2.0	7.3×10^{-1}	8.4×10^{-1}	87%
5.0	3.2×10^{1}	3.3×10^{1}	98%

Tabela supresji mocy.

Konsekwencja astrofizyczna. Przy $m_{\rm sp} = \sqrt{\beta}$, orbita o okresie T nie promieniuje jeśli:

$$T > T_{\rm crit} = \frac{2\pi}{m_{\rm sp}} = \frac{2\pi}{\sqrt{\beta}}. \tag{45}$$

Dla $\beta = 1$: $T_{\rm crit} \approx 6.28$ (w jednostkach TGP). Układy o długich okresach orbitalnych (galaktyki, gromady) są *dokładnie* stabilne w TGP — nie tracą energii na promieniowanie.

12.3 Podsumowanie: trzy nowe przewidywania TGP

1. **Próg wielociałowy** $N_{\rm crit}$ (wynik dokładny): Skupiska $N > N_{\rm crit}$ ciał TGP wykazują energię trójciałową porównywalną lub większą od parowej. Dla typowych parametrów ($C \sim 0.15$, $t \sim 1$):

$$N_{\rm crit} \approx 16 \quad (C = 0.15,\ t = 1). \tag{46}$$

   Brak odpowiednika w Newtonie i GR na tym poziomie przybliżenia.

2. **Zależność oddziaływania trójciałowego od kształtu**: Dwa trójkąty o tym samym obwodzie dają różne I_Y. Trójkąt zdegenerowany (bardzo skośny) ma I_Y o 19% większe niż równoboczny przy tym samym obwodzie. Przybliżenie siodłowe nie widziało tego efektu w ogóle (dawało identyczny wynik).

3. **Odcięcie promieniowania skalarnego**: Orbity z $\omega < m_{\rm sp}$ nie promieniują w TGP. Długookresowe układy (gromady, galaktyki) są dokładnie stabilne — inaczej niż w GR gdzie każdy układ traci energię przez promieniowanie grawitacyjne.

Status obliczeń. Wszystkie trzy wyniki uzyskano z dokładnej całki Feynmana (7) i dokładnego gradientu (36), bez przybliżeń. Poprzednie obliczenia trójciałowe TGP używały wzoru siodłowego z błędem 160–770% i brakiem zależności od kształtu.

13 Topological defect structure of TGP bodies

13.1 Motivation: what is a "body" in TGP?

In the N-body formulation of TGP, each body is modelled as a *point source* coupling to the scalar field Φ. This is analogous to classical electrodynamics, where electrons are treated as point charges without deriving their internal structure. Scenario C (the most ambitious interpretation) asks whether a *body* in TGP is actually a **topological defect** of the Φ field: a localised region where Φ departs from the vacuum value Φ_0 and approaches $\Phi \to 0$ (the "absolute nothing" sector $\mathcal{S}_0$).

13.2 Static spherically symmetric ODE

We seek a static, spherically symmetric profile $\Phi(r) = \Phi_0\, g(r)$, with $g(r) \to 1$ as $r \to \infty$ (vacuum) and $g(0) = g_0 < 1$ (defect core). With the kinetic function $f(g) = 1 + 2\alpha \ln g$ (the TGP nonlinear kinetic term, $\alpha = 2$) and the potential $V(\Phi_0 g)/\Phi_0^2 = \frac{\beta}{3} g^3 - \frac{\gamma}{4} g^4$, the Euler–Lagrange equation gives

$$f(g)\left[g'' + \frac{2}{r} g'\right] + \frac{\alpha}{g}(g')^2 = g^2(1-g), \qquad \beta = \gamma = 1. \tag{47}$$

The right-hand side $g^2(1-g) > 0$ for $g \in (0,1)$, so $g''(0) > 0$: solutions rise from g_0 toward $g = 1$. The **ghost boundary**

$$g^* = \exp\!\left(-\tfrac{1}{2\alpha}\right) = e^{-1/4} \approx 0.7788 \tag{48}$$

marks the minimum allowed core depth: for $g_0 < g^*$, the kinetic term $f(g) < 0$, rendering the theory unphysical (negative kinetic energy). The physical domain is $g_0 \in (g^*, 1)$.

13.3 Linearised far-field: oscillatory, not Yukawa

Setting $g = 1 - \delta$ with $|\delta| \ll 1$ and linearising Eq. (47):

$$\delta'' + \frac{2}{r}\delta' = -\delta \quad \Longrightarrow \quad (\nabla^2 + 1)\,\delta = 0. \tag{49}$$

The general solution regular at the origin is

$$\delta(r) = A\,\frac{\sin r}{r} + B\,\frac{\cos r}{r}. \tag{50}$$

This is *oscillatory* ($\sin r/r$), **not** a Yukawa tail (e^{-r}/r). The sign of the mass-squared term in the linearised field equation is $+1$ (from $V''(\Phi_0) = -\beta < 0$ evaluated at the vacuum), leading to $(\nabla^2 + m^2)\delta = 0$ with $m^2 = \beta > 0$. A Yukawa tail would require $(\nabla^2 - m^2)\delta = 0$ with $m^2 > 0$, which arises only at a *minimum* of V.

Key result (ex11): Numerical integration confirms that all solutions with $g_0 \in (g^*, 1)$ exhibit oscillatory far-field tails with amplitude $A \approx 0.05$–0.11 and residual from a Yukawa fit $\gtrsim 0.003$. The full nonlinear $f(g) \neq 1$ ODE does not change this conclusion.

13.4 Implication: Path B as external-source formulation

Since the free-field TGP vacuum emits oscillatory rather than Yukawa radiation, the Yukawa profile $\Phi_i(r) = C_i e^{-m_{sp} r}/r$ *cannot* be derived from the vacuum sector of the theory. It must be imposed as an external source:

$$(\nabla^2 - m_{sp}^2)\,\Phi = -4\pi\, C_i\, \delta^{(3)}(r - r_i), \tag{51}$$

whose solution is the Yukawa propagator. This is **Path B**: the Yukawa ansatz is a *phenomenological input*, not a derivable consequence of the TGP field equations.

13.5 Physical analogies

1. **Classical electrodynamics:** the point-charge model $(\nabla^2)\Phi = -4\pi q\, \delta^3(r)$ does not derive the electron radius; it is an axiomatic external source. TGP Path B is structurally identical.

2. **Chameleon/scalar-tensor gravity:** in many modified gravity theories, the scalar field couples to matter through an external source term; the vacuum propagator alone does not determine the particle profile.

3. **Fuzzy dark matter (FDM):** if Φ acquires a non-trivial background condensate, its density profile could provide dark-matter-like contributions. However, the FDM mass scale $m_{boson} \sim 10^{-22}$ eV requires $m_{sp} \approx 8 \times 10^{-51}$ [Planck], which is *below* the radiation-suppression threshold $m_{sp} > 3.4 \times 10^{-41}$ derived in [**?**] (see also the N-body predictions section).

13.6 Rotation-curve diagnostic (ex14)

A systematic comparison of TGP Path B Yukawa against the NGC 3198 rotation curve shows that the Yukawa modification $G_{\text{eff}}(r) = G \exp(-r/\lambda)$ *decreases* the gravitational force at large r, giving velocities *below* the Newton-only prediction. The flat observed rotation curve ($v \approx 147\,\text{km/s}$ at $r = 20\,\text{kpc}$) requires additional dark mass; TGP Yukawa alone makes the discrepancy *worse*, not better. Specifically:

Model	$v_{\text{rot}}(20\,\text{kpc})$	χ^2/N
Observed	147 km/s	—
Newton (baryons)	71 km/s	198
CDM (ISO halo)	197 km/s	84
MOND	≈ 138 km/s	58
TGP $\lambda = 100$ kpc	64 km/s	220
TGP $\lambda = 10$ kpc	26 km/s	387

Conclusion: TGP (Path B) is not a dark-matter replacement. It must coexist with standard CDM, contributing three-body corrections to the internal structure of dark-matter halos and producing an observable $G_{\text{eff}}(r)$ deviation at scales $r \sim \lambda$.

13.7 Summary of constraints on m_{sp}

Constraint	Bound on m_{sp} [Planck]	Source
Radiation suppr. (LIGO)	$m_{sp} > 3.4 \times 10^{-41}$	ex13
Gravity range >50 AU	$m_{sp} < 1.1 \times 10^{-46}$	ex13
FDM mass scale	$m_{sp} \approx 8 \times 10^{-51}$	ex14

The FDM value falls *below* the radiation-suppression bound, ruling out TGP as an ultralight FDM candidate unless the radiation coupling is suppressed by an additional symmetry.

The range $3.4 \times 10^{-41} < m_{sp} < 1.1 \times 10^{-46}$ is *empty* (the lower bound exceeds the upper), implying that no single value of m_{sp} simultaneously suppresses TGP radiation and allows the force to operate at Solar-System scales.

Resolution: TGP with $m_{sp} \gg 1$ (Planck-scale mass) gives $G_{\text{eff}} \approx 0$ everywhere except at sub-nuclear distances and zero radiation for all sub-Planck frequencies. In this limit, TGP is a theory of *quantum spacetime foam* at the Planck scale, decoupled from classical gravity observations entirely. Classical gravity is then described by GR alone, and TGP provides quantum corrections only at Planck energies.

14 False vacuum structure and spontaneous nucleation in TGP

14.1 The inverted potential: $V(g = 1)$ is a maximum

The TGP interaction potential, in dimensionless units $g = \Phi/\Phi_0$ with $\beta = \gamma = 1$, reads

$$V(g) = \frac{g^3}{3} - \frac{g^4}{4}, \qquad V'(g) = g^2(1-g), \qquad V''(g) = 2g - 3g^2. \tag{52}$$

The critical points satisfy $V' = 0$: these are $g = 0$ (inflection, $V''(0) = 0$) and $g = 1$ ($V''(1) = -1 < 0$, a **maximum**). There is no local minimum for $g \in (0, 1)$. The values at the structurally important points are:

Point	g	$V(g)$	Type
"Nothingness"	0	0	inflection
Ghost boundary	$g^* = e^{-1/4} \approx 0.7788$	0.0655	—
"Space" (vacuum)	1	$\frac{1}{12} \approx 0.0833$	**maximum**

Since $V(0) < V(g^*) < V(1)$, the vacuum $g = 1$ has the *highest* potential energy among the accessible field values. It is a **false vacuum**.

14.2 Linearised field equation and oscillatory tails

Setting $g = 1 - \delta$, $|\delta| \ll 1$, and linearising gives $(\nabla^2 + 1)\delta = 0$, whose spherically symmetric solution is $\delta(r) = A \sin r/r$ — oscillatory, not Yukawa. This follows *directly* from $V''(1) = -1 < 0$: the negative second derivative is the field-theoretic origin of the oscillatory far field.

14.3 Defect energy landscape

The total energy of a spherically symmetric defect profile $g(r)$ relative to the vacuum is

$$E_{\text{def}} = 4\pi \int_0^\infty r^2 \left[\tfrac{1}{2} f(g)(g')^2 + V(g) - V(1)\right] dr, \tag{53}$$

with the nonlinear kinetic function $f(g) = 1 + 2\alpha \ln g$ ($\alpha = 2$) and ghost boundary $g^* = e^{-1/4}$.

Numerical integration of the full ODE (47) with $g(0) = g_0$, $g'(0) = 0$ (ex16) yields the following energy landscape:

g_0	E_{def} [Planck]	Character
0.780	+0.019	unfavourable (E>0)
0.849	+0.003	near-critical
0.869	−0.000	**zero crossing**
0.914	−0.0043	**optimal (E minimum)**
0.958	−0.0020	shallow, nearly zero
0.999	$\to 0^-$	trivial (no defect)

The energy changes sign at $g_0 \approx 0.87$:

- $g_0 \in (g^*, 0.87)$: **deep defects,** $E > 0$ — energetically unfavourable; the kinetic cost at the ghost boundary dominates.
- $g_0 \in (0.87, 1)$: **shallow defects,** $E < 0$ — energetically favourable; the potential energy gain $V(g_0) - V(1) < 0$ dominates.

The **minimum-energy defect** has $g_0 \approx 0.91$, $E_{\min} \approx -0.0043\,\text{Planck}$.

14.4 Metastability and the nucleation barrier

The sign reversal at $g_0 \approx 0.87$ implies a **nucleation barrier**: to create a stable ($E < 0$) defect, a fluctuation must overshoot the $g_0 \approx 0.87$ threshold. The thin-wall approximation for a spherical bubble of radius R gives

$$E_{\text{bubble}}(R) = -\frac{4\pi}{3} R^3 \, |\varepsilon(g_0)| + 4\pi R^2 \, \sigma(g_0), \tag{54}$$

with $\varepsilon(g_0) = V(g_0) - V(1) < 0$ the volumetric energy gain and $\sigma(g_0)$ the wall surface energy. The critical radius and barrier height are

$$R_{\text{crit}} = \frac{2\sigma}{|\varepsilon|}, \qquad E_{\text{barrier}} = \frac{16\pi\sigma^3}{3\varepsilon^2}. \tag{55}$$

For the optimal depth $g_0 = 0.90$ with $\sigma_{\text{eff}} \approx 0.045$ and $|\varepsilon| \approx 0.0044$:

$$R_{\text{crit}} \approx 21\, l_{\text{Pl}}, \qquad E_{\text{barrier}} \approx 81\, E_{\text{Pl}}. \tag{56}$$

The vacuum lifetime is $\tau \sim \exp(E_{\text{barrier}}/\hbar) \sim \exp(81) \sim 10^{35}\, t_{\text{Pl}}$. In SI units, $10^{35}\, t_{\text{Pl}} \approx 5 \times 10^{-9}\,\text{s}$ — surprisingly short, but in Planck units enormous compared to the Planck time.

Important caveat on energy convergence. The oscillatory tail $g(r) - 1 \sim A \sin r/r$ contributes a kinetic term $(g')^2 \sim A^2/r^2$, giving an integrand $\sim r^2/r^2 = A^2$ that does not decay. Thus E_{def} as defined is *IR-divergent* for isolated defects. The physical implication is that **TGP bodies must appear in pairs (defect + anti-defect)**, whose oscillatory tails cancel at large r, leaving a finite pair energy. This mirrors the UV divergence of point charges in classical electrodynamics, and predicts a matter–antimatter pairing structure.

14.5 Cosmological scenario: matter nucleation from the vacuum

The false-vacuum structure of TGP suggests the following cosmological scenario:

1. **Initial state:** Universe at $g = 1$ everywhere — "pure space", the false vacuum. This corresponds to the axiom $\mathcal{N}_0$ of TGP.
2. **Quantum fluctuations:** At the Planck scale, field configurations with $g < 1$ are continuously generated. Fluctuations with $g_0 > 0.87$ are energetically favoured and grow.
3. **Critical bubble nucleation:** Bubbles reaching $R > R_{\text{crit}} \approx 21\, l_{\text{Pl}}$ grow without bound, converting the false vacuum locally to a defect state.
4. **Defect = body:** Each stable defect pair (defect + anti-defect) constitutes a TGP body with rest energy $E_{\text{pair}} \approx 2|E_{\min}| \approx 0.0086\, E_{\text{Pl}} \approx 1.9 \times 10^{-19}\,\text{kg} \cdot c^2$.
5. **Ghost boundary as stabiliser:** Defects cannot deepen below $g^* \approx 0.778$ because the kinetic term $f(g^*) = 0$ creates an impenetrable kinematic barrier. Bodies are "coated" by this ghost layer.

The analogy with Schwinger pair production in QED is close: just as a strong electromagnetic field can nucleate electron–positron pairs from the QED vacuum, the TGP vacuum (false vacuum at $g = 1$) can nucleate body–antibody pairs from its own instability.

14.6 Energy scale of nucleated bodies

With $\beta = \gamma = 1$ (natural Planck-scale TGP), the pair energy $E_{\text{pair}} \approx 0.0086\, E_{\text{Pl}}$, corresponding to a mass

$$m_{\text{TGP}} \approx 0.0086\, m_{\text{Pl}} \approx 1.9 \times 10^{-10}\,\text{kg}. \tag{57}$$

This is the mass of a dust grain — not a fundamental particle. For TGP to describe quarks or electrons, the coupling constants must satisfy $m_{\text{body}} \sim 10^{-19}\, m_{\text{Pl}}$, requiring $\beta, \gamma \ll 1$. The ratio

$$m_{\text{body}}/m_{\text{Pl}} \sim (\beta\gamma)^{1/2}/12 \tag{58}$$

relates the TGP coupling constants to the observed particle masses.

14.7 Summary of vacuum structure

Property	Value / Conclusion
Vacuum type	False vacuum ($V''(1) < 0$)
True minimum	$g = 0$ (nothingness), $V = 0$
Ghost boundary	$g^* = e^{-1/4} \approx 0.778$, $f(g^*) = 0$
Energy barrier crossing	$g_0^{\text{crit}} \approx 0.87$
Optimal defect depth	$g_0^{\text{opt}} \approx 0.91$
Min. defect energy	$E_{\text{min}} \approx -0.0043\, E_{\text{Pl}}$
Critical bubble radius	$R_{\text{crit}} \approx 21\, l_{\text{Pl}}$
Nucleation barrier	$E_{\text{bar}} \approx 81\, E_{\text{Pl}}$
IR convergence	Requires defect pairs (oscillatory tails cancel)

15 Exact three-body N-body observability: Feynman integral vs saddle-point

15.1 Setup

The three-body TGP interaction energy for bodies at positions $\mathbf{r}_1, \mathbf{r}_2, \mathbf{r}_3$ with coupling $C_i = m_i/(2\sqrt{\pi})$ [Planck units] is

$$V_3 = C_1 C_2 C_3\, I_Y(d_{12}, d_{13}, d_{23}; m_{\text{sp}}), \tag{59}$$

where I_Y is the Feynman two-parameter integral

$$I_Y = 2\int_{\Delta_2} \Delta^{-3/2}\, K_0\!\left(m_{\text{sp}}\sqrt{Q/\Delta}\right) d\alpha, \qquad Q = \alpha_2 d_{12}^2 + \alpha_1 d_{13}^2 + \alpha_3 d_{23}^2, \quad \Delta = \alpha_1\alpha_2 + \alpha_1\alpha_3 + \alpha_2\alpha_3. \tag{60}$$

The two-body Yukawa energy is $V_2 = C_i C_j\, e^{-m_{\text{sp}} d_{ij}}/d_{ij}$.

The observable quantity is the **three-to-two body energy ratio**

$$\mathcal{R} \equiv \frac{V_3}{V_2} = C\,\frac{I_Y(d, d, d; m_{\text{sp}})}{e^{-t}/d}, \qquad t = m_{\text{sp}}\, d, \tag{61}$$

evaluated for an equilateral triangle of side d and a single coupling C.

15.2 Numerical results

15.2.1 $\mathcal{R}$ as a function of $t = m_{\text{sp}} d$

Table 3 gives the exact ratio $\mathcal{R}$ computed by Gauss–Legendre quadrature on the Duffy-transformed simplex (60-point rule), together with the saddle-point approximation obtained by evaluating the integrand at $\boldsymbol{\alpha}^* = (1/3, 1/3, 1/3)$ with the asymptotic $K_0(u) \approx \sqrt{\pi/(2u)}\, e^{-u}$.
The saddle-point formula systematically *overestimates* $\mathcal{R}$ by a factor 1.6–5×.

Table 3: Exact vs saddle-point $\mathcal{R} = V_3/V_2$ for an equilateral triangle, $C = 0.20$ (i.e. $m_{\rm body} \approx 0.7\, m_{\rm Pl}$).

$t = m_{\rm sp} d$	$\mathcal{R}_{\rm exact}$	$\mathcal{R}_{\rm saddle}$	Error
0.5	0.580	0.929	+60%
1.0	0.484	0.898	+86%
1.5	0.357	0.767	+115%
2.0	0.253	0.627	+148%
3.0	0.121	0.373	+208%
5.0	0.026	0.112	+329%
6.0	0.011	0.055	+392%

15.2.2 Origin of the saddle-point error

The saddle-point relies on the asymptotic expansion

$$K_0(u) \approx \sqrt{\frac{\pi}{2u}}\, e^{-u}, \qquad u \gg 1. \tag{62}$$

For the equilateral triangle at $\boldsymbol{\alpha}^* = (1/3, 1/3, 1/3)$:

$$u^* = m_{\rm sp}\sqrt{Q^*/\Delta^*} = \sqrt{3}\, t. \tag{63}$$

The asymptotic expansion is reliable only for $u^* > 3$, i.e. $t > 1/\sqrt{3} \approx 1.73$. In the range $t < 1.73$ (Planck-scale close encounters) the expansion fails completely and the saddle-point overestimates by 50–400%.

15.3 Critical coupling $C_{\rm crit}$ for observability

Defining observability as $\mathcal{R} > 1\%$, the critical coupling is

$$C_{\rm crit}(t) = \frac{0.01\, V_2(d, m_{\rm sp})}{I_Y(d, d, d; m_{\rm sp})}. \tag{64}$$

Table 4: Critical coupling $C_{\rm crit}$ for $\mathcal{R} > 1\%$ and corresponding body mass via $m = 2\sqrt{\pi}\, C\, m_{\rm Pl}$.

$t = m_{\rm sp} d$	$C_{\rm crit}$ [Planck]	$m_{\rm crit}$ (kg)
0.5	3.4×10^{-3}	2.7×10^{-10}
1.0	4.1×10^{-3}	3.2×10^{-10}
2.0	7.9×10^{-3}	6.1×10^{-10}
4.0	3.5×10^{-2}	2.7×10^{-9}
5.0	7.6×10^{-2}	5.9×10^{-9}

Three-body TGP effects are therefore observable only for bodies with

$$m_{\rm body} \gtrsim 3 \times 10^{-10}\,{\rm kg} \quad (\approx 300\,{\rm ng}) \tag{65}$$

interacting at Planck distances.

15.4 Physical systems

The conclusion is definitive:

- For all *known* fundamental particles, $\mathcal{R} \ll 10^{-19}$ — three-body TGP interactions are completely negligible.
- Three-body effects become $O(1)$ only for hypothetical objects of Planck mass ($m \approx m_{\rm Pl} \approx 2 \times 10^{-8}\,{\rm kg}$) at Planck separation.

Table 5: $\mathcal{R} = V_3/V_2$ for known physical systems. C from Path B: $C = m_{\rm body}/(2\sqrt{\pi}\, m_{\rm Pl})$.

System	C [Pl]	$t = m_{\rm sp} d$	$\mathcal{R}$	Observable?
Proton (QCD scale)	2.2×10^{-20}	$\gg 1$	$\sim 10^{-20}$	No
Electron (Bohr)	1.2×10^{-23}	$\gg 1$	$\sim 10^{-23}$	No
Dust grain (100 nm)	1.3×10^{-14}	0.62 (at $m_{\rm sp} = 10^{-28}$)	3.7×10^{-14}	No
Hypothetical Planck object	0.282	1.0	0.68	**Yes**

15.5 Implication: saddle-point systematically underestimates 3-body effects

Because $\mathcal{R}_{\rm saddle} > \mathcal{R}_{\rm exact}$ (saddle overestimates I_Y but the ratio $C\, I_Y/V_2$ is *also* overestimated), the physical conclusion is unchanged: three-body TGP forces are unobservable for standard particles. However, the analytic approximation from Schwinger/saddle-point predicts **stronger** three-body effects than actually occur, meaning any analytic bound on observability based on saddle-point is *conservative* (correctly excludes detection).

Table 6: Summary of three-body observability analysis.

Property	Value / Conclusion
Exact integral method	Duffy-transformed 2D Gauss–Legendre (60 pts)
Saddle-point validity	$t = m_{\rm sp} d > 1.73$ only
Saddle error at $t = 1$	+86% (overestimate)
Saddle error at $t = 5$	+329% (overestimate)
Observability threshold ($\mathcal{R} > 1\%$)	$m_{\rm body} \gtrsim 300\,$ng at $d \sim l_{\rm Pl}$
Standard particles	$\mathcal{R} \sim 10^{-20}$ (undetectable)
Planck-scale objects	$\mathcal{R} \sim 0.7$ (dominant 3-body)

16 Three-body TGP scattering dynamics for Planck-scale objects

16.1 Setup

We simulate three Planck-scale objects with coupling $C = 1/(2\sqrt{\pi}) \approx 0.282$ [Planck] and scalar screening mass $m_{\rm sp} = 1\, m_{\rm Pl}^{-1}$. The interaction potential is

$$V = \underbrace{-\sum_{i<j} C^2 \frac{e^{-m_{\rm sp} r_{ij}}}{r_{ij}}}_{V_2} \underbrace{- \sum_{i<j<k} C^3\, I_Y(r_{ij}, r_{ik}, r_{jk};\, m_{\rm sp})}_{V_3}, \tag{66}$$

with I_Y the Feynman two-parameter integral (eq. 7). The equations of motion are integrated with the Velocity Verlet scheme ($\Delta t = 0.02\, t_{\rm Pl}$, $T_{\rm max} = 25\, t_{\rm Pl}$).

16.2 Static energetics

For an equilateral triangle of side d:

$$V_2 = -3C^2 \frac{e^{-d}}{d}, \qquad V_3 = -C^3\, I_Y(d, d, d;\, m_{\rm sp} = 1). \tag{67}$$

Key result: at Planck separations $d \sim 1$–$2\, l_{\rm Pl}$, the three-body interaction contributes 10–23% of the total potential energy. This is not a perturbative correction; it is a *leading-order effect*.

Table 7: Potential energies and ratio V_3/V_2 for equilateral triangle, $C = C_{\rm Pl} = 1/(2\sqrt{\pi})$, $m_{\rm sp} = 1$.

d [$l_{\rm Pl}$]	V_2 [$E_{\rm Pl}$]	V_3 [$E_{\rm Pl}$]	V_3/V_2
1.0	−0.0878	−0.0200	22.8%
1.5	−0.0355	−0.0060	16.8%
2.0	−0.0162	−0.0019	11.9%
2.5	−0.0078	−0.0007	8.3%
3.0	−0.0040	−0.0002	5.7%
4.0	−0.0011	−0.00003	2.7%

16.3 Collision dynamics: observer displacement

Configuration: bodies 1 and 2 collide head-on along the x-axis with $v_0 = 0.25\, l_{\rm Pl}/t_{\rm Pl}$; body 3 (observer) sits at impact parameter $b = 2\, l_{\rm Pl}$ on the y-axis.

By symmetry, body 3 moves only along y. The position discrepancy between the "2-body only" and "2-body+3-body" trajectories grows with time:

Table 8: Position of observer (body 3) under 2B vs 2B+3B forces.

t [$t_{\rm Pl}$]	y_3 (2B only)	y_3 (2B+3B)	$\lvert\delta y_3\rvert$
5	2.010	2.010	1.8×10^{-4}
10	2.087	2.082	4.1×10^{-3}
15	2.344	2.298	4.5×10^{-2}
20	2.764	2.621	1.4×10^{-1}
25	3.234	2.997	2.4×10^{-1}

At $t = 25\, t_{\rm Pl}$ the three-body force has displaced the observer by $|\delta y_3| \approx 0.24\, l_{\rm Pl}$ — a measurable Planck-scale deviation.

16.4 Triangle stability: three-body compactness

An equilateral triangle ($d = 2\, l_{\rm Pl}$, $\omega = 0.1$) is unbound under 2-body forces alone (total $E_0 > 0$). With three-body forces the binding energy deepens:

$$E_0^{(3B)} = E_0^{(2B)} + V_3 = 0.00885 - 0.00192 = 0.00693\, E_{\rm Pl}. \tag{68}$$

Even though the triangle is still unbound, the separation grows more slowly:

Table 9: Side length d_{12} of the expanding triangle: 2B vs 2B+3B.

t [$t_{\rm Pl}$]	d_{12} (2B)	d_{12} (2B+3B)	δd_{12}
5	2.49	2.45	−0.040
10	3.54	3.44	−0.099
15	4.78	4.63	−0.148
20	6.06	5.87	−0.190

The three-body interaction makes the triangle $\sim 3\%$ more compact throughout the evolution. For configurations with $E_0^{(2B)} < 0$ (bound), the three-body term would deepen the bound state energy by $\sim 12\%$ at $d = 2\, l_{\rm Pl}$.

16.5 Summary

Main conclusion: for hypothetical Planck-scale objects, three-body TGP interactions are *not a perturbative correction* but contribute at the 10–23% level to the total potential energy. The dynamical signature — a $\sim 0.24\, l_{\rm Pl}$ observer displacement over $25\, t_{\rm Pl}$ — is of the same order as the

Table 10: Summary of three-body scattering results for Planck-scale objects ($C \approx 0.282$, $m_{\rm sp} = 1$).

Observable	Result
V_3/V_2 at $d = 1\,l_{\rm Pl}$	22.8%
V_3/V_2 at $d = 2\,l_{\rm Pl}$	11.9%
Observer displacement at $t = 25$	$0.24\,l_{\rm Pl}$
Triangle compactness increase (2B+3B vs 2B)	$\approx 3\%$
Bound-state energy deepening	$\approx 12\%$ at $d = 2$
Standard particles ($C \sim 10^{-20}$)	$V_3/V_2 \sim 10^{-20}$ (undetectable)

Planck length and therefore, in principle, measurable within Planck-scale physics. For all known particles the effect is $\sim 10^{-20}$ and completely negligible.

17 Three-body induced binding in TGP

17.1 Static potential landscape

For an equilateral triangle of $N = 3$ Planck-scale objects ($C = 1/(2\sqrt{\pi})$, $m_{\rm sp} = 1$), the total potential energy is

$$V_{\rm tot}(d) = V_2(d) + V_3(d) = -3C^2\frac{e^{-d}}{d} - C^3 I_Y(d, d, d;\, m_{\rm sp} = 1), \tag{69}$$

where I_Y is the Feynman integral from Sec. 15.

Table 11: Potential energies for equilateral triangle, $C = C_{\rm Pl}$, $m_{\rm sp} = 1$.

d [$l_{\rm Pl}$]	V_2	V_3	$V_2 + V_3$	$\lvert V_3/V_2\rvert$
0.40	−0.400	−0.109	−0.509	27.3%
0.80	−0.134	−0.034	−0.168	25.1%
1.00	−0.088	−0.020	−0.108	22.8%
2.00	−0.016	−0.002	−0.018	11.9%
3.00	−0.004	−0.0002	−0.004	5.7%

$V_2 + V_3 < 0$ for all $d > 0$: the triangular configuration is always in an attractive potential well, with V_3 contributing 6–27% of the total depth.

17.2 Geometry: triangle vs linear configuration

Table 12: $V_2 + V_3$ for equilateral triangle vs collinear arrangement ($d_{12} = d_{23} = d$, $d_{13} = 2d$).

d	Triangle	Linear	Δ
0.5	−0.369	−0.271	−0.098
1.0	−0.108	−0.073	−0.035
2.0	−0.018	−0.012	−0.006
3.0	−0.0042	−0.0027	−0.0015

The equilateral triangle is the energetically preferred geometry: it sits 26–36% deeper in the potential well than the collinear arrangement. This is a selection rule imposed by V_3: among all three-body configurations, the symmetric triangle is favoured.

17.3 N-body binding: super-extensive deepening

A crucial feature of V_3 is that the number of three-body triples grows as $\binom{N}{3} = O(N^3)$ while the two-body terms grow as $\binom{N}{2} = O(N^2)$. Consequently, as N increases, V_3/V_2 *grows*, and the binding

energy per particle increases super-extensively.

Table 13: N objects on a regular polygon with side $d = 1.5\, l_{\rm Pl}$.

N	V_2	V_3	$V_2 + V_3$	$E_{\rm tot}/N$
3	-0.0355	-0.0060	-0.0415	-0.0138
4	-0.0563	-0.0151	-0.0714	-0.0179
5	-0.0737	-0.0229	-0.0966	-0.0193
6	-0.0887	-0.0284	-0.1171	-0.0195

Table 14: N objects on polygon, $d = 0.8\, l_{\rm Pl}$ (near optimal).

N	V_3/V_2	$E_{\rm tot}/N$ $[E_{\rm Pl}]$
2	—	-0.022
3	25.1%	-0.056
4	45.1%	-0.081
5	59.6%	-0.098

As N grows from 3 to 5, the binding energy per particle increases from 0.056 to $0.098\, E_{\rm Pl}$ — a 75% gain entirely due to three-body effects, since V_2/N varies only weakly with N for a fixed polygon side.

17.4 Virial theorem: exact three-body deepening (ex21)

The full virial energy including both V_2 and V_3 is

$$E_{\rm virial} = (V_2 + V_3) - \tfrac{1}{2}\big(\langle \mathbf{r} \cdot \nabla V_2 \rangle + \langle \mathbf{r} \cdot \nabla V_3 \rangle\big). \tag{70}$$

For $V_2 = -C^2 e^{-mr}/r$: $r\, \partial V_2/\partial r = -(1 + mr)\, V_2$. The V_3 gradient is computed by finite difference of $I_Y(d)$.

Table 15: Virial equilibrium energy vs coupling C for equilateral triangle, $m_{\rm sp} = 1$, $d = 0.3\, l_{\rm Pl}$ (hard-core limit).

C	$E_{\rm min}^{(2B)}$	$E_{\rm min}^{(2B+3B)}$	ΔE	$\lvert V_3/V_2 \rvert$
0.050	-0.031	-0.032	-0.001	4.7%
0.100	-0.122	-0.133	-0.011	9.3%
0.150	-0.275	-0.311	-0.036	14.0%
0.200	-0.489	-0.575	-0.086	18.6%
0.250	-0.764	-0.932	-0.168	23.3%
0.282	-0.972	-1.213	-0.241	26.2%

Key result: The Yukawa potential ($V_2 \sim -1/r$) is classically unbounded from below, so both $E^{(2B)}$ and $E^{(2B+3B)}$ are always negative — the system is always bound. Three-body forces *consistently deepen* the bound state by 5–26% depending on C.

Quantum correction: In a quantum treatment, zero-point kinetic energy $\Delta p^2/2m \sim \hbar^2/(2md^2)$ stabilises the system at a finite $d_{\rm eq}$. There the two-body Yukawa coupling has a critical value $C_{\rm crit}^{\rm QM}$ below which the quantum ground state is unbound; near this threshold, V_3 could create a purely *three-body-induced bound state* — a falsifiable prediction requiring a full quantum TGP calculation.

Table 16: Three-body bound-state analysis summary.

Result	Value
V_3/V_2 at $d = 0.8\,l_{\rm Pl}$, $N = 3$	25.1%
V_3/V_2 at $d = 0.8\,l_{\rm Pl}$, $N = 5$	59.6%
Preferred geometry	Equilateral triangle
$E_{\rm tot}/N$ improvement ($N\colon 3 \to 5$)	+75%
Virial (2B only, $d = 0.8$)	$E > 0$ (unbound without V_3)
3B-induced bound state	Exists near $C \lesssim C_{\rm Pl}$

17.5 Summary

18 Efimov-like three-body bound states in TGP

18.1 The question

Can three-body TGP forces create a *bound triangular state* that does not exist in the two-body theory alone? Specifically: is there a regime where

$$E_{\rm min}^{(2B+{\rm ZP})} > 0 \quad \text{(unbound pair)} \qquad \text{but} \qquad E_{\rm min}^{(2B+3B+{\rm ZP})} < 0 \quad \text{(bound triple)?} \tag{71}$$

The effective energy including zero-point pressure is

$$E_{\rm eff}(d; C, m_{\rm sp}) = \underbrace{\frac{3}{8d^2}}_{\rm ZP} + \underbrace{V_2(d; C, m_{\rm sp})}_{\text{2-body Yukawa}} + \underbrace{V_3(d; C, m_{\rm sp})}_{\text{3-body Feynman}}, \tag{72}$$

where d is the equilateral triangle side length and $V_3 = -C^3 I_Y(d, d, d;\, m_{\rm sp})$.

18.2 Critical couplings

Minimising $E_{\rm eff}$ over d defines $E_{\rm min}(C, m_{\rm sp})$. The **critical coupling** $C_{\rm crit}(m_{\rm sp})$ is where $E_{\rm min} = 0$: above this value the triangle is quantum-mechanically bound.

Table 17: Critical couplings for two-body ($C_{\rm crit}^{(2B)}$) and two-body+three-body ($C_{\rm crit}^{(3B)}$) binding vs scalar mass $m_{\rm sp}$. The *window* column gives the interval of C for which the triple is bound but no pair is bound.

$m_{\rm sp}$ $[l_{\rm Pl}^{-1}]$	$\lambda = 1/m_{\rm sp}$ $[l_{\rm Pl}]$	$C_{\rm crit}^{(2B)}$	$C_{\rm crit}^{(3B)}$	Window width
0.05	20.0	0.130	0.084	0.047
0.10	10.0	0.184	0.128	**0.057**
0.12	8.3	0.202	0.142	0.060
0.15	6.7	0.226	0.163	0.063
0.20	5.0	0.261	0.193	0.067
0.25	4.0	0.292	0.221	0.071
0.30	3.3	0.319	0.246	0.074

The window $C_{\rm crit}^{(3B)} < C < C_{\rm crit}^{(2B)}$ exists for all $m_{\rm sp} \lesssim 0.4\,l_{\rm Pl}^{-1}$ (range $\lambda \gtrsim 2.5\,l_{\rm Pl}$). Its width *grows* monotonically from $\Delta C \approx 0.047$ at $m_{\rm sp} = 0.05$ to $\Delta C \approx 0.074$ at $m_{\rm sp} = 0.30$ (see Table 17).

Numerical note: Values for $m_{\rm sp} \geq 0.15$ were computed using a dense two-step scan over $d \in [0.3,\, 25]\,l_{\rm Pl}$ (600 coarse + 500 refinement points) rather than scalar optimization, which fails to locate the shallow minimum at $d \sim 4$–$8\,l_{\rm Pl}$ for short-range Yukawa potentials.

18.3 The 3B-induced window at $m_{\rm sp} = 0.1$

For $m_{\rm sp} = 0.1\, l_{\rm Pl}^{-1}$ ($\lambda = 10\, l_{\rm Pl}$):

$$C_{\rm crit}^{(3B)} = 0.128, \qquad C_{\rm crit}^{(2B)} = 0.184. \tag{73}$$

In the interval $C \in (0.128,\, 0.184)$:

Table 18: 3B-induced bound states for $m_{\rm sp} = 0.1$, $C \in (C_{\rm crit}^{(3B)},\, C_{\rm crit}^{(2B)})$.

C	$E_{\rm min}^{(2B)}$	$E_{\rm min}^{(2B+3B)}$	$d_{\rm eq}$ $[l_{\rm Pl}]$	$E_{\rm bind}$ $[E_{\rm Pl}]$
0.141	+0.00032	−0.00248	5.74	0.0025
0.155	+0.00030	−0.00714	4.50	0.0071
0.169	+0.00027	−0.01452	3.70	0.0145
0.183	+0.00006	−0.02525	3.13	0.0253

The equilibrium separation $d_{\rm eq} \approx 3$–$6\, l_{\rm Pl}$ places the triangle deep in the sub-Planck regime.

18.4 Connection to Efimov physics

This phenomenon is the TGP analog of **Efimov states** [**?**]: in nuclear/atomic physics, three identical bosons can form an infinite tower of bound states even when their pairwise interaction is too weak to support a two-body bound state. The mechanism is a $1/R^2$ effective potential in the three-body channel (where R is the hyper-radius), which overcomes the centrifugal barrier.

In TGP, the analogous mechanism is:

- Two-body Yukawa $V_2 = -C^2 e^{-m_{\rm sp} d}/d$ cannot overcome quantum pressure for $C < C_{\rm crit}^{(2B)}$.
- The Feynman three-body integral $V_3 = -C^3 I_Y(d, d, d;\, m_{\rm sp})$ provides additional attractive energy scaling as C^3 (growing faster than $V_2 \sim C^2$ as C increases) which can *tip the balance*.
- Result: an equilateral triangle is bound at $d_{\rm eq} \sim 3$–$6\, l_{\rm Pl}$ in a range of C where no two-body bound state exists.

18.5 Physical parameters of the 3B-induced state

At $m_{\rm sp} = 0.1\, l_{\rm Pl}^{-1}$, $C = 0.155$ (midpoint of window):

$$d_{\rm eq} \approx 4.5\, l_{\rm Pl} \approx 7 \times 10^{-35}\,{\rm m}, \tag{74}$$

$$E_{\rm bind} \approx 7 \times 10^{-3}\, E_{\rm Pl} \approx 1.4 \times 10^{7}\,{\rm J}, \tag{75}$$

$$m_{\rm body} = 2\sqrt{\pi}\, C\, m_{\rm Pl} \approx 1.2 \times 10^{-8}\,{\rm kg} \approx 0.55\, m_{\rm Pl}. \tag{76}$$

The 3B-induced window in mass space (for $m_{\rm sp} = 0.1$):

$$m_{\rm body} \in [9.8 \times 10^{-9},\, 1.4 \times 10^{-8}]\,{\rm kg} \quad (= [0.45,\, 0.65]\, m_{\rm Pl}). \tag{77}$$

18.6 WKB level count: finite tower (ex24)

Using the WKB quantisation condition $n(E) = \frac{1}{\pi}\int_{d_L}^{d_R} \sqrt{2\mu(E - V_{\rm eff})}\, dd = n + \frac{1}{2}$, with $V_{\rm eff}(d) = \frac{3}{8d^2} + V_2 + V_3$ and $\mu = 1\, m_{\rm Pl}$:

For $C = 0.180$, the single WKB ground state has:

$$E_0 = -9 \times 10^{-4}\, E_{\rm Pl}, \qquad d_L = 1.9\, l_{\rm Pl}, \qquad d_R = 16.3\, l_{\rm Pl}, \qquad \Delta d = 14.4\, l_{\rm Pl}. \tag{78}$$

The state is very *diffuse* (spread over $\sim 14\, l_{\rm Pl}$) and weakly bound.

Table 19: WKB bound-state count in the 3B window, $m_{\rm sp} = 0.1$.

C	$N_{\rm states}$	E_0 $[E_{\rm Pl}]$	$d_{\rm eq}$ $[l_{\rm Pl}]$	
0.128–0.175	0	—	5–8	(below WKB threshold)
0.175	1	−0.00029	3.48	
0.180	1	−0.00090	3.26	
0.184	1	−0.00156	3.04	

Key finding: TGP supports *at most one* WKB-level three-body bound state in the window $C \in (C_{\rm crit}^{(3B)}, C_{\rm crit}^{(2B)})$. There is no infinite Efimov tower.

Why no tower: The classical Efimov effect arises from an exact $\sim 1/R^2$ hyper-radial potential, which supports an infinite geometric series of levels. The TGP three-body integral $V_3(d) \sim -C^3 I_Y(d)$ is *screened* (decays as Yukawa for $d \gtrsim \lambda = 1/m_{\rm sp}$), producing only a finite potential well and at most one or two bound levels.

Erratum (ex26, Sec. 20): The WKB result above is *invalid* for this regime. At $C = 0.180$ the de Broglie wavelength at the well centre is $\lambda_{\rm dB} \approx 44\, l_{\rm Pl} \gg L_{\rm well} \approx 14\, l_{\rm Pl}$, violating the WKB validity condition. Exact finite-difference diagonalisation (ex26) shows the state is *unbound* ($E_0^{\rm FD} = +1.8 \times 10^{-3}\, E_{\rm Pl} > 0$) for all $C \leq 0.184$. The true quantum Efimov window is $C \in (0.191, 0.310)$; see Sec. 20.

18.7 Falsifiability

This is a **falsifiable prediction** of TGP:

1. If $m_{\rm sp} \in (0.05,\, 0.4)\, l_{\rm Pl}^{-1}$ (scalar range $\lambda \in (2.5, 20)\, l_{\rm Pl}$), triangular bound states of Planck-mass objects should exist that are absent in the two-body theory.
2. The window width $\Delta C \approx 0.05$ translates to a mass window $\Delta m/m_{\rm Pl} \approx 0.35$ — not fine-tuned.
3. No such state should exist for ordinary matter ($C \sim 10^{-20} \ll C_{\rm crit}^{(3B)}$).

Table 20: Summary: TGP Efimov-analog bound states.

Property	Value
Window existence	$m_{\rm sp} < 0.4\, l_{\rm Pl}^{-1}$, $\lambda > 2.5\, l_{\rm Pl}$
Window width	$\Delta C \approx 0.05$ (independent of $m_{\rm sp}$)
$d_{\rm eq}$ (centre of window)	3–6 $l_{\rm Pl}$
$E_{\rm bind}$ (centre of window)	10^{-3}–$10^{-2}\, E_{\rm Pl}$
Mass range (window, $m_{\rm sp} = 0.1$)	$[9.8 \times 10^{-9},\, 1.4 \times 10^{-8}]$ kg
Analogy	Efimov states (nuclear/atomic physics)

19 Optimal three-body geometry

19.1 Question

For a fixed coupling C and scalar mass $m_{\rm sp}$, which spatial arrangement of three (or four) TGP bodies minimises the total effective energy? Is the equilateral triangle — assumed throughout the Efimov analysis — truly the ground-state geometry?

19.2 Setup

We parameterise each geometry by a single length scale d (the nearest-neighbour separation) and minimise the effective potential

$$E_{\text{eff}}(d) = \underbrace{\frac{N}{8d^2}}_{\text{ZP}} + V_2^{\text{tot}}(d; C, m_{\text{sp}}) + V_3^{\text{tot}}(d; C, m_{\text{sp}}), \tag{79}$$

with $N = 3$ for triangles and $N = 4$ for the square, using the same Feynman-integral convention as `ex23` (see Sec. 18). Geometries studied:

1. **Equilateral triangle**: $r_{12} = r_{13} = r_{23} = d$.
   $V_2^{\text{tot}} = -3C^2 e^{-m_{\text{sp}}d}/d, \quad V_3^{\text{tot}} = -C^3 I_Y(d, d, d;\, m_{\text{sp}})$.

2. **Linear chain**: $r_{12} = r_{23} = d$, $r_{13} = 2d$.
   $V_2^{\text{tot}} = -C^2(2e^{-m_{\text{sp}}d}/d + e^{-2m_{\text{sp}}d}/(2d)), \quad V_3^{\text{tot}} = -C^3 I_Y(d, 2d, d;\, m_{\text{sp}})$.

3. **Isosceles triangle**: $r_{12} = r_{13} = d$, $r_{23} = sd$.
   $V_2^{\text{tot}} = -C^2(2e^{-m_{\text{sp}}d}/d + e^{-m_{\text{sp}}sd}/(sd)), \quad V_3^{\text{tot}} = -C^3 I_Y(d, d, sd;\, m_{\text{sp}})$.
   Shape parameter s is varied.

4. **Square** ($N = 4$): side d, diagonal $d\sqrt{2}$.
   $V_2^{\text{tot}} = -C^2(4e^{-m_{\text{sp}}d}/d + 2e^{-m_{\text{sp}}d\sqrt{2}}/(d\sqrt{2}))$,
   four right-triangle V_3 contributions (side, side, diagonal).

19.3 Results at $m_{\text{sp}} = 0.1$, $C = 0.155$

Parameters correspond to the midpoint of the Efimov-window identified in Sec. 18.

Table 21: Binding energy vs geometry for $m_{\text{sp}} = 0.1$, $C = 0.155$.

Geometry	N	E_{min} $[E_{\text{Pl}}]$	d_{ref} $[l_{\text{Pl}}]$	Bound?
Equilateral triangle	3	-0.00706	4.52	Yes
Linear chain	3	-0.00026	6.25	Yes
Square	4	-0.06842	2.48	Yes

Key result: The equilateral triangle is the energetically preferred symmetric three-body geometry. The linear chain is *bound* at $C = 0.155$ but with binding energy $\sim 27\times$ weaker than the equilateral triangle.

19.4 Equilateral vs linear: 3B-only windows

Both geometries possess a coupling window in which pairs are unbound but the multi-body cluster is bound (the TGP Efimov-analog):

Table 22: 3B-only windows for equilateral and linear geometries, $m_{\text{sp}} = 0.1$.

Geometry	$C_{\text{crit}}^{(3B)}$	$C_{\text{crit}}^{(2B)}$	Window width
Equilateral	≈ 0.128	≈ 0.184	0.056
Linear	≈ 0.155	≈ 0.184	0.029

The linear-chain window is **twice as narrow** as the equilateral window because the Feynman integral $I_Y(d, 2d, d;\, m_{\text{sp}})$ is smaller than $I_Y(d, d, d;\, m_{\text{sp}})$ (the outer pair at $r_{13} = 2d$ is beyond the Yukawa screening length $\lambda = 10\, l_{\text{Pl}}$, so their three-body Feynman diagram is suppressed).

19.5 Isosceles deformation

Holding d fixed at the equilateral minimum $d_{\rm eq} = 4.52\, l_{\rm Pl}$ and varying the shape parameter $s = r_{23}/d$:

Table 23: Isosceles energy vs shape s at fixed $d = 4.52\, l_{\rm Pl}$, $C = 0.155$, $m_{\rm sp} = 0.1$.

$s = r_{23}/d$	r_{23} $[l_{\rm Pl}]$	$E_{\rm eff}$ $[E_{\rm Pl}]$
0.50	2.26	−0.01651
0.75	3.39	−0.01072
1.00	4.52	−0.00706 (equilateral)
1.25	5.64	−0.00440
1.50	6.77	−0.00232
2.00	9.03	+0.00077

The single-collective-coordinate ZP model predicts deeper binding as $s \to 0$ (third pair approaches $r_{23} \to 0$). This is an artefact of the 1D reduction: a full two-dimensional treatment would include a separate ZP term for the r_{23} degree of freedom. Within the present model, the *physically meaningful* comparison is at $s \simeq 1$; the equilateral configuration ($s = 1$) is the natural symmetric ground state.

19.6 Four-body square ($N = 4$)

The $N = 4$ square has $\binom{4}{3} = 4$ contributing three-body Feynman diagrams (each a right-triangle triple with legs d, d and hypotenuse $d\sqrt{2}$). Despite the larger ZP ($4/(8d^2)$), the additional V_3 contributions produce substantially deeper binding:

$$E_{\rm min}^{(N=4)} \approx -0.068\, E_{\rm Pl} \quad \text{at } d_{\rm side} \approx 2.5\, l_{\rm Pl}, \tag{80}$$

some $\sim 10\times$ deeper than the $N = 3$ equilateral triangle at the same C and $m_{\rm sp}$. This confirms the super-extensive growth of three-body binding with N: $\binom{N}{3}/N \sim N^2/6$ triples per body grows rapidly.

19.7 Summary

- Among symmetric three-body geometries, the **equilateral triangle** maximises V_3 (all pairs at equal separation, maximising the Feynman integral) and is the ground-state configuration.
- The **linear chain** is bound in a narrower window ($\Delta C \approx 0.029$ vs 0.056) and is $\sim 27\times$ more weakly bound than the equilateral at the window midpoint.
- Increasing N dramatically deepens binding via the $\binom{N}{3}$ growth of three-body terms.
- The falsifiable prediction is unaffected: triangular clusters are the dominant channel for three-body binding near Planck scale.

20 Quantum Efimov window: exact finite-difference result

20.1 Motivation: WKB validity

The WKB analysis of Sec. 18 located a single quasi-bound level in the classical Efimov window $C \in (0.128,\, 0.184)$ using the criterion $n(E) = \frac{1}{\pi}\int \sqrt{2\mu(E - V_{\rm eff})}\, dd \geq \frac{1}{2}$. The WKB approximation requires $\lambda_{\rm dB} \ll L_{\rm well}$, where $L_{\rm well}$ is the length scale of the potential well. At the WKB energy

$E_0^{\rm WKB} = -9\times10^{-4}\,E_{\rm Pl}$ (for $C = 0.180$), the de Broglie wavelength at the centre of the well ($d \approx 7$, $V_{\rm eff} \approx -0.011$) is

$$\lambda_{\rm dB} = \frac{2\pi}{\sqrt{2\mu(E_0^{\rm WKB} - V_{\rm eff})}} \approx \frac{2\pi}{\sqrt{2\times 0.010}} \approx 44\,l_{\rm Pl}, \tag{81}$$

far exceeding the well width $L_{\rm well} \approx 14\,l_{\rm Pl}$. **The WKB is invalid here.** A correct quantum treatment is required.

20.2 Finite-difference Hamiltonian

We solve the 1D Schrödinger equation in the collective coordinate d:

$$\hat{H}\psi = E\psi, \qquad \hat{H} = -\frac{1}{2\mu}\frac{d^2}{dd^2} + V_{\rm eff}(d), \tag{82}$$

with Dirichlet boundary conditions $\psi(d_{\rm lo}) = \psi(d_{\rm hi}) = 0$, $d_{\rm lo} = 0.4$, $d_{\rm hi} = 30\,l_{\rm Pl}$, $N = 3000$ grid points. The ZP term $3/(8d^2)$ is included in $V_{\rm eff}$ as the *hypercentrifugal* kinetic energy from the transverse degrees of freedom (the d-axis kinetic energy is added by $\hat{H}$).

We define:

$$V_{\rm 2B}(d; C, m_{\rm sp}) = \frac{3}{8d^2} - \frac{3C^2 e^{-m_{\rm sp}d}}{d}, \tag{83}$$

$$V_{\rm 3B}(d; C, m_{\rm sp}) = V_{\rm 2B}(d; C, m_{\rm sp}) - C^3\, I_Y(d, d, d;\, m_{\rm sp}). \tag{84}$$

The **classical Efimov window** (Sec. 18) is $\min_d V_{\rm 2B} > 0$ and $\min_d V_{\rm 3B} < 0$. The **quantum Efimov window** (this section) is $E_0^{\rm FD}(V_{\rm 2B}) > 0$ and $E_0^{\rm FD}(V_{\rm 3B}) < 0$, where $E_0^{\rm FD}$ is the lowest eigenvalue of $\hat{H}$.

20.3 Results: classical vs quantum windows

Table 24: Classical Efimov window (ex23) vs quantum window (ex26 FD), $m_{\rm sp} = 0.1\,l_{\rm Pl}^{-1}$.

Quantity	Classical (ex23)	Quantum FD (ex26)
$C_{\rm crit}^{(3B)}$ (3B binding threshold)	0.128	0.191
$C_{\rm crit}^{(2B)}$ (2B binding threshold)	0.184	0.310
Window width ΔC	0.056	**0.119**
Mass range [$m_{\rm Pl}$]	[0.45, 0.65]	[0.68, 1.10]

Both thresholds are shifted upward by $\Delta C \approx 0.063$–0.126 because the quantum kinetic energy in the d-mode ($E_{\rm kin}^{(d)} \approx 0.006\,E_{\rm Pl}$) must be overcome by the attractive potential. The window is **wider** in the quantum case ($\Delta C = 0.119$ vs 0.056) because the 3-body force adds steeper C^3 attraction, reaching the quantum threshold at lower C relative to the 2-body threshold.

Table 25: FD eigenvalues in the quantum Efimov window, $m_{\rm sp} = 0.1$.

C	$E_0^{\rm 2B}$ [$E_{\rm Pl}$]	$E_0^{\rm 3B}$ [$E_{\rm Pl}$]	$N_{\rm states}$	$d_{\rm peak}$ [$l_{\rm Pl}$]
0.128	+0.00722	+0.00618	0	—
0.184	+0.00608	+0.00122	0	—
0.192	+0.00587	−0.00027	1	8.87
0.207	+0.00540	−0.00419	1	7.12
0.280	+0.00220	−0.06637	2	3.56
0.310	+0.00051	−0.11036	2	—
0.312	−0.00016	−0.12838	2	—

20.4 Planck coupling is inside the quantum window

The TGP Planck coupling $C_{\rm Pl} = 1/(2\sqrt{\pi}) \approx 0.282$ lies *inside* the quantum Efimov window $C \in (0.191,\, 0.310)$:

$$\boxed{C_{\rm Pl} = 0.282 \in (C_Q^{(3B)},\, C_Q^{(2B)}) = (0.191,\, 0.310)} \tag{85}$$

At $C = C_{\rm Pl}$:

$$E_0^{\rm 2B}(C_{\rm Pl}) = +0.0022\, E_{\rm Pl} > 0 \quad \text{(triangle unbound by 2-body)}, \tag{86}$$

$$E_0^{\rm 3B}(C_{\rm Pl}) = -0.0664\, E_{\rm Pl} < 0 \quad \text{(triangle bound by 3-body!)}. \tag{87}$$

Planck-mass objects in TGP possess a quantum-mechanically bound 3B-induced triangular ground state.

20.5 Wavefunction at $C = 0.207$

At $C = 0.207$ (just above the 3B quantum threshold):

$$E_0 = -4.2 \times 10^{-3}\, E_{\rm Pl}, \quad d_{\rm peak} = 7.1\, l_{\rm Pl}, \quad \langle d \rangle = 9.8\, l_{\rm Pl}, \quad \sigma_d = 4.8\, l_{\rm Pl}. \tag{88}$$

The state is diffuse (spread over $\sim 5\, l_{\rm Pl}$) but well-localised compared to the box size ($d_{\rm hi} = 30\, l_{\rm Pl}$).

20.6 Correction to ex24 WKB

The WKB predicted a bound state at $E_0^{\rm WKB} = -9 \times 10^{-4}\, E_{\rm Pl}$ for $C = 0.180$. The exact FD gives $E_0^{\rm FD} = +1.8 \times 10^{-3}\, E_{\rm Pl}$ (unbound). The WKB was wrong because:

$$\lambda_{\rm dB} \approx 44\, l_{\rm Pl} \gg L_{\rm well} \approx 14\, l_{\rm Pl} \quad \text{(at } C = 0.180\text{)}. \tag{89}$$

The WKB quantisation condition $n + \frac{1}{2} = \frac{1}{\pi} \int \sqrt{2\mu(E - V)}\, dd$ overestimates binding for such wide, shallow wells.

20.7 Summary

Table 26: Summary: quantum Efimov window (ex26 FD), $m_{\rm sp} = 0.1$.

Property	Value
Quantum 3B threshold $C_Q^{(3B)}$	0.191
Quantum 2B threshold $C_Q^{(2B)}$	0.310
Quantum window width ΔC	0.119 (2.1× classical)
Mass range (quantum)	$[0.68,\, 1.10]\, m_{\rm Pl}$
$C_{\rm Pl}$ in window?	**Yes** ($C_{\rm Pl} = 0.282 \in (0.191, 0.310)$)
$E_0^{(3B)}(C_{\rm Pl})$	$-0.066\, E_{\rm Pl}$ (strongly bound)
WKB validity in classical window	No ($\lambda_{\rm dB} \gg L_{\rm well}$)
Max FD states in quantum window	2 (for $C \in (0.280, 0.310)$)

21 Quantum Efimov window versus scalar mass $m_{\rm sp}$

21.1 Question

The quantum Efimov window $C_Q^{(3B)} < C < C_Q^{(2B)}$ was computed at a fixed $m_{\rm sp} = 0.1\, l_{\rm Pl}^{-1}$ in Sec. 20. Here we ask: *over what range of $m_{\rm sp}$ does the Planck coupling $C_{\rm Pl} = 1/(2\sqrt{\pi}) \approx 0.282$ lie inside this window?*

This determines the physical parameter space for which Planck-mass objects would form quantum-mechanically bound triangular clusters solely through the three-body TGP interaction.

21.2 Method

For each $m_{\rm sp}$ in $[0.05,\ 1.00]\, l_{\rm Pl}^{-1}$:

1. Build the look-up table for $I_Y(d, d, d;\, m_{\rm sp})$ (Gauss–Legendre quadrature with 25 points per axis, $N = 350$ table nodes).
2. Solve the FD Schrödinger equation (Sec. 20) on a grid $d \in [0.4,\ 30\,]l_{\rm Pl}$, $N = 2000$ points.
3. Bisect in C to find $C_Q^{(3B)}$ (lowest C giving $E_0^{(3B)} < 0$) and $C_Q^{(2B)}$ (lowest C giving $E_0^{(2B)} < 0$).
4. Record whether $C_{\rm Pl} \in (C_Q^{(3B)},\, C_Q^{(2B)})$.

21.3 Results: window versus $m_{\rm sp}$

Table 27: Classical and quantum Efimov windows vs scalar mass $m_{\rm sp}$. $C_{\rm Pl} = 0.282$; "in?" $= C_{\rm Pl} \in (C_Q^{(3B)}, C_Q^{(2B)})$. "—" means no binding found up to $C = 0.70$.

$m_{\rm sp}$ $[l_{\rm Pl}^{-1}]$	λ $[l_{\rm Pl}]$	$C_{\rm cl}^{(3B)}$	$C_{\rm cl}^{(2B)}$	$\Delta C_{\rm cl}$	$C_Q^{(3B)}$	$C_Q^{(2B)}$	ΔC_Q	$C_{\rm Pl}$ in?
0.05	20.0	0.084	0.130	0.047	0.135	0.252	0.116	no
0.08	12.5	0.111	0.165	0.053	0.170	0.287	0.118	**YES**
0.10	10.0	0.128	0.184	0.057	0.191	0.310	0.119	**YES**
0.12	8.3	0.142	0.202	0.059	0.211	0.332	0.122	**YES**
0.15	6.7	0.163	0.226	0.063	0.239	0.365	0.126	**YES**
0.18	5.6	0.182	0.247	0.066	0.267	0.396	0.130	**YES**
0.20	5.0	0.193	0.261	0.067	0.284	0.416	0.132	no
0.25	4.0	0.221	0.291	0.071	0.326	0.464	0.139	no
0.30	3.3	0.246	0.319	0.074	0.365	0.510	0.145	no
0.40	2.5	0.290	0.369	0.078	0.440	0.598	0.157	no
0.50	2.0	0.331	0.412	0.082	0.512	0.680	0.168	no
0.60	1.7	0.367	0.452	0.084	0.583	—	—	no
0.80	1.2	0.433	0.521	0.089	—	—	—	no
1.00	1.0	0.491	0.583	0.092	—	—	—	no

Observations:

- The quantum window is *consistently* $\approx 2\times$ wider than the classical window ($\Delta C_Q/\Delta C_{\rm cl} \approx$ 2.0–2.5 for all $m_{\rm sp}$).
- Both thresholds *increase with* $m_{\rm sp}$: stronger coupling is needed to bind tighter (shorter-range) interactions.
- For $m_{\rm sp} \gtrsim 0.6\, l_{\rm Pl}^{-1}$ the two-body quantum threshold $C_Q^{(2B)} > 0.70$ leaves the scanned range; for $m_{\rm sp} \gtrsim 0.8$ even the three-body threshold disappears. The quantum window effectively *closes* at $m_{\rm sp} \sim 0.7\, l_{\rm Pl}^{-1}$ ($\lambda \sim 1.4\, l_{\rm Pl}$).

21.4 Critical scalar mass range for Planck objects

Exact bisection (30 iterations, $\delta C < 10^{-9}$) of the condition $C_Q^{(3B)}(m_{\rm sp}) = C_{\rm Pl}$ and $C_Q^{(2B)}(m_{\rm sp}) = C_{\rm Pl}$ gives:

$$\boxed{C_{\rm Pl} \in (C_Q^{(3B)},\, C_Q^{(2B)}) \iff m_{\rm sp} \in (0.076,\ 0.198)\, l_{\rm Pl}^{-1}} \tag{90}$$

or equivalently in terms of the scalar range $\lambda_{\rm sp} = 1/m_{\rm sp}$:

$$\lambda_{\rm sp} \in (5.1,\ 13.2)\, l_{\rm Pl} \quad \Longrightarrow \quad \text{Planck-mass 3-body quantum bound state exists.} \tag{91}$$

The window corresponds to a Compton wavelength of the scalar ϕ mediator in the range

$$\lambda_{\rm sp} \sim 5\text{–}13\, l_{\rm Pl} = 8\text{–}21 \times 10^{-35}\,\text{m}. \tag{92}$$

21.5 Significance: non-trivial intersection

That $C_{\rm Pl}$ lies inside the quantum window for a *finite, non-fine-tuned* range of $m_{\rm sp}$ is a non-trivial result:

1. At $m_{\rm sp} = 0$ (massless scalar) the Yukawa interaction becomes Coulomb-like; both thresholds shift to $C = 0$ and the window is not well-defined (all couplings bind).
2. At very small $m_{\rm sp} \ll 0.076$ (long-range Yukawa, $\lambda \gg 13\, l_{\rm Pl}$) the 2-body quantum threshold satisfies $C_Q^{(2B)} < C_{\rm Pl}$: *pairs* also bind quantum-mechanically, so there is no Efimov-type 3B-only window at the Planck coupling. Note that pairs also bind *classically* in this regime.
3. At $m_{\rm sp} \gg 0.198$ (short-range Yukawa) the 3-body quantum threshold $C_Q^{(3B)} > C_{\rm Pl}$: the Planck coupling is insufficient to sustain quantum 3-body binding. A *classical* Efimov window exists for $m_{\rm sp} \in (0.235,\ 0.38)$, but quantum zero-point energy destroys it.
4. In the intermediate range $m_{\rm sp} \in (0.076,\ 0.198)$, $\lambda_{\rm sp} \in (5.1,\ 13.2)\, l_{\rm Pl}$: the pair is classically bound but quantum-mechanically unbound, while the triple is quantum-mechanically bound. The triangular cluster is a *genuine quantum Efimov analog*: purely three-body, with no corresponding two-body bound state.

21.6 Comparison: classical versus quantum window membership

Table 28: Whether $C_{\rm Pl}$ is in the classical vs quantum Efimov window as a function of $m_{\rm sp}$.

$m_{\rm sp}$	λ	Classical	Quantum	Interpretation
< 0.076	> 13.2	no	no	pairs bind at $C_{\rm Pl}$ (classical & quantum)
0.076–0.198	5.1–13.2	no	**YES**	pairs classically bound; quantum 3B only
0.198–~ 0.235	~ 4.3–5.1	no	no	pairs classically bound; quantum unbound
~ 0.235–~ 0.38	~ 2.6–4.3	**YES**	no	classical Efimov; quantum unbound
$>\sim 0.38$	$<\sim 2.6$	no	no	neither binds at $C_{\rm Pl}$

Key observation: the quantum Efimov window and the classical Efimov window are *non-overlapping* in $m_{\rm sp}$. For $m_{\rm sp} \in (0.076,\ 0.198)$ the pair is *classically* bound but *quantum-mechanically unbound* (zero-point energy overcomes the shallow classical well); simultaneously, the three-body quantum state IS bound. For $m_{\rm sp} \in (0.235,\ 0.38)$ the classical picture gives an Efimov state (3B only) but quantum fluctuations destroy it. This demonstrates that classical and quantum analyses give qualitatively opposite conclusions across most of the parameter space, underscoring the necessity of the exact FD treatment.

21.7 Summary

Main conclusion: Planck-mass objects in TGP form quantum-mechanically bound three-body triangular clusters *if and only if* the scalar mediator range satisfies $\lambda_{\rm sp} \in (5.1,\ 13.2)\, l_{\rm Pl}$. This is a sharp, parameter-space-resolved prediction. For all known sub-Planck particles ($C \sim 10^{-20}$) the coupling is far below both thresholds, so the effect is absent.

Table 29: Key parameters: quantum Efimov window at $C_{\rm Pl}$.

Quantity	Value
$m_{\rm sp}$ window (lower)	$0.076\, l_{\rm Pl}^{-1}$ ($\lambda = 13.2\, l_{\rm Pl}$)
$m_{\rm sp}$ window (upper)	$0.198\, l_{\rm Pl}^{-1}$ ($\lambda = 5.05\, l_{\rm Pl}$)
Window width $\Delta m_{\rm sp}$	$0.122\, l_{\rm Pl}^{-1}$ (factor $\times 2.6$)
Ratio $\Delta C_Q/\Delta C_{\rm cl}$	≈ 2.0 (independent of $m_{\rm sp}$)
$C_{\rm Pl}$ inside quantum window?	Yes, for $\lambda_{\rm sp} \in (5.1, 13.2)\, l_{\rm Pl}$
Quantum window closes at	$m_{\rm sp} \approx 0.7\, l_{\rm Pl}^{-1}$ ($\lambda \approx 1.4\, l_{\rm Pl}$)

22 Polygon binding: N=3, 4, 5 bodies

22.1 Setup

We compare regular N-polygon clusters — equilateral triangle ($N = 3$), square ($N = 4$), and regular pentagon ($N = 5$) — each parameterised by the nearest-neighbour side length d. For each geometry the effective energy is

$$E_{\rm eff}(d;\, C, m_{\rm sp}) = \underbrace{\frac{N}{8d^2}}_{\rm ZP} + V_2(d;\, C, m_{\rm sp}) + V_3(d;\, C, m_{\rm sp}), \tag{93}$$

with the same zero-point convention $\mathrm{ZP} = N/(8d^2)$, two-body sum $V_2 = -C^2 \sum_{\rm pairs} e^{-m_{\rm sp} r_{ij}}/r_{ij}$, and three-body sum $V_3 = -C^3 \sum_{\rm triples} I_Y(r_{ij}, r_{ik}, r_{jk};\, m_{\rm sp})$.

Pentagon geometry. For $N = 5$ with side d the golden ratio $\varphi = 2\cos(\pi/5) \approx 1.618$ gives the two distinct inter-vertex distances:

$$\text{5 adjacent pairs:} \qquad r = d, \tag{94}$$

$$\text{5 diagonal pairs:} \qquad r = \varphi d. \tag{95}$$

The $\binom{5}{3} = 10$ triples split into two classes:

$$\text{Type A (consecutive, 5):} \qquad (d,\ d,\ \varphi d), \tag{96}$$

$$\text{Type B (skip-one, 5):} \qquad (d,\ \varphi d,\ \varphi d). \tag{97}$$

22.2 Binding energies at $C = 0.155$

Table 30 compares the classical energy minimum for $m_{\rm sp} = 0.1\, l_{\rm Pl}^{-1}$ at the midpoint of the equilateral 3B-only window.

Table 30: Binding energies for regular N-gons. $m_{\rm sp} = 0.1\, l_{\rm Pl}^{-1}$, $C = 0.155$. $E_{\rm min}^{(2B)}$ uses ZP+V_2 only; $E_{\rm min}^{(2B+3B)}$ adds V_3.

N	Geometry	Triples $\binom{N}{3}$	$E_{\rm min}^{(2B)}$ $[E_{\rm Pl}]$	$E_{\rm min}^{(3B)}$ $[E_{\rm Pl}]$	$d_{\rm eq}$ $[l_{\rm Pl}]$	V_3/V_2	Bound?
3	Equilateral	1	$+3.6 \times 10^{-4}$	-7.1×10^{-3}	4.52	1.51	YES
4	Square	4	$+4.5 \times 10^{-4}$	-6.9×10^{-2}	2.42	2.74	YES
5	Pentagon	10	$+3.6 \times 10^{-4}$	-2.3×10^{-1}	1.69	3.76	YES

All three geometries are in the 3B-only window at $C = 0.155$: the two-body energy is positive (pairs unbound) while the full two-body+three-body energy is negative (cluster bound). The binding energy grows by a factor ~ 32 from $N = 3$ to $N = 5$.

Physical origin. The dramatic growth comes from two cooperating effects:

1. **More triples**: $\binom{N}{3}$ grows as $N^3/6$, giving $1 \to 4 \to 10$ for $N = 3, 4, 5$.
2. **Shorter equilibrium separation**: $d_{\rm eq}$ decreases from $4.5 \to 2.4 \to 1.7\, l_{\rm Pl}$, pushing the cluster deeper into the high-I_Y short-distance regime.

The V_3/V_2 ratio — a measure of three-body dominance — rises from 1.5 to 3.8, confirming that the cluster becomes progressively more "three-body driven" as N increases.

22.3 Critical coupling thresholds

Table 31: Classical critical couplings and 3B-only windows for regular N-gons at $m_{\rm sp} = 0.1\, l_{\rm Pl}^{-1}$. $C_{\rm Pl} = 0.282$ lies outside all classical windows for this $m_{\rm sp}$.

N	Geometry	$C_{\rm cl}^{(2B)}$	$C_{\rm cl}^{(3B)}$	ΔC	$C_{\rm Pl}$ in?
3	Equilateral	0.184	0.128	0.057	no
4	Square	0.166	0.102	0.064	no
5	Pentagon	0.159	0.091	0.068	no

Key trend: both thresholds *decrease* with N, because a larger cluster has more 3-body interactions. Specifically:

- $C_{\rm cl}^{(3B)}$ decreases from $0.128 \to 0.102 \to 0.091$: it takes less coupling per triple to bind a larger polygon.
- $C_{\rm cl}^{(2B)}$ also decreases ($0.184 \to 0.166 \to 0.159$) because the additional Yukawa pairs in larger polygons aid two-body binding.
- The window width ΔC *increases* slightly: $0.057 \to 0.064 \to 0.068$.

The 3B-only window *shifts downward* as N increases. At $m_{\rm sp} = 0.1\, l_{\rm Pl}^{-1}$, the window lies in $C \in (0.091, 0.184)$ for any of the three polygon types, but $C_{\rm Pl} = 0.282$ lies well above $C_{\rm cl}^{(2B)}$ for all N. This means that at the Planck coupling *all* polygon geometries have their pairs classically bound; the purely-three-body classical window does not extend to $C_{\rm Pl}$ for $m_{\rm sp} = 0.1$.

22.4 Pentagon Feynman integrals

At the pentagon equilibrium ($d_{\rm eq} = 1.69\, l_{\rm Pl}$, $C = 0.155$):

$$I_Y^{(A)}(d, d, \varphi d) = 10.16, \tag{98}$$

$$I_Y^{(B)}(d, \varphi d, \varphi d) = 8.77, \tag{99}$$

$$\sum_{\rm triples} I_Y = 5 \times 10.16 + 5 \times 8.77 = 94.7. \tag{100}$$

For comparison, the equilateral triangle at its equilibrium ($d_{\rm eq} = 4.52\, l_{\rm Pl}$) gives $I_Y(d, d, d) = 4.10$. The pentagon integral is larger primarily because $d_{\rm eq}$ is smaller (shorter-range Yukawa is stronger at close range).

22.5 Comparison at $C = C_{\rm Pl}$

At $C_{\rm Pl} = 0.282$, all pair energies are negative; there is no 3B-only classical window. However, the absolute binding energy grows dramatically:

The 3-body contribution at $C_{\rm Pl}$ dominates: pentagon binding ($-3.73\, E_{\rm Pl}$) is 14× stronger than the equilateral ($-0.27\, E_{\rm Pl}$), while the 2-body energy ratio is only 5×.

Table 32: Binding energies at $C = C_{\rm Pl} = 0.282$, $m_{\rm sp} = 0.1\, l_{\rm Pl}^{-1}$. All geometries are classically bound (pairs also bind), so no Efimov-type 3B-only window exists at this coupling.

N	Geometry	$E_{\rm min}^{(2B)}$ $[E_{\rm Pl}]$	$E_{\rm min}^{(2B+3B)}$ $[E_{\rm Pl}]$
3	Equilateral	-0.018	-0.265
4	Square	-0.051	-1.350
5	Pentagon	-0.096	-3.729

22.6 Summary

Table 33: Polygon geometry comparison summary, $m_{\rm sp} = 0.1\, l_{\rm Pl}^{-1}$.

Property	$N = 3$	$N = 4$	$N = 5$
# pairs	3	6	10
# triples	1	4	10
$C_{\rm cl}^{(3B)}$	0.128	0.102	0.091
$C_{\rm cl}^{(2B)}$	0.184	0.166	0.159
ΔC window	0.057	0.064	0.068
$d_{\rm eq}$ $(C = 0.155)$	$4.5\, l_{\rm Pl}$	$2.4\, l_{\rm Pl}$	$1.7\, l_{\rm Pl}$
$E_{\rm bind}$ $(C = 0.155)$	7×10^{-3}	6.9×10^{-2}	2.3×10^{-1}
V_3/V_2 ratio	1.51	2.74	3.76

Main conclusion: TGP three-body forces create bound polygon clusters for all $N = 3, 4, 5$ in an Efimov-like 3B-only window. As N increases: (i) the binding energy grows superlinearly ($\sim N^3$), (ii) the equilibrium separation shrinks, (iii) the window shifts to lower C, and (iv) three-body forces become progressively more dominant. The pentagon cluster is $\sim 32\times$ more tightly bound than the equilateral triangle at the same coupling constant.

23 Ground-state wavefunction at $C = C_{\rm Pl}$

23.1 Setup

We solve the 1D Schrödinger equation in the collective coordinate d (equilateral triangle side length):

$$\left[-\frac{1}{2\mu}\frac{d^2}{dd^2} + V_{\rm eff}(d;\, C, m_{\rm sp})\right]\psi(d) = E\,\psi(d), \tag{101}$$

with $V_{\rm eff} = \frac{3}{8d^2} + V_2 + V_3$, $\mu = 1\, m_{\rm Pl}$, Dirichlet boundary conditions at $d = 0.4$ and $d = 30\, l_{\rm Pl}$, and $N = 3000$ finite-difference grid points (same as ex26/ex27).

23.2 Ground-state energy vs $m_{\rm sp}$

The quantum window $E_0^{(2B)} > 0$, $E_0^{(3B)} < 0$ at $C_{\rm Pl}$ spans exactly $m_{\rm sp} \in (0.076,\, 0.198)\, l_{\rm Pl}^{-1}$ (consistent with Sec. 21).

23.3 Detailed wavefunction at $m_{\rm sp} = 0.1$

Quantum delocalisation. The quantum mean position $\langle d \rangle = 4.52\, l_{\rm Pl}$ is 3.3 times larger than the classical equilibrium $d_{\rm cl} = 1.37\, l_{\rm Pl}$. This dramatic delocalisation is a direct consequence of the WKB invalidity: the wavefunction is spread over the entire attractive region of the potential (0.7–6 $l_{\rm Pl}$), and the de Broglie wavelength ($16\, l_{\rm Pl}$) exceeds the well width ($5.3\, l_{\rm Pl}$) by a factor ~ 3.

Table 34: FD ground-state energies at $C = C_{\rm Pl} = 0.282$. "in win?" $= E_0^{(2B)} > 0$ and $E_0^{(3B)} < 0$ (3B-only quantum window).

$m_{\rm sp}$ $[l_{\rm Pl}^{-1}]$	λ $[l_{\rm Pl}]$	$E_0^{(2B)}$ $[E_{\rm Pl}]$	$E_0^{(3B)}$ $[E_{\rm Pl}]$	Bound?	In win?
0.050	20.0	−0.00298	−0.2021	yes	no (2B also)
0.076	13.2	+0.000031	−0.1168	yes	**YES** (boundary)
0.080	12.5	+0.000416	−0.1073	yes	**YES**
0.100	10.0	+0.00207	−0.0696	yes	**YES**
0.120	8.3	+0.00336	−0.0435	yes	**YES**
0.150	6.7	+0.00477	−0.0183	yes	**YES**
0.180	5.6	+0.00572	−0.00449	yes	**YES**
0.198	5.1	+0.00614	+0.000050	no	no (boundary)
0.200	5.0	+0.00618	+0.000428	no	no
0.250	4.0	+0.00695	+0.00533	no	no

Table 35: Ground-state wavefunction properties at $C_{\rm Pl} = 0.282$, $m_{\rm sp} = 0.1\, l_{\rm Pl}^{-1}$, $\mu = 1\, m_{\rm Pl}$.

Quantity	Value
E_0	$-0.06964\, E_{\rm Pl} = -1.36 \times 10^8$ J
$d_{\rm peak}$ (density maximum)	$3.51\, l_{\rm Pl}$
$\langle d \rangle$ (mean separation)	$4.52\, l_{\rm Pl}$
$\sigma_d = \sqrt{\langle d^2 \rangle - \langle d \rangle^2}$	$2.10\, l_{\rm Pl}$
$\sigma_d / \langle d \rangle$	0.46
Classical minimum $d_{\rm cl}$	$1.37\, l_{\rm Pl}$
$\langle d \rangle / d_{\rm cl}$	3.30
d_L (left classical turning pt.)	$0.73\, l_{\rm Pl}$
d_R (right classical turning pt.)	$6.03\, l_{\rm Pl}$
$L_{\rm well} = d_R - d_L$	$5.29\, l_{\rm Pl}$
$\lambda_{\rm dB}$ at $d_{\rm peak}$	$16.0\, l_{\rm Pl}$
$\lambda_{\rm dB}/L_{\rm well}$	3.03 (WKB **invalid**)
Tunnelling fraction	20.8%
V_3/V_2 at $\langle d \rangle$	2.74
Number of bound states	2

Three-body dominance. At the mean position $\langle d \rangle = 4.52\, l_{\rm Pl}$:

$$V_2(\langle d \rangle) = -0.034\, E_{\rm Pl}, \qquad V_3(\langle d \rangle) = -0.092\, E_{\rm Pl}, \qquad V_3/V_2 = 2.74. \tag{102}$$

The three-body Feynman force contributes 2.7× more binding than the two-body Yukawa. This confirms that the cluster is sustained *primarily* by the three-body interaction.

Tunnelling. Approximately 21% of the probability density lies outside the classical turning points $[d_L,\, d_R] = [0.73,\, 6.03]\, l_{\rm Pl}$. This significant tunnelling fraction reflects the quantum-mechanical character of the state.

23.4 Second bound state

At $m_{\rm sp} = 0.1$, $C = C_{\rm Pl}$, the FD solver finds *two* bound states:

The ratio of spatial scales $d_{\rm peak}^{(1)}/d_{\rm peak}^{(0)} \approx 3.8$ is far below the classical Efimov ratio $e^{\pi/s_0} \approx 22.7$, confirming that TGP does *not* produce an infinite geometric tower. The Yukawa screening caps the number of bound levels at ≤ 2.

Table 36: Bound states at $C_{\rm Pl}$, $m_{\rm sp} = 0.1$.

n	E_n $[E_{\rm Pl}]$	$d_{\rm peak}$ $[l_{\rm Pl}]$	$\langle d\rangle$ $[l_{\rm Pl}]$	σ_d $[l_{\rm Pl}]$
0	-0.06964	3.51	4.52	2.10
1	-0.000651	13.30	14.42	5.14

23.5 Wavefunction evolution near window boundary

Table 37: Ground-state wavefunction moments vs $m_{\rm sp}$ at $C = C_{\rm Pl}$.

$m_{\rm sp}$ $[l_{\rm Pl}^{-1}]$	λ $[l_{\rm Pl}]$	E_0 $[E_{\rm Pl}]$	$d_{\rm cl}$ $[l_{\rm Pl}]$	$\langle d\rangle$ $[l_{\rm Pl}]$	σ_d $[l_{\rm Pl}]$	Tunnel [%]
0.08	12.5	-0.1073	1.36	4.17	1.87	19.6
0.10	10.0	-0.0696	1.36	4.52	2.10	20.8
0.12	8.3	-0.0435	1.40	4.98	2.40	22.5
0.15	6.7	-0.0183	1.44	6.04	3.17	26.3
0.18	5.6	-0.00449	1.48	8.05	4.47	32.1

As $m_{\rm sp}$ increases toward the upper boundary (0.198): the binding energy decreases, $\langle d\rangle$ grows (state expands), σ_d grows (state delocalises), and the tunnelling fraction increases — all characteristic signatures of a bound state approaching the dissociation threshold. The state at $m_{\rm sp} = 0.18$ ($E_0 = -0.0045\,E_{\rm Pl}$, $\langle d\rangle = 8.1\,l_{\rm Pl}$, tunnel= 32%) is at the verge of unbinding, confirming the sharp upper boundary of the quantum Efimov window.

24 Physical interpretation: window bounds and classical/quantum non-overlap

24.1 Why does the quantum Efimov window exist for $\lambda_{\rm sp} \in (5.1,\, 13.2)\,l_{\rm Pl}$?

The quantum Efimov window $C_{\rm Pl} \in (C_Q^{(3B)},\, C_Q^{(2B)})$ exists for scalar range $\lambda_{\rm sp} \in (5.1,\, 13.2)\,l_{\rm Pl}$. Each boundary has a clear physical origin.

Upper boundary $\lambda_{\rm sp}^{\rm upper} = 13.2\,l_{\rm Pl}$ ($m_{\rm sp} = 0.076\,l_{\rm Pl}^{-1}$): As the scalar range $\lambda_{\rm sp}$ increases, the Yukawa two-body potential becomes more long-range and the two-body quantum threshold $C_Q^{(2B)}(\lambda_{\rm sp})$ decreases. At $\lambda_{\rm sp} = 13.2\,l_{\rm Pl}$ the threshold reaches the Planck coupling:

$$C_Q^{(2B)}(m_{\rm sp} = 0.076) = C_{\rm Pl} = 0.282. \tag{103}$$

For $\lambda_{\rm sp} > 13.2\,l_{\rm Pl}$ (longer range), pairs quantum-mechanically bind at $C_{\rm Pl}$, so the three-body cluster is no longer a "Efimov-type" purely three-body state — pairs already bind by themselves.

Physical meaning: $\lambda^{\rm upper}$ is the Yukawa range at which the Planck coupling first becomes strong enough to form quantum-mechanically bound *pairs*. Above this range, two-body physics dominates.

Lower boundary $\lambda_{\rm sp}^{\rm lower} = 5.1\,l_{\rm Pl}$ ($m_{\rm sp} = 0.198\,l_{\rm Pl}^{-1}$): As the scalar range decreases (heavier mediator), the three-body Feynman integral $I_Y(d, d, d;\, m_{\rm sp})$ becomes more rapidly screened for $d \gg \lambda_{\rm sp}$, reducing the range over which three-body attraction acts. The three-body quantum threshold $C_Q^{(3B)}$ therefore increases. At $\lambda_{\rm sp} = 5.1\,l_{\rm Pl}$:

$$C_Q^{(3B)}(m_{\rm sp} = 0.198) = C_{\rm Pl} = 0.282. \tag{104}$$

For $\lambda_{\rm sp} < 5.1\, l_{\rm Pl}$ (shorter range), the Planck coupling is insufficient to bind a quantum three-body cluster — the zero-point kinetic energy $3/(8d^2)$ in the collective coordinate overcomes the screened attraction.

Physical meaning: $\lambda^{\rm lower}$ is the Yukawa range at which the three-body TGP interaction becomes too short-ranged to support a quantum bound cluster at $C_{\rm Pl}$. Below this range, zero-point pressure wins.

Summary of window origin. The window $\lambda_{\rm sp} \in (5.1,\, 13.2)\, l_{\rm Pl}$ is the range in which:

- The scalar is *long-range enough* that three-body attraction overcomes zero-point pressure (requires $\lambda_{\rm sp} > 5.1\, l_{\rm Pl}$).
- The scalar is *short-range enough* that pairs remain quantum-mechanically unbound at $C_{\rm Pl}$ (requires $\lambda_{\rm sp} < 13.2\, l_{\rm Pl}$).

The window width $\Delta\lambda = 8.1\, l_{\rm Pl}$ (factor $\times 2.6$ in $m_{\rm sp}$) is not fine-tuned.

Connection to wavefunction (ex29). The ground-state wavefunction at $C_{\rm Pl}$, $m_{\rm sp} = 0.1$ peaks at $d_{\rm peak} = 3.5\, l_{\rm Pl}$ with $\langle d \rangle = 4.5\, l_{\rm Pl}$. This is well within the Yukawa range $\lambda = 10\, l_{\rm Pl}$, explaining why the three-body force is effective: the cluster lives where $I_Y(d, d, d;\, m_{\rm sp})$ is large. At the lower boundary ($\lambda = 5.1\, l_{\rm Pl}$, $m_{\rm sp} = 0.198$), the wavefunction would need to sit at $d \lesssim \lambda = 5.1\, l_{\rm Pl}$ to feel the three-body force, but zero-point energy pushes it outward — the potential well is too narrow to contain the state (from section C of ex29: $\langle d \rangle$ grows from 4.2 to $8.1\, l_{\rm Pl}$ as $m_{\rm sp}$ approaches 0.198).

24.2 Why are the classical and quantum windows non-overlapping?

The central result of Sec. 21 is that the classical Efimov window for $C_{\rm Pl}$ (at $m_{\rm sp} \in (0.235,\, 0.38)$) and the quantum Efimov window (at $m_{\rm sp} \in (0.076,\, 0.198)$) are *mutually exclusive.* We now explain why.

24.2.1 The role of zero-point energy

The classical and quantum treatments of the collective coordinate d differ in how zero-point kinetic energy enters:

- **Classical:** The system minimises $E_{\rm eff}(d) = \frac{3}{8d^2} + V_2 + V_3$ over d. The ZP term acts as a static repulsive "pressure" at small d. Binding requires $\min_d E_{\rm eff} < 0$.
- **Quantum:** The operator $\hat{T} = -\frac{1}{2\mu}\frac{d^2}{dd^2}$ contributes kinetic energy that depends on the wavefunction curvature. The ground-state energy $E_0 = \langle\psi|\hat{H}|\psi\rangle$ includes both potential *and* kinetic contributions. Since ψ is spread over $\sigma_d \sim 2\, l_{\rm Pl}$ (ex29), it samples a range of $V_{\rm eff}$ values around the minimum, raising the effective energy.

The net effect is that the quantum criterion for binding ($E_0 < 0$) requires a *deeper and broader* potential well than the classical criterion ($E_{\rm eff}^{\rm min} < 0$). This raises both thresholds: $C_Q^{(3B)} > C_{\rm cl}^{(3B)}$ and $C_Q^{(2B)} > C_{\rm cl}^{(2B)}$, and the quantum window is shifted upward in C by ≈ 0.063 at $m_{\rm sp} = 0.1$.

24.2.2 Why the shift causes non-overlap

At fixed $C = C_{\rm Pl} = 0.282$:

1. **Classical window** ($m_{\rm sp} \in (0.235,\, 0.38)$, short-range Yukawa, $\lambda \in (2.6,\, 4.3)\, l_{\rm Pl}$): Here $C_{\rm cl}^{(3B)} < C_{\rm Pl} < C_{\rm cl}^{(2B)}$. The classical equilibrium separation is $d_{\rm cl} \sim 1.5$–$2\, l_{\rm Pl}$, well within the Yukawa range λ. However, at these short Yukawa ranges the potential well is very

narrow (width $\Delta d \sim 2\, l_{\rm Pl}$) and shallow. The quantum ground-state wavefunction in this shallow well has $\sigma_d \sim \Delta d$, so kinetic energy is large and $E_0 = E_{\rm eff}^{\rm min} + \langle \hat{T} \rangle \gg 0$ — zero-point energy destroys the binding. In ex29 at $m_{\rm sp} = 0.18$ (close to the classical window): $\langle d \rangle = 8.1\, l_{\rm Pl} \gg \lambda = 5.6\, l_{\rm Pl}$ and the three-body force is effectively absent.

2. **Quantum window** ($m_{\rm sp} \in (0.076,\, 0.198)$, longer-range, $\lambda \in (5.1,\, 13.2)\, l_{\rm Pl}$): Here $C_Q^{(3B)} < C_{\rm Pl} < C_Q^{(2B)}$. The potential well is broader (extends over $d \sim 1$–$6\, l_{\rm Pl}$ within the Yukawa range), so the wavefunction can be contained while zero-point energy remains manageable. The quantum bound state exists precisely because the broader range allows $\langle d \rangle \sim 4.5\, l_{\rm Pl} \lesssim \lambda = 10\, l_{\rm Pl}$ (ex29, $m_{\rm sp} = 0.1$). However, classically the pair threshold $C_{\rm cl}^{(2B)} \sim 0.18$ is already below $C_{\rm Pl}$, so pairs *classically* bind — the classical Efimov-type 3B-only window at $C_{\rm Pl}$ does not exist in this range of $m_{\rm sp}$.

24.2.3 Schematic

Table 38: Physical mechanism behind classical/quantum window separation.

$m_{\rm sp}$ range	$\lambda_{\rm sp}$	Classical binding	Quantum binding
< 0.076 ($\lambda > 13.2$)	Long-range	pairs + triples bind	pairs bind at $C_{\rm Pl}$
0.076–0.198 ($\lambda =$ 5–13)	Medium	pairs bind classically, no 3B-only window	3B-only window: **quantum Efimov**
0.198–0.235 ($\lambda =$ 4.3–5.1)	Medium-short	pairs bind classically	unbound
0.235–0.38 ($\lambda =$ 2.6–4.3)	Short	3B-only classical Efimov	unbound: ZP destroys it
> 0.38 ($\lambda < 2.6$)	Very short	neither binds	neither

24.2.4 Key insight: WKB invalidity as a diagnostic

A quantitative signature of the non-overlap is the WKB ratio $\lambda_{\rm dB}/L_{\rm well}$ at the quantum ground state (ex29):

$$\frac{\lambda_{\rm dB}}{L_{\rm well}} = \frac{16.0}{5.3} \approx 3.0 \quad (m_{\rm sp} = 0.1,\ C = C_{\rm Pl}), \tag{105}$$

which is $\gg 1$, confirming deep quantum mechanics. In the classical Efimov regime (short-range Yukawa), the well would be narrow enough that the WKB ratio approaches unity — but the state is then pushed out by zero-point energy. The two regimes are incompatible at $C_{\rm Pl}$.

24.3 Quantum state characterisation at $C = C_{\rm Pl}$

The ground state at $C_{\rm Pl} = 0.282$, $m_{\rm sp} = 0.1\, l_{\rm Pl}^{-1}$ (ex29) has the following properties:
The second bound state ($n = 1$) has $E_1 = -6.5 \times 10^{-4}\, E_{\rm Pl}$ and peaks at $d_{\rm peak}^{(1)} = 13.3\, l_{\rm Pl}$ — a very diffuse, weakly-bound excited state analogous to the first Efimov excited state in nuclear/atomic physics. The ratio of spatial scales $d_{\rm peak}^{(1)}/d_{\rm peak}^{(0)} \approx 3.8$ is much smaller than the classical Efimov ratio $e^{\pi/s_0} \approx 22.7$, confirming that TGP does not produce an infinite geometric tower. The Yukawa screening limits the well depth, allowing at most two bound levels at $C_{\rm Pl}$.

As $m_{\rm sp}$ approaches the upper boundary $0.198\, l_{\rm Pl}^{-1}$, the bound state becomes increasingly diffuse ($\langle d \rangle$ grows from 4.2 to $8.1\, l_{\rm Pl}$, tunnelling from 20% to 32%) — a classic signature of a state approaching the dissociation threshold.

Table 39: Ground-state wavefunction properties at $C_{\rm Pl}$, $m_{\rm sp} = 0.1$.

Quantity	Value
E_0	$-0.0696\, E_{\rm Pl} = -1.36 \times 10^8$ J
$d_{\rm peak}$	$3.51\, l_{\rm Pl} = 5.45 \times 10^{-35}$ m
$\langle d \rangle$	$4.52\, l_{\rm Pl}$
σ_d	$2.10\, l_{\rm Pl}$
$\sigma_d/\langle d \rangle$	0.46 (broad, non-classical)
$d_{\rm cl}$ (classical minimum)	$1.37\, l_{\rm Pl}$
$\langle d \rangle / d_{\rm cl}$	3.30 (quantum delocalisation)
$\lambda_{\rm dB}/L_{\rm well}$	3.03 (WKB invalid)
Tunnelling fraction	20.8%
V_3/V_2 at $\langle d \rangle$	2.74 (3-body dominant)
Number of bound states	2

24.4 Summary

1. The quantum Efimov window bounds at $\lambda \approx 5$ and $13\, l_{\rm Pl}$ are set by the two-body and three-body quantum thresholds coinciding with $C_{\rm Pl}$ — not by any fine-tuning.

2. The classical and quantum windows are non-overlapping because the zero-point energy operator raises both thresholds by ≈ 0.063 in C, shifting the quantum window to longer Yukawa ranges where the potential well is broad enough to contain the wavefunction.

3. The quantum ground state at $C_{\rm Pl}$ is deeply non-classical: $\langle d \rangle / d_{\rm cl} = 3.3$, WKB ratio = 3.0, tunnelling = 21%. Classical methods give qualitatively wrong predictions for the existence and location of bound states.

4. A second bound state exists at $m_{\rm sp} = 0.1$, $C = C_{\rm Pl}$, at $d \sim 13\, l_{\rm Pl}$. This analogue of the first Efimov excitation has no infinite tower — Yukawa screening limits the count to ≤ 2.

25 Triangle shape-space stability and the equilateral approximation

25.1 Question

Sections 18–23 solve the Schrödinger equation along the equilateral *breathing mode* $d_{12} = d_{13} = d_{23} = d$. Is the equilateral configuration actually a stable energy minimum in shape space, or is it a saddle point? The answer determines whether the 1D reduction is exact or approximate.

25.2 Parameterisation of shape deformations

We parameterise deviations from equilateral by two dimensionless parameters:

$$d_{12} = d(1 + \varepsilon_a), \quad d_{13} = d(1 + \varepsilon_b), \quad d_{23} = d. \tag{106}$$

The ZP term uses $\bar{d} = (d_{12} + d_{13} + d_{23})/3$. At $(\varepsilon_a, \varepsilon_b) = (0, 0)$ the triangle is equilateral.

By S_3 symmetry, the full Hessian in the 3-distance space is $H^{(3)}_{ij} = \alpha\, \delta_{ij} + \beta\, (1 - \delta_{ij})$, with two eigenvalue families:

$$\text{Breathing mode:} \quad \lambda_{\rm br} = \alpha + 2\beta \quad \text{(equilateral scaling)}, \tag{107}$$

$$\text{Shape modes (2-fold degenerate):} \quad \lambda_{\rm sh} = \alpha - \beta. \tag{108}$$

25.3 Numerical results at $C = C_{\rm Pl}$, $m_{\rm sp} = 0.1$

Classical equilibrium ($d_{\rm cl} = 1.37\, l_{\rm Pl}$):

$$H = \begin{pmatrix} -0.0144 & +0.1026 \\ +0.1026 & -0.0146 \end{pmatrix} \mathrm{E_{Pl}} \quad \Rightarrow \quad \lambda_{\rm sh} = -0.117, \quad \lambda_{\rm br}^{(2)} = +0.088\ \mathrm{E_{Pl}}. \tag{109}$$

Both eigenvalues of the reduced 2×2 matrix are identified: the *antisymmetric* shape mode (one bond shortens, another lengthens) has $\lambda_{\rm sh} = -0.117 < 0$ — the equilateral is a **saddle point** at its own classical minimum.

Table 40: Shape-mode Hessian eigenvalues vs d at $C_{\rm Pl}$, $m_{\rm sp} = 0.1$. Negative eigenvalue indicates local instability in that direction.

d [$l_{\rm Pl}$]	λ_1	λ_2	Classification
1.00	-0.160	$+0.278$	Saddle
1.37 (class. min.)	-0.117	$+0.088$	Saddle
1.84	0	-0.030	Transition
2.00	-0.081	-0.016	Fully unstable
4.52 (quant. $\langle d \rangle$)	-0.064	-0.037	Fully unstable
8.00	-0.051	-0.022	Fully unstable

The stability transition (where the larger shape eigenvalue crosses zero) occurs at $d^* \approx 1.84\, l_{\rm Pl}$. For $d > d^*$, the equilateral is a *local maximum* in both shape directions — the potential energy decreases in all shape-deformation directions.

At the quantum mean position $\langle d \rangle = 4.52\, l_{\rm Pl}$: Both eigenvalues are negative ($\lambda_1 = -0.064$, $\lambda_2 = -0.037$), confirming the instability extends through the entire classically accessible region of the quantum wavefunction.

25.4 Physical interpretation

The shape instability has a clear physical origin: at $C = C_{\rm Pl}$, the pair interaction $V_2 \propto e^{-m_{\rm sp} d}/d$ is non-negligible. An isosceles deformation creating one shorter bond $d_{\rm short} < d$ and one longer bond $d_{\rm long} > d$ with $d_{\rm short} + d_{\rm long} = 2d$ lowers V_2 because Yukawa is convex: e^{-mx}/x is a convex function for $mx \lesssim 1$, so the average over asymmetric bonds gives lower energy than at the symmetric point.

At $C = 0.282$ all pairs are classically bound ($E_0^{(2B)} < 0$, see ex29), so pair interactions drive isosceles deformation.

25.5 Consequence for quantum Efimov window

Classical level. The true classical ground state is a deformed (isosceles) triangle, not equilateral. Along the softer mode ($\varepsilon_a = -\varepsilon_b$), the energy decreases monotonically with deformation at fixed d:

$$E(d, \varepsilon) < E(d, 0) \quad \forall\, \varepsilon \neq 0. \tag{110}$$

The equilateral result $E_{\rm cl}^{\rm eq} = -0.265\, E_{\rm Pl}$ (at $d_{\rm cl} = 1.37$) is an *upper bound* on the true classical minimum.

Quantum level. In the quantum problem, the full shape space is the 2D hyperangle manifold. The equilateral breathing-mode calculation (ex26/ex29) restricts to the 1D submanifold $d_{12} = d_{13} = d_{23}$. Since the equilateral is a local energy maximum in the shape directions, the restriction gives an *upper bound* on E_0:

$$E_0^{\rm equil} \geq E_0^{\rm true}. \tag{111}$$

The true ground-state energy is lower (more bound).

Implication for window boundaries. A lower true E_0 means both thresholds $C_Q^{(3B)}$ and $C_Q^{(2B)}$ are *smaller* than computed in ex26/ex27:

- $C_Q^{(3B)}$ (lower window bound): shifts downward. Binding occurs at weaker coupling.
- $C_Q^{(2B)}$ (upper window bound): shifts downward. Two-body binding also occurs at weaker coupling.
- The window $m_{\rm sp} \in (0.076, 0.198)$ computed in ex27 is a *conservative estimate*. The true quantum window is *at least as wide* and may extend to slightly larger $m_{\rm sp}$ (requiring less coupling to bind the triple via the 3-body force).

Three-body only vs two-body. Whether the 3B-only Efimov window still exists for the full 2D problem depends on whether $C_Q^{(3B)}$ shifts faster than $C_Q^{(2B)}$ as the geometry deforms. Since V_3 (the three-body force) is also symmetric and may have different shape dependence than V_2, a full 2D calculation is required.

25.6 Normal mode frequencies

The radial (breathing) mode at $d_{\rm cl} = 1.37\, l_{\rm Pl}$ has curvature:

$$\left.\frac{d^2E}{dd^2}\right|_{\rm eq} = 0.305\ {\rm E_{Pl}}/l_{\rm Pl}^2. \tag{112}$$

With effective mass $\mu_r = m_{\rm Pl}/3$, the breathing frequency is $\omega_r = \sqrt{0.305 \times 3} \approx 0.957\, E_{\rm Pl}$.

The stiffer shape mode has $\lambda_2^{(2)}/d_{\rm cl}^2 = +0.047$, giving $\omega_{{\rm sh},2} = \sqrt{0.047 \times 3} \approx 0.375\, E_{\rm Pl}$ (about $0.39 \times \omega_r$). The softer shape mode has $\lambda_1^{(2)}/d_{\rm cl}^2 < 0$ (imaginary frequency): this is the unstable mode.

25.7 Shape stability vs C at quantum mean position

Table 41: Shape stability at $d = \langle d \rangle = 4.52\, l_{\rm Pl}$ vs coupling C. At $C < 0.19$ (near quantum Efimov threshold) the larger eigenvalue is still positive — saddle-type stability.

C	λ_1	λ_2	Type	Note
0.100	−0.0045	+0.0177	Saddle	below 3B threshold
0.128	−0.0074	+0.0123	Saddle	classical 3B threshold
0.150	−0.0102	+0.0066	Saddle	Efimov window midpoint
0.191	−0.0168	−0.0081	Fully unstable	quantum 3B threshold
0.282	−0.0637	−0.0374	Fully unstable	$C_{\rm Pl}$

The shape mode transitions from "saddle" to "fully unstable" near $C \approx 0.19$, coinciding with the quantum Efimov threshold $C_Q^{(3B)} = 0.191$. Below threshold, the stiff shape mode has a positive restoring force, partially stabilising the equilateral geometry. Above threshold, both modes are unstable.

25.8 Summary

1. The equilateral triangle is a **saddle point** in shape space for all C and d tested — never a stable minimum.
2. The 1D equilateral breathing-mode calculation (ex26/ex29) gives **upper bounds** on E_0 and on the window thresholds.

3. The true quantum ground state is likely a deformed (isosceles) triangle. By S_3 symmetry, the quantum state is a superposition of three equivalent isosceles configurations, restoring permutation symmetry.

4. The quantum Efimov window for $C_{\rm Pl}$ may be slightly wider than $(0.076,\, 0.198)\, l_{\rm Pl}^{-1}$ (more conservative: could start at lower $m_{\rm sp}$ or extend to higher $m_{\rm sp}$).

5. A full 2D hyperangle Schrödinger solver is required for quantitatively accurate window boundaries. This is identified as the primary open computational problem (ex31+).

26 Open theoretical questions

This section discusses three fundamental open questions in TGP Path B that require theoretical rather than numerical resolution.

26.1 Q1: Yukawa source mechanism

Question. Why does a body of mass m act as a source of the Yukawa scalar field ϕ with coupling $C = m/(2\sqrt{\pi}\, m_{\rm Pl})$? In TGP Path B, this is taken as an axiom. No derivation from first principles has been given.

Scalar-field formulation (standard . In a standard massive scalar field theory with Lagrangian $\mathcal{L} = -\frac{1}{2}(\partial_\mu\phi)^2 - \frac{1}{2}m_{\rm sp}^2\phi^2 - g\phi T$, the coupling to the trace of the energy-momentum tensor $T = -\rho$ (for non-relativistic matter) gives the field equation $(\Box - m_{\rm sp}^2)\phi = g\rho$, with the Yukawa solution $\phi = -\frac{g}{4\pi}\frac{me^{-m_{\rm sp}r}}{r}$ for a point mass m.

The two-body potential from scalar exchange is then $V_{12} = g^2\phi_1(r_{12})m_2 = -\frac{g^2 m_1 m_2}{4\pi}\frac{e^{-m_{\rm sp}r}}{r}$. Matching to TGP: $g^2/(4\pi) = C_1C_2/(m_1m_2) = 1/(4\pi m_{\rm Pl}^2)$, giving $g = 1/m_{\rm Pl}$ — this is the Planck-scale coupling.

Three-body vertex from ϕ^3 term. The three-body TGP interaction arises if the scalar Lagrangian contains a cubic self-interaction $\mathcal{L}_3 = -\lambda\phi^3/6$. The one-loop diagram with one ϕ^3 vertex and three external matter lines generates the Feynman integral I_Y computed in the preceding sections. The coupling scales as $C^3 = (gm/m_{\rm Pl})^3 = (m/m_{\rm Pl})^3/(4\pi)^{3/2}$.

Identification with Dilaton or Brans-Dicke scalar. A natural candidate is the dilaton field ϕ of string theory or the scalar partner in Brans-Dicke gravity. In Brans-Dicke theory, the scalar χ couples to matter as $g_{\rm BD}\chi T$, with $g_{\rm BD} \sim 1/m_{\rm Pl}$. The Yukawa mass $m_{\rm sp}$ would arise from a non-minimal scalar potential $V(\chi) \ni \frac{1}{2}m_{\rm sp}^2\chi^2$.

Status. The coupling structure of TGP Path B is consistent with a standard dilaton-like scalar with coupling $g = 1/m_{\rm Pl}$ and mass $m_{\rm sp}$. The derivation from a specific string or modified-gravity Lagrangian is an open problem. The key observation is that the TGP 2-body potential IS the Newtonian gravitational potential at $r \ll 1/m_{\rm sp}$, so the scalar-exchange mechanism is a completion of Newtonian gravity at sub-Planck scales.

26.2 Q2: Dark matter and rotation curves

Question. Can TGP with $m_{\rm sp} \sim 1/\text{kpc}$ explain galactic rotation curves without dark matter?

Answer: No, for ordinary matter. The TGP two-body coupling for ordinary baryonic matter is $C = m/(2\sqrt{\pi}\, m_{\rm Pl}) \sim 10^{-19}$–$10^{-20}$ (for masses $m \sim 10^{-27}$–10^{-26} kg). The two-body TGP potential is:

$$V_{\rm TGP}(r) = -\frac{G m_1 m_2}{r}\, e^{-m_{\rm sp} r}. \tag{113}$$

At $r \ll 1/m_{\rm sp}$ (sub-Yukawa range) this is Newtonian gravity. At $r \gg 1/m_{\rm sp}$ the potential is *exponentially suppressed*, giving *less* gravity than Newton. Flat rotation curves require *more* gravity at large r (not less), so Yukawa screening cannot explain dark-matter rotation curves for any $m_{\rm sp}$.

The three-body correction scales as $C^3 \sim 10^{-60}$ and is completely negligible for galactic dynamics.

TGP dark matter scenario. A TGP dark matter candidate would require particles with $C \sim 1$ (near-Planck-mass objects, $m \sim m_{\rm Pl}$) in sufficient density to contribute to galactic rotation. Such particles would:

- Interact gravitationally (via the same Yukawa) with baryons with $C \sim 10^{-19}$, giving effectively no observable coupling at galactic scales.
- Be gravitationally self-interacting with $C^2 \sim 1$, potentially forming bound clusters (the Efimov-type states of this work).
- Have no electroweak or strong interactions, consistent with collisionless dark matter at galaxy scales.

MOND connection. Modified Newtonian dynamics (MOND, Milgrom 1983) postulates a transition at acceleration $a_0 \approx 1.2 \times 10^{-10}\,{\rm m/s^2}$. TGP provides no derivation of a_0. The TGP three-body force gives a correction $\delta a/a \sim C \sim 10^{-19}$ for baryonic matter, far below $a_0/g_{\rm Newt} \sim 10^{-4}$ at galactic edges. TGP and MOND are presently independent frameworks; a connection would require a theoretical bridge identifying a_0 with some combination of C and $m_{\rm sp}$.

Numerical comparison (ex31). A rotation curve computation (see ex31) confirms that Yukawa-screened gravity (TGP) gives:

- Identical to Newtonian for $r \ll 1/m_{\rm sp}$ (as expected).
- Below Newtonian for $r \gtrsim 1/m_{\rm sp}$ — opposite to observation.
- CDM (NFW halo) and MOND both produce flat curves and fit observations; TGP alone does not.

26.3 Q3: Path C — quantum defects and the Yukawa derivation

Question. TGP Path C proposes that the Yukawa coupling of Path B can be *derived* from the quantum theory of topological defects of the scalar field ϕ. What is the mathematical structure of this derivation?

Topological defect review (ex10-ex11). The TGP scalar field satisfies a nonlinear equation with defect solutions (ex10-ex11). The linearised tail of the defect is not Yukawa but oscillatory:

$$\phi(r) \sim \frac{A\sin(r)}{r} + \frac{B\cos(r)}{r} \quad (r \to \infty). \tag{114}$$

This oscillatory tail arises from the Helmholtz-type linearisation $(\nabla^2 + 1)\delta = 0$ near $\phi = 1$. A Yukawa tail ($e^{-m_{\rm sp} r}/r$) would require a *massive* linearised equation $(\nabla^2 - m_{\rm sp}^2)\delta = 0$, which is not present in the current TGP defect ODE.

Sketch of Path C. Path C would require modifying the scalar Lagrangian (or adding a mass term) so that the defect tail becomes Yukawa. One approach:

1. Add a symmetry-breaking potential $V(\phi) = \frac{1}{2}m_{\rm sp}^2(\phi - 1)^2$ to the TGP Lagrangian.
2. The linearised equation becomes $(\nabla^2 - m_{\rm sp}^2)\delta = 0$ near $\phi = 1$, giving Yukawa decay.
3. The defect core acts as a Yukawa source with coupling C determined by the defect profile — this would *derive* the TGP Path B axiom.

This sketch requires: (a) A consistent modified TGP Lagrangian with both defect and Yukawa solutions, (b) A derivation of $C(m_{\rm sp})$ from the defect quantisation, (c) Consistency with the oscillatory solutions found in ex11 (which would emerge in the limit $m_{\rm sp} \to 0$).

Status. Path C is mathematically open. The key obstacle identified in ex11 is that the *standard* TGP defect ODE produces oscillatory (not Yukawa) tails. A modified Lagrangian is required. The construction of a consistent TGP Lagrangian connecting defect solutions to Yukawa couplings is the primary open theoretical problem.

26.4 Q4: Relation to general relativity

Question. TGP Path B is a scalar field theory; GR is a tensor theory. How do they coexist, and what are the observational signatures distinguishing them?

Coexistence. In the standard picture, two independent fields can mediate gravity:

- **Tensor (graviton):** mediates spin-2 gravity, responsible for GR effects (light deflection, perihelion precession, gravitational waves).
- **Scalar (ϕ, TGP):** mediates additional spin-0 attraction, responsible for the Yukawa correction $e^{-m_{\rm sp}r}$.

The post-Newtonian parameter framework (PPN) accommodates both:

$$\Phi(r) = -\frac{Gm}{r}\left(1 + \alpha_{\rm TGP}\, e^{-m_{\rm sp}r}\right), \tag{115}$$

where $\alpha_{\rm TGP} = C^2/(Gm/r) \sim 1$ is the ratio of TGP to Newtonian coupling. For TGP Path B with $C = m/(2\sqrt{\pi}m_{\rm Pl})$, one has $\alpha_{\rm TGP} \sim 1$ at $r \ll 1/m_{\rm sp}$, so TGP contributes $\sim 100\%$ of Newtonian gravity at short range.

Observational distinction from GR. Key signatures of the scalar sector:

1. **Fifth-force violation of equivalence principle:** Scalar coupling $g = 1/m_{\rm Pl}$ is universal (proportional to mass), so equivalence principle is preserved and TGP is equivalent to Newtonian for $r \ll 1/m_{\rm sp}$.
2. **Yukawa deviation from $1/r^2$ force:** At $r \gtrsim 1/m_{\rm sp}$, TGP predicts sub-Newtonian gravity. Laboratory and satellite tests of the inverse-square law constrain $m_{\rm sp} \gtrsim 10^{-3}\,{\rm m}^{-1}$ ($\lambda_{\rm sp} \lesssim 10^3\,{\rm m}$). Planck-scale $m_{\rm sp} \sim 1/l_{\rm Pl}$ is far above this bound.
3. **Scalar gravitational waves:** The TGP scalar ϕ can emit monopole/dipole radiation in addition to GR's quadrupole. The scalar wave energy is $P_{\rm scalar} \sim G^2m^2\ddot{m}_{\rm sp}^2/c^5$ — negligible for non-relativistic sources.
4. **Brans-Dicke parameter:** If TGP is identified with Brans-Dicke theory with parameter $\omega_{\rm BD}$, solar system tests require $\omega_{\rm BD} > 40{,}000$. This constrains the scalar coupling; TGP with $g = 1/m_{\rm Pl}$ and $m_{\rm sp} \gg H_0$ (screening at sub-solar-system scales) is consistent with $\omega_{\rm BD} \to \infty$ (no long-range scalar modification).

Summary. TGP Path B is consistent with GR and all solar-system tests if $m_{\rm sp} \gtrsim 1/(1\,\mathrm{AU}) \approx 7\times10^{-12}\,\mathrm{m}^{-1}$. For Planck-scale $m_{\rm sp} \sim l_{\rm Pl}^{-1} = 6\times10^{34}\,\mathrm{m}^{-1}$, the Yukawa range is $\lambda_{\rm sp} \sim 10^{-35}$ m — far below any observable scale except the Planck-mass N-body physics studied in this work. TGP and GR coexist without contradiction at all observationally accessible scales.

26.5 Summary of open questions

Table 42: Status of open theoretical questions in TGP Path B.

Q	Question	Status	Key obstacle
Q1	Yukawa source derivation	Open	No Lagrangian for TGP
Q2	Dark matter/rotation curves	Answered (negative)	TGP reduces gravity at large r; 3B negligible
Q3	Path C: defect → Yukawa	Open	Oscillatory defect tail; need modified Lagrangian
Q4	TGP vs GR	Addressed	Coexist; TGP is sub-Planck modification of gravity
Q5	Full 2D quantum Efimov	Open	Equilateral approx. is upper bound; need 2D FD solver

27 Galactic rotation curves: TGP vs CDM and MOND

27.1 Setup

We compute rotation curves for a Milky-Way-like galaxy under four scenarios: baryons only (Newtonian), TGP with Yukawa screening, CDM with NFW halo, and MOND.

Galaxy model. Exponential stellar disk ($M_{\rm disk} = 5\times10^{10}\,M_\odot$, $R_d = 3.5\,\mathrm{kpc}$), Hernquist bulge ($M_{\rm bulge} = 10^{10}\,M_\odot$, $r_b = 0.7\,\mathrm{kpc}$), gas disk ($M_{\rm gas} = 5\times10^{9}\,M_\odot$, $R_{\rm gas} = 7\,\mathrm{kpc}$). NFW halo: $M_{200} = 10^{12}\,M_\odot$, $c_{\rm NFW} = 15$, $r_s = 13.6\,\mathrm{kpc}$. MOND acceleration constant $a_0 = 1.2\times10^{-10}\,\mathrm{m\,s^{-2}}$.

TGP rotation curve. For TGP Path B, the 2-body coupling is $C = m/(2\sqrt{\pi}m_{\rm Pl})$, giving $C^2 = Gm_1m_2$ — identical to Newtonian gravity. The radial acceleration from TGP is

$$a_{\rm TGP}(r) = \frac{GM(r)}{r^2}\,e^{-m_{\rm sp}r}(1+m_{\rm sp}r), \tag{116}$$

where the Yukawa screening factor $f(r) = e^{-m_{\rm sp}r}(1+m_{\rm sp}r)$ satisfies $f(0) = 1$ and $f(r) \to 0$ exponentially for $r \gg 1/m_{\rm sp}$.

27.2 Results

27.3 Analysis

Why TGP cannot explain flat rotation curves. The Yukawa potential $e^{-m_{\rm sp}r}/r$ satisfies

$$\frac{e^{-m_{\rm sp}r}}{r} < \frac{1}{r} \quad \forall\, m_{\rm sp} > 0,\ r > 0. \tag{117}$$

Therefore TGP acceleration is *always less than or equal to* Newtonian. Flat rotation curves ($v =$ const) require $a(r) \propto 1/r$ — more gravity at large r than the baryonic Newtonian prediction. TGP provides systematically less gravity, moving in the *wrong direction.*

Table 43: Circular velocity $v_{\rm circ}$ [km/s] for each model. TGP with any $m_{\rm sp} > 0$ gives *less* gravity than Newton at large r — opposite to what flat rotation curves require.

r	Baryons (Newton)	TGP $\lambda = 1$ [kpc]	TGP $\lambda = 5$ [kpc]	TGP $\lambda = 15$ [kpc]	CDM +NFW	MOND kpc>kpc
1	138	118	137	138	157	139
5	170	34	145	166	222	182
10	165	4	105	153	239	195
15	145	0	65	124	235	194
20	126	0	38	99	228	190
30	100	0	13	64	217	185

Table 44: Yukawa screening factor $f(r) = e^{-m_{\rm sp} r}(1 + m_{\rm sp} r)$. At $r = 20\,$kpc, even $\lambda = 15\,$kpc gives $f = 0.62$ — a 38% reduction in gravity relative to Newton.

r [kpc]	$\lambda = 1\,$kpc	$\lambda = 5\,$kpc	$\lambda = 15\,$kpc
1	0.736	0.983	0.998
5	0.040	0.736	0.955
10	0.0005	0.406	0.856
20	≈ 0	0.092	0.615
30	≈ 0	0.017	0.406

CDM (NFW halo) works. The NFW halo provides an additional $\sim 100\,$km/s at $r = 20\,$kpc, sustaining a flat curve of $v \approx 220\,$km/s consistent with the observed Milky Way rotation velocity.

MOND works. In the deep-MOND limit ($a \ll a_0$), Milgrom's relation gives $a^2/a_0 = a_N \Rightarrow v^4 = GMa_0$ (Tully-Fisher relation). For $M_{\rm bar} = 6.5 \times 10^{10}\,M_\odot$, the MOND prediction is $v_{\rm TF} = (GM_{\rm bar}a_0)^{1/4} = 179\,$km/s, consistent with the computed curve $v(30\,{\rm kpc}) = 185\,$km/s.

Three-body TGP correction. For ordinary baryonic matter ($C \sim 10^{-19}$), the three-body coupling is $C^3 \sim 10^{-57}$, completely negligible for galactic dynamics. Even summing over $N \sim 10^{11}$ stellar triples within a 1 kpc sphere gives a relative correction $\delta a_{\rm 3B}/a \sim NC^3 \sim 10^{-46}$.

27.4 Conclusion

TGP Path B (for baryonic matter with $C \sim 10^{-19}$) is indistinguishable from screened Newtonian gravity. It:

1. Reproduces Newtonian gravity exactly at $r \ll \lambda_{\rm sp}$.
2. Predicts *less* gravity than Newton at $r \gtrsim \lambda_{\rm sp}$.
3. Cannot produce flat rotation curves for any $m_{\rm sp} > 0$.
4. Has negligible ($\sim 10^{-57}$) three-body corrections for baryons.

TGP cannot substitute for dark matter or MOND in explaining rotation curves. A TGP dark-matter candidate would require near-Planck-mass particles ($C \sim 1$, $m \sim m_{\rm Pl}$), but their interaction with baryons (coupling $\propto C_{\rm baryon} \cdot C_{\rm DM} \sim 10^{-19}$) would be observationally indistinguishable from zero.

The Yukawa screening is a generic property: any Lagrangian with a *positive* scalar mass $m_{\rm sp}^2 > 0$ yields a screened potential. Only a tachyonic scalar ($m_{\rm sp}^2 < 0$) or a massless scalar ($m_{\rm sp} = 0$) could produce long-range enhancement — but these lie outside TGP Path B.

28 2D isosceles quantum Schrödinger equation

28.1 Motivation

Section 25 (ex30) showed that the equilateral triangle is a **saddle point** in shape space: the antisymmetric deformation mode has a negative Hessian eigenvalue $\lambda_{\rm sh} = -0.117\, E_{\rm Pl}$. This means the 1D equilateral breathing-mode calculation of ex23–ex29 restricts to a saddle of the potential, giving *upper bounds* on E_0. The question is: by how much does the true ground-state energy differ?

To answer this, we implement the full 2D quantum Schrödinger equation on the *isosceles submanifold* $d_{12} = d_{13} = a$, $d_{23} = b$ (the natural complement to the equilateral line), and compare with the 1D result.

28.2 Classical isosceles minimum (ex32)

Before solving the quantum problem, we locate the classical energy minimum on the isosceles submanifold using the effective potential $E_{\rm eff}(a, b) = {\rm ZP}(\bar{d}) + V_2(a, a, b) + V_3(a, a, b)$ where $\bar{d} = (2a + b)/3$.

Table 45: Classical isosceles minimum vs equilateral minimum at $C = C_{\rm Pl}$, $m_{\rm sp} = 0.1\, l_{\rm Pl}^{-1}$. The isosceles minimum lies at $b \to 0$ (Yukawa singularity), confirming that the effective-potential landscape has no classical lower bound off the equilateral line.

Configuration	a	b	b/a	E $[E_{\rm Pl}]$
Equilateral min.	1.371	1.371	1.000	-0.265
Isosceles (grid)	2.293	0.300	0.131	-0.446
True limit $b \to 0$	any	0	0	$-\infty$

Physical interpretation. The Yukawa pair potential $V_2 = -C^2 e^{-m_{\rm sp} b}/b \to -\infty$ as $b \to 0$. The effective ZP term $3/(8\bar{d}^2)$ is finite as $b \to 0$ (with a fixed), so the effective potential is *unbounded below* in the isosceles direction.

This is a classical collapse singularity: the 3-body Yukawa potential has no hard core. The equilateral "minimum" found in ex23–ex29 is a local minimum only within the equilateral subspace $d_{12} = d_{13} = d_{23}$.

Quantum mechanics resolves the collapse: the kinetic energy operator $\sim -\nabla^2$ diverges as $b \to 0$ faster than the Yukawa attraction $\sim -1/b$, providing a quantum-mechanical hard core. The correct treatment therefore requires the full quantum kinetic energy, not the semi-classical ZP approximation.

28.3 2D Jacobi coordinates (ex33)

Coordinate system. For 3 particles of mass $m = m_{\rm Pl}$ in 1D, the Jacobi coordinates are:

$$\boldsymbol{\xi}_1 = \mathbf{r}_2 - \mathbf{r}_3, \quad |\boldsymbol{\xi}_1| = b = d_{23}, \tag{118}$$

$$\boldsymbol{\xi}_2 = \mathbf{r}_1 - \tfrac{1}{2}(\mathbf{r}_2 + \mathbf{r}_3), \quad |\boldsymbol{\xi}_2| = h = \sqrt{a^2 - b^2/4}, \tag{119}$$

with reduced masses $\mu_1 = m/2 = 0.5$ and $\mu_2 = 2m/3 \approx 0.667$.

Hamiltonian. The quantum Hamiltonian in (b, h) coordinates (no ZP term — kinetic energy is exact):

$$\hat{H} = -\frac{1}{2\mu_1}\frac{\partial^2}{\partial b^2} - \frac{1}{2\mu_2}\frac{\partial^2}{\partial h^2} + V(b, h) = -\frac{\partial^2}{\partial b^2} - \frac{3}{4}\frac{\partial^2}{\partial h^2} + V(b, h), \tag{120}$$

where the isosceles potential is

$$V(b,h) = V_2(a,a,b) + V_3(a,a,b)\,, \quad a = \sqrt{h^2 + b^2/4}. \tag{121}$$

No separate ZP term appears: the full quantum kinetic energy provides the natural repulsion against collapse.

Kinetic energy for the equilateral breathing mode. Along the equilateral line $b = d$, $h = d\sqrt{3}/2$, the effective kinetic energy coefficient for the breathing coordinate d is

$$T_{\rm br} = -\left[\frac{1}{2\mu_1}\left(\frac{db}{dd}\right)^2 + \frac{1}{2\mu_2}\left(\frac{dh}{dd}\right)^2\right]\frac{d^2}{dd^2} = -\left[1 + \frac{3}{4}\cdot\frac{3}{4}\right]\frac{d^2}{dd^2} = -\frac{25}{16}\frac{d^2}{dd^2}, \tag{122}$$

corresponding to effective mass $\mu_{\rm eff} = 8/25 = 0.32\,m_{\rm Pl}$. The 1D equilateral calculation (ex23–ex29) used $\mu = 1\,m_{\rm Pl}$, which is 3.1× too large — it *underestimates* the kinetic energy, overestimating binding.

28.4 Numerical results

Grid. We use a uniform (b,h) grid with $N_b \times N_h = 50 \times 50$ points, $b, h \in [0.15,\, 20]\,l_{\rm Pl}$, and Dirichlet boundary conditions ($\psi = 0$ on all edges). The sparse Hamiltonian (2500×2500, diagonal + nearest-neighbour off-diagonals) is diagonalised with `scipy.sparse.linalg.eigsh`.

Table 46: Ground-state energy comparison: 1D equilateral vs 2D Jacobi isosceles. At $C = C_{\rm Pl}$, $m_{\rm sp} = 0.1$.

Method	E_0 $[E_{\rm Pl}]$	$\langle b\rangle$	$\langle a\rangle$	b/a (peak)
1D equilateral (ex23–ex29)	-0.0698	—	4.52	1.000
2D Jacobi isosceles (ex33)	-0.0088	6.88	7.53	0.956
Difference ΔE	$+0.0610$ (87% of $\lvert E_0^{\rm 1D}\rvert$)			

Table 47: Ground-state energy vs $m_{\rm sp}$ from 2D Jacobi calculation ($N = 40 \times 40$, $C = C_{\rm Pl}$). For $m_{\rm sp} \geq 0.130$, the 2D calculation gives an unbound state ($E_0 > 0$).

$m_{\rm sp}$ $[l_{\rm Pl}^{-1}]$	λ $[l_{\rm Pl}]$	$E_0^{\rm 1D}$	$E_0^{\rm 2D}$	ΔE $[E_{\rm Pl}]$
0.076	13.2	-0.117	-0.049	$+0.068$
0.100	10.0	-0.070	-0.013	$+0.057$
0.130	7.7	-0.034	$+0.010$	$+0.044$
0.150	6.7	-0.018	$+0.019$	$+0.037$
0.180	5.6	-0.005	$+0.026$	$+0.031$
0.198	5.1	≈ 0	$+0.029$	$+0.029$

28.5 Physical interpretation

Wavefunction geometry. At $m_{\rm sp} = 0.1$, $C = C_{\rm Pl}$, the ground-state wavefunction peaks at $(b,h) = (5.01,\, 4.61)\,l_{\rm Pl}$, corresponding to $a = \sqrt{4.61^2 + 5.01^2/4} = 5.24\,l_{\rm Pl}$ and $b/a = 5.01/5.24 = 0.956$. The equilateral value is $b/a = 1$, so the peak is only 4.4% off the equilateral line. The quantum ground state is effectively equilateral in shape.

Kinetic energy discrepancy. The 2D Jacobi calculation gives $E_0^{\rm 2D} = -0.0088$ vs $E_0^{\rm 1D} = -0.0698$. The discrepancy ($\Delta E = +0.061$, 87% of $\lvert E_0^{\rm 1D}\rvert$) arises primarily from the effective mass:

- 1D equilateral (ex23): effective mass $\mu_{\rm eff} = 1\,m_{\rm Pl}$ (kinetic coefficient 1/2), ZP $= 3/(8d^2)$.

- 2D Jacobi (ex33): kinetic coefficients 1 (b-direction) and 3/4 (h-direction), no ZP term.
- Along equilateral breathing mode: the 2D kinetic coefficient is $25/16 = 1.5625$, versus $1/2$ in the 1D model — a factor $3.1\times$ larger.

The larger kinetic energy in the 2D Jacobi model provides stronger quantum pressure, reducing binding by a factor ~ 3.

Efimov window in 2D Jacobi. The 2D Jacobi calculation predicts the 3-body bound state threshold at $m_{\rm sp} \approx 0.12\, l_{\rm Pl}^{-1}$ (below which $E_0 < 0$, above which $E_0 > 0$), compared with $m_{\rm sp} = 0.198$ in the 1D equilateral model. This shift reflects the stronger kinetic energy in the correct Jacobi treatment.

28.6 Conclusion and open problems

The 2D Jacobi Schrödinger equation for the isosceles submanifold gives a less-bound ground state than the 1D equilateral approximation, due to the correct Jacobi reduced masses. Key conclusions:

1. The ground-state wavefunction is near-equilateral in shape ($b/a \approx 0.96$), confirming the equilateral as the preferred geometry.
2. The equilateral approximation of ex23–ex29 *overestimates* binding because it uses an effective mass $\mu = 1$ instead of the correct $\mu_{\rm eff} = 0.32$ for the equilateral breathing mode.
3. The corrected (2D Jacobi) ground-state energy is $E_0^{\rm 2D} = -0.0088\, E_{\rm Pl}$ at $C_{\rm Pl}$, $m_{\rm sp} = 0.1$.
4. The Efimov window upper boundary shifts from $m_{\rm sp} = 0.198$ (1D) to $m_{\rm sp} \approx 0.12$ (2D Jacobi).
5. A full 3D calculation including all angular degrees of freedom (not just the isosceles submanifold) is required for the physically correct Efimov spectrum. This is the primary open computational problem (Q5).

Status of Q5 (full 2D quantum Efimov). The 2D Jacobi calculation of ex33 gives a first quantitative estimate of the true (beyond equilateral-approximation) quantum ground state. The result: $E_0^{\rm true} \approx -0.009\, E_{\rm Pl}$ at $C_{\rm Pl}$, $m_{\rm sp} = 0.1$, and the Efimov window narrows to $m_{\rm sp} \lesssim 0.12\, l_{\rm Pl}^{-1}$. The next step is a full 3D Faddeev or hyperspherical calculation.