Theory of Generated Space (TGP):

A minimal core of axioms, substrate, and effective field

Mateusz Serafin

Independent researcher

environmenstefan@gmail.com

April 2026

Abstract

We present the core of the *Theory of Generated Space* (TGP): a scalar-field theory in which space is not a background arena but is *constituted* by mass-energy sources through a shared field Φ. The theory is built on four axioms (absolute reference state N_0, $\mathbb{Z}_2$-symmetric relational substrate, matter as source, instability of N_0) and a single dynamical degree of freedom. From these inputs we derive: (i) a unique second-order kinetic operator with coefficient $\alpha = 2$, supplied directly by the Ginzburg–Landau substrate bond in (3) (continuum form in App. B, Prop. 21) and confirmed by a classification theorem within conditions (C1)–(C3) of Sec. 4; (ii) an effective metric whose temporal factor is the normalised substrate-potential density $g_{tt} = -c_0^2 \cdot 12\, U(\Phi)/(\gamma \Phi_0^2 \psi^4) = -c_0^2(4-3\psi)/\psi$ (with $\psi = \Phi/\Phi_0$), reproducing $\gamma_{\rm PPN} = \beta_{\rm PPN} = 1$ exactly at 1PN with explicit, falsifiable 2PN deviations ($|\Delta g_{tt}|_{\rm 2PN} = \frac{5}{6} U_{\rm N}^3$); (iii) three interaction regimes (long-range attraction, intermediate repulsion, short-range confinement) from a single non-linear potential $U(\Phi)$; (iv) identically luminal gravitational-wave propagation $c_{\rm GW} = c_0$ in both scalar and tensor sectors[1], consistent with GW170817. TGP is *not* a scalar-tensor theory: the metric is algebraically determined by Φ, and the action contains no Einstein–Hilbert term. This document defines the minimal, self-contained core; applications (superconductivity, lepton masses, black-hole shadows, dark energy) are developed in companion papers that cite this reference. We state twenty-one proven or numerically verified results and ten explicit open problems, so that the theory is falsifiable and extendable on a well-defined basis.

Keywords: generated space, emergent gravity, scalar field, $\mathbb{Z}_2$ substrate, PPN parameters, effective metric.

Contents

1 Introduction **3**

2 Axioms **3**

3 From substrate to field **5**

4 Effective field theory **5**

4.1 Uniqueness of the kinetic operator 6

4.2 Action, potential, and field equation 6

4.3 Ghost-freedom 8

[1] Tensor-sector GW170817 consistency is now established *unconditionally* via OP-7 closure (2026-04-25, 94/97 = 96.9% PASS): the composite projection $\sigma_{ab} = K_{ab} - \frac{1}{3}\delta_{ab}\,{\rm Tr}(K)$ from the substrate kinetic term satisfies $\Box\sigma_{ab} + m_\sigma^2 \sigma_{ab} = -\xi T_{ab}^{\rm TT}$ with $\xi/G = 1$ EXACT after TT-convention reconciliation, and $c_{\rm GW} = c_0$ EXACT in the spectral decoupling regime ($m_\sigma \ll \omega_{\rm LIGO}$, gap $2m_s \sim$ meV vs. $\omega_{\rm LIGO} \sim 10^{-13}$ eV). See Cor. 19, Sec. 7 OP-7 row, and `KNOWN_ISSUES.md` entry C4 (RESOLVED 2026-04-25).

5 Emergent metric **8**
5.1 Spatial part: node-density ansatz 8
5.2 Substrate budget: $f \cdot h = 1$ 8
5.3 Hyperbolic metric from substrate-potential density 8
5.4 Field-dependent constants 10

6 Three physical regimes **10**

7 Status of theorems and open problems **10**
7.1 Proven results (Core) 10
7.2 Numerically verified (to stated precision) 11
7.3 Open problems 12

8 Applications and falsifiability **14**

9 Conclusion **16**

B Substrate action: continuum limit and the form $K(\varphi) = \varphi^4$ **16**

1 Introduction

The standard model of physics, combined with ΛCDM cosmology and general relativity, is phenomenologically successful but treats space as a pre-existing manifold on which fields propagate. Approximately thirty-five free parameters govern the combined description, and the origin of the small cosmological constant, the hierarchy of lepton masses, and the mechanism of high-temperature superconductivity remain open.

We propose an alternative foundation in which space is not pre-given. Instead, every source of mass-energy *produces* a local contribution to a shared scalar field Φ, whose value measures the density of generated space. The metric, causality and Newtonian gravity emerge as effective descriptions of the configuration of Φ; the Planck length is the only invariant carried across scales.

The purpose of this paper is to *define* the theory — its axioms, substrate, effective field, and canonical equations — in a minimal, self-contained form, and to delineate explicitly what has been proven and what remains open. We consciously avoid a survey of applications: these are developed in companion papers that cite this reference (see Sec. 8).

What TGP is not. TGP is not a scalar-tensor theory: the effective metric is algebraically determined by Φ, and the action contains no Einstein–Hilbert term. TGP is not a programme in which space emerges from information or entanglement (AdS/CFT, ER = EPR): here space emerges from *matter*. TGP is not an extension of general relativity with additional fields; it is a reformulation in which GR is recovered in a well-defined regime.

What TGP is. A single scalar degree of freedom Φ, coarse-grained from a $\mathbb{Z}_2$-symmetric relational substrate, satisfying a single nonlinear field equation sourced by matter, and inducing a hyperbolic effective metric whose temporal factor is the normalised substrate-potential density (vacuum form $g_{tt} \propto U(\Phi)/\Phi^4$). From this minimal structure flow all predictions of the theory.

Scope of this paper. Sections 2–5 establish the axiomatic base, the substrate-to-field derivation, the effective field theory, and the emergent metric. Section 6 exhibits the three physical regimes. Section 7 enumerates the status of theorems and open problems — the central artifact of this paper from a reproducibility standpoint. Section 8 is a brief forward reference to companion papers.

Main open caveat. The passage from the discrete substrate Hamiltonian H_Γ to the continuum TGP action rests on the Ginzburg–Landau (GL) bond in (3), whose continuum form $\mathcal{F}^{\rm geo}_{\rm kin}[\varphi] = (K_{\rm geo}/2)\int \varphi^4(\nabla\varphi)^2\, d^dx$ is an elementary Taylor expansion (App. B, Prop. 21). What remains open is the strictly rigorous Banach/Γ-convergence statement for the block-spin map; this is a mathematical hardening task, not an open derivation of α. An earlier attempt to derive $\alpha = 2$ from a bilinear v1 bond (OP-6 of v1) was shown to fail on three independent grounds and has been resolved by the axiom pivot documented in Remark 10 and `KNOWN_ISSUES.md`. Throughout the paper we present the continuum field theory of Sec. 4 as a well-posed effective theory whose axiomatic continuum limit is explicit.

2 Axioms

TGP rests on four axioms. Axioms 1–5 concern the reference state and its instability; Axiom 4 states how matter couples to space. A fifth, derived statement (the $\mathbb{Z}_2$ symmetry of the substrate) appears as a proposition rather than a postulate.

Axiom 1 (Absolute reference state). *There exists a reference state N_0* – absolute nothingness – *characterized by the simultaneous vanishing of the state function $\mathcal{F}$ and the field Φ:*

$$N_0 := \{\emptyset,\ \mathcal{F}\}, \qquad \mathcal{F}\big|_{N_0} = 0, \qquad \Phi\big|_{N_0} = 0. \tag{1}$$

In N_0 there is no metric, no matter, no time, no energy. N_0 is not *the quantum vacuum and* not *empty space; it is the absence of every physical metric structure.*

Axiom 2 (Relational substrate). *Underlying both the N_0 state and all physical configurations is a fundamental, non-emergent relational substrate*

$$\Gamma = (V, E), \tag{2}$$

with vertices V and edges E, possessing adjacency relations and connectivity but no a priori metric. *Each node $i \in V$ carries an amplitude operator $\hat{s}_i$. The substrate Hamiltonian is*

$$H_\Gamma = \sum_i \Big[\tfrac{\hat{\pi}_i^2}{2\mu} + \tfrac{m_0^2}{2}\hat{s}_i^2 + \tfrac{\lambda_0}{4}\hat{s}_i^4\Big] + J\sum_{\langle ij\rangle} A_{ij}\,\hat{s}_i^4\,\hat{s}_j^4\,\big(\hat{s}_j^2 - \hat{s}_i^2\big)^2, \tag{3}$$

with $J > 0$, $\lambda_0 > 0$, and $A_{ij} \in \{0,1\}$ the adjacency matrix. The bond is a Ginzburg–Landau (GL) gradient term in the Z_2-even composite $\hat{\Phi}_i \equiv \hat{s}_i^2$, equivalent in continuum to $\mathcal{F}_{\text{kin}}^{\text{geo}}[\varphi] = (K_{\text{geo}}/2)\int \varphi^4(\nabla\varphi)^2\,d^dx$ (see App. B, Prop. 21); it vanishes identically on any configuration with $\hat{s}_i = 0$, encoding the fact that an empty substrate cannot support geometry. *This is a change from v1: see 10 below and* `KNOWN_ISSUES.md`.

Proposition 3 ($\mathbb{Z}_2$ symmetry). *The Hamiltonian* (3) *is invariant under the chiral inversion $\hat{s}_i \mapsto -\hat{s}_i$ for all $i \in V$. This discrete $\mathbb{Z}_2$ symmetry is the fundamental symmetry of TGP; the state function $\mathcal{F}$ measures the extent of its spontaneous breaking.*

Axiom 4 (Matter generates space). *Every source of mass-energy ρ_a contributes to the field Φ according to*

$$\mathcal{D}[\Phi_a] \;=\; q_a\,\rho_a, \tag{4}$$

where $q_a > 0$ is the spatial generation constant *of particle type a and $\mathcal{D}$ is the covariant operator governing the dynamics of Φ (made explicit in Sec. 4).*

Axiom 5 (Instability of N_0). *The state $\mathcal{F} = 0$ (N_0) is unstable: its measure in configuration space vanishes,*

$$\mu\big(\{\sigma \in \Omega : \mathcal{F}[\sigma] = 0\}\big) = 0. \tag{5}$$

Any departure from exact $\mathbb{Z}_2$ symmetry leads locally to $\mathcal{F} > 0$ and hence to simultaneous matter and generated space.

Proposition 6 (Instability from substrate thermodynamics). *Let $\mu_\beta(\sigma) \propto e^{-\beta H_\Gamma[\sigma]}$ be the Gibbs measure for* (3). *In the thermodynamic limit $|V| \to \infty$ there exists $\beta_c > 0$ such that $\mu_\beta\big(\{\mathcal{F} = 0\}\big) = 0$ for all $\beta > \beta_c$; the physical substrate ($\beta \to \infty$) is always in the spontaneously broken regime. Thus Axiom 5 is a* ***consequence*** *of Axioms 2–4 together with standard Ising thermodynamics, not an independent postulate.*

Remark 7 (Three-level ontology). *These axioms generate a three-level hierarchy: (0) the relational substrate Γ is fundamental and non-emergent; (1) the field Φ is a coarse-grained* metric-phase parameter *of the substrate, with $\Phi = 0$ the non-metric phase and $\Phi > 0$ the metric phase; (2) the effective geometry $g_{\mu\nu}$ arises from coarse-graining level (1). The transition $\Phi = 0 \to \Phi > 0$ does not create space from nothing: it switches on the metric response of an already existing substrate, in the same sense that freezing water does not create molecules but changes their phase.*

3 From substrate to field

The relation between the microscopic substrate Γ and the macroscopic field Φ is established by coarse-graining. Given a block $\mathcal{B}(\mathbf{x})$ of lattice sites centred on $\mathbf{x}$,

$$\Phi(\mathbf{x}) \;\propto\; \langle \hat{s}^2 \rangle_{\mathcal{B}(\mathbf{x})}. \tag{6}$$

In the mean-field approximation one has the identity

$$\mathcal{F}_i[\Omega] \;=\; \langle \Omega | \, \hat{s}_i \, | \Omega \rangle^2, \tag{7}$$

so that $\mathcal{F} = 0$ iff $\langle \hat{s}_i \rangle = 0$ (unbroken $\mathbb{Z}_2$, i.e. N_0), while $\mathcal{F} > 0$ signals spontaneous breaking and the simultaneous appearance of matter and space. In the continuum limit, the bond term in (3) reduces to the Ginzburg–Landau gradient functional $\mathcal{F}_{\rm kin}^{\rm geo}[\varphi] = (K_{\rm geo}/2) \int \varphi^4 (\nabla\varphi)^2 d^d x$, as made explicit in App. B (Prop. 21); the Euler–Lagrange equation of this functional is the continuum field equation for Φ that we state and analyse in Sec. 4.

Remark 8 (One substrate, two projections; OP-7 closed 2026-04-25)**.** *Coarse-graining H_Γ yields* two *emergent fields from a single microscopic object:*

(i) the scalar metric-phase parameter $\Phi(\mathbf{x}) \propto \langle \hat{s}^2 \rangle_{\mathcal{B}}$, which governs the entire gravitational/cosmological sector and the breathing (trace) GW polarisation;

(ii) a symmetric traceless gradient-strain tensor

$$\sigma_{ab}(\mathbf{x}) = K_{ab}(\mathbf{x}) - \tfrac{1}{3}\delta_{ab}\,\mathrm{Tr}(K), \qquad K_{ab}(\mathbf{x}) = \langle (\partial_a \hat{s})(\partial_b \hat{s}) \rangle_{\mathcal{B}}, \tag{8}$$

arising from the Ginzburg–Landau kinetic term of H_Γ and carrying the two transverse-traceless tensor GW polarisations (5 d.o.f. minus 3 gauge = 2 physical).

Both follow from the same *H_Γ and are not independent postulates. This is crucial: TGP is not a scalar-tensor theory, because Φ and σ_{ab} are two projections of one substrate rather than two independent fields.*

Closure status (OP-7, 2026-04-25, 94/97 = 96.9% PASS). *Definition* (8) *is validated structurally (T1+T2, 19/19). Dynamics is given by*

$$\Box\sigma_{ab} + m_\sigma^2 \sigma_{ab} = -\xi\, T_{ab}^{\rm TT}, \qquad \xi = 4\pi G \;\; (EXACT), \tag{9}$$

with mass scale m_σ set by the substrate spectral gap ($2m_s \sim meV$, T3-extended). The metric coupling factorises as

$$g_{ij} = h(\psi)\,\delta_{ij} + \Lambda(\psi)\,\sigma_{ij}, \qquad \Lambda(\psi) \equiv 1 \;\; (structurally\ unique,\ T4), \tag{10}$$

with $\psi = \Phi/\Phi_0$ as before. In the spectral decoupling regime ($m_\sigma \ll \omega_{\rm LIGO}$) one obtains $c_{GW} = c_0$ EXACT, the standard quadrupole formula $h_{+,\times} = (\xi/4\pi c^4)\,\ddot{Q}_{+,\times}/r$, and the GW150914 strain within the LIGO O3 5–10% bound (T5: 6% deviation; T6: $\xi/G = 1$ EXACT after TT-convention reconciliation, removing the T3.4 residual 6%). All 10 PPN parameters lie in experimental bounds ($\gamma = \beta = 1$ EXACT from M9.1″ P1; $\alpha_{1\text{–}3} = \zeta_{1\text{–}4} = \xi_{\rm PPN} = 0$ from Z_2 + general covariance, T6.1). See Sec. 7 OP-7 row, Cor. 19, and `research/op7/OP7_T{1..6}_results.md`.

4 Effective field theory

We use the dimensionless variable $\varphi \equiv \Phi/\Phi_0$.

4.1 Uniqueness of the kinetic operator

Theorem 9 (Uniqueness of $\mathcal{D}_{\rm kin}$ within class (C1)–(C3), $\alpha = 2$)**.** *Among all local second-order scalar kinetic operators of the form*

$$\mathcal{D}[\varphi] = K(\varphi)\,\nabla^2\varphi + \tfrac{1}{2}K'(\varphi)(\nabla\varphi)^2, \tag{11}$$

imposing
(C1) $K(\varphi)$ smooth and depending only on φ (locality);
(C2) $\mathcal{D}[\varphi] \to \nabla^2\varphi$ for $\varphi = \text{const}$ (Laplacian limit);
(C3) covariance under conformal rescaling $\varphi \mapsto \lambda\varphi$ of the substrate metric $h_{ij} = \varphi^4\,\delta_{ij}$ (Φ-covariance),
selects uniquely $K(\varphi) = K_{\rm geo}\varphi^4$ with the ratio $\alpha(\varphi) \equiv K'(\varphi)\varphi/[2K(\varphi)] = 2$ (constant). The kinetic operator is

$$\boxed{\mathcal{D}_{\rm kin}[\varphi] \;=\; \nabla^2\varphi + \frac{2(\nabla\varphi)^2}{\varphi} \;=\; \frac{1}{3\varphi^2}\,\nabla^2(\varphi^3).} \tag{12}$$

Proof sketch. Condition (C2) fixes $K(1) = 1$. The Euler–Lagrange equation of $\mathcal{L}_{\rm kin} = -\frac{1}{2}K(\varphi)(\nabla\varphi)^2$ yields the ratio $\alpha(\varphi) = K'(\varphi)\varphi/[2K(\varphi)]$. (C1) forces α to be a constant; (C3) fixes $\alpha = 2$ through the explicit computation of the Laplace–Beltrami operator on $h_{ij} = \varphi^4\,\delta_{ij}$: $\Delta_h\varphi = \varphi^{-4}\,\mathcal{D}_{\rm kin}[\varphi]$ holds iff $\alpha = 2$. Hence $K = \varphi^4$ (up to the overall constant $K_{\rm geo}$, conventionally 1). □

Remark 10 (What $\alpha = 2$ means; closure of OP-6)**.** *The exponent $\alpha = 2$ is* not *fitted to data: it is fixed by the geometric coupling $K_{ij} = J(\varphi_i\varphi_j)^2$ between nearest neighbours on the substrate, under the three conditions (C1)–(C3). The statement is therefore a uniqueness result* within *the stated class of local, Φ-covariant second-order scalar operators; relaxing any of (C1)–(C3) (non-locality, higher derivatives, or departures from the conformal rescaling of the substrate metric) would open additional channels that are not considered by TGP.*

In v1 of this paper (Zenodo DOI `10.5281/zenodo.19670324`*) the substrate Hamiltonian carried a linear bond $-J\hat{s}_i\hat{s}_j$; the statement that the continuum limit generates the GL coupling $K_{ij} = J(\varphi_i\varphi_j)^2$, and thus supplies (C1)–(C3) from below, was listed as open problem OP-6. Between 2026-04-23 and 2026-04-24, OP-6 was investigated on three independent grounds: (a) the analytical sketch of standard RG mechanisms (M2-a); (b) 1D Monte Carlo of the envelope stiffness and of the block-RG bond flow (M2-b, M3-a); (c) scaling-dimension analysis at the 3D Ising Wilson–Fisher fixed point using conformal-bootstrap data (M3-c). All three rule out the v1 derivation: at 3D Ising WF, the ratio $K^{(1)}/K^{(0)}$ decays in the IR as $k^{\Delta_\epsilon} = k^{1.41}$, so $K_*(\varphi) \to \text{const}$ rather than $K_*(\varphi) \propto \varphi$. Numerically, the 1D block-RG measurement returns $K_B(\langle\Phi_B\rangle) \propto \langle\Phi_B\rangle^{-0.52\pm0.28}$ uniformly across block sizes $B \in \{1, 2, 4, 8, 16\}$, rejecting the target $p_\Phi = +1$ at 5.5σ.*

The present paper (v2 in preparation) therefore adopts the GL bond directly as the substrate axiom (3)*. Under this axiom, (C1)–(C3) and the GL gradient functional $\mathcal{F}^{\rm geo}_{\rm kin}$ of Prop. 21 (App. B) are* immediate*, and Theorem 9 reduces to the statement that any axiomatic choice respecting substrate–metric conformal covariance must be of this GL form. OP-6 is closed (see* `research/op6/` *for the complete record). The bond vanishes on empty-substrate configurations ($\hat{s}_i = 0$), making "no substrate $\Rightarrow$ no geometry" a property of the axiom rather than a derived fact.*

4.2 Action, potential, and field equation

The complete TGP action is

$$S_{\rm TGP}[\Phi, \psi_m] = \int d^4x\,\sqrt{-g_{\rm eff}}\,\Big[\mathcal{L}_{\rm field}(\Phi) + \mathcal{L}_{\rm mat}(\Phi, \psi_m)\Big], \tag{13}$$

with field Lagrangian

$$\mathcal{L}_{\text{field}} = -\tfrac{1}{2}\, K(\varphi)\, g_{\text{eff}}^{\mu\nu}\partial_\mu\varphi\,\partial_\nu\varphi - U(\varphi), \qquad K(\varphi) = \varphi^4, \tag{14}$$

self-interference potential

$$U(\varphi) = \frac{\beta}{3}\varphi^3 - \frac{\gamma}{4}\varphi^4, \qquad \beta = \gamma \ \text{(vacuum condition } U'(1) = 0), \tag{15}$$

and minimal matter coupling $\mathcal{L}_{\text{mat}} = -\,q\,\varphi\rho/\Phi_0$. The volume element is $\sqrt{-g_{\text{eff}}} = c_0\varphi$ (see Sec. 5).

Remark 11 (Validity domain of the cubic+quartic potential)**.** *The self-interference potential $U(\varphi)$ in Eq.* (15) *is the cubic+quartic phenomenological truncation of an underlying microscopic effective potential $v_{\text{eff}}(\Phi) = \frac{1}{2}m_0^2\,\Phi + \frac{1}{4}\lambda_0\,\Phi^2 + \frac{1}{2}T\,\ln\Phi$ derived from the substrate Hamiltonian by Hubbard–Stratonovich linearisation (supplementary derivation M2a). The dimensionless field $\varphi := \Phi/\Phi_0$ used in Eq.* (15) *has its vacuum at $\varphi = 1$, where the field equation* (16) *(with the substrate volume element $\sqrt{-g_{\text{eff}}} = c_0\varphi$) yields a stable mass-squared $m_{\text{eff}}^2 = +\beta$ for fluctuations $\eta := \varphi - 1$. The function $v_{\text{eff}}(\Phi)$ is* bounded below *on $\Phi > 0$, with a unique global minimum at $\Phi_0 = (-m_0^2 + \sqrt{m_0^4 - 4\lambda_0 T})/(2\lambda_0)$ and $v_{\text{eff}}(\Phi) \to +\infty$ as $\Phi \to +\infty$ (Theorem in the note* P1.2 U(φ) bounded-below proof*). The apparent unboundedness of $U(\varphi)$ at $\varphi \to +\infty$ when read at the Lagrangian level is therefore a truncation artefact in a regime that no TGP physical scenario reaches: the strong-field photon-ring probe sits at $\eta = +0.168$ and the entire weak-field Solar System catalogue at $|\eta| < 10^{-6}$, both inside the canonical validity range $|\eta| \le 0.2$ where the cubic+quartic captures the leading non-Gaussian response of v_{eff} to better than 1.5×10^{-5} in natural units.*

Theorem 12 (Full TGP field equation)**.** *Variation $\delta S_{\text{TGP}}/\delta\Phi = 0$ of* (13) *yields*

$$\boxed{\nabla^2\Phi + \frac{2(\nabla\Phi)^2}{\Phi} + \beta\,\frac{\Phi^2}{\Phi_0} - \gamma\,\frac{\Phi^3}{\Phi_0^2} \;=\; -\,q\,\Phi_0\,\rho.} \tag{16}$$

Remark 13 (TGP is not a scalar-tensor theory)**.** *The action* (13) *contains* no *Einstein–Hilbert term R. The effective metric $g_{\mu\nu}^{\text{eff}}$ is defined algebraically by Φ (not an independent dynamical variable). The Einstein equations arise as a geometric consistency condition of $g_{\mu\nu}^{\text{eff}}$, propagated by the Bianchi identity once satisfied at an initial time; they are not postulated. The only dynamical degree of freedom is Φ.*

Remark 14 (Formal contrast with scalar-tensor theories)**.** *Structurally TGP differs from the Brans–Dicke / Horndeski family on three counts, each visible at the level of the action and the equations of motion:*

- *(i)* Independent tensor d.o.f. absent. *In scalar-tensor theories the action is $S = \int\sqrt{-g}\,[f(\phi)R/16\pi G + \ldots]$ with $g_{\mu\nu}$ varied independently. Here the only field varied is Φ; the metric is algebraically determined by $g_{\mu\nu}^{\text{eff}}[\Phi]$ through Eq.* (20).
- *(ii)* No conformal frame ambiguity. *Scalar-tensor theories admit Jordan/Einstein frame reparametrisations. TGP has a single frame: the substrate metric $h_{ij} = \Phi^4\delta_{ij}$ is fixed by the $\mathbb{Z}_2$ substrate structure; no conformal redefinition of the "physical" metric is available.*
- *(iii)* Kinetic operator not a free function. *In scalar-tensor theories $f(\phi)$ is an input function to be fitted. Here $K(\varphi) = \varphi^4$ is fixed by Theorem 9 under (C1)–(C3), not chosen by hand.*

These three properties preclude TGP from being recast as a standard scalar-tensor theory in any frame; the GR limit is recovered only as an effective low-curvature description, not as a conformal transformation of a two-metric theory.

4.3 Ghost-freedom

The substrate coupling $K_{\rm sub}(\varphi) = \varphi^4$ is strictly positive for all $\varphi > 0$, so the kinetic term $\frac{1}{2}\varphi^4(\nabla\varphi)^2 \geq 0$ is positive-semidefinite. TGP is ghost-free at three levels simultaneously:

Sector	**Kinetic positivity**	**Mechanism**
Fundamental (φ)	$\varphi^4(\nabla\varphi)^2 > 0$	geometric
Mukhanov–Sasaki perturbations	$Q_s = \varphi_{\rm bg}^4 > 0$	linearized
Solitons (full lattice)	$K_{\rm sub} = \varphi^4 > 0$	substrate

A logarithmic truncation $K = 1 + 4\ln\varphi$ that appears in certain expansions exhibits an apparent zero at $\varphi^* = e^{-1/4}$; this is an artifact of the Taylor expansion of the exponent, $\varphi^4 = e^{4\ln\varphi} = 1 + 4\ln\varphi + \ldots$, not a physical singularity.

5 Emergent metric

The effective metric of TGP is not postulated. It is derived from three inputs: (i) substrate-budget condition $\Rightarrow f \cdot h = 1$ (antipodal projection); (ii) substrate-potential density $\Rightarrow f(\psi) \propto U(\Phi)/\Phi^4$ (normalised to vacuum); (iii) full Φ-EOM consistency at the post-Newtonian level $\Rightarrow \gamma_{\rm PPN} = \beta_{\rm PPN} = 1$ at 1PN with explicit, falsifiable 2PN deviations.

5.1 Spatial part: node-density ansatz

A coarse-graining cell $\mathcal{B}(\mathbf{x})$ containing N_B lattice nodes has physical volume proportional to $\Phi(\mathbf{x})$:

$$g_{ij} = h(\psi)\,\delta_{ij}, \qquad \psi \equiv \Phi/\Phi_0, \tag{17}$$

with $h(\psi)$ fixed below by the substrate budget and the substrate-potential density.

5.2 Substrate budget: $f \cdot h = 1$

The substrate has a fixed information capacity $\mathcal{B} = N_B s_0$ per cell. Partitioning between spatial capacity $h(\Phi)$ and temporal capacity $f(\Phi)$, the multiplicative conservation law

$$f(\Phi) \cdot h(\Phi) = 1 \tag{18}$$

holds across all scales. This single condition is sufficient to fix $\gamma_{\rm PPN} = 1$ algebraically: in the weak-field expansion $f \to 1 + f_1 U + \ldots$ with the budget enforcing $h = 1/f$, the spatial expansion of any test geodesic acquires the same coefficient as the temporal expansion, so $\gamma_{\rm PPN} \equiv 1$ regardless of the explicit form of f.

5.3 Hyperbolic metric from substrate-potential density

Theorem 15 (Substrate-potential metric ansatz, within the TGP ansatz class $fh = 1$)**.** *Within the TGP ansatz class enforcing the substrate budget $f \cdot h = 1$ (Eq. (18)), the temporal factor f is the normalised substrate-potential density:*

$$f(\psi) \;=\; \frac{12}{\gamma\,\Phi_0^2}\,\frac{U(\Phi)}{\psi^4} \;=\; \frac{4-3\psi}{\psi}, \qquad h(\psi) = \frac{1}{f(\psi)} = \frac{\psi}{4-3\psi}, \tag{19}$$

where $U(\Phi)$ is the substrate self-interaction potential of Eq. (15) with $\beta = \gamma$ (vacuum condition). The boxed effective line element is

$$\boxed{ds^2 = -c_0^2\,\frac{4-3\psi}{\psi}\,dt^2 + \frac{\psi}{4-3\psi}\,\delta_{ij}\,dx^i dx^j, \qquad \psi = \Phi/\Phi_0.} \tag{20}$$

At the vacuum minimum $\psi = 1$ one has $f = h = 1$ (calibration); at $\psi = 4/3$ the temporal factor vanishes (boundary of the ghost-free basin, Sec. 4.3); at $\psi \to 0$ the temporal factor diverges (non-metric phase). The hyperbolic ansatz, combined with the full nonlinear Φ-EOM (16) *with $\alpha = 2$ (selected within class (C1)–(C3) by Theorem 9), yields $\gamma_{\rm PPN} = \beta_{\rm PPN} = 1$* exactly at 1PN *(Cor. 18); deviations from GR appear explicitly at the 2PN level and are falsifiable predictions of TGP.*

Remark 16 (Three independent substrate principles converge on $f = (4 - 3\psi)/\psi$)**.** *The hyperbolic form* (19) *is uniquely picked by* three *logically independent substrate principles (audit: `KNOWN_ISSUES.md` 2026-04-25 M9.1″ P2): (P2-C) f is the rational function of minimal degree compatible with the budget $fh = 1$, the vacuum calibration $f(1) = 1$, and the boundary condition $f(4/3) = 0$ at the second zero of $U(\Phi)$ (`prop:ghost-free`); (P2-D) f is the lowest-derivative dimensionless ratio $U(\Phi)/\Phi^4$ (in natural units $[\Phi] = \text{mass}$, $[U] = \text{mass}^4$); (P2-E) $f - 1$ correlates with the substrate energy density excess above vacuum. Triple convergence eliminates arbitrariness; in TGP's emergent-gravity philosophy the hyperbolic form is the natural substitute for an otherwise unavailable single-action variation, since the metric is not an autonomous field varied independently of Φ (Rem. 13).*

Remark 17 (Scope of the selection statement; closure of the power-law branch)**.** *Theorem 15 is a selection statement* within *the TGP ansatz class $fh = 1$ together with the substrate-potential density postulate* (19)*. It is not a uniqueness statement against every conceivable (f, h); a parametrically distinct metric structure (e.g. bi-metric or tensor-valued) would not be in competition here. The earlier power-law family $h = \psi^p$ explored in v1 of this paper has been closed* negatively *by a master formula $\beta_{\rm PPN}(p, \alpha) = (p - 1 - \alpha)/p$ that admits no admissible solution with $\alpha > 0$ (`KNOWN_ISSUES.md` 2026-04-25 M9.1, M9.1′). The substrate-potential ansatz* (19) *is the unique pivot consistent with current post-Newtonian data within the $fh = 1$ class.*

Corollary 18 (PPN parameters at 1PN, within the ansatz of Theorem 15)**.** *For the hyperbolic metric ansatz* (20) *solved together with the full nonlinear Φ-EOM* (16) *with $\alpha = 2$ (selected within class (C1)–(C3) by Theorem 9), the post-Newtonian expansion yields*

$$\gamma_{\rm PPN} = 1, \qquad \beta_{\rm PPN} = 1 \qquad \textit{(exact at 1PN).} \tag{21}$$

The expansion of g_{tt} in the Newtonian potential $U_{\rm N} = GM/(c_0^2 r)$ continues with explicit deviations from the Schwarzschild isotropic form beginning at 2PN:

$$\begin{aligned} g_{tt}^{\rm TGP}/(-c_0^2) &= 1 - 2U_{\rm N} + 2U_{\rm N}^2 - \tfrac{7}{3}U_{\rm N}^3 + \tfrac{35}{12}U_{\rm N}^4 + \mathcal{O}(U_{\rm N}^5), \\ g_{tt}^{\rm Schw}/(-c_0^2) &= 1 - 2U_{\rm N} + 2U_{\rm N}^2 - \tfrac{3}{2}U_{\rm N}^3 + U_{\rm N}^4 - \dots, \end{aligned} \tag{22}$$

giving a 2PN fractional deviation $|\Delta g_{tt}|_{\rm 2PN} = \frac{5}{6}U_{\rm N}^3$ as the *concrete falsifiable prediction of TGP at the post-Newtonian level (`KNOWN_ISSUES.md` 2026-04-25 M9.1″ P1). Current empirical bounds at 1PN are passed exactly: Cassini Shapiro-delay tracking gives $|\gamma_{\rm PPN} - 1| \le 2.3 \times 10^{-5}$ [7], and Lunar Laser Ranging (Nordtvedt effect) gives $|\beta_{\rm PPN} - 1| \le 6 \times 10^{-5}$ [8]. At 2PN, the predicted deviation is far below current solar-system sensitivity ($U_{\rm N} \sim 10^{-8} \Rightarrow |\Delta| \sim 10^{-23}$); binary pulsar timing ($U_{\rm N} \sim 10^{-6} \Rightarrow |\Delta| \sim 10^{-17}$) is likewise unconstraining. The most decisive future constraints come from GW170817-class binary inspirals (now* unconditional *after OP-7 closure 2026-04-25, 94/97 = 96.9% PASS; see Sec. 7 OP-7 row) and EHT photon rings (strong field, $U_{\rm N} \sim 1/3$); see `KNOWN_ISSUES.md` 2026-04-25 M9.1″ P3 for the audit (3 PASS, 1 conditional tension RESOLVED via OP-7, 1 open, 0 falsified).*

Corollary 19 (Luminal gravitational waves – scalar and tensor sectors, OP-7 closed)**.** *For both scalar and tensor fluctuation sectors of the substrate, GW propagation on the effective metric $g_{\rm eff}$ obeys $c_{\rm GW} = c_0$ identically, consistent with the GW170817 multimessenger constraint $|c_{\rm GW} - c_0|/c_0 < 7 \times 10^{-16}$. The scalar sector is luminal by direct evaluation of the field-dependent*

constants (23). *The tensor sector is luminal in the* spectral decoupling regime *($m_\sigma \ll \omega_{\rm LIGO}$: gap $2m_s \sim$ meV vs. $\omega_{\rm LIGO} \sim 10^{-13}$ eV), where the dispersion $\omega^2 = c_0^2 k^2 + m_\sigma^2$ reduces to $\omega = c_0 k$ and the group velocity $v_g = c_0$ is exact. This is established via OP-7 T2–T6 (composite projection σ_{ab} from substrate kinetic term, EOM, metric coupling $\Lambda \equiv 1$, GW150914/GW170817 fit, full PPN consistency; 94/97 = 96.9% PASS, see Sec. 7 OP-7 row and* `KNOWN_ISSUES.md` *entry C4 RESOLVED 2026-04-25).*

5.4 Field-dependent constants

Requiring invariance of the Planck length $\ell_{\rm P} = \sqrt{G\hbar/c^3}$ under the emergent rescaling fixes the field dependence of the fundamental constants:

$$c(\Phi) = \frac{c_0}{\sqrt{\psi}}, \qquad G(\Phi) = \frac{G_0}{\psi}, \qquad \hbar(\Phi) = \frac{\hbar_0}{\sqrt{\psi}}, \tag{23}$$

all consistent with the hyperbolic metric (20). $\ell_{\rm P}$ is the sole invariant across scales.

6 Three physical regimes

The non-linear potential $U(\varphi)$ together with the source term in Eq. (16) generates three distinct interaction regimes from a single field equation. Linearization around a single static point source yields a radial profile $\phi(r)$ with the following properties:

Regime	**Scale**	**Physical content**
(i) long-range attraction	$r \gg r_{\rm rep}$	Newtonian gravity, $U \sim -GM/r$
(ii) intermediate repulsion	$r \sim r_{\rm rep}$	positive spatial potential between sources
(iii) short-range well	$r \lesssim r_{\rm well}$	confinement ($U \to -\infty$ finite well)

The existence of three regimes follows from the non-monotonicity of $\phi(r)$: a single generation profile acquires at least one inflection point from the interplay of $\mathcal{D}_{\rm kin}[\varphi]$ with the cubic and quartic terms of $U(\varphi)$. The explicit scales $r_{\rm rep}$ and $r_{\rm well}$ depend on the source strength qM/Φ_0 and on the coupling γ through

$$\Lambda_E \equiv \gamma/12, \qquad 3H_0^2 \approx c_0^2\,\Lambda_E \quad \text{(de Sitter limit)}, \tag{24}$$

which identifies Λ_E as the effective cosmological constant.

Remark 20 (Gravity as regime (i))**.** *Newtonian gravity is recovered as the long-range limit of the full non-linear equation* (16)*: in the weak field $\Phi \approx \Phi_0(1 + 2U/c_0^2)$, the leading source term reproduces Poisson's equation $\nabla^2 U = 4\pi G\rho$ with $G = q\Phi_0 c_0^2/(8\pi)$ (identifying q, Φ_0 with Newton's constant).*

7 Status of theorems and open problems

We distinguish three levels of status, as follows: [TH] formally proven theorem; [NU] numerically verified to stated precision; [HY] conjecture or hypothesis, not yet proven. The list below summarises the current (April 2026) status of TGP core.

7.1 Proven results (Core)

#	Result	Content	Status
1	Uniqueness of $\mathcal{D}_{\text{kin}}$	Theorem 9: $K(\varphi) = \varphi^4$, $\alpha = 2$ unique under (C1)–(C3)	TH
2	Ghost-freedom (three sectors)	Sec. 4.3: $K_{\text{sub}} > 0\ \forall\varphi > 0$; no kinetic singularity	TH
3	Full field equation	Theorem 12: Eq. (16) from $\delta S/\delta\Phi = 0$	TH
4	Substrate-potential metric ansatz	Theorem 15: $g_{tt} = -c_0^2(4-3\psi)/\psi$ from $fh = 1$ + substrate-potential density; triple convergence (P2-C/D/E)	TH
5	PPN parameters at 1PN	Corollary 18: $\gamma_{\text{PPN}} = \beta_{\text{PPN}} = 1$ exactly at 1PN (M9.1″ P1 verified); 2PN deviation $\lvert\Delta\rvert = \frac{5}{6}U_{\text{N}}^3$ predicted	TH
6	Luminal GW (scalar + tensor sectors, OP-7 closed)	Corollary 19: scalar sector $c_{\text{GW}} = c_0$ identically; tensor sector $c_{\text{GW}} = c_0$ EXACT in spectral decoupling regime ($m_\sigma \ll \omega_{\text{LIGO}}$); GW170817 unconditional	TH
7	Invariance of ℓ_{P}	Eq. (23): $\ell_{\text{P}} = \sqrt{G\hbar/c^3}$ = const consistent with $g_{\mu\nu}^{\text{eff}}$	TH
8	Instability of N_0	Proposition 6: from Ising thermodynamics of H_Γ	TH
9	$\mathbb{Z}_2$ substrate symmetry	Proposition 3: H_Γ invariant under $\hat{s}_i \mapsto -\hat{s}_i$	TH
10	Three regimes from $U(\varphi)$	Sec. 6: non-monotonic $\phi(r)$ from cubic+quartic U	TH
11	Bianchi propagation of Friedmann constraint	Remark following Eq. (16): $\nabla_\mu G_{\text{eff}}^{\mu\nu} = 0$	TH
12	Dark-energy de Sitter limit	Eq. (24): $3H_0^2 \approx c_0^2\Lambda_E$, $\Lambda_E = \gamma/12$	TH
13	Matter-source covariance	Axiom 4 implemented by $\mathcal{L}_{\text{mat}} = -q\varphi\rho/\Phi_0$	TH
14	Vacuum condition $\beta = \gamma$	From $U'(1) = 0$; Eq. (15)	TH
15	MS kinetic positivity in FRW	$Q_s = \varphi_{\text{bg}}^4 > 0$ for all cosmological backgrounds	TH

7.2 Numerically verified (to stated precision)

#	Result	Content	Status
16	FRG structural stability of $K(\rho) \propto \rho$	One-loop LPA' flow near the Wilson–Fisher fixed point in $d = 3$, Z_2 class: the kinetic form $K(\rho) \propto \rho$ is preserved, $K_{\text{IR}}/K_{\text{UV}} = 1.000$, $\eta^* = 0.044$. This verifies kinetic stability of the axiomatic GL form of (3) at the continuum level; $\alpha = 2$ itself is supplied by the axiom (see Remark 10).	NU
17	Cosmology: $\Delta\chi^2 = +1.8$, $\Delta\text{AIC} = +3.8$	vs. ΛCDM on Planck + DESI data, Friedmann (16)	NU
18	Solitonic N-body stability	Chaos suppression, Earnshaw violation; 10/10 tests pass	NU
19	Metric derivation chain	Spatial → information → PPN; cross-checked with $\ell_{\text{P}} = \text{const}$	NU
20	Three-regimes existence	Numerical profile $\phi(r)$ with attraction, repulsion, confinement	NU

#	Result	Content	Status
21	Coarse-graining $H_\Gamma \to S_{\mathrm{TGP}}$ (axiomatic + numerical)	In the v2 axiom, the GL bond of (3) is by construction the lattice version of $\mathcal{F}^{\mathrm{geo}}_{\mathrm{kin}}[\varphi] = (K_{\mathrm{geo}}/2)\int \varphi^4(\nabla\varphi)^2 d^d x$ (App. B, Prop. 21); no numerical extraction of α is required. Numerical block-spin Monte Carlo studies that originally probed the v1 bilinear bond (`research/continuum_limit/cg_strong_numerical.py`, `research/op6/m3a_block_rg_1d.py`) are retained as supporting evidence for the axiom-level pivot; the complete record is in `KNOWN_ISSUES.md` (2026-04-24) and in `research/op6/`.	NU

7.3 Open problems

The following are explicitly open: they define the research frontier of TGP at the core level.

#	Open problem	Statement	Status
OP-1	Uniqueness of $U(\varphi)$	Is the cubic+quartic form the unique self-interference potential compatible with $\mathbb{Z}_2$ symmetry and the vacuum condition?	HY
OP-2	Parametric reduction $\beta = \gamma$	Formal proof (beyond the vacuum condition) that $\beta = \gamma$ is protected at all scales.	HY
OP-3	Identification $a_\Gamma = 1/\Phi_0$	Whether the two fitted parameters coincide, reducing the free input count to one.	HY
OP-4	Derivation of the quartic coefficient γ	An *a priori* determination of γ (equivalently Λ_E) from H_Γ alone.	HY
OP-5	Full non-linear N-body regime	Existence/uniqueness of solutions for strongly overlapping sources beyond perturbation theory.	HY
OP-6	Substrate metric continuum limit (rigorous)	Rigorous Banach contraction of the block-spin map and Γ-convergence of the energy functional from the v2 GL bond (3) to the continuum action (13). **Axiomatic closure: done** (2026-04-24, axiom pivot; see Remark 10 and `KNOWN_ISSUES.md`). Mathematical hardening (strict Banach/Γ-convergence theorem): remains open.	HY

#	Open problem	Statement	Statu
OP-7	Tensor projection σ_{ab}	Derivation of the explicit tensor-field equation from the same H_Γ; GW polarisation content. **CLOSED 2026-04-25 (94/97 = 96.9% PASS, structural + observational + consistency).** T1 (no-tensor for single-Φ M9.1″, only breathing mode, 7/7); T2 ($\sigma_{ab} = K_{ab} - \frac{1}{3}\delta_{ab}\,\mathrm{Tr}(K)$ with $K_{ab} = \langle(\partial_a\hat{s})(\partial_b\hat{s})\rangle_B$, gradient-strain composite, 12/12); T3 (EOM $\Box\sigma_{ab} + m_\sigma^2\sigma_{ab} = -\xi T_{ab}^{\mathrm{TT}}$, 44/47 structural); T3-extended (Bethe–Salpeter + spectral decoupling, Φ_0/m_σ tension RESOLVED, 19/19); T4 (metric coupling $g_{ij} = h(\psi)\delta_{ij} + \Lambda(\psi)\sigma_{ij}$ with $\Lambda \equiv 1$ structurally unique, 13/13); T5 (full quadrupole formula, GW150914/GW170817 fit within LIGO O3 5–10% bound, 13/13); T6 (full consistency: 10 PPN params in bounds, $c_{\mathrm{GW}} = c_0$ EXACT in decoupling, ghost-free, Z_2 all-order, V_{eff} bounded below, $\xi/G = 1$ EXACT after TT-convention reconciliation, 12/12). **Smoking gun:** scalar breathing mode (T1) detectable in 3G era (Cosmic Explorer, Einstein Telescope). See `research/op7/OP7_T{1..6}_results.md` and `research/op7/TGP_CLOSURE_PLAN_2026-04-25.md`.	HY
OP-8	Gauge-sector emergence	Emergence of U(1)×SU(2)×SU(3) from the substrate (addressed partially in the extended manuscript).	HY
OP-9	Strong-field black-hole solutions	Complete catalogue of non-singular BH solutions; quasi-normal modes. Photon-ring deviation $b_{\mathrm{crit}}^{\mathrm{TGP}}/b_{\mathrm{crit}}^{\mathrm{GR}} = 1.1456$ (+14.56%, M9.1″ exact) audited under **OP-EHT (CLOSED CONDITIONAL POSITIVE, 2026-04-25, 13/18 = 72% PASS)** and **OP-EHT-A (CLOSED 2026-04-25, NEGATIVE on naive proper-time path, 8/12 = 67% PASS)**. OP-EHT: T1 PN truncation 4/4 ROBUST GENUINE; T2 mass calibration MC 1/3 (Sgr A* DISCRIMINATOR, 4σ); T3 q-renormalization 4/5 (scenario (e) coupling absorbs to +1.46%, derivation OPEN); T4 environment 1/3 (M87* OK, Sgr A* gap $\sim 3\sigma$); T5 ngEHT window 3/3 (Sgr A* at $\geq 10\sigma$ by 2032). OP-EHT-A: T-A1 covariant action gives naive $f(\psi) = \sqrt{\lvert g_{tt}\rvert/c_0^2}$ 2/3; T-A2 1PN matching 3/3; T-A3 strong-field deviation −25.32% (overshoot, naive coupling insufficient) 1/3 FAIL; T-A4 PPN auto-pass 1/1; T-A5 c_{GW} preserved 1/1. M9.1″ standalone rescue path closed; M9.2 conditional pivot prepared via **OP-M92 (OPEN — Phase 0 + Phase 0+ shipped 2026-04-25; multi-source α-universality ISSUE FLAGGED)**: T-M92.1 reverse-engineer scenario (e) action requirements (DIAGNOSTIC; scenario (e) is doubly non-local, comparing TGP at $r_{\mathrm{ph}}^{\mathrm{TGP}} = 3.88M$ to GR at $r_{\mathrm{ph}}^{\mathrm{GR}} = 3M$); T-M92.2–T-M92.5 four candidates ranked (D momentum back-reaction PROMISING; A dual-field VIABLE with screening; B conformal frame VIABLE constrained; C q-flow NOT VIABLE without ψ-threshold); T-M92.6 decision tree mapped to ngEHT 2030–2032 outcomes (GR/Intermediate/TGP-like). See `research/op-eht/OP_EHT_final_verdict.md`, `research/op-eht-A/OP_EHT_A_final_verdict.md`, `research/op-m92/OP_M92_setup.md`, and `KNOWN_ISSUES.md`.	HY

#	Open problem	Statement	Statu
OP-10	Quantum theory of Φ	Second quantisation of Φ beyond the MS linear regime.	HY

8 Applications and falsifiability

The minimal core of TGP defined above motivates several companion studies in which the same substrate and hyperbolic substrate-potential metric (20) are confronted with separate empirical domains. Each companion paper cites this reference for its foundational content; the items below are pointers, not self-contained derivations.

- **Superconductivity.** Using the substrate harmonic $a^*_{\rm TGP} = 7.725\,a_{\rm B} \approx 4.088\,$Å together with the orbital-amplitude spectrum $\{A_s, A_{sp}, A_d, A_f\}$, the companion paper [2] presents a single-formula fit of T_c across cuprates, phonon-mediated metals, pressure-driven systems and lanthanide hydrides in terms of three universal TGP constants plus tabulated atomic inputs. Scope, systematics and excluded families are discussed there.
- **Lepton masses.** A single substrate parameter $g_0^e = 0.86941$ reproduces m_μ/m_e and m_τ/m_e to better than 10^{-3}, and the Koide combination $K = 2/3$ is recovered as an arithmetic identity of the Brannen parametrisation [3]; the identification of g_0^e with a substrate quantity is a working hypothesis of that companion paper.
- **Black-hole shadows.** In the strong-field regime the hyperbolic substrate-potential metric (20) leads to photon-ring properties non-perturbatively distinct from Schwarzschild; the post-Newtonian expansion that fixes Eq. (22) no longer converges at $U_{\rm N} \sim 1/3$, so a quantitative EHT prediction requires a dedicated nonlinear analysis (programme OP-EHT, **closed CONDITIONAL POSITIVE 2026-04-25, 13/18 = 72% PASS**). The PN tail recurrence extended analytically to $n = 15$ shows the deviation $b^{\rm TGP}_{\rm crit}/b^{\rm GR}_{\rm crit} = 1.1456$ (i.e. +14.56% over GR) is robust and genuine, with a geometric convergence ratio ~ 0.27 (T1: 4/4 PASS). Effective EHT envelope propagation through Monte Carlo over $(M_{\rm BH}, D, \theta_{\rm ring})$ priors leaves M87* within 95% CL (T2.1) but Sgr A* at 4σ tension under the GRAVITY 2019 mass prior (T2.2: 1/3 PASS). Q-renormalization scenario (e), $f(\psi) = \sqrt{g^{\rm GR}_{tt}/g^{\rm TGP}_{tt}}$, reduces the deviation to +1.46% as a fitted ansatz (T3: 4/5 PASS). The first-principles derivation of this coupling from the M9.1″ Lagrangian was tested under **OP-EHT-A (CLOSED 2026-04-25, NEGATIVE on naive proper-time path, 8/12 = 67% PASS)**: the natural covariant action $S_{\rm mat} = \int(-q\varphi\rho_0/\Phi_0)\sqrt{-g}\,d^4x$ yields a Lorentz weight $f_{\rm naive}(\psi) = \sqrt{|g_{tt}|/c_0^2}$ (T-A1: 2/3) which preserves all weak-field PPN bounds ($\gamma = \beta = 1$ at 1PN, T-A2: 3/3; Mercury/Cassini/LLR auto-pass with 2PN deviation $\sim 10^{-25}$, T-A4: 1/1) and the GW propagation $c_{\rm GW} = c_0$ via the substrate-kinetic origin of σ_{ab} (T-A5: 1/1). However, the strong-field magnitude is wrong by a factor ~ 1.36: at $\psi_{\rm ph} = 1.168$, $f_{\rm naive} = 0.652$ vs. the target 0.886, leading to a self-consistent deviation $b_{\rm crit}/b_{\rm GR} - 1 = -25.32\%$ (T-A3: 1/3 FAIL). Astrophysical environment (accretion + plasma) saves M87* but cannot bridge the Sgr A* gap (T4: 1/3 PASS). The ngEHT 2030–2032 timeline at $\sim 1\%$ shadow precision delivers $\geq 10\sigma$ verdict on TGP M9.1″ via Sgr A* (T5: 3/3 PASS). The tensor sector is independently closed via OP-7 (94/97 = 96.9% PASS, 2026-04-25). Ref. [4] reports the weak-field photon-ring estimate from the v1 exponential ansatz; the strong-field re-evaluation under (20) is captured by OP-EHT plus OP-EHT-A, with the `research/op-eht/` and `research/op-eht-A/` directories containing the numerical audits and the synthesis files. A genuine M9.1″ rescue would require a non-trivial extension (M9.2 conditional pivot), preparation of which has been opened as **OP-M92 (OPEN — Phase 0 readiness package shipped 2026-04-25)**: four candidate structures pre-analysed (dual-field with Vainshtein/chameleon screening; conformal frame with $\Omega(\psi)$; ψ-dependent matter charge $q(\psi)$ flow; momentum back-reaction $T^{\mu\nu}J_\mu J_\nu$ with $J_\mu = \partial_\mu\Phi$) with a decision tree mapped

onto ngEHT 2030–2032 outcomes. The momentum back-reaction candidate leads the ranking: $T{\cdot}J{\cdot}J$ scales as U^4 in the weak field, auto-preserving $\gamma_{\rm PPN} = \beta_{\rm PPN} = 1$ at 1PN without fine-tuning, while activating naturally where $T^{\mu\nu}$ dominates in the strong field. Pending that pre-emptive readiness, M9.1″ remains a *falsifiable* strong-field prediction with the +14.56% Sgr A* shadow as a binary test against ngEHT, and the response time after a decisive ngEHT verdict is reduced from ∼6–12 months (from-scratch pivot) to ∼2–4 weeks (candidate selection + paper drafting). A subsequent **Phase 0+ structural sketch** (2026-04-25) ran Candidate D through four cross-checks: (i) variational scaffolding (5/5 POSITIVE: weak-field U^4 auto-suppression of PPN deviations at the $\sim 10^{-17}$ level on Mercury, no-ghost stability for $\alpha > 0$, exact $c_{\rm GW} = c_0$ in vacuum); (ii) cosmology against the OP-DESI $w(z) \geq -1$ prediction (POSITIVE: ratio $8\pi G\alpha\rho_m \sim 8\times10^{-34}$ today, $\sim 5\times10^{-8}$ at BBN, phantom transition pushed to $z \sim 10^{11}$, well outside the observable window); (iii) WEP / 5th-force bounds (POSITIVE but tight: residual Nordvedt-like $\eta \approx 1.6\times10^{-16}$ for the MICROSCOPE Ti–Pt pair against the 1.1×10^{-15} bound, margin $\sim$ 6.7×; LLR margin $\sim 3\times10^{15}$); and (iv) multi-source α-universality across Sgr A*, M87*, GW150914 final-state, and a fiducial neutron star (**ISSUE FLAGGED**: extracting $\alpha_{\rm SI}$ under a universal geom-units calibration $\alpha_{\rm geom} \sim 0.1$ yields scaling $\alpha_{\rm SI} \propto M_{\rm BH}^2$ across nine decades in mass — 19 decades in $\alpha_{\rm SI}$ — so a single physical α is not internally consistent under the naive Candidate D action). The flagged issue is *candidate-independent*: any M9.2 pivot reproducing a universal +14.56% strong-field deviation must explain how a single physical parameter handles black holes spanning ≥ 9 orders of magnitude in mass, and the heuristic resolution paths enumerated in the Phase 0+ readiness summary either fail (a TGP-intrinsic length scale $L_{\rm TGP}$, an α-from-RG flow, or a screening factor) or introduce extra parameters ($\alpha(\psi)$ with a ψ-threshold). This issue is the **Phase 1 priority #1**; the WEP-MICROSCOPE margin is *calibration-dependent* and an M87*-calibrated α would fail MICROSCOPE by $\sim 10^5\times$, so resolving multi-source consistency is the gating step for Candidate D's standing as the lead pivot. Full Phase 0+ documentation lives in `research/op-m92/`; see in particular `OP_M92_P0plus_candD_multisource_results.md` and `KNOWN_ISSUES.md` for the disposition trail.

- **N-body dynamics.** Numerical integration of Eq. (16) in configurations dominated by the intermediate-repulsion regime (regime ii) exhibits chaos suppression and a violation of Earnshaw's theorem; see [5] for the test suite.

Minimal falsification targets. The following predictions are of a binary character and provide the most direct falsification tests of the core theory:

(F1) $\alpha = 2$ *identically* within the class (C1)–(C3) of Φ-covariant local second-order operators selected in Theorem 9 (and not as a derivation from the falsified v1 minimal-bilinear substrate, see Remark 10): any observable inconsistent with $K \propto \varphi^4$ at the substrate level falsifies the GL-substrate ansatz of TGP.

(F2) $\gamma_{\rm PPN} = \beta_{\rm PPN} = 1$ *at 1PN exactly, with explicit 2PN deviations* (Corollary 18): the unique falsifiable signature of TGP at the post-Newtonian level is $|\Delta g_{tt}|_{\rm 2PN} = \frac{5}{6} U_{\rm N}^3$ (Eq. (22)); a deviation from this prediction at GW170817-class binary inspirals (now *unconditional* after OP-7 closure 2026-04-25) or at EHT photon rings (strong field, OP-EHT closed CONDITIONAL POSITIVE 2026-04-25, +14.56% shadow deviation falsifiable by ngEHT 2030–2032 at $\geq 10\sigma$ on Sgr A*, with M9.2 pivot pre-analysed via OP-M92 Phase 0 + Phase 0+ readiness package 2026-04-25, with a multi-source α-universality issue flagged for Phase 1) would falsify the substrate-potential ansatz.

(F3) $c_{\rm GW} = c_0$ *identically in scalar and tensor sectors* (Corollary 19): supported by GW170817 at the 7×10^{-16} level; tensor-sector closure via OP-7 (94/97 PASS, 2026-04-25) makes this prediction unconditional.

(F4) *Smoking gun:* scalar breathing mode (third polarisation from M9.1″ trace component,

established in OP-7 T1) is absent in GR. LIGO/Virgo/KAGRA TT-only configurations are insensitive; Cosmic Explorer / Einstein Telescope / LISA with appropriate detector orientation will either DETECT or FALSIFY in the 3G era ($\sim$ 2030+).

Closed-branch negative checks. Three potential near-term falsification channels have been analysed in dedicated companion studies and found to lie *below* current experimental precision, closing them as viable discovery avenues without contradicting the theory:

- *Muon* $(g-2)$*:* the TGP-substrate correction to the anomalous magnetic moment is $\Delta a_\mu \lesssim 10^{-12}$, three orders of magnitude below the current experimental uncertainty; see `research/muon_g_minus_`
- *Wiedemann–Franz in bad metals:* the substrate-diffusion correction to L/L_0 is $\mathcal{O}(10^{-4})$ at 300 K, three to four orders of magnitude smaller than observed deviations in cuprates and heavy-fermion compounds, which are therefore *not* of TGP origin; see `research/thermal_transport_mole`
- *Casimir force in MOFs:* the TGP correction is at the 10^{-4}–10^{-3} level, below current torsion-balance precision; see `research/casimir_mof/`.

These null results are stated not as corroborations but as explicit constraints on where a TGP-specific signature could plausibly emerge with present instrumentation.

9 Conclusion

We have presented a minimal, self-contained core of the Theory of Generated Space: four axioms, one scalar field with a geometrically fixed kinetic operator ($\alpha = 2$, selected within class (C1)–(C3) by Theorem 9; see Remark 10 for the v1→v2 retraction trail), a single non-linear potential with two coefficients related by a vacuum condition, and an algebraically determined effective metric whose temporal factor is the normalised substrate-potential density $U(\Phi)/\Phi^4$, reproducing general relativity exactly at 1PN with explicit, falsifiable 2PN-level deviations. Fifteen core results are formally proven, six more are numerically verified, and ten explicit open problems are stated so that the theory is both well-defined and falsifiable on a clear basis.

This core is intended to serve as a common reference for applications to superconductivity, particle masses, black-hole phenomenology and cosmology, developed in companion papers.

Data and reproducibility. All numerical scripts underlying the results of this paper are available in the accompanying source repository [1]. The present paper deposits a fixed, versioned snapshot on Zenodo (DOI assigned at publication).

Acknowledgements. The author thanks the community of open-source physics for the tools that made this work possible.

B Substrate action: continuum limit and the form $K(\varphi) = \varphi^4$

This appendix records the derivation that anchors the continuum geometric kinetic functional $\mathcal{F}^{\rm geo}_{\rm kin}[\varphi] = (K_{\rm geo}/2)\int \varphi^4 (\nabla\varphi)^2\, d^d x$ in the Ginzburg–Landau (GL) bond of the substrate Hamiltonian (3). The argument is a standard block-averaging / gradient-expansion calculation; the only point that requires care is the bookkeeping of the composite field $\hat{\Phi}_i \equiv \hat{s}_i^2$, whose lattice powers translate into integer powers of the continuum block-averaged field $\Phi(x) \equiv \langle \hat{\Phi} \rangle_{\mathcal{B}(x)}$.

Proposition 21 (Continuum limit of the substrate GL bond)**.** *Let H_Γ be the substrate Hamiltonian* (3) *on a regular hypercubic graph with spacing a and coordination $z = 2d$, with the bond term*

$$H_\Gamma^{\rm bond} \;=\; J \sum_{\langle ij \rangle} A_{ij}\, \hat{s}_i^4\, \hat{s}_j^4 \left(\hat{s}_j^2 - \hat{s}_i^2\right)^2. \tag{25}$$

Let $\Phi(x)$ be the block-averaged composite field and $\varphi(x) \equiv \Phi(x)/\Phi_0$ its dimensionless rescaling. Assume that φ is smooth on scales $\gg a$ and that the substrate respects translational and rotational invariance at the coarse-grained scale (axioms A2–A3). Then

$$\langle H_\Gamma^{\text{bond}} \rangle \;=\; \frac{K_{\text{geo}}}{2} \int \varphi^4 (\nabla\varphi)^2 \, d^d x \;+\; \mathcal{O}\big(a^{4-d}\big), \qquad K_{\text{geo}} \;=\; 2J\, a^{2-d}\, \Phi_0^6. \tag{26}$$

In particular the geometric kernel takes the form $K(\varphi) = \varphi^4$ (up to the overall normalisation K_{geo}), in agreement with the continuum exponent $\alpha = 2$ of Theorem 9 and the closure Remark 10.

Proof. Identify lattice powers of $\hat{s}$ with the composite field via $\hat{\Phi}_i = \hat{s}_i^2$, so that $\hat{s}_i^4 = \hat{\Phi}_i^2$. Block-averaging the bond gives

$$\hat{s}_i^4\, \hat{s}_j^4 \left(\hat{s}_j^2 - \hat{s}_i^2\right)^2 \;\longmapsto\; \Phi_i^2\, \Phi_j^2\, (\Phi_j - \Phi_i)^2. \tag{27}$$

For a smooth Φ and $j = i + a\hat{n}$ with $\hat{n}$ a lattice unit vector,

$$\Phi_j \;=\; \Phi_i + a\, \hat{n}\cdot\nabla\Phi(x_i) + \mathcal{O}(a^2), \tag{28}$$

hence $\Phi_i^2 \Phi_j^2 = \Phi^4(x_i) + \mathcal{O}(a)$ and $(\Phi_j - \Phi_i)^2 = a^2 (\hat{n}\cdot\nabla\Phi)^2 + \mathcal{O}(a^3)$. Summing over all bonds, each lattice site contributing $z/2 = d$ directed nearest-neighbour pairs, and using the rotational identity $\sum_{\hat{n}\in\{\pm\hat{e}_\mu\}} (\hat{n}\cdot\nabla\Phi)^2 = 2|\nabla\Phi|^2$,

$$\sum_{\langle ij\rangle} A_{ij}\, \Phi_i^2 \Phi_j^2 (\Phi_j - \Phi_i)^2 \;=\; a^2 \sum_i \Phi^4(x_i)\, |\nabla\Phi(x_i)|^2 + \mathcal{O}(a^3). \tag{29}$$

Converting the lattice sum to an integral via $\sum_i \to a^{-d} \int d^d x$,

$$H_\Gamma^{\text{bond}} \;=\; J\, a^{2-d} \int \Phi^4 (\nabla\Phi)^2 \, d^d x + \mathcal{O}(a^{3-d}). \tag{30}$$

Rescaling $\Phi = \Phi_0\, \varphi$ introduces a factor $\Phi_0^{4+2} = \Phi_0^6$, giving the announced result with $K_{\text{geo}} = 2J\, a^{2-d} \Phi_0^6$. The next-order correction in (26) is dimensionally a^{4-d} once one expands the Taylor remainder and applies parity to drop odd-a terms; on $d = 3$ it is $\mathcal{O}(a)$, vanishing under the continuum limit $a \to 0$ at fixed K_{geo}.

The identification $K(\varphi) = \varphi^4$ is then immediate: the continuum bond is $(K_{\text{geo}}/2) \int K(\varphi) (\nabla\varphi)^2 d^d x$ with $K(\varphi) = \varphi^4$, matching the exponent $\alpha = K'\varphi/(2K) = 2$ singled out by the closure conditions (C1)–(C3) of Theorem 9. □ □

Remark 22 (Why $\hat{s}^4 \hat{s}^4$ and not $\hat{s}^2 \hat{s}^2$)**.** *A naive bond $\hat{s}_i^2 \hat{s}_j^2 (\hat{s}_j^2 - \hat{s}_i^2)^2$ would Taylor-expand to $K(\varphi) = \varphi^2$, reproducing the v1 result $\alpha = 1$ and contradicting the v2 fixed-point analysis of Theorem 9. The two extra powers of $\hat{s}$ on each end of the bond are precisely what is required so that the continuum kernel matches the Z_2-even composite scaling $\hat{\Phi}_i = \hat{s}_i^2$ carried into both endpoints* and *into the gradient: schematically, $\hat{s}^4 = \hat{\Phi}^2$ on each end yields Φ^4 overall, while $\hat{s}_j^2 - \hat{s}_i^2 = \Phi_j - \Phi_i$ under the same identification, producing a single gradient. This is a v1→v2 correction noted in `KNOWN_ISSUES.md` and recorded in the present paper at* (3) *and in Remark 10.*

References

[1] M. Serafin, *Theory of Generated Space — Core paper source repository*, `https://github.com/Stefan13610/tgp-core-paper` (2026).

[2] M. Serafin, *Unified superconductivity from the TGP substrate*, companion paper, Zenodo (2026).

[3] M. Serafin, *Lepton masses and the Koide relation from a single TGP parameter*, companion paper (2026).

[4] M. Serafin, *Black-hole shadows in TGP: weak-field estimate (v1 exponential ansatz); strong-field analysis under the substrate-potential ansatz captured in OP-EHT (CLOSED CONDITIONAL POSITIVE 2026-04-25, 13/18 PASS, falsifiable by ngEHT 2030–2032 at* $\geq 10\sigma$ *on Sgr A*)*, companion paper (2026).

[5] M. Serafin, *N-body dynamics in TGP: chaos suppression and Earnshaw violation*, companion paper (2026).

[6] C. M. Will, *The confrontation between general relativity and experiment*, Living Rev. Relativity **17**, 4 (2014).

[7] B. Bertotti, L. Iess, and P. Tortora, *A test of general relativity using radio links with the Cassini spacecraft*, Nature **425**, 374 (2003).

[8] J. G. Williams, S. G. Turyshev, and D. H. Boggs, *Progress in lunar laser ranging tests of relativistic gravity*, Phys. Rev. Lett. **93**, 261101 (2004).

[9] Planck Collaboration, *Planck 2018 results. VI. Cosmological parameters*, A&A **641**, A6 (2020).

[10] LIGO/Virgo Collaboration, *Gravitational waves and gamma-rays from a binary neutron star merger: GW170817 and GRB 170817A*, ApJL **848**, L13 (2017).