


Particle Markov chain Monte Carlo methods

Vortrag am X.09.2010

Stefan Harinko
Bachelor-Kolloquium Finanzmarktökonomie SS 2010
Uni Bonn



Outline

SCM methods

Particle Independent
Metropolis Hastings


Particle Marginal
Metropolis Hastings

Particle Gibbs

Outline

- paper by Andrieu, Doucet, Holenstein: **Particle Markov chain Monte Carlo methods**
- goal: inference in State Space Models
- how: combination of sequential Monte Carlo (SMC) and Markov chain Monte Carlo (MCMC) methods

- 1 Outline
- 2 SCM methods
- 3 Particle Independent Metropolis Hastings
- 4 Particle Marginal Metropolis Hastings
- 5 Particle Gibbs



Outline

SCM methods

Particle Independent
Metropolis Hastings

Particle Marginal
Metropolis Hastings

Particle Gibbs

state space models: Hidden Markov Models

state assumptions


- states: $\{X_n; n \geq 1\} \subset \mathcal{X}^{\mathbb{N}}$
- $X_1 \sim \mu_{\theta}(\cdot)$
- $X_{n+1} | (X_n = x) \sim f_{\theta}(\cdot | x)$
- parameters: $\theta \in \Theta$

observation assumptions

- observations: $\{Y_n; n \geq 1\} \subset \mathcal{Y}^{\mathbb{N}}$
- $Y_{n+1} | (X_1, \dots, X_n = x, \dots, X_m) \sim g_{\theta}(\cdot | x)$

interested in:

- $p_{\theta}(x_{1:n} | y_{1:n})$
- even more: $p(\theta, x_{1:n} | y_{1:n})$



Outline

SCM methods

Particle Independent
Metropolis Hastings

Particle Marginal
Metropolis Hastings

Particle Gibbs

Bayesian inference conditional on observations $y_{1:T}$

Bayesian inference

- $p_{\theta}(x_{1:n}|y_{1:n}) \propto p_{\theta}(x_{1:n}, y_{1:n})$
- with


$$p_{\theta}(x_{1:T}, y_{1:T}) = \mu_{\theta}(x_1) \prod_{n=2}^T f_{\theta}(x_n|x_{n-1}) \prod_{n=1}^T g_{\theta}(y_n|x_n)$$

- also: $p_{\theta}(x_{1:T}|y_{1:T}) \propto p_{\theta}(x_{1:T-1}, y_{1:T-1}) f_{\theta}(x_T|x_{T-1}) g_{\theta}(y_T|x_T)$

particle filter

- weights and particles: $(w_n^{(i)}, x_{1:n}^{(i)})_{i=1}^N \sim p_{\theta}(x_{1:n}|y_{1:n})$
- new particles: $x_{n+1}^{(i)} \sim q(\cdot|x_{1:n}^{(i)}, y_{n+1})$
- remember Importance Sampling: correct for "wrong distribution"
-

$$w_{n+1}^{(i)} = w_n^{(i)} \frac{f_{\theta}(x_{n+1}^{(i)}|x_n^{(i)}) g_{\theta}(y_{n+1}|x_{n+1}^{(i)})}{q(x_{n+1}|x_{1:n}^{(i)}, y_{n+1})}$$



Outline



SCM methods

Particle Independent
Metropolis Hastings

Particle Marginal
Metropolis Hastings

Particle Gibbs

bootstrap filter illustration



Outline

SCM methods

Particle Independent
Metropolis Hastings


Particle Marginal
Metropolis Hastings

Particle Gibbs

- we get particles and weight $\sim p_\theta(x_{1:T}|y_{1:T})$
- we also get marginal likelihood:

$$\prod_{n=1}^T \left(\frac{1}{N} \sum_{i=1}^N w_n^i \right) \xrightarrow{N \rightarrow \infty} p_\theta(y_{1:T}) := \hat{Z}$$

- Particle MCMC idea: use SMC as proposal in Metropolis-Hastings algorithms
- use SMC distributions as conditional distributions in Gibbs algorithm



Outline

SCM methods

Particle Independent
Metropolis Hastings

Particle Marginal
Metropolis Hastings

Particle Gibbs

Particle Independent Metropolis Hastings algorithm

- run SMC, sample


$$x_{1:T}^* \sim \hat{p}_\theta(dx_{1:T}|y_{1:T})$$

- compute simulated likelihood

$$\hat{Z}^* = p_\theta(\hat{y}_{1:T})$$

- accept with

$$1 \wedge \frac{\hat{Z}^*}{\hat{Z}^i}$$



Outline

SCM methods

Particle Independent Metropolis Hastings

Particle Marginal Metropolis Hastings

Particle Gibbs

nonlinear toy example


-

$$X_n = \frac{X_{n-1}}{2} + 25 \frac{X_{n-1}}{1 + X_{n-1}^2} + 8 \cos(1.2n) + V_n$$

-

$$Y_n = \frac{X_n^2}{20} + W_n$$

- $X_1 \sim^{IID} \mathcal{N}(0, 5)$
- $V_n \sim^{IID} \mathcal{N}(0, \sigma_v^2)$
- $W_n \sim^{IID} \mathcal{N}(0, \sigma_w^2)$



Outline

SCM methods

Particle Independent Metropolis Hastings

Particle Marginal Metropolis Hastings

Particle Gibbs

single bootstrap filter run





Figure: true states x and Bootstrap median estimates marked with points



Outline

SCM methods

Particle Independent Metropolis Hastings

Particle Marginal Metropolis Hastings

Particle Gibbs





Figure: PIMH $N=50$ particles, $n=50$ observations, 10000 iterations





Outline

SCM methods

Particle Independent Metropolis Hastings

Particle Marginal Metropolis Hastings

Particle Gibbs



Outline

SCM methods

Particle Independent Metropolis Hastings

Particle Marginal Metropolis Hastings

Particle Gibbs

Figure: PIMH $N=6000$ particles, $n=50$ observations, 10000 iterations

- interested in θ and $x_{1:T}$ from
 $p_{\theta}(\theta, x_{1:T}|y_{1:T}) = p(\theta|y_{1:T})p_{\theta}(x_{1:T}|y_{1:T})$
- Andrieu, Doucet, Holenstein, proved to be the correct MH ratio:


$$1 \wedge \frac{\hat{p}_{\theta^*}(y_{1:T})p(\theta^*)q\{\theta(i-1)|\theta^*\}}{\hat{p}_{\theta(i-1)}(y_{1:T})p(\theta)q\{\theta^*|\theta(i-1)\}}$$

- applied to a basic stochastic volatility (SV) model:

$$y_t = e^{h_t/2}\varepsilon_t$$

$$h_t = \mu + \phi h_{t-1} + \tau \eta_t$$

where $\varepsilon_t, \eta_t \sim \mathcal{N}(0, 1)$, $\mathbb{E}(\varepsilon_t \eta_{t+h}) = 0$ for all h and
 $\mathbb{E}(\varepsilon_t \varepsilon_{t+l}) = \mathbb{E}(\eta_t \eta_{t+l}) = 0$ for all $l \neq 0$





Outline

SCM methods

Particle Independent
Metropolis Hastings

Particle Marginal
Metropolis Hastings

Particle Gibbs



Outline



SCM methods

Particle Independent Metropolis Hastings

Particle Marginal Metropolis Hastings

Particle Gibbs

Figure: SV model $N=1000$ Particles $n=100$ observations 10000 iterations



Outline

SCM methods

Particle Independent Metropolis Hastings


Particle Marginal Metropolis Hastings

Particle Gibbs

Figure: SV model N=1500 particles 250 observations 40000 mcmc iterations

Particle Gibbs

- iteratively sample from $p(\theta|x_{1:T}, y_{1:T})$ and with SMC from $p_{\theta}(x_{1:T}|y_{1:T})$
- important in the Particle Gibbs: conditioning on last particle-path



Outline

SCM methods

Particle Independent
Metropolis Hastings

Particle Marginal
Metropolis Hastings

Particle Gibbs





Figure: conditionalSMC starting with dashed lineage



Outline

SCM methods

Particle Independent
Metropolis Hastings

Particle Marginal
Metropolis Hastings

Particle Gibbs





Figure: conditionalSMC N-1 new path conditioned on dashed path




Outline

SCM methods

Particle Independent
Metropolis Hastings

Particle Marginal
Metropolis Hastings

Particle Gibbs



Proofs

- formalize SMC on extended space including all random variables
- for Particle MH show: acceptance ratios lead to target distribution
- for Particle Gibbs show full conditional densities and "collapsed Gibbs"-step



Outline

SCM methods

Particle Independent
Metropolis Hastings

Particle Marginal
Metropolis Hastings

Particle Gibbs



Outline



SCM methods

Particle Independent Metropolis Hastings

Particle Marginal Metropolis Hastings

Particle Gibbs

Figure: Particle Gibbs with MH step for nonlinear model, 40.000 iterations



Outline

SCM methods

Particle Independent Metropolis Hastings

Particle Marginal Metropolis Hastings

Particle Gibbs

Figure: Particle Gibbs for nonlinear model, 40.000 iterations





Figure: toy model N250 particles n250 observations



Outline

SCM methods

Particle Independent
Metropolis Hastings

Particle Marginal
Metropolis Hastings

Particle Gibbs




Figure: toy model N250 particles n300 observations



Outline

SCM methods

Particle Independent
Metropolis Hastings

Particle Marginal
Metropolis Hastings

Particle Gibbs





Figure: toy model N250 particles n350 observations





Outline

SCM methods

Particle Independent
Metropolis Hastings

Particle Marginal
Metropolis Hastings

Particle Gibbs



Outline

SCM methods

Particle Independent Metropolis Hastings

Particle Marginal Metropolis Hastings

Particle Gibbs

Figure: toy model N250 particles n350 observations 70000 mcm iterations





Figure: toy model N500 particles n350 observations



Outline

SCM methods

Particle Independent Metropolis Hastings

Particle Marginal Metropolis Hastings

Particle Gibbs




Figure: PH with MH toy model N500 particles n350 observations





Outline

SCM methods

Particle Independent Metropolis Hastings

Particle Marginal Metropolis Hastings

Particle Gibbs



Outline

SCM methods

Particle Independent Metropolis Hastings

Particle Marginal Metropolis Hastings

Particle Gibbs

Figure: PG toy model N700 particles n350 observations





Figure: PG for toy model (10,10) 10000 iterations



Outline

SCM methods

Particle Independent Metropolis Hastings

Particle Marginal Metropolis Hastings

Particle Gibbs





Figure: PMMH for toy model (10,10) 10000 iterations



Outline

SCM methods

Particle Independent Metropolis Hastings

Particle Marginal Metropolis Hastings

Particle Gibbs