

User testing

Experimental Design, Basic Statistics

Experimental Design

Example Experiment



How much do people
like these crisps?



Which crisps do they
like more?

Dependent variables

- What you want to measure
- For example, for the crisps could have 2:
 - *User satisfaction:*

How much they say they like them, say on a scale from 1 to 7 (1=I really hate them, 7=I love them)
 - *Amount eaten:*

How many (in grams) they eat if left alone with a bowl of crisps, without knowing this is what we are interested in

Independent variables

- Experimental factors the effect of which you want to measure
- For example, for the crisps:
 - The brand of crisp (Brand A, Brand B)Or
 - The crisp flavouring (neutral, cheese, chilli)

Aside: brand in this example has 2 so-called *levels* (Brand A and B), whilst crisp flavouring has 3 levels.

Another Example Experiment



How good are these
running shoes?



Which running shoes
are better?

Question: Variables

- What dependent variable(s) may we be interested in for running shoes?
- What independent variable(s)?

Often in User-Testing: Compare A to B

Compare

the performance (on the dependent variable)
of a group of users who experienced a
certain level of the independent variable (A)

with

the performance of a group of users who
experienced another level of the
independent variable (B)

Examples

- Compare two systems:
 - Is Virgin's on-line shopping site more usable than Amazon's?
 - Is the new system more usable than the old one?
- Compare two designs:
 - Is my system more usable with drop-down menus or with lists?

Aim of experimental design



Highlight effect of independent variable
while avoiding undesired effects
by strictly controlling the influence of
irrelevant variables

Irrelevant variables (1)

- User variables
 - Age
 - Sex
 - Education
 - Cultural background
 - Experience with computers
 - Etc. etc.


(unless this is the purpose of your experiment)

Is this correct?




Is this correct?

Group 1



Group 2





Let's try making groups

- I need two volunteers
- I want you to select two groups of 4 people, so that both groups are as similar as possible

Within subjects design

Use the same subjects in each group
(also called Repeated Measures)



Within subjects design (2)

Advantage

- Subject variables (like age, sex, education, intelligence) are the same for both groups

Disadvantage

- More time needed per subject
- Order effect due to practice, fatigue, etc

Within subjects design (3)

Counterbalancing:


- half the subjects perform the tasks in one order (A-B)
- half the subjects perform the tasks in the reverse order (B-A)

Disadvantage

- Statistical analysis more difficult

Matched pairs design

- Divide subjects in pairs of similar subjects
- For each pair: randomly assign one subject to group 1 and the other to group 2



Matched pairs design (2)



- When analyzing, compare pair-wise

Disadvantages:

- Difficult to match pairs
- Statistics more difficult

Between subjects design

Use two different groups of subjects, allocate subjects randomly



Between subjects design (2)


Advantages

- No order effects
- Less time needed per subject

Disadvantages

- More subjects needed
- Less powerful
- Need to carefully select subjects

Choosing the design



Irrelevant variables (2)

- Situational variables
 - Physical characteristics of the experimental room, like light, temperature, noise,...

- Equipment



- Experimenter

Real-life versus laboratory



Control vs Realism

- Control: Ensuring 'irrelevant' variables do not influence the outcome
- Realism: Ensuring the results hold in the real world

This is a trade-off, cannot always do both...

Basic statistics

Misconception

- “With statistics you can prove anything....”

No, you cannot, but it is easy to confuse people with numbers.

I want you to be able to:

- criticize other people’s misuse of numbers
- understand relation with experimental design

Critical thinking required!

- I will show you some ‘stories’ I have heard, which use numbers.
- Tell me what is wrong with them!

On the radio....

- “In 65% of domestic violence cases, the offender is drunk. Alcoholism causes violence.”

In the train..

- “I work for the council, providing information to people in my area about the environment and recycling. The glass-recycling containers in my area are *much fuller* than in other areas. This shows how good I am at my job!”

In the newspaper

- The newspapers have published a *ranking list* for schools: a school is higher in the ranking if a higher percentage of its pupils do well at the national exams.

A mother sees this ranking and decides that her daughter should go to a particular school because it is higher in the ranking than other schools in the area.

How does this relate to experimental design?

- The misconception in the stories came from someone overlooking some variables that may have influenced the measurements.
- These should have been controlled for through good experimental design.

Hypothesis testing

A hypothesis should be

- Stated in clear language
- A question with yes or no answer
- Answerable using a small set of independent and dependent variables.

Examples:

H_1 People will say they like crisps of Brand A more than crisps of Brand B.

H_2 The amounts people eat from A and B will differ

Statistical Hypothesis

Because of the way statistics works, statisticians want to *reject* a hypothesis!

Therefore, a statistician will state a hypothesis (called null hypothesis) in the opposite way.

H_0 There will be no significant difference in how much

1. People say they like crisps of Brand A compared to B
2. Is eaten from A and B

Two types of errors stats can make

- Type I (also called false positive):
Null Hypothesis rejected while actually it is true
(so, original hypothesis 'proven' while actually it is false) This is very bad..
- Type II (also called false negative):
Null Hypothesis not rejected while it is false.
(so, original hypothesis not 'proven' though true)
This is not so bad.. Found not enough evidence yet..
May need more participants!

p-value

- p-value is the probability of a Type I error, so the probability that the effect you are seeing is due to chance
- Normally, you want $p < 0.05$ to say that something is *statistically significant*
- You run a statistical test on your data, and it tells you the p-value.

What statistics to use? (1)

One independent variable, two treatments

(Example: brand of crisps, one group A, other B)

Use T-test*.

Computed based on the difference in means of the two groups and how spread out the data is (standard deviation).

T-test is available as a function in Excel.

* If data is normally distributed.

Otherwise non-parametric test (Mann-Whitney U)

T-Test in Excel

Tails: 2 if test whether different

1 if test larger/smaller

Type: Paired: Within subject

Two-sample: Between subject

The screenshot shows an Excel spreadsheet with two columns of data: Group1 and Group2. The data is as follows:

	Group1	Group2
1	10	1
2	8	2
3	9	1
4	9	3
5	7	1
6	8	3
7	10	2
8	9	1

In cell A11, the text "T-test" is entered. In cell B11, the formula "=TTEST()" is entered. The "Function Arguments" dialog box for the TTEST function is open, showing the following fields:

- Array1: (empty)
- Array2: (empty)
- Tails: (empty)
- Type: (empty)

The dialog box also includes a description: "Returns the probability associated with a Student's t-Test." and a note: "Type is the kind of t-test: paired = 1, two-sample equal variance (homoscedastic) = 2, two-sample unequal variance = 3." The "Formula result =" field is empty. The "OK" and "Cancel" buttons are at the bottom right.

Result is p-value

What statistics to use? (2)

One independent variable, more than two treatments.

Use one-way ANOVA*.

Two independent variables, any number of treatments for each one.

Use two-way ANOVA.

*If data is normally distributed. Otherwise Kruskal-Wallis.

Example User Test: Transport Project

Does our journey planner result in more sustainable use of transport?

Main issues

Transport use / experience depends on:

- people's circumstances, such as home address, bus routes, having children
=> Used within-subject design
- people's normal modes of transport used
=> Used stratified sampling
- weather, events on specific days
=> Used staggering & Added control group