

Physics Concepts and Fundamentals

Generic introduction to some concepts in physics

July 24, 2026

Contents

1	Generic Mathematical Tools	3
1.1	Inner Product	3
1.2	Tensor Product	3
1.3	Legendre Transform	4
1.4	Differentials	4
1.5	Integrals	4
2	Newtonian Mechanics	5
3	Lagrangian Mechanics	5
3.1	Mass Matrix	5
3.2	Lagrange's Equations	6
4	Hamiltonian Mechanics	7
4.1	Phase Space	8
5	Statistical Mechanics	8
5.1	Canonical Ensemble	8
5.2	Grand Canonical Ensemble	10
6	Fourier Analysis	10
7	Information Theory	11
8	Game Theory	12
8.1	Shapley Value	12
9	Network Theory	12
9.1	Types of Matrixes	12
9.2	Types of Networks	15
9.3	Centralities	15
9.4	Diffusion on Graphs (analytical)	19
9.5	Experimental Diffusion	20
9.6	Path Finding	21
9.7	Link Prediction	23

10 Markov Processes	23
11 Time Series	24
11.1 Theoretical Description	24
11.2 Technical Difficulties	25
11.3 Algorithms	25
12 Likelihood Function	25
12.1 Probability Functions	25
12.2 Maximum Likelihood Estimation	26
13 P-Value	27
14 Monte Carlo Simulation	28
15 Quantum Mechanics	28
15.1 Hilber Space	28
15.2 Nomenclature	28
15.3 State Vectors, Basis and Descriptions	29
15.4 Define Observable Operators	32
15.4.1 Commutator	32
15.4.2 Generating Observables	33
15.5 Schrödinger Equation	33

1 Generic Mathematical Tools

1.1 Inner Product

Let V be a space vector and F be the field on which the vector space is defined. The inner product – or dot/scalar product – is a bilinear transformation $\langle \cdot, \cdot \rangle$ such that:

$$\begin{aligned}\langle \cdot, \cdot \rangle : V \times V &\rightarrow F \\ \vec{v}, \vec{u} &\mapsto \langle \vec{v}, \vec{u} \rangle = \sum_{i=1}^N v_i u_i\end{aligned}$$

In vector form we can write the inner product as:

$$u^\top v = \begin{bmatrix} u_1 & u_2 & \cdots & u_n \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ \vdots \\ v_n \end{bmatrix} = \sum_{i=1}^N v_i u_i$$

1.2 Tensor Product

Let V be an N -dimensional vector space over a field $\mathbb{F}$. The tensor product is a bilinear map

$$\otimes : V \times V \longrightarrow V \otimes V,$$

defined by

$$(\vec{u}, \vec{v}) \longmapsto \vec{u} \otimes \vec{v}.$$

Let $\{\vec{e}_1, \dots, \vec{e}_N\}$ be a basis of V (canonical basis), and write

$$\vec{u} = \sum_{i=1}^N u_i \vec{e}_i, \quad \vec{v} = \sum_{j=1}^N v_j \vec{e}_j.$$

By bilinearity, their tensor product is

$$\vec{u} \otimes \vec{v} = \sum_{i=1}^N \sum_{j=1}^N u_i v_j \vec{e}_i \otimes \vec{e}_j.$$

The tensors

$$\{\vec{e}_i \otimes \vec{e}_j\}_{1 \leq i, j \leq N}$$

form a basis of $V \otimes V$. Therefore,

$$\dim(V \otimes V) = N^2.$$

Once a basis is chosen, the tensor $\vec{u} \otimes \vec{v}$ can be represented by the coefficient matrix

$$[\vec{u} \otimes \vec{v}] = \begin{bmatrix} u_1 v_1 & u_1 v_2 & \cdots & u_1 v_N \\ u_2 v_1 & u_2 v_2 & \cdots & u_2 v_N \\ \vdots & \vdots & \ddots & \vdots \\ u_N v_1 & u_N v_2 & \cdots & u_N v_N \end{bmatrix}.$$

In matrix notation, this is the outer product

$$[\vec{u} \otimes \vec{v}] = \vec{u}\vec{v}^\top.$$

More generally, an arbitrary tensor $\mathbf{T} \in V \otimes V$ has the form

$$\mathbf{T} = \sum_{i=1}^N \sum_{j=1}^N T_{ij} \vec{e}_i \otimes \vec{e}_j,$$

and is represented, in the chosen basis, by the matrix

$$[\mathbf{T}] = \begin{bmatrix} T_{11} & \cdots & T_{1N} \\ \vdots & \ddots & \vdots \\ T_{N1} & \cdots & T_{NN} \end{bmatrix}.$$

Thus, a simple tensor $\vec{u} \otimes \vec{v}$ is represented by a rank-one outer-product matrix $\vec{u}\vec{v}^\top$, whereas a general tensor in $V \otimes V$ is represented by an arbitrary $N \times N$ matrix.

1.3 Legendre Transform

Method to rewrite a convex function as a different one replacing one variable with the derivative of the original function:

$$L_T(f(\vec{x}))(\vec{p}) = \sup_{\vec{x} \in \mathbb{R}^N} [\langle \vec{p}, \vec{x} \rangle - f(\vec{x})] , \quad (1)$$

Where it emerges that the sup is obtained exactly at a certain $\vec{x}^* = \nabla f(\vec{x}^*) \equiv \vec{p}$.

This transform is crucial in Hamiltonian mechanics to switch between a second order differential equation, to a system of first order differential equations (Hamilton equations).

1.4 Differentials

Let $f(\vec{x})$ be a multivariate function. The differential of f can be defined as the overall infinitesimal displacement of f in any direction:

$$df(\vec{x}) = \frac{\partial f}{\partial x_1} dx_1 + \frac{\partial f}{\partial x_2} dx_2 + \cdots + \frac{\partial f}{\partial x_n} dx_n . \quad (2)$$

The differential is a linear operator in the dual space of the tangent plane of f in $\vec{x}$ on the manifold M where the function lives:

$$df_{\vec{x}} : T_{\vec{x}}M \rightarrow \mathbb{R} , \quad df_{\vec{x}} \in T_{\vec{x}}^*M$$

$$\vec{v} \mapsto df_{\vec{x}}(\vec{v}) = \sum_{i=1}^n \left. \frac{\partial f}{\partial x_i} \right|_{\vec{x}} v_i$$

1.5 Integrals

Generalization of an infinite sum of a function over an interval divided in infinitesimal steps equally spaced. We start by considering the finite steps Δx ..

Complex integral ...

2 Newtonian Mechanics

Solve the dynamics of a system by modeling it to a vectorial problem and applying the principles in order to get a second order differential equation.

- $\vec{r}(t)$: position $\longrightarrow \frac{d}{dt}(\vec{r}(t)) = \vec{v}(t) \equiv \dot{\vec{x}}(t)$: velocity
- $\vec{v}(t)$: velocity $\longrightarrow \frac{d}{dt}(\vec{v}(t)) = \vec{a}(t) \equiv \dot{\vec{v}}(t) = \ddot{\vec{x}}(t)$: acceleration

The second principle of Newtonian mechanics is:

$$\vec{F}(t) = m\vec{a}(t) \quad (3)$$

therefore we set the differential system:

$$\begin{cases} \vec{F}(t) = m\vec{a}(t) = m\frac{d}{dt}(\vec{v}(t)) \\ \vec{v}(t) = \frac{d}{dt}(\vec{r}(t)) \\ \vec{v}(0) = \vec{v}_0 \\ \vec{r}(0) = \vec{r}_0 \end{cases} \Rightarrow m\frac{d^2}{dt^2}(\vec{r}(t)) = F(\vec{r}, t)$$

Hooke example..

3 Lagrangian Mechanics

Solve the dynamics of a system using only algebraic procedures to get the same second order differential equation system coming from Newtonian mechanics – but it is general even to non relativistic mechanics.

3.1 Mass Matrix

Before setting the Lagrangian system, we need to define the *mass matrix* so that we can generalize the concept of **kinetic energy**. We define the kinetic energy as a scalar quantity in the i -th direction:

$$d\mathcal{T}_i = dr_i F_i = dr_i m_i \ddot{r}_i = \frac{dr_i}{dt} m_i d\dot{r}_i ,$$

that we can rewrite using $d(\dot{r}_i \cdot \dot{r}_i) = 2\dot{r}_i d\dot{r}_i$ as:

$$\begin{aligned} d\mathcal{T}_i &= m_i \frac{1}{2} d(\dot{r}_i^2) \\ \Rightarrow \mathcal{T}_i &= \int_{\dot{r}_0}^{\dot{r}_1} \frac{1}{2} m_i d(\dot{r}_i^2) = \frac{1}{2} m_i (\dot{r}_{i,1}^2 - \dot{r}_{i,0}^2) = \frac{1}{2} m_i \dot{r}_i^2 \end{aligned}$$

We can write all the contributions to the energy in a vectorial form:

$$\mathcal{T} = \frac{1}{2} \dot{\vec{r}}^\top M(\vec{r}, t) \dot{\vec{r}} ,$$

with $M(\vec{r}, t)$ being the **mass matrix**. In the euclidean coordinates, $M(\vec{r}, t) \equiv M_{eu}$ is diagonal filled with the single masses – the difficulty is in defining the velocities. Using the generalized coordinates instead $\vec{r} = \vec{r}(\vec{q})$ we get:

$$\frac{d\vec{r}(\vec{q})}{dt} = \sum_i \frac{\partial \vec{r}_k}{\partial q_i} \dot{q}_i + \frac{\partial \vec{r}(\vec{q})}{\partial t} = J_q \dot{\vec{q}} + \frac{\partial \vec{r}(\vec{q})}{\partial t} = \dot{\vec{r}}(\vec{q}) .$$

Substituting this into E and reordering the terms we get:

$$\mathcal{T} = \frac{1}{2} \left(\dot{\vec{q}}^\top M(\vec{q}, t) \dot{\vec{q}} + \vec{M}_v(\vec{q}, t) \cdot \dot{\vec{q}} + M_0(\vec{q}, t) \right) ,$$

where the details are the following:

$$M(\vec{q})_{i,j} = \sum_k m_k \frac{\partial \vec{r}_k}{\partial q_i} \frac{\partial \vec{r}_k}{\partial q_j} = \left(J_q^\top M_{eu} J_q \right)_{i,j} \longrightarrow \text{Generalized inertia}$$

$$\vec{M}_v(\vec{q})_i = \sum_k m_k \frac{\partial \vec{r}_k}{\partial q_i} \frac{\partial \vec{r}_k}{\partial t} = \left(J_q^\top M_{eu} \frac{\partial \vec{r}}{\partial t} \right)_i \longrightarrow \text{Linear velocity coefficient}$$

$$M_0(\vec{q}) = \sum_k m_k \left(\frac{\partial \vec{r}_k}{\partial t} \right)^2 \longrightarrow \text{Time dependent potential-like term}$$

which for the systems in which the coordinates transformation is not affected by the time, then we have:

$$\mathcal{T} = \frac{1}{2} \dot{\vec{q}}^\top M \dot{\vec{q}} . \quad (4)$$

Clearly if we are dealing with points of the same mass (or a single point) $M = m_{tot} I$, with I being the identity matrix.

3.2 Lagrange's Equations

We start by defining one of the most profound empirical observations in physics: the **principle of Least Action**. This fundamental principle states that, for any dynamical system, the path followed by the system between two states is the one that minimizes a quantity called **action** S , which is defined as:

$$S = \int_{t_0}^{t_1} \mathcal{L}(\vec{q}, \dot{\vec{q}}, t) dt , \quad (5)$$

where $\mathcal{L}(\vec{q}, \dot{\vec{q}}, t)$ is the a specific function embedding the dynamics of the system. This function is called **Lagrangian** and it must be a function of $\vec{q}$ and $\dot{\vec{q}}$ the generalized coordinates describing the system.

The action depends on the path followed by the system, so we cannot write it as a differential. However, if we consider small variations in the path – not time – keeping the endpoints fixed we can write:

$$\delta S = \int_{t_0}^{t_1} \left(\frac{\partial \mathcal{L}}{\partial \vec{q}} \delta \vec{q} + \frac{\partial \mathcal{L}}{\partial \dot{\vec{q}}} \delta \dot{\vec{q}} \right) dt . \quad (6)$$

Integrating by parts the second term:

$$\begin{aligned}\delta S &= \int_{t_0}^{t_1} \frac{\partial \mathcal{L}}{\partial \vec{q}} \delta \vec{q} dt + \left[\frac{\partial \mathcal{L}}{\partial \dot{\vec{q}}} \delta \vec{q} \right]_{t_0}^{t_1} - \int_{t_0}^{t_1} \frac{d}{dt} \left(\frac{\partial \mathcal{L}}{\partial \dot{\vec{q}}} \right) \delta \vec{q} dt = \\ &= \int_{t_0}^{t_1} \left(\frac{\partial \mathcal{L}}{\partial \vec{q}} \delta \vec{q} - \frac{d}{dt} \frac{\partial \mathcal{L}}{\partial \dot{\vec{q}}} \right) \delta \vec{q} dt\end{aligned}$$

where we used the fact that $\delta \vec{q}(t_0) = \delta \vec{q}(t_1) = 0$, since the endpoints are fixed. For the principle of least action to hold, we need $\delta S = 0$ for any arbitrary variation $\delta \vec{q}$. This is possible only if the integrand is zero, thus:

$$\frac{d}{dt} \left(\frac{\partial \mathcal{L}}{\partial \dot{\vec{q}}} \right) - \frac{\partial \mathcal{L}}{\partial \vec{q}} = 0, \quad (7)$$

which is the **Lagrange's equation**.

Now we want to find a relation between the Lagrangian $\mathcal{L}(\vec{q}, \dot{\vec{q}}, t)$ with the known Newtonian mechanics. We observe a similarity between the Lagrange's equation and the Newton's second law:

$$\begin{aligned}m \ddot{\vec{r}} &= -\nabla V(\vec{r}, t) \\ \frac{d}{dt} \left(\frac{\partial \mathcal{L}}{\partial \dot{\vec{q}}} \right) &= \frac{\partial \mathcal{L}}{\partial \vec{q}} \\ \Rightarrow \begin{cases} \frac{\partial \mathcal{L}}{\partial \vec{q}} \sim -\nabla V(\vec{r}, t) \\ \frac{\partial \mathcal{L}}{\partial \dot{\vec{q}}} \sim m \dot{\vec{r}} \end{cases}\end{aligned}$$

From this analogy, we can define the Lagrangian as:

$$\mathcal{L}(\vec{q}, \dot{\vec{q}}, t) = \mathcal{T}(\vec{q}, \dot{\vec{q}}, t) - V(\vec{q}, t) + g(t), \quad (8)$$

where $\mathcal{T}$ is the kinetic energy expressed in generalized coordinates, V is the potential energy of the system and $g(t)$ is a generic function of time that doesn't affect the dynamics. With this definition, we can recover exactly the Newtonian dynamics using the Lagrange's equation, avoiding all the geometrical considerations about forces.

This definition of Lagrangian is very general, and holds also for relativistic mechanics if we define all the quantities properly.

4 Hamiltonian Mechanics

Hamiltonian mechanics arises from Lagrangian mechanics by rewriting the second order differential equation as a system of first order differential equations. In particular, for this purpose we use the **Legendre transform** as follows:

$$\begin{aligned}L_T(\mathcal{L}(q_i, \dot{q}_i, t))(q_i, p_i, t) &= \sum_i \dot{q}_i \frac{\partial \mathcal{L}}{\partial \dot{q}_i} - \mathcal{L}(q_i, \dot{q}_i, t) \\ \Rightarrow \mathcal{H}(q_i, p_i, t) &= \sum_i \dot{q}_i p_i - \mathcal{L}(q_i, \dot{q}_i, t)\end{aligned}$$

The dynamics of the system is now described by the system:

$$\begin{cases} \frac{dq_i}{dt} = \frac{\partial \mathcal{H}}{\partial p_i} \\ \frac{\partial \mathcal{L}}{\partial q_i} = \frac{d}{dt} \left(\frac{\partial \mathcal{L}}{\partial \dot{q}_i} \right) = \frac{d}{dt}(p_i) = -\frac{\partial \mathcal{H}}{\partial q_i} \end{cases}$$

which gives the Hamilton equations:

$$\begin{cases} \frac{dq_i}{dt} = \frac{\partial \mathcal{H}}{\partial p_i} \\ \frac{dp_i}{dt} = -\frac{\partial \mathcal{H}}{\partial q_i} \end{cases}$$

Hamilton equations are crucial when describing the dynamics of a mechanical system because they are directly connected to the concept of **phase space**. The phase space is the (p, q) plane that encapsulate all the dynamic of the system in terms of energy and possible trajectories. In addition, almost the totality of numerical integrators requires first order differential equations, which is exactly what we obtained from the Legendre transform.

4.1 Phase Space

The phase space is an extremely useful tool to extrapolate quantitative/qualitative information about dynamical systems, in particular about the energy of systems. Once we managed to write the Hamiltonian of the system $\mathcal{H}(\vec{q}, \vec{p})$ we can apply hamilton equation to get $\dot{\vec{q}}$ and $\dot{\vec{p}}$ and solve the dynamics of the system.

The graph in the phase plane can be usually drawn directly from $\mathcal{H}(\vec{q}, \vec{p})$ by writing p in function of q .

In the example below we see:

$$\mathcal{H}(\vec{q}, \vec{p}) = \frac{\vec{p}^2}{2} + \frac{\vec{q}^2}{2}$$

starting from different initial conditions (which means different energies). Each closed curve in the phase space corresponds to a specific energy configuration of the system.

5 Statistical Mechanics

Statistical Mechanics is the main tool to describe high degrees of freedom systems, in particular by extracting generic propertis from statistical considerations. Many fields are based on statistical mechanics, from material physics to social network analysis.

Assume to have a system s that can occupy discrete energy states ε_i . The system s interacts with a reservoir R , so that the total energy $E_{tot} = \varepsilon_i + \varepsilon_R$.

5.1 Canonical Ensemble

In the canonical configuration, the systems s and R exchange only energy, not matter. This means that each energy ε_i can be obtained in $g(\varepsilon_i)$ number of ways. The function




Figure 1: Phase-space trajectories of the harmonic oscillator for different energies.

$g(\varepsilon)$ is called **degeneracy** of the system s . The reservoir degeneracy is $g_R(E_{tot} - \varepsilon_i)$. We assume that all the states i of the whole system $s + R$ at an energy E_{tot} are *equally probable*, which is reasonable.

Since R is large compared to s , any change in R is a smooth function of ε , which means that we do not have sharp jumps in the physical quantities.

We define a quantity called **entropy** as:

$$S(E) = k_B \log(g(E))$$

which is a transformation of the degeneracy, in other words it is a measure of the probabilistic configuration of the system. With the entropy being defined, we can invert the relation:

$$g(E) = \exp\left(\frac{S(E)}{k_B}\right),$$

which means that the number of possible states at an energy ε_i for the whole system $s + R$ is:

$$N(\varepsilon_i) = g(\varepsilon_i) g_R(E_{tot} - \varepsilon_i).$$

Since each microstate has the same probability to occur, that will be proportional to:

$$P(\varepsilon_i) \propto N(\varepsilon_i) = g(\varepsilon_i) \exp\left(\frac{S(E_{tot} - \varepsilon_i)}{k_B}\right),$$

and since we know that $s \ll R$ we can expand with Taylor $S(E_{tot} - \varepsilon_i)$ around $(E_{tot} - \varepsilon_i) \sim E_{tot}$:

$$S(E_{tot} - \varepsilon_i) = S(E_{tot}) - \varepsilon_i \left. \frac{dS(E)}{dE} \right|_{E_{tot}} + \frac{\varepsilon_i^2}{2} \left. \frac{d^2 S(E)}{dE^2} \right|_{E_{tot}} + \dots$$

If we call the quantity

$$\left. \frac{dS(E)}{dE} \right|_{E_{tot}} \equiv \frac{1}{T}$$

keeping only the first order term we can rewrite the probability $P(\varepsilon_i)$ as:

$$P(\varepsilon_i) \propto g(\varepsilon_i) \cdot \exp\left(\frac{S(E_{tot})}{k_B}\right) \cdot \exp\left(\frac{-\varepsilon_i}{k_B T}\right) \cdot \dots$$

$$\Rightarrow P(\varepsilon_i) \propto g(\varepsilon_i) \exp(-\beta \varepsilon_i) , \quad \beta = \frac{1}{k_B T}$$

and since the relation $\sum_i P(\varepsilon_i) = 1$ must hold, we get that the probability to find the system in an energy configuration ε_i is:

$$P(\varepsilon_i) = \frac{g(\varepsilon_i) \exp(-\beta \varepsilon_i)}{\mathcal{Z}(\beta)} , \quad \mathcal{Z}(\beta) = \sum_j g(\varepsilon_j) \exp(-\beta \varepsilon_j)$$

where $\mathcal{Z}(\beta)$ is called **canonical partition function** and it's the normalization coefficient of the probability.

For dynamical systems we use the **Hamiltonian** as the energy:

$$\mathcal{Z}(\beta) = \sum_j g(\varepsilon_j) \exp(-\beta \mathcal{H}_i(\vec{q}, \vec{p}))$$

5.2 Grand Canonical Ensemble

...

6 Fourier Analysis

Fourier formalism is based on the concept that L^2 has a basis of periodic functions $\{e^{inx}\}$ in an interval $[-T/2, T/2]$, which means that:

$$\int_{-T/2}^{T/2} e^{inx} e^{imx} = T \delta_{mn} ,$$

so that we can actually write any function $f \in L^2$ as:

$$f(x) = \sum_{n=-\infty}^{+\infty} F_n e^{inx} .$$

The values of the coefficients F_n can be found as:

$$\int_{-T/2}^{T/2} dx f(x) e^{-imx} = \sum_{n=-\infty}^{+\infty} F_n \int_{-T/2}^{T/2} dx e^{inx} e^{-imx} = T F_m$$

$$\Rightarrow F_n = \frac{1}{T} \int_{-T/2}^{T/2} dx f(x) e^{inx} ,$$

where T is the periodicity of the function. F_n are the amplitudes of the waves generating $f(x)$, so in other words the contribution to f given by a specific frequency.

We now want to take the limit for the period T that goes to infinity. We rewrite the decomposition of f within T and we take the limit $T \rightarrow +\infty$. Considering the limit, we

Quantity	T -finite	T $\rightarrow +\infty$
Allowed Frequencies	$k_n = \frac{2\pi n}{T}$	$k \in \mathbb{R}$
Frequency Distance	$\Delta k = \frac{2\pi}{T}$	dk
Amplitude Coefficients	$F_n = \frac{1}{T} \int_{-T/2}^{T/2} dx f(x) e^{ik_n x}$	$F(n) = \lim_{n \rightarrow +\infty} \frac{F_n}{T}$
Function Decomposition	$f(x) = \sum_{n=-\infty}^{+\infty} F_n e^{ik_n x}$	$f(x) = \int_{-\infty}^{+\infty} dk F(k) e^{ikx}$

can finally explicitly write the amplitude coefficients:

$$\begin{aligned}
F_n &= \sum_{n=-\infty}^{+\infty} F_n e^{ik_n x} \cdot \frac{\Delta k}{\Delta k} = \sum_{n=-\infty}^{+\infty} \left(F_n \frac{T}{2\pi} \right) e^{ik_n x} \Delta k \\
&\rightarrow F(k_n) \equiv F_n \frac{T}{2\pi} = \frac{1}{T} \int_{-T/2}^{T/2} dx f(x) e^{-ik_n x} \cdot \frac{T}{2\pi} \xrightarrow{T \rightarrow +\infty} \frac{1}{T} \int_{-\infty}^{+\infty} dx f(x) e^{-ikx} \\
&\Rightarrow F(\nu) = \frac{1}{2\pi} \int_{-\infty}^{+\infty} dx f(x) e^{-i\nu x}
\end{aligned}$$

which is the **Fourier Transform** of the function f . In other words, $F(\nu)$ is the contribution of the frequency ν when generating the function $f(x)$. This is used in any field of science, from signal processing, to quantum mechanics.

7 Information Theory

Information theory aims to quantify the concept of "information" of a system configuration. Suppose to have a set of events $\{x_i\}$ each with a probability $P(x_i) = P_i$ to occur. We want to find a quantity related to x_i such that it must be *non negative*, *higher for rare events* and *additive*. We define **self-information** of x_i the quantity:

$$I(x_i) = -k \log(P(x_i))$$

and we notice that

- $I(P) \leq 0$, $\forall P \in [0, 1]$
- $I(P_1) > I(P_2)$, if $P_1 > P_2$
- $I(P_1 \cdot P_2) = I(P_1) + I(P_2)$

$I(P)$ gives information about a single element. If we have a set $\{x_i\}$ then it's natural to find the *expected information value* if a generic element of the set as:

$$E[I(P(x))] = \sum_i P_i I(P(x_i)) = -k \sum_i P(x_i) \log(P(x_i)) \equiv S(x) .$$

The quantity $S(x)$ is called **Shannon entropy** and we note that it satisfies:

- **Additivity** : $S(x, y) = E[I_x, I_y] = E[I_x] + E[I_y] = S(x) + S(y)$

- **Increasing Monotonically** : $S(\{x_i\}) \leq S(\{x_i\}^n + x_{n+1})$ (so if we add an element to the set, the total entropy increases)
- **Maximal for uniform probabilities** : $S(\{x_i\}_{\text{uniform}}) \geq S(\{x_i\}_{\text{any}})$, $P_i = \frac{1}{N}$
- **Continuous in P** : Any change in $P(x_i)$ will affect S accordingly.

Also, Shannon uniqueness theorem tells us that $S(x)$ is the only function that satisfies all the four conditions. It's useful to know that the base of the logarithm gives us the possibility to model the system information on the specific problem. For instance, in the case of PC programming, where we store data in bits (0,1), we use the base 2 logarithm so that $I(x_i) = \log_2(P(x_i))$ is the number of bytes required to store x_i . This also implies that $S_{\max}(x)$ is the maximum compression obtainable to store the sequence of elements $\{x_i\}$.

8 Game Theory

8.1 Shapley Value

Shapley value is an algorithm to estimate the proper contribution of a generic player in a specific game. The game should have a "score" function that gives a quantitative output.

$$\psi(p, \mathbf{f}) = \frac{1}{|N|!} \sum_{C_p \in P(N_p)} (\mathbf{f}(C_p \cup \{p\}) - \mathbf{f}(C_p))$$

where $\psi(p, \mathbf{f})$ is the Shapley value of the element p considering $\mathbf{f}$ as the score function of the game, $N_p = N \setminus \{p\}$ is the set of players in the game excluded p , $P(N_p)$ is its power set and C_p is an element of the power set.

This algorithm is widely used also in machine learning in order to estimate the relevance of a feature during a model prediction. It works great with tree-based models.

9 Network Theory

We define $G(V, E)$ as a graph. V is the set of nodes, E is the set of edges. The number of nodes is $|V| = N$, while the number of edges is $|E| = m$.

9.1 Types of Matrixes

Adjacency Matrix

We define adjacency matrix A the structure that identifies if a link exists:

$$A_{i,j} = \begin{cases} 1, & \text{if } (i, j) \in E \\ 0, & \text{otherwise} \end{cases}$$

For undirected graphs, $A = A^T$, which is clearly not true for directed ones. For weighted graphs, the adjacency matrix becomes W , where we simply insert the edge weight attribute instead of 0/1.

Degree Matrix

We define the degree matrix as D :

$$D_{i,j} = \begin{cases} \sum_{k=1}^N A_{i,k} , & \text{if } i=j \\ 0 , & \text{if } i \neq j \end{cases}$$

which is a diagonal matrix. The same definition considering the weighted adjacency matrix allows us to find the weighted degree matrix D_W .

Now, in the definition we wrote we slide over k , keeping i fixed. This is ok for symmetric adjacency matrixes, since $A_{i,j} = A_{j,i}$, but not for directed graphs! In fact, we expand our definition with an **in-degree** and **out-degree** matrix

$$D_{i,j}^{out} = \begin{cases} \sum_{k=1}^N A_{i,k} , & \text{if } i=j \\ 0 , & \text{if } i \neq j \end{cases}$$
$$D_{i,j}^{in} = \begin{cases} \sum_{k=1}^N A_{k,j} , & \text{if } i=j \\ 0 , & \text{if } i \neq j \end{cases}$$

These two concepts are crucial for steady state probabilities (page rank, diffusion, etc..). Usually, we use the notation ν_u to identify the degree of the node u .

Neighborhood

Given a graph $G(V, E)$, the neighborhood N of a node u is defined as

$$N(u) = \{i \in V \mid (u, i) \in E\} \cup \{u\}$$

If we remove the node itself we get the neighbors nodes of u .

Transition Matrix

For an unweighted graph, we define the transition matrix (or Markov matrix) as

$$P_{i,j} = \frac{A_{i,j}}{\sum_k A_{i,k}} = (D^{-1}A)_{i,j}$$

Similarly, for an weighted graph, we define the transition matrix as

$$P_{i,j} = \frac{W_{i,j}}{\sum_k W_{i,k}} = (D_W^{-1}W)_{i,j}$$

Again, we pay attention to the fact that we are not specifying if we are evaluating the *in* or *out* degree. Both have a well defined physical meaning, one is the probability to get to the node, one to leave the node. It's more intuitive the out-degree transition matrix, since the other has like a backward logic.

It's important to notice that the transition matrix rows (for out-degree) or columns (for in-degree) must add up to 1

$$\sum_i^N P_{i,j}^{in} = 1$$

$$\sum_j^N P_{i,j}^{out} = 1$$

This will allow us to exploit some important theorems for Markov matrices.

Path Matrix

Once we know the adjacency matrix A , we can actually take any power l of that to get the number of paths of length l between a pair of the nodes

$$A_{i,j}^l = \text{number of paths of length } l \text{ between } i \text{ and } j$$

This can be easily shown by taking the matrix power

$$(A^l)_{i,j} = \sum_{k=1}^N A_{i,k}^{l-1} \cdot A_{k,j}$$

which is equal to 0 if there is no path $i \rightarrow k$ in the graph identified by A^{l-1} and also if there is no path $k \rightarrow j$ in the graph identified by A (the starting network), otherwise it's just a sum of ones (since A is a 0/1 matrix).

Now, if we iterate this l times we clearly see that, in the end, we are tracing all the paths of length l ($u \rightarrow v \rightarrow \dots \rightarrow i \rightarrow k \rightarrow j$) possible in the network. This shows an elegant property of the adjacency matrix, with some important implications (like cycle finding on the diagonal).

It is also worth noticing that we are doing no assumption on the adjacency matrix symmetry, so this is valid for both directed and undirected networks.

Sparse Matrix Formalism

todo, formalism to use to save memory when dealing with matrices with many zeros. don't know how to use it in practice tho.

Hub

Hub is an informal way to identify an highly connected node in a network. This concept emerges frequently in peer reviews and social network analysis. It is usually used in **directed** networks to identify a node with high in-degree, without considering the out-degree.

In fact, an hub can be an editor (i.e. PubMed), an influencer (i.e. Cristiano Ronaldo), but if we define the network according to our use case, an hub can be a professor, a parent etc.. Ideally, all the nodes are hubs, that's why the definition is not rigorous, however we can determine how much a node behaves like an hub.

NOTE: In Kleinberg centrality we will introduce the concept of **authority**. For Kleinberg authority is what we mean by hub, and hub is a node that points to many good authorities, which is just stupid, so I will always invert them. Sadly, network theory is quite a new field, so it hasn't developed a properly unified nomenclature. Just the name itself, graph or network, is astonishingly stupid (cause a graph is a damn graph, and a network is waaay to many things).

9.2 Types of Networks

Here we present some of the recurrent network structures that appear in letterature and that have been studied. We always consider a generic graph $G(V, E)$.

- **Complete Graph** : All the nodes are connected to every other node (K_N).
- **Regular Graph** : All the nodes have the same degree (K_n).
- **Cycle Graph** : All the nodes have degree 2 and forms a perfect cycle.
- **Tree Graph** : Acyclic graph and has $m = N - 1$ edges (a star graph is a special case of tree).
- **Bipartite Graph** : Graph that has two different set of nodes V_1 and V_2 , each with no inner connection between the nodes. A tree is also bipartite. We can also have n -partite graphs, but usually it's only convenient the bipartite.
- **Complete Bipartite** : Graph that has two different set of nodes V_1 and V_2 , each with no inner connection between the nodes and every node of set V_1 is connected to all the nodes of V_2 and viceversa.
- **Lattice Graph** : Network that can be arranged in a regular grid, so that forms a regular tiling if embedded into an euclidean space.

Now we present *toy networks* that are usually used to randomly generate graphs. There are others, but a these two are the only ones that can resable something useful in the real world.

- **Erdős–Rényi Graph** : Random graph, each node with a probability p to have a link with any other.
- **Barabási–Albert** : Iterative graph, that exhibits power law behavior. Each time, add a new node u with a probability $p_v = \nu_v / \sum \nu_j$ to be connected with the node v .

9.3 Centralities

Centrality measures are the most useful method to get both local and globa information regarding nodes and its neighbors. There are many of them out there, all serving different purposes. In general, networks are very complex structures and we may don't have a general recipe to complete a task, or extracting specific information. It's usually a trade-off between accuracy required and performance costs. All the following discussions will consider a network $G(V, E)$, with N number of nodes and m number of edges.

Degree Centrality

Degree centrality is by far the easiest yet surprisingly effective centrality there is. It is defined as

$$\vec{C}_d = \frac{1}{N-1} D \vec{1}$$

where D is the unweighted degree matrix and $\vec{1}$ is the unit vector. It can be defined both for D^{out} and D^{in} , and it shows how much a node is connected in the network.

If we consider the weighted degree matrix D_W , we get the *strength centrality*, which combined with the degree centrality can give us an insight on the most effective nodes existing in the network, or precious hints on finding the potential ones (i.e. a node with many weak connections, if improved, can become crucial).

Betweenness Centrality

Betweenness centrality is a quantity used to determine the most crucial nodes for the network connectivity, in other words, those nodes that are visited by the most number of shortest paths. The definition of betweenness centrality C_b for a node u

$$C_b(u) = \sum_{v \neq u \neq w} \frac{N_{sp}(u, w|u)}{N_{sp}(u, w)}$$

where $N_{sp}(a, b)$ gets the number of shortest paths between the nodes a and b , and $N_{sp}(a, b|c)$ requires those paths to pass from c . The complexity is clearly high, in fact it scales as $O(N^3)$ for a naive solution, and $O(N \cdot m)$ in case of unweighted graphs, or $O(N \cdot m + N^2 \cdot \log N)$ for positive weights. Usually, we use approximated methods, mostly sampling based methods.

Closeness Centrality

Closeness centrality quantifies how near a node is to all other nodes in a network. It reflects how efficiently information can spread from that node to others and can, for instance, support recommendation systems based on the proximity of nodes' interests. It is defined as:

$$C_c(u) = \frac{N - 1}{\sum_{v \neq u} d(v, u)}$$

where $d(v, u)$ is the length of the shortest path between v and u (the geodesic distance). It's important to notice that, for directed networks, $d(u, v) \neq d(v, u)$, so we have to keep in mind what we are asking. This centrality is defined only for connected graphs, since $d(u, v) = +\infty$ for disconnected components.

Harmonic Centrality

Harmonic centrality also measures how close a node is to all others but provides a more robust formulation for disconnected networks. It is defined as:

$$C_c(u) = \sum_{v \neq u} \frac{1}{d(v, u)}$$

where $d(v, u)$ is the length of the shortest path between v and u (the geodesic distance). Again, notice that for directed networks, $d(u, v) \neq d(v, u)$. This centrality addresses the problem of closeness centrality for unconnected graphs, where $d(u, v) = +\infty$ would get all to zero. The use cases are basically the same as the closeness (in reality, there is no use to use closeness, just use this)

Eigenvector Centrality

This centrality is a bit tricky, so let's start by recalling the eigenvector problem, in terms of adjacency matrix, and $\vec{x}$ being an eigenvector

$$A\vec{x} = \lambda\vec{x}$$

Now let's recall a few theorems. First, **Perron-Frobenius theorem** assures us that, if A is a real square matrix, then the eigenvector associated to the largest eigenvalue has all positive components. This is interesting if we want to exploit this for a centrality measure (it should be a positive quantity for physical interpretation).

So, let's assume that A is an adjacency matrix for a weighted graph with non negative weights. At this point, we can evaluate the largest eigenvector solving the eigenvalue problem

$$A\vec{x}_L = \lambda_L\vec{x}_L$$

and we can rewrite this emphasizing the i -th component of the largest eigenvector $\vec{x}_L$ as

$$x_{L,i} = \frac{1}{\lambda} \sum_j A_{i,j} x_{L,j}$$

By definition, we evaluated the eigenvector centrality measure of the i -th node. Now, we can give it a physical meaning as follows: The eigenvector centrality quantify how much a node is important, and how important its neighborhood is too, which affects the total score. We can then rename for consistency

$$\vec{x} = \vec{C}_e$$

This centrality is quite easy to compute, and it tells particularly well the *rank* system of the nodes in the network. In particular, it is used for *undirected* networks.

Kleinberg Centrality

Kleinberg centrality tackles the problem of recognizing the importance of hubs and authorities in directed graphs. As we introduced earlier, *hubs* are nodes that have high in-degree connectivity, but we can improve this concept by quantifying the "quality" of these connections. Here we introduce the concept of *authority*, which are nodes that points to many "good" hubs. We define hubs $\vec{h}$ and authorities $\vec{r}$ as:

$$\vec{h} = A^T \vec{r}$$

$$\vec{r} = A \vec{h}$$

which in the case of an *undirected* network we have that $A^T = A$, therefore it makes no sense considering $\vec{h} \neq \vec{r}$, thus we can impose $\vec{h} = \vec{r}$ and we proceed with the traditional eigenvalue problem.

$$\vec{x} = A\vec{x}$$

In this context, *hubs* are nodes that are pointed by many good authorities, while *authorities* are nodes that points many good *hubs*. In other words, authorities build bridges between good hubs.

This algorithm is used when dealing with *sub-networks* or query-dependent networks, like topic related graphs.

NOTE: To keep consistency with the initial definition of hubs, I inverted Kleinberg definition of hubs and authorities. So, whenever you find this algorithm, remember to switch them.

PageRank Centrality

PageRank algorithm is the eigenvector centrality with the addition of a random jumping factor. We have to keep in mind that we are studying a graph $G(V, E)$ with N nodes and m edges. The network G is described by the weighted adjacency matrix W .

We don't start from the eigenvalue problem, but we start by setting up the node presence probability equation. We define the page rank equation:

$$\vec{C}_{pr} = \alpha P^{in} \vec{C}_{pr} + (1 - \alpha) \vec{v}$$

which states exactly: *Given any node i , node i PageRank score depends on which other nodes point to i^* , and we find it as a steady-state probability. We use the steady state probability because we want to study the equilibrium state, otherwise $\vec{C}_{pr} \in \mathbb{R}^N$ would be different from left and right.

The stochastic vector $\vec{v} \in \mathbb{R}^N$ is exactly the probability of jumping to a node in the network. In fact, $(1 - \alpha)$ is the probability to jump, but where to jump is given by $\vec{v}$. Usually, we simply define the stochastic vector as

$$\vec{v}_i = \frac{1}{N}$$

We recall the definition of the *in-degrees* stochastic matrix $P \in (\mathbb{R}^N \times \mathbb{R}^N)$ (we will drop the "in" later):

$$P_{i,j}^{in} = \frac{1}{\sum_k A_{k,j}} = \frac{W_{i,j}}{\sum_k W_{k,j}}$$

Each column adds up to 1, which is fundamental for *Perron-Frobenius theorem* in the case of normalized stochastic matrixes.

With all of this being said, using the fact that $\vec{1}^\top \vec{C}_{pr} = 1$, we can manipulate the page rank equation so that we can write:

$$\begin{aligned} \vec{C}_{pr} &= \alpha P \vec{C}_{pr} + (1 - \alpha) \vec{v} (\vec{1}^\top \vec{C}_{pr}) = \\ &= [\alpha P + (1 - \alpha) \vec{v} \vec{1}^\top] \vec{C}_{pr} \end{aligned}$$

Thus we get the Google Matrix

$$G = \alpha P + (1 - \alpha) \vec{v} \vec{1}^\top$$

with which we can set the eigenvalue problem

$$G \vec{C}_{pr} = \vec{C}_{pr}$$

where *Perron-Frobenius theorem* assures us that for a stochastic matrix, the largest eigenvalue is $\lambda = 1$, precisely our case.

From here on, it's just the eigenvector centrality evaluation. PageRank is used for ranking the pages importance. We must note that W is not required to be symmetric, just stochastic. This means that PageRank algorithm is perfect for *directed networks*. In particular, the random jump factor allows to manage also the *dead-end* nodes (no outgoing link), that cannot be treated by the simple eigenvector centrality.

Katz Centrality

Katz centrality consider the relevance of a node according to the total number of walks that pass through a given node. The definition is straight forward:

$$C_K(u) = \sum_{k=1}^{+\infty} \sum_{i=1}^N \beta^k (A^k)_{i,u}$$

which finds all the possible paths to u of all possible lengths. The value β is called attenuator factor, and is there to grant us the convergence of the series. In fact, it cannot be any number. For all the nodes we can use the unit vector $\vec{1}$:

$$\vec{C}_K = \sum_{k=1}^{+\infty} \beta^k (A^k)^\top \vec{1} = \sum_{k=1}^{+\infty} (\beta A^\top)^k \vec{1}$$

and at this point we recall Neumann Matrix expansion:

$$(I - A)^{-1} = \sum_{n=0}^{+\infty} A^n = I + \sum_{n=1}^{+\infty} A^n$$

with the spectral radius $\rho(A) < 1$ (the module of the maximum eigenvalue). We can rewrite Katz centrality as

$$\vec{C}_K = ((I - \beta A^\top)^{-1} - I) \vec{1}$$

with the requirement that

$$\beta < \frac{1}{\lambda_{max}}$$

in order for the series to converge.

9.4 Diffusion on Graphs (analytical)

The study of how a quantity will go through a structure is particularly interesting in many situations. This because it allows to predict the behavior of certain phenomena, like money distribution or power shortages.

Any diffusion process follows the diffusion equation

$$\frac{\partial}{\partial t} u(\vec{x}, t) = \nabla^2 (D(\vec{x}, t) \cdot u(\vec{x}, t))$$

which can be easily obtained by considering the continuity equation and the Fick's law

$$\frac{\partial}{\partial t} u(\vec{x}, t) + \nabla J(\vec{x}, t) = 0$$

$$J(\vec{x}, t) = -\nabla (D(\vec{x}, t) \cdot u(\vec{x}, t))$$

In a graph context, $\vec{u}$ are the quantities present in each node, and the diffusion equation can be written as

$$\frac{d}{dt} \vec{u}(t) = -L_D \vec{u}(t)$$

obtained as:

$$\begin{aligned}
\frac{d}{dt}u_i(t) &= \sum_{j=1}^N W_{i,j}(u_j(t) - u_i(t)) = \\
&= \sum_{j=1}^N W_{i,j}u_j - u_i \sum_{j=1}^N W_{i,j} = \\
&= \sum_{j=1}^N W_{i,j}u_j - u_i D_i = \sum_{j=1}^N W_{i,j}u_j - \sum_{j=1}^N \delta_{i,j}u_i D_i = \\
&= \sum_{j=1}^N u_j(W_{i,j} - \delta_{i,j}D_i)
\end{aligned}$$

thus we can consider all the nodes at the same time, obtaining the diffusion equation

$$\frac{d}{dt}\vec{u}(t) = (W - D)\vec{u}(t) = -L_D\vec{u}(t)$$

Note that here we explicitly consider the fact that $\vec{u}$ is a vector, since the concept of position in a network is different from the continuum space. Here each node is a single point, so we consider all at the same time.

The new matrix found L_D is called **Laplacian** matrix:

$$L_D = D - W$$

This reasoning is true for both *directed* and *undirected* graphs, with a very important difference. We have considered D at the node i , but without specifying if that is an "in-degree" or "out-degree" matrix. According to what we want to diffuse, we will need to specify it for directed networks.

9.5 Experimental Diffusion

In general, if we can't evaluate the Laplacian etc.. we do have **random walks** and **transition matrixes**. Random walks allow to simulate a diffusion process, according if we deploy a proper amount of independent walkers (probably central limit theorem or something like that). This is useful cause we can isolate a specific portion of the graph and perform many random walk simulations in order to monitor how the quantities changes. For Markov Ergodic Theorem (i think) we know that simulating infinite random walks on a given network, it will reproduce the steady state distribution.

Transition matrixes on the other hand provide a statistical description of where the quantities will go, according to the connectivity of the nodes. Both in the limit of many steps, will end up in the Laplacian case. The transition matrix, as we saw, is defined as

$$P_{i,j} = \frac{W_{i,j}}{\sum_k W_{i,k}} = (D_W^{-1}W)_{i,j}$$

again considering the incoming or outgoing links. We can use the transition matrix to perform step by step the diffusion of any quantity in a network by simply evaluating:

$$\vec{x}(t+1) = P\vec{x}(t)$$

which will eventually converge to the steady state distribution.

Cover Time

We define cover time τ as the expected number of steps required for a random walk to visit all the nodes in a network, starting from a random node u :

$$\tau = E[n_{steps}] = \lim_{n \rightarrow +\infty} \frac{\sum_{i=0}^n \tau_i}{n}$$

Hitting Time

We define hitting time $\vec{\tau}_u$ the expected number of steps required for a random walk to reach each node v in the network, starting from the node u :

$$\tau_{u,v} = E[n_{steps,i}|u] = \lim_{n \rightarrow +\infty} \frac{\sum_{i=0}^n n_i(v|u)}{n}$$

Note that $\tau_{u,v} \neq \tau_{v,u}$ even for undirected graph, in fact if we define the hitting time matrix τ it won't be symmetric unless the network is **regular** (all nodes have the same degree).

We can define the **commute time** as $C_{\tau_{u,v}} = \tau_{u,v} + \tau_{v,u}$ so that the commute matrix C_τ is actually symmetric.

9.6 Path Finding

Path finding is a fundamental topic in network theory and it is used in many fields, from mathematical modeling to video games characters behaviors. There are different algorithms, but A* is surely the most used.

Breath First Search (BFS)

Simple algorithm for graph traversal, in particular is used to find a path from a node u to a node v . We don't find shortest paths or stuff, just a path, depending on how you queue the nodes.

The idea is very simple: We define 2 structures, the **queue nodes** and the **visited nodes**, which are two different set of nodes. The *visited nodes* start empty, while the *queue nodes* start with the first node we want to visit.

- Start from node $u \rightarrow$ remove it from *queue*, add it to *visited*.
- Get the neighborhood $N(u)$ and insert every node in that set that is not in *visited* inside *queue*.
- Proceed with the next node w in *queue* $\rightarrow$ remove it from *queue*, add it to *visited*.
- Repeat 2 and 3.
- Stop when you find the required node v .

Dijkstra's Algorithm

Algorithm used to find the shortest path in a weighted graph. It adapts the BFS algorithm by providing the concept of **priority queue**, that selects the following node according to the minimum of distance from the starting one. We need some structures:

- **priority queue nodes** : where you select the node with smallest distance from the starting one. This is needed to get the shortest path length.
- **visited nodes** : set of visited nodes, and that will never be analyzed again. This is needed to avoid loops.
- **previous nodes** : map between any node and its previous. The previous is chosen according to the smallest distance found (it will be update during the algorithm). This is needed to reconstruct the path.

We initialize all the distances to any node (except the starting one) with $+\infty$. The idea is:

1. We always visit the node in the priority queue. If a node is given by the priority queue, that node goes into the **visited nodes** and it will never be visited again. This assures that we reached the node in the smallest path possible, because of how the priority queue is constructed (prioritizes always the step with smallest distance, independently on which node we are).
2. Given a node i we evaluate the distance to all the neighbors nodes j so that we can get the distance from the starting node $d(u, j) = d(u, i) + W_{i,j}$. If $d(u, j)$ is smaller than the previous distance $d_{prev}(u, j)$ (recall that we initialized it to $+\infty$) then we update it.
3. We also assign the previous node to j , as the one with the smallest distance $d(i, j)$.

Once the priority queue gives us the target node v , then we stop. To reconstruct the path, we simply use the *previous nodes* map, starting from the *target node*, and arriving the starting one, ensuring the shortest path.

A* Algorithm

A* is the algorithm mostly used for path search, in videogames, gps, and many other fields. It is based on Dijkstra Algorithm, but it adds an *"educated"* estimation on how distant the target node is, so that it manages to avoid searching for evidently off paths (imagine going to a place, but moving in the opposite direction, it's pointless).

This works well if we are in a metric space in which the nodes are located, it's harder if we don't have a correspondence between nodes and metric place. In fact, GPS is the perfect example, since nodes are cities, and they are mapped to a specific point in the metric space.

We still initialize all the nodes distances with $+\infty$, but during the priority queue evaluation we also consider an additional term, the **heuristic distance**, from the incoming node and target node, exactly as a potential function. Assuming that u is the starting node and v is the target node, the priority queue will process a distance for the incoming node j :

$$f(j) = h(j, v) + d(u, j)$$

where $d(u, j)$ is the shortest distance from u to j .

There are many ways to define $h(x, y)$, usually we can use the *"euclidean distance"* if we are in $\mathbb{R}^N$, otherwise the *"arc"* distance if we are on an hypersphere.

9.7 Link Prediction

We can exploit similarities in order to predict nodes connections according to some pre-defined threshold. Clearly, each similarity express some specific node relation, the weights affects these relations too, so every case has its own peculiarities that must be taken in considerations.

Jaccard Similarity

Jaccard similarity is the fraction of elements in common between two neighborhoods, defined as:

$$S_J(u, v) = \frac{|N(u) \cap N(v)|}{|N(u) \cup N(v)|} \in [0, 1]$$

This gives a straightforward representation of similarity between two nodes, without considering any second order relations.

Adamic-Adar Similarity

Adamic-Adar is a refinement of Jaccard similarity. It quantifies the amount of elements somewhat in common between two nodes as:

$$S_{AA}(u, v) = \sum_{i \in (N(u) \cap N(v)) \setminus \{u, v\}} \frac{1}{\log(D_W(i))} \geq 0$$

with $D_W(i)$ being the value of the weighted degree matrix for the node i . Again we notice that D_W can represent both *in* or *out* degrees.

Random walk Similarity

Random walk similiarity is a cool concept that can be obtained through simulations too. Given a node u in a graph $G(V, E)$ we define random walk similiarity vector:

$$\vec{S}_R(u) = \vec{\tau}(u)^{-1}$$

with $\vec{\tau}(u)$ being the **hitting time** vector from u to all the nodes in the network. The hitting time is defined as the average number of steps required to get from a starting node u to a target node v .

10 Markov Processes

A Markov process is a stochastical sequence of events such that:

$$P(x_{n+1} = x | x_n, x_{n-1}, \dots, x_0) = P(x_{n+1} | x_n) ,$$

which means that the probability to have a new state x_{n+1} depends only on the current state in which my system is x_n . If the sequence of events $\{x_i\}$ is finite, we call it **Markov chain**, if instead $x \in \mathbb{R}$ then it's called **Brownian motion**.

We describe the Markov process using the **Transition matrix** T such that:

$$T : T_{i,j} = P(x_{n+1} = j | x_n = i) , \sum_j T_{i,j} = 1 ,$$

so we are asking T to be *row-stochastic*. Given a generic system, containing many nodes with some quantities, the evolution of the system at a step t is:

$$\vec{s}_t = \vec{s}_0 T^t ,$$

where $t \in \mathbb{N}$ and $\vec{s}_0$ is the initial distribution of a generic quantity in a system. Clearly, the stationary distribution is obtained when nothing changes in the system when the transition matrix is applied:

$$\vec{s} = \vec{s} T$$

11 Time Series

Data can be collected in many different ways. Time series are a way to collect data by emphasizing the concept of "time" in which the data has been collected. This produces a list of peculiarities and difficulties that are a bit different from generic (x, y) points.

In broader terms, a time series is simply a collection of multidimensional points $\{(\vec{p}, t_i)\}_i$ where the order of acquisition t_i must be considered. This usually means that any point $\vec{p}_i$ depends in some way on the previous points, so $\{\vec{p}_i\}_i$ are **not independent**.

A **stationary time series**, or stationary process, is a series that do not change its statistical variables if shifted in times, meaning that mean, variance, behaviors, remain always constant. In real world application, time series are never stationary, but they can be useful as null hypothesis.

11.1 Theoretical Description

A time series is basically a Markov process $P(x_{n+1} = x | x_n, x_{n-1}, \dots, x_0) = P(x_{n+1} | x_n, \dots, x_{n-m})$ of order m , where in general the effect of further terms gets smaller. The value x_t is called *process*, and the set $\{x_t\}_0^n$ represents the steps taken in the process, or *path*.

The **expected value** of the process is:

$$\mu_t = E[x_t] = \int_{-\infty}^{+\infty} z f_t(z) dz ,$$

where $f_t(z)$ is the probability density distribution of x_t , z is the set of possible values that x_t can take. For a *stationary process* we have that $\mu_t = \mu$, $\forall t$.

The **variance** of the process is:

$$\sigma_{x,t} = \text{Var}[x_t] = E[(x_t - \mu_t)^2] ,$$

and again for a *stationary process* we have $\sigma_t = \sigma$, $\forall t$.

One very important quantity is the **autocorrelation**, because it allows to understand the influence of previous points on the newer ones:

$$R_x(t, j) = \frac{\text{Cov}[x_t, x_{t-j}]}{\sqrt{\text{Var}[x_t] \text{Var}[x_{t-j}]}} = \frac{E[(x_t - \mu_t)(x_{t-j} - \mu_{t-j})]}{\sqrt{\sigma_{x,t} \sigma_{x,t-j}}}$$

We understand that *white noise* $\{\varepsilon_t\}_t$ will contribute to these quantities as:

$$\begin{aligned} E[\varepsilon_t] &= 0 \\ \sigma_{\varepsilon,t} &= \sigma_\varepsilon \\ R_\varepsilon(t, j) &= 0, \forall t \neq j \end{aligned}$$

If a stationary time series has white noise, it will lose the stationarity properties, since:

$$\sigma_{x,t} = t\sigma^2 ,$$

so it will grow with time.

11.2 Technical Difficulties

The timestamps are usually in a format that is not easily manageable by programming languages. In fact, the timestamp dataformat is not quite supported, we usually need to convert it to integers, or floats, while keeping all the spacing between the points unchanged.

11.3 Algorithms

Here we list a series of algorithm used in time series analysis.

Moving Average

Moving average is an algorithm to smooth the curve according to a moving windows, defined as:

$$M_k(x') = \frac{1}{k} \sum_{i=n-k+1}^n x_i$$

so we understand that we need to sacrifice $k - 1$ points from the original curve, in order to create a more stable function. Normally, the window k is an odd number below 10, but it really depends on the type of data acquired.

ARIMA

ARIMA models (AutoRegressive Integrated Moving Average) are a family of models that takes in consideration the autocorrelation terms ..

12 Likelihood Function

The Likelihood function is a fundamental concept in statistics that flips the perspective of probability of a random variable given some parameters, to the probability of having some parameters given some specific observations of the random variable.

12.1 Probability Functions

A probability function of a generic process describes how likely is to observe any event among the possible ones. It's crucial to distinguish between *discrete* and *continuous* probability functions.

A **discrete probability function** describes a process that can assume only a finite or countable number of values. The probability function is called **probability mass function** and it is defined as:

$$P_\chi(x) = P(\chi = x) \tag{9}$$

so it is the probability for the random variable χ to assume exactly the value x . It must satisfy the following properties:

- $P_\chi(x) \geq 0 \forall x$
- $\sum_x P_\chi(x) = 1$
- $P(A) = \sum_{x \in A} P_\chi(x)$

For a **continuous probability function** instead it doesn't make sense to ask how much likely is to observe a specific value, since we are dealing with a continuous random variable. For this reason we define the **probability density function** as:

$$\rho_\chi(x) : P(a \leq \chi \leq b) = \int_a^b \rho_\chi(x) dx \quad (10)$$

where $P(\chi)$ is the cumulative distribution function. Again we require:

- $\rho_\chi(x) \geq 0 \forall x$
- $\int_{-\infty}^{+\infty} \rho_\chi(x) dx = 1$
- $P(a \leq \chi \leq b) = \int_a^b \rho_\chi(x) dx$

12.2 Maximum Likelihood Estimation

Given some random variable $\chi \sim f_\chi(x|\vec{\theta})$ where $\vec{\theta}$ are eventual fixed parameters, assuming the observations are independent and identically distributed we define the **likelihood** as:

$$\mathcal{L}(\vec{\theta}|\vec{x}) = \prod_{i=1}^N f_\chi(x_i|\vec{\theta}), \quad (11)$$

which corresponds to the joint probability density (or mass, for discrete variables) of observed data $\vec{x} = \{x_1, x_2, \dots, x_N\}$ as a function of the parameters $\vec{\theta}$. Now this definition makes perfect sense for *discrete* random variables, but also for *continuous* ones, since we can consider the joint probability densities instead of the probabilities themselves.

We defined $\mathcal{L}(\vec{\theta}|\vec{x})$ so that if we maximize it around the observed data $\vec{x}$, we would get the parameters $vec\theta$ that better explain the data. This method is called **Maximum Likelihood Estimation**; for the sake of calculations usually we consider the **log-likelihood** instead:

$$\ln \mathcal{L}(\vec{\theta}|\vec{x}) = \sum_{i=1}^N \ln f_\chi(x_i|\vec{\theta}) \quad (12)$$

and we maximize it with respect to $\vec{\theta}$:

$$\begin{aligned} \hat{\vec{\theta}} &= \arg \max_{\vec{\theta}} \ln \mathcal{L}(\vec{\theta}|\vec{x}) \\ \Rightarrow \left(\frac{d\mathcal{L}(\vec{\theta}|\vec{x})}{d\vec{\theta}} \right)_{\vec{\theta}=\hat{\vec{\theta}}} &= \vec{0}. \end{aligned}$$

13 P-Value

The p-value is a fundamental tool used to quantify how much something is likely to be true. We start from a statistical assumption, called **null hypothesis** H_0 , and we want to estimate how much H_0 fits the observed data. The formal definition is "*Assuming that H_0 is true, the p-value is the probability of obtaining a test statistic as extreme as at least the one measured from the experimental data*":

$$p \equiv P(t > t_{obs} | H_0), \quad t \sim f(t).$$

Now we need to define H_0 and the test statistic t .

- H_0 : My data comes from the distribution $f(\vec{\theta})$, or "my data has mean in μ ", etc..
- **Test Statistics** : Quantity strictly related to the null hypothesis, and measured from the observed data. It should be tailored on the null hypothesis and on what we want to achieve.

Each test statistics has its particular scope in mind. Here we list few of them.

- **t-test** : Tests if the data has average in μ_0 . Variance unknown, small dataset.

$$t = \frac{\bar{x} - \mu_0}{s/\sqrt{N}}, \quad s^2 = \frac{\sum^N (x_i - \bar{x})^2}{N - 1}.$$

- **Pearson's Chi-squared** : Tests the goodness of fit. $\hat{x}_i$ predicted, x_i observed.

$$\chi^2 = \sum_{i=1}^N \frac{(x_i - \hat{x}_i)^2}{\hat{x}_i}.$$

- **F-test (ANOVA)** : Compare the variance between two groups. Used to see if two groups are different or not.

$$f = \frac{s_1^2}{s_2^2}.$$

In ANOVA notation, s_1^2 is the within group variance, s_2^2 is the between groups variance.

the distribution of p varies depending on if H_0 is true or not. In fact, if H_0 is true, p must be distributed uniformly:

- The test statistics follows a distribution $f(t)$
- Sampling from any cumulative distribution means that any percentile has the same probability to occur, since it is the interval $[0, 1]$.
- If we define the cumulative of t as $F_{H_0}(T \leq t | H_0) = P(t)$, then $p \sim P(t)$ must be uniform.

14 Monte Carlo Simulation

15 Quantum Mechanics

Review of the fundamental concepts and result of quantum mechanics. Quantum mechanics is based on wave functions and operators, so we will introduce the main mathematical tools to describe them and we will follow by defining the physics that uses them.

15.1 Hilber Space

We define Hilbert Space H a space with the following properties:

1. H is a vector space
2. H has inner product $\langle f, g \rangle$
3. Norm $\|f\| = \sqrt{\langle f, f \rangle}$
4. H is complete, so every Cauchy sequence converges in the space H .

In practice, we will always consider $H \sim L^2(a, b)$, where $L^2(a, b) = \{f : \int_a^b |f(x)|^2 dx < +\infty\}$ such that:

1. $L^2(a, b)$ is a vector space
2. Inner product $\langle f, g \rangle = \int_a^b \overline{f(x)} g(x) dx$
3. Norm $\|f\| = \langle f, f \rangle = \int_a^b |f(x)|^2 dx$
4. Complete with respect to the norm $\|f\|_2 = \sqrt{\int_a^b |f(x)|^2 dx}$.

This is useful for wave functions, since in quantum mechanics the squared wave function is a probability density function, so the wave functions live in an Hilbert space.

15.2 Nomenclature

The concepts expressed in quantum mechanics are mostly the same of linear algebra. However, since the elements of the vector space are usually functions (wave functions), they live in an infinite dimensional space. Considering Fourier decomposition, we know that we can express any function as an infinite sum of waves:

$$f(x) = \sum_{n=0}^{+\infty} F_n e^{inx} .$$

Here n is an integer that, in some sense, limits the number of allowed wavelengths according to their spacing. In the limit of the spacing going to zero, the same function can be expressed as an integral:

$$f(x) = \int_{-\infty}^{+\infty} F(\nu) e^{i\nu x} d\nu ,$$

with $\nu \in \mathbb{R}$ being the wave frequencies that are now continuous. As we will see, these two cases represents in quantum mechanics a confined observable that becomes discretized

due to the dependency on the wavelength (like the energy of a particle), and an observable that is not affected by the quantization (like the position of a particle) respectively.

With this being said, the generators of the vector space can be either a numberable set of basis vectors (still infinite, but countable), or an uncountable set of functions that span the space via integrals. For this reason, a new nomenclature is used to pay proper attention to this fact, and to simplify the representation of some recurrent calculations.

- **column vector** : $\vec{v} = \begin{pmatrix} v_0 \\ \vdots \\ v_N \end{pmatrix} \xrightarrow{qm} |v\rangle$
- **row conjugate vector** : $\vec{v}^\top = (v_0^*, \dots, v_N^*) \xrightarrow{qm} \langle v|$
- **inner product** : $\langle v, u \rangle = \sum_{i=0}^N v_i^* u_i \xrightarrow{qm} \langle v|u \rangle = \int_a^b \overline{v(x)} u(x) dx$
- **matrix** : $A \xrightarrow{qm} \hat{A}$. Matrices are also called *operators*, since they are usually applied to functions.
- **Matrix-Vector multiplication** $\vec{u} = A\vec{v} \xrightarrow{qm} |u\rangle = \hat{A}|v\rangle$
- **Quadratic form** $\vec{u}^\top A\vec{v} \xrightarrow{qm} \langle u|\hat{A}|v \rangle$. The result is a scalar.

Since L^2 is a vector space, we can find an orthonormal basis $\{f_n(x)\}$ such that:

$$f(x) = \sum_{n=0}^{+\infty} F_n f_n(x) \quad (13)$$

whose coefficients F_n can be found exploiting the orthonormality:

$$\begin{aligned} \langle f_n(x) | f_m(x) \rangle &= \delta_{mn} \\ \Rightarrow F_n &= \langle f_n(x) | f(x) \rangle \quad (\text{Fourier method}) \end{aligned}$$

where we recall $f_n(x), f(x)$ are vectors in Hilbert space.

15.3 State Vectors, Basis and Descriptions

Once $L^2(a, b)$ and Hilbert spaces are defined, we ask what a vector in such space is. We call the *normalized* vector $|\Psi(t)\rangle \in H$ a **quantum state**, whose components depend on a variable t – usually the time. All the information about the physics of the system is embedded inside the quantum state. The reason why we want it to be normalized is related to **Born's rule** that we will see later.

Now, what is a quantum state $|\Psi\rangle$ exactly? It is an abstract object that lives in Hilbert space and represents a generic configuration of the particle. We need some basis of the space that has physical properties so that it can describe the state with concrete implications (energy, position, momentum, ...).

In particular, once we fix a vector basis $\{|q_i\rangle\}$ we are able to rewrite any possible state Ψ as a linear combination of that base – or superposition in this context, since we are dealing with probability waves.

The idea is simple: find a basis that represents something physical, then write the state of the particle according to that physical quantity. This is possible because we are in L^2 and so our states can be seen as wave functions.

Now we want to define this basis $\{|q_i\rangle\}$. Since we want real quantities, we know that **Hermitian** operators have real eigenvalues. This is useful, because now we can define the Hermitian operator $\hat{Q}$ such that:

$$\hat{Q}|q\rangle = q|q\rangle \quad (14)$$

where we ask that if a system is in a state $|q\rangle$, then a measurement of the physical quantity Q will yield the value q with probability 1. We must be smart to define $\hat{Q}$ such that it actually represents physical quantities and that they would eventually recover the classical counterpart in the classical limit.

If we manage to define $\hat{Q}$, then to get $\{|q_i\rangle\}$, we can write any state in Hilbert space as:

$$|\Psi\rangle = \sum_{n=0}^{+\infty} c_n |q_n\rangle \quad (15)$$

Here we have to pause for a second to highlight a crucial point about Hilbert space and basis. A quantum state $|\Psi\rangle$ can be given by a linear combination of *numerable* vectors $\{|q_i\rangle\}$. However, since we are in L^2 , we can also rewrite any vector $|\Psi\rangle$ as a decomposition of functions using continuous integrals, exactly as Fourier decomposition:

$$f(x) = \sum_{n=-\infty}^{+\infty} F_n e^{inx}$$

$$f(x) = \int_{-\infty}^{+\infty} F(l) e^{ilx} dl$$

Depending on the physical situation we are in, we might end up in a situation, in another, or even in both. Some quantities in quantum mechanics depend on the **wavelength** of the particle, like the momentum and the energy. Others do not, like the position of the particle. If we consider a particle in a box, so in a potential well, it will have a boundary condition $\Psi(0) = \Psi(L) = 0$. This condition means that only some wavelengths are permitted, thus only some energies. The position is also limited within 0 and L , but in a continuous way. This means that not all observables have basis that lives in the Hilbert space! This doesn't affect that much in terms of discussion, but it is a profound property of L^2 spaces. The set of eigenvectors of an operator generate the **spectrum** of the operator. We will refer to **discrete spectrum** when the basis is numerable, and **continuous spectrum** when our state must be expressed as an integral – i.e. when the quantity depends on the wavelength or not.

Now, as we do in physics, we want to predict what could be the outcome of a real measure of the quantity Q . For the discrete spectrum, we know that the state is given as a superposition of vector basis, each representing a possible outcome of the measure by definition. So it's reasonable to assume that the projection of the state $|\Psi\rangle$ on the basis vector $|q_n\rangle$ gives us the probability amplitude to measure the value q_n when measuring Q . From this assumption, and from the fact that the basis is *orthonormal*, we can obtain one of the most important rules of quantum mechanics: **Born's rule**, which quantifies

the probability of measuring a specific value among the available ones given by the basis as:

$$P(Q = q_i) = |\langle q_i | \Psi \rangle|^2 = \frac{|c_i|^2}{|k|^2} = |c_i|^2 \quad (16)$$

with k being the length of the state vector, that we asked to be normalized exactly for this reason.

For the continuous spectrum, the reasoning is the same, but as we did for Fourier decomposition, at this point we need to work with a continuous function of coefficients $\{c_n\} \rightarrow \Psi(q)$ such that:

$$\begin{aligned} \Psi(q) &= \langle q | \Psi \rangle \\ \rightarrow |\Psi\rangle &= \int_a^b \Psi(q) |q\rangle dq \end{aligned}$$

and so the probability to measure the quantity Q with value between a range of values $[q_0, q_1]$ is:

$$\begin{aligned} \rho(q) &= |\Psi(q)|^2 \\ \rightarrow P(q_0 \leq Q \leq q_1) &= \int_{q_0}^{q_1} \rho(q) dq \end{aligned}$$

where $\rho(q)$ is the probability density function to measure a physical quantity of value q . Here we also understand that the wave function valued for a certain basis q will produce a scalar (or a complex number):

$$\psi : \mathcal{Q} \longrightarrow \mathbb{C}, \quad q \longmapsto \psi(q) = \langle q | \Psi \rangle.$$

This is exactly the same thing that happens in Fourier analysis: we can rewrite any function $f(x)$ as a superposition of different waves with different amplitudes, which in the limit of the distance between two consecutive wavelengths going to zero, it becomes an integral over all possible wavelengths. So we rewrite the function $f(x)$ in terms of a new continuous function $F(\nu)$, with ν being a variable in a continuous domain (**frequency domain**); in quantum mechanics we rewrite our state Ψ in terms of a new continuous function $\Psi(q)$, where q is a continuous variable defined in a specific physical space ($q = x \rightarrow$ **space domain**, $q = p \rightarrow$ **momentum domain**, ...).

Note that each wave function Ψ_p denoting the state of particle p lives in a Hilbert space H_p . If we consider two non-interacting particles x and y , the joint state will be described as:

$$|\Psi_{x,y}\rangle = |\Psi_x\rangle \otimes |\Psi_y\rangle$$

which lives in a new Hilbert space $H_x \otimes H_y$ and if we fix a basis q we do get, as we will see later:

$$\Psi_{x,y}(q_x, q_y) = \Psi_x(q_x) \cdot \Psi_y(q_y) .$$

It is required that the particles do not interact, so that the Hamiltonian is separable, which is rarely happening due to EM interactions. Also pay attention to the fact that we are doing tensor product, not cross products – otherwise we wouldn't be able to reproduce all the spaces!

So to summarize:

- A Hermitian operator $\hat{Q}$ is chosen to be associated to a real observable Q .

- According to the eigenvector basis we use to represent our quantum state $|\Psi\rangle$ we give a specific description of the state (position, momentum, energy, ...).
- The quantum state $|\Psi\rangle$ can be written as a linear combination of the eigenvectors $|q_i\rangle$ of $\hat{Q}$. Measurements of Q yield the eigenvalues associated with these eigenvectors, with some probability to occur.
- Born's rule allows to find the probability of measuring specific values of Q .
- According to the type of observable we are considering (quantized or not), Born's rule will get a *probability* ($P(Q = q_i)$) or a *probability density* ($\rho(q)$). Continuous spectra allows continuous quantities, while discrete spectra allows only the eigenvalues of the eigenvectors associated to the quantized observable.

All of this is supported by experimental evidences, which is kinda incredible to think that this actually holds.

15.4 Define Observable Operators

The only thing left to do is how to define the operators representing physical quantities.

15.4.1 Commutator

We start by the assumption that we can describe any physical observable by finding its Hermitian operator $\hat{Q}$, whose eigenvalues are the possible results of a measurement. Then we need to introduce the concept of **commutator operator**:

$$[\hat{A}, \hat{B}] = \hat{A}\hat{B} - \hat{B}\hat{A}$$

which tells us if we can switch the order of application of two operators. Now, since we said:

$$\hat{Q}|q\rangle = q|q\rangle$$

if two operators commute, they will share the same orthonormal basis of vectors $\{|q_i\rangle\}$, in fact:

$$\begin{aligned} [\hat{A}, \hat{Q}]|q\rangle &= (\hat{Q}\hat{A} - \hat{A}\hat{Q})|q\rangle = 0 \\ \Rightarrow \hat{Q}\hat{A}|q\rangle &= \hat{A}\hat{Q}|q\rangle = q\hat{A}|q\rangle \end{aligned}$$

which means that $\hat{A}$ must have the same eigenvectors of $\hat{Q}$ in order to commute. If two operators commute, we usually write the state of the system in terms of the eigenvalues of both operators for comodity:

$$|q, a\rangle : \{\hat{Q}|q, a\rangle = q|q, a\rangle, \hat{A}|q, a\rangle = a|q, a\rangle\}$$

since they share the same eigenvector basis. We said that an eigenvalue q is the certain result of a measurement of an observable Q when the system is exactly in the state $|q\rangle$. This means that for two commuting operators, we understand that we can measure both the observables Q and A with certainty if the system is in the state $|q, a\rangle$.

If the two operators do not commute, so if they have a non-zero commutator $[\hat{A}, \hat{B}] \neq 0$, then an important property arises: the **uncertainty principle**. This means that it doesn't exist a state $|\Psi\rangle$ in which we can measure at the same time the two observables Q and A with certainty, since they have different eigenvectors.

15.4.2 Generating Observables

The process of generating the "observable operator" for a specific physical quantity is non trivial and quite empirical. We usually need to start from physical relations in the observable space, then we manipulate them in order to extract a linear operator relationship, and with that we define the observable.

The energy operator is a good example to understand the process. We start with the one dimensional wave function:

$$\Psi(x, t) = e^{i(kx - \omega t)}$$

and we consider its time derivative:

$$\frac{\partial \Psi(x, t)}{\partial t} = -i\omega e^{i(kx - \omega t)} = -i\omega \Psi(x, t) .$$

We can use De Broglie relation $E = \hbar\omega$ to rewrite:

$$\frac{\partial \Psi(x, t)}{\partial t} = -i \frac{E}{\hbar} \Psi(x, t) ,$$

that can finally be written as:

$$E\Psi(x, t) = i\hbar \frac{\partial \Psi(x, t)}{\partial t} .$$

At this point, we notice that we are actually applying an "operator" to the wave function $\Psi(x, t)$ that extracts the *energy* of the function. We can rewrite it as:

$$E\Psi(x, t) = \hat{E}\Psi(x, t) , \quad \hat{E} = i\hbar \frac{\partial}{\partial t} .$$

In this way, we defined the **energy operator** for a system that can be described by a wave function $\Psi(x, t)$ (which are all systems in quantum mechanics).

Here is the list of all most notable observables operators:

- **Position Operator** : $\hat{q} = q$
- **Momentum Operator** : $\hat{p} = -i\hbar\nabla$
- **Angular Momentum Operator** : $\hat{L} = q \times -i\hbar\nabla$
- **Energy Operator** : $\hat{E} = i\hbar \frac{\partial}{\partial t}$

Now that we defined all the operators, we can actually prove **Heisenberg Uncertainty**

15.5 Schrödinger Equation

Schrödinger equation the exact equivalent of Newtons's second law of Dynamics for quantum systems. It comes from empirical observations regarding both the wave-like probabilistic behavior of quantum particles, and de Broglie particle wavelength assumption – in addition to the already known photon's wavelength.

The idea is to define for a single particle a function $\Psi(x, t)$ that For a single non-relativistic particle of mass m , moving in a potential $V(\mathbf{r}, t)$, the time-dependent Schrödinger equation is given by:

$$i\hbar \frac{\partial \Psi}{\partial t}(x, t) = \left[-\frac{\hbar^2}{2m} \nabla^2 + V(x, t) \right] \Psi(x, t)$$

which if we consider a steady state – thus we remove the time dependency – we get:

$$-\frac{\hbar^2}{2m} \nabla^2 \psi(x) + V(x) \psi(x) = E \psi(x)$$

where we decomposed the wave function in two pieces $\Psi(x, t) = \psi(x) \phi(t)$.

Since we are assuming that the wave function is a probability density function, it should be normalized in L^2 space:

$$\int_{-\infty}^{+\infty} |\Psi(x, t)|^2 dx = 1$$

In Dirac notation we rewrite all the relations above:

$$i\hbar \frac{\partial}{\partial t} |\Psi(t)\rangle = \hat{H}(t) |\Psi(t)\rangle \longrightarrow \text{Schrödinger Equation} \quad (17)$$

$$\hat{H} = \left[-\frac{\hbar^2}{2m} \nabla^2 + V(x) \right] \longrightarrow \text{Hamiltonian Operator} \quad (18)$$

$$\langle \Psi(t) | \Psi(t) \rangle = 1 \longrightarrow \text{Normalization Condition} \quad (19)$$

Dirac notation is useful to express quantum states as elements of Hilbert space, without having to refer to a specific basis.

It's important to notice that given a system of two particles p_1 and p_2 the system is described by the equation (17). However, the system is not separable into single pieces as $|\Psi(t)\rangle = |\Psi_{p_1}(t)\rangle \otimes |\Psi_{p_2}(t)\rangle$, because of the interaction terms between the two particles in the Hamiltonian, expressed as potential: