

Fundamentals of Ecology and Evolutionary Biology

BIOS 20153 Lecture Notes

Stefano Allesina

2026-10-06

Table of contents

1	Introduction	1
1.1	Ecology	1
1.2	Approach	3
1.3	Topics	3
2	Unbounded growth	5
2.1	Overview	5
2.2	Historical models: geometric growth	5
2.3	Exponential growth	12
2.4	Life tables, population pyramids and demographic transition	21
2.4.1	The Euler-Lotka equation	23
2.5	Structured populations	26
3	Density dependence	27
3.1	Overview	27
3.2	Limits to growth: the logistic model	27
3.3	Density dependence in the real world	27
3.4	The logistic map	27
4	Pathogens	31
4.1	Overview	31
4.2	Compartmental models	31
4.3	Susceptible-Infected-Recovered (SIR) model	32
4.3.1	Dynamics of the SIR model	34
4.4	Number of people experiencing infection over time	35
4.5	Vaccination	37

Table of contents

4.6 Other compartmental models	40
5 Predator-prey dynamics	47
5.1 Overview	47
References	49

1 Introduction

These are the lecture notes for **BIOS 20153 Fundamentals of Ecology and Evolutionary Biology**, Academic Year 2026-27.

- *Who am I:* Stefano Allesina, Professor in the Department of Ecology & Evolution; at UofC since 2009.
- *Main interests:* Theoretical and community ecology, dynamical systems, mathematical biology.
- *Email:* sallesina@uchicago.edu
- *Office:* Zoology 403a (next to Botany Pond)
- *Office hours:* We can setup a time once per week, or you can setup a meeting/zoom via email.

1.1 Ecology

Ecology is the study of the **distribution and abundance of organisms**.

It considers organisms at the individual, population, community, ecosystem, and biosphere levels. In this class, we are going to focus on the population and community levels. Ecology overlaps with the closely related sciences of biogeography, evolutionary biology, genetics, ethology, and natural history.

1 Introduction




Figure 1.1: Levels of biological organization relevant to ecology. Image: [Wikimedia Commons](#), CC BY-SA 3.0.

1.2 Approach

Many factors determine the abundance and distribution of the organisms of a species, including:

- The number of individuals in the population
- The distribution of their age, sex, and size
- The availability of resources
- The presence of other species, including predators, prey, parasites, etc.
- The environmental conditions

Keeping track of these factors, and determine how the abundance of a species will change in time is very hard. Historically, the best strategy has been to take a **quantitative approach**, and write **mathematical models of population dynamics**.

Therefore, you will see a fair number of **equations** in this class. The prerequisites are single-variable calculus (and possibly some multivariate calculus)—all the rest will be explained in detail. Importantly:

- you will **not** have to perform extensive calculations on the final, but
- you will need to be able to **interpret equations that are provided**

1.3 Topics

1. Unbounded growth
 - a. Demography, age structure
 - b. Endangered species
2. Density-dependent growth
 - a. Logistic growth
 - b. Discrete generations & cycles

1 Introduction

- c. Fisheries management

3. Pathogens

- a. Consequences of vaccination
- b. Effects of diseases on animal populations

4. Population Cycles

- a. Predator-prey
- b. Host-pathogen

2 Unbounded growth

2.1 Overview

2.2 Historical models: geometric growth

Among the first examples showing how growth can rapidly lead to large populations is an exercise found in the *Liber abaci* (1202) by Leonardo Da Pisa, known as Fibonacci.

A certain man had one pair of rabbits together in a certain enclosed place, and one wishes to know how many are created from the pair in one year when it is the nature of them in a single month to bear another pair, and in the second month those born to bear also.

What is a bit unclear from the description: each pair of rabbits matures in one month, and once mature it produces a new pair of immature rabbits per month. Moreover, rabbits do not die! Then, calling R_t the number of rabbits at month t we obtain the famous Fibonacci's sequence:

$$1, 1, 2, 3, 5, 8, 13, \dots$$

By defining $R_1 = 1$ and $R_2 = 1$, we can write this implicitly as:

$$R_t = R_{t-1} + R_{t-2} \quad t \geq 3$$

2 Unbounded growth

or solve the recursion to obtain:

$$R_t = \frac{\phi^n - \psi^n}{\sqrt{5}}$$

where we have defined the golden ratio $\phi = \frac{1}{2}(1+\sqrt{5})$ and its conjugate $\psi = \frac{1}{2}(1-\sqrt{5})$. Thus, after one year (12 months) we will have 144 rabbits, and after 3 years $R_{36} = 14,930,352$. Let's plot the growth of the population in time




Figure 2.1: Fibonacci's rabbits: number of pairs R_t over 36 months.

Because the numbers grow so rapidly, it is convenient to plot $\log R_t$ vs t

You can see that, after some jiggling, the curve in the logarithmic scale approaches a line; this is because, for $n \gg 1$ we have

2.2 Historical models: geometric growth




Figure 2.2: Log-scale plot of Fibonacci's rabbits: $\log R_t$ grows nearly linearly, reflecting exponential growth.

2 Unbounded growth

$$R_t \approx \phi R_{t-1}$$

and thus the number of rabbits will grow of approximately 62% per month. Soon, rabbits would take over the world!

i Rabbits in Australia

On Christmas day 1859, Thomas Austin released 24 breeding rabbits (*Oryctolagus cuniculus*) on his estate as game for shooting parties. The rabbits soon expanded across the continent, growing to concerning numbers and with devastating impacts on both agriculture and wildlife.




Figure 2.3: Rabbit expansion in Australia. Figure 1 from Alves et al. (2022).

Estimates vary widely (quantitative ecology was born in the late 1920s, and statistical ecology even later), but by 1910 there were about 1 billion rabbits roaming the continent. A recent genetic analysis (Alves et al. (2022)) has shown that they all descended from the original rabbits from Thomas Austin!

2.2 Historical models: geometric growth

In 1748, Leonard Euler published *Introductio in analysin infinitorum*, and included the following example:

If we assume that the human race was re-established by six individuals after the Flood—let us say, over a period of two hundred years—and that the population grew to one million, the question arises as to the rate at which the number of people must have increased annually.

He started from the equation:

$$P_{t+1} = (1 + x)P_t$$

with x representing the annual increase of the population. Iterating:

$$\begin{aligned}P_1 &= (1 + x)P_0 \\P_2 &= (1 + x)P_1 = (1 + x)(1 + x)P_0 = (1 + x)^2P_0 \\P_3 &= (1 + x)P_2 = (1 + x)^3P_0 \\&\dots\dots \\P_n &= (1 + x)P_{n-1} = (1 + x)^nP_0\end{aligned}$$

Then one needs to solve (note—we write $\log$ as the natural logarithm):

2 Unbounded growth

$$\begin{aligned}P_{200} &= (1+x)^{200}P_0 \\10^6 &= (1+x)^{200}6 \\ \frac{10^6}{6} &= (1+x)^{200} \\ \log \frac{10^6}{6} &= 200 \log(1+x) \\ 12.02 &\approx 200 \log(1+x) \\ 0.06 &\approx \log(1+x) \\ 1.062 &\approx 1+x \\ x &\approx 0.062\end{aligned}$$

He found the number reasonable, but mused:

But if, by the same rate of increase, the human population had grown over a period of 400 years, it would have risen to a number—16,666,666,666—which the entire earth would have been wholly inadequate to sustain.

We will return to this thought in the next chapter.

Geometric growth

These are all cases of **geometric growth**, in which the population in the next period is given by:

$$P_{t+1} = (1+x)P_t$$

If $x > 0$, the population will expand; if $x < 0$ it will contract; if it is exactly $x = 0$ it will remain constant. When the population is expanding, it will expand very rapidly; when it contracts, it will rapidly approach zero.

2.2 Historical models: geometric growth

The explosive nature of growth and decline has important consequences for natural populations, the spread of diseases, and even human society. Thomas Malthus, in the abrasive *An Essay on the Principle of Population* (Malthus (1798)) wrote that

Population, when unchecked, increases in a geometrical ratio. Subsistence increases only in an arithmetical ratio. A slight acquaintance with numbers will shew the immensity of the first power in comparison of the second.

He concluded that, as a result, population growth naturally outpaces the food supply, inevitably leading to poverty, famine, and population reduction unless checked by moral restraint, disease, war, or starvation. These bleak conclusions attracted praise and harsh critique, making this slim treatise one of the most influential books of the last few centuries.

Among the critics, Marx and Engels were among the most vitriolic. Engels described Malthus's thesis as (Engels (1844)):

the crudest, most barbarous theory that ever existed, a system of despair which struck down all those beautiful phrases about love thy neighbour and world citizenship

In the realm of biology, the *struggle for existence* implicit in the competition for scarce resources inspired the founders of evolutionary theory, and provided a mechanism for natural selection. Darwin wrote in *The Origin of Species* (Darwin (1859)):

This is the doctrine of Malthus, applied with manifold force to the animal and vegetable kingdoms, for in this case there can be no artificial increase of food, and no prudential restraint from marriage

and in his autobiography:

2 Unbounded growth

In October 1838 [...] I happened to read for amusement ‘Malthus on Population’ [...] it at once struck me that under these circumstances favourable variations would tend to be preserved, and unfavourable ones to be destroyed. The result of this would be the formation of new species.

Similarly, Wallace wrote in his autobiography (Wallace (2016)):

But perhaps the most important book I read was Malthus’s Principles of Population [...] It was the first great work I had yet read treating of any of the problems of philosophical biology, and its main principles remained with me as a permanent possession, and twenty years later gave me the long-sought clue to the effective agent in the evolution of organic species

2.3 Exponential growth

The geometric growth model is well suited for populations where reproduction occurs in discrete time steps (such as seasonally breeding plants or animals). However, many populations, including humans and bacteria, reproduce continuously. To account for this, we can extend the discrete model into continuous time.

Suppose a population has an annual per-capita growth rate r . If reproduction occurs not just once a year, but is split into k discrete reproductive periods per year, the growth rate per interval is $\frac{r}{k}$. The population size at time t years becomes:

$$N_t = N_0 \left(1 + \frac{r}{k}\right)^{kt}$$

Now we need to remember a key result from algebra: as k increases, the term $\left(1 + \frac{r}{k}\right)^k$ approaches e^r . This is a consequence of the definition of Euler’s number e :

2.3 Exponential growth

$$e = \lim_{x \rightarrow \infty} \left(1 + \frac{1}{x}\right)^x = 1 + \frac{1}{2} + \frac{1}{3!} + \frac{1}{4!} + \dots = 2.71828 \dots$$

Now consider the change of variables:

$$x = \frac{k}{r} \implies k = rx$$

and therefore

$$\left(1 + \frac{r}{k}\right)^k = \left(1 + \frac{1}{x}\right)^{rx} = \left[\left(1 + \frac{1}{x}\right)^x\right]^r$$

Because $\lim_{x \rightarrow \infty} \left(1 + \frac{1}{x}\right)^x = e$, we have

$$\lim_{k \rightarrow \infty} \left(1 + \frac{r}{k}\right)^k = e^r$$

In summary, taking the limit as $k \rightarrow \infty$ transforms the discrete compounding model into the continuous exponential growth equation:

$$N(t) = N(0)e^{rt}$$

Here, $N(t)$ denotes population size at any time $t \geq 0$.

! Key Distinction

In discrete growth, population size N_t is only defined at specific intervals ($t = 0, 1, 2 \dots$). In continuous growth, $N(t)$ is defined for **all** $t \geq 0$.

To find how fast the population is growing at any instant, we take the derivative of $N(t)$ with respect to time:

2 Unbounded growth

$$\frac{dN(t)}{dt} = \frac{d}{dt} [N(0)e^{rt}]$$

Using the exponential derivative rule ($\frac{d}{dt}e^{rt} = re^{rt}$):

$$\frac{dN(t)}{dt} = r \cdot \underbrace{N(0)e^{rt}}_{N(t)} = rN(t)$$

Biological Meaning

This reveals that the instantaneous growth rate ($\frac{dN}{dt}$) is simply the per-capita rate r multiplied by the current population size $N(t)$.

The interpretation of r is the difference between birth and death rates per individual. If $r > 0$, the population grows; if $r < 0$, it declines; if $r = 0$, it remains constant.

If we divide both sides of the equation by $N(t)$, we compute the per-capita growth rate:

$$\frac{1}{N(t)} \frac{dN(t)}{dt} = r$$

This shows that the per-capita growth rate is constant, independent of population size. This is a key assumption of the exponential growth model. We say that the **growth of the population is independent of its density**.

Are births uniformly distributed throughout the year?

The exponential growth model assumes a constant per-capita birth rate. But are births actually spread evenly across the calendar? The figure below shows the average number of births per calendar day in the United States, based on Social Security Administration records

2.3 Exponential growth

for 2000–2014.




Figure 2.4: Average daily births in the US (SSA data, 2000–2014). Red points mark major public holidays.

Notice the sharp dips on Christmas Day, New Year's Day, and Independence Day. Why might that be? And what does it imply about the assumption of a constant birth rate in our model?

Consider again the solution

$$N(t) = N(0)e^{rt}$$

and take the natural logarithm of both sides:

$$\log N(t) = \log N(0) + rt$$

2 Unbounded growth

This means that if we plot $\log N(t)$ against time t , we should see a straight line with slope r and intercept $\log N(0)$. This is a useful property for analyzing population growth data: a semi-log plot reveals the behavior of the model at a glance, because growth ($r > 0$) appears as a line with positive slope, decline ($r < 0$) as a line with negative slope, and a constant population ($r = 0$) as a horizontal line.




Figure 2.5: Semi-log plot of $N(t)$ for three values of r , with $N(0) = 1$. Growth, constancy, and decline appear as straight lines with positive, zero, and negative slope, respectively.

On the original (linear) scale, the same three trajectories make the asymmetry between growth and decline vivid: growing populations accelerate away rapidly, while declining ones approach zero gradually.

2.3 Exponential growth




Figure 2.6: Population size $N(t)$ on a linear scale for the same three values of r as Figure 2.5.

2 Unbounded growth

i Human population growth

The world human population at the end of the last Ice Age (~10,000 BCE) stood at roughly 4–5 million people. It took more than ten millennia to reach one billion (around 1800 CE), yet only two more centuries to multiply eightfold. The figure below shows this history on a log scale using estimates compiled by Our World in Data (2024) from HYDE, Gapminder, and UN data.




Figure 2.7: World human population from 10,000 BCE to 2023 on a log scale (Our World in Data (2024)). The slope is the intrinsic rate of increase r ; note how it steepens after the Industrial Revolution.

On the log scale, a constant r would appear as a straight line. The curve is not straight: r was low and roughly constant for millennia, then rose sharply after ~1800. What changed? Agricultural revo-

2.3 Exponential growth

lutions, industrialization, and advances in public health all reduced death rates while birth rates remained high — a transient but enormous surge in r that Malthus could not have anticipated.

Another notable feature of the exponential growth model is that if we divide both sides of the equation by $N(0)$, we find that the right-hand side depends only on the product rt , and not the population density.

$$\frac{N(t)}{N(0)} = e^{rt}$$

Now suppose that the population has doubled in size after τ years. Then we have:

$$\frac{N(\tau)}{N(0)} = 2 = e^{r\tau}$$

How long does it take for a population to double in size? We can solve for τ :

$$\log 2 = r\tau \implies \tau = \frac{\log 2}{r}$$

The value of $\log 2$ is approximately 0.693, so we can approximate the doubling time as:

$$\tau \approx \frac{0.693}{r} \approx \frac{70}{100r}$$

This is known as the “rule of 70”: a population with a constant growth rate of 1% will double in size after approximately 70 years, while a population with a constant growth rate of 10% will double in size after approximately 7 years.

2 Unbounded growth

Naturally, a population can double in time only if $r > 0$. If $r < 0$, the population will halve in size after $\tau = \frac{\log 2}{|r|}$ years, instead.

i Radiocarbon dating and Ötzi the Iceman

When $r < 0$, the exponential model describes decay rather than growth. One of the most consequential applications is **radiocarbon dating**, developed by Willard Libby at the University of Chicago — an achievement recognized with the Nobel Prize in Chemistry in 1960.

All living organisms continuously exchange carbon with the atmosphere, maintaining a fixed ratio of radioactive ^{14}C to stable ^{12}C . Upon death, exchange stops and ^{14}C decays exponentially. Its half-life is $t_{1/2} = 5730$ years, giving

$$r = -\frac{\log 2}{5730} \approx -0.000121 \text{ yr}^{-1}$$

By measuring the fraction of ^{14}C remaining in a sample, one can invert the exponential equation and solve for the time elapsed since death.

In September 1991, a mummified man — now known as Ötzi the Iceman — was found in a glacier at the Hauslabjoch on the Austrian–Italian border. Nine independent radiocarbon measurements of bone, tissue, and grass from his clothing were carried out by five laboratories (Prinother-Fornwagner and Niklaus 1994). The figure shows these measurements together with their weighted average.

2.4 Life tables, population pyramids and demographic transition




Figure 2.8: Radiocarbon measurements of Ötzi the Iceman from nine laboratory analyses (Prineth-Fornwagner and Niklaus 1994). Error bars show $\pm 2\sigma$; the dashed line and shaded band give the weighted mean 4546 ± 17 yr BP.

The weighted mean of 4546 ± 17 yr BP — years before present, where “present” is conventionally 1950 CE — places Ötzi’s death between approximately 3350 and 3100 BCE, during the Copper Age.

2.4 Life tables, population pyramids and demographic transition

In the model above, the growth of a population is determined by the per-capita growth rate r , which is the difference between the birth and death rates. However, there are different ways to achieve the same value of r .

2 Unbounded growth

For example, a population with a high birth rate and a high death rate can have the same r as a population with a low birth rate and a low death rate.

Demographers use **life tables** to summarize the age-specific birth and death rates of a population.




Figure 2.9: The demographic transition: as countries develop, death rates fall first (reducing r of mortality), then birth rates follow. Figure from Our World in Data / Wikimedia Commons.

2.4 Life tables, population pyramids and demographic transition

2.4.1 The Euler-Lotka equation

To connect age-structured demography to the exponential growth model, we need two schedules:

- l_x — the **survival function**: the probability that a newborn female survives to exact age x (by convention $l_0 = 1$).
- m_x — the **maternity function**: the expected number of female births produced per female per unit time at age x .

The **net reproductive rate** R_0 is the mean number of daughters produced by a female over her lifetime:

$$R_0 = \sum_x l_x m_x \cdot \Delta x$$

where the sum runs over reproductive ages and Δx is the width of each age interval. If $R_0 > 1$ the population grows on average; if $R_0 < 1$ it declines; $R_0 = 1$ gives exact replacement.

The **Euler-Lotka equation** links the age-specific schedules to the intrinsic rate of increase r :

$$1 = \sum_x l_x m_x e^{-rx} \cdot \Delta x \quad (2.1)$$

When $r = 0$, Equation 2.1 reduces to $R_0 = 1$. For $r \neq 0$, the exponential weight e^{-rx} discounts (or inflates) contributions from late ages relative to early ones, so a population growing faster ($r > 0$) discounts late reproduction more heavily.

A useful approximation is obtained by defining the **mean generation time** $T = \frac{\sum_x x l_x m_x}{\sum_x l_x m_x}$ and noting that $r \approx \frac{\log R_0}{T}$.

2 Unbounded growth




Figure 2.10: Survival function l_x for US females (CDC life table, 2020). The value at each age is the probability that a newborn survives to that age.

2.4 Life tables, population pyramids and demographic transition




Figure 2.11: Maternity function m_x for US females (CDC NCHS, 2018): female births per female per year, accounting for the sex ratio at birth (≈ 0.4886 female).

2 Unbounded growth

The figures below show l_x and m_x for US females, using the 2020 CDC life table and 2018 age-specific fertility rates.

Applying Equation 2.1 to the US data gives $R_0 = 0.831$, a mean generation time of $T \approx 29.5$ years, and an intrinsic rate of increase of $r \approx -0.0063 \text{ yr}^{-1}$. The negative r reflects that US fertility has been below replacement ($R_0 < 1$) since the early 1970s — a pattern consistent with the demographic transition shown in Figure 2.9.

2.5 Structured populations

A key assumption of the exponential growth model is that all individuals are identical, and thus have the same birth and death rates. In reality, populations are structured by age, size, or stage of development, and these differences can have profound effects on population dynamics.

3 Density dependence

3.1 Overview

3.2 Limits to growth: the logistic model

3.3 Density dependence in the real world

3.4 The logistic map

The **logistic map** is the simplest discrete-time model that produces both stable dynamics and chaos:

$$N_{t+1} = r N_t (1 - N_t)$$

where $N_t \in [0, 1]$ is the population scaled by its maximum and $r \in [0, 4]$ is the growth parameter. For $r < 1$ the population collapses to zero; for $1 < r < 3$ it converges to a stable equilibrium $N^* = 1 - 1/r$; beyond $r = 3$ the equilibrium destabilises through a cascade of period-doubling bifurcations that culminates in deterministic chaos near $r \approx 3.57$.

3 Density dependence




Figure 3.1: Bifurcation diagram of the logistic map. For each value of r , the attractor values of N_t are plotted after discarding a 500-step transient. Interactive version: [<>#sec-logistic](#).

3.4 The logistic map




Figure 3.2: Logistic map dynamics N_t starting from $N_0 = 0.5$ at four representative values of r , illustrating fixed-point, two-cycle, and chaotic regimes.

4 Pathogens

4.1 Overview

4.2 Compartmental models

Simple models for the spread of infectious diseases are often based on the idea of **compartments**. Individuals in a population are classified into different compartments based on their disease status, and the model describes how individuals move between compartments over time.

Common compartments include: Susceptible (available for infection, S), Infected (and Infectious!, I), and Recovered (or Removed, R). The dynamics of the disease can be described using differential equations that govern the rates of change between these compartments.

In most cases, one considers the total population fixed (the dynamics of the disease are rapid compared to the dynamics of the population), so that $S + I + R = N$ is constant, and $N = 1$ without loss of generality (i.e., compartments represent the proportion of the population in each state). The simplest model is the **SIR model**, which assumes that individuals move from being susceptible to infected, and then to recovered. This model was proposed by Kermack and McKendrick in 1927, and is still widely used today, despite being exactly a century old!

Many variations on this basic framework exist, including models that account for births and deaths, vaccination, waning immunity, and more. The

SIR model is a foundational tool in epidemiology and provides insights into the dynamics of infectious diseases.

4.3 Susceptible-Infected-Recovered (SIR) model

The population is divided into three compartments: Susceptible (S), Infected (I), and Recovered (R). Susceptible individuals can become infected upon contact with infected individuals, and infected individuals can recover from the disease, thereby gaining long-lasting immunity. The model is best understood using boxes and arrows to represent the flow of individuals between compartments:




Figure 4.1: Flow diagram of the SIR model. Susceptible individuals (S) become infected at rate βSI upon contact with infectious individuals (I), who recover at rate γI .

The dynamics of the SIR model can be described by the following set of ordinary differential equations (ODEs):

4.3 Susceptible-Infected-Recovered (SIR) model

$$\begin{aligned}\frac{dS}{dt} &= -\beta SI \\ \frac{dI}{dt} &= \beta SI - \gamma I \\ \frac{dR}{dt} &= \gamma I\end{aligned}$$

The dynamics of this model are governed by two key parameters: the transmission rate β and the recovery rate γ . β can be thought of as the rate at which susceptible individuals become infected upon contact with infectious individuals, while γ is the rate at which infected individuals recover and move to the recovered compartment (for example, the reciprocal of the average infectious period). The trajectories of the compartments cannot be solved analytically in general, but can be simulated numerically. However, we can make sense of the dynamics by considering that a) the number of susceptible individuals can only decrease over time, and b) the number of infected individuals can increase or decrease depending on the relative sizes of the susceptible and infected compartments. In particular:

$$\frac{1}{I} \frac{dI}{dt} = \beta S - \gamma$$

and thus the number of infected individuals will increase when $\beta S/\gamma > 1$. Suppose that the population is initially entirely susceptible, so that $S \approx 1$. Then the condition for the number of infected individuals to increase is $\beta/\gamma > 1$, which leads to the concept of the **basic reproduction number** $R_0 = \beta/\gamma$, which determines whether an infection will spread in a population. If $R_0 > 1$, the infection can invade and cause an epidemic; if $R_0 < 1$, the infection will die out. This quantity can be interpreted as the average number of secondary infections produced by a single infected individual in a completely susceptible population. The basic reproduction number is a fundamental concept in epidemiology and is used to assess the potential for disease outbreaks and to inform public health interventions.

4 Pathogens

With these qualitative insights, we can now explore the dynamics of the SIR model in more detail, including the effects of varying parameters and initial conditions on the spread of infectious diseases.

4.3.1 Dynamics of the SIR model




Figure 4.2: SIR model dynamics for three initial susceptible fractions ($S_0 = 0.25, 0.65, 0.95$), with $I_0 = 0.01$, $\beta = 2$, $\gamma = 1$ ($\mathcal{R}_0 = 2$). When $S_0 < \gamma/\beta = 0.5$ the infection dies out immediately; when $S_0 > 0.5$ an epidemic develops. Interactive version: <https://your-course-site#sec-topic3-sir-dynamics>.

4.4 Number of people experiencing infection over time

The SIR model cannot be solved analytically, but we can determine what fraction of the population was infected over the entire course of an epidemic — the **final attack rate** — without integrating the differential equations. As $t \rightarrow \infty$ the epidemic subsides: $I(\infty) = 0$, so all individuals are either susceptible or recovered, and $S(\infty) + R(\infty) = 1$.

To find $S(\infty)$, we eliminate time from the ODEs by computing

$$\frac{dS}{dR} = \frac{dS/dt}{dR/dt} = \frac{-\beta SI}{\gamma I} = -\frac{\beta}{\gamma} S = -R_0 S.$$

This separable equation integrates to $S(t) = S_0 \exp(-R_0(R(t) - R(0)))$. Starting from a nearly fully susceptible population ($S_0 \approx 1$, $R(0) = 0$), and taking $t \rightarrow \infty$ with $R(\infty) = 1 - S(\infty)$, we obtain the **final size relation**:

$$S(\infty) = \exp(-R_0(1 - S(\infty))). \quad (4.1)$$

This is a fixed-point equation: the right-hand side depends on $S(\infty)$ itself, so it cannot be solved algebraically. It can, however, be solved numerically for any given R_0 . The final attack rate $1 - S(\infty)$ is plotted as a function of R_0 in Figure 4.3.

A striking feature of Equation 4.1 is that the final attack rate is entirely determined by R_0 , regardless of the individual values of β and γ . Two diseases with different transmission and recovery rates but the same R_0 will infect the same total fraction of the population — even though their time courses may look very different.

To see how R_0 influences this quantity, we can use the approximation:




Figure 4.3: Final attack rate $1 - S(\infty)$ as a function of R_0 , obtained by solving Equation 4.1 numerically. Below the epidemic threshold $R_0 = 1$ (red dashed line) the infection always dies out; above it a non-zero fraction of the population is infected. For $R_0 = 2$, approximately 80% of a fully susceptible population is eventually infected before the epidemic ends.

$$\log S(\infty) \approx -R_0(1 - S(\infty)) \approx -R_0$$

when R_0 is small

4.5 Vaccination

Explain how vaccination can be incorporated into the SIR model, and how it affects the dynamics of disease spread. Discuss herd immunity and the critical vaccination threshold.

here it would be good to have a timeline for the eradication of smallpox and the near eradication of polio

i The resurgence of measles in the United States

Measles is one of the most transmissible human pathogens, with R_0 typically estimated between 12 and 18, so that vaccination coverage of roughly 95% is needed to prevent outbreaks. The figure below shows the number of reported cases in the United States since 1938.




Figure 4.4: Reported measles cases in the US, 1938–2026 (log scale). Data from the CDC and earlier public health reports, compiled by Our World in Data. Dashed lines mark the licensure of the first measles vaccine (1963) and the declaration that measles was eliminated from the US (2000). The 2026 count (open circle) is incomplete: it covers cases through October 1.

Before the vaccine, hundreds of thousands of cases were reported every year. The introduction of the vaccine in 1963 led to a dramatic decline in measles cases.

Note the spike in the number of cases in the period 1989–1991. The cause of this spike was attributed to low vaccination coverage among low-income, inner-city preschool-aged children who had not received their single recommended dose due to healthcare access barriers, underfunded public health infrastructure, and missed vaccination op-

portunities. In 1989, the CDC recommended adding a second dose of MMR before school entry. This was introduced because ~2–5% of individuals failed to develop immunity after a single dose (primary vaccine failure). This allowed to reach the minimum number of cases in recorded history, and in 2000 endemic transmission had stopped. Vaccination coverage among kindergartners has been reported by the CDC since the 2009–10 school year (Figure 4.5). National coverage with MMR hovered around the 95% needed to block transmission until 2019–20, and has since fallen steadily, while the fraction of children with an exemption from one or more vaccines has risen to its highest level on record.

Since then, imported cases have found pockets of unvaccinated individuals, and the numbers have risen again: 2025 saw 2,289 cases, and by October 2026 the count was the highest since 1991. Why does a small decline in vaccination coverage, which may look harmless, have such a large effect on a disease with a very high R_0 ?




Figure 4.5: MMR vaccination coverage (left) and exemption rate (right) among US kindergartners, school years 2009–10 to 2025–26. The dashed line marks 95% coverage. Data: CDC SchoolVaxView (state-reported); the 2010–11 school year was not reported. Note that coverage is measured at school entry.

4.6 Other compartmental models

A common variation on the SIR model is the inclusion of demography. Newborns enter the population as susceptible individuals, and all individuals experience a natural death rate. This leads to the **SIR model with demography**, which can be described by the following set of ODEs:

4.6 Other compartmental models




Figure 4.6: SIR model with demography. Newborns enter the susceptible class at constant rate Λ ; all compartments experience natural mortality at per-capita rate μ .

$$\begin{aligned}\frac{dS}{dt} &= \Lambda - \beta SI - \mu S \\ \frac{dI}{dt} &= \beta SI - \gamma I - \mu I \\ \frac{dR}{dt} &= \gamma I - \mu R\end{aligned}$$

In this model if $\Lambda = \mu$, the total population size remains constant, and the dynamics of the disease are similar to those of the basic SIR model. However, if $\Lambda > \mu$, the population grows over time, while if $\Lambda < \mu$, the population declines.

A simpler version of the model is the **SIS model**, which assumes that individuals do not gain long-lasting immunity after infection. In this case, individuals move from the Susceptible compartment to the Infected compartment and back again, without ever entering the Recovered compartment. The dynamics of the SIS model can be described by the following

4 Pathogens

set of ODEs:



Figure 4.7: Flow diagram of the SIS model. Recovered individuals return immediately to the susceptible class at rate γI , conferring no lasting immunity.

$$\begin{aligned}\frac{dS}{dt} &= -\beta SI + \gamma I \\ \frac{dI}{dt} &= \beta SI - \gamma I\end{aligned}$$

For diseases that do not confer long-lasting immunity, individuals may move from the Recovered compartment back to the Susceptible compartment. This leads to the **SIRS model**, which includes a waning immunity term. Other models may include additional compartments, such as Exposed (E) for diseases with a latent period, leading to the SEIR model, leading to the following set of ODEs:

4.6 Other compartmental models




Figure 4.8: SEIR model with waning immunity. Exposed individuals (E) progress to infectious at rate σE ; recovered individuals (R) lose immunity and re-enter the susceptible class at rate δR .

$$\begin{aligned}
 \frac{dS}{dt} &= -\beta SI + \delta R \\
 \frac{dE}{dt} &= \beta SI - \sigma E \\
 \frac{dI}{dt} &= \sigma E - \gamma I \\
 \frac{dR}{dt} &= \gamma I - \delta R
 \end{aligned}$$

You can find more complex models in which some compartments represent a disease vector (e.g., mosquitoes) or a pathogen (e.g., free-living bacteria) in the environment. These models are often referred to as **vector-borne** or **environmental transmission** models, respectively. While the number of compartments and the complexity of the models can vary widely, the basic principles of compartmental modeling remain the same: individuals

4 Pathogens

move between compartments based on rates that depend on the current state of the system.

Figure 4.9 shows a more complex example from Roosa and Chowell (2019): a model of **Zika virus transmission** that couples two interacting populations — humans and mosquitoes. The human population is divided into susceptible (S_h), exposed (E_h), asymptotically infectious (A_h), symptomatically infectious (I_{h1}), convalescent (I_{h2}), and recovered (R_h) classes. Mosquitoes are divided into susceptible (S_v), exposed (E_v), and infectious (I_v) classes. Transmission is bidirectional: infectious mosquitoes (I_v) infect susceptible humans (S_h), and infectious humans (A_h, I_{h1}) infect susceptible mosquitoes (S_v).

4.6 Other compartmental models




Figure 4.9: Compartmental diagram for the Zika virus transmission model of Roosa and Chowell (2019). The human population (top) is divided into six classes; the mosquito (vector) population (dark blue outline) into three. Arrows are labelled with the rates governing transitions between compartments.

5 Predator-prey dynamics

5.1 Overview

References

- Alves, Joel M, Miguel Carneiro, Jonathan P Day, John J Welch, Janine A Duckworth, Tarnya E Cox, Mike Letnic, Tanja Strive, Nuno Ferrand, and Francis M Jiggins. 2022. “A Single Introduction of Wild Rabbits Triggered the Biological Invasion of Australia.” *Proceedings of the National Academy of Sciences* 119 (35): e2122734119.
- Darwin, CR. 1859. *On the Origin of Species*. Harvard University Press, Cambridge, Massachusetts.
- Engels, Friedrich. 1844. “Outlines of a Critique of Political Economy.” *Deutsch-Französische Jahrbücher* 1 (1): 7.
- Malthus, Thomas Robert. 1798. “Essay on Principle of Population.” Readex Microprint.
- Our World in Data. 2024. “Population.” <https://ourworldindata.org/grapher/population>.
- Prinoth-Fornwagner, R., and Th.R. Niklaus. 1994. “The Man in the Ice: Results from Radiocarbon Dating.” *Nuclear Instruments and Methods in Physics Research B* 92: 282–90.
- Roosa, Kimberlyn, and Gerardo Chowell. 2019. “Assessing Parameter Identifiability in Compartmental Dynamic Models Using a Computational Approach: Application to Infectious Disease Transmission Models.” *Theoretical Biology and Medical Modelling* 16 (1): 1. <https://doi.org/10.1186/s12976-018-0097-6>.
- Wallace, Alfred Russel. 2016. *My Life a Record of Events and Opinions*. Read Books Ltd.

