

**Exercise 1.4.7:** Finish the proof of Theorem 1.4.5 by showing that the assumption  $\alpha^2 > 2$  contradicts the assumption that  $\alpha = \sup A$ .

*Proof.* Consider the set,

$$A = \{a \in \mathbb{R} : a^2 < 2\}.$$

Let  $\alpha = \sup A$ . Suppose to the contrary that  $\alpha^2 > 2$ . Consider an element of  $A$  that is smaller than  $\alpha$ , like  $(\alpha - \frac{1}{n})$ , where  $n > \frac{2\alpha}{\alpha^2 - 2}$ .

$$\begin{aligned} (\alpha - \frac{1}{n})^2 &= \alpha^2 - \frac{2\alpha}{n} + \frac{1}{n^2}, \\ &> \alpha^2 - \frac{2\alpha}{n}, \\ &> \alpha^2 - (\alpha^2 - 2), \\ &= 2. \end{aligned}$$

Thus we have shown that  $(\alpha - \frac{1}{n})$  is greater than  $a$  for all  $a \in A$  and therefore  $(\alpha - \frac{1}{n})$  is an upper bound. Since  $(\alpha - \frac{1}{n}) < \alpha$  we have contradicted  $\alpha = \sup A$ .

□

**Exercise Supplemental 1:** Give a from-scratch proof of the following facts:

(a) If  $f : A \rightarrow B$  has an inverse function  $g$ , then  $f$  is injective.

(b) If  $f : A \rightarrow B$  has an inverse function  $g$ , then  $f$  is surjective.

*Proof (a).* Suppose  $f : A \rightarrow B$ , whose inverse is  $g : B \rightarrow A$ , now consider  $a_i, a_j \in A$  such that  $f(a_i) = f(a_j)$ . Using  $g$  as an intermediary we get the equality,

$$\begin{aligned} f(a_i) &= f(a_j), \\ g(f(a_i)) &= g(f(a_j)), \\ a_i &= a_j. \end{aligned}$$

Thus we have shown  $f$  is an injective function.

□

*Proof (b).* Suppose  $f : A \rightarrow B$ , whose inverse is  $g : B \rightarrow A$ . Consider some  $b \in B$ , by definition of  $g$  we know that there exists some  $a \in A$  such that,  $g(b) = a$ . Taking the inverse of both sides we get,

$$\begin{aligned} f(g(b)) &= f(a) \\ b &= f(a). \end{aligned}$$

Since  $a \in A$  we have shown that for every  $b \in B$  there exists some  $a \in A$  such that  $f(a) = b$  thus  $f$  is surjective.

□

**Exercise Supplemental 2:** Show that the sets  $[0, 1)$  and  $(0, 1)$  have the same cardinality.

**Exercise 1.5.10 (a) (c):** (Wait until after Wednesday to start this one)

- (a) Let  $C \subseteq [0, 1]$  be uncountable. Show that there exists  $a \in (0, 1)$  such that  $C \cap [a, 1]$  is uncountable.
- (c) Determine, with proof, if the same statement remains true replacing uncountable with infinite.

*Proof (a).* Suppose  $C \subseteq [0, 1]$  is uncountable. Suppose the sake of contradiction that for all  $a \in (0, 1)$ ,  $C \cap [a, 1]$  is countable. Let  $a = \frac{1}{n}$ . Note that  $C \cap [\frac{1}{n}, 1]$  is countable. By Theorem 1.5.8 we now have since  $C \cap [\frac{1}{n}, 1]$  the infinite union is also countable. Through set theory

$$\begin{aligned} \bigcup_{n=1}^{\infty} C \cap [\frac{1}{n}, 1] &= C \cap (\bigcup_{n=1}^{\infty} [\frac{1}{n}, 1]) \\ &= C \cap (0, 1] \end{aligned}$$

Therefore,  $C \cap (0, 1] \cup \{0\} = C$  is also countable.  $\square$

*Proof (c).* Suppose the countably infinite set  $C = \{\frac{1}{n}, n \in \mathbb{N}\}$ . By Archimedean Principle we know that for all  $a \in (0, 1)$  we can find some  $\frac{1}{n} < a$ , and therefore we force the set  $C \cap [a, 1]$  to be finite.  $\square$

**Exercise Supplemental 3:** (Wait until after Wednesday to start this one) Suppose for each  $k \in \mathbb{N}$  that  $A_k$  is at most countable. Use the fact that  $\mathbb{N} \times \mathbb{N}$  is countably infinite to show that  $\bigcup_{k=1}^{\infty} A_k$  is at most countable. Hint: take advantage of surjection.

*Proof.* Suppose for each  $k \in \mathbb{N}$  that  $A_k$  is at most countable. Recall that since  $A_k$  is at most countable and therefore has an imposable total order letting us reference elements by index. Consider the function  $f : \mathbb{N} \times \mathbb{N} \rightarrow \bigcup_{k=1}^{\infty} A_k$  defined such that,  $f(n, m) = A_n[m]$ . Let  $a \in \bigcup_{k=1}^{\infty} A_k$  and by definition we know that  $a$  must exist in some set  $A_i$ , where  $i \in \mathbb{N}$ . Furthermore we can index the set  $A_i$  and let the index for  $a$  be  $j \in \mathbb{N}$ . Therefore we know that  $f(i, j) = A_i[j]$  where  $i, j \in \mathbb{N}$ . Thus  $f$  is a surjection, and it follows that since  $\mathbb{N} \times \mathbb{N}$  is countably infinite then  $\bigcup_{k=1}^{\infty} A_k$  is at most countable.  $\square$