

1. Prove that every ball $B_r(x)$ in a metric space (X, d) is an open set.

Solution:

Proof. Let $a \in B_r(x)$, and choose $\delta = r - d(x, a)$. Now consider $B_\delta(a)$ and note that to show $B_r(x)$ is open we must prove that $B_\delta(a) \subseteq B_r(x)$. Let $y \in B_\delta(a)$ and note that since (X, d) is a metric space the following expression comes from the triangle inequality,

$$d(x, y) \leq d(x, a) + d(a, y).$$

By the definition of $B_\delta(a)$ we know that $d(a, y) < r - d(x, a)$ and by substitution we get that $d(x, y) < r$. Thus by definition $y \in B_r(x)$ and therefore we get that $B_\delta(a) \subseteq B_r(x)$. Therefore $B_r(x)$ is open.

□

2. Let V be a subset of a metric space (X, d) . The set of limit points of V are those points x that can be written as the limit of a sequence of points in V . Show that a set $V \subseteq X$ is closed if and only if it contains its limit points.

Solution:

Proof. Suppose that $V \subseteq X$ is a closed set. By definition we know that V^c is open. For the sake of contradiction suppose V does not contain its limit points, therefore there exists a limit point of the set V , x such that $x \in V^c$. By the previous result there exists an open ball $B_r(x)$ such that $B_r(x) \subseteq V^c$. Note that since $x_n \in V$ and $x_n \rightarrow x$, we can choose n such that $d(x_n, x) < r$ and therefore $x_n \in B_r(x)$. Thus $x_n \in V^c$ a contradiction.

We will prove the reverse direction via contrapositive. Suppose V is not closed and therefore V^c is not open. Thus there exists some point $x \in V^c$ such that for all $r > 0$, $B_r(x) \not\subseteq V^c$. Choose the set $\{B_{\frac{1}{n}}(x) : i = \frac{1}{n}\}$ and by the previous result we know that inside each $B_{\frac{1}{n}}(x)$ there exists some $x_i \in V$. We will proceed by showing that $x_i \rightarrow x$. Let $\epsilon > 0$, and choose N such that $\frac{1}{N} < \epsilon$. Then for all $n \geq N$ we get $d(x_i, x) \leq \frac{1}{n} \leq \frac{1}{N} < \epsilon$. Therefore the sequence $\{x_i\} \in V$ converges to a limit point $x \in V^c$, so V does not contain all its limit points. □

3. Let d_1 and d_2 be two metrics on a set X . Show that the following conditions are equivalent.

- For every sequence $\{p_i\}_{i=1}^\infty$, if $p_i \xrightarrow{d_2} p$ then $p_i \xrightarrow{d_1} p$.
- For every function $f : X \rightarrow \mathbb{R}$, if f is continuous with respect to d_1 then f is continuous with respect to d_2 .
- For every set V , if V is closed with respect to d_1 then V is closed with respect to d_2 .
- For every set U , if U is open with respect to d_1 then U is open with respect to d_2 .

Solution:

1. ($a \iff b$) Suppose that for every sequence $\{p_i\}_{i=1}^\infty$, if $p_i \xrightarrow{d_2} p$ then $p_i \xrightarrow{d_1} p$. Let $f : X \rightarrow \mathbb{R}$ be a continuous function with respect to d_1 . Consider a sequence $\{p_i\}_{i=1}^\infty$ such that $p_i \xrightarrow{d_2} p$. By our supposition since $p_i \xrightarrow{d_2} p$ we know that $p_i \xrightarrow{d_1} p$. Since f is continuous with respect to d_1 , $p_i \xrightarrow{d_1} p$ implies that $f(p_i) \rightarrow f(p)$ in \mathbb{R} . Thus f is continuous with respect to d_2 .

For the converse we will make use of the following lemmas.

Lemma 1 (Reverse Triangle Inequality): Let X be a metric space, then for all $a, b, c \in X$

$$|d(a, c) - d(c, b)| \leq d(a, b).$$

Proof. Since X is a metric space, by the triangle inequality we know that

$$\begin{aligned} d(a, c) &\leq d(a, b) + d(c, b), \\ d(a, c) - d(c, b) &\leq d(a, b). \end{aligned}$$

Case 1: Suppose $d(a, c) - d(c, b) \geq 0$. Then it follows that $|d(a, c) - d(c, b)| \leq d(a, b)$.

Case 2: Suppose $d(a, c) - d(c, b) < 0$. By triangle inequality we also get that,

$$\begin{aligned} d(c, b) &\leq d(a, b) + d(a, c), \\ d(c, b) - d(a, c) &\leq d(a, b). \end{aligned}$$

Since $d(a, c) - d(c, b) < 0$ we know that $d(c, b) - d(a, c) \geq 0$ and therefore it follows that $|d(c, b) - d(a, c)| \leq d(a, b)$ or equivalently, $|d(a, c) - d(c, b)| \leq d(a, b)$. \square

Lemma 2: Let X be a metric space with a fixed point x^* . Then the function $f : X \rightarrow \mathbb{R}$ defined by $f(x) = d(x, x^*)$ is continuous.

Proof. Suppose $a \in X$ and let $\epsilon > 0$. Choose $\delta = \epsilon$ such that $d(x, a) < \delta$ implies that,

$$\begin{aligned} |f(x) - f(a)| &= |d(x, x^*) - d(a, x^*)|, \\ &\leq d(x, a), \\ &< \delta = \epsilon. \end{aligned}$$

\square

Suppose that for every function $f : X \rightarrow \mathbb{R}$, if f is continuous with respect to d_1 then f is continuous with respect to d_2 . Let $\{p_i\}_{i=1}^\infty$ be a sequence such that $p_i \xrightarrow{d_2} p$. Consider the function $f : X \rightarrow \mathbb{R}$ defined by $f(x) = d_1(x, p)$. By Lemma 2 we know that f is a continuous function with respect to d_1 and therefore by assumption f is continuous with respect to d_2 . Therefore it follows that since $p_i \xrightarrow{d_2} p$ we know that $d_1(x, p_i) \rightarrow 0$ which implies $p_i \xrightarrow{d_1} p$.

2. ($a \implies c$) Suppose that for every sequence $\{p_i\}_{i=1}^\infty$, if $p_i \xrightarrow{d_2} p$ then $p_i \xrightarrow{d_1} p$ and let V be a set that is not closed with respect to d_2 . Since V is not closed with respect to d_2 there exists a contact point $p \notin V$ where the sequence $\{p_i\}_{i=1}^\infty \in V$ $p_i \xrightarrow{d_2} p$. By assumption if $p_i \xrightarrow{d_2} p$ then $p_i \xrightarrow{d_1} p$ and since $\{p_i\}_{i=1}^\infty \in V$ and $p \notin V$, V is not closed with respect to d_1 .
3. ($c \implies d$) Suppose that for every set V , if V is closed with respect to d_1 then V is closed with respect to d_2 , and U is an open set with respect to d_1 . Note that if U is open with respect to d_1 then U^c is closed with respect to d_1 . It then follows that U^c is closed with respect to d_2 and thus U is open with respect to d_2 .
4. ($d \implies a$) Suppose that for every set U , if U is open with respect to d_1 then U is open with respect to d_2 and let $\{p_i\}_{i=1}^\infty \in X$ such that $p_i \xrightarrow{d_2} p$. Let $r > 0$ and consider an open ball $B_r^1(p)$. Since $B_r^1(p)$ is open with respect to d_1 , then by assumption $B_r^1(p)$ is open with respect to d_2 . It then follows that there exists some \hat{r} such that $B_{\hat{r}}^2(p) \subseteq B_r^1(p)$. Note that since $p_i \xrightarrow{d_2} p$ we can choose an N such that for all $n \geq N$, $p_n \in B_{\hat{r}}^2(x) \subseteq B_r^1(p)$. Thus $p_i \xrightarrow{d_1} p$.

Proof.

□

4. Let X be an infinite set.

a) Show that

$$\mathcal{T}_1 = \{U \subseteq X : U = \emptyset \text{ or } U^c \text{ is finite}\}$$

is a topology on X , called the finite complement topology.

Solution:

Proof. Note that by definition $\emptyset \in \mathcal{T}_1$ and since $X^c = \emptyset$ a finite set we get that $X \in \mathcal{T}_1$. Let $\{U_i\}_I$ be a set of open subsets of X . By definition we know that $\{U_i^c\}_I$ are all finite and therefore their intersection $\bigcap_{i \in I} U_i^c$ is also finite. By De Morgan's Law we conclude that,

$$\bigcap_{i \in I} U_i^c = \left(\bigcup_{i \in I} U_i \right)^c.$$

Thus $\cup_{i \in I} U_i$ is open. Let $\{U_i\}_I$ be a finite set of open subsets of X . Note that $\cup_{i \in I} U_i^c$ is a finite union of finite sets and is therefore also a finite set. By De Morgan's law we conclude that,

$$\bigcup_{i \in I} U_i^c = \left(\bigcap_{i \in I} U_i \right)^c.$$

Thus $\cup_{i \in I} U_i^c$ is open. □

b) Show that

$$\mathcal{T}_2 = \{U \subseteq X : U = \emptyset \text{ or } U^c \text{ is countable}\}$$

is a topology on X , called the countable complement topology.

Solution:

Proof. Note that by definition $\emptyset \in \mathcal{T}_2$. Since $X^c = \emptyset$ is a countable set we get that $X \in \mathcal{T}_2$. Let $\{U_i\}_I$ be a set of open subsets of X . By definition we know that $\{U_i^c\}_I$ are all countable and therefore their intersection $\cap_{i \in I} U_i^c$ is also countable. By De Morgan's Law we conclude that,

$$\bigcap_{i \in I} U_i^c = \left(\bigcup_{i \in I} U_i \right)^c.$$

Thus $\cup_{i \in I} U_i$ is open. Let $\{U_i\}_I$ be a finite set of open subsets of X . Note that $\cup_{i \in I} U_i^c$ is a finite union of countable sets which is also countable. Thus $\cup_{i \in I} U_i^c$ and by De Morgans law

$$\bigcup_{i \in I} U_i^c = \left(\bigcap_{i \in I} U_i \right)^c.$$

Thus $\cup_{i \in I} U_i^c$ is open. □

c) Let p be a fixed point in X , show that

$$\mathcal{T}_3 = \{U \subseteq X : U = \emptyset \text{ or } p \in U\}$$

is a topology on X , called the particular point topology.

Solution:

Proof. Note that by definition $\emptyset \in \mathcal{T}_3$. Since $p \in X$ we get that $X \in \mathcal{T}_3$. Let $\{U_i\}_I$ be a set of open subsets of X . By definition we know that each set in $\{U_i\}_I$ must be nonempty and exclude p and therefore either $\cap_{i \in I} U_i^c$ is empty or nonempty and excluding p . By De Morgans law we know that $(\cup_{i \in I} U_i)^c$ also excludes p and thus $p \in \cup_{i \in I} U_i$, an open set.

Let $\{U_i\}_I$ be a finite set of open subsets of X . Note that $\cup_{i \in I} U_i^c$ again is either empty or nonempty and excluding p . By De Morgans law we know that $\cup_{i \in I} U_i^c = (\cap_{i \in I} U_i)^c$ and therefore $p \in \cap_{i \in I} U_i$ an open set. □

d) Let p be a fixed point in X , show that

$$\mathcal{T}_4 = \{U \subseteq X : U = X \text{ or } p \notin U\}$$

is a topology on X , called the excluded point topology.

Solution:

Proof. Note that by definition $X \in \mathcal{T}_4$. Since $p \notin \emptyset$ we get that $\emptyset \in \mathcal{T}_4$. Let $\{U_i\}_I$ be a set of open subsets of X . By definition each set in $\{U_i\}_I$ must be nonempty and must include p , therefore either $\cap_{i \in I} U_i^c$ is empty, or nonempty and including p . By De Morgans law we know that $(\cup_{i \in I} U_i)^c$ also includes p and thus $p \notin \cup_{i \in I} U_i$, an open set.

Let $\{U_i\}_I$ be a finite set of open subsets of X . Again each $\{U_i^c\}_I$ must be nonempty and must include p , therefore either $\cup_{i \in I} U_i^c$ is empty, or nonempty and including p . By De Morgans law we know that $(\cap_{i \in I} U_i)^c$ also includes p and thus $p \notin \cap_{i \in I} U_i$, an open set. \square

e) Determine whether,

$$\mathcal{T}_5 = \{U \subseteq X : U = X \text{ or } U^c \text{ is infinite}\}$$

is a topology on X .

Solution:

For a counter example suppose \mathcal{T}_5 is a topology on \mathbb{R} . By definition the intervals $(-\infty, 0)$ and $(0, \infty)$ are open and therefore $A = (-\infty, 0) \cap (0, \infty)$ is open, yet $A \neq X$ and $A^c = 0$ is a finite set.

5. Let X be a set, and suppose $\{\mathcal{T}_\alpha\}_{\alpha \in A}$ is a collection of topologies on X . Show that the intersection $\mathcal{T} = \cap_{\alpha \in A} \mathcal{T}_\alpha$ is a topology on X .

Solution:

Proof. Let X be a set, and suppose $\{\mathcal{T}_\alpha\}_{\alpha \in A}$ is a collection of topologies on X . Note that for all $\alpha \in A$, $X, \emptyset \in \mathcal{T}_\alpha$ since each \mathcal{T}_α is a topology on X . So it follows $X, \emptyset \in \mathcal{T}$.

Now let $\{U_i\}_I$ be a set of subsets from \mathcal{T} . Note that by definition, for all $\alpha \in A$, $\{U_i\}_I \subseteq \mathcal{T}_\alpha$. Since \mathcal{T}_α are topologies on X it follows that $\cup_{i \in I} U_i \in \mathcal{T}_\alpha$. Thus $\cup_{i \in I} U_i \in \mathcal{T}$.

Now let $\{U_i\}_I$ be a finite set of subsets from \mathcal{T} . Note that by definition, for all $\alpha \in A$, $\{U_i\}_I \subseteq \mathcal{T}_\alpha$. Since \mathcal{T}_α are topologies on X it follows that $\cap_{i \in I} U_i \in \mathcal{T}_\alpha$. Thus $\cap_{i \in I} U_i \in \mathcal{T}$. \square