

**Exercise Supplemental 1:** Suppose  $(a_n) \rightarrow a$  and  $a \neq 0$ . Show that there exists  $N \in \mathbb{N}$  such that if  $n \geq N$ , then  $a_n \neq 0$ .

*Proof.* Suppose that the sequence  $(a_n) \rightarrow a$  and  $a \neq 0$ . Since the sequence  $(a_n)$  converges we know that for all,  $\epsilon \in \mathbb{R}$ , where  $\epsilon < 0$  there exists an  $N \in \mathbb{N}$  such that for all  $n \geq N$ ,

$$|a_n - a| < \epsilon.$$

Consider an  $\epsilon < a$  then there exists some  $N \in \mathbb{N}$  such that for all  $n \geq N$ ,

$$\begin{aligned} |a_n - a| &< \epsilon, \\ a - \epsilon &< a_n < a + \epsilon, \\ 0 &< a_n < a + \epsilon. \end{aligned}$$

Thus we have shown that there exists  $N \in \mathbb{N}$  such that if  $n \geq N$ , then  $a_n \neq 0$ .  $\square$

**Exercise Supplemental 2:** 1. Show that if  $a, b \geq 0$  and  $a > b$ , then  $\sqrt{a} > \sqrt{b}$ .

*Proof.* Let that  $a, b \geq 0$ , now suppose  $\sqrt{a} \leq \sqrt{b}$ . Through some algebra,

$$\begin{aligned} a &= \sqrt{a} \sqrt{a} \\ &\leq \sqrt{a} \sqrt{b} \\ &\leq \sqrt{b} \sqrt{b} \\ &= b \end{aligned}$$

Thus we have shown that  $a \leq b$ , and thus by contrapositive if  $a, b \geq 0$  and  $a > b$ , then  $\sqrt{a} > \sqrt{b}$ .  $\square$

2. Exercise 2.3.1(a) If  $(x_n) \rightarrow 0$ , show that  $\sqrt{(x_n)} \rightarrow 0$

*Proof.* Suppose the convergent sequence  $(x_n)$  such that  $(x_n) \rightarrow 0$ . Recall by the definition of convergent for all  $\epsilon > 0$  we know that there exists an  $N \in \mathbb{N}$  such that when  $n \geq N$ ,

$$|x_n| < \epsilon.$$

Note that since this inequality is true for all  $\epsilon > 0$ , its also true for  $\epsilon^2$  which leaves us with,

$$\begin{aligned} x_n &< \epsilon^2 \\ \sqrt{x_n} &< \epsilon \end{aligned}$$

Thus we have shown that  $\sqrt{(x_n)} \rightarrow 0$ .  $\square$

**Exercise 2.3.3:** Show that if  $x_n \leq y_n \leq z_n$  for all  $n \in \mathbb{N}$ , and if  $\lim x_n = \lim z_n = l$  then  $\lim y_n = l$  as well.

*Proof.* Suppose that  $x_n \leq y_n \leq z_n$  for all  $n \in \mathbb{N}$ , and that  $\lim x_n = \lim z_n = l$ . Let  $\epsilon > 0$ , since both  $x_n$  and  $z_n$  converge we know that there exists  $N_x, N_z \in \mathbb{N}$  such that for all  $n_x \geq N_x$ ,  $n_z \geq N_z$ , the following are true,

$$|x_{n_x} - l| \leq \epsilon$$

$$|z_{n_z} - l| \leq \epsilon$$

Now let  $N = \max\{N_x, N_z\}$ , to ensure that the above inequalities apply. Therefore for all  $n \geq N$ ,

$$-\epsilon < x_n - l < z_n - l < \epsilon.$$

Recall, that through algebra we get,

$$x_n \leq y_n \leq z_n,$$

$$x_n - l \leq y_n - l \leq z_n - l.$$

Therefore the following is true,

$$-\epsilon < x_n - l \leq y_n - l \leq z_n - l < \epsilon,$$

$$-\epsilon < y_n - l < \epsilon,$$

$$|y_n - l| < \epsilon.$$

Thus we have shown that  $\lim y_n = l$ . □

**Exercise 2.3.10:** Consider the following list of conjectures. Provide a short proof for those that are true and a counterexample for any that are false.

1. If  $\lim(a_n - b_n) = 0$ , then  $\lim a_n = \lim b_n$

*Proof.* Consider  $a_n = (-1)^{n+1}$  and  $b_n = (-1)^n$ . Clearly the following equation is true over all values of  $n$ ,

$$a_n - b_n = 0.$$

Therefore  $\lim(a_n - b_n) = 0$ , yet  $\lim a_n \neq \lim b_n$ . □

2. If  $(b_n) \rightarrow b$ , then  $|b_n| \rightarrow |b|$

*Proof.* Suppose  $(b_n) \rightarrow b$ . Consider that through the triangle inequality we know that (Exercise 1.2.6d),

$$|b_n - b| \geq ||b_n| - |b||.$$

Since  $(b_n) \rightarrow b$  we know that for all  $\epsilon > 0$  there exists an  $N \in \mathbb{N}$  such that for all  $n \geq N$ ,

$$|b_n - b| < \epsilon.$$

Thus it follows simply that,

$$||b_n| - |b|| < \epsilon.$$

Thus we have shown that  $|b_n| \rightarrow |b|$ . □

3. If  $(a_n) \rightarrow a$  and  $(b_n - a_n) \rightarrow 0$ , then  $(b_n) \rightarrow a$ .

*Proof.* Suppose  $(a_n) \rightarrow a$  and  $(b_n - a_n) \rightarrow 0$ . Rewriting the expression  $|b_n - a|$ ,

$$|b_n - a| = |b_n - a_n + a_n - a|.$$

By the triangle inequality,

$$|b_n - a_n + a_n - a| \leq |b_n - a_n| + |a_n - a|.$$

Since  $(a_n) \rightarrow a$  and  $(b_n - a_n) \rightarrow 0$  we know that for all  $\epsilon > 0$  there exists a  $N \in \mathbb{N}$  such that for all  $n \geq N$ ,

$$|a_n - a| < \frac{\epsilon}{2},$$

$$|a_n - b_n| < \frac{\epsilon}{2}.$$

Therefore,

$$\begin{aligned} |b_n - a| &\leq |b_n - a_n| + |a_n - a|, \\ &< \frac{\epsilon}{2} + \frac{\epsilon}{2}, \\ &< \epsilon. \end{aligned}$$

Thus we have shown that,  $(b_n) \rightarrow a$ . □

4. If  $a_n \rightarrow 0$  and  $|b_n - b| \leq a_n$  for all  $n \in \mathbb{N}$  then  $(b_n) \rightarrow b$ .

*Proof.* Suppose  $a_n \rightarrow 0$  and  $|b_n - b| \leq a_n$  for all  $n \in \mathbb{N}$ . Since  $a_n \rightarrow 0$  we know that for all  $\epsilon > 0$  there exists an  $N \in \mathbb{N}$  such that for all  $n \geq N$ ,

$$|a_n - 0| = |a_n| < \epsilon.$$

Therefore we chain these inequalities and get,

$$|b_n - b| \leq |a_n| < \epsilon$$

Thus we have shown that,  $(b_n) \rightarrow b$ . □

**Exercise Supplemental 3:** Show that if  $|b_n| \rightarrow 0$ , then  $b_n \rightarrow 0$ . Then show that this statement is false if we replace 0 with any other real number.

*Proof.* Suppose that  $|b_n| \rightarrow 0$ , Therefore for all  $\epsilon > 0$  there exists an  $N \in \mathbb{N}$  such that  $n \geq N$  where,

$$||b_n| - 0| = ||b_n|| = |b_n| = |b_n - 0| \leq \epsilon.$$

Thus we have shown that if  $|b_n| \rightarrow 0$ , then  $b_n \rightarrow 0$ .

$$||b_n| - a| = ||b_n|| = |b_n| = |b_n - a| \leq \epsilon.$$

□

$$||b_n| - b| \geq ||b_n| - |b||.$$

**Exercise 2.3.10 (c):**

**Exercise C:** Consider the series  $\sum_{n=1}^{\infty} 1/n^2$ . Give a careful proof by induction that the partial sums

$$s_k = \sum_{n=1}^k 1/n^2$$

satisfy  $s_k < 2 - 1/k$ .

**Exercise 2.4.3(a):** Hint: Use the Monotone Convergence Theorem!