Computational Physics Laboratory report

Stefano Ge

Winter semester 2025

Contents

0 Introduction **2**

1 Error analysis **2**
- 1.0.1 Approximating exponentials . . . 2
- 1.2 Floating-point arithmetic and roundoff errors . . . 3
  - 1.2.1 Computing the Basel problem . . . 3
- 1.3 Error propagation and condition number . . . 5
  - 1.3.1 Computing statistical momenta . . . 5
  - 1.3.2 Condition number: study of a simple algorithm . . . 5

2 Linear systems **5**
- 2.1 Forward- and back-substitution . . . 5
- 2.2 LUP Decomposition . . . 6
- 2.3 Conditioning of linear systems . . . 7
- 2.4 Overdetermined linear systems . . . 8
  - 2.4.1 Least squares through normal equations . . . 8
  - 2.4.2 Fitting of Kepler's Law . . . 8
  - 2.4.3 Fitting of exp(sine) function . . . 9
- 2.5 QR decomposition . . . 11
  - 2.5.1 Least squares through QR . . . 11
  - 2.5.2 QR decomposition for eigenvalues and eigenvectors . . . 11

3 Interpolation **11**
- 3.1 Polynomial interpolation by solving the Vandermonde matrix linear system . . . 11
- 3.2 Lagrange-Waring interpolation . . . 12
  - 3.2.1 Evenly spaced nodes . . . 12
  - 3.2.2 Chebyshev nodes . . . 12
  - 3.2.3 Chebyshev vs evenly spaced nodes: error study . . . 13
  - 3.2.4 Polynomial interpolation over the real line with Chebyshev nodes . . . 13
  - 3.2.5 Chebyshev nodes interpolation of non-smooth functions . 13

4 Roots of nonlinear equations **14**
4.1 Bisection method 14
4.2 Newton method 15
4.2.1 Convergence rate study 15
4.2.2 Which root gets found by Newton's method? 17
4.2.3 Eros' true anomaly 17
4.3 Secant method 18
4.4 Inverse quadratic interpolation 19

5 Numerical integration **19**
5.1 Newton-Cotes formulas 19
5.1.1 Trapezoidal rule 20
5.1.2 Simpson's rule 21
5.2 Gauss-Legendre rule 22
5.2.1 Nodes and weights of Gauss-Legendre rule 22
5.2.2 Integrals with Gauss-Legendre rule 22
5.3 Double exponential rule 24
5.3.1 Comparison of double exponential and Gauss-Legendre rules 24
5.3.2 Improper integrals with double exponential rule 26

6 Ordinary differential equations **27**
6.1 Euler's method 27
6.1.1 First order application 27
6.1.2 Second order application 27
6.2 Runge-Kutta methods 27
6.2.1 Improved Euler's and Runge-Kutta fourth order methods 27
6.2.2 Non-linear equation application of Runge-Kutta 27
6.2.3 Predator-prey model 27

7 Schrödinger equation **27**
7.1 Time-dependant Schrödinger equation 27
7.1.1 Gaussian packet in an infinite well 27
7.1.2 Moving gaussian packet 28
7.1.3 Finite barrier scattering 28
7.1.4 Pöschl-Teller potential 28
7.2 Time-independant Schrödinger equation 28

0 Introduction

1 Error analysis

1.0.1 Approximating exponentials

Results




Figure 1: Truncation errors when approximating e^x through the truncated power series Σ_0^N. N is the degree of the polinomyal used for the approximation. The plot of $x^{N+1}/(N+1)$ has been added for reference.

Remarks

- Horner's method was used in polynomial evaluation, because it is faster and reduces roundoff errors. A naive implementation of a polynomial computation requires more operations ($O(N^2)$) and, for small values of x, adds together numbers of order 1 and x^N, while Horner's method is $O(N)$ and keeps the additions only between terms of order 1 and x.
- By Taylor's theorem, if $\hat{f}(x)$ is the power series of e^x truncated after the N-th term, the error is of order $O(x^{N+1})$ as x approaches 0. Computing an additional term of the series shows that:

$$|e^x - \hat{f}(x)| = \frac{x^{N+1}}{(N+1)!} + o(x^{N+1})$$

  It is then not surprising that the approximation error in Figure 1 is indistinguishable from $x^{N+1}/(N+1)$ for small values of x. For larger values, the contribution of higher order terms in the series becomes ever more important, and hence the divergence from $x^{N+1}/(N+1)$.

1.2 Floating-point arithmetic and roundoff errors

1.2.1 Computing the Basel problem

Results

Remarks




Figure 2: Approximation error when computing $\pi^2/6$ with the Basel problem truncated series $\Sigma_0^N \frac{1}{n^2}$, with single precision floating variables. Here N is the number of terms used when calculating the series.




Figure 3: Same as above, but now using double precision floating variables. the plot of $1/x$ has been added for reference.

- The algorithm has been implemented with two mathematically equivalent forms, which, however, resulted in remarkably different numerical results. In the first the terms of the series are summed in increasing order of the summation index, i.e. decreasing of the value of the addends. Since small addends are summed to the accumulating partial sum, it gets to a point where the next addend is smaller than $\epsilon_{mach} \cdot sum \approx \epsilon_{mach}$, so the addition of it and subsequent values does not change the value of the partial sum.
  This can be seen in the error plots in Figure 2 and 3, since the error stops decreasing and remains constant after $1/\mathrm{N}^2 = \epsilon_{\mathrm{mach}}$, which is 2^{-24} for single precision ($\mathrm{N} = 4096$) and 2^{-53} for double precision ($\mathrm{N} \approx 9.5 \times 10^7$).

  In the second implementation the terms of the series are summed in decreasing order of the summation index, i.e. increasing the value of the addends. Therefore, the aforementioned effect is not seen, and the numerical results correspond to the mathematical expectation, in the range considered.

- The truncation error of this algorithm is asymptotically equivalent to k/N, with k real constant. This can be from the integral associated to $\Sigma_{n=0}^{\infty}$:

$$\int_N^{\infty} \frac{1}{n^2} = \frac{1}{N}$$

  therefore,

$$\Sigma_{n=N}^{\infty} \sim_{N\to\infty} \frac{1}{N}$$

  which is confirmed from the agreement between the computed values and the $1/x$ reference line in Figure 3.

1.3 Error propagation and condition number

1.3.1 Computing statistical momenta

1.3.2 Condition number: study of a simple algorithm

2 Linear systems

2.1 Forward- and back-substitution

Results

$$\begin{bmatrix} -2 & 0 & 0 \\ 1 & -1 & 0 \\ 3 & 2 & 1 \end{bmatrix} \mathbf{X} = \begin{bmatrix} -4 \\ 2 \\ 1 \end{bmatrix} \qquad \mathbf{X} = \begin{bmatrix} 2 \\ 0 \\ -5 \end{bmatrix}$$

$$\begin{bmatrix} 4 & 0 & 0 & 0 \\ 1 & -2 & 0 & 0 \\ -1 & 4 & 4 & 0 \\ 2 & -5 & 5 & 1 \end{bmatrix} \mathbf{X} = \begin{bmatrix} -4 \\ 1 \\ -3 \\ 5 \end{bmatrix} \qquad \mathbf{X} = \begin{bmatrix} -1 \\ -1 \\ 0 \\ 2 \end{bmatrix}$$

$$\begin{bmatrix} 3 & 1 & 0 \\ 0 & -1 & -2 \\ 0 & 0 & 3 \end{bmatrix} \mathbf{X} = \begin{bmatrix} 1 \\ 1 \\ 6 \end{bmatrix} \qquad \mathbf{X} = \begin{bmatrix} 2 \\ -5 \\ 2 \end{bmatrix}$$

$$\begin{bmatrix} 3 & 1 & 0 & 6 \\ 0 & -1 & -2 & 7 \\ 0 & 0 & 3 & 4 \\ 0 & 0 & 0 & 5 \end{bmatrix} \mathbf{X} = \begin{bmatrix} 4 \\ 1 \\ 1 \\ 5 \end{bmatrix} \qquad \mathbf{X} = \begin{bmatrix} -3.\overline{3} \\ 8 \\ -1 \\ 1 \end{bmatrix}$$

The results obtained through my routine were verified using LAPACK.

2.2 LUP Decomposition

Results

$$\begin{bmatrix} 2 & 3 & 4 \\ 4 & 5 & 10 \\ 4 & 8 & 2 \end{bmatrix} = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 1 & 1 & 0 \\ 0.5 & 0.\overline{3} & 1 \end{bmatrix} \begin{bmatrix} 4 & 8 & 2 \\ 0 & -3 & 8 \\ 0 & 0 & 0.\overline{3} \end{bmatrix}$$

Determinant is 4.

$$\begin{bmatrix} 6 & -2 & -4 & 4 \\ 3 & -3 & -6 & 1 \\ -12 & 8 & 21 & -8 \\ -6 & 0 & -10 & 7 \end{bmatrix} = \begin{bmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0.5 & 1 & 0 & 0 \\ -0.25 & 0.25 & 1 & 0 \\ -0.5 & -0.5 & -0.857143 & 1 \end{bmatrix} \begin{bmatrix} -12 & 8 & 21 & -8 \\ 0 & -4 & -20.5 & 11 \\ 0 & 0 & 4.375 & -3.75 \\ 0 & 0 & 0 & 2.28571 \end{bmatrix}$$

Determinant is -480.

$$\begin{bmatrix} 1 & 4 & 5 & -5 \\ -1 & 0 & -1 & -5 \\ 1 & 3 & -1 & 2 \\ 1 & -1 & 5 & -1 \end{bmatrix} = \begin{bmatrix} 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ 1 & 0.8 & 1 & 0 \\ -1 & -0.2 & -0.\overline{6} & 1 \end{bmatrix} \begin{bmatrix} 1 & -1 & 5 & -1 \\ 0 & 5 & 0 & -4 \\ 0 & 0 & -6 & 6.2 \\ 0 & 0 & 0 & -2.\overline{6} \end{bmatrix}$$

Determinant is 80.

Matrix $\mathbf{A} = \mathbf{T}(3, -1)\mathbf{R}(\pi/5)\mathbf{T}(-3, 1)$:

$$A = \begin{bmatrix} 2.3 & 1.6 & -2.3 \\ -1.1 & 0.48 & 2.7 \\ 0.5 & 0.\overline{3} & -0.\overline{6} \end{bmatrix}$$

$$\mathbf{b} = \mathbf{Az} = \mathbf{A} \begin{bmatrix} 2 \\ 2 \\ 1 \end{bmatrix} = \begin{bmatrix} 5.45 \\ 1.51 \\ 1.5 \end{bmatrix}$$

Factorization of $\mathbf{A}$ is:

$$\mathbf{A} = \begin{bmatrix} 1 & 0 & 0 \\ -0.471 & 1 & 0 \\ 0.217 & -0.00857 & 1 \end{bmatrix} \begin{bmatrix} 2.31 & 1.59 & -2.34 \\ 0 & 1.22 & 1.64 \\ 0 & 0 & 0.354 \end{bmatrix}$$

2.3 Conditioning of linear systems

Results In the case study proposed, first we construct numerically:

$$\mathbf{b} = \mathbf{A}\mathbf{x} = \begin{bmatrix} -\epsilon & 1 \\ 1 & -1 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

Where ϵ is a real parameter. We then apply the LU decomposition (without pivoting) to solve for x given $\mathbf{A}$ and the $\mathbf{b}$ just computed.

For $\epsilon = 10^{-12}$ we find:

$$\mathbf{A} = \mathbf{L}\mathbf{U} = \begin{bmatrix} 1 & 0 \\ -1\text{e}{-12} & 1 \end{bmatrix} \begin{bmatrix} -\text{e}{-12} & 1 \\ 0 & 1\text{e}12 \end{bmatrix}$$

$$\mathbf{x}_{\text{computed}} - \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} -2.2\text{e}{-5} \\ 0 \end{bmatrix}$$

As a test, we find correctly:

$$\mathbf{A} - \mathbf{L}\mathbf{U} = \mathbf{0}$$

However, if we set $\epsilon = 10^{-12}$, we find:

$$\mathbf{A} = \mathbf{L}\mathbf{U} = \begin{bmatrix} 1 & 0 \\ -1\text{e}{-20} & 1 \end{bmatrix} \begin{bmatrix} -\text{e}{-20} & 1 \\ 0 & 1\text{e}20 \end{bmatrix}$$

$$\mathbf{x}_{\text{computed}} - \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} -1 \\ 0 \end{bmatrix}$$

$$\mathbf{A} - \mathbf{L}\mathbf{U} = \begin{bmatrix} 0 & 0 \\ 0 & -1 \end{bmatrix}$$

Remarks

- The exact LU-factorization of $\mathbf{A}$ is:

  $$\mathbf{A} = \begin{bmatrix} 1 & 0 \\ -\epsilon^{-1} & 1 \end{bmatrix} \begin{bmatrix} -\epsilon & 1 \\ 0 & \epsilon^{-1} - 1 \end{bmatrix}$$

  If $\epsilon = 10^{-20} \ll \epsilon_{\text{mach}}$, in the bottom-right entry of $\mathbf{U}$ only ϵ^{-1} gets stored, since the relative variation of adding -1 to ϵ^{-1} is less than ϵ_{mach}. But when $\mathbf{L}$ and $\mathbf{U}$ get multiplied together, in the addition resulting into the bottom-right entry of $\mathbf{A}$ two ϵ^{-1} terms cancels out, and the exact result is the -1 that is lost in numerical LU-decomposition.

- Exact $\mathbf{b}$ is:

  $$\begin{bmatrix} 1 - \epsilon \\ 0 \end{bmatrix}$$

  but ϵ does not get registered if it is lower than ϵ_{mach}. Because the condition number of the matrices $\mathbf{L}$ and $\mathbf{U}$ is very high (ϵ^{-2} in ∞-norm), the tiny variation of $\mathbf{b}_{\text{calc}}$ over $\mathbf{b}_{\text{exact}}$ is allowed to propagate tremendously and results in the wrong result seen before.

  Doing the calculations shows that the error on $\mathbf{U}$ as no impact on $\mathbf{b}_{\text{calc}}$, and that the final error on $\mathbf{b}_{\text{calc}}$ is due entirely on the ill-conditioning of the linear systems $\mathbf{L}\mathbf{z} = \mathbf{b}$ and $\mathbf{U}\mathbf{x} = \mathbf{z}$.

- By contrast, the condition number of the starting matrix $\mathbf{A}$ is only ≈ 4. Therefore, in principle it should be possible to solve the system for $\mathbf{x}$ even if $\mathbf{b}$ is slightly nudged: the problem is the stability of the algorithm adopted.

  LU-decomposition without row-pivoting is much less numerically stable than LUP-decomposition with row-pivoting where the maximum value entry in a column is selected as the pivot.

  Indeed, if we perform LU-decomposition on the following matrix, obtained by permuting the two rows of matrix $\mathbf{A}$, as prescribed by row-pivoting, we find the correct results:

$$\mathbf{PA} = \begin{bmatrix} 1 & -1 \\ -\epsilon & 1 \end{bmatrix} = \mathbf{LU} = \begin{bmatrix} 1 & 0 \\ -1\text{e}{-20} & 1 \end{bmatrix} \begin{bmatrix} ta1 & -1 \\ 0 & 1 \end{bmatrix}$$

$$\mathbf{x}_{\text{computed}} - \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\mathbf{A} - \mathbf{LU} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$$

2.4 Overdetermined linear systems

2.4.1 Least squares through normal equations

Results After implementing a routine for finding a the vector $\mathbf{X}$ that best fits an overdetermined linear system, we tested it on the following linear system:

$$\begin{bmatrix} 2 & -1 \\ 0 & 1 \\ -2 & 2 \end{bmatrix} \mathbf{X} = \begin{bmatrix} 1 \\ -5 \\ 6 \end{bmatrix}$$

and found:

$$\mathbf{X} = \begin{bmatrix} -2 \\ 1 \end{bmatrix}$$

2.4.2 Fitting of Kepler's Law

We want to extrapolate Kepler's third law from experimental data through the algorithm just introduced. First we linearize the following relationship by taking the logarithm at both sides:

$$\text{T} = \text{CD}^{\alpha} \implies \log(\text{T}) = \log(\text{C}) + \alpha \log(\text{D})$$

where T is the orbital periods of the planets, D is their average distance from the Sun and alpha and C are two costants to be determined through the method of least squares. This is equivalent to finding x_1 and x_2 that minimize:

$$\log(\mathbf{T}) = x_1\mathbf{1} + x_2 \log(\mathbf{D}) = \begin{bmatrix} \mathbf{1} & \log(\mathbf{D}) \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

where quantities in bold are vectors, leading to an overdetermined linear system.

Results We therefore find that the values that fit the aforementioned system are:

$$\alpha = 1.49865 \qquad C = 0.201522$$

where alpha is consistent with Kepler's law, which predicts $\alpha = 1.5$.

The graph containing the fitted curve and experimental data can be found in Figure 4




Figure 4: Orbital peirods vs distances from the Sun of the planets.

2.4.3 Fitting of exp(sine) function

Results We want to approximate the function $\exp(\sin(t-1))$, first through a Taylor-like polynomial:

$$y(t) = c_1 + c_2 t + ... + c_7 t^6$$

and then through a periodic Fourier-like approximation:

$$y(t) = d_1 + d_2 \cos(t) + d_3 \sin(t) + d_4 \cos(2t) + d_5 \sin(5t)$$

Results are shown in Figure 5 and in Figure 6.

Remarks

- Not surpringly, the function is better approximated by the first terms of the Fourier basis than by a sixth degree polynomial. The Fourier basis is an orthonormal complete set of $L^2([0, 2\pi])$, therefore it is guaranteed to be the best approximation in L^2 norm, among expansions on sets spanned by it. Moreover, sines and cosines are themselves periodic and are thus well suited for approximating a periodic function like $\exp^{\sin(t-1)}$, producing a nice approximation over the whole domain $\mathbb{R}$. On the other hand, monic




Figure 5: Least square fit of $\exp(\sin(t-1)$ with 6th degree polynomial




Figure 6: Least square fit of $\exp(\sin(t-1)$ with Fourier periodic functions

monomyals, though complete, are neither an orthonormal set of $L^2([0, 2\pi])$ nor are well suited for approximating periodic functions, as they do blow up at infinity and do not approximate well outside of the points employed in the fit.

2.5 QR decomposition

In this section we implement algorithms to find the QR decomposition of any matrices. First the modified Gram-Schmidt procedure (MGS).

Results We test the modified Gram-Schmidt procedure on the following matrices:

$$\begin{bmatrix} 2 & -1 \\ 0 & 1 \\ -2 & 2 \end{bmatrix} = \begin{bmatrix} 0.707 & 0.408 \\ 0.000 & 0.816 \\ -0.707 & 0.408 \end{bmatrix} \begin{bmatrix} 2.828 & -2.121 \\ 0 & 1.225 \end{bmatrix}$$

$$\begin{bmatrix} 2 & 3 & 4 \\ 4 & 5 & 10 \\ 4 & 8 & 2 \end{bmatrix} = \begin{bmatrix} 0.333 & -0.104 & 0.937 \\ 0.667 & -0.677 & -0.312 \\ 0.667 & 0.729 & -0.156 \end{bmatrix} \begin{bmatrix} 6.000 & 9.667 & 9.333 \\ 0 & 2.134 & -5.726 \\ 0 & 0 & 0.312 \end{bmatrix}$$

$$\begin{bmatrix} 2 & -1 & 4 \\ 0 & 3 & 5 \\ 7 & 2 & -6 \\ 1 & -4 & 0 \end{bmatrix} = \begin{bmatrix} 0.272 & -0.241 & 0.733 \\ 0.000 & 0.559 & 0.617 \\ 0.953 & 0.179 & -0.233 \\ 0.136 & -0.773 & 0.163 \end{bmatrix} \begin{bmatrix} 7.348 & 1.089 & -4.627 \\ 0 & 5.368 & 0.752 \\ 0 & 0 & 7.418 \end{bmatrix}$$

Computing the product $\mathbf{A} = \mathbf{QR}$ gives us the matrices we have decomposed, confirming the correctness of the decomposition. Also we computed $\mathbf{Q}^\intercal\mathbf{Q}$ and we confirmed that the computed $\mathbf{Q}$'s are indeed ONC matrices.

2.5.1 Least squares through QR

We implemented QR algorithm to solve least-squares problem, and applied it to the problem of fitting the function $\exp^{\sin(4t)}$ with a 14th degree polynomial.

Results

2.5.2 QR decomposition for eigenvalues and eigenvectors

3 Interpolation

3.1 Polynomial interpolation by solving the Vandermonde matrix linear system

Interpolating polynomials in the power basis by constructing the Vandermonde matrix and solving the related linear system is numerically unstable. Tha matrix thus constructed is likely to be ill-conditioned, due to the fact that its entries are the nth powers of x, and can become quickly extremely large or small as the system scales in size.

In our case study, we considered equidistant point in the interval [-1,1] generated by the function $\cos(kx)$.

In Figure AGGIUNGI REFERENCE we can see that the condition number scales exponentially with the number of points considered in the fit, which is also the size of the linear system.

AGGIUNGI FIGURA

In our study we always considered $4n + 1$ equidistant points, including the extrema of the interval -1 and 1. As a consequence, the middle point considered is always (0,1), since $\cos(0) = 1$. Therefore, we must have:

$$P_n(0) = a_0 + a_1 \cdot 0 + ... + a_n \cdot 0 = a_0 = 1$$

CERCA DI CAPIRE COME MOSTRARE I VALORI

If we were operating in exact arithmetic, this would produce a value of a_0 always equal to 1, since it would lead to a linear system where a row of the Vandermonde matrix (the one corresponding to point x=0) which is equivalent to equation $a_0 = 1$. However, the LUP decomposition algorithm contaminates several times the aforementioned row of the linear system, relying later on cancellation of possibly large values to recover the information of $a_0 = 1$. Therefore, in finite-precision arithmetic the value of a_0 can differ from 1, as we can see ... . This is closely related to the ill-conditioning of the Vandermonde matrix.

3.2 Lagrange-Waring interpolation

3.2.1 Evenly spaced nodes

Results In Figure ADD 4 GRAPHS AND REFERENCES we show the polynomials resulting from the Lagrange-Waring interpolation of the the functions $\log(x)$, $tanh(x)$, $cosh(x)$ and $|x|$, using equally spaced nodes.

The weights of the points were evaluated through the analytic formula for equally spaced weights. Some lines of code have been added to the code in order to show that the special formula agrees with the generic one.

Remarks

- The Lagrange-Waring interpolation implementation has been organised around a struct, `BarFit`, which stores points, weights, etc.

  This OOP-style design is flexible, and allows reusage of the weights already calculated for other functions.

3.2.2 Chebyshev nodes

Results We have tested our routine for Lagrange interpolation using Chebyshev nodes of the second kind by fitting the function $cosh(sin(x))$ using 40 points in the interval $[0, 2\pi]$.

The interpolating polynomial, together with the nodes, can be visualized in ADD REFERENCE

In the testing program we also included a a comparison between the weights calculated through the analytical formula for Chebyshev nodes and the generic nodes formula. It confirmed that the weights are the same aside from roundoff errors.

3.2.3 Chebyshev vs evenly spaced nodes: error study

We study the convergence of Lagrange intrpolation wit equidistant an Chebyshev points for the functions $1/(25x^2 + 1)$, $\tanh(5x + 2)$, $\cosh(\sin(x))$ and $\sin(\cosh(x))$ over the inerval [-1:1].

The results are shown in ADD REFERENCS AND FIGURES

Remarks

- Interpolation theory has proven that, for analytical functions bounded in the interpolation interval, $||f - p||_\infty \leq CK^n$ for some constants C and K, i.e. that Lagrange's interpolation with Chebyshev nodes converges at least exponentially. In our plot we see this exact pattern. We can even estimate the constant K:

f(x)	K
$1/(25x^2 + 1)$	1.22
$\tanh(5x + 2)$	1.39
$\cosh(\sin x)$	5.70
$\sin(\cosh x)$	4.81

- When we use equidistant points as nodes convergence is not guaranteed for every analytical and bounded functions. Instead, as the number of nodes grows, $||f - p||_\infty$ can grow exponentially, that is Runge's phenomenon. We observe this phenomenon in our plot.

3.2.4 Polynomial interpolation over the real line with Chebyshev nodes

By means of a change of variable we now interpolate the function $1/(z^2-2z+2)$ over an unbounded interval [-∞, +∞]. We used Chebyshev nodes of the first kind to avoid evaluation at x = $\pm\infty$.

As change of variable we used the function $\phi(x) = \frac{2x}{1-x^2}$ which maps the interval [-1, 1] to the real line.

Results are shown in ADD REFERENCE AND FIGURE

3.2.5 Chebyshev nodes interpolation of non-smooth functions

Polynomial interpolants on Chebyshev nodes are guaranteed to converge exponentially to the reference function if the latter is analytic, as the number of nodes approaches infinity. If it is not, convergence might not be guaranteed or it might be slower.

We can observe this slowdown by studying the convergence for the family of functions $|x|^m$. These functions are not smooth on the interval [-1, 1]. They can be differentiated $m - 1$ times, but not m times, since $|m|$ is not differentiable in $x = 0$. Hence they are not analytic. The results are shown in ADD REFERENCE AND FIGURES

Remarks

- As previously said, $|x|^m$ is not analytical. However, their derivatives are continuously differentiable up to $f^{(m-2)}$, thus absolutely continuous. $f^{(m-1)} = C|x|$, which is Lipschitz continuous and as such it is absolutely continuous. On the other hand:

$$f^{(m)}(x) = \begin{cases} +C, & x > 0 \\ -C, & x < 0 \end{cases}$$

  for some C>0. This function is monotonic, hence it is of bounded variation. This proves a theorem that states that if the above hypotheses are verified (absolute continuity up to $m-1$-th derivative and bounded variation of m-th derivative), then the Chebyshev polynomial interpolant converges to the function in sup-norm as $\mathcal{O}(n^{-m})$, as the number of nodes n approaches infinity.

- To compare this prediction with our results, we interpolated the sup-norm error using algorithm from ADD REFERENCES By choosing $Y = \log_{10}(||f-p||_\infty) = \log_{10}(c) - K\log_{10}(n)$ and $X = \log_{10}(n)$ we interpolated the relation:

$$Y = C - K\,X$$

  where -K is the exponent in the asymptotic behaviour $\mathcal{O}(n^{-m})$ of $||f-p||_\infty$. The fitting lines can are visualised in ADD REFERENCE as dashed lines. Our results are reported in the following table and are in accordance to the prediction above:

m	K
1	1.04
3	3.13
5	5.15
7	7.26
9	9.41
11	11.7

4 Roots of nonlinear equations

4.1 Bisection method

In order to test our routine for rootfinding with the bisection method, we searched for the roots of the function $x^3 - 7x^2 + 14x - 6$ in the interval [0, 1] and found $x = 0.5858$. The plot (which we omitted from the report, but can be visualized during code execution) of the function and the found root confirms that it is the only root in the interval.

Given an iterative rootfinding algorithm, if we define $\epsilon_n = x_n - r$, where r is the root that we are going to find and x_n is the root estimate at step n, then if

$$\lim_{n\to+\infty} \left| \frac{\epsilon_{n+1}}{\epsilon_n^q} \right| = C \tag{1}$$

for $q \geq 1$ and $0 < C < \infty$, we say that the algorithm converges to the root with order of convergence q and asymptotic constant C.

It is very easy to see that for the bisection algorithm $q = 1$ and $C = \frac{1}{2}$. Under this assumption, we get the asymptotic relation

$$|\epsilon_{n+1}| \approx |\frac{1}{2}\epsilon_n|$$

which is the asymptotic recurrence relation of a geometric sequence $\epsilon_n = K(\frac{1}{2})^n$.

Defining $a_n = -\log_{10}(|\epsilon_n|)$ we get:

$$a_n = K' + n\log_{10}(2)$$

In ADD FIGURE AND REFERENCE we plotted the a_n as a function of n. The curve is a line, confirming that the order of convergence is 1. Fitting the curve with a line through least squares yields a slope of 1.992, which implies an asymptotic error constant $C = 0.5018$, in accordance with bisection method's convergence theory.

As a termination condition of the loop in the algorithm we check if:

$$|x_1 - x_2| <= \text{tol} + \epsilon_{\text{mach}} \max(|a|, |b|)$$

Where $|a|$ and $|b|$ are the boundaries of the starting interval and x_1, x_2 are the boundaries of the interval at current step.

`tol` can be specified by the caller or instead be 10ϵmach if the caller inserts the macro `DEF_TOL` as parameter.

$\epsilon_{\text{mach}} \max(|a|, |b|)$ is added because it is approximately the distance between the largest of the two starting boundaries (in absolute value). The algorithm is not always able to bisect the algorithm further when it is of the order of the quantity above, so it is a safety check.

4.2 Newton method

4.2.1 Convergence rate study

We seek to check the correctness of our routine and study the convergence rate of Newton's method by applying it to the functions $f_1(x) = x^2 - \exp(-x)$, $f_2(x) = \tan(x) - 2x$ and $f_3(x) = \exp(x + 1) - x - 2$.

By calculating $a_n = -log_{10}|x_n - r|$ we can calculate how many digits of precision we are gaining at each step, and estimate order of convergence and asymptotic error constant.

Results In ADD FIGURES AND REFERENCES we plotted the functions and the roots found.

In table 1 we quote the roots found and the numerical estimates for the values of q and C.

q is estimated as the asymptotic value of a_{n+1}/a_n, since it holds $a_{n+1} \sim qa_n$.

C is estimated as

$$|\epsilon_{n+1}/\epsilon_n^q|$$

where we use for q the value that we calculated analytically (see Remarks).

f(x)	Root	≈q	≈C
$x^2 - \exp(-x)$	0.703	2.08	0.40
$\tan(x) - 2x$	0.000	3.07	0.67
	1.166	1.92	3.38
$\exp(x+1) - x - 2$	-1.000	1.06	0.50

Remarks

- If a root r of a function $f(x)$ satisfies $f'(r) \neq 0$ and $f''(r) \neq 0$, then Newton's method's convergence order is quadratic for that root and the asymptotic error constant is:

$$C = \frac{1}{2}\left|\frac{f''(r)}{f'(r)}\right|$$

  That is the case for the root of f_1 and for root $x = 1.166$ of f_2. In both cases the estimated convergence order is close to 2 and the estimated asymptotic error constant agrees with the one calculated analitically by 4+ significant digits. Hence, we do not include separately the analitical values in the report and we refer the reader to the program `newton_method_error_study.c`.

- If the root is not simple, i.e. $f'(r) = 0$, but $f''(x) \neq 0$, then it can be shown that convergence is linear with asymptotic error constant $C = 0.5$. Root the root of f_3 satisfies this condition. In our analysis we see that most ratios a_{n+1}/a_n agree wih 0.5, except for the last few values.

  This is the same order of convergence and asymptotic error constant of bisection method. However, in bisection method only a single function evaluation is required at each step, while in Newton method both the function and its derivative must be evaluated. Thus, in the case $f'(r) = 0$, for the same precision gained at each step, Newton method is twice as expensive, compared to bisection method, if we assume the evaluation of the derivative to be comparable to evaluation of the function in computational cost.

- For root $r = 0$ of f_3 the estimated order of convergence is about 3. Seeing that for this function $f''(r) = 0$, this prompted us to check with a small computation whether $f''(r) = 0$ implies order of convergence 3. We start from the relation:

$$\epsilon_{k+1} = \epsilon_k - \frac{f(r+\epsilon_k)}{f'(r+\epsilon_k)}$$

  which is a rewrite of Newton's algorithm. Expanding with Taylor's:

$$\epsilon_{k+1} = \epsilon_k - \frac{f(r) + f'(r)\epsilon_k + \frac{1}{2}\epsilon_k^2 f''(r) + \frac{1}{6}\epsilon_k^3 f'''(r) + O(\epsilon_k^4)}{f'(r) + f''(r)\epsilon_k + \frac{1}{2}\epsilon_k^2 f'''(r) + O(\epsilon_k^3)} =$$

$$= \epsilon_k - \frac{f'(r)\epsilon_k + \frac{1}{6}\epsilon_k^3 f'''(r) + O(\epsilon_k^4)}{f'(r) + \frac{1}{2}\epsilon_k^2 f'''(r) + O(\epsilon_k^3)} =$$

Collecting $f'(r)$:

$$= \epsilon_k \left[1 - \left(1 + \frac{1}{6}\epsilon_k^2 \frac{f'''(r)}{f'(r)} \right) \left(1 + \frac{1}{2}\epsilon_k^2 \frac{f'''(r)}{f'(r)} \right)^{-1} \right] + O(\epsilon_k^4) =$$

By doing some simple algebra:

$$= \frac{1}{3}\epsilon_k^3 \frac{f'''(r)}{f'(r)} + O(\epsilon_k^4)$$

i.e. cubic convergence, with asymptotic error constant $C = \frac{1}{3}f'''(r)/f'(r)$. By executing `newton_method_error_study`, the reader can see that the numerically estimated and the analitically calculated one agree by 4 significant digits, after only 6 steps have been taken by Newton's method.

4.2.2 Which root gets found by Newton's method?

Newton method, in general, is not globally convergent. Moreover, if two or more roots are present, if convergent there is no guarantee that the root that gets found is the one "closest" to the starting value for the algorithm.

In this paragraph we study the relationship between the starting value and the root that gets found by the method. We use as case study the function $f(x) = x^{-2} + \sin(x)$ and as starting values for the algorithm $x = 1, 2, ..., 7$.

In ADD FIGURE AND REFERENCES we have plotted $f(x)$, along with the starting values and the related roots that get found.

We see that the initial guesses 2 and 5 produce return roots that are not the closest to the starting points. In order to understand what is going on, we have drawn the tangent line to the graph at each of those two initial guesses. Newton method approximates the function to that line at the first iteration. We can see from the graph the tangent lines drawn intersect the x axis "far away" from the starting guess, since at both points the function has a slope small in absolute value.

If we include bracketing in our algorithm, then we can be sure that the root that gets found will be inside of the interval from which we started.

ADD FIGURE AND REFERENCE shows the root found inside the arbitrality chosen intervals [0.5, 2], [2, 5], [5, 8] and [8, 10].

4.2.3 Eros' true anomaly

As an application of Newton method, we seek to evaluate the *true anomaly* $\theta(t)$ of asteroid Eros, defined as:

$$\tan\frac{\theta}{2} = \sqrt{\frac{1+\epsilon}{1-\epsilon}} \tan\frac{\psi}{2}$$

For that, we need to evaluate first the *eccentric anomaly* $\psi(t)$ which can be found by solving the equaton:

$$f(\psi) = \psi - \epsilon \sin\psi - \frac{2\pi t}{\tau} = 0 \tag{2}$$

For asteroid Eros, we have *orbial period* $\tau = 1.7610$ and *elliptical eccentricity* $\epsilon = 0.2230$.

Derivative of equation 2 with respect to ψ is:

$$f'(\psi) = 1 - \epsilon \cos \psi$$

Since $\epsilon < 1$, $f'(\psi) > 0$ for any ψ, the function is monotonic and we need not worry about finding multiple roots. Also, by plotting function 2 we see that it is reasonably well-behaved, and it might be even reasonable to expect global convergence. Nonetheless, we can also simply observe that for ψ such that $f(\psi) = 0$:

$$\left| \psi - \frac{2\pi t}{\tau} \right| = |\epsilon \sin \psi| \leq \epsilon$$

that is, if we safeguard Newton method with bracketing and we take as starting interval $[\frac{2\pi t}{\tau} - \epsilon, \frac{2\pi t}{\tau} + \epsilon]$, then we are guaranteed to find the root.

In ADD FIGURE AND REFERENCES we have plotted the graph of $\theta(t)$ thus obtained. $\theta(t)$ was chosen to be in the interval $[0, 2\pi]$ to ensure continuity.

4.3 Secant method

We seek to test our implementation of the secant method and study its convergence rate by searching the roots of the same functions examined in 4.2.1.

Results In ADD FIGURES AND REFERENCES we plotted the functions and the roots found.

In table 1 we quote the roots found and the numerical estimates for the values of q, estimated as in 4.2.1.

f(x)	Root	≈q
$x^2 - \exp(-x)$	0.703	1.65
$\tan(x) - 2x$	0.000	1.86
	1.166	1.58
$\exp(x+1) - x - 2$	-1.000	1.03

The values of q quoted above must not be taken to be asymptotically accurate estimates. In contrast with our analysis of Newton method, the estimated values were not settling around a value. With our precision we likely were not able to reach accurately the asymptotic behaviour; yet our rough estimation confirms a rate of convergence between linear and quadratic for the first 3 roots examined. In the case of $f_3(x)$, $f'(r) = 0$, therefore convergence is not guaranteed to be superlinear and appears to be linear instead.

Secant method is the inverse interpolation method with degree 1 We prove the above statement as an exercise.

Suppose that we have calculated the approximations x_k and x_{k-1} for a root of the function $f(x)$ by using the inverse interpolation method of $m = 1$. Next approximation is $q(0)$, where $q(x)$ is a polynomial of degree 1 such that $q(f(x_k) = x_k$ and $q(f(x_{k-1}) = x_{k-1}$. But a polynomial of degree 1 is just a line. We can write it explicitely as:

$$q(y) - x_k = \frac{x_k - x_{k-1}}{f(x_k) - f(x_{k-1})}(y - f(x_k))$$

By substituting 0 in place of y in the previous expression we find $x_{k+1} = q(0)$. We see that it is the same x_{k+1} that would be calculated by the secant method.

4.4 Inverse quadratic interpolation

We now repeat the same study with the inverse quadratic interpolation method.

Results In ADD FIGURES AND REFERENCES we plotted the functions and the roots found.

In table 1 we quote the roots found and the numerical estimates for the values of q, estimated as in 4.2.1.

f(x)	Root	≈q
$x^2 - \exp(-x)$	0.703	1.6-2.0
$\tan(x) - 2x$	0.000	1.6-2.0
	1.166	1.6-1.8
$\exp(x+1) - x - 2$	-1.000	1.04

We confirm again an order of convergence between 1.6 and 2.0.

5 Numerical integration

5.1 Newton-Cotes formulas

In the following paragraphs we will test different integration rules on the integrals listed down here:

1) $f_1(x)$: $\int_0^1 x \log(1+x)\, dx = \frac{1}{4}$

2) $f_2(x)$: $\int_0^1 x^2 \tan^{-1} x\, dx = \frac{\pi - 2 + 2\log 2}{12}$

3) $f_3(x)$: $\int_0^{\pi/2} e^x \cos x\, dx = \frac{e^{\pi/2} - 1}{2}$

4) $f_4(x)$: $\int_0^1 \frac{\tan^{-1}(\sqrt{2+x^2})}{(1+x^2)\sqrt{2+x^2}} dx = \frac{5\pi^2}{96}$

5) $f_5(x)$: $\int_0^1 \sqrt{x} \log(x)\, dx = -\frac{4}{9}$

6) $f_6(x)$: $\int_0^1 \sqrt{1-x^2}\, dx = \frac{\pi}{4}$

We will sometimes refer to the definite integrals above simply by the integrand functions.

5.1.1 Trapezoidal rule

Results In ADD FIGURE AND REFERENCES, we plot the error ΔI between the exact value of the integrals and the approximation obtained using the trapezoidal rule, as a function of the number of subintervals used.

Since the spacing between the nodes h is inversely proportional to the number of nodes n, if ΔI is asymptotically $O(h^m)$, then it must also be asymptotically $O(\frac{1}{n^m})$.

Given this, we can estimate m by linearizing the asymptotic relationship between ΔI and n. m will be the opposite of the slope of the line thus obtained. In Table 4 we list the values of m found for every integral.

f	m
$f_1(x)$	2.00
$f_2(x)$	2.00
$f_3(x)$	2.00
$f_4(x)$	2.00
$f_5(x)$	1.37
$f_6(x)$	1.50

Table 4

Remarks

- All integrals except those of functions f_5 and f_6 to the exact value at second order with respect to the subinterval length. Indeed, the integrand functions are all smooth in the integration interval. This assures by expanding with Taylor that the truncation error is quadratic in h as h goes to zero.

- f_5 is not differentiable in 0, hence the linear interpolation fails at producing a "good" approximation of the function in $x = 0$. We can try to estimate the truncation error analitically. We will consider only the subinterval $[0, \epsilon]$, since in all others the function is smooth and will converge quadratically. We have:

$$\begin{aligned} R &= \int_0^\epsilon \sqrt{x}\log(x)\mathrm{d}x - \frac{\epsilon^{\frac{3}{2}}}{2}\log(\epsilon) = \\ &= \left[\frac{2}{3}\epsilon^{\frac{3}{2}}\log(x)\right]_0^\epsilon - \frac{2}{3}\int_0^\epsilon \sqrt{x}\mathrm{d}x - \frac{\epsilon^{\frac{3}{2}}}{2}\log(\epsilon) = \\ &= \epsilon^{\frac{3}{2}}\log(\epsilon)\left(\frac{1}{6}\log(\epsilon) - \frac{4}{9}\right) \end{aligned}$$

  Hence the truncation error is $O(h^{\frac{3}{2}}\log(h))$. Note that as h goes to zero, $\log(|R|) \sim m\log(\epsilon)$, hence the estimated m value should be asymptotically 1.5.

- f_5 is not differentiable in 1. As in previous point we can try to estimate

the truncation error analitically:

$$
\begin{aligned}
R &= \int_{1-\epsilon}^{1} \sqrt{1-x^2}\mathrm{d}x - \frac{\epsilon}{2}\sqrt{1-(1-\epsilon)^2} = \\
&= \int_{0}^{\epsilon} \sqrt{2y-y^2}\mathrm{d}y - \frac{\epsilon}{2}\sqrt{2\epsilon-\epsilon^2} = \\
&= \int_{0}^{\epsilon} \sqrt{2y}(1+O(y))\mathrm{d}y - \frac{\epsilon}{2}\sqrt{2\epsilon}(1+O(y)) = \\
&= \frac{2\sqrt{2}}{3}\epsilon^{\frac{3}{2}} - \frac{\sqrt{2}\epsilon^{\frac{3}{2}}}{2} + O(y^{\frac{5}{2}}) = \\
&= \frac{\sqrt{2}}{6}\epsilon^{\frac{3}{2}} + O(\epsilon^{\frac{5}{2}})
\end{aligned}
$$

Hence the truncation error is $O(h^{\frac{3}{2}})$, as hinted in the Results paragraph.

- Since the trapezoidal rule approximates the integrand function with linear interpolations, intervals in which the function is convex produce underestimates, while intervals where the function is concave produce overstimated integrals. In fact, f_1, f_2 and f_5 are convex within the bounds of integration: the error is negative. On the other hand, f_3 and f_6 are concave: the error is positive.

5.1.2 Simpson's rule

Exactly like in previous section, we test our implementation of Simpson's rule on the functions in 5.1.

In ADD FIGURE AND REFERENCES we show again the truncation error as a function of the number of subintervals used.

In Table 5 we report the estimated values for the orders of m.

f	m
$f_1(x)$	4.00
$f_2(x)$	4.00
$f_3(x)$	4.00
$f_4(x)$	4.01
$f_5(x)$	1.40
$f_6(x)$	1.50

Table 5

The first four functions are smooth in the integration interval, hence for Simpson's rule fourth-order convergence is guaranteed.

Calculations analogous to those in previous section would show that the truncation errors of $f_5(x)$ and $f_6(x)$ are, respectively, $O(h^{\frac{3}{2}\log(h)}$ and $O(h^{\frac{3}{2}}$.

One might point out that the estimated value of m for function $f_5(x)$ is very close to $\frac{7}{2}$. It is a coincidence. If we changed the program `simpson_rule.c` in order to remove some initial points from the fit, we would see that the value of m would get closer and closer to 1.5.

5.2 Gauss-Legendre rule

5.2.1 Nodes and weights of Gauss-Legendre rule

In this section we designed and tested some routines for evaluating Legendre polynomials, their roots and the weights in Gauss-Legendre integration.

Results In Table 6 we report the values of roots and weights Legendre polynomials with degree 2–4 that we found with our routines. In the program `gauss_legendre_weights.c` we also compare them to the exact values, to which they agree with an error lower than the tolerance requested in the rootfinding routine.

In ADD FIGURE AND PLOT we also show the plot of the first 6 Legendre polynomials calculated with our routines.

Degree	Root index	Root value	Weight
2	1	0.577	1.000
	2	-0.577	1.000
3	1	0.775	0.556
	2	0.000	0.889
	3	-0.775	0.556
4	1	0.861	0.348
	2	0.340	0.652
	3	-0.340	0.652
	4	-0.861	0.348

Table 6

Remarks

- For evaluating the nth Legendre polynomial for $L_n(x)$, the values for all lower degrees must be computed recursively for the same x. As a micro-optimization, these are stored inside an `static` array. Since `static` arrays in C are conserved between function calls, this saves time if $L_m(x)$ for the same x is requested in the following function call. Our routine will compute values of $L_n(x)$ only if they are not already stored, otherwise it will use the values already present to evaluate a higher degree polynomial. The array size count resets only when the caller asks for a different value of x.

- For this reason, evaluating derivatives is extremely cheap and justifies the use of Newton method for finding the roots. The algorithm has complexity $O(nk)$, where n is the degree of the polynomial and k is the number of steps in Newton method. It should be a low number, reasonably independent from n, and most likely less than 10.

5.2.2 Integrals with Gauss-Legendre rule

We now test our routine for calculating integrals with Gauss-Legendre rule and we study the convergence rate. We compute the integrals of the functions:

1) $\int_{-1}^{1} e^{-4x}\mathrm{d}x$

2) $\int_{-1}^{1} e^{-9x^2}\mathrm{d}x$

3) $\int_{-1}^{1} \mathrm{sech}(x)\mathrm{d}x$

4) $\int_{-1}^{1} \frac{1}{1+9x^2}\mathrm{d}x$

5) $\int_{\pi/2}^{\pi} x^2 \sin(8x)\mathrm{d}x$

In ADD FIGURE AND REFERENCE we plot the error in integral estimation as a function of the degree of the interpolant polynomial.

The convergence is very fast and is exponential or superexponential. But why do some functions appear to converge much faster than others?

The remainder term in Gauss-Legendre rule is guaranteed to be:

$$R_P[f] = \frac{2^{2n+1}(n!)^4}{(2n+1)[(2n)!]^3} f^{(2n)}(\xi) \qquad \xi \in (-1,1) \tag{3}$$

where f is the integrand function after we have changed variables in the integration interval.

The function $f_4 = \frac{1}{1+9x^2}$ converges as the slowest. In fact:

$$\begin{aligned}
\frac{\mathrm{d}^n}{\mathrm{d}x^n}\frac{1}{1+9x^2} &= 3^n \frac{\mathrm{d}^n}{\mathrm{d}y^n}\frac{1}{1+y^2} = \\
&= \frac{(-1)^n 3^n n!}{2i}\left(\frac{1}{(y-i)^{n+1}} - \frac{1}{(y+i)^{n+1}}\right) \\
&= \frac{(-1)^n 3^n n!}{2i(y^2+1)^{\frac{n+1}{2}}}\left(e^{i(n+1)\theta} - e^{-i(n+1)\theta}\right) \\
&= \frac{(-1)^n 3^n n!}{(y^2+1)^{\frac{n+1}{2}}}\sin\left((n+1)\arctan\left(\frac{1}{y}\right)\right) \\
&= \frac{(-1)^n 3^n n!}{(9x^2+1)^{\frac{n+1}{2}}}\cos\left((n+1)\arctan 3x - \frac{n\pi}{2}\right)
\end{aligned}$$

Since we took $\theta = \arctan(\frac{1}{y})$, this technically would be valid only if $\theta \in [-\frac{\pi}{2}, \frac{\pi}{2}]$, i.e. if $y = 3x > 0$, but this will not matter since all derivatives are either even or odd, and we are interested only in the absolute value of them. Plugging this into Equation 3 we find, after some simplifications:

$$R_P[f] \approx k\left(\frac{3}{2}\right)^{2n}(-1)^n \frac{\cos\left((2n+1)\arctan(3\xi)\right)}{(9\xi^2+1)^{n+\frac{1}{2}}} \qquad \xi \in (-1,1)$$

where k is a constant factor. This... is not great help, since, depending on the value that ξ takes, the above expression can be, in absolute value, anything between 0 and $\left(\frac{3}{2}\right)^{2n}$. The point is that in this case the remainder theorem does not guarantee a superexponential convergence as we will see in other cases, and it has to do with the the magnitude of the derivatives of $\frac{1}{1+3x^2}$.

Nonetheless, f_4 is analytic and bounded in a Bernstein ellipse E_ρ with $\rho < 3$. Therefore convergence is guaranteed to be exponential, but it is not faster than that, as we can see from our plot.

With $f(x) = e^{-4x}$ the calculations are easier and convergence is much faster. We have $f^{(n)}(x) = (-4)^n e^{-4x}$. Plugging into equation 3 we find the remainder:

$$\begin{aligned} R_P[f] =& \approx \frac{\sqrt{n} e^{-4\xi}}{(n!)^2} \qquad \xi \in (-1, 1) \\ & \leq k \frac{\sqrt{n} e^4}{[(n)!]^2} \end{aligned}$$

With k constant. That is, we have squared-factorial convergence guaranteed!

5.3 Double exponential rule

5.3.1 Comparison of double exponential and Gauss-Legendre rules

We now test our routine for the double-exponential rule on the integrals:

1) $\int_0^1 \sqrt{x} \log(x) \mathrm{d}x = -\frac{4}{9}$

2) $\int_0^1 \sqrt{1 - x^2} \mathrm{d}x = \frac{\pi}{4}$

3) $\int_1^0 (\log x)^2 \mathrm{d}x = 2$

4) $\int_{\pi/2}^{\pi} \log(\cos x) \mathrm{d}x = -\frac{\pi}{2} \log 2$

5) $\int_{\pi/2}^{\pi} \sqrt{\tan x} \mathrm{d}x = \frac{\pi}{\sqrt{2}}$

and we study the the error as a function of n, comparing it with that of the Gauss-Legendre rule. Here n is the number of nodes in the Gauss-Legendre rule and the number of subintervals in the double-exponential rule.

In ADD FIGURE AND REFERENCES we plot the error as a function of n.

Remarks

- Our routine is highly modular in the sense that it is possible to inject into `double_exp_base` the double exponential diffeomorphism suited for the needs of the caller.

  In `intg_double_exp` it is possible to choose between four different diffeomorphisms that we implemented, one for bounded intervals, the other two for the whole real line and half line.

- The double-exponetial rule maps the integration interval I into the real line, and calculates the resulting integral with the trapezoidal rule. Determing where to truncate the latter integral requires some care.

  Let $t_M > 0$ so that the transformed integral is truncated to the interval $[-t_M, t_M]$. In order to determine t_M, our routine sets initially its value to 0.1, and then adds 0.1 until both $|g(t_M)|$ and $|g(-t_M)|$ are lower than a

certain `tol`, where g is the integrand function of the transformed integral. We also check against `NaN` that could be returned by some functions on singular points.

Since we are dealing with double exponential functions, which grow extremely fast, it is necessary to safeguard for overflow and underflow cases. Even for just $t_M = 6.6$, $e^{e^{t_M}} \approx 10^{319}$, which overflows double precision floating representation. The sinh-tanh quadrature rule is even more delicate. As t becomes "big" (when dealing with double exponentials "big" numbers are reached very fast...) we have:

$$\phi(t) = \tanh\left(\frac{\pi}{2}\sinh t\right) \approx 1 - 2e^{-\frac{\pi}{2}e^t}$$

when $t_M = 3.2$, $2e^{-\frac{\pi}{2}e^{t_M}} \approx 3.7 \times 10^{-17} < \epsilon_{\text{mach}}$, therefore we have an underflow and $\psi(t_M) = 1$. A proper and implementation should take into account the possibilities of underflow and overflow for every diffeomorphism used. However, due to the overall design and time constraints, we only check the overflow for integrals over real line, and we leave to the caller the responsibility to guard against overflows in the tanh-sinh case, by adjusting the `tol` required and therefore allowing the routine to settle on a lower value of t_M.

- All the integrand functions of the integrals considered are either infinite or non differentiable at the bounds of integration. Unsurprisingly, convergence is slow with the Gauss-Legendre rule. In fact, all functions have unbounded derivatives, and functions 3-5 are not even bounded themselves. Although the integrals appear to converge nonetheless, it is yet very slow and likely algebraic.

  On the contrary, the approximations with double-exponential quadrature rule converge exponentially with n. For this rule, fast convergence is guaranteed even if the integrand is not analytic at the limits of integration, as it is the case for the integrand functions above.

- The only exceptions appear to be f_3 and f_5, which, after a phase of exponential convergence, the errors appear to stop shrinking and start growing slowly at 10^{-8} and 10^{-13}. The problem here is not the theoretical convergence of double-exponential rule, but the truncation error due to considering the transformed integral only in $[-t_M, t_M]$. In these two cases, indeed, we had to lower manually the tolerance `tol` in the routine for determining t_M, as otherwise it would push for a value of t_M too high, resulting in an underflow. Setting a high tolerance in truncation determination has of naturally the effect of adding a constant error which cannot be lowered by adding more points.

  A smart way to achieve more accuracy for the integral of function f_5 is to use a change of variables:

$$\int_0^{\frac{\pi}{2}} \sqrt{\tan(x)}\mathrm{d}x = \int_0^{\infty} \frac{\sqrt{y}}{1+y^2}\mathrm{d}y$$

  As observed above, the double-exponential rule for infinite intervals is more safeguarded from the risk of overflow. A quick function call in the

program shows indeed that the estimate reaches ϵ_{mach} levels of precision by using just $n = 36$ nodes.

5.3.2 Improper integrals with double exponential rule

We now test our implementation of the double-exponential rule on integrals over infinite intervals, and study the convergence:

1) $\int_{-\infty}^{\infty} \frac{1}{1+x^2+x^4}\,\mathrm{d}x = \frac{\pi}{\sqrt{3}}$

2) $\int_{-\infty}^{\infty} e^{-x^2}\cos(x)\,\mathrm{d}x = e^{-1/4}\sqrt{\pi}$

3) $\int_{-\infty}^{\infty} (1+x^2)^{-2/3}\,\mathrm{d}x = \frac{\sqrt{\pi}\,\Gamma(1/6)}{\Gamma(2/3)}$

4) $\int_{0}^{\infty} \frac{1}{1+x^2}\,\mathrm{d}x = \frac{\pi}{2}$

5) $\int_{0}^{\infty} \frac{e^{-x}}{\sqrt{x}}\,\mathrm{d}x = \sqrt{\pi}$

For integrals 1-3 we use as change of variables:

$$I = \int_{-\infty}^{\infty} f(x)\,\mathrm{d}x \qquad\qquad x = \sinh\left(\frac{\pi}{2}\sinh(t)\right)$$

For integral 4 we use:

$$I = \int_{0}^{\infty} f(x)\,\mathrm{d}x \qquad\qquad x = \exp\left(\frac{\pi}{2}\sinh(t)\right)$$

For integral 5 we use:

$$I = \int_{0}^{\infty} f(x)e^{-x}\,\mathrm{d}x \qquad x = \exp\left(t - \exp(-t)\right)$$

after putting $f(x) = \sqrt{x}$.

In ADD FIGURE AND REFERENCES we show the plot of the absolute value of the error as a function of the number of function evaluations n. Despite oscillations, the errors appears to decay linearly with n in the semi-log scale, for all the above integrals. We then conclude that it shrinks exponentially. The oscillations are due to the fact that the transformed integrand functions have varying concavity, producing therefore both positive and negative errors that can "accidentally" cancel out, resulting in a numerical estimate more precise than the subsequent estimate with more nodes.

6 Ordinary differential equations

6.1 Euler's method

6.1.1 First order application

6.1.2 Second order application

6.2 Runge-Kutta methods

6.2.1 Improved Euler's and Runge-Kutta fourth order methods

6.2.2 Non-linear equation application of Runge-Kutta

6.2.3 Predator-prey model

7 Schrödinger equation

In the following paragraphs we solve the one-dimensional Time-Dependant Schrödinger Equation (TDSE) and Time-Independant Schrödinger Equation (TISE).

To discretize the Hamiltonian, we first discretize space in equally spaced nodes and use the following finite difference approximation for the second derivative:

$$\left.\frac{d^2\psi}{dx^2}\right|_{x_j} \approx \frac{-\psi_{j+2} + 16\psi_{j+1} - 30\psi_j + 16\psi_{j-1} - \psi_{j-2}}{12(\Delta x)^2} + \mathcal{O}(\Delta x^4)\,.$$

We use Dirichlet boundary condition, that is, we consider fixed at 0 the wavefunction outside the simulation interval. This is equivalent to putting the system inside an infinite potential well.

7.1 Time-dependant Schrödinger equation

For solving TDSE, we use a fourth-order Runge-Kutta algorithm specialized for time-independant potentials. This time discretization is equivalent to a fourth order expansion of time evolution operator. In natural units:

$$\begin{aligned}
\langle\psi(t+\mathrm{d}t)| &= U(\mathrm{d}t)\,\langle\psi(t)| \\
&= e^{-i\mathrm{H}\mathrm{d}t}\,\langle\psi(t)| \\
&= \left(\mathrm{I} + i\mathrm{H}\mathrm{d}t - \frac{1}{2}\mathrm{H}^2\mathrm{d}t^2 - i\frac{1}{6}\mathrm{H}^3\mathrm{d}t^3 + \frac{1}{24}\mathrm{H}^4\mathrm{d}t^4\right)\langle\psi(t)|
\end{aligned}$$

7.1.1 Gaussian packet in an infinite well

We first test our algorithm by studying time evolution of a free Gaussian wave packet with average momentum 0. The result can be visualised in the `plot` folder of the program as `.gif` animation file. Considering only the square modulus of the wave function, as expected the packet first spreads out, its peak falling, while its tails expanding. Then it gets reflected back when it hits the walls of the well, giving rise to standing waves.

Note that in our test $\Delta x = 0.1$ and $\Delta t = 0.01$ satisfies the stability condition for the free case

$$\Delta t \lesssim= \frac{m(\Delta x)^2}{\hbar}$$

Raising Δt even so slightly as to 0.02 causes the solution to explode.

Norm conservation Schrödinger equation preserves probabilities over time, and we want to check how good our numerical solver reproduces this property. To evaluate the square norm we just integrate the square modulus of the wave function with a midpoint rule, considering even the values at the endpoints since the function should ideally continue beyond them if we were not to truncate space.

First we evaluate the square norm of the packet at time $t = 0$. We find 0.995. This is sensibly different than the expected value 1. Is it because the quadrature rule that we used was not accurate enough with these many grid points or we cut off the tails of the gaussian? Definitely the latter, as repeating the calculation with a grid 100 times larger yields a square norm value of 0.99995.

Using $|\langle\psi|\,|^2 = 0.995$ as norm reference value, we can compare at each time the difference between the square norm at that time and the reference value. At time $t = 20$ in time evolution the difference is maximum at 0.006, i.e. 0.6%. Our numerical solver then preserves quite well probabilities for the parameters and time scales provided.

7.1.2 Moving gaussian packet

7.1.3 Finite barrier scattering

7.1.4 Pöschl-Teller potential

7.2 Time-independant Schrödinger equation