

Linking environmental stochasticity with animal space use using  
continuous-time stochastic processes

Master's Thesis

Stefano Mezzini

April 2023

***NOTES:***

- refer to the style guideline at <https://github.com/jepa/ubcdown>
- UBC-V style: <https://faculty.washington.edu/mforbes/projects/ubcthesis/>
- TU Delft, Netherlands, template <https://www.tudelft.nl/huisstijl/downloads>
- contact Sunil and Kristy Baxter once I've made a template

The following individuals certify that they have read, and recommend to the College of Graduate Studies for acceptance, a thesis/dissertation entitled: Click or tap here to enter text. submitted by Click or tap here to enter text. in partial fulfillment of the requirements of the degree of Click or tap here to enter text. .

Click or tap here to enter text. Supervisor Click or tap here to enter text. Supervisory Committee Member Click or tap here to enter text. Supervisory Committee Member Click or tap here to enter text. University Examiner Click or tap here to enter text. External Examiner

Additional Committee Members include: Click or tap here to enter text. Supervisory Committee Member Click or tap here to enter text. Supervisory Committee Member.

# Abstract

## Lay Summary

## Preface

# Contents

<b>Abstract</b>	<b>ii</b>
<b>Lay Summary</b>	<b>iii</b>
<b>Preface</b>	<b>iv</b>
<b>Acknowledgements</b>	<b>viii</b>
<b>Dedication</b>	<b>ix</b>
<b>1 Chapter 1: Introduction</b>	<b>1</b>
1.1 Timescale of stochasticity . . . . .	1
1.2 Objectives . . . . .	2
<b>2 Chapter 2: Two-Eyed Seeing</b>	<b>4</b>
2.1 Importance and recognition of Traditional Indigenous Knowledge . . . . .	4
<b>3 Chapter 3: Movement analyses</b>	<b>6</b>
3.1 Movement simulations . . . . .	6
3.2 Stochasticity map . . . . .	6
3.3 Movement analysis . . . . .	6
<b>4 Chapter (?): Conclusion</b>	<b>7</b>
<b>Bibliography</b>	<b>8</b>
<b>Appendix 1</b>	<b>11</b>

## List of Tables



## List of Figures

- 1 Simulations depicting the effects of resource availability and stochasticity on spatial needs. Animals moved from the circles to nearby tiles until satiated. The labels indicate how many steps animals took to reach satiety. Note the higher spatial needs of animals in more unpredictable or resource-scarce environments. . . . . 2

## Acknowledgements

## Dedication

# 1 Chapter 1: Introduction

animals with short lives (e.g. mice) will be impacted more by a stochastic year than larger animals (e.g. elephant) See figures in *Southwood 1977*

- spend time reading papers initially to do a review, get a review paper out of it
- make predictions/hypotheses for project
- what influences stochasticity: what is it and what is affecting it
- very little testing has been done on effects of stochasticity

Environmental productivity is tightly linked to the amount of space that animals need to cover to obtain the resources they needed to survive and reproduce (Lucherini & Lovari, 1996; Relyea, Lawrence & Demarais, 2000). While resource availability is often considered in conservation decision-making, an environment's heterogeneity, stochasticity, and how the two change over time are rarely accounted for. In addition, environmental stochasticity, including extreme events, can reduce a landscape's energetic balance (Chevin, Lande & Mace, 2010), which, in turn, would decrease animals' fitness. Therefore, we expect animals living in unpredictable environments to require more space than those in stable environments (Fig. 1). Although this hypothesis is supported by a few recent studies (Morellet *et al.*, 2013; Nandintsetseg *et al.*, 2019; Riotte-Lambert & Matthiopoulos, 2020), many of them are limited in their analytical depth and geographic and taxonomic scales, so there remains a need for developing a more complete understanding of how animals' spatial needs change with environmental stochasticity. These stresses are compounded by climate change, which exposes species to increasingly common stochastic events (IPCC, 2018). Furthermore, anthropogenic structures reduce the habitat available to terrestrial species (Wilson *et al.*, 2016), who struggle to move in fragmented (Fahrig, 2007), human-dominated landscapes (Tucker *et al.*, 2018). As the impacts of habitat loss and climate change will worsen in the future (Hansen *et al.*, 2013; IPCC, 2018), it is imperative that we better understand spatial requirements of taxa to protect wildlife existence and biodiversity. Environmental safeguarding is also essential for Reconciliation with Indigenous People in Canada (Truth and Reconciliation Commission of Canada, 2015).

## 1.1 Timescale of stochasticity

Organisms are most affected by stochastic events and processes which occur on time scales which are shorter than the organism's life span (ref?). Weekly heavy rains which alter a lake's salinity (ref?) are more likely to affect the lake's inhabitants than a multi-centennial drought, and high-salinity conditions may be perceived as the (stressful) standard by organisms which were born during periods of drought. However, stochastic




Figure 1: Simulations depicting the effects of resource availability and stochasticity on spatial needs. Animals moved from the circles to nearby tiles until satiated. The labels indicate how many steps animals took to reach satiety. Note the higher spatial needs of animals in more unpredictable or resource-scarce environments.

processes and events which occur on time scales that are longer than an organism's lifespan may still cause significant effects on a population's fitness and stability. Droughts which occur on the time scale of centuries or millennia (Haig *et al.* (2013)) are unlikely to affect organisms directly, but such events could still alter the population's habitat or breeding grounds enough to cause a population collapse or prevent individuals from reproducing in their habitual breeding grounds (or reproduce altogether).

For an event or process to be recognized as deterministic by an individual, it must occur multiple times during the individual's lifetime (but the converse is not true). *some animals can develop memory* (Foley, Pettorelli & Foley, 2008)

Trees that have drought resilience have higher mortality (DeSoto *et al.*, 2020) *check the stats & causation from this reference*

## 1.2 Objectives

This project will produce a global raster of a new environmental stochasticity index, and new quantitative methods for animal movement. Findings will inform design of protected areas and assist in conserving Canada's wildlife, particularly in light of Canada's pledge to protect 30% of its landmass and oceans by 2030(Ryan Patrick Jones, 2021), although only 13% of Canada's area is currently protected(Government of

Canada, 2021). Local Indigenous groups will be included in the research process, in the hope of forming long-lasting co-operative relationships.

I will use simulation studies and an unprecedented and conservation-relevant animal tracking dataset (>1500 animals, 77 globally-distributed species) to provide the most detailed investigation into how animal spatial needs change with environmental stochasticity to date. This work has four key objectives: (i) estimating individuals' spatial requirements in a way which is insensitive to variation in sampling protocols and data quality; (ii) quantifying environmental stochasticity and its effects; (iii) estimating between-species trends using models that are robust to commonly-found issues (e.g., correlations within species); and (iv) understanding how Traditional Indigenous Knowledge can be integrated into large-scale ecological research and conservation planning within a framework that acknowledges both Traditional Indigenous Knowledge and Western science(Kutz & Tomaselli, 2019).

To achieve these aims: (i) I will use continuous-time models (Johnson *et al.*, 2008) that do not depend on sampling frequency. Such models will allow me to use the entirety of the data rather than aggregated data (e.g., daily averages), as aggregated data contain less information on sample variance and can lead to biased estimates. (ii) I will produce a global, time-varying raster of environmental stochasticity which accounts for productivity(Nilsen, Herfindal & Linnell, 2005), weather, and climate (e.g., precipitation, temperature), as well as the frequency of extreme events (e.g., heat domes, flooding, forest fires). I will then use the raster to estimate the effect of stochasticity on home range sizes. (iii) I will use a hierarchical approach(Pedersen *et al.*, 2019) to estimate common trends and variances within and between populations, species, and data collection methods. (iv) I will collaborate closely with various Indigenous groups and include any Traditional Knowledge and practices they wish to include in my project. The research will be published in open-access journals and all code will be publicly available.

## 2 Chapter 2: Two-Eyed Seeing

### 2.1 Importance and recognition of Traditional Indigenous Knowledge

The ancestral and traditional Knowledge of Indigenous and colonized Peoples is often dismissed, ignored, and contradicted by Western institutions (Smith, 2012). ... The development of Western science is often assumed to clash with the sacred Knowledge many colonized People hold. Western science is also often viewed as more objective, methodical, and unbiased than traditional Knowledge, and as such it is believed by Western institutions to be superior to Indigenous Knowledge (Smith, 2012). However, it is common for Western institutions to (reluctantly) reach similar, if not identical, conclusions as those held by Indigenous people (ref?). The refusal to recognize traditional Knowledge and cooperate with non-Western institutions often results in a loss of time, resources, and funds to the Western institutions and severe damage to the Land the institution operated on, as well as to the people who's ancestors inhabited the region for millennia. The development of Western science at the exclusion of Indigenous Peoples perpetuates colonialism and brings harm all parties involved.

The concept of *two-eye seeing* (ref?) refers to an approach to knowledge and growth that braids Indigenous Knowledge and science together with Western science (Kutz & Tomaselli, 2019; Kimmerer, 2020). Since Traditional Indigenous Knowledge tends to be qualitative, while biological sciences tend to be quantitative, connecting the two is not always simple. One possibility, however, is to use Traditional Knowledge to create well-informed Bayesian priors (ref?). The validity of the priors can be ensured using prior predictive modeling to select priors that align with the Traditional Knowledge. This practice is not new (Girondot & Rizzo, 2015; Bélisle *et al.*, 2018), but it is rarely used, despite it aligning well with the philosophy of Bayesian statistics.

- Inform priors and simulation distributions using Indigenous Traditional Knowledge:
  - A Bayesian framework with Indigenous Knowledge-informed priors (Girondot & Rizzo, 2015)
  - *Local knowledge in ecological modeling* (Bélisle *et al.*, 2018)
  - <https://focus.science.ubc.ca/stats-660805dd930a>
- *A-spatial-overview-of-the-global-importance-of-Indigenous-lands-for-conservation*
- *Native knowledge for native ecosystems*
- *Searching for synergy integrating traditional and scientific ecological knowledge in environmental science education*
- *The Role of Indigenous Burning in Land Management*
- calls to action for scientists: Wong *et al.* (2020)

- weaving TIK and Western knowledge: Tengö *et al.* (2017)
- *Weaving Indigenous knowledge systems and Western sciences in terrestrial research, monitoring and management in Canada: A protocol for a systematic map* (Henri *et al.*, 2021)
- *Indigenizing the North American Model of Wildlife Conservation* (Hessami *et al.*, 2021)
- fish conservation, Indigenous perspectives (Bowles *et al.*, 2021)
- Vertebrate biodiversity in Indigenous-managed lands in Australia, Brazil, and Canada is equal to or greater biodiversity in protected areas (Schuster *et al.*, 2019)



## **3 Chapter 3: Movement analyses**

### **3.1 Movement simulations**

- Inform priors and simulation distributions using Indigenous Traditional Knowledge

### **3.2 Stochasticity map**

- currently there's no raster of stochasticity => paper / product
- PCA on main drivers/causes of stochasticity

### **3.3 Movement analysis**

- add HFI to analysis (but it's temporally static)
- look at HPAMs?

## 4 Chapter (?): Conclusion

- why is this work important?
- so what?
- now what?

## Bibliography

- Bélisle A.C., Asselin H., LeBlanc P. & Gauthier S. (2018). Local knowledge in ecological modeling. *Ecology and Society* **23**, art14. <https://doi.org/10.5751/ES-09949-230214>
- Bowles E., Marin K., MacLeod P. & Fraser D.J. (2021). A three-pronged approach that leans on Indigenous knowledge for northern fish monitoring and conservation. *Evolutionary Applications* **14**, 653–657. <https://doi.org/10.1111/eva.13146>
- Chevin L.-M., Lande R. & Mace G.M. (2010). Adaptation, Plasticity, and Extinction in a Changing Environment: Towards a Predictive Theory. *PLoS Biology* **8**, e1000357. <https://doi.org/10.1371/journal.pbio.1000357>
- DeSoto L., Cailleret M., Sterck F., Jansen S., Kramer K., Robert E.M.R., *et al.* (2020). Low growth resilience to drought is related to future mortality risk in trees. *Nature Communications* **11**, 545. <https://doi.org/10.1038/s41467-020-14300-5>
- Fahrig L. (2007). Non-optimal animal movement in human-altered landscapes. *Functional Ecology* **21**, 1003–1015. <https://doi.org/10.1111/j.1365-2435.2007.01326.x>
- Foley C., Pettorelli N. & Foley L. (2008). Severe drought and calf survival in elephants. *Biology Letters* **4**, 541–544. <https://doi.org/10.1098/rsbl.2008.0370>
- Girondot M. & Rizzo A. (2015). Bayesian Framework to Integrate Traditional Ecological Knowledge into Ecological Modeling: A Case Study. *Journal of Ethnobiology* **35**, 337–353. <https://doi.org/10.2993/etbi-35-02-337-353.1>
- Government of Canada (2021). *Canadian Protected and Conserved Areas Database*.
- Haig H.A., Kingsbury M.V., Laird K.R., Leavitt P.R., Laing R. & Cumming B.F. (2013). Assessment of drought over the past two millennia using near-shore sediment cores from a Canadian boreal lake. *Journal of Paleolimnology* **50**, 175–190. <https://doi.org/10.1007/s10933-013-9712-z>
- Hansen M.C., Potapov P.V., Moore R., Hancher M., Turubanova S.A., Tyukavina A., *et al.* (2013). High-Resolution Global Maps of 21st-Century Forest Cover Change. *Science* **342**, 850–853. <https://doi.org/10.1126/science.1244693>
- Henri D.A., Provencher J.F., Bowles E., Taylor J.J., Steel J., Chelick C., *et al.* (2021). Weaving Indigenous knowledge systems and Western sciences in terrestrial research, monitoring and management in Canada: A protocol for a systematic map. *Ecological Solutions and Evidence* **2**. <https://doi.org/10.1002/2688-8319.12057>
- Hessami M.A., Bowles E., Popp J.N. & Ford A.T. (2021). Indigenizing the North American Model of Wildlife Conservation. *FACETS* **6**, 1285–1306. <https://doi.org/10.1139/facets-2020-0088>

- IPCC (2018). *Global Warming of 1.5°C. An IPCC Special Report on the impacts of global warming of 1.5°C above pre-industrial levels and related global greenhouse gas emission pathways, in the context of strengthening the global response to the threat of climate change, sustainable development, and efforts to eradicate poverty.*
- Johnson D.S., London J.M., Lea M.-A. & Durban J.W. (2008). Continuous-time correlated random walk model for animal telemetry data. *Ecology* **89**, 1208–1215. <https://doi.org/10.1890/07-1032.1>
- Kimmerer R.W. (2020). *Braiding sweetgrass: Indigenous wisdom, scientific knowledge, and the teachings of plants*, Second hardcover edition. Milkweed Editions, Minneapolis.
- Kutz S. & Tomaselli M. (2019). “Two-eyed seeing” supports wildlife health. *Science* **364**, 1135–1137. <https://doi.org/10.1126/science.aau6170>
- Lucherini M. & Lovari S. (1996). Habitat richness affects home range size in the red fox *Vulpes vulpes*. *Behavioural Processes* **36**, 103–105. [https://doi.org/10.1016/0376-6357\(95\)00018-6](https://doi.org/10.1016/0376-6357(95)00018-6)
- Morellet N., Bonenfant C., Börger L., Ossi F., Cagnacci F., Heurich M., *et al.* (2013). Seasonality, weather and climate affect home range size in roe deer across a wide latitudinal gradient within Europe. *Journal of Animal Ecology* **82**, 1326–1339. <https://doi.org/10.1111/1365-2656.12105>
- Nandintsetseg D., Bracis C., Leimgruber P., Kaczensky P., Buuveibaatar B., Lkhagvasuren B., *et al.* (2019). Variability in nomadism: Environmental gradients modulate the movement behaviors of dryland ungulates. *Ecosphere* **10**. <https://doi.org/10.1002/ecs2.2924>
- Nilsen E.B., Herfindal I. & Linnell J.D.C. (2005). Can intra-specific variation in carnivore home-range size be explained using remote-sensing estimates of environmental productivity? *Écoscience* **12**, 68–75. <https://doi.org/10.2980/i1195-6860-12-1-68.1>
- Pedersen E.J., Miller D.L., Simpson G.L. & Ross N. (2019). Hierarchical generalized additive models in ecology: An introduction with mgcv. *PeerJ* **7**, e6876. <https://doi.org/10.7717/peerj.6876>
- Relyea R.A., Lawrence R.K. & Demarais S. (2000). Home Range of Desert Mule Deer: Testing the Body-Size and Habitat-Productivity Hypotheses. *The Journal of Wildlife Management* **64**, 146. <https://doi.org/10.2307/3802984>
- Riotte-Lambert L. & Matthiopoulos J. (2020). Environmental Predictability as a Cause and Consequence of Animal Movement. *Trends in Ecology & Evolution* **35**, 163–174. <https://doi.org/10.1016/j.tree.2019.09.009>
- Ryan Patrick Jones (2021). PM commits up to \$55 million to reduce land degradation at virtual biodiversity summit. *CBC News*
- Schuster R., Germain R.R., Bennett J.R., Reo N.J. & Arcese P. (2019). Vertebrate biodiversity on indigenous-managed lands in Australia, Brazil, and Canada equals that in protected areas. *Environmen-*

- tal Science & Policy* **101**, 1–6. <https://doi.org/10.1016/j.envsci.2019.07.002>
- Smith L.T. (2012). *Decolonizing methodologies: Research and indigenous peoples*, Second edition. Zed Books, London.
- Tengö M., Hill R., Malmer P., Raymond C.M., Spierenburg M., Danielsen F., *et al.* (2017). Weaving knowledge systems in IPBES, CBD and beyond—lessons learned for sustainability. *Current Opinion in Environmental Sustainability* **26-27**, 17–25. <https://doi.org/10.1016/j.cosust.2016.12.005>
- Truth and Reconciliation Commission of Canada (2015). *Honouring the truth, reconciling for the future: Summary of the final report of the Truth and Reconciliation Commission of Canada*. Truth; Reconciliation Commission of Canada.
- Tucker M.A., Böhning-Gaese K., Fagan W.F., Fryxell J.M., Van Moorter B., Alberts S.C., *et al.* (2018). Moving in the Anthropocene: Global reductions in terrestrial mammalian movements. *Science* **359**, 466–469. <https://doi.org/10.1126/science.aam9712>
- Wilson M.C., Chen X.-Y., Corlett R.T., Didham R.K., Ding P., Holt R.D., *et al.* (2016). Habitat fragmentation and biodiversity conservation: Key findings and future challenges. *Landscape Ecology* **31**, 219–227. <https://doi.org/10.1007/s10980-015-0312-3>
- Wong C., Ballegooyen K., Ignace L., Johnson M.J.(Gùdia). & Swanson H. (2020). Towards reconciliation: 10 Calls to Action to natural scientists working in Canada. *FACETS* **5**, 769–783. <https://doi.org/10.1139/facets-2020-0005>

## Appendix 1

```
plot(1:10, pch = 1:10)
```

