

Generalized Threshold Graph MILP

Baihan Liu


University of Colorado Denver

Independent k -set

Definition

Definition. An **independent k -set** is a k -element vertex set $U \subseteq V$ with no edges between vertices of U .

Example:



$S = \{\{1, 2, 3\}, \{1, 2, 4\}, \{1, 2, 5\}\}$ is a family of independent 3-sets in the graph on vertex set $V = \{1, 2, 3, 4, 5\}$.

* Throughout this talk, all graphs are simple (no loops or multiple edges).

Independent k -set question

We are interesting in the inverse question...

Question

Given a family S of k -sets with $k \geq 2$, does there exist a graph G whose independent k -sets are exactly the sets in S ?

Let the vertex set of our graph to be

$$V(G) = V = \bigcup_{U \in S} U = \{1, \dots, n\}.$$

Additional vertices do not affect whether the graph G exist.

Integer program for independent k -set question

Edge variables:

$$x_{ij} = \begin{cases} 1, & \text{if } \{i, j\} \text{ is an edge of } G, \\ 0, & \text{otherwise,} \end{cases} \quad (1 \leq i < j \leq n).$$

Integer program:

min 0

subject to

$$x_{ij} = 0 \quad \forall U \in S, \forall \{i, j\} \subseteq U,$$

$$\sum_{\{i, j\} \subseteq U} x_{ij} \geq 1 \quad \forall U \in \binom{V}{k} \setminus S,$$

$$x_{ij} \in \{0, 1\} \quad \forall 1 \leq i < j \leq n.$$

LP relaxation

Edge variables:

$x_{ij} > 0$, if $\{i, j\}$ is an edge of G .

$x_{ij} = 0$, otherwise. $(1 \leq i < j \leq n)$.

LP relaxation program:

min 0

subject to

$$x_{ij} = 0 \quad \forall U \in S, \forall \{i, j\} \subseteq U,$$

$$\sum_{\{i, j\} \subseteq U} x_{ij} \geq 1 \quad \forall U \in \binom{V}{k} \setminus S.$$

Equivalent integer program: find complement graph (S is the family of k -cliques)

Edge variables:

$$x_{ij} = \begin{cases} 1, & \text{if } \{i, j\} \text{ is an edge of } G, \\ 0, & \text{otherwise,} \end{cases} \quad (1 \leq i < j \leq n).$$

Integer program:

min 0

subject to

$$x_{ij} = k \quad \forall U \in S, \forall \{i, j\} \subseteq U,$$

$$\sum_{\{i, j\} \subseteq U} x_{ij} \leq k - 1 \quad \forall U \in \binom{V}{k} \setminus S,$$

$$x_{ij} \in \{0, 1\} \quad \forall 1 \leq i < j \leq n.$$

Shifted family (up to relabeling)

Definition

Definition. A family of k -sets S is **shifted** (up to relabeling) if whenever $U \in S$ and $i < j$ with $i \notin U, j \in U$, then

$$(U \setminus \{j\}) \cup \{i\} \in S.$$


Example: $S = \{\{1, 2, 3\}, \{1, 2, 4\}, \{1, 2, 5\}\}$ is shifted (up to relabeling).

Question

Given a family S of k -sets with $k \geq 2$, is it shifted?

Shifted family (up to relabeling)

- The family can be encoded by a nondecreasing sequence (s_1, \dots, s_k) .
- Every $U = (u_1, \dots, u_k) \in S$ satisfies $u_1 < \dots < u_k$ and $u_i \leq s_i$ for all i .



Red Path: $(s_1 = 2, s_2 = 5, s_3 = 8)$

Blue Path: $(u_1 = 1, u_2 = 2, u_3 = 4)$

Shifted \iff k -sum-threshold

Theorem

Let $S \subseteq \binom{V}{k}$ with $V = \bigcup_{U \in S} U$. Then the following are equivalent:

- **Shifted (up to relabeling):** whenever $U \in S$ and $i < j$ with $i \notin U, j \in U$, also $(U \setminus \{j\}) \cup \{i\} \in S$.
- **k -sum-threshold:** there exist weights $w : V \rightarrow \mathbb{R}$ and $t \in \mathbb{R}$ such that

$$S = \left\{ U \in \binom{V}{k} : \sum_{u \in U} w(u) \leq t \right\}.$$



Proof: k -sum-threshold \Rightarrow shifted

Assume \mathcal{S} is k -sum-threshold.

There exist weights $w : V \rightarrow \mathbb{R}$ and a threshold $t \in \mathbb{R}$ such that

$$\mathcal{S} = \left\{ U \in \binom{V}{k} : \sum_{u \in U} w(u) \leq t \right\}.$$

Relabel $V = \{1, \dots, n\}$ so that

$$w(1) \leq w(2) \leq \dots \leq w(n).$$

Let $U \in \mathcal{S}$ and suppose $i < j$ with $j \in U$ and $i \notin U$. Define

$$U' := (U \setminus \{j\}) \cup \{i\}.$$

Then

$$\sum_{u \in U'} w(u) = \sum_{u \in U} w(u) - w(j) + w(i) \leq \sum_{u \in U} w(u) \leq t.$$

Hence $U' \in \mathcal{S}$, so \mathcal{S} is shifted with respect to $1 < \dots < n$.

Shifted \Rightarrow k -sum-threshold (I)

Assume $\mathcal{S} \subseteq \binom{V}{k}$ is shifted (w.r.t. $1 < \dots < n$).

By the upper bound sequence representation, there is a strictly increasing sequence (s_1, \dots, s_k) such that, for

$$U = \{u_1 < \dots < u_k\}, \quad U \in \mathcal{S} \iff u_i \leq s_i \text{ for all } i = 1, \dots, k.$$

Choose a large base $B > 1$ and define the weights and sum:

$$w(i) := B^i \quad (i = 1, \dots, n), \quad \phi(U) := \sum_{u \in U} w(u).$$

Let the “maximal” set be

$$S^* := \{s_1, \dots, s_k\}, \quad t := \phi(S^*) = \sum_{i=1}^k B^{s_i}.$$

Goal: show

$$U \in \mathcal{S} \iff \phi(U) \leq t \iff \sum_{u \in U} w(u) \leq t.$$

Shifted \Rightarrow k -sum-threshold (II)

$$w(i) = B^i, \quad \phi(U) = \sum_{u \in U} B^u, \quad t = \phi(S^*) = \sum_{i=1}^k B^{s_i}.$$

1. If $U \in \mathcal{S}$, write $U = \{u_1 < \dots < u_k\}$. Then $u_i \leq s_i$ for all i , so

$$\phi(U) = \sum_{i=1}^k B^{u_i} \leq \sum_{i=1}^k B^{s_i} = t.$$

2. If $U \notin \mathcal{S}$, let j be the first index with $u_j > s_j$ (so $u_i \leq s_i$ for all $i < j$). The term B^{u_j} is strictly larger than B^{s_j} , and for sufficiently large B this “first violation” dominates all later terms. Concretely, we can choose B so large that

$$B^{u_j} - B^{s_j} > \sum_{i>j} (B^{s_i} - B^{u_i}),$$

which implies

$$\phi(U) > \phi(S^*) = t.$$

Thus, for such a choice of B ,

$$U \in \mathcal{S} \iff \phi(U) \leq t \iff \sum_{u \in U} w(u) \leq t,$$

so \mathcal{S} is k -sum-threshold.

Integer program via shifted definition

Given $V = \{1, \dots, n\}$, fixed k , and a family $S \subseteq \binom{V}{k}$.

Variables:

$$p_{i\ell} \in \{0, 1\} \quad (i \in V, \ell = 1, \dots, n), \quad \text{pos}_i \in \mathbb{Z} \quad (i \in V).$$

Objective:

$$\min 0 \quad (\text{feasibility problem}).$$

Permutation constraints:

$$\sum_{\ell=1}^n p_{i\ell} = 1 \quad \forall i \in V, \quad \sum_{i=1}^n p_{i\ell} = 1 \quad \forall \ell = 1, \dots, n.$$

Integer program via shifted definition

Position constraints:

$$\text{pos}_i = \sum_{\ell=1}^n \ell p_{i\ell} \quad \forall i \in V.$$

Shifted constraints:

For all $U \in S$ and all pairs (i, j) with $j \in U, i \notin U$:

- Let $U' = (U \setminus \{j\}) \cup \{i\}$.
- If $U' \notin S$, impose

$$\text{pos}_i \geq \text{pos}_j + 1,$$

i.e.

$$\sum_{\ell=1}^n \ell p_{i\ell} \geq \sum_{\ell=1}^n \ell p_{j\ell} + 1.$$

Linear programming via k -sum threshold definition

Goal: Find a positive margin δ that separate S with its complement.

Variables:

$$w_i \quad (i \in V), \quad t, \quad \delta \geq 0.$$

Linear program:

$$\max \delta$$

subject to

$$\sum_{i \in U} w_i \leq t - \delta \quad \forall U \in S,$$

$$\sum_{i \in U} w_i \geq t + \delta \quad \forall U \in \binom{V}{k} \setminus S,$$

$$0 \leq w_i \leq 1 \quad \forall i \in V,$$

$$0 \leq t \leq k,$$

$$\delta \geq 0.$$

Interpreting the LP

- If the optimal value satisfies $\delta^* > 0$, then S is shifted/ k -sum threshold.
- If the LP is infeasible or $\delta^* = 0$, then S is not shifted/not k -sum threshold.
- For any k -set $U \in S$, the sum of weights $\sum_{i \in U} w_i$ lies *within* (below) the threshold.
- For any k -set $U \notin S$, the sum of weights $\sum_{i \in U} w_i$ lies *beyond* (above) the threshold.
- We bound w_i and t to prevent unboundedness.

Comparison of the two formulations

Comparison (variables and types)

● IP via shifted definition (permutation-based):

- Number of variables: $n^2 + n$.
- Types: n^2 *binary* variables $p_{i\ell}$ and n *integer* variables pos_i .
- Integer Program(IP); much more combinatorial and typically harder to solve.

● LP via k -sum-threshold:

- Number of variables: $n + 2$.
- Types: all variables are *continuous* (real-valued).
- Complexity driven mainly by the number of constraints over all k -sets.

● Summary:

- IP is larger ($O(n^2)$ variables) and uses integrality to encode the permutation explicitly.
- LP is small ($O(n)$ variables) and purely continuous.

k -sum threshold graph question

What if S is a family of independent k -sets of a graph G and S is also shifted?

Then there exist $w : V(G) \rightarrow \mathbb{R}$ and $t \in \mathbb{R}$ such that

$$U \subseteq V(G), |U| = k \text{ is independent in } G \iff \sum_{u \in U} w(u) \leq t.$$

combinatorial structure defined by edges \iff linear information on vertices

We call graph G is a k -sum-threshold graph.

Question

Given a family S of k -sets with $k \geq 2$, does there exist a k -sum-threshold graph G whose independent k -sets are exactly the sets in S ?

Mixed Integer Linear Program

Variables:

$$x_{ij} \in \{0, 1\} \quad (1 \leq i < j \leq n), \quad w_i \in [0, 1] \quad (i \in V), \quad t \in [0, k], \quad \delta \geq 0.$$

MILP: max δ , subject to

$$\sum_{\{i,j\} \subseteq E(U)} x_{ij} = 0 \quad \forall U \in S,$$

$$\sum_{\{i,j\} \subseteq E(U)} x_{ij} \geq 1 \quad \forall U \in \binom{V}{k} \setminus S,$$

$$x_{ij} \in \{0, 1\} \quad \forall 1 \leq i < j \leq n.$$

$$\sum_{i \in U} w_i \leq t - \delta \quad \forall U \in S,$$

$$\sum_{i \in U} w_i \geq t + \delta \quad \forall U \in \binom{V}{k} \setminus S,$$

$$0 \leq w_i \leq 1 \quad \forall i \in V,$$

$$0 \leq t \leq k,$$

$$\delta \geq 0.$$

AMPL examples

Example 1: shifted, 3-sum-threshold family

$$S = \{\{1, 2, 3\}, \{1, 2, 4\}, \{1, 2, 5\}\}$$

- The AMPL model finds feasible weights w , threshold t , and edge variables x .
- $\delta^* \neq 0$. We obtain a 3-sum-threshold graph whose independent 3-sets are exactly the sets in S .

Example 2: non-shifted, not 3-sum-threshold

$$S = \{\{1, 2, 3\}, \{1, 2, 4\}, \{1, 3, 5\}\}$$

- There exists a graph G with independent 3-sets being S .
- However, $\delta^* = 0$. There is no 3-sum-threshold graph with independent 3-sets exactly S .

MILP complexity and feasible region

- Since there are $\binom{n}{k}$ many k -sets, the formulation has exponentially many constraints, and the resulting MILP is NP-hard to solve in general. It is therefore not very practical for large n and k . For example, $n = 50$ and $k = 5$.
- NP-hard algorithm is still useful. For small values of n and k , we can search over families of independent k -sets that correspond to k -sum-threshold graphs, observe combinatorial patterns, and then formally prove the resulting patterns. In fact, this analysis is being done in my article.
- LP analysis is useful since its feasible solutions correspond to all possible graphs that satisfies the given constraints.

Independence polytopes and degree-sequence polytopes

- **Independence polytope:**

$$P_{\text{ind}}(G) = \text{conv}\{\chi^U : U \subseteq V \text{ independent}\},$$


$$\text{where } (\chi^U)_i = \begin{cases} 1, & i \in U, \\ 0, & i \notin U \end{cases}$$

and independent k -sets are the integer points with $\sum_{i=1}^n x_i = k$.

- **Degree-sequence polytope:**

$$P_{\text{deg}}(n) = \text{conv}\{d(G) = (d_1, \dots, d_n) : G \text{ is a simple graph on } \{1, \dots, n\}\},$$

whose extreme points are exactly the degree sequences of threshold graphs.



Bibliography



B. Liu and L. Xue.

Shifted Total Cut Complex.

In preparation, 2025.



M. Bayer, M. Denker, M. J. Milutinović, R. Rowlands, S. Sundaram, and L. Xue.

Total cut complexes of graphs.

Discrete & Computational Geometry, 73(2):500–527, 2025.



C. J. Klivans.

Combinatorial properties of shifted complexes.

PhD thesis, MIT, 2003.

Thank you!

