

Find the drinkers who frequent **only** bars serving a beer they like

$\forall x \in R \ \phi(x) \equiv \neg \exists x \in R \ \neg \phi(x)$

$\{d : \text{drinker} \mid \exists f \in \text{frequents} \ (f(\text{drinker}) = d(\text{drinker})) \wedge$
 $\forall y \in \text{frequents} [y(\text{drinker}) = f(\text{drinker}) \rightarrow$
 $\exists s \in \text{serves} \exists l \in \text{likes} (s(\text{bar}) = y(\text{bar}) \wedge$
 $s(\text{beer}) = l(\text{beer}) \wedge l(\text{drinker}) = y(\text{drinker}))]\}$

$\{d : \text{drinker} \mid \exists f \in \text{frequents} \ (f(\text{drinker}) = d(\text{drinker})) \wedge$
 $\neg \exists y \in \text{frequents} [y(\text{drinker}) = f(\text{drinker}) \wedge$
 $\neg \exists s \in \text{serves} \exists l \in \text{likes} (s(\text{bar}) = y(\text{bar}) \wedge$
 $s(\text{beer}) = l(\text{beer}) \wedge l(\text{drinker}) = y(\text{drinker}))]\}$

```
select f.drinker
from frequents f
where not exists
  (select *
   from frequents y
   where y.drinker = f.drinker and not exists
     (select *
      from serves s, likes l
      where s.bar = y.bar
      and s.beer = l.beer
      and l.drinker = y.drinker))
```

```
select drinker
from frequents where drinker not in
  (select f.drinker
   from frequents f
   where f.bar not in
     (select bar
      from serves, likes
      where serves.beer = likes.beer
      and likes.drinker = f.drinker))
```

List the days when no red boat is reserved

$\{d : \text{day} \mid \text{weekday}(d) \wedge \forall r \in \text{reservation} \ \forall b \in \text{boat}$
 $((r(\text{day}) = d(\text{day}) \wedge r(\text{bname}) = b(\text{bname})) \rightarrow b(\text{color}) \neq \text{red})\}$

$\{d : \text{day} \mid \text{weekday}(d) \wedge \neg \exists r \in \text{reservation} \ \exists b \in \text{boat}$
 $(r(\text{day}) = d(\text{day}) \wedge r(\text{bname}) = b(\text{bname}) \wedge b(\text{color}) = \text{red})\}$

```
select d.day from weekday d
where not exists
  (select * from reservation r, boat b
   where r.day = d.day and r.name = b.bname AND b.color = 'red')
```

Find actors playing in every movie by Berto

$\{a : \text{actor} \mid \exists y \in \text{movie} \ [a(\text{actor}) = y(\text{actor}) \wedge$
 $\forall m \in \text{movie} \ [m(\text{director}) = \text{"Berto"} \rightarrow \exists t \in \text{movie} \ (m(\text{title}) =$
 $t(\text{title}) \wedge t(\text{actor}) = y(\text{actor}))]\}$

$\{a : \text{actor} \mid \exists y \in \text{movie} \ [a(\text{actor}) = y(\text{actor}) \wedge$
 $\neg \exists m \in \text{movie} \ [m(\text{dir}) = \text{"Berto"} \wedge \neg \exists t \in \text{movie} \ (m(\text{title}) = t(\text{title})$
 $\wedge t(\text{actor}) = y(\text{actor}))]\}$

```
CREATE VIEW berto-movie AS
SELECT title FROM movie WHERE director = 'bertolucci'

CREATE VIEW not-all-ber to AS
SELECT m.actor FROM movies m, berto-movies
WHERE berto-movies.title NOT IN
  (SELECT title FROM movies
   WHERE actor = m.actor)

SELECT actor FROM movies WHERE actor NOT IN
  (SELECT * FROM not-all-ber to)
```

List the directors that every actor is cast in one of his movies

$\{d : \text{director} \mid \exists m \in \text{movie} \ [d(\text{director}) = m(\text{director}) \wedge$
 $\forall a \in \text{movie} \ \exists x \in \text{movie} \ \exists y \in \text{movie} (x(\text{director}) = m(\text{director}) \wedge x(\text{title}) = y(\text{title})$
 $\wedge y(\text{actor}) = a(\text{actor}))]\}$

$\{d : \text{director} \mid \exists m \in \text{movie} \ [d(\text{director}) = m(\text{director}) \wedge$
 $\neg \exists a \in \text{movie} \neg \exists x \in \text{movie} \ \exists y \in \text{movie} (x(\text{director}) = m(\text{director}) \wedge x(\text{title}) = y(\text{title})$
 $\wedge y(\text{actor}) = a(\text{actor}))]\}$

Propositional logic

- $\neg(\neg p) = p$
- $p \rightarrow q = \neg p \vee q$ (definition of implication)
- $\neg(p \wedge q) = \neg p \vee \neg q$ (De Morgan's Law 1)
- $\neg(p \vee q) = \neg p \wedge \neg q$ (De Morgan's Law 2)
- $\neg(p \rightarrow q) = \neg(\neg p \vee q) = p \wedge \neg q$

Relational calculus

Below R is a relation and Q, Q_1, Q_2 are calculus formulas.

- $\forall x \in R \ (Q(x)) = \neg (\exists x \in R \ (\neg Q(x)))$
- $\exists x \in R \ (Q(x)) = \neg (\forall x \in R \ (\neg Q(x)))$
- $\exists x \in R \ (Q_1(x) \vee Q_2(x)) = \exists x \in R \ (Q_1(x)) \vee \exists x \in R \ (Q_2(x))$ (\exists distributes over \vee)
- $\forall x \in R \ (Q_1(x) \wedge Q_2(x)) = \forall x \in R \ (Q_1(x)) \wedge \forall x \in R \ (Q_2(x))$ (\forall distributes over \wedge)

Note that \exists does not distribute over \wedge and \forall does not distribute over \vee (think of some counter-examples!). However, suppose x occurs in Q_1 but not in Q_2 . Then:

- $\exists x \in R \ (Q_1(x) \wedge Q_2) = \exists x \in R \ (Q_1(x)) \wedge Q_2$
- $\forall x \in R \ (Q_1(x) \vee Q_2) = \forall x \in R \ (Q_1(x)) \vee Q_2$

Also note the following useful facts that follow from the previous ones:

- $\forall x \in R \ (Q_1(x) \rightarrow Q_2(x)) = \neg (\exists x \in R \ (Q_1(x) \wedge \neg Q_2(x)))$
- $\forall x \in R \ \forall y \in R \ (Q(x, y)) = \neg (\exists x \in R \ \exists y \in R \ (\neg Q(x, y)))$

If Q_2 does not contain x then:

- $[\exists x \in R \ (Q_1(x)) \rightarrow Q_2] = \forall x \in R \ [Q_1(x) \rightarrow Q_2]$

Basic

- Testing if an attribute A is null: **IS null, IS NOT null**
- Arithmetic operations: involves any null return null
- Comparison: involves any null return **unknown**

Truth tables involving **unknown**

- AND: false then must false, else **unknown**
- OR: true then must true, else **unknown**
- Aggregate functions
 - Usually ignores tuples with null
 - Returns null if there is no non-null amount
 - All except the COUNT(*) ignore tuples with null values on the aggregate attributes

Foreign key references the **primary key(must have)** of the target or **null**.

```
CREATE TABLE branch(bra_name char(15) not null, bra_city char(30),
PRIMARY KEY(dnumber), UNIQUE(dname),
FOREIGN KEY (mgrssn) REFERENCES emp);
DROP TABLE branch; ALTER TABLE r ADD attribute dom
```

<attribute> **LIKE** <pattern>/(<NOT>UNIQUE <query>/ **ORDER BY** (default asc)

NOT monotonic (**CANNOT BE FLATTENED**)

- A **op ANY/ALL** <query>, If any/all X of the result of query satisfies A op X.
- **Close world** assumption: If a tuple is missing in database, then it's not true.

```
INSERT INTO r(attribute, att) VALUES (v1,v2,v3,..)
DELETE FROM r WHERE, UPDATE r SET WHERE
```

update views without aggregates, nesting, group-by or tuple alias, defined on a **single table**

N/A	pro	con
virtual	no need to maintain correspondence with base	inefficient for views defined via complex queries
materialized	Expect many queries, fast	Cost of space, refresh every time DB update

Find the drinkers who frequent **only** bars serving a beer they like

$\forall x \in R \ \varphi(x) \equiv \neg \exists x \in R \ \neg \varphi(x)$

$\{d : \text{drinker} \mid \exists f \in \text{frequents} (f(\text{drinker}) = d(\text{drinker}) \wedge$
 $\forall y \in \text{frequents}[y(\text{drinker}) = f(\text{drinker}) \rightarrow$
 $\exists s \in \text{serves} \exists l \in \text{likes}(s(\text{bar}) = y(\text{bar}) \wedge$
 $s(\text{beer}) = l(\text{beer}) \wedge l(\text{drinker}) = y(\text{drinker}))]\}$

$\{d : \text{drinker} \mid \exists f \in \text{frequents} (f(\text{drinker}) = d(\text{drinker}) \wedge$
 $\neg \exists y \in \text{frequents}[y(\text{drinker}) = f(\text{drinker}) \wedge$
 $\neg \exists s \in \text{serves} \exists l \in \text{likes}(s(\text{bar}) = y(\text{bar}) \wedge$
 $s(\text{beer}) = l(\text{beer}) \wedge l(\text{drinker}) = y(\text{drinker}))]\}$

```
select f.drinker
from frequents f
where not exists
  (select *
   from frequents y
   where y.drinker = f.drinker and not exists
     (select *
      from serves s, likes l
      where s.bar = y.bar
            and s.beer = l.beer
            and l.drinker = y.drinker))
```

```
select drinker
from frequents where drinker not in
  (select f.drinker
   from frequents f
   where f.bar not in
     (select bar
      from serves, likes
      where serves.beer = likes.beer
            and likes.drinker = f.drinker))
```

List the days when no red boat is reserved

$\{d : \text{day} \mid \text{weekday}(d) \wedge \forall r \in \text{reservation} \ \forall b \in \text{boat}$
 $((r(\text{day}) = d(\text{day}) \wedge r(\text{bname}) = b(\text{bname})) \rightarrow b(\text{color}) \neq \text{red})\}$

$\{d : \text{day} \mid \text{weekday}(d) \wedge \neg \exists r \in \text{reservation} \ \exists b \in \text{boat}$
 $(r(\text{day}) = d(\text{day}) \wedge r(\text{bname}) = b(\text{bname}) \wedge b(\text{color}) = \text{red})\}$

```
select d.day from weekday d
where not exists
  (select * from reservation r, boat b
   where r.day = d.day and r.name = b.bname AND b.color= 'red')
```

Find actors playing in every movie by Berto

$\{a : \text{actor} \mid \exists y \in \text{movie} [a(\text{actor}) = y(\text{actor}) \wedge$
 $\forall m \in \text{movie} [m(\text{director}) = \text{"Berto"} \rightarrow \exists t \in \text{movie} (m(\text{title}) =$
 $t(\text{title}) \wedge t(\text{actor}) = y(\text{actor}))]\}$

$\{a : \text{actor} \mid \exists y \in \text{movie} [a(\text{actor}) = y(\text{actor}) \wedge$
 $\neg \exists m \in \text{movie} [m(\text{dir}) = \text{"Berto"} \wedge \neg \exists t \in \text{movie} (m(\text{title}) = t(\text{title})$
 $\wedge t(\text{actor}) = y(\text{actor}))]\}$

List the directors that every actor is cast in one of his movies

$\{d : \text{director} \mid \exists m \in \text{movie} [d(\text{director}) = m(\text{director}) \wedge$
 $\forall a \in \text{movie} \exists x \in \text{movie} \exists y \in \text{movie} (x(\text{director}) = m(\text{director}) \wedge x(\text{title}) = y(\text{title})$
 $\wedge y(\text{actor}) = a(\text{actor}))]\}$

$\{d : \text{director} \mid \exists m \in \text{movie} [d(\text{director}) = m(\text{director}) \wedge$
 $\neg \exists a \in \text{movie} \neg \exists x \in \text{movie} \exists y \in \text{movie} (x(\text{director}) = m(\text{director}) \wedge x(\text{title}) = y(\text{title})$
 $\wedge y(\text{actor}) = a(\text{actor}))]\}$

```
select m.director from movie m
where not exists
  (select * from movie a
   where not exists
     (select * from movie x, movie y
      where x.director = m.director and x.title = y.title and y.actor = a.actor ))
```

Propositional logic

- $\neg(\neg p) = p$
- $p \rightarrow q = \neg p \vee q$ (definition of implication)
- $\neg(p \wedge q) = \neg p \vee \neg q$ (De Morgan's Law 1)
- $\neg(p \vee q) = \neg p \wedge \neg q$ (De Morgan's Law 2)
- $\neg(p \rightarrow q) = \neg(\neg p \vee q) = p \wedge \neg q$

Relational calculus

Below R is a relation and Q_1, Q_2 are calculus formulas.

- $\forall x \in R (Q(x)) = \neg (\exists x \in R (\neg Q(x)))$
- $\exists x \in R (Q(x)) = \neg (\forall x \in R (\neg Q(x)))$
- $\exists x \in R (Q_1(x) \vee Q_2(x)) = \exists x \in R (Q_1(x)) \vee \exists x \in R (Q_2(x))$ (\exists distributes over \vee)
- $\forall x \in R (Q_1(x) \wedge Q_2(x)) = \forall x \in R (Q_1(x)) \wedge \forall x \in R (Q_2(x))$ (\forall distributes over \wedge)

Note that \exists does not distribute over \wedge and \forall does not distribute over \vee (think of some counter-examples!). However, suppose x occurs in Q_1 but not in Q_2 . Then:

- $\exists x \in R (Q_1(x) \wedge Q_2) = \exists x \in R (Q_1(x)) \wedge Q_2$
- $\forall x \in R (Q_1(x) \vee Q_2) = \forall x \in R (Q_1(x)) \vee Q_2$

Also note the following useful facts that follow from the previous ones:

- $\forall x \in R (Q_1(x) \rightarrow Q_2(x)) = \neg(\exists x \in R (Q_1(x) \wedge \neg Q_2(x)))$
- $\forall x \in R \forall y \in R (Q(x, y)) = \neg(\exists x \in R \exists y \in R (\neg Q(x, y)))$

If Q_2 does not contain x then:

- $[\exists x \in R (Q_1(x)) \rightarrow Q_2] = \forall x \in R [Q_1(x) \rightarrow Q_2]$
- $[Q_2 \rightarrow \exists x \in R (Q_1(x))] = \exists x \in R [Q_2 \rightarrow Q_1(x)]$

Basic

- Testing if an attribute A is null: **IS null, IS NOT null**
- Arithmetic operations: involves any null return null
- Comparison: involves any null return **unknown**

Truth tables involving **unknown**

- AND: false then must false, else **unknown**
- OR: true then must true, else **unknown**

- Aggregate functions
 - Usually ignores tuples with null
 - Returns null if there is no non-null amount
 - All except the COUNT(*) ignore tuples with null values on the aggregate attributes

Foreign key references the **primary key(must have)** of the target or **null**.

```
CREATE TABLE branch(bra_name char(15) not null, bra_city char(30),
                     PRIMARY KEY(dnumber), UNIQUE(dname),
                     FOREIGN KEY (mgrssn) REFERENCES emp);
DROP TABLE branch; ALTER TABLE r ADD attribute dom
```

<attribute> LIKE <pattern>/ (NOT) UNIQUE <query>/ ORDER BY (default asc)

NOT monotonic (CANNOT BE FLATTENED)

- A op ANY/ALL <query>, If any/all X of the result of query satisfies A op X.
- **Close world** assumption: If a tuple is missing in database, then it's not true.

```
INSERT INTO r(attribute, att) VALUES (v1,v2,v3,...)
DELETE FROM r WHERE, UPDATE r SET WHERE
```

update views without aggregates, nesting, group-by or tuple alias, defined on a **single table**

N/A	pro	con
virtual	no need to maintain correspondence with base	inefficient for views defined via complex queries
materialized	Expect many queries, fast	Cost of space, refresh every time DB update