

Learning Objectives


After this next lesson, you should be able to:

- How to conduct linear regression modeling
- Use interaction effects and binary categorical variables (also called dummy variables)
- Understand model complexity, underfitting, right fit, and overfitting
- Define regularization and error metrics for regression problems

Review

- Simple and Multiple Linear Regressions
- Common regression assumptions; how to check for them
- OLS (Ordinary Least Squares)
- How to interpret the model's parameters
- Variable Transformations
- Inference, Fit, R^2 (r-squared), and \bar{R}^2 (adjusted R^2)
- Multicollinearity

Today we keep our focus on the **REFINE** the data and **BUILD** a model steps but with (1) a focus on linear regression modeling and (2) what the inferential statistics tell us about the fit of these linear models



Today (cont.)

Research Design and Data Analysis	Research Design	Data Visualization in <i>pandas</i>	Descriptive Statistics for Exploratory Data Analysis	Exploratory Data Analysis in <i>pandas</i>
			Inferential Statistics for Model Fit	
Foundations of Modeling	Linear Regression	Classification Models	Evaluating Model Fit	Presenting Insights from Data Models
Data Science in the Real World	Decision Trees and Random Forests	Time Series Data	Natural Language Processing	Databases

Here's what's happening today:

- Announcements and Exit Tickets
- Review
- **⑤ Refine the Data and ⑥ Build a Model | Linear Regression**
 - F-statistic
 - Backward selection or “how to conduct linear regression modeling”
 - Linear Regression Modeling with sklearn (scikit-learn)
- *statsmodels* vs. *sklearn*
- Interaction Effects
- Underfitting and overfitting; Training and generalization errors
- (Complexity and Regularization)
- Binary (dummy) categorical variables
- Lab – Introduction to Regression and Model Fit, Part 2
- Review

DS

Linear Regression

Model's F-statistic

What β_i would make our multiple linear regression model useless?

- (the multiple linear regression model again)

$$y = \beta_0 + \beta_1 \cdot x_1 + \cdots + \beta_k \cdot x_k + \varepsilon$$

- Answer: If $\beta_0 = \beta_1 = \cdots = \beta_k = 0$, we don't have a model
 - ($y = 0$ isn't very exciting, is it?)

Model's F-statistic Hypothesis Test

- The *null hypothesis* (H_0) represents the status quo; that all β_i are zeros.

$$H_0: \beta_0 = \beta_1 = \cdots = \beta_k = 0$$

- The *alternate hypothesis* (H_a) represents the opposite of the null hypothesis (that at least one β_i is not zero) and holds true if H_0 is found to be false:

$$H_a: \exists i: \beta_i \neq 0$$

DS

Linear Regression

Activity | Model's F-statistic

Two-step guidance on how to conduct linear regression modeling

① Model's significance

- Always start with the F-statistics for the whole model; only then check individual variables

② Regressors' significance

- Prefer to work solely with significant variables: if you observe insignificant variables you *usually* need to get rid of them and rerun your regression modeling without those
- Backward selection method
 - If you have insignificant variables, start dropping the most insignificant variable. If after removing that variable you still have insignificant variables, drop them one by one, until you are left with no insignificant variables

A black circle containing the white text "DS".

DS

Linear Regression

Linear Regression Modeling with sklearn (scikit-learn)

Linear Modeling with *sklearn*

- When modeling with *sklearn* (scikit-learn), you'll use the following base principles:
 - All *sklearn* modeling classes are based on the base estimator `sklearn.base.BaseEstimator`
 - This means that all *sklearn* models take a similar form
 - All estimators take a matrix X (a *pandas* DataFrame), either sparse or dense
 - Supervised estimators also take a vector y (the response) (a *pandas* Series)
 - Estimators can be customized through setting the appropriate parameters

General format for *sklearn* model classes and methods

❶ `model = base_models.AnySKLearnObject()`

`# create an instance of an estimator class`

❷ `model.fit(train_X, train_y)`

`# train your model; also called “fitting your data”`

❸ `model.score(train_X, train_y)`

`# score your model using the training data using the
default scoring method (recommended to use the metrics
module in the future)`

❹ `model.predict(test_X)`

`# predict your test data`

❺ `model.score(test_X, test_y)`


`# score your model using your test data`

❻ `model.predict(new_X)`


`# make predictions for a new set of data`

q-q plots of residuals. Are they normally distributed?


Sales ~ TV



Sales ~ Radio




Sales ~ Newspaper




Scatterplots of residuals against advertising budget. Are they randomly distributed?


Sales ~ TV



Sales ~ Radio



Sales ~ Newspaper



DS

Linear Regression

Multiple Linear Regression / $Sales \sim TV + Radio + Newspaper$

$$Sales \sim TV + Radio + Newspaper$$

Dep. Variable:	Sales	R-squared:	0.895
Model:	OLS	Adj. R-squared:	0.894
Method:	Least Squares	F-statistic:	553.5
Date:		Prob (F-statistic):	8.35e-95
Time:		Log-Likelihood:	-383.24
No. Observations:	198	AIC:	774.5
Df Residuals:	194	BIC:	787.6
Df Model:	3		
Covariance Type:	nonrobust		

	coef	std err	t	P> t	[95.0% Conf. Int.]
Intercept	2.9523	0.318	9.280	0.000	2.325 3.580
TV	0.0457	0.001	32.293	0.000	0.043 0.048
Radio	0.1886	0.009	21.772	0.000	0.171 0.206
Newspaper	-0.0012	0.006	-0.187	0.852	-0.014 0.011

Omnibus:	59.593	Durbin-Watson:	2.041
Prob(Omnibus):	0.000	Jarque-Bera (JB):	147.654
Skew:	-1.324	Prob(JB):	8.66e-33
Kurtosis:	6.299	Cond. No.	457.

DS

Linear Regression

Multiple Linear Regression / $Sales \sim TV + Radio$

Sales ~ TV + Radio. Are we done yet?


Dep. Variable:	Sales	R-squared:	0.895
Model:	OLS	Adj. R-squared:	0.894
Method:	Least Squares	F-statistic:	834.4
Date:		Prob (F-statistic):	2.60e-96
Time:		Log-Likelihood:	-383.26
No. Observations:	198	AIC:	772.5
Df Residuals:	195	BIC:	782.4
Df Model:	2		
Covariance Type:	nonrobust		

	coef	std err	t	P> t	[95.0% Conf. Int.]
Intercept	2.9315	0.297	9.861	0.000	2.345 3.518
TV	0.0457	0.001	32.385	0.000	0.043 0.048
Radio	0.1880	0.008	23.182	0.000	0.172 0.204


Omnibus:	59.228	Durbin-Watson:	2.038
Prob(Omnibus):	0.000	Jarque-Bera (JB):	145.127
Skew:	-1.321	Prob(JB):	3.06e-32
Kurtosis:	6.257	Cond. No.	423.

$Sales \sim TV + Radio$. What do you observe? Are we done yet?


Residuals q-q plot



Residuals against TV



Residuals against $Radio$



$Sales \sim TV + Radio$

$$\triangleright Sales = \underbrace{2.93}_{\hat{\beta}_0} + \underbrace{.0457}_{\hat{\beta}_1} \times TV + \underbrace{.188}_{\hat{\beta}_2} \times Radio$$

- This model assumes that the effect on sales of increasing one media (e.g., *TV*) is independent of the amount spent on the other media (e.g., *Radio*)
- More specifically, the model states that the average effect on sales of a one-unit increase (\$1,000) in *TV* is always $(\underbrace{.0457}_{\hat{\beta}_1} \times \underbrace{.\$1,000}_{TV} = \$45.7)$, regardless of the amount spend on *Radio*

DS

Linear Regression

Interaction Effects

Interaction effects

- But suppose that spending money on radio advertising actually increases the effectiveness of *TV* advertising
 - the slope term for *TV* should increase as *Radio* increases
- E.g., given a fixed budget of \$100,000, spending half on TV and half on radio may increase sales more than allocating the entire amount to either TV or radio
- This is known as a synergy effect in marketing; in statistics it is referred to as an interaction effect

A black circle containing the white text 'DS' in a bold, sans-serif font.

DS

Linear Regression

Codealong – Part B
Interaction Effects

$$\text{Sales} \sim \text{TV} + \text{Radio} + \text{TV} * \text{Radio}$$

Dep. Variable:	Sales	R-squared:	0.968
Model:	OLS	Adj. R-squared:	0.967
Method:	Least Squares	F-statistic:	1934.
Date:		Prob (F-statistic):	3.19e-144
Time:		Log-Likelihood:	-267.07
No. Observations:	198	AIC:	542.1
Df Residuals:	194	BIC:	555.3
Df Model:	3		
Covariance Type:	nonrobust		

	coef	std err	t	P> t	[95.0% Conf. Int.]
Intercept	6.7577	0.247	27.304	0.000	6.270 7.246
TV	0.0190	0.002	12.682	0.000	0.016 0.022
Radio	0.0276	0.009	3.089	0.002	0.010 0.045
TV:Radio	0.0011	5.27e-05	20.817	0.000	0.001 0.001

Omnibus:	126.182	Durbin-Watson:	2.241
Prob(Omnibus):	0.000	Jarque-Bera (JB):	1151.060
Skew:	-2.306	Prob(JB):	1.12e-250
Kurtosis:	13.875	Cond. No.	1.78e+04

Interaction effects (cont.)

- $Sales = \underbrace{6.76}_{\hat{\beta}'_0} + \underbrace{.0190}_{\hat{\beta}'_1} \times TV + \underbrace{.0276}_{\hat{\beta}'_2} \times Radio + \underbrace{.0011}_{\hat{\beta}'_3} \times TV \times Radio$
- The interaction is important
 - β'_3 is statistically significant
 - R^2 with this model went up to 96.8% up from 89.5% for the model without interaction. This that $1 - \frac{1-.968}{1-.895} = .70 = 70\%$ of the unexplained variability in the previous model has been explained by the interaction term

Activity | Interaction effects



EXERCISE

DIRECTIONS (10 minutes)

1. Our TV budget is \$50,000 that we consider increasing it by \$5,000. What would be the corresponding increase in sales based on different levels of radio budget?
2. When finished, share your answers with your table

DELIVERABLE

Answers to the above questions

Activity | Interaction effects (cont.)

EXERCISE

Radio budget	Model without interactions	Model with interactions
Formula	$\underbrace{.0457}_{\hat{\beta}_1} \times \Delta TV$	$\left(\underbrace{.0190}_{\hat{\beta}'_1} + \underbrace{.0011}_{\hat{\beta}'_3} \times Radio \right) \times \Delta TV$
\$15,000	$.0457 \times 5 = .228 = \229	$(.0190 + .0011 \times 15) \times 5$ $= .178 = \$178$
\$10,000	\$229	$(.0190 + .0011 \times 10) \times 5$ $= .150 = \$150$
\$5,000	\$229	$(.0190 + .0011 \times 5) \times 5$ $= .123 = \$123$

Hierarchy Principle

- ▶ Sometimes an interaction term $x_i \cdot x_j$ is significant, but one or both of its main effects (in this case x_i and/or x_j) are not
- ▶ The hierarchy principle
 - ▶ If we include an interaction in a model, we should also include the main effects, even if they aren't significant

A black circle containing the white text "DS".

DS


Linear Regression

Underfitting and overfitting

Training and generalization errors

Polynomial regressions

- ▶ Polynomial regressions ($y = \beta_0 + \beta_1 \cdot x + \beta_2 \cdot x^2 + \dots + \beta_k \cdot x^k + \varepsilon$) allow us to fit very complex curves (nonlinear relationships) to the data
- ▶ *(For now, we will gloss over the multicollinearity issue we mentioned in the previous lecture)*



Training and generalization errors

- Training error


- Error rate (e.g., $\|\varepsilon\|^2$ for OLS) derived from the training set ($x = [x_{i,j}]_{\substack{1 \leq i \leq n \\ 0 \leq j \leq k}}$) when estimating $\hat{\beta}$

- Generalization error


- Error rate when estimating \hat{y} for unknown data points (data points that haven't been used to estimate $\hat{\beta}$)

How low can we push the training error?


- Down to zero (effectively “memorizing” the entire training set)
- However, the model is now not only too complex but it will also not generalize well to data that was not used during training
 - This is called overfitting



Error rates, model complexity, and fit




Error rates, model complexity, and fit (cont.)



- Underfitting


- Model is too simple and cannot represent the desired behavior very well
 - Both its training and generalization error are poor

Error rates, model complexity, and fit (cont.)



- Good fit
 - Model has the right level of complexity
 - It performs well on the training set (low training error) and generalize well to unknown data points (low generalization error)

Error rates, model complexity, and fit (cont.)



- ▶ **Overfitting**
 - ▶ **Model is too complex**
 - ▶ **It performs very well on the training set (low training error) but does not generalize well to unknown data points (high generalization error)**

Activity | Underfitting, good fit, and overfitting


EXERCISE

DIRECTIONS (10 minutes)

1. Classify the following polynomial regressions according to their fit:
 1. Underfitting
 2. Good fit
 3. Overfitting
2. When finished, share your answers with your table

DELIVERABLE

Answers to the above questions



DS

Linear Regression

Complexity and Regularization

FYI | How do we define complexity?

- E.g., as a function of the size of the coefficients
 - $\|\beta\|_1 = \sum_{j=0}^k |\beta_j|$ (L1-norm)
 - $\|\beta\|_2^2 = \sum_{j=0}^k \beta_j^2$ (L2-norm)
 - (with $\beta = (\beta_0, \dots, \beta_k)$)

FYI | Regularization prevents overfitting by explicitly controlling model complexity

- These definitions of complexity lead to the following regularization techniques

- $\min \left(\underbrace{\|y - x \cdot \beta\|^2}_{OLS \text{ term}} + \underbrace{\lambda \|\beta\|_1}_{\text{regularization term}} \right)$ (L1 regularization; a.k.a., Lasso)

- $\min(\|y - x \cdot \beta\|^2 + \lambda \|\beta\|_2^2)$ (L2 regularization; a.k.a., Ridge)

- (You will use the gradient descent technique discussed earlier to train your model)
- This formulation reflects the fact that there is a cost associated with regularization that we want to minimize

A black circle containing the white text 'DS' in a bold, sans-serif font.

Linear Regression

Binary (a.k.a., Dummy) (Categorical) Variables

Back to the SF housing dataset and the issue of bed and bath counts

- ▶ So far, we've considered *BedCount* and *BathCount* as ratio variables
 - ▶ Namely that the price premium between a property with 1 bathroom and another with 2 bathrooms was the same between a property with 3 bathrooms and another with 4 bathrooms
- ▶ Does this make sense?

Dep. Variable:	SalePrice	R-squared:	0.137
Model:	OLS	Adj. R-squared:	0.136
Method:	Least Squares	F-statistic:	146.6
Date:		Prob (F-statistic):	1.94e-31
Time:		Log-Likelihood:	-1690.7
No. Observations:	929	AIC:	3385.
Df Residuals:	927	BIC:	3395.
Df Model:	1		
Covariance Type:	nonrobust		

	coef	std err	t	P> t	[95.0% Conf. Int.]
Intercept	0.3401	0.099	3.434	0.001	0.146 0.535
BathCount	0.5242	0.043	12.109	0.000	0.439 0.609

Omnibus:	1692.623	Durbin-Watson:	1.582
Prob(Omnibus):	0.000	Jarque-Bera (JB):	2167434.305
Skew:	12.317	Prob(JB):	0.00
Kurtosis:	238.345	Cond. No.	5.32

Back to the SF housing dataset and the issue of bed and bath counts

- ▶ Let's test this hypothesis and convert *BathCount* to a nominal variable (indeed, we won't even assume an order) and then encode it to “dummy” categorical variables

m (# bathrooms)	$Bath = \begin{pmatrix} Bath_1, \\ Bath_2, \\ Bath_3, \\ Bath_4 \end{pmatrix}$ (encoding)
1	(1, 0, 0, 0)
2	(0, 1, 0, 0)
3	(0, 0, 1, 0)
4	(0, 0, 0, 1)

Activity | Binary (categorical) variables



EXERCISE

DIRECTIONS (10 minutes)

1. Complete the codealong by
 - a. Run 4 regressions, one for each of the case highlighted in the handout (Each case only include 3 out of the 4 dummy variables we created) (we give you the first one...)
 - b. What are the coefficients for the different β s?
 - c. How do you interpret the β s?
 - d. Why do we only need three dummy variables, not four?
2. When finished, share your answers with your table

DELIVERABLE

Answers to the above questions

Activity | Binary (categorical) variables (cont.)



EXERCISE

$$\text{SalePrice} = \beta_1 + \beta_{1,2} \cdot \text{Bath}_2 + \beta_{1,3} \cdot \text{Bath}_3 + \beta_{1,4} \cdot \text{Bath}_4$$

(don't include Bath_1)

$$\text{SalePrice} = \beta_2 + \beta_{2,1} \cdot \text{Bath}_1 + \beta_{2,3} \cdot \text{Bath}_3 + \beta_{2,4} \cdot \text{Bath}_4$$

(don't include Bath_2)

$$\text{SalePrice} = \beta_3 + \beta_{3,1} \cdot \text{Bath}_1 + \beta_{3,2} \cdot \text{Bath}_2 + \beta_{3,4} \cdot \text{Bath}_4$$

(don't include Bath_3)

$$\text{SalePrice} = \beta_4 + \beta_{4,1} \cdot \text{Bath}_1 + \beta_{4,2} \cdot \text{Bath}_2 + \beta_{4,3} \cdot \text{Bath}_3$$

(don't include Bath_4)

Activity | Four linear regressions to run (cont.)

$$\text{SalePrice} = \beta_1 + \beta_{1,2} \cdot \text{Bath}_2 + \beta_{1,3} \cdot \text{Bath}_3 + \beta_{1,4} \cdot \text{Bath}_4$$

```
formula = 'SalePrice ~ Bath_2 + Bath_3 + Bath_4'
```

$$\text{SalePrice} = \beta_2 + \beta_{2,1} \cdot \text{Bath}_1 + \beta_{2,3} \cdot \text{Bath}_3 + \beta_{2,4} \cdot \text{Bath}_4$$

```
formula = 'SalePrice ~ Bath_1 + Bath_3 + Bath_4'
```

$$\text{SalePrice} = \beta_3 + \beta_{3,1} \cdot \text{Bath}_1 + \beta_{3,2} \cdot \text{Bath}_2 + \beta_{3,4} \cdot \text{Bath}_4$$

```
formula = 'SalePrice ~ Bath_1 + Bath_2 + Bath_4'
```

$$\text{SalePrice} = \beta_4 + \beta_{4,1} \cdot \text{Bath}_1 + \beta_{4,2} \cdot \text{Bath}_2 + \beta_{4,3} \cdot \text{Bath}_3$$

```
formula = 'SalePrice ~ Bath_1 + Bath_2 + Bath_3'
```

Activity | Four linear regressions to run (cont.)

Dep. Variable:	SalePrice	R-squared:	0.043
Model:	OLS	Adj. R-squared:	0.039
Method:	Least Squares	F-statistic:	11.78
Date:		Prob (F-statistic):	1.49e-07
Time:		Log-Likelihood:	-1314.2
No. Observations:	794	AIC:	2636.
Df Residuals:	790	BIC:	2655.
Df Model:	3		
Covariance Type:	nonrobust		

	coef	std err	t	P> t	[95.0% Conf. Int.]
Intercept	0.9914	0.070	14.249	0.000	0.855 1.128
Bath_2	0.2831	0.099	2.855	0.004	0.088 0.478
Bath_3	0.4808	0.142	3.383	0.001	0.202 0.760
Bath_4	1.2120	0.232	5.231	0.000	0.757 1.667

Omnibus:	1817.972	Durbin-Watson:	1.867
Prob(Omnibus):	0.000	Jarque-Bera (JB):	8069883.811
Skew:	19.917	Prob(JB):	0.00
Kurtosis:	495.280	Cond. No.	5.79

Dep. Variable:	SalePrice	R-squared:	0.043
Model:	OLS	Adj. R-squared:	0.039
Method:	Least Squares	F-statistic:	11.78
Date:		Prob (F-statistic):	1.49e-07
Time:		Log-Likelihood:	-1314.2
No. Observations:	794	AIC:	2636.
Df Residuals:	790	BIC:	2655.
Df Model:	3		
Covariance Type:	nonrobust		

	coef	std err	t	P> t	[95.0% Conf. Int.]
Intercept	1.2745	0.071	18.040	0.000	1.136 1.413
Bath_1	-0.2831	0.099	-2.855	0.004	-0.478 -0.088
Bath_3	0.1977	0.143	1.386	0.166	-0.082 0.478
Bath_4	0.9290	0.232	4.003	0.000	0.473 1.384

Omnibus:	1817.972	Durbin-Watson:	1.867
Prob(Omnibus):	0.000	Jarque-Bera (JB):	8069883.811
Skew:	19.917	Prob(JB):	0.00
Kurtosis:	495.280	Cond. No.	5.84

Dep. Variable:	SalePrice	R-squared:	0.043
Model:	OLS	Adj. R-squared:	0.039
Method:	Least Squares	F-statistic:	11.78
Date:		Prob (F-statistic):	1.49e-07
Time:		Log-Likelihood:	-1314.2
No. Observations:	794	AIC:	2636.
Df Residuals:	790	BIC:	2655.
Df Model:	3		
Covariance Type:	nonrobust		

	coef	std err	t	P> t	[95.0% Conf. Int.]
Intercept	1.4722	0.124	11.881	0.000	1.229 1.715
Bath_1	-0.4808	0.142	-3.383	0.001	-0.760 -0.202
Bath_2	-0.1977	0.143	-1.386	0.166	-0.478 0.082
Bath_4	0.7313	0.253	2.886	0.004	0.234 1.229

Omnibus:	1817.972	Durbin-Watson:	1.867
Prob(Omnibus):	0.000	Jarque-Bera (JB):	8069883.811
Skew:	19.917	Prob(JB):	0.00
Kurtosis:	495.280	Cond. No.	7.52

Dep. Variable:	SalePrice	R-squared:	0.043
Model:	OLS	Adj. R-squared:	0.039
Method:	Least Squares	F-statistic:	11.78
Date:		Prob (F-statistic):	1.49e-07
Time:		Log-Likelihood:	-1314.2
No. Observations:	794	AIC:	2636.
Df Residuals:	790	BIC:	2655.
Df Model:	3		
Covariance Type:	nonrobust		

	coef	std err	t	P> t	[95.0% Conf. Int.]
Intercept	2.2035	0.221	9.969	0.000	1.770 2.637
Bath_1	-1.2120	0.232	-5.231	0.000	-1.667 -0.757
Bath_2	-0.9290	0.232	-4.003	0.000	-1.384 -0.473
Bath_3	-0.7313	0.253	-2.886	0.004	-1.229 -0.234

Omnibus:	1817.972	Durbin-Watson:	1.867
Prob(Omnibus):	0.000	Jarque-Bera (JB):	8069883.811
Skew:	19.917	Prob(JB):	0.00
Kurtosis:	495.280	Cond. No.	11.7

Activity | What are the β s' coefficient? (cont.)

β_1 0.9914		$\beta_{1,2}$ 0.2831	$\beta_{1,3}$ 0.4808	$\beta_{1,4}$ 1.212
β_2 1.2745	$\beta_{2,1}$ -0.2831		$\beta_{2,3}$ 0.1977	$\beta_{2,4}$ 0.9290
β_3 1.4722	$\beta_{3,1}$ -0.4808	$\beta_{3,2}$ -0.1977		$\beta_{3,4}$ 0.7313
β_4 2.2025	$\beta_{4,1}$ -1.212	$\beta_{4,2}$ -0.9290	$\beta_{4,3}$ -0.7313	

Activity | What are the β s' coefficient? (cont.)

β_i	Value (Sale's price) of a property in SF with i bathrooms
$\beta_{i,j}$ when $j > i$	Increase of value for a property when increasing the number of bathrooms from i to j (while keeping the rest of the same...)
$\beta_{i,j}$ when $j < i$	Decrease of value for a property when decreasing the number of bathrooms from i to j (while keeping the rest of the same...)
$\beta_{i,j} = -\beta_{j,i}$	Going from i to j bathrooms has the opposite effect of going from j bathrooms to i bathrooms
$\beta_j = \beta_i + \beta_{i,j}$ for any i and j	E.g., $\beta_4 = \beta_1 + \beta_{1,4}$. I.e., the value of a 4 bathrooms can be derived from a 1 bedroom house and by increasing the number of bathrooms for 1 to 4
$\beta_{i,j} = \beta_{i,k} + \beta_{k,j}$ for any i, j and k	E.g., $\beta_{1,4} = \beta_{1,2} + \beta_{2,4}$. I.e., the increase in value from a 1 bathroom house to a 4 bathrooms house is identical to going from upgrading from 1 bathroom to 2 bathrooms and then from upgrading from 2 bathrooms to 4 bathrooms

DS

Review

Review

- Linear Regressions
 - Simple and Multiple
 - Regression assumptions; how to check for them
- Variables
 - Variable Transformations; dummy categorical variables; Interaction effects and the hierarchy principle
 - How to interpret the model's parameters
- Inference and Fit
 - F-statistic
- R^2 (r-squared), and \bar{R}^2 (adjusted R^2)
- Guidance on how to conduct linear regression modeling
 - Backward selection
- Estimating the β s and model complexity
 - OLS (Ordinary Least Squares)
 - Underfitting and overfitting, training and generalization errors, and regularization

Review

You should now be able to:

- How to conduct linear regression modeling
- Use interaction effects and dummy categorical variables
- Understand model complexity, underfitting, right fit, and overfitting
- Define regularization and error metrics for regression problems

DS

Q & A

Next Class

Introduction to Classification

Learning Objectives

After the next lesson, you should be able to:

- Define class label and classification
- Build a K-Nearest Neighbors using the *sklearn* library
- Evaluate and tune model by using metrics such as classification accuracy/error



DS

Exit Ticket

Don't forget to fill out your exit ticket [here](#)