



# OPEN ACCESS

Citation: Ramachandran A, Radhapriya P, Jayakumar S, Dhanya P, Geetha R (2016) Critical Analysis of Forest Degradation in the Southern Eastern Ghats of India: Comparison of Satellite Imagery and Soil Quality Index. PLoS ONE 11(1): e0147541. doi:10.1371/journal.pone.0147541

**Editor:** Madhur Anand, University of Guelph, CANADA

Received: August 14, 2015

Accepted: January 4, 2016

Published: January 26, 2016

Copyright: © 2016 Ramachandran et al. This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Data Availability Statement:** The data is deposited in the dryad doi:10.5061/dryad.1f6m9.

**Funding:** This study received intramural funding from the Anna University and the Pondicherry University. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

**Competing Interests:** The authors have declared that no competing interests exist.

RESEARCH ARTICLE

# Critical Analysis of Forest Degradation in the Southern Eastern Ghats of India: Comparison of Satellite Imagery and Soil Quality Index

Andimuthu Ramachandran<sup>1</sup>\*, Parthasarathy Radhapriya<sup>1</sup>\*, Shanmuganathan Jayakumar<sup>2</sup>, Praveen Dhanya<sup>1</sup>, Rajadurai Geetha<sup>1</sup>

- 1 Centre for Climate Change and Adaptation Research, Anna University, Chennai, Tamil Nadu, India,
- 2 Environmental Informatics and Spatial Modelling Lab (EISML), Department of Ecology & Environmental Sciences, School of Life Sciences, Pondicherry University, Puducherry, India
- \* andimuthu.ramachandran@gmail.com (AR); radhapriya.cc@gmail.com (PR)

# **Abstract**

India has one of the largest assemblages of tropical biodiversity, with its unique floristic composition of endemic species. However, current forest cover assessment is performed via satellite-based forest surveys, which have many limitations. The present study, which was performed in the Eastern Ghats, analysed the satellite-based inventory provided by forest surveys and inferred from the results that this process no longer provides adequate information for quantifying forest degradation in an empirical manner. The study analysed 21 soil properties and generated a forest soil quality index of the Eastern Ghats, using principal component analysis. Using matrix modules and geospatial technology, we compared the forest degradation status calculated from satellite-based forest surveys with the degradation status calculated from the forest soil quality index. The Forest Survey of India classified about 1.8% of the Eastern Ghats' total area as degraded forests and the remainder (98.2%) as open, dense, and very dense forests, whereas the soil quality index results found that about 42.4% of the total area is degraded, with the remainder (57.6%) being non-degraded. Our ground truth verification analyses indicate that the forest soil quality index along with the forest cover density data from the Forest Survey of India are ideal tools for evaluating forest degradation.

#### Introduction

India, a country with rich biodiversity, has different forest types in different climatic zones, each with a unique floristic composition [1, 2]. The forests are mainly distributed in the Himalayas, Western Ghats, Eastern Ghats (EG), and Vindhya ranges. Indian forests have historically experienced a wide variety of management practices because of the differing policies of diverse peoples and governances [2, 3]. In particular, the protected forest reserves of the southern EG, an area in Tamil Nadu [4, 5], had experienced considerable need-based forestry practices (e.g. felling of timber for construction, fuel, or the creation of grazing lands and real estate) for more than 150 years until 1980 [6–10]. As a result, the floristic composition of the forests in the EG underwent massive changes [10, 11], including the emergence of secondary and tertiary vegetation. Furthermore, severe, negative effects on soil health were also reported [8]. Therefore,



effective monitoring and inventory strategies are crucial for the continued conservation of these forests.

Currently, the Forest Survey of India (FSI) periodically monitors forest status based on canopy density, which is obtained via satellite data with ground truth verification. Canopy density is classified into four categories: (i) very dense forest (VDF) with >70% canopy density, (ii) moderately dense forest with 40%–70%, canopy density, (iii) open forest with 10%–40% canopy density, and (iv) degraded forest with <10% canopy density [12]. This classification is based on a normalised difference vegetation index (NDVI), which does not distinguish between crown density and ground density [13]. However, research has shown that the NDVI is not detailed enough to provide reliable data and may underestimate the degradation status [14–17]. Menon and Bawa [18], who estimated the deforestation rates in India and identified large disparities in the forest definitions, also highlighted the need for improved remote sensing and ground truth verification data to discriminate native and plantation forests. Moreover Aziz et al. [19] clearly inferred that the NDVI could not capture the difference between natural forests and rubber plantations in Malaysian forests.

Soil is a vital component of forest ecosystems and is responsible for processes that support biomass production and carbon sequestration [20]. Forest soil health is sensitive to numerous natural and anthropogenic factors, and a deterioration of soil quality typically leads to a deterioration of site quality [21]. Thus, it is important to establish appropriate soil quality indices for measuring the various properties of soil. A soil quality index is defined as the minimum set of parameters that, when interrelated, provides numerical data on the capacity of a soil to carry out one or more functions. A soil quality indicator is a measurable property that influences the capacity of a soil to carry out a given function [22]. A soil quality index, to some extent, should be use-dependent, so that it can be applicable on a larger scale. SQI will help us to select a minimum set of indicators that can address the overall soil quality of a particular region [23]. Foresters have always relied on their knowledge of the chemical, physical, and biological properties of soils to assess the capacity of a site to support productive forests [24]. Recently, the need to assess soil properties has expanded because of growing public interest in determining the consequences of management practices on the quality of soil relative to the sustainability of forest ecosystem functions in addition to plant productivity [25]. The concept of soil quality includes the assessment of soil properties and processes as they relate to the soil's ability to function effectively as a component of a healthy ecosystem [26]. The specific functions and subsequent values provided by forest ecosystems are variable and rely on numerous soil properties and processes [27]. Thus, an understanding of soil properties and soil biota is imperative to the accurate classification of the forest degradation level [8].

Forest soil quality assessment is well established in several developed countries, such as the United States [28], New Zealand [29], and numerous European nations [30], including the United Kingdom [31]. In India, because the majority of soil quality studies are performed in agricultural and horticultural sectors [32, 33], very little data are available on forest soil quality and biota [8]. Therefore, the aim of this study was to identify the degradation status of forests in the EG by performing a detailed soil quality assessment, and then using geospatial technology to compare the soil quality results with those obtained from FSI forest density classes.

#### **Materials and Methods**

#### Study area

The EG (between 10°00′00″–13°00′00″N and 77°50′00″–79°10′00″E) is a discontinuous mountain range running almost parallel to the east coast of India and constitutes the watersheds of many rivers. In Tamil Nadu, the EG start from the Jawadi Hills and extend up to the Alagar






Fig 1. Study area—Eastern Ghats of Southern India.

doi:10.1371/journal.pone.0147541.g001

Hills (Fig 1), comprising 13 major hills in total (Jawadi, Elagiri, Shevaroy, Chitteri, Kalrayan, Bodamalai, Kolli, Pachaimalai, Semmalai, Aiyalur, Karandamalai, Sirumalai, and Alagar). The hills of the EG in Tamil Nadu range from 70 km² to 1860 km² in area, and the altitude in this region ranges from 180 m above mean sea level (MSL) to 1650 m above MSL. The mean minimum and maximum temperatures are 17°C and 33°C, respectively, and the average annual rainfall is 800–1600 mm. The major soil types are entisols, inceptisols, and alfisols. Geologically, the EG consist of charnockite with minor bands of pyroxin granulate and magnetite quartzite. The tributaries of many perennial rivers like the Pennaiyar, Palar, Vellar, Cauvery, and Vaigai originate from these hills. These hills are still rich in biodiversity, despite heavy exploitation, and they contain more than 960 species of angiosperm and gymnosperm. There are many endemic, endangered, and medicinally important species distributed within these hills [34, 35].

The forests are mostly degraded due to the repeated felling of trees in the past; presently, three major types of forests exist, with different compositions and canopy densities, as explained in detail below.



Tropical dry evergreen forest. This forest type is referred to locally as 'shola forest' and it occurs in six hills: Jawadi, Elagiri, Sherveroy, Kolli, Pachaimalai, and Sirumalai, which are generally located 1000 m above MSL. The general composition includes *Memecylon edule*, *Neolitsea scrobiculata*, *Persea macrantha*, *Memecylon umbellatum*, *Elaeocarpus serratus*, *Syzygium cumini*, *Canarium strictum*, *Artocarpus heterophyllus*, *Artocarpus hirsuta*, and *Alangium salvifolium*, while the dominant shrub species are *Psychotria subintegra*, *Mesa indica*, *Glycosmis mauritiana*, *Phyllanthus wightianus*, *Tarenna asiatica*, *Lantana camara*, and *Clausena dentata*, among others.

**Tropical dry deciduous forest.** This forest type is widely distributed throughout the 13 hills and is found in all types of topography, such as valleys, plateaus, and foothills from 400 m up to 1000 m above MSL. The species composition in the upper and middle slopes consists of *Albizia odoratissima*, *Bridelia retusa*, *Chloroxylon swietenia*, *Terminalia chebula*, *T. bellirica*, *Tectona grandis*, *Schleichera oleosa*, *Sapindus emarginatus*, *Emblica officinalis*, *Pterocarpus marsupium*, *Dalbergia latifolia*, *Gyrocarpus asiaticus*, *and Moringa oleifera*, among others, while the dominant shrubs are *G. mauritiana*, *L. camara*, *Pterolobium hexapetalum*, *Ziziphus oenoplia*, and *Acacia pennata*, among others.

**Southern tropical thorn forest.** This forest type is the most affected and degraded in nature and is distributed throughout the entire lower hills and foothills at <400 m above MSL, as it appears near adjoining villages. The main tree species are *Acacia planifrons*, *Acacia leucophloea*, *Albizia amara*, *Azadirachta indica*, *Streblus asper*, and *Wrightia tinctoria*, among others, and this forest type also contains heavy impenetrable thorny species such as *Carissa carandas*, *Dichrostachys cineraria*, *Pterolobium hexapetalum*, *P. indicum*, *Randia dumetorum*, *Toddalia asiatica*, *Euphorbia antiquorum*, and *Z. oenoplia*, among others.

Permission to conduct experiments was obtained from the Principle Chief Conservator of Forest, Department of Forestry, Tamil Nadu, India. Care was taken to collect soil and floristic samples without damaging the existing forest.

## Forest cover density

Forest cover density (FCD) maps of the study region were provided by the FSI as  $1^{\circ} \times 1^{\circ}$  tiles for academic and research purposes. Three FCD tiles of the EG in 2012 were procured on payment for this study. The spatial resolution is 24 m.

## Soil sampling and analysis

Forests were divided into 5 km  $\times$  5 km grids, excluding private land and rocky patches. In each grid, a one-soil sample location was randomly chosen and its geographical coordinate was noted. In total, 408 soil samples were collected during January–February 2012. All samples were shade-dried, pulverised with an agate mortar, and then sieved (0.2 mm). The samples were analysed following standard methods (Table 1). For microbiological studies, soil samples were collected in sterile bags and transported to the laboratory, where they were stored at 4°C until needed for further processing.

# Forest soil quality index

In this study, the forest soil quality was examined using 22 soil physico-chemical and biological properties. These 22 parameters were then used to calculate the forest soil quality index (FSQI). The FSQI can effectively interpret multivariate data sets, which are characteristic of soil data; a single score is able to identify an overall trend among potentially conflicting indicators [49]. The FSQI was calculated using the following steps: 1) identification of a minimum data set (MDS) of indicators, or variables that best represent soil functions associated with the



Table 1. Methods for soil physico-chemical and microbiological analysis.

Estimation	Method
Physical Properties	
Bulk Density (BD)	Core method [36]
Chemical Properties	
Soil pH	Potentiometry 1: 25 (soil-water) [37]
Electrical Conductivity (EC)	Conductometry (1:2 soil-water suspension) [37]
Total Nitrogen (TN)	CHSN/O Elemental Analyser [38]
Total Potassium (TK)	Flame Photometry [39]
Total Phosphorus (TP)	Pemberton method [40]
Total Trace Elements (Fe, Mn, Zn, Cu)	HCl extract in Atomic Absorption Spectrophotometer (Perkin Elmer model 380) [41–43]
Biological properties	
Soil Organic Carbon (SOC)	CHSN/O Elemental Analyser [38]
Available Nitrogen (Ava N)	Alkaline permanganate method [44]
Available Phosphorus (Ava P)	Olsen's extractant (0.5 M NaHCO3) at pH 8.5 [45]
Available Potassium (Ava K)	Neutral N NH4OAc method [46]
Microbial Biomass Carbon (MBC)	Fumigation extraction method [47]
Microbial Biomass Nitrogen (MBN)	Fumigation extraction method [47]
<b>Total and Functional Microbes</b>	Pour plate method [48]

doi:10.1371/journal.pone.0147541.t001

selected management goal, 2) normalisation of the MDS indicators, and 3) integration of the indicator scores into an overall index of soil quality, which are each described below.

MDS identification. Principal component analysis (PCA) was employed in IBM SPSS version 20.1 (IBM SPSS statistics for windows, version 22.0. Armonk, NY: IBM Corp) [50] as a data reduction technique to identify the appropriate indicators without losing the vital information [51, 52]. The principle components (PCs) are linear combinations of variables that account for the maximum variance in a data set and are more accurate descriptors of the data than the variables themselves. For this study, PCs with eigenvalues  $\geq$ 1.0 (>10% of the total variance explained) was taken into consideration for indexing. Highly weighed variables (eigenvectors  $\geq$  ± 0.7) within each PC were considered significant and retained for the FSQI [24, 53].

**Normalisation of the indicators.** As the selected soil parameters were measured in heterogeneous units, they were standardised to unit-less values between 0 and 1 [32]. Depending on whether a higher value was constructive or detrimental to soil quality, indicators were arranged in ascending or descending order [32, 33]. The resultant outputs were called *Indicator Scores*. These normalised scores were calculated using the following two formulas. Eq 1 was used for indicators that tended to increase soil quality as the values increased:

$$S_{i \text{ normalized}} = \frac{S_i - S_{i \text{ min}}}{S_{i \text{ max}} - S_{i \text{ min}}} \tag{1}$$

<u>Eq 2</u> was used for indicators that tended to decrease soil quality as the values increased:

$$S_{i \text{ normalized}} = \frac{S_{i \text{ max}} - S_{i}}{S_{i \text{ max}} - S_{i \text{ min}}}$$
 (2)

where  $Si_{max}$  is the maximum value of the soil indicator variable, Si is the actual observed value, and  $Si_{min}$  is the minimum value of the soil indicator variable.



Table 2.	Forest cover	density of the	Eastern Ghats.
I abic 2.	I OICGLOOVCI	actionly of the	Lastern Gnats.

Class Names	Forest cover area (ha)	Percentage
VERY DENSE FOREST	66547.5	15.7
MODERATELY DENSE FOREST	160585.0	38.0
OPEN FOREST	142630.4	33.8
DEGRADED FOREST	7611.0	1.8
OTHERS	45256.7	10.7
TOTAL	422630.6	100.0

doi:10.1371/journal.pone.0147541.t002

**Calculating the PCA-FSQI.** After normalisation, the MDS variable of each observed value was weighed using the PCA-derived outcome. The weighing factor was calculated via dividing the percentage of each selected MDS variable by the total percentage of variation in all PCs with eigenvectors >1. The scored indicator for each observation was calculated with the following equation:

$$PCA FSQI = \sum Wi * Si$$
 (3)

where *Wi* is the PCA-derived weighed factor and *Si* is the normalised scores of the indicator *i*. Higher index scores indicate better soil quality.

# Spatial data analysis

Spatial autocorrelation was performed using a semi-variogram that considered all sampling points to test the spatial autocorrelation between points. The global Moran's I test was used to estimate the overall degree of spatial autocorrelation and the analysis returns a single value, which is applicable to the entire study area [54].

## FSQI threshold and spatial mapping

To fix the FSQI threshold, the mean and 95% confidence interval (CI) of the 408 samples were calculated [55,56]:

$$FSQI \ threshold \ value = Mean - CI$$
 (4)

The FSQI value obtained for each sample was interpolated using the inverse distance weighted method in Arc GIS 9.3. The interpolated values were grouped into two categories: degraded (FSQI < threshold value) and non-degraded (FSQI > threshold value).

## Matrix analysis and ground truth verification

Matrix analysis is a spatial process where two raster data can be verified pixel-by-pixel for its agreement. Each pixel in the resultant image will carry information from both raster. Through this method the FSI forest cover density map and the soil quality index maps were analyzed. Each pixel of resultant image showed the forest cover density and its corresponding SQI index value.

The matrix analysis between the FSQI and the FCD of FSI was performed using ERDAS IMAGINE 2011 software (Erdas Imagine, Hexgon geospatial, ink GA,USA) [57] Ground truth verification was also performed for both the FCD and FSQI to verify whether the classified areas conformed to the field data across all forest density classes.






Fig 2. Forest cover density of the Eastern Ghats.

doi:10.1371/journal.pone.0147541.g002



Table 3. Soil physico-chemical and biological characteristics of the hills of the Eastern Ghats.

Properties	Alakar	Aiyalur	Bodamalai	Chitteri	Elagiri	Jawadi	Kairayan	Karandai	Kolli	Pachai	Semmalai	Shevaroy	Sirumalai
C (%)	0.05-2.12	0.25-1.18	0.31–1.62	0.17-3.58	0.06-2.03	0.32-2.31	0.6–3.74	0.32-2.02	0.15–3.8	0.16–3.16	0.18-1.81	0.09-3.63	0.13-2.41
(%) N	0.1-0.45	0.07-0.53	0.04-1.02	0.09-1.33	0.11-1.17	0.1-0.9	0.11-1.43	0.04-0.85	0.02-0.85	0.01-0.65	0.11-0.43	0.05-1.30	0.07-1.02
MBC (mg kg <sup>-1</sup> )	72.9-616	142-381	80-348	40-634	72–601	72–616	111–515	110–381	83-532	83-288	132-515	24-617	130-522
MBN (mg kg <sup>-1</sup> )	67-340	26–386	57-493	39–501	43-471	45-494	23–501	45-452	24-424	56-471	94-361	11–501	56-469
TB (cfu mL <sup>-1</sup> )	1.2×10-3- 7×10 <sup>7</sup>	1.8×10 <sup>3</sup> -3.4×10 <sup>5</sup>	1.8×10 <sup>3</sup> -3.4×10 <sup>5</sup> 1.6×10 <sup>3</sup> -4.5×10 <sup>6</sup>	1.9×10 <sup>3</sup> -4.5×10 <sup>5</sup>	2.5×10 <sup>3</sup> –1.2×10 <sup>5</sup>	2.4×10 <sup>3</sup> –4.5×10 <sup>5</sup>	1×10 <sup>4</sup> -4×10 <sup>6</sup>	4.2×10³-8.6×10 <sup>5</sup>	2.7×10 <sup>3</sup> –1.2×10 <sup>5</sup>	1.5×10³–3.1×10 <sup>5</sup>	2.4×10 <sup>3</sup> –6.5×10 <sup>4</sup>	1.4×10³-5.2×10 <sup>5</sup>	2.9×10³–5×10 <sup>6</sup>
TF (cfu mL <sup>-1</sup> )	2.5×103- 1.4×10 <sup>4</sup>	2.6×10³-3.5×10 <sup>5</sup>	2.6x10 <sup>3</sup> -3.5x10 <sup>5</sup> 1.1x10 <sup>3</sup> -3.5x10 <sup>5</sup> 1.2x10 <sup>3</sup> -3.1x10 <sup>5</sup>	1.2×10³-3.1×10⁵	1×10 <sup>4</sup> –3.1×10 <sup>6</sup>	3.2×10 <sup>3</sup> –4.6×10 <sup>5</sup>	2.2×10 <sup>3</sup> -3.6×10 <sup>5</sup>	3.1×10³-3.6×10 <sup>5</sup>	3.1×10 <sup>3</sup> –3.6×10 <sup>5</sup>	3.2×10³–6.6×10 <sup>5</sup>	3.1×10 <sup>3</sup> –8.6×10 <sup>5</sup>	4.2×10³-3.6×10 <sup>5</sup>	4.2×10³–2.1×10⁵
TA (cfu mL <sup>-1</sup> )	1×10-3- 3.1×10 <sup>5</sup>	1.5×10³–3×10⁴	1×10 <sup>3</sup> -4×10 <sup>5</sup>	1×10³–5×10⁴	2×10³-1×10⁴	4×10³–5×10⁵	1×104-4×10 <sup>5</sup>	2×10³-6×10⁴	2×10³-1×10⁴	1×10³–3×10⁴	4×10³–5×10⁴	4×10³–2×10 <sup>5</sup>	2×10³-5×10⁵
NFB (cfu mL-1)	2.5×103- 1.4×10 <sup>4</sup>	2.6×10³-3.5×10 <sup>5</sup>	2.6×10³–3.5×10 <sup>5</sup> 1.1×10³–3.5×10 <sup>5</sup>	1×10³–3×10 <sup>5</sup>	1×10 <sup>4</sup> –3×10 <sup>6</sup>	3×10³-4×10 <sup>5</sup>	2×10³-3×10 <sup>5</sup>	3×10³-4×10 <sup>5</sup>	1×10³–3×10 <sup>5</sup>	2×10³-6.6×10 <sup>5</sup>	1×10³–6×10 <sup>5</sup>	2×10³–3×10 <sup>5</sup>	2×10³-1×10⁵
CM (cfu mL <sup>-1</sup> )	1.2×10-3- 7×10 <sup>5</sup>	1.8×10 <sup>3</sup> -3.4×10 <sup>4</sup>	1.8×10 <sup>3</sup> -3.4×10 <sup>4</sup> 1.6×10 <sup>3</sup> -4.5×10 <sup>5</sup>	3×10 <sup>3</sup> -4.5×10 <sup>4</sup>	5×10³-1.2×10⁴	4×10³-4×10⁵	1×10 <sup>4</sup> -4×10 <sup>5</sup>	1.2×10³-2.3×10 <sup>5</sup>	2×10³-1×10⁴	1×10³–3×10⁵	2×10³–6×10⁴	1×10³–2×10⁴	9×10³-5×10⁴
PSB (cfu mL-1)	2.5×103- 1.4×10 <sup>4</sup>	2.6×10³-3.5×10 <sup>5</sup>	2.6×10³–3.5×10 <sup>5</sup> 1.1×10³–3.5×10 <sup>5</sup>	1.2×10 <sup>3</sup> –3.1×10 <sup>4</sup>	1×10 <sup>4</sup> -3.1×10 <sup>5</sup>	3.2×10 <sup>3</sup> -4.6×10 <sup>5</sup>	2.2×10 <sup>3</sup> –3.6×10 <sup>4</sup>	3.1 ×101 <sup>3</sup> -3.6×10 <sup>4</sup>	3.1×10 <sup>3</sup> -3.6×10 <sup>4</sup>	3.2×10³-6.6×10⁴	3.1×10 <sup>3</sup> –8.6×10 <sup>5</sup>	4.2×10³–3.6×10⁴	4.2×10³-2.1×10 <sup>5</sup>
Hd	6.2-6.7	6.1–6.7	6.1–6.9	6.1–6.7	6.5-6.71	6.3–6.8	6.1–6.7	6.1–6.5	6.0-6.4	6.2–6.7	6.2–6.7	6.3–6.9	6.0-6.4
EC	0.11-0.12	0.12-0.13	0.12-0.14	0.11-0.13	0.11-0.13	0.11-0.12	0.12-0.14	0.11-0.12	0.11-0.13	0.11-0.12	0.11-0.13	0.14-0.16	0.14-0.16
TK (%)	9-21	10-21	10-25	21–23	9-19	12–23	13-31	10-34	20-30	16–24	17–31	12–23	16-20
TP (%)	15-20	17–30	21–34	19–27	11–23	14–17	12–28	15–32	12–29	16–31	14-17	13-31	12-21
Fe (mg kg <sup>-1</sup> )	2.06-4.01	2.05-4.91	1.73–3.4	1.7–2.56	1.12-4.5	1.23-5.6	1.78-2.39	1.63-4.51	1.34-3.67	.1.29–3.8	1.45–2.6	1.6-4.11	1.57-4.78
Mn (mg kg <sup>-1</sup> )	4.7-9.3	5.1–9.8	3.89-9.0	1.9–10.1	2.3-8.9	3.4-9.81	2.46-8.91	3.13-8.93	3.6–8.9	4.12-9.23	3.81-8.92	3.79-10.1	4.51–7.67
Zn (mg kg <sup>-1</sup> )	1.23-2.45	1.11–2.89	1.45–2.39	1.34-2.91	1.11–3.1	1.9–2.7	1.3-2.0	1.3–2.1	1.4–2.5	1.4–3.6	1.3-2.2	1.31–2.6	1.5-2.7
Cu (mg kg <sup>-1</sup> )	1.2-2.1	1.2–2.4	1.24–2.5	1.31-4	1.34–2.1	1.21.2.1	1.67–2.3	1.11–2.2	1.31–2.0	1.1–2.0	1–2.4	1.3–3.2	1.4-2.4
Ava N (kg ha <sup>-1)</sup>	121-392	153-212	158-234	123–301	213-312	124–312	214-378	156-301	145-294	156-345	156-291	239-315	120-234
Ava P (kg ha <sup>-1</sup> )	15-25	17.9–59	23-34	28–79	19-67	14–65	34.5–98	13.62-78.1	11.34-70.23	19.4–72.3	23.1–78	32.4-90.1	12.7–36.7
Ava K (kg ha <sup>-1</sup> )	123-234	189–210	120–534	149–560	137–589	131–601	123-467	290-601	312-421	172–390	138-560	278–670	145–346

SOC: soil organic carbon, TN: total nitrogen, MBC: microbial biomass carbon, MBN: microbial biomass nitrogen, TB: total bacteria, TF: total fungi, TA: total actinomycetes, NFB: nitrogen fixing bacteria, CM: cellulytic microbes, PSB: phosphate solubilising bacteria, EC: electrical conductivity, TK: total potassium, TP: total phosphorus, Fe: iron, Mn: manganese, Zn: zinc, Cu: copper, Ava N: available nitrogen, Ava P: available phosphorus, Ava K: available potassium.

doi:10.1371/journal.pone.0147541.t003



Table 4. Soil parameters incorporated in the principal component analysis.

Properties			Component		
	1	2	3	4	5
Eigen Value	3.538	1.607	1.086	1.009	0.871
Variance (%)	39.31	17.86	12.067	11.217	8.910
Cumulative (%)	39.31	57.17	69.237	80.454	89.364
Eigen Vectors					
Bulk Density	.011	.060	.341	.213	.324
SOC	.905	.211	142	115	.211
TN	.718	532	.313	.175	.456
MBC	.826	.168	126	041	.567
MBN	.680	561	.263	.249	.021
ТВ	.751	.368	259	113	.121
TF	.212	.031	.191	.304	.291
TA	.447	.152	.596	437	.128
NFB	.310	.775	200	.527	.235
CM	239	.011	.456	.435	.146
PSB	.230	.111	.231	.342	.417
pH	.141	.212	.303	003	.239
EC	.165	.342	574	411	.454
TK	.344	.136	.343	.291	.157
TP	.034	.458	.023	.359	.478
Fe	.450	122	228	452	.234
Mn	211	.042	245	.127	.458
Zn	220	.124	.284	.116	.171
Cu	.175	.571	.466	.376	.342
Ava N	.178	.377	.457	.125	.561
Ava P	.132	.375	.399	.274	.231
Ava K	.231	.452	.371	.312	.341

SOC: soil organic carbon, TN: total nitrogen, MBC: microbial biomass carbon, MBN: microbial biomass nitrogen, TB: total bacteria, TF: total fungi, TA: total actinomycetes, NFB: nitrogen fixing bacteria, CM: cellulytic microbes

PSB: phosphate solubilising bacteria, EC: electrical conductivity, TK: total potassium, TP: total phosphorus, Fe: iron, Mn: manganese, Zn: zinc, Cu: copper, Ava N: available nitrogen, Ava P: available phosphorus, Ava K: available potassium.

doi:10.1371/journal.pone.0147541.t004

In total there were 8 classes and for each class 25 points were randomly created and were verified on the ground using GPS. Of the total 200 quadrates, each location the vegetation data on trees and shrubs were enumerated through quadrate of 20X 20 m size covering 13 hills.

## **Results and Discussion**

## FCD status in the EG

Table 2 details the areas of the different FCD categories. Of the site's 422630.6 ha, approximately 45256.7 ha (10.7%) is non-forest. The effective forest area covers 377373.9 ha, with the majority categorised as moderately dense or open. Ground truth verification revealed considerable similarities between the open and the degraded forest cover categories; the floristic compositions were frequently identical, with sprawling growth of shrubs and bushes indicating heavy



degradation (<u>Table 2</u>, <u>Fig 2</u>). Additionally, most of the areas occupied by scrub forests were classified as dense or moderately dense, indicating that the FSI data may be misleading and may fail to provide an accurate description of forest degradation.

#### **FSQI**

Soil physico-chemical and biological analysis. The physico-chemical and biological properties of the 408 soil samples are provided in <u>Table 3</u>. While most of the measured properties exhibited high variation, the soil pH was uniformly low. Across the 13 hills, Aiyalur displayed the lowest values for soil organic carbon (SOC), total nitrogen (TN), microbial biomass carbon (MBC), microbial biomass nitrogen (MBN), total bacteria (TB), and functional microbes. In contrast, the highest values of the measured properties were found in the Chitteri samples, which also exhibited greater variation in biological properties. Our results support previous findings that indicate that the variability in these factors is due to changes in climate, season, geographical location, and anthropogenic interference [24, 58–62].

# Spatial data analysis

A fitted semi-variogram of the FSQI in the study area had a nugget effect of 0.55 and a total sill of 0.74. The high nugget-sill ratio of 74% reveals that strong local scale variation prevails. The spatial autocorrelation that was performed based on the global Moran's I of the FSQI was 0.216, which was low but still statistically significant (p = 0.000014). This indicates that spatial clusters of the FSQI may be found across the study area with similar values (S1 Fig).

**MDS.** The results of our PCA on the 22 soil parameters (<u>Table 3</u>) revealed that the first four PCs explained approximately 80.45% of the soil variability. Further, the first two PCs accounted for approximately 57.17% of the variance, strongly indicating that the highly weighed variables in these two components can be used to assess the soil quality of this area.

Under PC1, these variables were SOC, MBC, TB, and TN; under PC2, NFB was the highest weighed value. These five properties indicate the level of soil disturbance since biological properties react relatively quickly to small changes in the soil environment and it has been broadly

Table 5. Status of the forest soil quality index (FSQI) and areas of different forest types in the hills of the Eastern Ghats.

Hills	Total Area (ha)	Forest Co	ver Density (ha)			FSQI	
		Degraded	Non Degraded	Mean	Range	Degraded	Non Degraded
						(ha)	(ha)
Alagar	6141.7	408	5733.7	0.38	0.13-0.67	4237.3	1904.5
Aiyalur	10424.1	1686	8738.1	0.32	0.21-0.50	10022.6	401.7
Bodamalai	7560.3	256.5	7303.8	0.36	0.13-0.77	4975.8	2584.5
Chitteri	49329.6	597.5	48732.1	0.69	0.17-1.24	5518.6	43810.6
Elagiri	6433.2	9.2	6424	0.46	0.13-1.29	2413.6	4019.6
Jawadi	146446.6	1788.5	144658.1	0.43	0.13-0.87	63062.9	83383.8
Kalrayan	41238.4	203.3	41035.1	0.56	0.20-1.38	14783.2	26455.2
Karandai	7877.8	360.2	7517.6	0.37	0.16-0.78	6448	1429.7
Kolli	23406.2	348.1	23058.1	0.42	0.09-0.81	12025	11381.2
Pachaimalai	34917.2	177.8	34739.4	0.41	0.10-1.05	17356.2	17560.9
Semmalai	6246.5	441.2	5805.3	0.39	0.16-0.58	3354.2	2892.3
Shevaroy	23957.2	919.2	23038	0.46	0.01-1.42	11330.6	12626.8
Sirumalai	13395.1	415.5	12979.6	0.46	0.17-1.20	4590.5	8804.6
Total	377373.9	7611	369762.9			160118.5	217255.4

doi:10.1371/journal.pone.0147541.t005






Fig 3. Forest soil quality index of the Eastern Ghats.

doi:10.1371/journal.pone.0147541.g003



reported that any alteration in soil management and land use is reflected through soil properties [63–64]. Thus, this study has considered these five biological properties as the MDS for identifying the soil quality status in all 13 hills (<u>Table 4</u>, <u>S2 Fig</u>). We then used <u>Eq 3</u> to calculate the weighting factors for the variables in each PC, which allowed us to obtain the FSQI.

$$\begin{aligned} \text{PCA} - \text{FSQI} &= \left[ (0.393 \text{ SOC} + 0.393 \text{ TN} + 0.393 \text{ MBC} + 0.393 \text{ TB} + 0.178 \text{ NFB}) / 1.75 \right] \\ \text{PCA} - \text{FSQI} &= 0.224 \text{ SOC} + 0.224 \text{ TN} + 0.224 \text{ MBC} + 0.224 \text{TB} + 0.101 \text{ NFB} \end{aligned} \tag{5}$$

As previously stated, low FSQI values indicate degraded soil and high values indicate non-degraded soil. Soil from Chitteri Hill exhibited the highest FSQI (0.69), followed by soil from Kalrayan Hill (Table 5). Aiyalur and Bodamalai possessed soils with the lowest FSQI values: 0.32 and 0.36, respectively (Fig 3). The mean FSQI of all EG sites in Tamil Nadu was only 0.44, very close to the threshold value. Based on the FSQI, approximately 42.4% of the study area consists of degraded forests and 57.6% consists of non-degraded forests.

However, ground truth verification revealed a  $\pm 13\%$  deviation from the FSQI-classified status, due to variation in vegetation within a particular forest density class. For instance, the presence of riparian flora along streams in the open and degraded forests meant they were apparently classified as dense forests.

Based on the FSQI, the highest soil degradation (>96.1%) was found in Aiyalur forests, followed by Bodamalai and Karandamalai (81.9%) and Alagar (69%) forests (Table 5). Furthermore, the lowest soil degradation occurred in Chitteri, with 11.2% of the area classified under the degraded category. In addition, the soil quality of Sirumalai, Kalrayan, and Elagiri was found to be less affected by forest degradation, with 34.3%, 35.8%, and 37.5% of their total areas classified as degraded, respectively (Table 5).

The matrix analysis of the FCD and FSQI (Table 6) revealed more details about the degradation status. Forests in all four FCD categories have both degraded and non-degraded soil. In the degraded forest category, 74.2% possesses degraded soil, and in the open forest cover category, approximately 52% possesses degraded soil. In the moderately dense and very dense categories, degraded soil occurs in 38.9% and 26.7% of the forest areas, respectively. Across all forests, only 154474.2 ha (40.9%) were classified as non-degraded by both the FCD and FSQI (Fig 4). Based on previous reports of the EG, soil quality degenerated wherever severe anthropogenic exploitation slowly modified primary forest vegetation to secondary vegetation [8, 10]. Such forests tend to consist of patchily distributed, low scrub that degenerates into spare bushes with occasional, stunted tree growth predominantly occupied by *L. camara* bushes that suppress the natural regeneration of other species [3, 13, 65–68].


Table 6. Matrix analysis between the forest cover density and forest soil quality index.

Class Name		FSI and FSQI Matrix			
	cover (ha)	Degraded (ha)	Non-Degraded (ha)	Degraded (%)	Non Degraded (%)
VERY DENSE FOREST	66547.5	17741.4	48806.1	26.7	73.3
MODERATELY DENSE FOREST	160585.0	62515.5	98069.5	38.9	61.1
OPEN FOREST	142630.4	74217.3	68413.1	52.0	48.0
DEGRADED	7611.0	5644.3	1966.7	74.2	25.8
TOTAL	377373.9	160118.5	217255.4	42.4	57.6

FSI: Forest Survey of India.

doi:10.1371/journal.pone.0147541.t006





 $\label{eq:Fig-4.} \textbf{Fig-4. Matrix} \ \text{analysis between the forest cover density and forest soil quality index.}$ 

doi:10.1371/journal.pone.0147541.g004



Table 7. Matrix-based species composition in degraded and non-degraded forests located in the lower, middle, and upper slopes, as verified through ground truth surveys.

Forest types	Species Composition					
	Degraded (Low FSQI)	Non-degraded (High FSQI)				
Thorn forest	Tree species: Moringa oleifera,	Tree species: Azadirachta				
(Lower elevation	Gyrocarpus americanus, Wrightia	indica, Acacia sundra,				
<400 m)	tinctoria, Acacia ferruginea, acacia	Hardwickia binata, Albizia				
	leucophloea, Acacia sundra,	amara, Chloroxylon swietenia,				
	Acacia planifrons, Albizia	Acacia leucophloea, Zizyphus				
	Odoratissima	xylopyrus, Moringa oleifera,				
	Shrubs: Lantana camara, Strychnos	Wrightia tinctoria, Feronia				
	potatorum, Pterolobium indicum,	elephantum, Streblus asper				
	Acacia concinna, Acacia latronum,	Shrubs: Dichrostachys Randia				
	Acacia intsia, Dichrostachys cineraria,	cineraria, Dodonaea viscosa,				
	Dodonaea viscosa, Randia dumetorum,	dumetorum, Clausena dentata,				
	Clausena dentata, Commiphora	Commiphora caudata, Carissa				
	Caudate	carandas, Acacia intsia, Acacia				
		Latronum				
Deciduous forest	Tree species: Albizia odoratissima,	Tree species: Shorea talura,				
Middle elevation	Canthium didymum, Anogeissus	Anogeissus latifolia, Terminalia				
(400–1000 m)	latifolia, Feronia elephantum, Ziziphus	tomentosa, Terminalia				
,	xylopyrus, Erythroxylon monogynum	paniculata, Albizia lebbeck,				
		Albizia odoratissima, Terminalia				
	Shrubs: Lantana camara, Strychnos	arjuna, Hardwickia binata,				
	nuxvomica, Citronella grass,	Diospyros montana,				
	Pterolobium indicum, Acacia intsia,	Stereospermum suaveolens,				
	Pavetta indica, Acacia pennata,	Pterocarpus marsupium, Adina				
	Pterolobium indicum, Acacia intsia,	cordifolia, Bridelia retusa,				
	Ziziphus oenoplia	Dalbergia latifolia, Ficus				
		microcarpa, Grewia tiliifolia,				
		Premna tomentosa, Santalum				
		album, Sapindus emarginatus,				
		Schleichera oleosa				
		Shrubs: Pterolobium indicum,				
		Acacia intsia, Ziziphus oenoplia,				
		Semecarpus anacardium,				
		Ziziphus mauritiana, Lantana				
		Camera				
Evergreen and	Tree species: Memecylon edule,	Tree species: Manilkara hexandra,				
semi-evergreen	Memecylon umbellatum, Persea	Memecylon edule, Alseodaphne				
(Upper elevation >	macrantha, Melia dubia, Syzygium	semecarpifolia, Alstonia scholaris,				
1000 m)	montana, Trema orientalis	Bischofia javanica, Celtis tetrandra,				
1000 111,	montana, monta onemaile	Chukrasia tabularis, Cinnamomum				
	Shrubs: Lantana camara, Embelia indica,	macrocarpum, Elaeocarpus				
	basaal, Atlantia monophylla, Clausena	lanceifolius, Litsea oleoides, Mallot				
	dentata, Glycosmis mauritiana, Mesa	·				
	Psychotria subintegra, Phyllanthus	philippensis, Mangifera indica, Meliosma simplicifolia, Memecylon				
		Meliosma simplicifolia, Memecylon				
	wightianus, Tarenna asiatica,	edule, Memecylon umbellatum,				
	Adenostemma lavenia, Balanophora	Myristica dactyloides, Neolitsea				

(Continued)



Table 7. (Continued)

Forest types	Species Composition					
	Degraded (Low FSQI)	Non-degraded (High FSQI)				
	fungosa, Dorstenia indica	scrobiculata, Nothopegia				
		colebrookeana				
		Shrubs: Gymnema kollimalayanum,				
		Embelia basaal, Atlantia monophylla,				
		Clausena dentata, Glycosmis				
		Mauritiana				

FSQI: forest soil quality index.

doi:10.1371/journal.pone.0147541.t007

Although there is some correlation between the forest density and the percentage of degradation estimated by the FSI, the present study clearly addresses the inconsistencies between the density classes of the FSI and FSQI (<u>Table 6</u>). Forests classified by the FSI as very dense forests, moderately dense forests, and open forests contain soils below the FSQI threshold, while high FSQI values occurred in forests classified as degraded by the FSI. Overall, an imprecise relationship exists between the FSI density classes and the FSQI. We conclude that the NDVI-based forest cover assessment cannot comprehensively describe the real degradation status of forests.

The species composition of various slope categories in the EG (<u>Table 7</u>) reveals two distinct patterns that correspond to the FSQI rather than the FCD. Wherever the FSQI value is below the threshold, thorny scrubs instead of trees generally dominate the floristic composition. Moreover, when the FSQI value is higher than the threshold, dense tree growth dominates, with a very limited occurrence of herbs and shrubs (<u>Table 7</u>). A large contributor to these patterns is the invasive species *L. camara* [65–68], which was extremely prominent in areas with low FSQI values and appeared to prevent further proliferation of native species. Other factors that aggravate soil degradation and hinder plant growth include anthropogenic activity, environmental factors such as erratic rainfall and extreme heat during summer, as well as geological features like steep slopes [10].

In summary, our study lends further support to previous findings that demonstrate the impreciseness of NDVI-based forest cover assessments, which fail to differentiate between natural forests and plantations [69]. Our results strongly suggest that satellite-based assessment of canopy cover gives an incomplete idea of forest status. However, we also found that the FSQI can be an effective tool for categorising forest degradation with greater accuracy.

#### Conclusions

This study pioneers the use of the FSQI for estimating forest degradation in India, which currently depends on satellite-based canopy cover data to assess forest status. Using the FSQI, degradation was found in all FSI forest cover density classifications. Combining the FSQI with the FSI density classes resulted in a more detailed picture of forest degradation. Hence, the FSQI is an ideal tool for studying the degradation status distributed across different forest types and density classes. The present study reveals that the FSQI categorised approximately 42.4% of the total forests examined as degraded, whereas the FCD only categorised approximately 1.8% of the forests as degraded. Therefore, relying on satellite data alone may no longer provide an accurate representation of forest status, given the increase in environments that may appear as undisturbed forest without a more sensitive investigation. We suggest that the FSQI can be used as supplemental data along with the FSI forest cover density to efficiently categorise forest



degradation and to address this continuing environmental issue, especially in developing countries. Management of degraded forests involves meticulous reforestation work to restore the forests, which could be achieved only by prioritising the area based on the FSQI.

# Supporting Information

S1 Fig. Semi- variogram model of Eastern Ghats FSQI distribution using ordinary kriging based on 408 samples collected from the Eastern Ghats.

(TIF)

S2 Fig. Biplots for PCA Analysis.

(TIF)

# **Acknowledgments**

The authors sincerely thank the Tamil Nadu forest department for permitting us to perform field surveys and soil samplings. We are extremely thankful for the forest staff for their valuable assistance and their continuous support during the study period. Thanks are due to the Forest Survey of India for providing the forest density map for the entire Eastern Ghats. We thank Anna University for providing the laboratory facilities to perform the soil analysis.

## **Author Contributions**

Conceived and designed the experiments: AR SJ. Performed the experiments: PR. Analyzed the data: PR. Contributed reagents/materials/analysis tools: PR RG. Wrote the paper: AR PR PD SI.

### References

- 1. Cleghorn H. The forests and gardens of south India. London: W.H. Allen & Company; 1861.
- 2. Puri GS, Meher-Homij VM, Gupta RK, Puri S. Forest ecology. Oxford: IBH Publishing Company; 1960.
- 3. Adolphus WHW. Consolidated working plan for the Salem forest division. Government Press. Madras Presidency, Madras. 1932.
- Thankam K. Working plan for the Chengalpattu forest division. Government of Tamil Nadu. Government Press, Madras. 1959.
- Harikrishnan I. Working plan for the Salem forest division. Government of Tamil Nadu. Government Press. Madras. 1967.
- 6. Bhat DM, Murali KS, Ravindranath NH. Formation and recovery of secondary forests in India: a particular reference to Western Ghats in South India. J Trop For Sci. 2001; 13: 601–620.
- Prasad R. Sustainable forest management for dry forests of South Asia. In: Cheng TL, Durst PB, editors. Development of national-level criteria and indicators for the sustainable management of dry forests of Asia: background papers. Thailand: RAP Publishers/FAO; 2000. p. 27–74.
- Ramachandran A, Jayakumar S, Haroon ARM, Bashkaran A. Carbon management in forest floor–An agenda of 21st century in Indian forestry scenario. Indian For. 2007; 133: 25–40.
- 9. Stebbing EP. The forests of India, vol I. London: John Lane/The Bodley Head Limited; 1922.
- Jayakumar S, Ramachandran A, Baskaran G, Heo J. Forest dynamics in the Eastern Ghats of Tamil Nadu, India. Environ Manage. 2009; 43: 326–345. doi: 10.1007/s00267-008-9219-y PMID: 18953598
- Jayakumar S, Ramachandran A, Bhaskaran G, Heo J, Kim WS. Site suitability assessment for Joint Forest Management (JFM): a geospatial approach. Korean J Remote Sens. 2007; 23: 473–481.
- Ministry of Environment and Forests. Forest survey of India: State of forest report. Dehradun: Government of India; 2003.
- Ramachandran A, Jayakumar S, Arockiasamy DI. Need for reclassification of forest types in India using high-resolution satellite data and field study for better management. Indian For. 2005; 131: 3–12.
- Sagar R, Singh JS. Local plant species depletion in a tropical dry deciduous forest of northern India. Environ Conserv. 2004; 31: 55–62.



- Turner W, Spector S, Gardiner N, Fladeland M, Sterling E, Steininger M. Remote sensing for biodiversity science and conservation. Trends Ecol Evol. 2003; 18: 306–314.
- Chambers JQ, Asner GP, Morton DC, Anderson LO, Saatchi SS, Espírito-Santo FD, et al. Regional ecosystem structure and function: ecological insights from remote sensing of tropical forests. Trends Ecol Evol. 2007; 22: 414–423. PMID: 17493704
- 17. Puyravaud JP, Davidar P, Laurance WF. Cryptic loss of India's native forests. Science. 2010; 329: 32.
- Menon S, Bawa KS. Deforestation in the tropics: reconciling disparities in estimates for India. Ambio. 1998; 27: 576–577.
- Aziz SA, Laurance WF, Clements R. Forests reserved for rubber? Frontiers Ecol Environ. 2010; 8: 178.
- 20. Moffat AJ. Indicators of soil quality for UK forestry. Forestry. 2003; 76: 1–22.
- Satyam V, Jayakumar S. Impact of forest fire on physical, chemical and biological properties of soil: A review. Proc Int Acad Ecol Environ Sci. 2012; 2: 168–176.
- 22. Acton DF, Padbury GA. A conceptual framework for soil quality assessment and monitoring. In: Acton DF, editor. A program to assess and monitor soil quality in Canada: soil quality evaluation program summary (interim). Ottawa, Canada: Centre for Land and Biological Resources Research, No. 93–49, Agriculture Canada; 1993.
- 23. Bastida F, Zsolnay A, Hernández T, García C. Past, present and future of soil quality indices: a biological perspective. Geoderma. 2008; 147: 159–171.
- Tesfahunegn GB. Soil quality assessment strategies for evaluating soil degradation in northern Ethiopia. Appl Environ Soil Sci. 2014; 2014, Article ID 646502, 14 pages. <a href="http://dx.doi.org/10.1155/2014/646502">http://dx.doi.org/10.1155/2014/646502</a>
- Schoenholtz SH, Van Miegroet H, Burger JA. A review of chemical and physical properties as indicators
  of forest soil quality: challenges and opportunities. Forest Ecology and Management. 2000; 138: 335

  356.
- USDA NRCS. United States Department of Agriculture Natural Resources Conservation Service. Soil
  Quality Enhancement Activity—SQL07—Forest stand improvement for soil quality. 2010.
- Blecker SW, Stillings LL, Amacher MC, Ippolito JA, DeCrappeo NM. Development of vegetation based soil quality indices for mineralized terrane in arid and semi-arid regions, Ecol Indic. 2012; 20: 65–74.
- Doran JW, Parkin TB. Quantitative indicators of soil quality: a minimum data set. In: Doran JW, Jones
  AJ, editors. Methods for assessing soil quality. Madison: Soil Science Society of America, Inc.; 1996.
- 29. Lilburne L, Sparling G, Hewitt A, Campbell S, Stephens P. Sindi v3: improvements to a web-based tool for assessing soil quality. In: Stephens P, Callaghan J, Austin A, editors. Proceedings of Soil Quality and Sustainable Land Management Workshop; 2002 Apr 3–5; Palmerston North, New Zealand. Palmerson North: Landcare Research; 2002. p. 47–49.
- Council of Europe. Soil Protection. Recommendation to the Council of Ministers. Strasbourg: The Council; 1992.
- Royal Commission on Environmental Pollution. Nineteenth report–Sustainable Use of Soil. London: HMSO: 1996.
- **32.** Bhardwaj AK, Jasrotia P, Hamilton SK, Robertson GP. Ecological management of intensively cropped agro-ecosystems improves soil quality with sustained productivity. Agric Ecosyst Environ. 2011; 3: 419–429.
- **33.** Masto RE, Chhonkar PK, Singh D, Patra AK. Soil quality response to long-term nutrient and crop management on a semi-arid inceptisol. Agric Ecosyst Environ. 2007; 118: 30–142.
- Ramachandran A, Natarajan N, Nagamurugan N, Balaguru B, Soosairaj S, Britto SJ. Assessment of tree species diversity in semi-evergreen forests of Chitteri hills, Eastern Ghats of Tamil Nadu, South India. Indian Journal of Forestry. 2008; 31: 175–180.
- **35.** Balaguru B, Britto SJ, Nagamurugan N, Natarajan D, Soosairaj S. Identifying conservation priority zones for effective management of tropical forests in Eastern Ghats of India Volume 2. Forest Diversity and management. 2006; 469–483.
- **36.** Blake GR, Hartge KH. Bulk Density. In: Klute A, editor. Methods of soil analysis. Agronomy monograph 9, Part I. Madison, WI: America Society of Agronomy; 1986. pp. 363–375.
- Tandon HLS. Methods of analysis of soils plants water and fertilizers. New Delhi: Fertilizer Development and Consultation Organisation; 1995.
- **38.** Wang D, Anderson DW. Direct measurement of organic carbon content in soils by the Leco CR- 12 carbon analyzer. Commun Soil Sci Plant Anal. 1998; 29: 15–21.
- **39.** Toth SJ, Prince AL. Estimation of cation exchange capacity and exchangeable calcium, potassium, and sodium contents of soils by flame photometer techniques. Soil Sci. 1949; 67: 439–445.



- Anderson JM, Inagram JSI. Tropical Soil Biology and Fertility, A handbook of Methods. Wallingford, UK: CAB International; 1993.
- Pratt PF. Potassium. In: Page AL, editor. Methods of soil analysis. Part 2. Chemical and microbiological properties. Madison: Soil Science Society of America, Inc.; 1956: 1022–1030.
- 42. Jackson ML. Soil chemical analysis. New Delhi: Prentice Hall Inc.; 1973.
- **43.** Lindsay WL, Norvell WA. Development of a DTPA soil test for zinc, iron, manganese and copper. Soil Sci Soc Am J. 1978; 42: 421–428.
- Subbiah B, Asija GL. A rapid procedure for estimation of available nitrogen in soils. Curr Sci. 1956; 25:
   8.
- **45.** Olsen SR, Cole CV, Watanabe FS, Dean LA. Estimation of available phosphorus soil by extraction with sodium bicarbonate. Washington: USDA; 1954. 18 p.
- 46. Stanford S, English L. Use of flame photometer in rapid soil tests of K and Ca. Agron J. 1949; 41: 446–447.
- **47.** Vance ED, Brookes PC, Jenkinson DS. An extraction method for measuring soil microbial biomass. Soil Biol Biochem. 1987; 19: 703–707.
- **48.** Owen JS, Wang MK, Wang CH, King HB, Sun HL. Net N mineralization and nitrification rates in a forested ecosystem in north-eastern Taiwan. For Ecol Manage. 2003; 176: 519–530.
- Andrews SS, Flora CB, Mitchell JP, Karlen DL. Growers' perception and acceptance of soil quality indices. Geoderma. 2003; 114: 187–213.
- 50. IBM Carp. IBM SPSS statistics for windows, version 22.0. Armonk, NY: IBM Corp; 2013.
- Andrews SS, Karlen DL, Mitchel JP. A comparison of soil quality indexing methods for vegetable production systems in Northern California. Agric Ecosyst Environ. 2002; 90: 25–45.
- **52.** Andrews SS, Carroll CR. Designing a soil quality assessment tool for sustainable agroecosystem management. Ecol Appl. 2001; 11: 1573–1585.
- Brejda JJ, Moorman TB, Karlen DL, Dao TH. Identification of regional soil quality factors and indicators.
   I. Central and Southern High Plains. Soil Sci Soc Am J. 2000; 64: 2115–2124.
- 54. Tan SY, Li J. An exploratory spatial analysis of soil organic carbon distribution in Canadian Ecoregions. The International Archives of the Photogrammetry, Remote sensing and spatial information sciences, Volume XL-2, 2014. ISPRS Technical Commission II Symposium, 6–8 October 2014, Toronto. Canada.
- Wang S, Zhou C, Liu J, Tian H, Li K, Yang X. Carbon storage in northeast China as estimated from vegetation and soil inventories. Environ Pollut. 2002; 116: S157–S165. PMID: 11833904
- Ramachandran A, Jayakumar S, Haroon RM, Bhaskaran A, Arockiasamy DI. Carbon sequestration: estimation of carbon stock in natural forests using geospatial technology in the Eastern Ghats of Tamil Nadu, India. Curr Sci. 2007; 92: 323–331.
- Erdas 2011. Erdas Imagine, Hexgon geospatial, ink GA, USA.
- Liebig MA, Varvel G, Doran J. A simple performance based index for assessing multiple agroecosystem functions. Agron J. 2001; 93: 313–318.
- 59. Leiros MC, Trasar-Cepeda C, Seoane S, Gil-Sotres F. Biochemical properties of acid soils under climax vegetation (Atlantic oakwood) in an area of the European temperate-humid zone (Galicia, NW Spain): general parameters. Soil Biol Biochem. 2000; 32: 733–745.
- Chen CR, Xu ZH, Blumfield TJ, Hughes JM. Soil microbial biomass during the early establishment of hoop pine plantation: seasonal variation and impacts of site preparation. Forest Ecol Manage. 2003; 186: 213–225
- **61.** Gil-Sotres F, Trasar-Cepeda C, Leiros MC, Seoane S. Different approaches to evaluating soil quality using biochemical properties. Soil Biol Biochem. 2005; 37: 877–887.
- Miralles I, Ortega R, Sanchez-Maranon M, Leiros MC, Trasar-Cepeda C, Gil-Sotres F. Biochemical properties of range and forest soils in Mediterranean mountain environments. Biol Fertil Soils. 2007; 43: 721–729.
- **63.** Ndiaya EL, Sandeno JM, McGrath D, Dick RP. Integrative biological indicators for detecting change in soil quality Ameri J Alternat Agricult. 2000; 15: 26–36.
- 64. Ramaswami G, Sukumar R. Long-term environmental correlates of invasion by Lantana camara (Verbenaceae) in a seasonally dry tropical forest. PLoS One. 2013; 8: e76995. doi: 10.1371/journal.pone. 0076995 PMID: 24167555
- Raizada P, Sharma GP, Raghubanshi AS. Ingress of Lantana in dry tropical forest fragments: edge and shade effects. Curr Sci. 2006; 94:180–182.



- **66.** Joshi AA, Mudappa D, Shankar Raman TR. Invasive alien species in relation to edges and forest structure in tropical rainforest fragments of the Western Ghats. Trop Ecol. 2015; 56: 233–244.
- 67. Ramaswami G, Sukumar R. Lantana camara L. (Verbenaceae) invasion along streams in a heterogeneous landscape. J Biosci. 2014; 39: 717–726. PMID: <u>25116626</u>
- Goyal N, Pardha-Saradhi P, Sharma GP. Can adaptive modulation of traits to urban environments facilitate *Ricinus communis* L. invasiveness? Environ Monit Assess. 2014; 186: 7941–7948. doi: <a href="https://doi.org/10.1007/s10661-014-3978-0">10.1007/s10661-014-3978-0</a> PMID: 25103212
- **69.** Puyravaud JP, Davidar P, Laurance WF. Cryptic destruction of India's native forests. Conserv Lett. 2010; 3: 90–394.