
7 A particular piece of music was played by 91 pianists and for each pianist, the number of incorrect notes was recorded. The results are summarised in the table.

| Number of incorrect notes | 1 – 5 | 6 – 10 | 11 – 20 | 21 – 40 | 41 – 70 |
|---------------------------|-------|--------|---------|---------|---------|
| Frequency | 10 | 5 | 26 | 32 | 18 |

(a) Draw a histogram to represent this information.

[5]



| ` ′ | State which class interval contains the lower quartile and which class interval contains the upper quartile. | | | | | |
|-----|--|---|--|--|--|--|
| | Hence find the greatest possible value of the interquartile range. [2] |] | | | | |
| | | • | | | | |
| | | | | | | |
| | | • | | | | |
| | | • | | | | |
| | | • | | | | |
| | | • | | | | |
| | | • | | | | |
| | | • | | | | |
| | | • | | | | |
| | | | | | | |
| (c) | Calculate an estimate for the mean number of incorrect notes. [3 |] | | | | |
| (c) | Calculate an estimate for the mean number of incorrect notes. [3 |] | | | | |
| (c) | |] | | | | |
| (c) | |] | | | | |
| (c) | | | | | | |
| (c) | | | | | | |
| (c) | | | | | | |
| (c) | | | | | | |
| (c) | | | | | | |