

# Numerical Optimization

## Lecture Notes #3

### Convergence; Line Search Methods

Fall 2024

# Outline

## 1 Introduction

- Recap
- Fundamentals: Rate of Convergence

## 2 Line Search Methods

- Search Direction: Steepest Descent, Newton, or Other?!?
- Step Length Selection — 1D Minimization
- Step Length Selection — The Wolfe Conditions

## Quick Recap: Last Time

Some fundamental building blocks of unconstrained optimization:

### Theorem (Taylor)

For some  $t \in (0, 1)$ , we have

$$f(\bar{\mathbf{x}} + \bar{\mathbf{p}}) = f(\bar{\mathbf{x}}) + \bar{\mathbf{p}}^T \underbrace{\nabla f(\bar{\mathbf{x}})}_{\text{gradient}} + \frac{1}{2} \bar{\mathbf{p}}^T \underbrace{\left[ \nabla^2 f(\bar{\mathbf{x}} + t\bar{\mathbf{p}}) \right]}_{\text{Hessian}} \bar{\mathbf{p}}.$$

**4 theorems** relating  $f(\bar{\mathbf{x}})$  and its derivatives to optimal solutions.

- [1]  $\bar{\mathbf{x}}^*$  optimal  $\Rightarrow \nabla f(\bar{\mathbf{x}}^*) = 0$ .
- [2]  $\bar{\mathbf{x}}^*$  optimal  $\Rightarrow \nabla f(\bar{\mathbf{x}}^*) = 0$ , and  $\nabla^2 f(\bar{\mathbf{x}}^*)$  positive semi-definite.
- [3]  $\nabla f(\bar{\mathbf{x}}^*) = 0$ , and  $\nabla^2 f(\bar{\mathbf{x}}^*)$  positive definite  $\Rightarrow \bar{\mathbf{x}}^*$  optimal.
- [4a]  $f$  convex, and  $\bar{\mathbf{x}}^*$  local optimum  $\Rightarrow \bar{\mathbf{x}}^*$  global optimum.
- [4b]  $f$  convex, and  $\nabla f(\bar{\mathbf{x}}^*) = 0 \Rightarrow \bar{\mathbf{x}}^*$  global optimum.

**Note:** The complete statement of the theorems require sufficient smoothness (existence) of derivatives of  $f$ .

## Key Concepts

## Rate of Convergence

## Definition (Rate of Convergence, Sequences)

Suppose the sequence  $\beta = \{\beta_n\}_{n=1}^{\infty}$  converges to zero, and  $\bar{\mathbf{x}} = \{\bar{\mathbf{x}}_n\}_{n=1}^{\infty}$  converges to a point  $\bar{\mathbf{x}}^*$ .

If  $\exists K > 0$ :  $\|\bar{\mathbf{x}}_n - \bar{\mathbf{x}}^*\| < K\beta_n$ , for  $n > N$  (i.e. for  $n$  large enough), then we say that  $\{\bar{\mathbf{x}}_n\}_{n=1}^{\infty}$  converges to  $\bar{\mathbf{x}}^*$  with a **Rate of Convergence**  $\mathcal{O}(\beta_n)$  ("Big Oh of  $\beta_n$ ").

We write

$$\bar{\mathbf{x}}_n = \bar{\mathbf{x}}^* + \mathcal{O}(\beta_n).$$

**Note:** The sequence  $\beta = \{\beta_n\}_{n=1}^{\infty}$  is usually chosen to be e.g.

$$\beta_n = n^{-p}, \quad \text{for some value of } p.$$

## Rates of Convergence: Example

What does the sequence  $1 + (0.5)^k$  converge to? What is the convergence rate?

Let  $\bar{\mathbf{x}} = \{\bar{\mathbf{x}}_n\}_{n=1}^{\infty}$  be a sequence converging to  $\bar{\mathbf{x}}^*$ , the convergence rate is said to be

**Q-linear** (quotient-linear) if  $\exists r \in (0, 1)$  and

$$\frac{\|\bar{\mathbf{x}}_{k+1} - \bar{\mathbf{x}}^*\|}{\|\bar{\mathbf{x}}_k - \bar{\mathbf{x}}^*\|} \leq r, \quad \text{for } k \text{ sufficiently large}$$

## Rates of Convergence

Let  $\bar{\mathbf{x}} = \{\bar{\mathbf{x}}_n\}_{n=1}^{\infty}$  be a sequence converging to  $\bar{\mathbf{x}}^*$ , the convergence rate is said to be

**Q-linear** (quotient-linear) if  $\exists r \in (0, 1)$  and

$$\frac{\|\bar{\mathbf{x}}_{k+1} - \bar{\mathbf{x}}^*\|}{\|\bar{\mathbf{x}}_k - \bar{\mathbf{x}}^*\|} \leq r, \quad \text{for } k \text{ sufficiently large}$$

**Q-superlinear** if

$$\lim_{k \rightarrow \infty} \frac{\|\bar{\mathbf{x}}_{k+1} - \bar{\mathbf{x}}^*\|}{\|\bar{\mathbf{x}}_k - \bar{\mathbf{x}}^*\|} = 0.$$

**Q-quadratic** if  $\exists M \in \mathbb{R}^+$  and

$$\frac{\|\bar{\mathbf{x}}_{k+1} - \bar{\mathbf{x}}^*\|}{\|\bar{\mathbf{x}}_k - \bar{\mathbf{x}}^*\|^2} \leq M, \quad \text{for } k \text{ sufficiently large}$$

## Line Search Methods

Consider that we want to solve  $\min_{\bar{\mathbf{x}} \in \mathbb{R}^n} f(\bar{\mathbf{x}})$ .

We can reduce this n-dimensional problem to a one-dimensional problem that can be solved iteratively via a line search method. Key steps for a line search method: (i) pick a **search direction**  $\bar{\mathbf{p}}_k$  and, then (ii) solve the one-dimensional problem

$$\min_{\alpha_k > 0} f(\bar{\mathbf{x}}_k + \alpha_k \bar{\mathbf{p}}_k).$$

The solution gives us an optimal value for  $\alpha_k$ , so the next point is given by

$$\bar{\mathbf{x}}_{k+1} = \bar{\mathbf{x}}_k + \alpha_k \bar{\mathbf{p}}_k,$$

where  $\alpha_k$  is known as the **step length**. In order for a line search method to work well, we need good choices of the direction  $\bar{\mathbf{p}}_k$  and the step length  $\alpha_k$ .

How we choose  $\bar{\mathbf{p}}_k$ 

Line Search

We can choose  $\bar{\mathbf{p}}_k$  for a line search method by using

- the **Steepest Descent Method** or
- the **Newton Method**.

In the next couple of slides, we will derive the Steepest Descent Method and the Newton method by using Taylor expansion.

Note that once we obtain  $\bar{\mathbf{p}}_k$  we can solve the one-dimensional problem

$$\min_{\alpha_k > 0} f(\bar{\mathbf{x}}_k + \alpha_k \bar{\mathbf{p}}_k).$$

to obtain the optimal  $\alpha_k$ . Given a starting point  $\bar{\mathbf{x}}_0$ , the subsequent points can be computed iterative:

$$\bar{\mathbf{x}}_{k+1} = \bar{\mathbf{x}}_k + \alpha_k \bar{\mathbf{p}}_k.$$



## Steepest Descent Direction

Line Search

The intuitive choice for  $\bar{\mathbf{p}}_k$  is to move in the direction of steepest descent, *i.e.* in the negative gradient direction.

Going back to the Taylor expansion

$$f(\bar{\mathbf{x}} + \alpha \bar{\mathbf{p}}) = f(\bar{\mathbf{x}}) + \alpha \bar{\mathbf{p}}^T \nabla f(\bar{\mathbf{x}}),$$

we immediately see that the direction of most rapid decrease gives


$$\min_{\|\bar{\mathbf{p}}\|=1} \bar{\mathbf{p}}^T \nabla f(\bar{\mathbf{x}}) = \min_{\theta \in [0, 2\pi]} \cos \theta \|\nabla f(\bar{\mathbf{x}})\| = -\|\nabla f(\bar{\mathbf{x}})\|,$$

which is achieved when  $\theta = \pi \Leftrightarrow \bar{\mathbf{p}} = -\nabla f(\bar{\mathbf{x}}) / \|\nabla f(\bar{\mathbf{x}})\|$ .


**Recall:**  $\bar{\mathbf{v}}^T \bar{\mathbf{w}} = \cos \theta \|\bar{\mathbf{v}}\| \cdot \|\bar{\mathbf{w}}\|$ , where  $\theta$  is the angle between the vectors  $\bar{\mathbf{v}}$  and  $\bar{\mathbf{w}}$ .

## Steepest Descent Direction

Line Search



**Figure:** The steepest descent direction  $\bar{p}_k$  is perpendicular to the contour lines of the objective.



**Figure:**  $\bar{v}^T \bar{w} = \cos \theta \|\bar{v}\| \cdot \|\bar{w}\|$ .

## Newton Direction

If  $f$  is smooth enough and the Hessian is positive definite, we can select  $\bar{\mathbf{p}}_k$  to be the “Newton direction.” We write down the second order Taylor expansion:

$$f(\bar{\mathbf{x}} + \bar{\mathbf{p}}) \approx f(\bar{\mathbf{x}}) + \bar{\mathbf{p}}^T \nabla f(\bar{\mathbf{x}}) + \frac{1}{2} \bar{\mathbf{p}}^T \left[ \nabla^2 f(\bar{\mathbf{x}}) \right] \bar{\mathbf{p}}.$$

We seek the minimum of the right-hand-side by computing the derivative with respect to  $\bar{\mathbf{p}}$  and set the result to zero

$$\nabla f(\bar{\mathbf{x}}) + \left[ \nabla^2 f(\bar{\mathbf{x}}) \right] \bar{\mathbf{p}} = 0,$$

which gives the Newton direction

$$\bar{\mathbf{p}}^N = - \left[ \nabla^2 f(\bar{\mathbf{x}}) \right]^{-1} \nabla f(\bar{\mathbf{x}}).$$

## Newton Direction

Line Search

Recall Taylor expansion:  $f(\bar{\mathbf{x}} + \alpha \bar{\mathbf{p}}) = f(\bar{\mathbf{x}}) + \alpha \bar{\mathbf{p}}^T \nabla f(\bar{\mathbf{x}}),$

As long as the Hessian is positive definite,  $\bar{\mathbf{p}}^N$  is a descent-direction:

$$\bar{\mathbf{p}}^N \nabla f(\bar{\mathbf{x}}) = - \underbrace{\nabla f(\bar{\mathbf{x}})^T \left[ \nabla^2 f(\bar{\mathbf{x}}) \right]^{-1}}_{\text{Pos. Def.}} \nabla f(\bar{\mathbf{x}}) < 0$$

**Note:** Clearly, the Newton direction is more “expensive” than the steepest descent direction — we must compute the Hessian matrix  $\nabla^2 f(\bar{\mathbf{x}})$ , and invert it (*i.e.* solve an  $n \times n$  linear system).

**Note:** The convergence rate for steepest descent methods is **linear** and for Newton methods it is **quadratic**, hence there is a lot to gain by finding the Newton direction.

Example: NW<sup>1st</sup>-2.2, p 30.

1 of 3

**Problem:** Show that the function  $f(x) = 8x + 12y + x^2 - 2y^2$  has only one stationary point, and that it is neither a maximum nor a minimum, but a saddle point. Sketch the contours for  $f$ .


**Solution:** The gradient of  $f$  is

$$\nabla f = \begin{bmatrix} 8 + 2x \\ 12 - 4y \end{bmatrix}$$


which has the stationary point  $(x, y) = (-4, 3)$ . Since the Hessian

$$\nabla^2 f = \begin{bmatrix} 2 & 0 \\ 0 & -4 \end{bmatrix}$$

has both positive and negative eigenvalues, the stationary point must be a saddle point.



**Figure:** The contour lines for  $f(x)$ .



**Figure:** The function  $f(x)$  around the stationary point.

Example: NW<sup>1st</sup>-2.2, p 30.

2 of 3

If we start an iteration in  $(x_0, y_0) = (0, 0)$ :

The steepest descent direction is


$$\bar{\mathbf{p}}_0^{\text{SD}} = -\nabla f = - \begin{bmatrix} 8 + 2x \\ 12 - 4y \end{bmatrix} = - \begin{bmatrix} 8 \\ 12 \end{bmatrix}$$

and the Newton direction is

$$\bar{\mathbf{p}}_0^N = -[\nabla^2 f]^{-1} \nabla f = - \begin{bmatrix} 2 & 0 \\ 0 & -4 \end{bmatrix}^{-1} \begin{bmatrix} 8 \\ 12 \end{bmatrix} = \begin{bmatrix} -4 \\ 3 \end{bmatrix}$$


Example: NW<sup>1st</sup>-2.2, p 30.

3 of 3



**Figure:** The **Newton** and **Steepest Descent** directions starting in  $(0, 0)$ . Note that the Newton method is heading to the saddle point, but the Steepest descent method will, in general, not converge to a non-minimum stationary point.

## Modified (Convexified) Example



**Figure:** Convexification of the silly book problem. Same point of interest,  $\nabla f = [8 + 2x, 12 - 4y + 2/3y^3]^T$ ,  $\nabla^2 f = \begin{bmatrix} 2 & 0 \\ 0 & -4 + 2y^2 \end{bmatrix}$ . Now, both the steepest descent and Newton directions are descent directions.



## Line Search Methods — Directions

Method	Search Direction	Convergence
<b>Steepest Descent</b>	$p_k = -\nabla f(\bar{\mathbf{x}}_k) / \ \nabla f(\bar{\mathbf{x}}_k)\ $	Linear
<b>Quasi-Newton</b>	$p_k = -H_k^{-1} \nabla f(\bar{\mathbf{x}}_k)$	Super-Linear
<b>Newton</b>	$p_k = -[\nabla^2 f(\bar{\mathbf{x}}_k)]^{-1} \nabla f(\bar{\mathbf{x}}_k)$	Quadratic

**Table:** Summary of search directions for different schemes. In **Quasi-Newton** schemes we do not explicitly compute the Hessian  $\nabla^2 f(\bar{\mathbf{x}}_k)$  in each iteration, instead we use an approximation  $H_k \approx \nabla^2 f(\bar{\mathbf{x}}_k)$  which is updated in some clever way [TO BE EXPLORED IN GREAT DETAIL LATER] (lecture 18  $\rightarrow$  ...).

We will return to the selection of  $\bar{\mathbf{p}}_k$ , but let's consider the computation of the step length  $\alpha_k$ ...

## Line Search Methods: Step Length Selection

Given a descent direction  $\bar{\mathbf{p}}_k$  we would like to find the global minimizer  $\alpha_k^*$  of


$$\min_{\alpha_k > 0} f(\bar{\mathbf{x}}_k + \alpha_k \bar{\mathbf{p}}_k).$$

As this is just one of possible many steps in the iteration, it is not wise to expend too much time in finding  $\alpha_k$ . We are faced with a trade-off:

- We want an  $\alpha_k$  so that we get a **substantial reduction** in the objective  $f$ .
- We want to find  $\alpha_k$  **fast**.

In practice we perform an **inexact line search** — settling for an  $\alpha_k$  which gives **adequate reduction** in the objective.

# What is “adequate reduction?”



**Figure:** Consider the objective  $f(x) = \sqrt{x^2 + 10^{-8}}$ , if we let  $x_k = \{1, -0.8, 0.7, -0.65, 0.625, -0.6125, 0.60625, \dots\}$ , then the descent directions are given by  $p_k = \{-1, 1, -1, 1, -1, 1, -1, \dots\}$ , so this generates a decreasing sequence  $f(x_k + \alpha_k p_k) < f(x_k)$ . However, with the current choice of  $\alpha_k = \{1.8, -1.5, 1.35, -1.275, 1.2375, -1.21875, \dots\}$  the convergence rate is less than spectacular. Note that y-axis represents  $f(x)$  and x-axis represents  $x$  on both graphs.

Clearly, we need a stronger condition than  $f(x_k + \alpha_k p_k) < f(x_k)$ .

## The Wolfe Conditions

1 of 2

There are many ways to enforce reduction in the objective, e.g.

## Armijo Condition

(Wolfe Condition #1)

The **Armijo Condition**

$$f(\bar{\mathbf{x}}_k + \alpha \bar{\mathbf{p}}_k) \leq f(\bar{\mathbf{x}}_k) + c_1 \alpha \bar{\mathbf{p}}_k^T \nabla f(\bar{\mathbf{x}}), \quad c_1 \in (0, 1),$$

requires the reduction to be proportional to the step length  $\alpha$ , as well as the directional derivative  $\bar{\mathbf{p}}_k^T \nabla f(\bar{\mathbf{x}})$ . **In practice**  $c_1$  is usually set to be quite small, e.g.  $\sim 10^{-4}$ .

Armijo Condition requires that the step length cause a sufficient decrease in the objective function value.

## The Wolfe Conditions

2 of 2

To rule out unacceptably short steps, we additionally enforce

Curvature Condition

(Wolfe Condition #2)

The **Curvature Condition**

$$\bar{\mathbf{p}}_k^T \nabla f(\bar{\mathbf{x}}_k + \alpha \bar{\mathbf{p}}_k) \geq c_2 \bar{\mathbf{p}}_k^T \nabla f(\bar{\mathbf{x}}_k), \quad c_2 \in (c_1, 1).$$

It prevents us from stopping when more progress can be made by moving further (increasing  $\alpha$ ).


Together these two conditions are known as the **Wolfe conditions**.

## The Wolfe Conditions: Part I — The Armijo Condition

### The **Armijo Condition**

$$f(\bar{\mathbf{x}}_k + \alpha \bar{\mathbf{p}}_k) \leq f(\bar{\mathbf{x}}_k) + c_1 \alpha \bar{\mathbf{p}}_k^T \nabla f(\bar{\mathbf{x}}), \quad c_1 \in (0, 1)$$

requires the reduction to be proportional to the step length  $\alpha$ , as well as the directional derivative. **In practice**  $c_1$  is usually set to be quite small, e.g.  $\sim 10^{-4}$ .



Here  $y$ -axis represents  $f(x)$  and  $x$ -axis represents  $\alpha$ .


## The Wolfe Conditions: Part II — The Curvature Condition

To rule out unacceptable short steps, the **curvature condition**

$$\bar{\mathbf{p}}_k^T \nabla f(\bar{\mathbf{x}}_k + \alpha \bar{\mathbf{p}}_k) \geq c_2 \bar{\mathbf{p}}_k^T \nabla f(\bar{\mathbf{x}}_k), \quad c_2 \in (c_1, 1)$$

— it prevents us from stopping when more progress can be made by moving further (increasing  $\alpha$ ). Typical values:  $c_2^{\text{N,QN}} = 0.9$ ,  $c_2^{\text{CG}} = 0.1$ .

— The curvature condition requires that the directional derivative at the next iterate be shallower.




Here y-axis represents  $f(x)$  and x-axis represents  $\alpha$ .

## The Wolfe Conditions: Part I+II — Acceptable Step

1/2

Together, the Armijo and Curvature conditions constitute the Wolfe Conditions.






## The Wolfe Conditions: Part I+II — Acceptable Step

2/2

Together, the Armijo and Curvature conditions constitute the Wolfe Conditions.



## The Strong Wolfe Conditions

A step length  $\alpha$  may satisfy the **Wolfe Conditions**

$$\begin{aligned} f(\bar{\mathbf{x}}_k + \alpha \bar{\mathbf{p}}_k) &\leq f(\bar{\mathbf{x}}_k) + c_1 \alpha \bar{\mathbf{p}}_k^T \nabla f(\bar{\mathbf{x}}), & c_1 &\in (0, 1) \\ \bar{\mathbf{p}}_k^T \nabla f(\bar{\mathbf{x}}_k + \alpha \bar{\mathbf{p}}_k) &\geq c_2 \bar{\mathbf{p}}_k^T \nabla f(\bar{\mathbf{x}}_k), & c_2 &\in (c_1, 1) \end{aligned}$$

even though it is far from a minimizer of  $f(\bar{\mathbf{x}}_k + \alpha \bar{\mathbf{p}}_k)$ , the **Strong Wolfe Conditions**

$$\begin{aligned} f(\bar{\mathbf{x}}_k + \alpha \bar{\mathbf{p}}_k) &\leq f(\bar{\mathbf{x}}_k) + c_1 \alpha \bar{\mathbf{p}}_k^T \nabla f(\bar{\mathbf{x}}), & c_1 &\in (0, 1) \\ |\bar{\mathbf{p}}_k^T \nabla f(\bar{\mathbf{x}}_k + \alpha \bar{\mathbf{p}}_k)| &\leq c_2 |\bar{\mathbf{p}}_k^T \nabla f(\bar{\mathbf{x}}_k)|, & c_2 &\in (c_1, 1) \end{aligned}$$

further disallows values of

$$\left[ \bar{\mathbf{p}}_k^T \nabla f(\bar{\mathbf{x}}_k + \alpha \bar{\mathbf{p}}_k) \right]$$

which are “too positive,” thus excluding point that are far from the stationary points of  $\bar{\mathbf{p}}_k^T \nabla f(\bar{\mathbf{x}}_k + \alpha \bar{\mathbf{p}}_k)$ .

## Are the Wolfe Conditions too Restrictive?

It can be shown (see NW<sup>2nd</sup> pp.35–36) that there **exist** step lengths  $\alpha$  which satisfy the Wolfe Conditions (and the Strong Wolfe Conditions) **for every** function  $f$  which is smooth and bounded below.

Formally —

### Theorem (Existence of Acceptable $\alpha$ )

*Suppose  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  is continuously differentiable. Let  $\bar{\mathbf{p}}_k$  be a descent direction at  $\bar{\mathbf{x}}_k$ , and assume that  $f$  is bounded below along the line  $\{\bar{\mathbf{x}}_k + \alpha\bar{\mathbf{p}}_k : \alpha > 0\}$ . Then if  $0 < c_1 < c_2 < 1$ , there exist intervals of step lengths satisfying the Wolfe conditions and the strong Wolfe conditions.*

See also “Goldstein Conditions” (NW<sup>2nd</sup> p.36.)

## Algorithm: Backtracking Line Search

### Algorithm: Backtracking Line Search

```
[0] Find a descent direction  $\bar{\mathbf{p}}_k$ 
[1] Set  $\bar{\alpha} > 0$ ,  $\rho \in (0,1)$ ,  $c \in (0,1)$ , set  $\alpha = \bar{\alpha}$ 
[2] While  $f(\bar{\mathbf{x}}_k + \alpha \bar{\mathbf{p}}_k) > f(\bar{\mathbf{x}}_k) + c\alpha \bar{\mathbf{p}}_k^T \nabla f(\bar{\mathbf{x}}_k)$ 
[3]    $\alpha = \rho\alpha$ 
[4] End-While
[5] Set  $\alpha_k = \alpha$ 
```

If an algorithm selects the step lengths appropriately (e.g. backtracking), we do not have to check the second inequality of the Wolfe conditions.

The algorithm above is especially well suited for use with Newton method ( $\bar{\mathbf{p}}_k = \bar{\mathbf{p}}_k^N$ ), where  $\bar{\alpha} = 1$ . It is less successful for quasi-Newton and CG-based approaches.

The value of the **contraction factor**  $\rho$  can be allowed to vary at each iteration of the line search. (To be revisited)

# Index

- Armijo condition, 20
- backtracking linesearch, 29
- convergence
  - rate of, 4
    - linear, 6
    - quadratic, 6
    - superlinear, 6
- curvature condition, 21
- Newton direction, 11
- steepest descent direction, 9
- Wolfe conditions, 21
  - strong, 27