


Pediatric Febrile Infant Clinical Pathway (8-60 days old)

29-60 days

HSV Uncommon In This Age Group

HSV most likely to present within first 3 weeks of age, but may occur in first 4-6 weeks.

See AAP Red Book for further info.



Key Evidence Summary:

In this age group, febrile infants have a 0.12-0.32% risk of bacterial meningitis, a 1.1-2.2% risk of bacteremia, and a 5-10% risk of UTI.

If all IMs neg, UA neg, CSF (if obtained) neg, risk of bacteremia is ~0.1%.

If all IMs are normal, an additional 4000 successful LPs would be required to diagnose 1 additional case of bacterial meningitis.

from CPG. Numbers map to KAS in CPG. Colored circles map to REVISE II Project measures.

s include temp, PCT, CRP, ANC. IMs considered (+) if any elevated: temperature ($\geq 38.5^{\circ}\text{C}$), PCT ≥ 0.5 ng/mL, CRP ≥ 20 mg/L, or ANC > 4000 -5200 per mm^3 .

nd CSF for cell count, gram stain, glucose, protein, CSF culture, and CSF pathogen panel.

Observation at home reasonable if reliable phone and transportation, parent willingness to observe and communicate changes in condition, and agreement that infant be evaluated in 24 hours. Most infants 29-60 days old with negative IMs and UA can safely be observed at home, though hospital observation is an option when there are risks to follow-up.