1 **(a)** 24 **[B1]**

(b) Identifies $\angle OAP = 90°$ (tangent perpendicular to radius) **[M1]**

$OP^2 = 7^2 + 24^2 = 625 \implies OP = 25$ **[A1]**

(c) Let M be the intersection of OP and AB. Identifies $\triangle OAM \sim \triangle OPA$ OR uses Area of $\triangle OAP$ OR uses $\cos(\angle AOP)$ **[M1]**

$\cos(\angle AOP) = \dfrac{7}{25}$ and $OM = 7 \times \cos(\angle AOP)$ **[M1]**

$OM = \dfrac{49}{25}$ or 1.96 **[A1]**

[Total Marks: 6]

2 **(a)** Tangents from an external point are equal in length **[B1]**

(b) 70 **[B1]**

(c) Angle $OAP = 90°$ OR Angle $OBP = 90°$ **[M1]**

140 **[A1]**

[Total Marks: 4]

3 **(a)** $x + 3 = 3x - 5$ **[M1]**

$x = 4$ **[A1]**

(b) Perpendicular distance = 7 (can be implied) **[B1]**

$\sqrt{25^2 - 7^2}$ OR 24 **[M1]**

48 **[A1]**

[Total Marks: 5]

4 **(a)** States or uses $OM \perp AB$ (perpendicular bisector of chord passes through centre), so $AM = 7$ **[B1]**

Uses Pythagoras' Theorem: $OM = \sqrt{25^2 - 7^2} = \sqrt{625 - 49} = \sqrt{576} = 24$ **[B1]**

(b) States $ON = OM = 24$ (equal chords are equidistant from centre) **[B1]**

Uses Cosine rule: $MN^2 = 24^2 + 24^2 - 2(24)(24)\cos 100°$ OR uses $MN = 2 \times 24 \times \sin 50°$ **[M1]**

36.8 (accept $36.77\ldots$) **[A1]**

(c) $\dfrac{1}{2} \times 24 \times 24 \times \sin 100°$ **[M1]**

284 (accept $283.6\ldots$) **[A1]**

[Total Marks: 7]

5 **(a)** Identifies $\angle OAT = 90°$ and $\angle AOT = 90° - x$ **[M1]**

Uses $AM = r\sin(90° - x) = r\cos x$ and perpendicular bisector property $AB = 2AM$ to get $2r\cos x$ **[A1]**

(b) $r\sin x$ (equal chords are equidistant from centre, distance $= OM = r\cos(90° - x)$) **[B1]**

(c) Area of $OAB = \frac{1}{2} \times AB \times OM = \frac{1}{2}(2r\cos x)(r\sin x) = r^2 \sin x \cos x$ **[M1]**

$TM = \frac{AM}{\tan x} = \frac{r\cos x}{\tan x} = \frac{r\cos^2 x}{\sin x}$ **[M1]**

Area of $ATB = \frac{1}{2} \times AB \times TM = \frac{1}{2}(2r\cos x)\left(\frac{r\cos^2 x}{\sin x}\right) = \frac{r^2\cos^3 x}{\sin x}$ **[M1]**

Equates areas: $r^2 \sin x \cos x = \frac{r^2\cos^3 x}{\sin x} \implies \sin^2 x \cos x = \cos^3 x \implies \tan^2 x = 1$ **[M1]**

45 **[A1]**

[Total Marks: 8]

6 (a) The perpendicular bisector of a chord passes through the centre of the circle **[B1]**

(b) $MX = 16 \div 2 = 8$ **[B1]**

$OX = \sqrt{10^2 - 8^2} = 6$ **[M1]**

$10 - 6 = 4$ **[A1]**

(c) Distance of PQ from $O = 11 - 6 = 5$ **[B1]**

Half length of $PQ = \sqrt{10^2 - 5^2}$ **[M1]**

$\sqrt{75}$ or 8.66... **[A1]**

$2 \times \sqrt{75} = 17.3$ (17.32...) **[A1]**

[Total Marks: 8]