IGCSE 0607 Practice Questions — Combined Questions & Markscheme

Total Questions: 1301 — **Total Marks:** 7771 Marks

Table of Contents & Quick Navigation

- **E1.1** Types of number
  → **Paper 2 (Non-Calculator)** *(12 Qs)* · **Paper 4 (Calculator)** *(6 Qs)*
- **E1.2** Sets
  → **Paper 2 (Non-Calculator)** *(7 Qs)* · **Paper 4 (Calculator)** *(8 Qs)*
- **E1.3** Powers and roots
  → **Paper 2 (Non-Calculator)** *(10 Qs)* · **Paper 4 (Calculator)** *(5 Qs)*
- **E1.4** Fractions, decimals and percentages
  → **Paper 2 (Non-Calculator)** *(10 Qs)* · **Paper 4 (Calculator)** *(5 Qs)*
- **E1.5** Ordering
  → **Paper 2 (Non-Calculator)** *(9 Qs)* · **Paper 4 (Calculator)** *(6 Qs)*
- **E1.6** The four operations
  → **Paper 2 (Non-Calculator)** *(10 Qs)* · **Paper 4 (Calculator)** *(5 Qs)*
- **E1.7** Indices I
  → **Paper 2 (Non-Calculator)** *(11 Qs)* · **Paper 4 (Calculator)** *(1 Q)*
- **E1.8** Standard form
  → **Paper 2 (Non-Calculator)** *(9 Qs)* · **Paper 4 (Calculator)** *(6 Qs)*
- **E1.9** Estimation
  → **Paper 2 (Non-Calculator)** *(13 Qs)* · **Paper 4 (Calculator)** *(5 Qs)*
- **E1.10** Ratio and proportion
  → **Paper 2 (Non-Calculator)** *(7 Qs)* · **Paper 4 (Calculator)** *(8 Qs)*
- **E1.11** Rates
  → **Paper 2 (Non-Calculator)** *(6 Qs)* · **Paper 4 (Calculator)** *(12 Qs)*
- **E1.12** Percentages
  → **Paper 2 (Non-Calculator)** *(6 Qs)* · **Paper 4 (Calculator)** *(12 Qs)*
- **E1.13** Using a calculator
  → **Paper 4 (Calculator)** *(18 Qs)*
- **E1.14** Time
  → **Paper 2 (Non-Calculator)** *(6 Qs)* · **Paper 4 (Calculator)** *(12 Qs)*
- **E1.15** Money
  → **Paper 2 (Non-Calculator)** *(5 Qs)* · **Paper 4 (Calculator)** *(13 Qs)*

- **E1.16** Exponential growth and decay
  → **Paper 2 (Non-Calculator)** *(2 Qs)* · **Paper 4 (Calculator)** *(16 Qs)*

- **E1.17** Surds
  → **Paper 2 (Non-Calculator)** *(36 Qs)*

- **E2.1** Introduction to algebra
  → **Paper 2 (Non-Calculator)** *(16 Qs)* · **Paper 4 (Calculator)** *(11 Qs)*

- **E2.2** Algebraic manipulation
  → **Paper 2 (Non-Calculator)** *(21 Qs)* · **Paper 4 (Calculator)** *(6 Qs)*

- **E2.3** Algebraic fractions
  → **Paper 2 (Non-Calculator)** *(16 Qs)* · **Paper 4 (Calculator)** *(11 Qs)*

- **E2.4** Indices II
  → **Paper 2 (Non-Calculator)** *(22 Qs)* · **Paper 4 (Calculator)** *(5 Qs)*

- **E2.5** Equations
  → **Paper 2 (Non-Calculator)** *(26 Qs)* · **Paper 4 (Calculator)** *(19 Qs)*

- **E2.6** Inequalities
  → **Paper 2 (Non-Calculator)** *(27 Qs)* · **Paper 4 (Calculator)** *(18 Qs)*

- **E2.7** Sequences
  → **Paper 2 (Non-Calculator)** *(10 Qs)* · **Paper 4 (Calculator)** *(14 Qs)*

- **E2.8** Proportion
  → **Paper 2 (Non-Calculator)** *(10 Qs)* · **Paper 4 (Calculator)** *(13 Qs)*

- **E3.1** Graphs of functions
  → **Paper 2 (Non-Calculator)** *(20 Qs)* · **Paper 4 (Calculator)** *(19 Qs)*

- **E3.2** Sketching graphs on a calculator
  → **Paper 4 (Calculator)** *(15 Qs)*

- **E3.3** Functions
  → **Paper 2 (Non-Calculator)** *(20 Qs)* · **Paper 4 (Calculator)** *(16 Qs)*

- **E3.4** Finding a quadratic function using given information
  → **Paper 2 (Non-Calculator)** *(19 Qs)* · **Paper 4 (Calculator)** *(17 Qs)*

- **E3.5** Asymptotes
  → **Paper 2 (Non-Calculator)** *(7 Qs)* · **Paper 4 (Calculator)** *(8 Qs)*

- **E3.6** Transforming graphs of functions
  → **Paper 2 (Non-Calculator)** *(8 Qs)* · **Paper 4 (Calculator)** *(4 Qs)*

- **E3.7** The logarithmic function
  → **Paper 2 (Non-Calculator)** *(12 Qs)* · **Paper 4 (Calculator)** *(24 Qs)*

- **E4.1** Coordinates
  → **Paper 2 (Non-Calculator)** *(8 Qs)* · **Paper 4 (Calculator)** *(4 Qs)*

- **E4.2** Gradient of linear graphs
  → **Paper 2 (Non-Calculator)** *(10 Qs)* · **Paper 4 (Calculator)** *(5 Qs)*

- **E4.3** Length and midpoint
  → **Paper 2 (Non-Calculator)** *(8 Qs)* · **Paper 4 (Calculator)** *(7 Qs)*

- **E4.4** Equations of linear graphs
  → **Paper 2 (Non-Calculator)** *(10 Qs)* · **Paper 4 (Calculator)** *(5 Qs)*

- **E4.5** Parallel lines
  → **Paper 2 (Non-Calculator)** *(8 Qs)* · **Paper 4 (Calculator)** *(7 Qs)*

- **E4.6** Perpendicular lines
  → **Paper 2 (Non-Calculator)** *(10 Qs)* · **Paper 4 (Calculator)** *(5 Qs)*

- **E5.1** Geometrical terms
  → **Paper 2 (Non-Calculator)** *(13 Qs)* · **Paper 4 (Calculator)** *(8 Qs)*

- **E5.2** Angle measurement in degrees
  → **Paper 2 (Non-Calculator)** *(10 Qs)* · **Paper 4 (Calculator)** *(11 Qs)*

- **E5.3** Similarity
  → **Paper 2 (Non-Calculator)** *(13 Qs)* · **Paper 4 (Calculator)** *(20 Qs)*

- **E5.4** Symmetry
  → **Paper 2 (Non-Calculator)** *(8 Qs)* · **Paper 4 (Calculator)** *(7 Qs)*

- **E5.5** Angles
  → **Paper 2 (Non-Calculator)** *(7 Qs)* · **Paper 4 (Calculator)** *(8 Qs)*

- **E5.6** Circle theorems I
  → **Paper 2 (Non-Calculator)** *(20 Qs)* · **Paper 4 (Calculator)** *(16 Qs)*

- **E5.7** Circle theorems II
  → **Paper 2 (Non-Calculator)** *(6 Qs)* · **Paper 4 (Calculator)** *(9 Qs)*

- **E6.1** Units of measure
  → **Paper 2 (Non-Calculator)** *(4 Qs)* · **Paper 4 (Calculator)** *(8 Qs)*

- **E6.2** Area and perimeter
  → **Paper 2 (Non-Calculator)** *(4 Qs)* · **Paper 4 (Calculator)** *(8 Qs)*

- **E6.3** Circles, arcs and sectors
  → **Paper 2 (Non-Calculator)** *(5 Qs)* · **Paper 4 (Calculator)** *(7 Qs)*

- **E6.4** Surface area and volume
  → **Paper 2 (Non-Calculator)** *(4 Qs)* · **Paper 4 (Calculator)** *(8 Qs)*

- **E6.5** Compound shapes and parts of shapes
  → **Paper 2 (Non-Calculator)** *(10 Qs)* · **Paper 4 (Calculator)** *(20 Qs)*

- **E7.1** Pythagoras' theorem
  → **Paper 2 (Non-Calculator)** *(7 Qs)* · **Paper 4 (Calculator)** *(5 Qs)*

- **E7.2** Right-angled triangles
  → **Paper 4 (Calculator)** *(9 Qs)*

- **E7.3** Exact trigonometric values
  → **Paper 2 (Non-Calculator)** *(9 Qs)*

- **E7.4** Trigonometric functions
  → **Paper 2 (Non-Calculator)** *(7 Qs)* · **Paper 4 (Calculator)** *(5 Qs)*

- **E7.5** Non-right-angled triangles
  → **Paper 2 (Non-Calculator)** *(2 Qs)* · **Paper 4 (Calculator)** *(28 Qs)*

- **E7.6** Pythagoras' theorem and trigonometry in 3D
  → **Paper 4 (Calculator)** *(3 Qs)*

- **E8.1** Transformations
  → **Paper 2 (Non-Calculator)** *(10 Qs)* · **Paper 4 (Calculator)** *(5 Qs)*

- **E8.2** Vectors in two dimensions
  → **Paper 2 (Non-Calculator)** *(5 Qs)* · **Paper 4 (Calculator)** *(7 Qs)*

- **E8.3** Magnitude of a vector
  → **Paper 2 (Non-Calculator)** *(5 Qs)* · **Paper 4 (Calculator)** *(4 Qs)*

- **E9.1** Introduction to probability
  → **Paper 2 (Non-Calculator)** *(6 Qs)* · **Paper 4 (Calculator)** *(3 Qs)*

- **E9.2** Relative and expected frequencies
  → **Paper 2 (Non-Calculator)** *(3 Qs)* · **Paper 4 (Calculator)** *(6 Qs)*

- **E9.3** Probability of combined events
  → **Paper 2 (Non-Calculator)** *(18 Qs)* · **Paper 4 (Calculator)** *(21 Qs)*

- **E10.1** Classifying statistical data
  → **Paper 2 (Non-Calculator)** *(5 Qs)* · **Paper 4 (Calculator)** *(7 Qs)*

- **E10.2** Interpreting statistical data
  → **Paper 2 (Non-Calculator)** *(5 Qs)* · **Paper 4 (Calculator)** *(7 Qs)*

- **E10.3** Discrete and continuous data
  → **Paper 2 (Non-Calculator)** *(5 Qs)* · **Paper 4 (Calculator)** *(7 Qs)*

- **E10.4** Averages and measures of spread
  → **Paper 2 (Non-Calculator)** *(3 Qs)* · **Paper 4 (Calculator)** *(6 Qs)*

- **E10.5** Averages on a calculator
  → **Paper 4 (Calculator)** *(9 Qs)*

- **E10.6** Statistical charts and diagrams
  → **Paper 2 (Non-Calculator)** *(3 Qs)* · **Paper 4 (Calculator)** *(6 Qs)*

- **E10.7** Scatter diagrams
  → **Paper 2 (Non-Calculator)** *(1 Q)* · **Paper 4 (Calculator)** *(8 Qs)*

- **E10.8** Cumulative frequency diagrams
  → **Paper 2 (Non-Calculator)** *(2 Qs)* · **Paper 4 (Calculator)** *(7 Qs)*

1 **(a)** Express 504 as a product of its prime factors.

..... **[2]**

(b) Find the smallest positive integer k such that $504k$ is a square number.

$k =$ ..... **[2]**

2 **(a)** Write 84 as a product of its prime factors.

..... **[2]**

(b) Find the lowest common multiple (LCM) of 84 and 126.

..... **[2]**

3 Here is a list of numbers.

2 9 12 23 36 45 64 81

(a) Write down a number from the list which is:

(i) a multiple of 12,

..... **[1]**

(ii) a cube number,

..... **[1]**

(iii) both a prime number and a factor of 46.

..... **[1]**

(b) Find the lowest common multiple (LCM) of 36 and 64.

..... **[2]**

4 Here is a list of numbers.

$$0.25 \quad \sqrt{7} \quad 8 \quad 9 \quad 13 \quad -5 \quad 21 \quad 64$$

(a) From the list, write down a prime number.

..... **[1]**

(b) From the list, write down an irrational number.

..... **[1]**

(c) From the list, write down an integer less than zero.

..... **[1]**

(d) From the list, write down the reciprocal of 4.

..... **[1]**

(e) From the list, write down a number that is both a square number and a cube number.

..... **[1]**

5 Here is a list of numbers.

8 15 23 27 36 49 64

(a) From the list, write down

(i) a prime number,

..... **[1]**

(ii) a cube number,

..... **[1]**

(iii) a number that is both a square number and a cube number.

..... **[1]**

(b) Express 36 as a product of its prime factors.

..... **[2]**

6 Here is a list of numbers.

2 9 15 16 23 27 64 72

(a) From the list, write down

(i) a prime number,

..... **[1]**

(ii) a cube number,

..... **[1]**

(iii) a number that is both a square number and a cube number,

..... **[1]**

(iv) a triangle number.

..... **[1]**

(b) Express 72 as a product of its prime factors.

..... **[2]**

7 Here is a list of numbers.

$$-5 \qquad -1 \qquad \frac{3}{8} \qquad 8 \qquad 13 \qquad 25 \qquad \sqrt{30} \qquad \pi$$

(a) From this list, write down:

(i) a prime number,

..... **[1]**

(ii) a cube number,

..... **[1]**

(iii) an irrational number,

..... **[1]**

(iv) an integer that is not a natural number.

..... **[1]**

(b) Find the reciprocal of 0.8.
Give your answer as a fraction in its simplest form.

..... **[2]**

8 Two numbers, A and B, are written as products of their prime factors.

$A = 2^3 \times 3^2 \times 5$

$B = 2^2 \times 3^4 \times 7$

(a) Find the highest common factor (HCF) of A and B.

..... **[2]**

(b) Find the lowest common multiple (LCM) of A and B.
Give your answer as a product of prime factors.

..... **[2]**

(c) The integer C is such that:

- the HCF of A and C is 6
- the LCM of A and C is $2^3 \times 3^2 \times 5 \times 11$

Find the value of C.

$C =$ ..... **[3]**

9 **(a)** Let x be a rational number and y be an irrational number.

State whether $x + y$ is rational or irrational.

..... **[1]**

(b) Give an example of an irrational number y to show that y^2 can be a rational number.

[1]

(c) The nth term of a sequence is given by the expression

$$n^2 + n + 41$$

(i) Work out the 3rd term of this sequence, and state whether it is a prime number.

..... **[2]**

(ii) Explain why the 41st term of this sequence is not a prime number.

..... **[2]**

10 The number N is written as a product of its prime factors as $N = 2^3 \times 3^x \times 5^2$.

(a) The highest common factor (HCF) of N and 540 is 180.
Find the value of x.

$x =$ ..... **[3]**

(b) The lowest common multiple (LCM) of N and a number M is $2^4 \times 3^3 \times 5^2 \times 7$.
Given that $M = 2^y \times 3^z \times 7$, find the value of y and the value of z.

$y =$ ..... , $z =$ ..... **[2]**

(c) Find the smallest positive integer k such that Nk is a cube number.

$k =$ ..... **[2]**

11 **(a)** The highest common factor (HCF) of x and 12 is 4.
The lowest common multiple (LCM) of x and 12 is 120.
Find the value of x.

$x =$ ..... **[2]**

(b) **(i)** Express 1008 as a product of its prime factors.

..... **[2]**

(ii) An integer N is a perfect cube.
The highest common factor (HCF) of N and 1008 is 72.
Find the smallest possible positive value of N.

$N =$ ..... **[3]**

12 Let $m = p^3q$ and $n = pq^3$, where p and q are distinct prime numbers.

(a) Express, in terms of p and q:

(i) the highest common factor (HCF) of m and n,

..... **[1]**

(ii) the lowest common multiple (LCM) of m and n.

..... **[1]**

(b) The HCF of m and n is 15.
It is given that $m > n$.
Find the value of p and the value of q.

$p =$ ..... , $q =$ ..... **[3]**

(c) Show algebraically that the product $m \times n$ is a square number.

[2]

(d) Using your values of p and q, find the smallest positive integer k such that $m \times k$ is a cube number.

$k =$ ..... **[2]**

13 Three numbers A, B and C are given as products of their prime factors.

$$A = 2^3 \times 3^2 \times 5 \qquad B = 2^2 \times 3^x \times 5^2 \qquad C = 2^y \times 3 \times 7$$

The highest common factor (HCF) of A and B is 180.
The lowest common multiple (LCM) of B and C is 12600.

Find the value of x and the value of y.

$x =$ ..... , $y =$ ..... **[4]**

14 Two numbers, A and B, are given as a product of their prime factors.

$$A = 2^3 \times 3^2 \times 5 \qquad B = 2^2 \times 3^3 \times 7$$

(a) Find the highest common factor (HCF) of A and B.

..... **[1]**

(b) Find the lowest common multiple (LCM) of A and B.
Give your answer as a product of its prime factors.

..... **[2]**

(c) $A \times k$ is a perfect square.
Find the smallest positive integer value of k.

$k =$ ..... **[2]**

15 Number A and Number B are given as products of their prime factors.

$A = 2^2 \times 3^3 \times 5$

$B = 2^x \times 3 \times 7$

(a) The highest common factor (HCF) of A and B is 6.
Find the value of x.

$x =$ ..... **[2]**

(b) Number $C = 2^2 \times 3^y \times 5^2$.
The lowest common multiple (LCM) of A and C is 8100.
Find the value of y.

$y =$ ..... **[3]**

(c) Here is a list of numbers.

$$\sqrt{11} \qquad -3.5 \qquad 4\pi \qquad \frac{7}{9} \qquad \sqrt{144} \qquad 0.\dot{4}$$

Write down all the irrational numbers from this list.

..... **[2]**

16 **(a)** Find the highest common factor (HCF) of 84 and 120.

..... **[2]**

(b) Find the lowest common multiple (LCM) of 84 and 120.

..... **[2]**

(c) Write down the reciprocal of 0.8 as a fraction in its simplest form.

..... **[2]**

(d) State whether $\sqrt{120}$ is rational or irrational.
Give a reason for your answer.

..... **[2]**

17 **(a)** x, y and z are positive integers.

- x is a prime number.
- y is a cube number greater than 1.
- z is a triangle number.

It is given that $x + y = z$ and $x < 10$.

Find two possible sets of values for x, y and z.

(i) First set of values.

$x =$ ..... , $y =$ ..... , $z =$ ..... **[2]**

(ii) Second set of values.

$x =$ ..... , $y =$ ..... , $z =$ ..... **[2]**

(b) p and q are two different irrational numbers.

Write down an example of p and q such that:

(i) $p + q$ is a rational number.

$p =$ ..... , $q =$ ..... **[1]**

(ii) $p \times q$ is a rational number.

$p =$ ..... , $q =$ ..... **[1]**

18 **(a)** The number 168 can be written as $2^3 \times 3 \times 7$.
The number N can be written as $2^x \times 3^2 \times 7^y$, where x and y are integers.
The highest common factor (HCF) of 168 and N is 84.
The lowest common multiple (LCM) of 168 and N is 3528.

Find the value of x and the value of y.

$x =$ ..... , $y =$ ..... **[3]**

(b) Find the exact value of N.


$N =$ ..... **[1]**

(c) Two flashing lights, A and B, are turned on at the same time.
Light A flashes every 45 seconds.
Light B flashes every 72 seconds.
They both flash together at 10:15:00.
Work out the next time they flash together.

..... **[3]**

19 The universal set U contains all integers from 1 to 10 inclusive.
Set A contains all the even numbers in U.
Set B contains all the prime numbers in U.

(a) Complete the Venn diagram to show the elements of these sets.



[2]

(b) List the elements of $A \cap B$.

..... **[1]**

(c) Find $n(A \cup B)'$.

..... **[1]**

20 $\mathsf{U} = \{x : x \text{ is an integer and } 1 \leq x \leq 10\}$
$A = \{x : x \text{ is a prime number}\}$
$B = \{x : x \text{ is a factor of } 12\}$

(a) List the members of set A.

..... **[1]**

(b) List the members of set B.

..... **[1]**

(c) Find $n(A \cup B)$.

..... **[1]**

(d) Write down the elements of $A \cap B$.

..... **[1]**

(e) Write down a number that is in U but is not in $A \cup B$.

..... **[1]**

21 The universal set U and sets A and B are defined as follows:

- $\mathsf{U} = \{1, 2, 3, 4, 5, 6, 7, 8, 9, 10\}$
- $A = \{2, 3, 5, 7\}$
- $B = \{1, 3, 5, 7, 9\}$

(a) List the elements of $A \cap B$.

..... **[1]**

(b) Find $n((A \cup B)')$.

..... **[2]**

(c) Write down the correct set notation symbol $(\in, \notin, \subset, \cup, \cap)$ to complete the statement:

$2 \ldots A$

[1]


(d) Write down the correct set notation symbol to complete the statement:

$\{1, 9\} \ldots B$

[1]

22 Let U be the set of integers from 1 to 10 inclusive.
Let A be the set of prime numbers.
Let B be the set of odd numbers.

(a) Complete the Venn diagram to show the elements of these sets.



[2]


(b) List the elements of $A \cap B'$.

..... **[1]**

(c) Find $n(A \cup B)$.

..... **[1]**

(d) Write down the set notation for the shaded region in the Venn diagram above.



..... **[1]**

23 The universal set U and sets P and F are defined as follows:

- $\mathsf{U} = \{x : x \text{ is an integer} 1 \leq x \leq 10\}$
- $P = \{x : x \text{ is a prime number}\}$
- $F = \{x : x \text{ is a factor of } 18\}$

(a) Write down the elements of set F.

$F = \{$ ..... $\}$ **[1]**

(b) List the elements of $P \cap F$.

$P \cap F = \{$ ..... $\}$ **[1]**

(c) Find $n((P \cup F)')$.


..... **[2]**

(d) Write down the correct set symbol, $\in$ or $\notin$, to make this statement true:

$5 \ldots\ldots F$

[1]

24 The Venn diagram shows the number of elements in each region for sets A, B, and C.



(a) Given that $n(\mathsf{U}) = 35$, find the value of x.

$x =$ ..... **[2]**

(b) Find $n(A \cap C')$.

..... **[1]**


(c) Use set notation to describe the set of elements whose number is 4.

..... **[1]**

(d) Find $n(A \cup (B \cap C))$.

..... **[2]**

25 The Venn diagram shows the number of elements in sets A and B, in terms of x.
U is the universal set and $n(\mathsf{U}) = 52$.




(a) Form an equation in x and solve it.

$x =$ ..... **[3]**

(b) Find $n(A \cup B')$.

..... **[2]**


(c) On the Venn diagram below, shade the region $A \cap B'$.



[1]

26 There are 45 students in a class.
Let U be the set of students in this class.
Let P be the set of students who study Physics.
Let C be the set of students who study Chemistry.
$n(P) = 28$
$n(C) = 22$
$n(P \cap C) = x$
$n((P \cup C)') = 4$

(a) Complete the Venn diagram in terms of x.



[2]


(b) Form an equation in x and solve it to find the value of x.

$x =$ ..... **[2]**

(c) Find $n(P \cap C')$.

..... **[1]**

27 The Venn diagram shows the number of elements in each region for the sets P, Q and R.




(a) The total number of elements in the universal set is 45.
Form an equation in terms of x and solve it to find the value of x.

$x =$ ..... **[3]**

(b) Find $n(P \cup Q)$.

..... **[1]**

(c) On the Venn diagram below, shade the region $P \cap (Q \cup R)'$.




[1]

28 In a class of 40 students:

- 25 study French (F)
- 18 study Spanish (S)
- 5 study neither French nor Spanish

(a) Complete the Venn diagram to show this information.



[3]

(b) Find $n(F \cap S')$.

..... **[1]**

(c) A student is chosen at random from the class.
Find the probability that they study exactly one of the two languages.

..... **[2]**

(d) Two students are chosen at random from those who study French.
Find the probability that both also study Spanish.

..... **[1]**


29 In a survey of 100 people, information was collected about the pets they own.

Let U be the set of people surveyed.
Let C be the set of people who own a cat.
Let D be the set of people who own a dog.
Let B be the set of people who own a bird.

- 8 people own all three pets.
- 12 people own a cat and a dog but not a bird.
- 7 people own a cat and a bird but not a dog.
- 10 people own a dog and a bird but not a cat.
- 3 people do not own any of these pets.

The number of people who own only a cat is x.
The number of people who own only a bird is also x.
The number of people who own only a dog is $2x$.

(a) Complete the Venn diagram to show this information.



[3]

(b) Find the value of x.


$x =$ ..... **[2]**

(c) Find $n((C \cup D) \cap B')$.

..... **[2]**

30 There are 60 people in a sports club.

- 30 play Tennis (T)
- 28 play Badminton (B)
- 25 play Squash (S)
- 12 play both Tennis and Badminton
- 9 play both Tennis and Squash
- 11 play both Badminton and Squash
- 4 play all three sports



(a) Complete the Venn diagram.

[3]

(b) Find $n(T \cup B \cup S)'$.

..... **[1]**

(c) Find $n(T \cap (B \cup S))$.

..... **[1]**

(d) Use set notation to describe the region containing the number 9 (the number of people who play Badminton only).


..... **[2]**

31 In a sports club, the universal set U represents all the students.
The set T represents students who play Tennis.
The set C represents students who play Cricket.

You are given the following information:

- $n(\mathsf{U}) = 50$
- $n(T) = 3x$
- $n(C) = 4x$
- $n(T \cap C) = x$
- $n((T \cup C)') = 2$

(a) Complete the Venn diagram, writing the number of students in each region in terms of x where appropriate.



[3]


(b) Form an equation in x and solve it to find the value of x.

$x =$ ..... **[2]**

(c) Find the number of students who play Tennis but not Cricket.

..... **[1]**

(d) On the Venn diagram below, shade the region $T' \cup C$.




[2]

32 In a survey of 103 people, the following information was recorded about their use of three types of transport: Bus (B), Train (T), and Car (C).

- 15 people use all three types.
- 28 people use both a Train and a Car.
- 22 people use both a Bus and a Car.
- 30 people use both a Bus and a Train.
- x people use only a Bus.
- The number of people who use only a Train is twice the number of people who use only a Bus.
- The number of people who use only a Car is 5 more than the number of people who use only a Bus.
- 8 people do not use any of these types of transport.

(a) Complete the Venn diagram in terms of x to show this information.



[3]

(b) Find the value of x.

$x =$ ..... **[2]**

(c) Find $n(T \cup C)$.

..... **[1]**


(d) A person is chosen at random from those who use a Train.
Find the probability that this person also uses a Car but does not use a Bus.

..... **[2]**

33 In a year group of 120 students, they can choose to study Music (M), Drama (D), and Art (A).

- 12 students study all three subjects.
- 25 students study Music and Drama.
- 32 students study Drama and Art.
- 28 students study Music and Art.
- 50 students study Music.
- 65 students study Drama.
- x students study Art only.
- 15 students study none of these three subjects.

(a) Use this information to complete the Venn diagram.



[3]


(b) Find the value of x.

$x =$ ..... **[2]**

(c) Find $n(M \cup (D \cap A))$.

..... **[2]**

(d) On the Venn diagram below, shade the region $M \cap (D \cup A)'$.



[1]

34 **(a)** Write down the value of $\sqrt{121}$.

..... **[1]**

(b) Work out the value of

$$4^3 - \sqrt[3]{27}$$

..... **[2]**

35 **(a)** Work out the value of

$$\sqrt{169} + 2^3$$

..... **[2]**

(b) Work out the value of

$$4^2 \times \sqrt[3]{125}$$

..... **[2]**

36 **(a)** Work out the value of

$3^3 - \sqrt{144}$

..... **[2]**

(b) Work out the value of

$2^4 \times \sqrt[3]{125}$

..... **[2]**

37 **(a)** Write down the value of

$\sqrt{196}$

..... **[1]**

(b) Work out the value of

$4^3 - \sqrt[3]{125}$

..... **[2]**

(c) Calculate the exact value of

$\sqrt{13^2 - 12^2}$

..... **[2]**

38 **(a)** Work out the value of

$$5^2 - \sqrt{121}$$

..... **[1]**

(b) Work out the value of

$$\sqrt[3]{27} \times 2^4$$

..... **[2]**

(c) Work out the exact value of

$$\frac{6^2 + \sqrt{64}}{\sqrt[3]{125}}$$

..... **[2]**

39 **(a)** Work out the exact value of

$$\frac{5^2 + \sqrt{144}}{2^3 - \sqrt[3]{1}}$$

..... **[2]**

(b) Find the value of k when

$$3^k = \sqrt{81} \times 3^2$$

$k =$ ..... **[2]**

40 **(a)** Calculate the value of

$$\frac{4^3 + \sqrt{225}}{\sqrt[3]{64}}$$

Give your answer as a decimal.

..... **[3]**

(b) Given that $p = 5$ and $q = -2$, work out the value of

$$p^2 - q^3$$

..... **[2]**

41 **(a)** A cube has a volume of 125 cm^3.

Find the length of one edge of the cube.

..... cm **[1]**

(b) Work out the total surface area of this cube.

..... cm^2 **[2]**

(c) Given that $\sqrt{N} = 3^3 - 4^2$, find the value of N.

$N =$ ..... **[2]**

42 Solve the following equations.

(a)

$\sqrt[3]{3x+4} = 4$

$x =$ ..... **[3]**

(b)

$x^2 = 4^3 + 6^2$

$x =$ ..... , or $x =$ ..... **[2]**

(c)

$\sqrt{x} + 2\sqrt{x} = 21$

$x =$ ..... **[2]**

43 **(a)** Find the integer k such that

$$\sqrt[3]{k} = 4^2 - 3^2 - 2$$

$k =$ ..... **[2]**

(b) Calculate the exact value of

$$\frac{\sqrt{10^2 - 6^2}}{\sqrt[3]{125} - 3^2}$$

..... **[3]**

(c) Given that $y > 0$ and $\sqrt[3]{y} + \sqrt[3]{y} = 6$, work out the value of y^2.

..... **[3]**

44 **(a)** Calculate the value of

$$\frac{\sqrt[3]{450} + 12.5^2}{3.4^3 - \sqrt{80}}$$

Write down all the figures on your calculator display.

..... **[2]**

(b) Write your answer to part (a) correct to 3 decimal places.

..... **[1]**

(c) Find the largest integer n such that $n^4 < 2000$.

$n =$ ..... **[2]**

45 **(a)** **(i)** Calculate the value of

$$\sqrt[3]{8.5^2 - 4.1^3}$$

Write down your full calculator display.

..... **[2]**

(ii) Write your answer to part (a)(i) correct to 3 significant figures.

..... **[1]**

(b) Calculate the exact value of

$$\frac{4.8^2}{\sqrt{0.0576}}$$

..... **[2]**

46 **(a)** Calculate the value of

$$\frac{\sqrt[3]{45.6} + 2.1^4}{3.8^2}$$ Give your answer correct to 3 significant figures.

..... **[2]**

(b) A solid cube has a volume of 4913 cm^3.
Calculate the total surface area of the cube.

..... cm^2 **[3]**

47 **(a)** Calculate the value of

$\sqrt[5]{7.59375}$

..... **[1]**

(b) Find the two possible values of x in the equation

$2x^4 - 17 = 1233$

$x =$ ..... , $x =$ ..... **[3]**

(c) It is given that

$$y = \sqrt{\frac{3a^2 + b}{c}}$$

Work out the value of y when $a = 4.2$, $b = -5.7$ and $c = 1.8$.
Give your answer correct to 2 decimal places.

$y =$ ..... **[3]**

48 **(a)** Calculate the value of

$$\frac{\sqrt[4]{150000}}{4.5^3 - \sqrt{75}}$$ Give your answer correct to 3 decimal places.

..... **[2]**

(b) x is a positive integer such that $\sqrt[3]{x^2} = 16$.
Find the value of x.

$x =$ ..... **[3]**

(c) The volume of a sphere is equal to the volume of a cube with edge length 12 cm.
Calculate the radius of the sphere.
The volume, V, of a sphere with radius r is $V = \frac{4}{3}\pi r^3$.

..... cm **[3]**

49 **(a)** Write the following in order of size, starting with the smallest.

$$0.65 \qquad \frac{2}{3} \qquad 64\% \qquad \frac{5}{8}$$

..... **[2]**

(b) Write 0.85 as a fraction in its simplest form.

..... **[1]**

50 **(a)** Write $0.\dot{7}$ as a fraction.

..... **[1]**

(b) Write $0.\dot{3}\dot{6}$ as a fraction in its simplest form.

..... **[2]**

51 **(a)** Write 45% as a fraction in its simplest form.

..... **[1]**

(b) Write $\frac{9}{25}$ as a decimal.

..... **[1]**

(c) Write 2.04 as a mixed number in its simplest form.

..... **[2]**

52 **(a)** Write 0.45 as a fraction in its simplest form.

..... **[1]**

(b) Write $\frac{3}{8}$ as a decimal.

..... **[1]**

(c) Write these in order of size, starting with the smallest.

$$\frac{5}{12} \qquad 0.4 \qquad 42\% \qquad \frac{3}{7}$$

..... **[2]**

53 **(a)** Write 0.075 as a fraction in its simplest form.

..... **[2]**

(b) Write $3\frac{2}{5}$ as a decimal.

..... **[1]**

(c) Write the following in order of size, starting with the smallest.

$$62\% \qquad \frac{5}{8} \qquad 0.6 \qquad \frac{7}{11}$$

..... **[2]**

54 The length of a rectangle is increased by $33.\dot{3}\%$.
The width of the rectangle is decreased by $x\%$.
The overall area of the rectangle increases by $16.\dot{6}\%$.
Calculate the value of x.

$x =$ ..... **[5]**

55 **(a)** A tank is initially $\frac{1}{5}$ full of water.
When 24 litres of water are added, the tank is exactly 60% full.
Work out the total capacity of the tank.

..... litres **[3]**

(b) A bottle contains x millilitres of juice.
30% of the juice is poured out.
Then $\frac{1}{4}$ of the remaining juice is poured out.
The bottle now contains 210 millilitres of juice.
Work out the value of x.

$x =$ ..... **[3]**

56 A bag contains red counters, blue counters and green counters.
The number of red counters is x.
The number of blue counters is $2x + 5$.
The number of green counters is $3x - 1$.

(a) Exactly 35% of the counters in the bag are blue.
Work out the value of x.

$x =$ ..... **[4]**

(b) Work out the percentage of the counters in the bag that are green.
Write your answer as a mixed number.

..... % **[3]**

57 **(a)** Write the recurring decimal $0.2\dot{4}\dot{5}$ as a fraction in its simplest form.
You must show all your working.

..... **[3]**

(b) The fraction $\frac{N}{42}$, where N is a positive integer, is in its simplest form.
Explain why the decimal equivalent of $\frac{N}{42}$ is a recurring decimal.

.....

..... **[2]**

(c) The recurring decimal $0.\dot{x}\dot{y}$, where x and y are distinct digits, can be written as a fraction $\frac{k}{11}$, where k is an integer.
Find the value of $x + y$.

$x + y =$ ..... **[2]**

58 **(a)** By converting the recurring decimal to a fraction, show that

$$0.2\dot{7} = \frac{5}{18}$$

[2]

(b) In a school, $\frac{2}{5}$ of the students are boys and the rest are girls.
45% of the boys play a sport.
30% of the girls play a sport.

(i) Work out the percentage of all students in the school who play a sport.

..... % **[3]**

(ii) There are 144 students who play a sport.
Work out the total number of students in the school.

..... **[2]**

59 In a school of 840 students, 35% are in the lower school, $\frac{3}{8}$ are in the middle school, and the rest are in the upper school.

(a) Work out the number of students in the middle school.

..... **[1]**

(b) Work out the percentage of students in the upper school.

..... % **[2]**

(c) 15% of the middle school students play a musical instrument.
Work out the fraction of the total number of students who are middle school students that play a musical instrument.
Give your answer in its simplest form.

..... **[2]**

60 In a school, $\frac{3}{8}$ of the students are in the Junior section, 45% are in the Middle section, and the rest are in the Senior section.
There are 140 students in the Senior section.

(a) Work out the total number of students in the school.

..... **[3]**

(b) The number of boys in the Junior section is 150% of the number of girls in the Junior section.
Work out the number of girls in the Junior section.

..... **[3]**

61 **(a)** Write the following in order of size, starting with the smallest.

$$\frac{5}{7} \qquad 71\% \qquad 0.714 \qquad \frac{3}{4}$$

Smallest ..... **[2]**

(b) Find the fraction which is exactly halfway between $\frac{2}{5}$ and $\frac{5}{8}$.
Give your answer in its simplest form.

..... **[2]**

(c) A fraction is equivalent to $\frac{4}{9}$.
The difference between the denominator and the numerator is 35.
Find the fraction.

..... **[2]**

62 A company has three departments: Sales, Production, and Administration.

- $\frac{2}{5}$ of the employees are in Sales.
- 28% of the employees are in Production.
- The remaining 24 employees are in Administration.

(a) Work out the total number of employees in the company.

..... **[3]**

(b) 60% of the Sales employees are women.
$\frac{4}{7}$ of the Production employees are men.
Half of the Administration employees are women.
Work out the percentage of the whole company who are women.

..... % **[4]**

63 **(a)** A television costs \$450. In a sale, the price is reduced by 18%.
Calculate the sale price of the television.

\$ ..... **[2]**

(b) In the same sale, a laptop is sold for \$629 after a 15% reduction.
Calculate the original price of the laptop.

\$ ..... **[3]**

(c) The price of a tablet is increased by $\frac{1}{5}$.
Later, this new price is decreased by 20%.
Calculate the overall percentage decrease in the price of the tablet.

..... % **[2]**

64 Insert one of the symbols $<$, $>$ or $=$ to make each statement correct.

(i) $0.45 \ldots\ldots \dfrac{2}{5}$

(ii) $-3 \ldots\cdots -7$

[2]

65 Write the following in order of size, starting with the smallest.

$$\frac{5}{8} \qquad 0.62 \qquad 61.5\% \qquad \frac{3}{5}$$

..... **[2]**

66 Write $<, >$ or $=$ to make each statement true.

(a) $0.45 \ldots\ldots\ldots \frac{4}{9}$

[1]

(b) $2^{-3} \ldots\ldots\ldots 0.125$

[1]

(c) $-1.5 \ldots\ldots\ldots \cdots -\frac{7}{5}$

[1]

67 **(a)** Insert one of the symbols $<, =$ or $>$ to make each statement true.

(i) $\frac{5}{8} \ldots 0.65$

[1]

(ii) $72\% \ldots \frac{18}{25}$

[1]

(b) Write the following in order of size, starting with the smallest.

$$0.55 \qquad \frac{6}{11} \qquad 54\% \qquad \frac{5}{9}$$

..... **[2]**

68 **(a)** Insert $<$, $>$ or $=$ to make each statement correct.

(i)

$$0.4 \ldots\ldots \frac{2}{5}$$

[1]

(ii)

$$-7 \ldots\ldots -9$$

[1]

(iii)

$$2^3 \ldots\ldots \sqrt{25}$$

[1]

(b) Write these in order of size, starting with the smallest.

$$33\% \qquad \frac{1}{3} \qquad 0.3 \qquad \frac{3}{8}$$

..... **[2]**

69 **(a)** Given that $0 < p < 1$, write the following expressions in order of size, starting with the smallest.

$$p \qquad p^2 \qquad \sqrt{p} \qquad \frac{1}{p}$$

..... **[2]**

(b) Given that $-1 < q < 0$, write $<, >$ or $=$ to make each statement true.

(i) $q^3 \ldots\ldots\ldots q$

[1]

(ii) $\frac{1}{q} \ldots\ldots\ldots q$

[1]

70 **(a)** Given that $0 < x < 1$, write the following expressions in order of size, starting with the smallest.

$$x \qquad x^2 \qquad \frac{1}{x} \qquad \sqrt{x}$$

..... **[2]**

(b) Given that $x < -1$, write the following expressions in order of size, starting with the largest.

$$x \qquad x^3 \qquad x^{-2} \qquad \frac{1}{x}$$

..... **[2]**

71 **(a)** Given that $x = -0.4$, write the following in order of size, starting with the smallest.

$$x \qquad x^2 \qquad x^3 \qquad \frac{1}{x}$$

..... **[2]**

(b) Insert the correct symbol ($<, =$, or $>$) to make the statement true.
Explain your reasoning.

$5\sqrt{3} \ldots 4\sqrt{5}$

[2]

(c) Find the integer m that satisfies the inequality

$$\frac{m}{7} < \frac{5}{9} < \frac{m+1}{7}$$

$m =$ ..... **[3]**

72 **(a)** It is given that $0 < p < 1$.
Write the following values in order of size, starting with the smallest.

$$p \qquad \sqrt{p} \qquad p^2 \qquad \frac{1}{p}$$

..... **[2]**

(b) Write the correct symbol $<$, $>$ or $=$ to complete each logical statement.

(i) If $x < 0$, then $x^3 \ldots\ldots x^2$.

[1]

(ii) If $x > 1$, then $x^{-1} \ldots\ldots 1$.

[1]

(c) Given $a = 2^{50}$, $b = 3^{30}$ and $c = 5^{20}$, write a, b and c in order of size, starting with the smallest.
You must show all your working.

..... **[3]**

73 Write the following in order of size, starting with the smallest.

$$\frac{4}{33} \qquad 0.12 \qquad 12.1\% \qquad \sqrt{0.015}$$

..... **[2]**

74 Write the following masses in order of size, starting with the largest.

5.2×10^{-4} kg 0.04 g 4.5×10^{3} mg 6×10^{-7} tonnes

..... **[3]**

75 **(a)** Here are four numerical expressions.

$a = 3.1 \times 10^3$
$b = \sqrt{9.8 \times 10^6}$
$c = 5^5$
$d = 56^2$

Work out the value of each expression and write them in order of size, starting with the largest.

..... **[3]**

(b) State the integer n such that

$$n < \sqrt{250} < n + 1$$

$n =$ ..... **[1]**

76 **(a)** Write these numbers in order of size, starting with the smallest.

$$5^{-3} \qquad 8.5 \times 10^{-3} \qquad 0.00805 \qquad \frac{1}{120}$$

..... **[2]**

(b) The mass, m kilograms, of a bag of rice is 12 kg, correct to the nearest whole kilogram. Write down the inequality representing the possible values of m.

..... **[2]**

77 **(a)** Write the following numbers in order of size, starting with the smallest.

$$\frac{7}{8} \qquad 86\% \qquad 0.865 \qquad \frac{5}{6}$$

..... **[2]**

(b) Write down a fraction, f, such that $\frac{1}{3} < f < \frac{1}{2}$.

..... **[2]**

78 **(a)** Given that $x > 0$ and $y < 0$, insert the correct symbol $<$, $>$ or $=$ to make the statement correct.

$x^2y \ldots\ldots 0$

[1]

(b) Write these expressions in order of size, starting with the smallest, when $a = 0.5$.

$$a^2 \qquad a \qquad \sqrt{a} \qquad \frac{1}{a}$$

..... **[2]**

(c) State whether the following statement is True or False.
Give a reason for your answer.
If $p > q$, then $\frac{1}{p} < \frac{1}{q}$ for all non-zero real numbers p and q.

..... **[2]**

79 **(a)** The temperature at midnight was $-5°C$.
By noon, the temperature had risen by $14°C$.
Write down the temperature at noon.

..... **[1]**

(b) Work out the difference between $-8°C$ and $17°C$.

..... **[1]**

(c) Work out the value of

$$24 - 12 \div (3 - 7)$$

..... **[2]**

80 **(a)** Work out the value of

$$12 - 2 \times (-3) + 4^2$$

..... **[2]**

(b) Work out the value of

$$\frac{3}{4} + \frac{1}{6} \times \frac{3}{2}$$

Give your answer in its simplest form.

..... **[2]**

81 **(a)** Work out the value of

$$12 - 3 \times (-4) + (15 \div -3)$$

..... **[2]**

(b) Work out the value of

$$\frac{2}{5} \div \frac{4}{7} - \frac{1}{10}$$

Give your answer as a fraction in its simplest form.

..... **[2]**

82 **(a)** Work out the value of

$$15 - 3 \times (4 - (-2))$$

..... **[2]**

(b) Work out the value of

$$2\frac{1}{4} \div \frac{3}{8} - 1\frac{1}{2}$$

Give your answer in its simplest form.

..... **[3]**

83 **(a)** Work out the value of

$12 - 4 \times (5 - 9)$

..... **[2]**

(b) Work out the value of

$1.2 \div 0.04$

..... **[2]**

(c) The temperature at 04:00 was $-5.2°C$.
By 14:00, the temperature had risen by $8.7°C$.
Work out the temperature at 14:00.

..... °C **[1]**

84 **(a)** Work out the exact value of

$\frac{5}{6} + \frac{2}{3} \times \frac{9}{10}$ Give your answer as a mixed number in its simplest form.

..... **[3]**

(b) Work out the exact value of

$2\frac{1}{4} \div \left(1\frac{1}{8} - \frac{3}{4}\right)$

..... **[3]**

85 **(a)** Calculate the exact value of

$$2\frac{2}{5} \div \left(1\frac{1}{3} - \frac{5}{6}\right)$$

Show all your working and give your answer as a mixed number in its simplest form.

..... **[4]**

(b) Work out the exact value of

$$\frac{0.45 \times 1.2}{0.06}$$

..... **[3]**

86 Solve the equation, giving your answer as a fraction in its simplest form.

$$\frac{x+\frac{1}{3}}{2-0.4} = -1\frac{1}{4}$$

$x =$ ..... **[5]**

87 Without using a calculator, work out

$$\frac{3\frac{2}{3}+2\frac{1}{4}\times 1\frac{1}{3}}{4\frac{1}{2}-1\frac{5}{6}}$$

You must show all your working and give your answer as a fraction in its simplest form.

..... **[5]**

88 **(a)** Work out the value of

$$2\frac{2}{3} \div \left(1\frac{1}{4} - \frac{5}{6}\right)$$

Give your answer as a mixed number in its simplest form.

..... **[4]**

(b) A rectangular field has a length of $12\frac{1}{2}$ m and an area of $83\frac{1}{3}$ m^2.
Work out the perimeter of the field.
Give your answer as a mixed number in its simplest form.

..... m **[4]**

89 **(a)** At 06:00, the temperature in a city is $-4.5°C$.
By 14:00, the temperature has risen by $11.2°C$.
By 22:00, the temperature has dropped by $8.1°C$ from the temperature at 14:00.

Work out the temperature at 22:00.

..... **[2]**

(b) Use your calculator to work out the value of

$$\frac{4.72^2 - \sqrt[3]{18.5}}{3.1 \times (0.45 + 1.2)}$$

Give your answer correct to 3 significant figures.

..... **[2]**

90 A water tank has a capacity of 85.5 litres.
It is initially completely empty.
A hose pumps water into the tank at a rate of $\frac{3}{4}$ of a litre per minute.
At the same time, a leak in the tank loses water at a rate of 0.45 litres per minute.

Work out how long it takes to completely fill the tank.
Give your answer in hours and minutes.

hours ..... , minutes ..... **[4]**

91 The temperature of a liquid is $-4.5°C$.
It is heated so that its temperature rises by $2.8°C$ every minute for 15 minutes.

(a) Work out the temperature of the liquid after the first 15 minutes.

..... **[2]**

(b) After the 15 minutes, the liquid is cooled so its temperature falls by $1.2°C$ every minute for m minutes.
After the m minutes of cooling, the temperature of the liquid is $9.9°C$.
Work out the value of m.

$m =$ ..... **[3]**

92 A large container has a total capacity of 1000 litres.

(a) The container is initially $\frac{2}{5}$ full of water.
Water is pumped into the container at a constant rate of 18.5 litres per minute for 20 minutes.
Work out the volume of water in the container after these 20 minutes.

..... litres **[3]**

(b) Water then leaks from the container at a constant rate of 4.5 litres per minute.
Calculate the time taken for the container to become exactly half full.

..... minutes **[3]**

(c) Once the container is half full, the leak is fixed.
A different pump then fills the container completely in 1 hour and 20 minutes.
Calculate the average rate at which this pump fills the container.
Give your answer in litres per minute.

..... litres per minute **[3]**

93 **(a)** A tank contains 450.5 litres of oil.
The oil is poured into bottles.
Each bottle holds exactly 1.25 litres.
Work out the number of completely filled bottles and the volume of oil left over.

Number of bottles = ... , Volume left over = ... **[3]**

(b) A machine pumps water at a rate of 1800 litres per minute.
It is used to fill a rectangular pool of length 24.5 m, width 12 m and depth 1.8 m.
($1\ m^3 = 1000$ litres)

Calculate the time taken to fill the pool completely.
Give your answer in hours and minutes.

hours, minutes **[4]**

(c) A freezer is set at $-18°C$.
During a power cut, the temperature inside the freezer rises by $1.5°C$ per hour.
The power cut lasts for 6 hours and 20 minutes.
Work out the temperature in the freezer when the power is restored.

..... °C **[3]**

94 **(a)** Find the value of 7^0.

..... **[1]**

(b) Find the value of $81^{\frac{1}{2}}$.

..... **[1]**

(c) Write as a single power of x.

$x^5 \times x^{-2}$

..... **[1]**

95 **(a)** Work out the value of

$125^{\frac{1}{3}}$

..... **[1]**

(b) Work out the value of

6^{-2}

..... **[1]**

(c) Simplify.

$t^3 \times t^5$

..... **[1]**

96 **(a)** Simplify.

$m^5 \times m^2$

..... **[1]**

(b) Find the value of

5^{-2}

..... **[1]**

(c) Find the value of

$49^{\frac{1}{2}}$

..... **[1]**

97 **(a)** Find the value of

3^{-2}

..... **[1]**

(b) Find the value of

$\left(\frac{8}{27}\right)^{\frac{1}{3}}$

..... **[1]**

(c) Simplify.

$x^4 \times x^{-2}$

..... **[1]**

(d) Simplify.

$(y^3)^4$

..... **[1]**

98 **(a)** Work out the value of

$$16^{-\frac{3}{4}}$$

..... **[2]**

(b) Simplify.

$$\frac{12w^6y^3}{3w^{-2}y^5}$$

..... **[3]**

99 **(a)** Simplify.

$(4x^3y)^2$

..... **[2]**

(b) Simplify.

$$\frac{18a^6b^2}{6a^2b^5}$$

..... **[2]**

(c) Work out the value of

$27^{-\frac{2}{3}}$

..... **[2]**

100 **(a)** Find the exact value of each of the following.

(i) $64^{\frac{2}{3}}$

..... **[2]**

(ii) 2^{-4}

..... **[1]**

(b) **(i)** Simplify fully.

$$\frac{15a^6b^3}{3a^2b^7}$$

..... **[2]**

(ii) Simplify fully.

$$(4m^6n^{-2})^{\frac{1}{2}}$$

..... **[2]**

101 **(a)** Find the exact value of

$16^{-\frac{3}{4}}$

..... **[2]**

(b) Simplify fully.

$(5x^3y^{-2})^2$

..... **[2]**

(c) Solve the equation.

$2^{3x-1} \times 4^x = 32$

$x =$ ..... **[3]**

102 **(a)** Simplify.

$$\left(\frac{25x^4}{16y^8}\right)^{-\frac{1}{2}}$$

..... **[3]**

(b) Solve the equation.

$$2^{3x-1} \times 4^{x+2} = 8^{x-3}$$

$x =$ ..... **[4]**

103 **(a)** Simplify fully.

$$\left(\frac{16x^8}{y^4}\right)^{-\frac{1}{2}}$$

..... **[3]**

(b) Solve the equation.

$$4^x - 2^{2x+1} + 8 = 0$$

$x =$ ..... **[4]**

104 **(a)** Simplify fully.

$$\left(\frac{16x^{12}}{81y^{8}}\right)^{-\frac{3}{4}}$$

..... **[3]**

(b) Find the value of x when

$$8^{x} = 4^{x+3}$$

$x =$ ..... **[3]**

(c) Solve the equation.

$$5 \times 25^{k} = 125$$

$k =$ ..... **[3]**

105 **(a)** Find the value of k when $5^k = \dfrac{1}{\sqrt{125}}$.

$k =$ ..... **[3]**

(b) Simplify fully.

$$\left(\frac{16x^{12}}{81y^8}\right)^{-\frac{3}{4}}$$

..... **[3]**

(c) Solve the equation.

$$2^{2x-1} \times 4^x = 32$$

$x =$ ..... **[3]**

106 **(a)** Write 450000 in standard form.

..... **[1]**

(b) Write 3.08×10^{-4} as an ordinary number.

..... **[1]**

(c) Work out.

$(2 \times 10^3) \times (4 \times 10^5)(0)$

Give your answer in standard form.

..... **[1]**

107 **(a)** Write 452000 in standard form.

..... **[1]**

(b) Write 6.07×10^{-3} as an ordinary number.

..... **[1]**

(c) Work out $(4 \times 10^4) \times (3 \times 10^5)$.
Give your answer in standard form.

..... **[2]**

108 **(a)** Write 0.00714 in standard form.

..... **[1]**

(b) Write 5.02×10^4 as an ordinary number.

..... **[1]**

(c) Work out the value of

$$(3 \times 10^5) \times (6 \times 10^4)$$

Give your answer in standard form.

..... **[2]**

109 Consider the following numerical values.

(a) Convert the following numbers as instructed.

(i) Write 0.00034 in standard form.

..... **[1]**

(ii) Write 5.2×10^4 as an ordinary number.

..... **[1]**

(b) Work out, giving your answer in standard form:

$(2 \times 10^3) \times (4 \times 10^5)$

..... **[2]**

110 **(a)** Write 0.000452 in standard form.

..... **[1]**

(b) Write 3.71×10^5 as an ordinary number.

..... **[1]**

(c) Calculate, giving your answer in standard form.

$(2 \times 10^4) \times (4 \times 10^3)$

..... **[2]**

111 The mass of the Earth is 5.97×10^{24} kg.
The mass of the Moon is 7.35×10^{22} kg.

(a) Calculate the difference between the mass of the Earth and the mass of the Moon.
Give your answer in standard form.

..... kg **[2]**

(b) The mass of the Sun is 1.99×10^{30} kg.
Find the value of k when

mass of the Sun = k ×mass of the Earth
Give your answer to 3 significant figures.

$k =$ ..... **[3]**

112 **(a)** Work out, giving your answer in standard form.

$$\frac{3.6 \times 10^{-4}}{8 \times 10^{-7}}$$

..... **[3]**

(b) Work out, giving your answer in standard form.

$(5 \times 10^8) + (7 \times 10^7)$

..... **[2]**

(c) Given that $x = 2 \times 10^m$ and $y = 8 \times 10^n$, where m and n are integers.
Find an expression for x^2y in standard form, in terms of m and n.

..... **[2]**

113 **(a)** Work out the exact value of

$$\frac{5.4 \times 10^5 + 6 \times 10^4}{2 \times 10^{-3}}$$

Give your answer in standard form.

..... **[3]**

(b) Evaluate $\sqrt{2.5 \times 10^9}$.
Give your answer in standard form.

..... **[2]**

(c) Given that $p = 2 \times 10^x$ and $q = 8 \times 10^y$, find an expression for $p^2 \div q$.
Give your answer in standard form in terms of x and y.

..... **[3]**

114 $p = 5 \times 10^m$ and $q = 8 \times 10^n$, where m and n are integers.

(a) Find an expression, in terms of m and n, for $p \times q$.
Give your answer in standard form.

..... **[2]**

(b) Find an expression, in terms of m and n, for $\frac{p^2}{q}$.
Give your answer in standard form.

..... **[3]**

(c) Given that $p + q = 5.08 \times 10^m$, work out the exact value of $m - n$.

$m - n =$ ..... **[3]**

115 The distance from the Earth to the Sun is 1.5×10^8 km.

(a) Light travels at a speed of 3.0×10^5 km/s.
Calculate the time taken for light to travel from the Sun to the Earth.
Give your answer in seconds.

..... seconds **[2]**

(b) A spacecraft travels at a constant speed of 2.5×10^4 km/h.
Work out the time it takes the spacecraft to travel from the Earth to the Sun.
Give your answer in hours, in standard form.

..... hours **[2]**

116 The mass of the Sun is 1.99×10^{30} kg.
The mass of Jupiter is 1.90×10^{27} kg.

(a) Calculate how many times greater the mass of the Sun is than the mass of Jupiter.
Give your answer in standard form correct to 3 significant figures.

..... **[3]**

(b) A comet has a mass of 4.5×10^{13} kg.
Calculate the mass of the comet in grams.
Give your answer in standard form.

..... g **[2]**

117 **(a)** Calculate, giving your answer in standard form.

$3.4 \times 10^7 + 2.1 \times 10^6$

..... **[2]**

(b) A computer can do a calculation in 4×10^{-9} seconds.
Work out how many of these calculations the computer can do in 5 minutes.
Give your answer in standard form.

..... **[3]**

118 The mass of the Earth is approximately 5.97×10^{24} kg.
The mass of the Moon is approximately 7.35×10^{22} kg.

(a) Calculate the total mass of the Earth and the Moon.
Give your answer in standard form.

..... kg **[2]**

(b) Calculate how many times greater the mass of the Earth is than the mass of the Moon.
Give your answer correct to the nearest integer.

..... **[2]**

(c) The mass of a planet is $p \times 10^{25}$ kg.
The mass of this planet is 300 times the mass of the Earth.
Work out the value of p, giving your answer correct to 3 significant figures.

$p =$ ..... **[2]**

119 **(a)** The mass of Earth is 5.97×10^{24} kg and the mass of the Moon is 7.35×10^{22} kg.
Calculate the difference between the mass of Earth and the mass of the Moon.
Give your answer in standard form.

..... kg **[2]**

(b) The distance from the Sun to Earth is 1.50×10^{8} km.
The speed of light is 3.00×10^{5} km/s.

Calculate the time it takes for light to travel from the Sun to Earth.
Give your answer in standard form.

..... s **[2]**

(c) The Earth is modelled as a sphere with radius $r = 6.37 \times 10^{6}$ m.
Calculate the surface area of the Earth.
Give your answer in standard form, correct to 3 significant figures.
The surface area, A, of a sphere with radius r is $A = 4\pi r^2$.

..... m^2 **[3]**

120 A large cylindrical water tank has a base radius of 4.5×10^2 cm and a height of 1.2×10^3 cm.

(a) Calculate the volume of the water tank in cm^3.
Give your answer in standard form correct to 4 significant figures.

..... cm^3 **[3]**

(b) The empty tank is being filled with water at a constant rate of 1.5×10^{-4} m^3 per second.
Note that $1\ m^3 = 1 \times 10^6\ cm^3$.

Calculate the time taken to fill the tank completely.
Give your answer in hours, written in standard form correct to 3 significant figures.

..... hours **[4]**

121 Consider the number 47.5381.

(a) Write 47.5381 correct to:

(i) 2 decimal places.

..... **[1]**

(ii) 2 significant figures.

..... **[1]**

(iii) the nearest integer.

..... **[1]**

122 Write the following numbers correct to the specified degree of accuracy.

(a) 782.51 correct to the nearest ten.

..... **[1]**

(b) 0.03561 correct to 2 significant figures.

..... **[1]**

(c) 4.098 correct to 2 decimal places.

..... **[1]**

123 **(a)** Write 48.975 correct to 2 decimal places.

..... **[1]**

(b) Write 0.003058 correct to 2 significant figures.

..... **[1]**

(c) By writing each number correct to 1 significant figure, estimate the value of

$$\frac{7.84 \times 31.6}{0.385}$$

..... **[2]**

124 **(a)** Round the following numbers to the specified degree of accuracy.

(i) 34.567 correct to 2 decimal places.

..... **[1]**

(ii) 0.070381 correct to 2 significant figures.

..... **[1]**

(b) By writing each number correct to 1 significant figure, estimate the value of

$$\frac{61.3 \times 3.92}{0.485}$$

..... **[2]**

125 **(a)** Write 0.04758 correct to 2 significant figures.

..... **[1]**

(b) Write 384.6 correct to the nearest 10.

..... **[1]**

(c) By writing each number correct to 1 significant figure, estimate the value of

$$\frac{58.7 \times 4.12}{19.6 - 2.87}$$

..... **[2]**

126 **(a)** Write 203.498 correct to

(i) 2 decimal places.

..... **[1]**

(ii) 3 significant figures.

..... **[1]**

(iii) the nearest ten.

..... **[1]**

(b) By writing each number correct to 1 significant figure, estimate the value of

$$\frac{6.89 \times 41.2}{0.513}$$

..... **[2]**

127 By writing each number correct to 1 significant figure, estimate the value of

$$\frac{58.4 \times 3.09}{0.187}$$

..... **[3]**

128 **(a)** By writing each number correct to 1 significant figure, estimate the value of

$$\frac{7.84 \times 31.6}{0.381}$$

..... **[2]**

(b) By writing each number correct to 1 significant figure, estimate the value of

$$\frac{\sqrt{98.6} + 19.3}{0.513^2}$$

..... **[3]**

129 **(a)** By writing each number correct to 1 significant figure, estimate the value of

$$\frac{\sqrt{98.5} \times 4.12^2}{0.187}$$

..... **[3]**

(b) A rectangular field has length 489 m and width 215 m.

Estimate the area of the field, in square metres, by rounding each length to 1 significant figure.

..... m^2 **[2]**

130 **(a)** By writing each number correct to 1 significant figure, estimate the value of

$$\frac{\sqrt{3.87 \times 10^6}}{0.0524 \times 21.3}$$

..... **[3]**

(b) Is your estimate in part (a) greater than or less than the exact value of the expression? Give a reason for your answer.

.....

..... **[2]**

131 **(a)** By writing each number correct to 1 significant figure, estimate the value of

$$\frac{\sqrt[3]{8.25} \times 19.3^2}{0.485}$$

..... **[3]**

(b) The number of bacteria in an experiment is given by the formula

$$N = 4.12 \times 10^5 \times (1.95)^t$$

where t is the time in hours.
By writing each base number correct to 1 significant figure, estimate the number of bacteria when $t = 3$.
Give your answer in standard form.

$N \approx$ ..... **[3]**

132 **(a)** By writing each number correct to 1 significant figure, estimate the value of

$$\frac{\sqrt{98.5} \times 4.1^3}{0.192 \times 48.7}$$

You must show all your working.

..... **[3]**

(b) Given that $y = \dfrac{ax}{b-c}$, find an estimate for the value of y when
$a = 3.92$, $x = 0.0514$, $b = 1.08$ and $c = 0.812$.
Write all numbers correct to 1 significant figure in your calculation.

$y =$ ..... **[3]**

133 **(a)** By writing each number correct to 1 significant figure, estimate the value of

$$\frac{38.7 \times 5.12^2}{\sqrt[3]{996}}$$

..... **[3]**

(b) A solid metal cylinder has radius 3.89 cm and height 11.2 cm.

By writing each dimension correct to 1 significant figure, estimate the volume of the cylinder.
Take the value of π to be 3.

..... cm^3 **[3]**

(c) A train travels a distance of 4210 km at an average speed of 585 km/h.

By writing both the distance and the speed correct to 1 significant figure, estimate the time taken for the journey.
Give your answer in hours and minutes.

hours, minutes **[3]**

134 A rectangular field has length 125 m and width 74 m.

(a) Calculate the area of the field.
Give your answer correct to 2 significant figures.

..... m^2 **[2]**

(b) A different field is a square of side x metres.
The area of this square field is equal to the area of the rectangular field.
Calculate the value of x.
Give your answer correct to 1 decimal place.

$x =$ ..... **[3]**

135 A solid metal cone has radius $r = 5.82$ cm and height $h = 11.4$ cm.

(a) Calculate the volume of the cone.
Give your answer correct to 3 significant figures.
The volume, V, of a cone with radius r and height h is $V = \frac{1}{3}\pi r^2 h$.

..... cm^3 **[2]**

(b) By writing each of the values 5.82, 11.4 and π correct to 1 significant figure, estimate the volume of the cone.
Show your working.

..... cm^3 **[2]**

(c) A worker packs these cones into boxes.
Each box holds exactly 12 cones.
The factory produces 850 cones a day.
Calculate the maximum number of boxes that can be completely filled each day.

..... **[2]**

136 **(a)** Calculate the value of

$$\frac{4.2^3 + \sqrt{18.5}}{0.75 \times 1.2}$$

Give your answer correct to 3 significant figures.

..... **[2]**

(b) A spherical water tank has a radius of 1.85 metres.

(i) Calculate the volume of the tank.
Give your answer correct to the nearest cubic metre.
The volume, V, of a sphere with radius r is $V = \frac{4}{3}\pi r^3$.

..... m^3 **[2]**

(ii) The tank is initially empty.
Water is pumped into the tank at a rate of 12.5 litres per minute.
Calculate the time required to fill the tank completely.
Give your answer in hours and minutes, correct to the nearest minute.

hours, minutes **[3]**

137 **(a)** Calculate the value of

$$p = \frac{28.7^2 - 11.2^2}{\sqrt{98.5}}$$

(i) Write down all the figures on your calculator display.

$p =$ ..... **[1]**

(ii) Write your answer to part (a)(i) correct to 3 significant figures.

$p =$ ..... **[1]**

(b) By writing each number in the original expression correct to 1 significant figure, estimate the value of p.

$p \approx$ ..... **[2]**

138 A solid metal cylinder has a base radius of 4.15 cm and a height of 19.8 cm. The metal has a density of 8.13 grams per cubic centimetre.

(a) By writing each number correct to 1 significant figure, estimate the mass of the cylinder in grams.
Use $\pi \approx 3$ for your estimation.
Show your working clearly.

..... g **[3]**

(b) Calculate the actual mass of the cylinder.
Give your answer correct to 3 significant figures.

..... g **[3]**

139 **(a)** Write 45 : 75 : 105 in its simplest form.

..... **[1]**

(b) Alice, Bob and Charlie share \$240 in the ratio of their ages. Alice is 8 years old, Bob is 10 years old and Charlie is 12 years old.
Work out how much Bob receives.

% ..... **[2]**

140 **(a)** Write the ratio 450 g to 1.2 kg in its simplest form.

..... **[2]**

(b) Anna, Ben and Chloe share \$420 in the ratio 3 : 4 : 5.
Calculate the amount Ben receives.

\$ ..... **[2]**

141 Alex, Ben and Charlie share \$840 in the ratio $2 : 3 : 5$.

(a) Calculate the amount of money Ben receives.

\$ ..... **[2]**

(b) Express Charlie's share as a fraction of the total amount, giving your answer in its simplest form.

..... **[2]**

142 Alex, Ben and Chloe share \$120 in the ratio 3 : 4 : 5.

(a) Show that Chloe receives \$50.

[2]

(b) Work out the amount Alex receives.

\$ ..... **[2]**

(c) Write down the ratio of Ben's share to Chloe's share in its simplest form.

..... **[1]**

143 **(a)** Write the ratio 45 minutes to 2 hours in its simplest form.

..... **[2]**

(b) A piece of wood of length 3.6 m is cut into three pieces in the ratio 2 : 3 : 7.
Calculate the length, in cm, of the longest piece.

..... cm **[3]**

144 A sports club has m male members and f female members.
When 30 new male members join the club, the ratio of male members to female members becomes 3 : 1.
When, instead, 20 new female members join the original members, the ratio of male members to female members becomes 1 : 1.

(a) Write down two simultaneous equations in m and f.

.....

..... **[2]**

(b) Solve the equations to find the value of m and the value of f.

$m =$ ..... , $f =$ ..... **[3]**

(c) Find the original ratio of male members to female members in its simplest form.

..... **[1]**

145 A bag contains red balls, blue balls and green balls.
The ratio of red balls to blue balls is 3 : 4.
The ratio of blue balls to green balls is 5 : 2.

(a) Find the ratio of red balls to blue balls to green balls in its simplest form.

..... **[2]**

(b) There is a total of 129 balls in the bag.
Calculate the number of green balls.

..... **[2]**

(c) y blue balls and y green balls are removed from the bag.
The new ratio of blue balls to green balls is 3 : 1.
Work out the value of y.

$y =$ ..... **[3]**

146 A map is drawn to a scale of 1 : 250 000.

(a) The distance between two towns on the map is 8.4 cm. Calculate the actual distance between the two towns in kilometres.

..... km **[3]**

(b) A forest has an actual area of 35 square kilometres.
Calculate the area of the forest on the map in square centimetres.

..... cm^2 **[3]**

147 A map is drawn to a scale of 1 : 250 000.

(a) The distance between two cities on the map is 16.4 cm.
Calculate the actual distance between the two cities in kilometres.

..... km **[2]**

(b) The actual area of a lake is 35 km^2.
Calculate the area of the lake on the map in square centimetres.

..... cm^2 **[3]**

(c) Another map is drawn such that a length of 5 cm represents an actual distance of 20 km.
Write the scale of this map in the form 1 : n.

1 : ..... **[1]**

148 **(a)** A map has a scale of 1 : 50000.
The actual distance between two towns is 14.5 km.
Calculate the distance, in cm, between the two towns on the map.

..... cm **[2]**

(b) An area of a forest on the map is 12.4 cm^2.
Calculate the actual area of the forest in km^2.

..... km^2 **[2]**

(c) A recipe for 8 people requires 240 g of flour and 150 g of butter.
Marco wants to make the recipe for 12 people.
He has 400 g of flour and 200 g of butter.
Determine if Marco has enough ingredients.
Show your working.

.....

..... **[3]**

149 A map is drawn to a scale of 1 : 250 000.

(a) The distance between two cities on the map is 16.8 cm.
Calculate the actual distance between the cities, in km.

..... km **[2]**

(b) A lake has an actual area of 56 km^2.
Calculate the area of the lake on the map, in cm^2.

..... cm^2 **[2]**

(c) On a different map of the same region, the area of the same lake is 14 cm^2.
Find the scale of this new map in the form 1 : n.

1 : ..... **[3]**

150 **(a)** The ratio of adults to children in a theater is 5 : 3.
There are 84 more adults than children.
Calculate the total number of people in the theater.

..... **[3]**

(b) A map is drawn to a scale of 1 : 25000.

(i) Two towns are 8.5 cm apart on the map.
Calculate the actual distance between the towns in kilometres.

..... km **[2]**

(ii) A forest has an actual area of 3.5 km^2.
Calculate the area of the forest on the map in square centimetres.

..... cm^2 **[3]**

151 In a factory, the number of male workers to female workers is in the ratio 5 : 3.
The ratio of male workers who work full-time to male workers who work part-time is 3 : 2.
The ratio of female workers who work full-time to female workers who work part-time is 4 : 1.

(a) Find the ratio of the total number of full-time workers to the total number of part-time workers in the factory.
Give your answer in its simplest form.

..... **[4]**

(b) There are 42 more full-time male workers than part-time female workers.
Calculate the total number of workers in the factory.

..... **[3]**

152 In a factory, the number of workers on the day shift and the number of workers on the night shift are in the ratio 5 : 3.

(a) There are 42 more workers on the day shift than on the night shift.
Calculate the total number of workers in the factory.

..... **[3]**

(b) The ratio of male workers to female workers on the day shift is 3 : 4.
The ratio of male workers to female workers on the night shift is 2 : 7.
Calculate the overall ratio of male workers to female workers in the factory.
Give your answer in its simplest form.

..... **[4]**

153 Three partners, Alex, Ben and Chris, share a total profit of $\$P$. The ratio of Alex's share to Ben's share is 3 : 4. The ratio of Ben's share to Chris's share is 5 : 6.

(a) Find the ratio of Alex's share to Ben's share to Chris's share in its simplest form.

..... **[2]**

(b) Chris receives \$16 200 more than Alex. Calculate the value of P.

\$ ..... **[3]**

(c) Alex invests his share of the profit in a bank account that pays compound interest at a rate of 4.2% per year.
Calculate the value of Alex's investment at the end of 5 years.
Give your answer correct to the nearest dollar.

\$ ..... **[3]**

154 A train departs at 08 : 15 and arrives at its destination at 09 : 45.
The train travels a distance of 120 km.
Calculate the average speed of the train.

..... km/h **[2]**

155 A car travels 120 km in 1 hour and 30 minutes.

Calculate the average speed of the car in km/h.

..... km/h **[3]**

156 **(a)** A car travels a distance of 108 km in 1 hour 30 minutes.
Calculate its average speed in km/h.

..... km/h **[2]**

(b) Change your answer in part (a) into m/s.

..... m/s **[2]**

157 **(a)** A train travels a distance of 120 km in 1 hour and 30 minutes.
Calculate the average speed of the train.

..... km/h **[2]**

(b) A car uses 8 litres of fuel for every 100 km travelled.
Work out how much fuel the car uses to travel 350 km.

..... litres **[2]**

158 **(a)** A train travels a distance of 180 km in 2 hours and 15 minutes.
Calculate the average speed of the train in km/h.

..... km/h **[2]**

(b) An empty tank is filled with water at a constant rate of 1.5 litres per second.
Work out the time it takes to fill the tank with 450 litres of water.
Give your answer in minutes.

..... minutes **[2]**

159 A cyclist travels 30 km at an average speed of v km/h.
He then travels a further 40 km at an average speed of $(v + 5)$ km/h.
The total time taken for the entire journey is 4 hours.

(a) Write down an equation in terms of v and show that it simplifies to $2v^2 - 25v - 75 = 0$.

[3]

(b) Solve the equation $2v^2 - 25v - 75 = 0$ to find the possible values of v.

$v =$ ..... , or $v =$ ..... **[3]**

(c) Find the time taken for the first 30 km of the journey.

..... hours **[1]**

160 A train travels a distance of 385 km.
The journey takes 4 hours 25 minutes.
Calculate the average speed of the train in km/h.
Give your answer correct to 1 decimal place.

..... km/h **[3]**

161 Water flows from a pipe into a rectangular tank at a rate of 45 litres per minute.

(a) Calculate the volume of water, in cubic centimetres (cm^3), that flows into the tank in 1 second.

..... cm^3 **[2]**

(b) The tank has a base measuring 1.5 m by 80 cm. Calculate the rate at which the water level is rising in the tank, giving your answer in cm/s.

..... cm/s **[3]**

162 A car uses fuel at an average rate of 6.4 litres per 100 km.

(a) Calculate the amount of fuel used for a journey of 350 km.

..... litres **[2]**

(b) The cost of fuel is 1.35 euros per litre.
The exchange rate is \$1 = 0.85 euros. Calculate the cost of the fuel for this journey in dollars (\$).
Give your answer correct to the nearest cent.

\$ ..... **[3]**

163 A swimming pool has a volume of 52 m^3.

(a) Water flows into the empty pool from a pipe at a rate of 1.6 litres per second.
Calculate the time taken to completely fill the pool.
Give your answer in hours and minutes, correct to the nearest minute.

hours, minutes **[3]**

(b) The cost of the water is \$2.15 per cubic metre. The exchange rate is \$1 = 0.84 euros.
Calculate the total cost of the water used to fill the pool in euros.
Give your answer correct to 2 decimal places.

euros ..... **[3]**

164 **(a)** A car travels 325 km.
It uses fuel at a rate of 8.4 litres per 100 km.
Fuel costs \$1.35 per litre.
Calculate the total cost of the fuel used for this journey.

\$ ..... **[3]**

(b) On a different journey, the car travels at an average speed of 75 km/h.
The journey takes 4 hours and 24 minutes.
Work out the distance travelled.

..... km **[3]**

165 Water flows from a pipe into a rectangular tank at a rate of 2.5 litres per second.
The tank has a length of 1.8 m, a width of 1.5 m, and a height of 1.2 m.
The tank is initially empty.

(a) Calculate the capacity of the tank in litres.

..... litres **[2]**

(b) Work out how long it takes to fill the tank completely.
Give your answer in minutes.

..... minutes **[2]**

(c) Another pipe empties the full tank at a rate of x litres per minute.
It takes 45 minutes to empty the tank completely.
Calculate the value of x.

$x =$ ..... **[2]**

166 **(a)** Water flows from a pipe into an empty tank at a rate of 0.35 litres per second.
The tank is a cuboid with length 2.5 m, width 1.4 m and height 1.2 m.

Calculate the time taken to fill the tank completely.
Give your answer in hours and minutes.

h ..... , min ..... **[4]**

(b) A mobile phone costs \$850 in the USA and €780 in Germany. The exchange rate is \$1 = €0.84.

Calculate the difference in price between the two countries.
Give your answer in dollars (\$).

\$ ..... **[3]**

167 A car travels from City A to City B, a distance of d km, at an average speed of v km/h.
The car then travels from City B to City C, a distance of $2d$ km, at an average speed of $(v + 15)$ km/h.
The total time taken for the entire journey from City A to City C is T hours.

(a) Show that the total time T can be written as

$$T = \frac{3dv + 15d}{v(v + 15)}$$

[3]

(b) Given that $d = 120$ and $T = 4.5$,

Form an equation in v and show that it simplifies to

$$3v^2 - 195v - 1200 = 0$$

[4]

168 An aeroplane flies from City A to City B, a distance of 2400 km.

(a) The aeroplane flies at an average speed of v km/h.

Write down an expression, in terms of v, for the time taken in hours.

..... hours **[1]**

(b) The aeroplane returns from City B to City A.
The average speed on the return journey is $(v + 40)$ km/h.
The total time taken for the two flights is 10 hours and 30 minutes.
Write down an equation in v and show that it simplifies to

$$7v^2 - 2920v - 192000 = 0$$

[4]

(c) Solve the equation $7v^2 - 2920v - 192000 = 0$ to find the average speed of the aeroplane from City A to City B.
Give your answer correct to the nearest integer.

..... km/h **[2]**

169 A cyclist travels from Town A to Town B, a distance of 48 km.
The cyclist travels the first 30 km at an average speed of x km/h. They travel the remaining distance at an average speed of $(x - 4)$ km/h. The total time taken for the journey is 2 hours and 30 minutes.

(a) Show that $5x^2 - 116x + 240 = 0$.

[4]

(b) Solve the equation $5x^2 - 116x + 240 = 0$ to find the possible values of x.
Show all your working and give your answers correct to 2 decimal places.

$x =$ ..... **[3]**

170 A man runs 15 km at an average speed of x km/h.
He then walks 12 km at an average speed of $(x - 4)$ km/h.
The total time taken for the journey is 3.5 hours.

(a) Show that x satisfies the equation

$$7x^2 - 82x + 120 = 0$$

[3]

(b) **(i)** Solve the equation

$$7x^2 - 82x + 120 = 0$$

$x =$ ..... , or $x =$ ..... **[3]**

(ii) Find the time taken for the man to run the 15 km.

..... hours **[2]**

171 A car travels 150 km at an average speed of x km/h.

(a) Write down an expression, in terms of x, for the time taken in hours.

..... hours **[1]**

(b) The car then travels a further 90 km at an average speed of $(x + 15)$ km/h.
The total time for the whole journey is 3 hours.
Show that $x^2 - 65x - 750 = 0$.

[4]

(c) Solve the equation $x^2 - 65x - 750 = 0$ to find the initial average speed of the car.

$x =$ ..... km/h **[3]**

172 A shopkeeper buys a television for \$320. He sells it for \$400.
Calculate his percentage profit.

..... % **[2]**

173 **(a)** A shop has a sale.
All items are sold with a 15% discount.
Calculate the sale price of a coat originally costing \$80.

\$ ..... **[2]**

(b) A car was bought for \$4000 and sold for \$3200.
Work out the percentage loss.

..... % **[2]**

174 **(a)** Write 24 as a percentage of 60.

..... % **[2]**

(b) A coat has a normal price of \$80. In a sale, the price is reduced by 15%.
Work out the sale price of the coat.

\$ ..... **[2]**

175 **(a)** Find 15% of \$320.

\$ ..... **[2]**

(b) In a sale, the original price of a shirt is reduced by 20%.
The original price of the shirt is \$45.
Work out the sale price.

\$ ..... **[3]**

176 A shop sells a jacket for \$120. During a sale, the shop offers a 15% discount on the price of the jacket.

(a) Calculate the sale price of the jacket.

\$ ..... **[2]**

(b) The original price of \$120 gives the shop a profit of 20% on the cost price.
Calculate the cost price of the jacket.

\$ ..... **[3]**

177 **(a)** A shop reduces its prices by 20% in a sale.
The original price of a book is \$15.
Calculate the sale price of the book.

\$ ..... **[2]**

(b) The sale price of a jacket in the same shop is \$56.
Calculate the original price of the jacket.

\$ ..... **[3]**

(c) A different shop buys a watch for \$30 and sells it for \$42.
Calculate the percentage profit.

..... % **[2]**

178 **(a)** Alice invests \$4500 at a rate of 3.2% per year compound interest.
Calculate the total value of her investment at the end of 5 years.

\$ ..... **[3]**

(b) A shopkeeper buys a television for \$320. He sells it for \$408.
Calculate his percentage profit.

..... % **[2]**

179 **(a)** In 2021, the population of a town was 45 000.
In 2022, the population increased by 4.2%.
Calculate the population in 2022.

..... **[2]**

(b) In a sale, the price of a television is reduced by 24%.
The sale price is \$342.
Calculate the original price of the television.

\$ ..... **[3]**

180 Amira invests \$4500 in a savings account that pays compound interest at a rate of 2.4% per year.

(a) Calculate the total amount in Amira's account at the end of 5 years.

\$ ..... **[3]**

(b) Ben invests \$4500 in a different savings account that pays simple interest at a rate of r% per year. At the end of 5 years, Ben has the same total amount in his account as Amira has in hers.

Calculate the value of r.

$r =$ ..... **[3]**

181 **(a)** Anja invests \$8000 in an account paying compound interest at a rate of 3.5% per year.

Calculate the total value of her investment at the end of 4 years.
Give your answer correct to the nearest cent.

\$ ..... **[3]**

(b) The population of a town decreases by 4% each year.
In 2026, the population is 45 000.
Calculate the population of the town in 2023.
Give your answer correct to the nearest integer.

..... **[3]**

182 **(a)** Amira invests \$4500 in a savings account paying compound interest at a rate of 2.4% per year.
Calculate the total value of her investment at the end of 5 years.

\$ ..... **[3]**

(b) In a sale, the price of a computer is reduced by 18%.
The sale price is \$557.60.
Calculate the original price of the computer.

\$ ..... **[3]**

183 **(a)** An investment of \$8000 earns compound interest at a rate of 3.5% per year. Calculate the value of the investment at the end of 4 years.
Give your answer correct to the nearest cent.

\$ ..... **[3]**

(b) A car is bought for \$24000. Its value decreases by 12% each year. Calculate the number of complete years it takes for the value of the car to fall below \$10000.

..... years **[3]**

(c) In 2022, the price of a house was \$150000. This was a 25% increase on its price in 2017. Calculate the price of the house in 2017.

\$ ..... **[3]**

184 **(a)** The population of a city increases by 4.5% each year.
The population is now 1.25×10^6.
Calculate the expected population in 10 years' time.
Give your answer in standard form, correct to 3 significant figures.

..... **[3]**

(b) In a sale, a washing machine is sold for \$299.25. This is a 14% reduction on the original price.
Calculate the original price.

\$ ..... **[2]**

(c) A car depreciates in value by 15% in the first year and by 10% in each subsequent year.
The value of the car after 3 years is \$17212.50.
Calculate the original value of the car.

\$ ..... **[2]**

185 The population of a city increases by $x\%$ each year.
In 2015, the population was 1 250 000.
In 2018, the population was 1 367 631.

(a) Show that the value of x is approximately 3.

[3]

(b) Assume the population continues to increase by exactly 3% each year.

Calculate the year in which the population will first exceed double its 2015 value.

..... **[4]**

186 **(a)** A shopkeeper buys a television for $\$p$. He sells it for \$960, making a percentage profit of 20% on his cost price.
Calculate the value of p.

$p = \$ \ldots\ldots\ldots\ldots\ldots$ **[2]**

(b) The shopkeeper buys another television for $\$q$. He wants to make a 15% profit on his cost price after allowing a 10% discount on the marked price.

Write down an expression, in its simplest form, for the marked price in terms of q.

$\$ \ldots\ldots\ldots\ldots\ldots\ldots\ldots\ldots\ldots\ldots\ldots\ldots\ldots\ldots\ldots\ldots\ldots\ldots$ **[3]**

(c) An investment of \$500 earns compound interest at a rate of r% per year. At the end of 10 years, the value of the investment is \$740.12.
Work out the value of r.

$r = \ldots\ldots\ldots\ldots\ldots$ **[3]**

187 **(a)** Ali invests \$5000 at a rate of 2.5% per year compound interest. Calculate the value of his investment at the end of 3 years.
Give your answer correct to the nearest cent.

\$ ..... **[2]**

(b) Beatrice invests a sum of money at a rate of r% per year compound interest.
After 5 years, the value of her investment has increased by 30%.
Calculate the value of r.

$r =$ ..... **[3]**

(c) The value of a car depreciates by 12% each year.
After 3 years, the value of the car is \$17 036.80.
Calculate the original value of the car.

\$ ..... **[3]**

188 **(a)** A car's value depreciates by 15% in the first year.
In each subsequent year, its value depreciates by 8%.
After exactly 3 years, the value of the car is \$17 986.
Calculate the original value of the car.

\$ ..... **[4]**

(b) Calculate the overall percentage decrease in the car's value over the 3 years.
Give your answer correct to 1 decimal place.

..... % **[2]**

(c) The population of a city is P.
The population increases by r% each year.
After 12 years, the population has exactly doubled.
Calculate the value of r.

$r =$ ..... **[3]**

189 **(a)** A population of bacteria increases by $r\%$ every hour.
The population doubles in 6 hours.
Calculate the value of r, giving your answer correct to 3 significant figures.

$r =$ ..... **[3]**

(b) A merchant buys goods for $\$X$. He sets the selling price such that he makes a 40% profit on the cost price. During a sale, he gives a 15% discount on this selling price. The final sale price is \$142.80.
Calculate the value of X.

$X =$ ..... **[3]**

(c) An amount of $\$P$ is invested at a rate of $x\%$ per year simple interest. The total simple interest earned in 5 years is \$250.
The same amount $\$P$ is invested at a rate of $x\%$ per year compound interest. The total compound interest earned in 2 years is \$102.50.
Find the value of P and the value of x.

$P = \$$ ..... , x = ... **[5]**

190 Calculate the value of

$$\frac{\sqrt{15.2 - 3.4^2}}{4.1 \times 0.25}$$

Write down your full calculator display.

..... **[2]**

191 Work out the value of

$$\frac{\sqrt[3]{25.6 - 3.2^2}}{1.8 \times 0.45}$$

(a) Write down all the figures on your calculator display.

..... **[2]**

(b) Write your answer to part (a) correct to 3 decimal places.

..... **[1]**

192 Consider the following calculation.

(a) Work out the value of

$$\frac{\sqrt{4.1^2 + 5.2}}{2.3^3 - 1.4}$$

Write down all the figures on your calculator display.

..... **[2]**

(b) Give your answer to part (a) correct to 3 significant figures.

..... **[1]**

193 **(a)** Work out the value of

$$\frac{\sqrt{15.2^2 - 8.5^2}}{3.2 \times 1.4}$$

Write down your full calculator display.

..... **[2]**

(b) Write your answer to part (a) correct to 3 significant figures.

..... **[1]**

(c) Calculate

$$\frac{4.5 \times 10^5}{1.2 \times 10^{-3}}$$

Give your answer in standard form.

..... **[1]**

194 **(a)** Work out the value of

$$\frac{4.5^2 - \sqrt{18.2}}{3.1 + 0.8^3}$$

..... **[2]**

(b) Calculate the value of

$$2\sin(48°) - \cos(35°)$$

Give your answer correct to 3 decimal places.

..... **[2]**

195 **(a)** A shopkeeper calculates the cost of an item using the expression

$$\frac{17.3 + 8.14}{5.3}$$

Work out the cost.
Give your answer as a money value in dollars, formatted correctly.

$ ..... **[2]**

(b) A movie lasts for 2 hours and 18 minutes.
Write this time as a decimal number of hours.

..... hours **[2]**

196 **(a)** Use your calculator to find the value of

$$3.5^4 + \sqrt[3]{87.6}$$

Give your answer correct to 2 decimal places.

..... **[2]**

(b) Calculate the value of x when

$$x = \frac{2 \times \sin(43°)}{\cos(18°)}$$

Give your answer correct to 3 significant figures.

$x =$ ..... **[2]**

197 A student uses a calculator to solve three different problems.

(a) The answer to a problem about money is displayed on the calculator as 8.4.
Write this amount appropriately in dollars and cents.

..... **[1]**

(b) The answer to a problem about time is displayed on the calculator as 5.15 hours.
Write this time appropriately in hours and minutes.

hours ..... , minutes ..... **[1]**

(c) A car travels 280 km at an average speed of 85 km/h. Calculate the time taken for this journey.
Give your answer in hours and minutes, correct to the nearest minute.

hours ..... , minutes ..... **[3]**

198 **(a)** Calculate the value of

$$\frac{3.5 + 1.2^2}{\sqrt{4.1 - 2.8}}$$

(i) Write down all the figures on your calculator display.

..... **[2]**

(ii) Write your answer to part (a)(i) correct to 3 significant figures.

..... **[1]**

(b) A calculator displays a journey time as 5.35 hours.
Convert this time to hours and minutes.

h ..... , min ..... **[2]**

199 **(a)** A circular swimming pool has a radius of 4.2 metres.
Calculate the area of the pool.
Give your answer correct to the nearest square metre.

..... m^2 **[2]**

(b) A train journey takes 4 hours and 23 minutes.
The average speed of the train is 78 km/h.

Calculate the distance travelled by the train.
Give your answer correct to 1 decimal place.

..... km **[4]**

200 A plumber charges a fixed fee of \$45 plus an hourly rate of \$32.50.

(a) The plumber works on a job for 3 hours and 40 minutes.
Calculate the total cost of this job.
Give your answer to the nearest cent.

\$ ..... **[3]**

(b) On another job, the total cost charged by the plumber is \$230.25.
Calculate the time the plumber worked, giving your answer in hours and minutes.

hours, minutes **[3]**

201 A metal sphere has a radius of 4.85 cm.

(a) Calculate the volume of the sphere.

The volume, V, of a sphere with radius r is $V = \frac{4}{3}\pi r^3$.

Give your answer correct to 1 decimal place.

..... cm^3 **[2]**

(b) The metal sphere is melted down and made into identical small cones.

Each cone has a radius of 1.2 cm and a height of 3.5 cm.

The volume, V, of a cone with radius r and height h is $V = \frac{1}{3}\pi r^2 h$.

Calculate the maximum number of full cones that can be made.

..... **[4]**

(c) Calculate the percentage of metal wasted from the original sphere.

..... % **[1]**

202 A water tank is in the shape of a sphere with radius 1.4 m.
The tank is completely full of water.

(a) Work out the volume of water in the tank.
Write down all the figures on your calculator display.

..... m^3 **[2]**

(b) Water leaks from the tank at a constant rate of 0.035 cubic metres per minute.
Calculate the time taken for the tank to become completely empty.
Do not round any intermediate values during your working.
Give your answer in hours and minutes, correct to the nearest minute.

hours ..... , minutes ..... **[4]**

203 **(a)** Calculate the exact value of

$$\left(3.2 \times 10^{5}\right) \times \left(4.5 \times 10^{-2}\right)$$

Give your answer in standard form.

..... **[2]**

(b) Work out the value of

$$\frac{1}{\sqrt{2\pi}} \times e^{-0.5^2}$$

Give your answer correct to 4 decimal places.

..... **[2]**

(c) A formula is given by $T = 2\pi\sqrt{\frac{l}{g}}$.

Find the value of l when $T = 1.5$ and $g = 9.81$.

Give your answer correct to 2 decimal places.

$l =$ ..... **[2]**

204 **(a)** Calculate

$$\frac{3.4 \times 10^5 + 2.7 \times 10^4}{8.1 \times 10^{-3}}$$

Give your answer in standard form.

..... **[3]**

(b) A solid metal cone has a base radius of r cm and a height of h cm.
$r = 5.2$ correct to 1 decimal place.
$h = 12.4$ correct to 1 decimal place.
Calculate the upper bound for the volume of the cone.
The volume, V, of a cone with radius r and height h is $V = \frac{1}{3}\pi r^2 h$.

..... cm^3 **[3]**

205 The formula for the period, T seconds, of a simple pendulum is given by

$$T = 2\pi\sqrt{\frac{l}{g}}$$

where l is the length of the pendulum in metres and g is the acceleration due to gravity, 9.81 m/s^2.

(a) A pendulum has a length of 85 cm.
Calculate the period T.
Give your answer correct to 3 decimal places.

$T =$ ..... seconds **[2]**

(b) Rearrange the formula to make l the subject.

$l =$ ..... **[2]**

(c) A different pendulum has a period of exactly 2.5 seconds.
Calculate the length of this pendulum in centimetres.
Give your answer correct to 1 decimal place.

..... cm **[3]**

206 The volume, V, of a spherical cap is given by the formula

$$V = \frac{1}{3}\pi h^2(3R - h)$$

where R is the radius of the sphere and h is the height of the cap.

(a) Calculate V when $R = 8.45$ cm and $h = 3.2$ cm. Give your answer correct to 1 decimal place.

$V =$ ..... cm^3 **[2]**

(b) Rearrange the formula to make R the subject.

$R =$ ..... **[3]**

(c) A different spherical cap has volume $V = 500$ cm^3 and height $h = 5.6$ cm. Calculate the radius, R, of this sphere.
Give your answer correct to 3 significant figures.

$R =$ ..... cm **[2]**

207 The volume, V, of a spherical cap is given by the formula

$$V = \frac{1}{3}\pi h^2(3R - h)$$

(a) Calculate the volume V when $R = 8.74$ and $h = 2.31$.
Give your answer correct to 1 decimal place.

$V =$ ..... **[3]**

(b) A student calculates V by first rounding the values of R and h to 1 decimal place before substituting them into the formula.
Calculate the difference between this incorrect answer and the exact volume.
Give your difference correct to 2 significant figures.

Difference = ... **[4]**

208 A train journey from Portsmouth to London is shown in the timetable.

Station	Time
Portsmouth	08 45
London	11 20

(a) Work out the journey time in hours and minutes.

hours ..... , minutes ..... **[1]**

(b) The return journey takes exactly the same amount of time.
The train leaves London at 3.35 p.m.

Write down the time it arrives in Portsmouth using the 24-hour clock.

..... **[2]**

209 A train timetable shows that a train leaves Paris at 09 55 and arrives in Lyon at 12 10.

(a) Work out the time taken for this journey in hours and minutes.

hours ..... , minutes ..... **[1]**

(b) Another train leaves Paris at 16 45.
This journey takes 2 hours and 35 minutes.
Work out the time this train arrives in Lyon.

..... **[2]**

210 A flight departs from London to Dubai at 21 : 45 local time.

The flight time is 6 hours and 50 minutes.
The local time in Dubai is 4 hours ahead of the local time in London.
Work out the local time in Dubai when the flight arrives.

..... **[3]**

211 A flight leaves Dubai at 14 30 local time on Friday and arrives in London at 19 15 local time on the same day.
The local time in Dubai is 4 hours ahead of the local time in London.

(a) Work out the duration of the flight in hours and minutes.

hours, minutes **[2]**

(b) A second flight leaves Dubai at 22 45 local time on Friday.
The duration of this flight is 7 hours and 20 minutes.
Work out the local time and the day it arrives in London.

Time = ... , Day = ... **[2]**

212 **(a)** A train departs from Paris at 22:50.
The journey to Rome takes 14 hours and 35 minutes.
Work out the arrival time in Rome using the 24-hour clock.

..... **[2]**

(b) Convert 3.4 hours into hours and minutes.

h ..... , min ..... **[2]**

213 **(a)** A movie starts at 19 35 and lasts for 115 minutes.
Work out the time the movie ends in the 24-hour clock.

..... **[2]**

(b) A plane leaves New York at 21 40 local time.
The flight to Paris takes 7 hours and 15 minutes.
The local time in Paris is 6 hours ahead of New York.
Find the local time in Paris when the plane arrives.

..... **[3]**

214 A flight departs from Dubai at 14 30 local time and arrives in Sydney.
The flight time is 14 hours and 15 minutes.
The local time in Sydney is 7 hours ahead of the local time in Dubai.
Work out the local time in Sydney when the flight arrives.

..... **[3]**

215 Light travels at 3×10^5 km/s.

(a) Calculate the distance light travels in one year (365 days).
Give your answer in standard form.

..... km **[2]**

(b) The distance from Earth to a star is 4.5×10^{13} km.
Calculate the time taken for light to travel from the star to Earth.
Give your answer in years, correct to 3 significant figures.

..... years **[2]**

216 A train travels from Station A to Station B. The total distance is 420 km.
The train departs Station A at 08 : 15 and arrives at Station B at 11 : 03.

(a) Calculate the time taken for the journey, in hours and minutes.

hours, minutes **[2]**

(b) Calculate the average speed of the train in km/h.

..... km/h **[3]**

217 Dubai is in the time zone UTC+4 and London is in the time zone UTC+0.

(a) An airplane departs Dubai at 02 15 local time on Tuesday and flies to London.
The flight takes 7 hours and 45 minutes.

Find the local time and day in London when the flight arrives.

Time ..... , Day ..... **[3]**

(b) An astronaut spends exactly 1.5 years in space.
Given that 1 year = 365 days, calculate the total number of minutes the astronaut spends in space.

Give your answer in standard form.

..... minutes **[3]**

218 **(a)** A flight from Tokyo to London takes 14 hours and 15 minutes.
Tokyo is 9 hours ahead of London.
The flight arrives in London at 16:30 local time on Tuesday.
Calculate the departure time from Tokyo in local time.

..... **[3]**

(b) A machine completes one cycle every 1 minute and 45 seconds.
Calculate the maximum number of full cycles it can complete in 8 hours.

..... **[3]**

219 **(a)** A marathon runner completes a 42.2 km race at an average speed of 12.5 km/h.

Work out the time taken to complete the race.
Give your answer in hours, minutes and seconds, correct to the nearest second.

hours = ... , minutes = ... , seconds = ... **[4]**

(b) Another runner finishes the race at 13 04.
She started the race at 08 35.
Work out how long she took to complete the race, in hours and minutes.

hours = ... , minutes = ... **[2]**

220 A communications satellite orbits the Earth.
It completes one full orbit every 94 minutes and 30 seconds.

(a) Work out the number of full orbits the satellite completes in a 30-day month.

..... **[3]**

(b) The satellite passes directly over an observatory at 05 50 on 1st March.
Find the time and date when it completes its 100th orbit after this point.

Time ..... , Date ..... **[4]**

221 A ship travels a distance of 920 km at an average speed of 24 km/h.

(a) Work out the time taken for the journey.
Give your answer in hours and minutes.

hours ..... , minutes ..... **[3]**

(b) The ship left the port on Wednesday at 17 50.
Determine the day and the time it arrives at its destination.

Day ..... , Time ..... **[4]**

222 An aircraft flies from City A to City B, a distance of 6600 km.
The local time at City B is 5 hours behind the local time at City A.
The aircraft leaves City A at 21 40 local time on Tuesday.
The average speed of the aircraft is 800 km/h.

(a) Calculate the time of the flight in hours and minutes.

hours, minutes **[2]**

(b) Find the local time and the day when the aircraft arrives at City B.

Time = ... , Day = ... **[2]**

(c) The aircraft stays at City B for 14 hours and 20 minutes before flying back to City A.
The return flight takes 7 hours and 45 minutes.
Find the local time and the day when the aircraft arrives back at City A.

Time = ... , Day = ... **[3]**

223 A cyclist travels 35 km at an average speed of x km/h.
She then travels a further 42 km at an average speed of $(x - 5)$ km/h.

(a) Write down an expression, in terms of x, for the total time taken in hours.

..... **[1]**

(b) The total time for the journey is 4 hours and 33 minutes.
Show that $13x^2 - 285x + 500 = 0$.

[4]

(c) Solve the equation to find the value of x.

$x =$ ..... **[3]**

224 Consider the following applications of time, speed, and dates.

(a) A satellite completes one orbit of the Earth in exactly 92 minutes.
Calculate the number of full orbits it completes in exactly one non-leap year (365 days).

..... orbits **[4]**

(b) A plane flies from City P to City Q. The distance is 11 440 km and the average speed is 880 km/h.

(i) Calculate the flight time.

..... hours **[1]**

(ii) City P is in the time zone UTC−4.
City Q is in the time zone UTC+9.
The plane departs from City P on Monday at 22 : 45 local time.
Find the local day and time of arrival in City Q.

Day = ... , Time = ... **[3]**

225 **(a)** A satellite takes 92 minutes and 45 seconds to complete one orbit of the Earth.
Work out the total time taken to complete 15 orbits.
Give your answer in hours, minutes and seconds.

h ..... , min ..... , s ..... **[2]**

(b) A flight departs City A at 09:45 local time.
The flight time is 8 hours and 20 minutes.
It arrives in City B at 02:05 local time the next day.
City B is h hours ahead of City A.

Find the value of h.

$h =$ ..... **[3]**

(c) Train P leaves Station X at 08:00 and travels to Station Y at a constant speed of 90 km/h.
Train Q leaves Station Y at 08:30 and travels to Station X at a constant speed of 120 km/h.
The distance between X and Y is 360 km.
Work out the exact time when the two trains pass each other.

..... **[4]**

226 Hassan travels from Paris to Dubai.
He changes €400 into United Arab Emirates Dirhams (AED).
The exchange rate is €1 = 4.20 AED.

(a) Calculate how many AED he receives.

..... AED **[1]**

(b) Hassan spends 1050 AED in Dubai.
He changes his remaining AED back into euros.
The new exchange rate is €1 = 4.50 AED.

Calculate how many euros he receives.

€ ..... **[2]**

227 John has \$400. He buys a new camera that costs 255 Euros (). The exchange rate is \$1 = 0.85.
Work out how much money John has left in dollars.

\$ ..... **[3]**

228 Aisha buys 3 pens costing \$1.20 each and 4 notebooks costing \$2.50 each.
She pays with a \$20 note.
Work out how much change she receives.

% ..... **[3]**

229 The exchange rate between pounds (£) and dollars ($) is £1 = $1.50.
Anna changes £300 into dollars.
She spends $210 on a tablet.
She changes the remaining dollars back into pounds at the same exchange rate.

Work out how many pounds she receives.

£ ..... **[3]**

230 **(a)** Adele buys 4 pens costing \$2.35 each and 3 notebooks costing \$4.50 each.
She pays with a \$50 note.
Work out how much change she receives.

% ..... **[3]**

(b) The exchange rate between euros () and US dollars (\$) is 1 = \$1.20.
Change 45 into US dollars.

% ..... **[1]**

231 Amira goes on holiday to Japan.
She changes \$850 into Japanese Yen (JPY). The exchange rate is \$1 = 148 JPY.

(a) Calculate how many JPY she receives.

..... JPY **[2]**

(b) In Japan, she spends 45 000 JPY on a camera and 31 500 JPY on food.
At the end of the holiday, she changes her remaining JPY back to dollars (\$). The new exchange rate is \$1 = 152 JPY.
Calculate how many dollars she receives.
Give your answer correct to the nearest cent.

\$ ..... **[3]**

232 A car dealer buys a car in Japan for 1 250 000 JPY.
The exchange rate is £1 = 145 JPY.

(a) Calculate the cost of the car in pounds (£).

£ ..... **[2]**

(b) The dealer pays shipping costs of £600.
He then sells the car in the UK for £12 500.

Calculate his percentage profit on the total cost of buying and shipping the car.

..... % **[3]**

233 **(a)** The exchange rate between euros () and US dollars ($) is 1 = $1.14.
Convert 450 into US dollars.

% ..... **[2]**

(b) The exchange rate between British pounds (£) and US dollars ($) is £1 = $1.32.
Calculate the exchange rate between British pounds and euros.
Give your answer in the form £1 = x, correct to 2 decimal places.

£1 = ... **[3]**

234 A shop sells laptops in London and in New York.
The same model of laptop costs £680 in London and $850 in New York.

(a) The exchange rate is £1 = $1.32.

Calculate the difference in the cost of the laptop in London and the laptop in New York. Give your answer in dollars ($).

$ ..... **[3]**

(b) A customer buys the laptop in New York for $850. They must also pay a local tax of 4% on the cost of the laptop. The customer pays the total amount in euros (€). The exchange rate is $1 = €0.85.

Calculate the total amount the customer pays in euros.

€ ..... **[3]**

235 **(a)** Tomas changes £450 into euros ().
The exchange rate is £1 = 1.18.
He spends 285 on his holiday.
He changes the remaining money back into pounds.
The new exchange rate is £1 = 1.22.
Calculate the amount of pounds he receives.
Give your answer correct to the nearest penny.

£ ..... **[4]**

(b) A television is priced at $850. In a sale, the price is reduced by 15%. A customer has a voucher that gives a further 5% discount off the sale price.
Calculate the final price paid for the television.

% ..... **[2]**

236 Amira travels from the UK to Japan.
She changes 850 into Japanese Yen (JPY).
The bank charges a 2% fee on the amount exchanged.
The exchange rate is 1 = 180 JPY.

(a) Calculate the amount of JPY she receives.

..... JPY **[3]**

(b) While in Japan, she spends 115000 JPY.
She changes her remaining JPY back to pounds ().
The new exchange rate is 1 = 184 JPY.
The bank charges a fixed fee of 500 JPY for this transaction.
Calculate the amount of pounds she receives.

..... **[4]**

237 Carlos buys a car in the USA for \$17 400. He pays a shipping fee of \$1 200.

(a) The exchange rate is £1 = \$1.24. Calculate the total cost of the car and shipping in pounds (£).

£ ..... **[2]**

(b) Carlos sells the car in the UK for £17 100.
Calculate his percentage profit on his total cost.

..... % **[3]**

(c) Carlos changes his profit of £2 100 into euros.
The exchange rate is £1 = 1.16 euros.
The bank charges a 2% commission fee for the exchange.
Calculate the amount he receives in euros.

..... euros **[3]**

238 A trader buys 1000 items for y each.
He pays a shipping fee of 250.
He sells all 1000 items in the USA for \$4.20 each.

The exchange rate is 1 = \$1.20.
The trader makes a profit of 40% on his total cost in Euros.

Calculate the value of y.

$y =$ ..... **[5]**

239 **(a)** Carlos changes \$800 into euros () when the exchange rate is \$1 = 0.85.
He spends 420 on holiday.
He changes the remaining euros back into dollars when the exchange rate is \$1 = 0.82.
Calculate the amount of dollars he receives.
Give your answer correct to the nearest cent.

% ..... **[4]**

(b) In a sale, a shop reduces all prices by 15%.
The sale price of a jacket is \$61.20.
Calculate the original price of the jacket.

% ..... **[2]**

240 A shop sells two brands of coffee, Brand A and Brand B.
Brand A costs \$12.50 per 500 grams. Brand B costs \$8.40 per 300 grams.

(a) Which brand is better value for money?
You must show all your working.

..... **[3]**

(b) The shop owner mixes x grams of Brand A with y grams of Brand B to make a new blend.
She makes 2 kg of the new blend.
The total cost of the coffee in the new blend is \$54.10.

Work out the value of x and the value of y.

$x =$ ..... , $y =$ ..... **[4]**

241 In Year 1, the exchange rate between dollars ($) and euros (EUR) was $\$1 = x$ EUR.
In Year 2, the exchange rate changed to $\$1 = (x - 0.05)$ EUR.

In Year 1, a laptop cost 400 EUR. In Year 2, the same model laptop cost 450 EUR.

The cost of the laptop in dollars increased by $100 from Year 1 to Year 2.

(a) Write down an expression in terms of x for the cost of the laptop in dollars in Year 1.

$\$ \ldots\ldots\ldots\ldots\ldots\ldots$ **[1]**

(b) Form an equation in x and show that it simplifies to $20x^2 - 11x - 4 = 0$.

[4]

(c) Solve the equation to find the value of x.

$x = \ldots\ldots\ldots\ldots\ldots\ldots$ **[3]**

242 **(a)** Car A costs \$24 000 in the USA. Car B costs €21 500 in Germany. The exchange rate is \$1 = €x.
The difference in the cost of the two cars is exactly €740.
Car B is more expensive than Car A.

Find the value of x.

$x =$ ..... **[3]**

(b) A customer changes a sum of dollars (\$) into British pounds (£). The exchange rate is £1 = \$1.28.
The bank deducts a 2.5% commission from the initial sum in dollars.
The remaining dollars are then changed into pounds.
The customer receives £780.

Calculate the initial sum of dollars the customer wanted to change.

\$ ..... **[3]**

(c) A company earns revenue in dollars (\$) which is then converted into euros (€). In 2022, the exchange rate was \$1 = €0.828.
In 2023, the company's revenue in dollars decreased by 8%.
However, the company's revenue in euros was exactly the same in 2023 as it was in 2022.

Calculate the exchange rate in 2023, giving your answer in the form \$1 = €k.

$k =$ ..... **[3]**

243 A shop sells laptops.
In 2022, the price of a laptop was \$840. In 2023, the price increased by 15%. In 2024, the price decreased by r% from the 2023 price. The price of the laptop in 2024 was \$821.10.

(a) Calculate the value of r.

$r =$ ..... **[3]**

(b) The shop buys 450 of these laptops from a supplier in China.
The total cost is 2 430 000 Chinese Yuan (CNY).
The exchange rate is \$1 = 7.2 CNY.

Calculate the profit the shop makes on each laptop when sold at the 2022 price of \$840.

\$ ..... **[3]**

(c) A customer in the UK buys one of these laptops in 2022 for \$840. They pay with a credit card that charges a 2.5% foreign transaction fee. The exchange rate is £1 = \$1.32.
Calculate the total amount the customer pays in pounds (£).
Give your answer correct to the nearest penny.

£ ..... **[3]**

244 A population of bacteria increases exponentially at a constant rate of $p\%$ per hour.
At 09 : 00, the population is 8 000.
At 12 : 00 on the same day, the population is 27 000.

(a) **(a)** Work out the value of p.

$p =$ ..... **[3]**

(b) Work out the population of the bacteria at 14 : 00 on the same day.

..... **[2]**

245 **(a)** The value of a painting increases exponentially by $r\%$ each year.
After 2 years, the total percentage increase in the value of the painting is 44%.
Work out the value of r.

$r =$ ..... **[3]**

(b) The value of a rare stamp increases exponentially.
Its value doubles every 15 years.
The current value of the stamp is \$80.

Work out the value of the stamp after 45 years.

% ..... **[2]**

246 A new car is bought for \$24 000. The value of the car depreciates exponentially at a rate of 8% per year.

Calculate the value of the car after 5 years.
Give your answer correct to the nearest dollar.

\$ ..... **[3]**

247 The population of a city is 125 000.
The population increases exponentially at a rate of 1.8% per year.
Calculate the population of the city after 8 years.
Give your answer correct to the nearest integer.

..... **[3]**

248 A new car is bought for \$26000. The value of the car depreciates exponentially at a rate of 15% each year.

Work out the value of the car after 4 years.
Give your answer correct to the nearest dollar.

% ..... **[3]**

249 A rare stamp is bought for \$450. Its value increases exponentially at a rate of 6.5% per year. Calculate the value of the stamp after 12 years.

\$ ..... **[3]**

250 A machine is bought for \$32 000. It depreciates exponentially at a rate of 12% each year.

Calculate the value of the machine after 5 years.
Give your answer correct to the nearest dollar.

\$ ..... **[3]**

251 The population of a town is 65000.
The population increases exponentially at a rate of 2.8% per year.

(a) Calculate the expected population of the town after 5 years.
Give your answer correct to the nearest integer.

..... **[2]**

(b) Calculate the overall percentage increase in the population over the 5 years.
Give your answer correct to 1 decimal place.

..... % **[2]**

252 A population of bacteria in a dish grows exponentially.

(a) Initially, there are 1500 bacteria in the dish.
The population increases by 8% every hour.
Work out the population of bacteria after 12 hours.
Give your answer correct to the nearest whole number.

..... **[2]**

(b) Work out the least number of complete hours it will take for the initial population of 1500 bacteria to be greater than 3000.

..... hours **[3]**

253 The population of a town was 85 000 on 1 January 2010.
The population increases exponentially at a constant rate of $r\%$ per year.
On 1 January 2022, the population was 112 000.

(a) **(a)** Calculate the value of r, correct to 3 significant figures.

$r =$ ..... **[3]**

(b) Assuming the rate of increase remains the same, calculate the expected population on 1 January 2030.
Give your answer correct to the nearest thousand.

..... **[2]**

254 A car is purchased for \$28 000. Its value depreciates exponentially at a rate of r% each year. After 4 years, its value is \$16 150.

(a) Calculate the value of r.
Give your answer correct to 3 significant figures.

$r =$ ..... **[3]**

(b) The value of the car continues to depreciate at the same rate.
Calculate the value of the car after a further 3 years.
Give your answer correct to the nearest dollar.

\$ ..... **[2]**

255 A machine costs \$45000. Its value depreciates exponentially at a rate of $x\%$ per year. After 4 years, the value of the machine is \$26000.

(a) Calculate the value of x.
Give your answer correct to 3 significant figures.

$x =$ ..... **[3]**

(b) The machine will be replaced when its value first falls below \$10000.
Find the number of complete years before the machine is replaced.

..... years **[3]**

256 **(a)** The population of a city is 125 000.
It grows exponentially at a rate of 2.4% per year.

Calculate the expected population after 10 years.
Give your answer correct to the nearest hundred.

..... **[3]**

(b) The population of another city was 85 000 in the year 2015.
By 2023, the population had decreased exponentially to 68 000.

Calculate the annual percentage decrease.

..... % **[3]**

257 Consider the depreciation of two different vehicles over time.

(a) A car is purchased for \$28 500. Its value depreciates exponentially at a rate of 12% per year.
Calculate the value of the car after 5 years, giving your answer correct to the nearest dollar.

\$ ..... **[3]**

(b) A van, initially purchased for \$35 000, depreciates exponentially to a value of \$15 200 after 4 years.
Calculate the annual percentage rate of depreciation for this van.

..... % **[4]**

258 The mass, M grams, of a radioactive substance decays exponentially at a rate of $r\%$ per day.
After 2 days, the mass is 80 grams.
After 5 days, the mass is 40.96 grams.

(a) Calculate the value of r.

$r =$ ..... **[3]**

(b) Find the initial mass of the substance.

..... grams **[2]**

(c) Calculate the number of complete days from the start when the mass is first less than 10 grams.

..... days **[3]**

259 A radioactive substance decays exponentially.

(a) The mass of the substance decreases by 12% every day.
The initial mass is 500 grams.

Calculate the mass remaining after 1 week (7 days).

..... g **[2]**

(b) After d complete days, the mass of the substance falls below 50 grams.

Find the smallest integer value of d.

$d =$ ..... **[3]**

(c) A different radioactive substance decays exponentially such that its mass halves every 15 days.

Calculate the percentage decrease in its mass each day.

..... % **[3]**

260 Alex and Ben each invest \$4000 for 10 years. Alex invests his \$4000 in an account paying compound interest at a rate of 4.5% per year.
Ben invests his \$4000 in an account paying compound interest at a rate of r% per year.

(a) Calculate the total amount in Alex's account after 10 years.

\$ ..... **[2]**

(b) After 10 years, Ben has \$500 more than Alex.
Calculate the value of r.

$r =$ ..... **[4]**

(c) A different investment of \$2500 grows exponentially. It doubles in value every 8 years.
Calculate its value after 20 years.

\$ ..... **[2]**

261 The population of a species of insect increases exponentially.
Initially, there are 2500 insects.
After 6 weeks, there are 9100 insects.

(a) Calculate the percentage increase in the population per week.

..... % **[4]**

(b) Calculate the insect population after 10 weeks, giving your answer correct to the nearest 100.

..... **[2]**

(c) Find the number of complete weeks it takes for the initial population to increase to over 50 000.

..... weeks **[3]**

262 Simplify

$$\sqrt{75} - \sqrt{27} + \sqrt{12}$$

..... **[2]**

263 Simplify.

$\sqrt{75} + \sqrt{48} - \sqrt{27}$

..... **[2]**

264 Simplify fully.

$\sqrt{72} + \sqrt{50} - \sqrt{18}$

..... **[2]**

265 Simplify

$$\sqrt{75} - \sqrt{27} + \sqrt{12}$$

..... **[2]**

266 Simplify.

$\sqrt{75} + \sqrt{48} - \sqrt{27}$

..... **[2]**

267 **(a)** Simplify.

$\sqrt{45}$

..... **[1]**

(b) Simplify.

$\sqrt{20} + \sqrt{80}$

..... **[2]**

268 Simplify

$$\sqrt{75} + \sqrt{27} - \sqrt{12}$$

..... **[3]**

269 Simplify the following expression, leaving your answer in the form $k\sqrt{3}$.

$\sqrt{48} - \sqrt{12}$

..... **[3]**

270 **(a)** Simplify.

$$\sqrt{75} + \sqrt{12}$$

..... **[2]**

(b) Rationalise the denominator.

$$\frac{14}{\sqrt{7}}$$

..... **[2]**

271 **(a)** Simplify.

$\sqrt{75}$

..... **[2]**

(b) Simplify.

$2\sqrt{18} - \sqrt{8}$

..... **[2]**

272 **(a)** **(i)** Simplify.

$$\sqrt{45} + \sqrt{20}$$

..... **[2]**

(ii) Rationalise the denominator.

$$\frac{18}{\sqrt{3}}$$

..... **[2]**

273 **(a)** Simplify.

$\sqrt{45}$

..... **[1]**

(b) Work out the value of the following expression, giving your answer in the form $k\sqrt{2}$.

$\sqrt{72} + \sqrt{50} - \sqrt{18}$

..... **[2]**

(c) Rationalise the denominator.

$\dfrac{21}{\sqrt{7}}$

..... **[2]**

274 **(a)** Expand and simplify.

$$(4 - \sqrt{3})(2 + \sqrt{3})$$

..... **[2]**

(b) Rationalise the denominator and simplify.

$$\frac{21}{\sqrt{7}}$$

..... **[2]**

275 **(a)** Expand and simplify

$$(3 - \sqrt{5})^2$$

..... **[2]**

(b) Rationalise the denominator.

$$\frac{14}{\sqrt{7}}$$

..... **[2]**

276 **(a)** Expand and simplify.

$$(3 - 2\sqrt{5})(4 + \sqrt{5})$$

..... **[2]**

(b) Rationalise the denominator and simplify.

$$\frac{14}{\sqrt{7}}$$

..... **[2]**

277 **(a)** **(a)** Expand and simplify

$$(3 - 2\sqrt{5})(4 + \sqrt{5})$$

..... **[2]**

(b) Rationalise the denominator.

$$\frac{15}{\sqrt{3}}$$

..... **[2]**

278 **(a)** Expand and simplify

$$(3+\sqrt{2})(4-\sqrt{2})$$

..... **[2]**

(b) Rationalise the denominator.

$$\frac{22}{5-\sqrt{3}}$$

..... **[3]**

279 **(a)** Expand and simplify.

$$(3+\sqrt{2})^2$$

..... **[2]**

(b) Rationalise the denominator and simplify.

$$\frac{4}{3-\sqrt{5}}$$

..... **[3]**

280 **(a)** Expand and simplify.

$$(2\sqrt{5}-3)(2\sqrt{5}+3)$$

..... **[2]**

(b) Rationalise the denominator.

$$\frac{14}{3-\sqrt{2}}$$

..... **[3]**

281 **(a)** Expand and simplify.

$$(3\sqrt{2}-1)(2\sqrt{2}+5)$$

..... **[2]**

(b) Rationalise the denominator and simplify.

$$\frac{26}{4-\sqrt{3}}$$

..... **[3]**

282 **(a)** Rationalise the denominator and simplify fully.

$$\frac{15}{\sqrt{5}}$$

..... **[2]**

(b) Expand and simplify.

$$(2+\sqrt{3})(5-\sqrt{3})$$

..... **[3]**

283 **(a)** Expand and simplify.

$$(3+\sqrt{5})(2-\sqrt{5})$$

..... **[2]**

(b) Rationalise the denominator and simplify.

$$\frac{14}{\sqrt{7}}$$

..... **[3]**

284 **(a)** Expand and simplify.

$$(5 - 2\sqrt{7})(3 + \sqrt{7})$$

..... **[2]**

(b) Rationalise the denominator and simplify fully.

$$\frac{22}{5 - \sqrt{3}}$$

..... **[3]**

285 **(a)** Expand and simplify.

$$(4 - 3\sqrt{2})(5 + 2\sqrt{2})$$

..... **[2]**

(b) Rationalise the denominator and simplify.

$$\frac{11}{4 - \sqrt{5}}$$

..... **[3]**

(c) Solve the equation.

$$x\sqrt{27} - 8 = x\sqrt{3} + 10$$

Give your answer in the form $k\sqrt{3}$ where k is an integer.

$x =$ ..... **[3]**

286 A rectangle has an area of $5 + \sqrt{7}$ cm^2 and a width of $3 - \sqrt{7}$ cm.
Calculate the length of the rectangle.
Give your answer in the form $a + b\sqrt{7}$ where a and b are integers.

..... *cm* **[4]**


287 Solve the equation.

$$x\sqrt{3} - 2 = x + \sqrt{12}$$

Give your answer in the form $a + b\sqrt{3}$, where a and b are integers.

$x =$ ..... **[5]**

288 The diagram shows a rectangle with width $(2-\sqrt{3})$ cm and length $(a+b\sqrt{3})$ cm, where a and b are integers.
The area of the rectangle is $(1+4\sqrt{3})\,\text{cm}^2$.



NOT TO SCALE

Work out the value of a and the value of b.

$a =$ ..... , $b =$ ..... **[5]**

289 **(a)** Rationalise the denominator and simplify.

$$\frac{5+\sqrt{3}}{2-\sqrt{3}}$$

..... **[3]**

(b) Solve the equation.
Give your answer in the form $a+b\sqrt{2}$, where a and b are integers.

$$x(3-\sqrt{2})=14$$

$x =$ ..... **[3]**

290 **(a)** Express $\dfrac{5+\sqrt{3}}{2-\sqrt{3}}$ in the form $a+b\sqrt{3}$, where a and b are integers.

..... **[3]**

(b) A rectangle has an area of $12+4\sqrt{5}$ square centimetres and a width of $3+\sqrt{5}$ centimetres. Calculate the exact length of the rectangle, giving your answer in its simplest form.

Length = ... cm **[4]**

291 **(a)** Rationalise the denominator and simplify.

$$\frac{22}{5-\sqrt{3}}$$


..... **[3]**

(b) Solve the equation.

$x\sqrt{2} = x + \sqrt{8}$ Give your answer in the form $a + b\sqrt{2}$, where a and b are integers.

$x =$ ..... **[4]**

292 A right-angled triangle has a base of $(2+\sqrt{3})$ cm and an area of $(5+3\sqrt{3})$ cm^2.



(a) Show that the exact height, h, is $(2+2\sqrt{3})$ cm.

[4]

(b) The hypotenuse of the triangle is y cm.
Work out the exact value of y^2.
Give your answer in the form $a+b\sqrt{3}$, where a and b are integers.

$y^2 =$ ..... **[3]**

293 **(a)** A rectangle has an area of $(5 + 2\sqrt{3})$ cm^2 and a length of $(2 + \sqrt{3})$ cm.

Work out the width of the rectangle in the form $a + b\sqrt{3}$, where a and b are integers.

..... cm **[3]**

(b) Solve the equation.


$$x\sqrt{5} - 2 = 2x + \sqrt{20}$$

Give your answer in the form $c + d\sqrt{5}$, where c and d are integers.

Show all your working.

$x =$ ..... **[4]**

294 A rectangle has length $(3+\sqrt{5})$ cm and area $(14-2\sqrt{5})$ cm^2.



(a) Calculate the width of the rectangle.
Give your answer in the form $a+b\sqrt{5}$, where a and b are integers.

..... cm **[4]**

(b) The length of the diagonal of the rectangle is d cm.
Show that $d^2 = 308 - 124\sqrt{5}$.

[3]

295 **(a)** Rationalise the denominator and simplify fully.

$$\frac{6}{3-\sqrt{3}}$$

..... **[4]**

(b) Write the following expression in the form $a\sqrt{2}+b\sqrt{3}$, where a and b are integers.

$$\frac{\sqrt{24}+\sqrt{54}}{\sqrt{3}}-\frac{10}{\sqrt{2}}$$

..... **[3]**

296 A right-angled triangle has a base of length $(3+\sqrt{2})$ cm and an area of $(5+2\sqrt{2})$ cm^2.

(a) Show that the height of the triangle is $\dfrac{2(5+2\sqrt{2})}{3+\sqrt{2}}$ cm.

[2]


(b) Rationalise the denominator of $\dfrac{2(5+2\sqrt{2})}{3+\sqrt{2}}$ to express the height in the form $\dfrac{a+b\sqrt{2}}{c}$, where a, b and c are integers.

..... cm **[4]**

(c) Given instead that another expression for a length is $\dfrac{1}{\sqrt{7}-2}$, write this length in the form $\dfrac{\sqrt{7}+c}{d}$ where c and d are integers.

..... **[2]**

297 The diagram shows a right-angled triangle.



(a) Calculate the area of the triangle.

..... cm^2 **[3]**

(b) Show that the length of the hypotenuse is 4 cm.

[4]

(c) Another right-angled triangle is mathematically similar to this triangle.
The area of the similar triangle is 18 cm^2.
Calculate the length of its hypotenuse.

..... cm **[3]**

298 Work out the value of the expression

$$3x^2 - 5xy$$

when $x = -4$ and $y = 2$.

..... **[2]**

299 Work out the value of

$$5x^2 - 3y$$

when $x = -2$ and $y = 4$.

..... **[2]**

300 Find the value of $3a - 2b$ when $a = 4$ and $b = -5$.

..... **[2]**

301 Work out the value of

$$3x^2 - 4y$$

when $x = -3$ and $y = -5$.

..... **[2]**

302 Work out the value of

$$\frac{3x - y^2}{2z}$$

when $x = 5$, $y = -4$ and $z = 0.5$.

..... **[2]**

303 Given the formula $v = u + at$.

Find the value of v when $u = 5$, $a = -2$, and $t = 3$.

$v =$ ..... **[3]**

304 **(a)** Work out the value of $4x - 3y$ when $x = -2$ and $y = 5$.

..... **[2]**

(b) Work out the value of $\dfrac{p^2 - 7}{2q}$ when $p = 5$ and $q = -3$.

..... **[2]**

305 **(a)** Work out the value of $3a - 5b$ when $a = -2$ and $b = 4$.

..... **[2]**

(b) Work out the value of $x^2 + 4x$ when $x = -3$.

..... **[2]**

306 **(a)** Work out the value of $5x - 2y$ when $x = 3$ and $y = -4$.

..... **[2]**

(b) Evaluate

$$\frac{a^2 - b^3}{2c}$$

when $a = -4$, $b = 2$ and $c = 3$.

..... **[2]**

307 **(a)** **(a)** The cost of hiring a plumber is calculated using the formula

$C = 40 + 25h$ where C is the cost in dollars and h is the number of hours worked.
Work out the cost of hiring a plumber for 3.5 hours.

$C = \$ \ldots\ldots\ldots\ldots\ldots$ **[2]**

(b) Given that $v = u + at$, find the value of v when $u = -12$, $a = 9.8$ and $t = 3$.

$v = \ldots\ldots\ldots\ldots\ldots$ **[3]**

308 **(a)** A plumber charges a fixed fee of \$40 plus \$25 for each hour he works.
Write down an expression, in terms of h, for the total cost, in dollars, for h hours of work.

\$ ..... **[1]**

(b) Given the formula

$$P = \frac{k^3 - m^2}{\sqrt{n}}$$

Work out the exact value of P when $k = -2$, $m = -3$ and $n = \frac{1}{16}$.

$P =$ ..... **[3]**

309 Find the exact value of

$$W = \frac{p^{-2} + q^3}{\sqrt[3]{r}}$$

when $p = \frac{1}{2}$, $q = -2$ and $r = -\frac{1}{8}$.

$W =$ ..... **[4]**

310 The distance, d, between two points (x_1, y_1) and (x_2, y_2) is given by the formula

$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$


(a) Work out the value of d when $x_1 = 4$, $y_1 = -3$, $x_2 = -8$, and $y_2 = 2$.

$d =$ ..... **[2]**

(b) Given that $d = 17$, $x_1 = k$, $y_1 = 8$, $x_2 = 3$, and $y_2 = -7$, work out the two possible values of k.

$k =$ ..... , $k =$ ..... **[4]**

311 A cylindrical container of radius r and height h is open at the top.
The total exterior surface area of the container, A, consists of the curved surface area and the circular base.



(a) Show that the total surface area A can be written as

$$A = \pi r(2h + r)$$

[2]

(b) **(i)** Calculate the exact surface area A when $r = 4$ and $h = 6$.
Give your answer in terms of π.

$A =$ ..... **[2]**

(ii) The volume of the cylinder is V.
The formula for the volume is $V = \pi r^2 h$.
Express h in terms of V, π and r, and substitute this into the expression for A to show that

$$A = \frac{2V}{r} + \pi r^2$$

[3]

312 Consider the expression $P = \dfrac{2x^2 - 3y}{x - y}$.

(a) Calculate the value of P when $x = -2$ and $y = 4$.

$P =$ ..... **[3]**

(b) Rearrange the formula to make y the subject.

$y =$ ..... **[4]**

313 The mathematical operation $*$ is defined for all non-zero numbers a and b as

$$a * b = \frac{a^2 - b^2}{ab}$$

(a) Evaluate the following, giving each answer as a fraction in its simplest form.

(i) $6 * (-2)$

..... **[2]**

(ii) $\frac{1}{2} * 4$

..... **[3]**

(b) Given that $x * 2 = \frac{3}{2}$, find the positive value of x.

$x =$ ..... **[3]**

314 A plumber charges a fixed call-out fee of $\$C$ and an hourly rate of $\$R$.

(a) Write down an expression for the total cost, T dollars, for a job that takes h hours.

$T =$ ..... **[1]**

(b) The call-out fee is \$45 and the hourly rate is \$32.50.
Work out the total cost for a job that takes 3.5 hours.

\$ ..... **[2]**

315 A formula is given by

$$P = \frac{\sqrt{x - y}}{z^3}$$

Calculate the value of P when $x = 12.5$, $y = 3.84$, and $z = -1.2$.
Give your answer correct to 3 significant figures.

$P =$ ..... **[3]**

316 A solid shape has a volume, V, given by the formula

$$V = \pi r^2 h + \frac{2}{3}\pi r^3$$

Work out the value of V when $r = 4.5$ and $h = 8.2$.
Give your answer correct to 3 significant figures.

$V =$ ..... **[3]**

317 **(a)** A plumber charges a fixed fee of \$45 plus \$32 for each hour worked.
Write down a formula for the total charge, C dollars, when the plumber works for h hours.

$C =$ ..... **[1]**

(b) Work out the total charge when the plumber works for 3.5 hours.

\$ ..... **[1]**

(c) Work out the value of

$$x^3 - 4xy$$

when $x = -3$ and $y = 2.5$.

..... **[2]**

318 The volume, V, of a cone with radius r and height h is given by the formula

$$V = \frac{1}{3}\pi r^2 h$$

(a) Calculate the value of V when $r = 6.4$ and $h = 12.6$.
Give your answer correct to 1 decimal place.

$V =$ ..... **[2]**

(b) Work out the value of r when $V = 800$ and $h = 14$.
Give your answer correct to 3 significant figures.

$r =$ ..... **[3]**

319 The cost, C dollars, of hiring a campervan for d days and travelling m kilometres is given by the formula

$$C = 85d + 0.24m$$

(a) Calculate the cost of hiring the campervan for 7 days and travelling 450 kilometres.

$ ..... **[2]**

(b) A family hires the campervan for 14 days.
The total cost is $1475.60.
Work out the number of kilometres they travelled.

..... km **[3]**

320 The cost, C dollars, of hiring a plumber for h hours is given by the formula:

$C = 45 + 35h$

(a) Calculate the cost of hiring the plumber for 3.5 hours.

$C =$ ..... **[2]**

(b) The plumber charges a total of \$325 for a job.
Calculate the number of hours the plumber worked.

$h =$ ..... hours **[3]**

321 The volume, V, of a spherical cap of height h and base radius a is given by the formula

$$V = \frac{1}{6}\pi h(3a^2 + h^2)$$

(a) Calculate the volume of a spherical cap with height $h = 4.2$ cm and base radius $a = 6.5$ cm. Give your answer correct to 1 decimal place.

$V =$ ..... cm^3 **[2]**

(b) Rearrange the formula to make a^2 the subject.

$a^2 =$ ..... **[3]**

(c) Find the base radius, a, of a spherical cap with volume $V = 150$ cm^3 and height $h = 3$ cm. Give your answer correct to 3 significant figures.

$a =$ ..... cm **[1]**

322 The value of y is given by the formula

$$y = \frac{a - b^2}{c - \sqrt{d}}$$

Work out the value of y when $a = -4.5$, $b = -3.2$, $c = 1.8$ and $d = 0.64$.

$y =$ ..... **[3]**

323 A formula used in physics is

$$s = ut + \frac{1}{2}at^2$$

(a) **(a)** Calculate the value of s when $u = 5.2$, $t = 4.5$ and $a = -9.8$.

$s =$ ..... **[2]**

(b) Calculate the positive value of t when $s = -9.8$, $u = 4.9$ and $a = -9.8$.

$t =$ ..... **[3]**

324 The volume, V, and surface area, A, of a solid are given by the formulas below.

(a) The formula for the volume is

$$V = \frac{1}{3}\pi h(r^2 + rR + R^2)$$

Calculate the value of V when $h = 12.5$, $r = 4.2$, and $R = 7.8$.
Give your answer correct to 3 significant figures.

$V =$ ..... **[3]**

(b) The formula for the surface area is

$$A = \pi(R + r)\sqrt{(R - r)^2 + h^2} + \pi r^2 + \pi R^2$$

Calculate the value of A when $h = 12.5$, $r = 4.2$, and $R = 7.8$.
Give your answer correct to 1 decimal place.

$A =$ ..... **[3]**

325 Expand and simplify.

$(3x - 2)(2x + 5)$

..... **[2]**

326 **(a)** Expand and simplify.

$3(2x - 1) - 2(x + 4)$

..... **[2]**

(b) Factorise fully.

$15x^2y - 20xy^2$

..... **[2]**

327 **(a)** Expand and simplify.

$$4(2a - 3b) - 3(a - 5b)$$

..... **[2]**

(b) Factorise completely.

$$12x^2y - 18xy^2$$

..... **[2]**

328 **(a)** Expand and simplify.

$4(2x - 3y) - 3(x - 5y)$

..... **[2]**

(b) Factorise completely.

$14a^2b - 21ab^2$

..... **[2]**

329 **(a)** Expand and simplify

$$4(3x-2)-2(x-5)$$

..... **[2]**

(b) Factorise fully

$$15a^2b-10ab^2$$

..... **[2]**

330 **(a)** Expand and simplify.

$3(2x - 5) - 2(x + 4)$

..... **[2]**

(b) Factorise completely.

$8x^2y - 12xy^2$

..... **[2]**

331 **(a)** Expand and simplify

$$3(2x - 5) - 4(x - 2)$$

..... **[2]**

(b) Factorise fully

$$12x^2y - 18xy^2$$

..... **[2]**

332 **(a)** Simplify.

$3a - 5b + 2a + 8b$

..... **[2]**

(b) Factorise completely.

$12x^2y - 18xy^2$

..... **[2]**

333 **(a)** Expand and simplify.

$3(2x - 5) - 4(x - 3)$

..... **[2]**

(b) Factorise fully.

$14x^2y + 21xy^2$

..... **[2]**

334 **(a)** Factorise completely.

$12x^2 - 27y^2$

..... **[2]**

(b) Factorise completely.

$2ax - 6ay + bx - 3by$

..... **[2]**

335 **(a)** Expand and simplify.

$(2x - 3)(x + 5)$

..... **[2]**

(b) Factorise.

$y^2 - 16x^2$

..... **[1]**

(c) Factorise.

$p^2 - 4pq - 5q^2$

..... **[2]**

336 **(a)** Expand and simplify.

$(2x - 3)(x + 5)$

..... **[2]**

(b) Factorise.

$ac - 3bc + 2ad - 6bd$

..... **[2]**

(c) Factorise.

$16x^2 - 25y^2$

..... **[1]**

337 **(a)** Expand and simplify.

$(2x - 3)(x + 4)$

..... **[2]**

(b) Factorise completely.

$ax - bx - 3ay + 3by$

..... **[2]**

(c) Factorise.

$49x^2 - 16y^2$

..... **[2]**

338 **(a)** Factorise.

$x^2 - 8x + 15$

..... **[2]**

(b) Factorise.

$2x^2 - 5x - 12$

..... **[2]**

(c) Factorise.

$ax - 3a + 2bx - 6b$

..... **[2]**

339 **(a)** Expand and simplify.

$(2x - 5)(3x + 1)$

..... **[2]**

(b) Factorise.

$x^2 - 7x - 18$

..... **[2]**

(c) Factorise completely.

$18x^2 - 50y^2$

..... **[3]**

340 **(a)** Expand and simplify.

$(x-2)(2x+3)(x-5)$

..... **[3]**

(b) Factorise completely.

$12x^3 + 17x^2 - 5x$

..... **[3]**

341 **(a)** Factorise completely.

$18x^2 - 50y^2$

..... **[3]**

(b) Factorise completely.

$3x^3 + 10x^2 - 8x$

..... **[3]**

342 **(a)** Expand and simplify.

$$(x-2)(x+3)(2x-1)$$

..... **[3]**

(b) Factorise completely.

(i)

$$6x^2 - 11x - 10$$

..... **[2]**

(ii)

$$ax - ay + 3cx - 3cy$$

..... **[2]**

343 **(a)** Factorise

$$cx - 5dx + 3cy - 15dy$$

..... **[2]**

(b) Factorise fully

$$2x^3 - 18x$$

..... **[2]**

(c) Factorise fully

$$12x^3 - 14x^2 - 10x$$

..... **[3]**

344 **(a)** Expand and simplify.

$(x-4)(2x+1)(x+3)$

..... **[3]**

(b) Factorise completely.

$2ax-3by+2ay-3bx$

..... **[2]**

(c) Factorise.

$6x^2+7x-20$

..... **[2]**

345 **(a)** Factorise

$$6ax - 2ay - 3bx + by$$

..... **[2]**

(b) Factorise completely.

(i)

$$50x^2 - 8y^2$$

..... **[3]**

(ii)

$$2x^3 + 5x^2 - 12x$$

..... **[3]**

346 **(a)** Expand and simplify

$$(x-3)(2x+1)(x+4)$$

..... **[3]**

(b) Factorise

$$2x^2-7x-15$$

..... **[2]**

347 **(a)** Expand and simplify.

$(2x - 3)(x + 5)$

..... **[2]**

(b) Factorise.

$x^2 - 7x - 18$

..... **[2]**

(c) Factorise completely.

$4x^2 - 25y^2$

..... **[2]**

348 **(a)** Expand and simplify

$$(x-4)(2x+3)(x+2)$$

..... **[3]**

(b) Factorise

$$16x^2 - 49y^2$$

..... **[2]**

(c) Factorise

$$3x^2 - 10x - 8$$

..... **[2]**

349 **(a)** Expand and simplify.

$(x-4)(2x+3)(x+1)$

..... **[3]**

(b) Factorise completely.

$6x^2 - 11x - 10$

..... **[2]**

(c) Factorise completely.

$x^3 + 3x^2 - 4x - 12$

..... **[3]**

350 **(a)** Expand and simplify.

$(x-2)(2x+1)(x+3)$

..... **[3]**

(b) Factorise.

$6x^2 - 11x - 10$

..... **[2]**

(c) Factorise completely.

$3x^3 + 5x^2 - 2x$

..... **[3]**

351 **(a)** Expand and simplify.

$(x-2)(x+3)(2x-1)$

..... **[3]**

(b) Factorise.

$6x^2 - 11x - 10$

..... **[2]**

(c) Factorise completely.

$2x^3 - 18x$

..... **[3]**

352 Write as a single fraction in its simplest form.

$$\frac{2x}{5} - \frac{x-1}{3}$$

..... **[2]**

353 Write as a single fraction in its simplest form.

$$\frac{2x-1}{3}+\frac{x+4}{5}$$

..... **[2]**

354 Write as a single fraction in its simplest form.

$$\frac{3x}{4} - \frac{x-2}{5}$$

..... **[2]**

355 Write as a single fraction in its simplest form.

$$\frac{5}{x+2}+\frac{2}{3x-1}$$

..... **[3]**

356 Write as a single fraction in its simplest form.

$$\frac{x+3}{4} - \frac{2x-1}{5}$$

..... **[3]**

357 Write as a single fraction in its simplest form.

$$\frac{3x}{4} - \frac{x-2}{3}$$

..... **[3]**

358 Write as a single fraction in its simplest form.

$$\frac{2x}{3} - \frac{x-4}{5}$$

..... **[3]**

359 **(a)** Write as a single fraction in its simplest form.

$$\frac{2x}{3} + \frac{x-1}{4}$$

..... **[2]**

(b) Simplify fully.

$$\frac{15x^3y^2}{5xy^4}$$

..... **[2]**

360 **(a)** Simplify fully.

$$\frac{3x}{4} + \frac{2x}{5}$$

..... **[2]**

(b) Simplify fully.

$$\frac{5a^2b}{12} \div \frac{10ab^2}{9}$$

..... **[2]**

361 Simplify fully.

$$\frac{x^2 - 16}{x^2 - 5x + 4}$$

..... **[3]**

362 Simplify fully.

$$\frac{3x^2 - 12}{x^2 + 3x - 10}$$

..... **[4]**

363 **(a)** Write as a single fraction in its simplest form.

$$\frac{3}{x-2} - \frac{2}{x+5}$$

..... **[3]**

(b) Simplify fully.

$$\frac{x^2-16}{x^2-2x-8}$$

..... **[3]**

364 **(a)** Simplify fully.

$$\frac{x^2 - 9}{2x^2 + 5x - 3}$$

..... **[3]**

(b) Write as a single fraction in its simplest form.

$$\frac{4}{x - 2} - \frac{3}{x + 1}$$

..... **[3]**

365 Simplify fully.

$$\frac{2x^2 + 5x - 12}{4x^2 - 9}$$

..... **[4]**

366 Simplify fully.

$$\frac{3x^2 - 5x - 2}{x^2 - 4} \div \frac{3x + 1}{x^2 - x - 2}$$

..... **[6]**

367 **(a)** Simplify fully.

$$\frac{2x^2 - 5x - 3}{4x^2 - 9}$$

..... **[3]**

(b) Write as a single fraction in its simplest form.

$$\frac{3}{x-2} + \frac{4}{x+1} - \frac{2x}{x^2 - x - 2}$$

..... **[4]**

368 Simplify fully.

$$\frac{x^2 - 9}{2x^2 + 5x - 3}$$

..... **[3]**

369 Write as a single fraction in its simplest form.

$$\frac{5}{x-2} - \frac{3}{2x+1}$$

..... **[3]**

370 Simplify fully.

$$\frac{2x^2 - 9x + 9}{2x^2 - 18}$$

..... **[4]**

371 Simplify fully.

$$\frac{2x^2 - 8}{x^2 + x - 6}$$

..... **[5]**

372 **(a)** Write as a single fraction in its simplest form.

$$\frac{x+3}{2x-1} - \frac{x-2}{x+4}$$

..... **[3]**

(b) Simplify fully.

$$\frac{x^2-16}{x^2+4x}$$

..... **[2]**

373 Simplify fully.

$$\frac{x^2-25}{x^2-3x} \div \frac{x^2+6x+5}{2x-6}$$

..... **[4]**

374 **(a)** Simplify fully.

$$\frac{2x^2 - 8x}{x^2 - 16} \div \frac{6x}{3x + 12}$$

..... **[4]**

(b) Write as a single fraction in its simplest form.

$$\frac{5}{x - 3} + \frac{2x + 1}{x^2 - 9}$$

..... **[3]**

375 **(a)** Write as a single fraction in its simplest form.

$$\frac{4}{x-3} - \frac{3}{2x-1}$$

..... **[3]**

(b) Simplify fully.

$$\frac{x^2-25}{2x^2-9x-5} \div \frac{x+5}{4x+2}$$

..... **[4]**

376 **(a)** Write as a single fraction in its simplest form.

$$\frac{3}{x-2} - \frac{x+5}{x^2-4}$$

..... **[3]**

(b) Simplify fully.

$$\frac{4x^2-25}{2x^2-5x} \times \frac{6x^3}{4x^2+10x}$$

..... **[4]**

377 **(a)** Write as a single fraction in its simplest form.

$$\frac{5}{x-2}+\frac{4}{2x+3}$$

..... **[3]**

(b) Simplify fully.

$$\frac{2x^2-5x-12}{4x^2-9}\div\frac{x-4}{2x-3}$$

..... **[4]**

378 **(a)** Simplify fully.

$$\frac{2x^2 + 5x - 3}{4x^2 - 1}$$

..... **[4]**

(b) Express as a single fraction in its simplest form.

$$\frac{5}{x - 3} + \frac{x + 2}{x^2 - 4x + 3}$$

..... **[4]**

379 Simplify fully.

$4x^3y^{-2} \times 3x^5y^6$

..... **[2]**

380 **(a)** Simplify.

$$4x^2 \times 3x^{-5}$$

..... **[1]**

(b) Simplify.

$$\left(\frac{8x^9}{27}\right)^{\frac{1}{3}}$$

..... **[2]**

381 **(a)** Simplify.

$$4x^3 \times 5x^{-2}$$

..... **[1]**

(b) Simplify.

$$\left(\frac{2x^4}{y^3}\right)^2$$

..... **[2]**

(c) Solve the equation.

$$3^p = \frac{1}{27}$$

$p =$ ..... **[1]**

382 **(a)** Simplify.

$$4x^3y^{-2} \times 3x^5y^6$$

..... **[2]**

(b) Find the exact value of

$$16^{-\frac{1}{4}}$$

..... **[2]**

383 **(a)** Simplify.

$5x^4 \times 3x^{-2}$

..... **[2]**

(b) Find the value of

$16^{\frac{3}{4}}$

..... **[2]**

384 **(a)** Simplify.

$y^5 \times y^{-2}$

..... **[1]**

(b) Work out the value of

5^{-2}

..... **[1]**

(c) Simplify.

$(2x^4)^3$

..... **[2]**

385 **(a)** Find the value of x when

$$2^x = \frac{1}{8}$$

$x =$ ..... **[1]**

(b) Simplify

$$(4y^6)^{\frac{1}{2}}$$

..... **[2]**

(c) Work out the exact value of

$$16^{-\frac{3}{4}}$$

..... **[1]**

386 **(a)** Simplify.

$w^4 \times w^7$

..... **[1]**

(b) Simplify.

$(3x^3)^2$

..... **[2]**

(c) Simplify.

$$\frac{15y^8}{5y^{-2}}$$

..... **[2]**

387 **(a)** Simplify.

$p^4 \times p^7$

..... **[1]**

(b) Simplify.

$(3q^3)^2$

..... **[2]**

(c) Work out the value of $27^{\frac{2}{3}}$.

..... **[2]**

388 **(a)** Simplify fully

$$\frac{12x^5y^3}{3x^2y^{-1}}$$

..... **[2]**

(b) Solve the equation

$$27^{x-1} = 9^{2x+1}$$

$x =$ ..... **[3]**

389 **(a)** Work out the exact value of

$$32^{-\frac{2}{5}}$$

..... **[2]**

(b) Solve the equation.

$$5^{x+2} = 125^{x}$$

$x =$ ..... **[3]**

390 **(a)** Work out the exact value of

$$125^{\frac{2}{3}}$$

..... **[2]**

(b) Simplify fully.

$$\frac{15x^4y^{-3}}{5x^{-2}y^5}$$

..... **[3]**

391 **(a)** Solve the equation.

$$16^x = 2$$

$x =$ ..... **[1]**

(b) Work out the exact value of

$$125^{\frac{2}{3}} \times 5^{-2}$$

..... **[2]**

(c) Solve the equation.

$$5^{2x-1} = 125^x$$

$x =$ ..... **[3]**

392 **(a)** Simplify fully.

$$4x^{-3} \times \frac{3}{2}x^{\frac{1}{2}}$$

..... **[2]**

(b) Solve the equation.

$$3^{2x+1} = 27$$

$x =$ ..... **[2]**

(c) Simplify fully.

$$\left(\frac{8a^6}{b^3}\right)^{\frac{1}{3}}$$

..... **[2]**

393 **(a)** Simplify fully.

$$\left(\frac{2x^5}{3y^2}\right)^3$$

..... **[3]**

(b) Solve the equation.

$$9^x = 27^{x+1}$$

$x =$ ..... **[3]**

394 **(a)** Simplify fully.

$$\left(\frac{27x^9}{64y^{12}}\right)^{-\frac{1}{3}}$$

..... **[3]**

(b) Solve the equation.

$$2^{3x-1} = 32^x$$

$x =$ ..... **[3]**

395 **(a)** Simplify fully.

$$\left(\frac{16x^{12}}{y^8}\right)^{\frac{3}{4}}$$

..... **[3]**

(b) Solve the equation.

$$2^{x+3} = 8^x$$

$x =$ ..... **[3]**

396 **(a)** Solve the equation.

$$9^{2x} = 27^{x+1}$$

$x =$ ..... **[3]**

(b) Simplify fully.

$$\left(\frac{16a^8}{81b^{12}}\right)^{-\frac{3}{4}}$$

..... **[3]**

397 **(a)** Solve the equation.

$$16^{2x-1} = 8^{x+2}$$

$x =$ ..... **[3]**

(b) Simplify fully.

$$\left(\frac{81x^{12}}{16y^{8}}\right)^{-\frac{3}{4}}$$

..... **[3]**

398 **(a)** Simplify fully.

$$\frac{16x^{\frac{5}{2}}y^{-2}}{2x^{\frac{1}{2}}y^{3}} \div \frac{4x^{3}}{y^{4}}$$

..... **[3]**

(b) Solve the equation.

$$5^{2x-1} \times 25^{x+2} = 125^{x}$$

$x =$ ..... **[4]**

399 **(a)** Simplify fully.

$$\frac{4x^{-\frac{1}{2}} \times 3x^2}{6x^{\frac{1}{3}}}$$

..... **[3]**

(b) Solve the equation.

$$9^{2x-1} \times 3^{x+1} = 27^x$$

$x =$ ..... **[4]**

400 **(a)** Simplify fully.

$$\frac{15x^{\frac{2}{3}} \times 2x^{-\frac{1}{6}}}{5x^{\frac{3}{2}}}$$

..... **[3]**

(b) Solve the equation.

$$x^{-\frac{2}{3}} = \frac{4}{9}$$

$x =$ ..... **[3]**

(c) Solve the equation.

$$9^x \times 3^{2x-1} = 27^{x+1}$$

$x =$ ..... **[4]**

401 **(a)** Simplify fully.

$$\left(\frac{16x^{12}}{y^8}\right)^{\frac{3}{4}}$$

..... **[3]**

(b) Solve the equation.

$$4^{2x-1} = 8^{x+2}$$

$x =$ ..... **[3]**

402 **(a)** Simplify fully.

$$\frac{24x^{\frac{3}{2}}y^{-2}}{6x^{-\frac{1}{2}}y}$$

..... **[2]**

(b) Solve the equation.

$$9^x \times 27^{x-1} = 3\sqrt{3}$$

$x =$ ..... **[4]**

403 **(a)** Solve the equation.

$$4^x \times 8^{2x-1} = 16$$

$x =$ ..... **[3]**

(b) Simplify fully.

$$\frac{12x^{\frac{3}{2}} \times 2x^{-\frac{1}{2}}}{8x^{-3}}$$

..... **[3]**

404 **(a)** Simplify

$$\left(\frac{8x^9}{27y^6}\right)^{-\frac{2}{3}}$$

..... **[3]**

(b) Find the values of a and b given that

$$\frac{5^{a-2} \times 25^b}{125^{a+1}} = 5^4$$

$a =$ ..... , $b =$ ..... **[4]**

405 **(a)** Simplify fully.

$$\frac{12x^{\frac{3}{2}}y^{-2}}{4x^{-\frac{1}{2}}y^{4}}$$

..... **[3]**

(b) Solve the equation.

$$4^{2x-1} \times 8^{x+1} = 32^{x}$$

$x =$ ..... **[4]**

406 Solve the equation.

$$\frac{3x-1}{4} = \frac{x+5}{3}$$

$x =$ ..... **[3]**

407 Solve the equation.

$$\frac{2x-3}{4}+\frac{x+1}{3}=2$$

$x =$ ..... **[3]**

408 Solve the simultaneous equations.

$2x - y = 7$

$3x + 2y = 14$

$x =$ ..... , $y =$ ..... **[3]**

409 Solve the simultaneous equations.
You must show all your working.

$4x + 3y = 5$

$2x - 5y = 22$

$x =$ ..... , $y =$ ..... **[3]**

410 **(a)** Solve the equation.

$$4(x-3) = 2x + 10$$

$x =$ ..... **[2]**

(b) Rearrange the formula to make x the subject.

$$y = \frac{2x-1}{5}$$

$x =$ ..... **[2]**

411 **(a)** Solve the equation.

$$3(2x - 1) = 5x + 8$$

$x =$ ..... **[2]**

(b) Rearrange the formula to make x the subject.

$$y = 4x - 7$$

$x =$ ..... **[2]**

412 **(a)** Solve the equation.

$$3(2x - 5) = 4(x + 1)$$

$x =$ ..... **[3]**

(b) Make h the subject of the formula.

$$V = \frac{1}{3}\pi r^2 h$$

$h =$ ..... **[2]**

413 **(a)** Solve the equation.

$$3(2x - 1) = 5x + 8$$

$x =$ ..... **[2]**

(b) Make x the subject of the formula.

$$y = \frac{2x + 3}{5}$$

$x =$ ..... **[3]**

414 **(a)** Solve the equation.

$3(2x - 1) = 4x + 11$

$x =$ ..... **[2]**

(b) Solve the simultaneous equations.

$3x + 2y = 16$

$x - y = 2$

$x =$ ..... , $y =$ ..... **[3]**

415 **(a)** Solve

$$5x - 3 = 2(x + 6)$$

$x =$ ..... **[2]**

(b) Solve

$$\frac{x+2}{3} + \frac{2x-1}{4} = 3$$

$x =$ ..... **[3]**

416 **(a)** Solve the equation.

$$3(2x - 5) = 4x + 7$$

$x =$ ..... **[3]**

(b) Rearrange the formula to make x the subject.

$$y = \frac{2x + 3}{5}$$

$x =$ ..... **[2]**

417 **(a)** Solve the equation.

$$4(2x - 1) = 3x + 11$$

$x =$ ..... **[3]**

(b) Make x the subject of the formula.

$$y = \frac{3x - 5}{2}$$

$x =$ ..... **[3]**

418 Consider the following algebraic tasks.

(a) Solve the equation.

$$5(x-2) = 3x + 14$$

$x =$ ..... **[3]**

(b) Rearrange the formula to make b the subject.

$$A = \frac{h(a+b)}{2}$$

$b =$ ..... **[3]**

419 **(a)** Solve the equation.

$$3(2x - 5) = 4x + 7$$

$x =$ ..... **[3]**

(b) Solve the simultaneous equations.
You must show all your working.

$$2x + 3y = 13 \qquad 5x - 2y = 4$$

$x =$ ..... , $y =$ ..... **[3]**

420 **(a)** Solve.

$$4(x - 3) = 2x + 7$$

$x =$ ..... **[3]**

(b) Solve the simultaneous equations.
You must show all your working.

$$5x - 2y = 16$$

$$3x + 4y = 7$$

$x =$ ..... , $y =$ ..... **[3]**

421 Rearrange the formula to make x the subject.

$$y = \frac{3x - 2}{x + 4}$$

$x =$ ..... **[4]**

422 Rearrange the formula to make x the subject.

$$y = \frac{2x + 3}{x - 5}$$

$x =$ ..... **[4]**

423 Rearrange the formula to make x the subject.

$$y = \frac{3x - 5}{2x + 1}$$

$x =$ ..... **[4]**

424 **(a)** Solve the simultaneous equations.
You must show all your working.

$3x + 2y = 12$

$5x - 3y = 1$

$x =$ ..... , $y =$ ..... **[3]**

(b) Solve the equation by factorisation.

$2x^2 - 5x - 12 = 0$

$x =$ ..... , $x =$ ..... **[3]**

425 **(a)** Solve the simultaneous equations.
You must show all your working.

$$2x + 3y = 13$$
$$5x - 2y = 4$$

$x =$ ..... , $y =$ ..... **[3]**

(b) Make x the subject of the formula.

$$y = \frac{3x - 1}{x + 2}$$

$x =$ ..... **[4]**

426 **(a)** Rearrange the formula to make x the subject.

$$y = \frac{3x - 2}{x + 4}$$

$x =$ ..... **[4]**

(b) Solve the equation.

$$\frac{5}{x - 2} = \frac{2}{x + 1}$$

$x =$ ..... **[3]**

427 **(a)** Solve the simultaneous equations.
You must show all your working.

$3x + 2y = 12$

$5x - 3y = 1$

$x =$ ..... , $y =$ ..... **[4]**

(b) Solve the equation by factorisation.

$2x^2 - 7x - 15 = 0$

$x =$ ..... , or $x =$ ..... **[3]**

428 **(a)** Solve the simultaneous equations.
You must show all your working.

$3x - 2y = 19$

$4x + y = 18$

$x =$ ..... , $y =$ ..... **[3]**

(b) Rearrange the formula to make p the subject.

$$T = \frac{3p + 4}{p - 2}$$

$p =$ ..... **[4]**

429 **(a)** Solve the simultaneous equations.
You must show all your working.

$2x - y = 9$ $3x + 2y = 10$

$x =$ ..... , $y =$ ..... **[4]**

(b) Solve the equation.

$$\frac{2x}{3} - \frac{x-4}{4} = \frac{5}{2}$$

$x =$ ..... **[4]**

430 Solve the following equations.

(a) Solve the simultaneous equations.
You must show all your working.

$4x + 3y = 5$

$2x - 5y = 9$

$x =$ ..... , $y =$ ..... **[4]**

(b) Solve the equation by factorisation.

$3x^2 - 10x - 8 = 0$

$x =$ ..... or $x =$ **[4]**

431 Solve the equation.

$$\frac{2}{x-1}+\frac{1}{x-2}=1$$

Give your answers in the form $a \pm \sqrt{b}$, where a and b are integers.
You must show all your working.

$x =$ ..... **[6]**

432 **(a)** Tickets for a school show cost \$12 for an adult and \$8 for a child.
For one performance, a total of 150 tickets were sold and the total amount received was \$1480.
Let x be the number of adult tickets sold and y be the number of child tickets sold.

Write down a pair of simultaneous equations in x and y and solve them to find the value of x and the value of y.

$x =$ ..... , $y =$ ..... **[4]**

(b) Make m the subject of the formula.

$$T = 2\pi\sqrt{\frac{m}{k}}$$

$m =$ ..... **[3]**

433 **(a)** Rearrange the formula to make x the subject.

$$p = \frac{3x - 1}{x + 4}$$

$x =$ ..... **[4]**

(b) Solve the equation by factorising.

$$2x^2 - 5x - 12 = 0$$

$x =$ ..... , $x =$ ..... **[3]**

434 **(a)** A rectangle has length $(3x + 2)$ cm and width $(x - 1)$ cm.
The perimeter of the rectangle is 42 cm.

Work out the value of x.

$x =$ ..... **[3]**

(b) Make x the subject of the formula.

$$p = \sqrt{\frac{x-3}{x}}$$

$x =$ ..... **[4]**

435 **(a)** Make x the subject of the formula.

$$y = \frac{2x + 5}{x - 3}$$


$x =$ ..... **[3]**

(b) Solve the equation.
Show all your working and give your answers correct to 2 decimal places.

$$\frac{x + 1}{3} - \frac{2}{x - 2} = 1$$

$x =$ ..... , $x =$ ..... **[6]**

436 The diagram shows a rectangle with length $(2x + 3)$ cm and width $(x - 1)$ cm.
The area of the rectangle is $45\,\text{cm}^2$.



NOT TO SCALE

(a) Show that $2x^2 + x - 48 = 0$.

[2]

(b) Solve the equation $2x^2 + x - 48 = 0$.
You must show all your working and give your answers correct to 2 decimal places.

$x =$ ..... , $x =$ ..... **[4]**

(c) Make x the subject of the formula.

$$y = \frac{ax + 5}{x - 3}$$

$x =$ ..... **[4]**

437 Solve the equation.
Give your answers correct to 2 decimal places.

$$\frac{3}{x+2} + \frac{1}{x-1} = 2$$

$x =$ ..... , $x =$ ..... **[6]**

438 Solve the equation.
Show all your working and give your answers correct to 2 decimal places.

$$\frac{3}{x-2}+\frac{4}{2x+1}=2$$

$x =$ ..... , $x =$ ..... **[6]**

439 Solve the equation.
Show all your working and give your answers correct to 2 decimal places.

$$\frac{5}{x-2} - \frac{3}{2x+1} = 2$$

$x =$ ..... , $x =$ ..... **[7]**

440 **(a)** Show that $\frac{3}{x-1} - \frac{2}{x+3} = 1$ simplifies to $x^2 + x - 14 = 0$.

[4]

(b) Solve the equation $x^2 + x - 14 = 0$.
Show all your working and give your answers correct to 2 decimal places.

$x =$ ..... , $x =$ ..... **[3]**

441 **(a)** A car travels 120 km at an average speed of x km/h.
A train travels the same distance at an average speed of $(x + 10)$ km/h.
The time taken by the car is 36 minutes more than the time taken by the train.

(i) Show that $x^2 + 10x - 2000 = 0$.

[3]

(ii) Solve the equation $x^2 + 10x - 2000 = 0$.

$x =$ ..... or $x =$ ..... **[2]**

(b) Solve the equation.

$3x^2 - 7x - 5 = 0$ Give your answers correct to 2 decimal places.
You must show all your working.

$x =$ ..... or $x =$ ..... **[3]**

442 **(a)** A train travels 120 km at an average speed of x km/h.
It then travels a further 150 km at an average speed of $(x + 10)$ km/h.
The total time for the journey is 4 hours.
Write down an equation in x and show that it simplifies to $2x^2 - 115x - 600 = 0$.

[4]

(b) Solve the equation $2x^2 - 115x - 600 = 0$ to find the value of x.
Give your answer correct to 1 decimal place.

$x =$ ..... **[2]**

(c) Use your graphic display calculator to solve the equation.

$$2^x - 3 = \frac{5}{x}$$

Give your answers correct to 3 significant figures.

$x =$ ..... , $x =$ ..... **[2]**

443 A cyclist travels 40 km at an average speed of x km/h.

(a) Write down an expression, in terms of x, for the time taken in hours.

..... hours **[1]**

(b) A car travels the same 40 km journey at an average speed of $(x + 20)$ km/h.
The car takes 45 minutes less than the cyclist.
Form an equation in x and show that it simplifies to $3x^2 + 60x - 3200 = 0$.

[4]

(c) Solve the equation.

$$3x^2 + 60x - 3200 = 0$$

Give your answers correct to 2 decimal places.

$x =$ ..... , or $x =$ ..... **[3]**

444 **(a)** Solve the equation.

$$\frac{5}{x+2} + \frac{2}{x-3} = 1$$

Show all your working and give your answers correct to 2 decimal places.

$x =$ ..... , $x =$ ..... **[5]**

(b) Make x the subject of the formula.

$$y = \frac{2x-3}{5x+1}$$

$x =$ ..... **[4]**

445 **(a)** Solve the equation.

Give your answers in the exact form $\frac{p \pm \sqrt{q}}{2}$.

$$\frac{x}{x+2} = \frac{3}{x-6}$$

$x =$ ..... **[5]**

(b) Rearrange the formula to make x the subject.

$$y = \frac{2x-5}{3x+1}$$

$x =$ ..... **[4]**

446 **(a)** Solve the equation.

$$\frac{3}{x-1} + \frac{2}{2x+1} = 2$$

Show all your working and give your answers correct to 2 decimal places.

$x =$ ..... , or $x =$ ..... **[5]**

(b) The sum of the squares of two consecutive positive even numbers is 340.

Write down an equation in x and solve it to find the two numbers.

The numbers are ..... , and ..... **[4]**

447 A bus travels 100 km at an average speed of x km/h.
A car travels 100 km at an average speed of $(x + 15)$ km/h.
The car takes 30 minutes less than the bus.

(a) Write down an equation in x and show that it simplifies to $x^2 + 15x - 3000 = 0$.

[3]

(b) Solve the equation $x^2 + 15x - 3000 = 0$.
You must show all your working and give your answers correct to 2 decimal places.

$x =$ ..... , $x =$ ..... **[4]**

(c) Work out the time taken by the car to travel the 100 km.
Give your answer in hours and minutes, correct to the nearest minute.

hours, minutes **[3]**

448 **(a)** Make x the subject of the formula.

$$y = \frac{3x - 2}{x + 4}$$

$x =$ ..... **[4]**

(b) Solve the equation.
Show all your working and give your answers correct to 2 decimal places.

$$\frac{5}{x - 2} - \frac{3}{2x + 1} = 2$$

$x =$ ..... , or $x =$ ..... **[6]**

449 Consider the following algebraic equations and formulas.

(a) Rearrange the formula to make x the subject.

$$y = \frac{3x - 1}{x + 2}$$

$x =$ ..... **[4]**

(b) Solve the equation.
Give your answers correct to 2 decimal places.
Show all your working.

$$\frac{2}{x - 1} + \frac{3}{x + 2} = 1$$

$x =$ ..... or $x =$ **[6]**

450 A train travels 150 km at an average speed of x km/h.

(a) Write down an expression, in terms of x, for the time taken in hours.

..... hours **[1]**

(b) A bus travels the same distance of 150 km at an average speed of $(x - 15)$ km/h.
The bus takes 30 minutes longer than the train.
Show that x satisfies the equation $x^2 - 15x - 4500 = 0$.

[4]

(c) Solve the equation $x^2 - 15x - 4500 = 0$ to find the average speed of the train.

..... km/h **[3]**

(d) Rearrange the formula to make q the subject.

$$P = \frac{2q + 3}{q - 1}$$

$q =$ ..... **[4]**

451 Solve the inequality.

$$3x - 5 < 7x + 3$$

..... **[2]**

452 **(a)** Solve the inequality.

$$3(x-2) > 5x + 4$$

..... **[2]**

(b) Write down the largest integer that satisfies the inequality in part (a).

..... **[1]**

453 Solve the inequality.

$$4(3x - 1) - 5x \geq 2(x + 8)$$

..... **[3]**

454 **(a)** Solve the inequality

$$3(x-4) < 5x - 2$$

..... **[2]**

(b) Write down the smallest integer value of x that satisfies this inequality.


..... **[1]**

455 **(a)** Solve the inequality.

$$3(2x - 1) > 4x + 7$$

..... **[2]**

(b) On the number line, represent the solution to the inequality in part (a).



[1]

(c) Write down the smallest integer value of x that satisfies this inequality.

..... **[1]**

456 **(a)** Solve the inequality.

$$-5 < 3x + 1 \leq 13$$

..... **[2]**

(b) Write down all the integer values of x that satisfy this inequality.


..... **[2]**

457 **(a)** Solve the inequality.

$$-5 < 2x - 1 \leq 7$$

..... **[2]**

(b) Show your solution to part (a) on the number line.



[2]

458 **(a)** Solve the inequality.

$4x - 3 \leq 17$

..... **[2]**

(b) Write down the largest integer x that satisfies the inequality.

$3(x + 2) < 20$


$x =$ ..... **[2]**

459 **(a)** Solve the inequality.

$$4x - 3 < 2x + 9$$

..... **[2]**

(b) Write down the inequality shown on the number line.




..... **[2]**

460 **(a)** Solve the inequality.

$$-5 < 2x - 1 \leq 7$$

..... **[3]**

(b) Represent your answer to part (a) on the number line provided.




[2]

461 **(a)** Solve the inequality.

$$3(x-2) \leq 5x+8$$

. . . . . **[3]**

(b) Represent the solution to part (a) on the number line.



[2]

462 **(a)** Solve the inequality.

$3x - 2 > 5x + 7$


..... **[2]**

(b) Write down all the integer values of n that satisfy the inequality.

$-5 < 2n - 1 \leq 3$

..... **[3]**

(c) An inequality is shown on the number line above.
Write down the inequality.



..... **[1]**

463 **(a)** Solve the inequality.

$$4x - 3 \leq 7x + 12$$

..... **[2]**

(b) Solve the inequality.

$$-8 < 2x - 1 \leq 7$$

..... **[2]**

(c) Write down the largest integer and the smallest integer that satisfy the inequality in part (b).

Largest = ... , Smallest = ... **[2]**

464 **(a)** Solve the inequality.

$$2 - 3x < 11$$

..... **[2]**

(b) Solve the inequality.

$$4(2x - 1) \leq 5x + 17$$

..... **[3]**

(c) n is an integer such that $2 - 3n < 11$ and $4(2n - 1) \leq 5n + 17$.
Write down all the possible values of n.


..... **[2]**

465 **(a)** Solve the inequality.

$3x - 5 \leq 7x + 3$

..... **[2]**

(b) On the number line, show the solution to the inequality in part (a).




[2]

(c) List all the integers that satisfy the inequality.

$-2 < 3x - 5 \leq 4$

..... **[3]**

466 The unshaded region R in the diagram is defined by three inequalities.




Write down these three inequalities.

.....

.....

..... **[3]**

467 The diagram shows a region R, which is unshaded.
The unwanted regions are shaded.




Write down the three inequalities that define region R.

.....

.....

..... **[3]**

468 The diagram shows a region R defined by three inequalities.
The unwanted regions are shaded.




Write down the three inequalities that define the unshaded region R.

.....

..... **[3]**

469 The unshaded region R is defined by three inequalities.
The boundaries of the region are shown on the coordinate grid.



Write down the three inequalities that define the unshaded region R.

.....

.....


..... **[3]**

470 **(a)** Solve the inequality.

$$-4 < 3x + 2 \leq 14$$

..... **[2]**

(b) Represent your answer to part (a) on the number line.




[2]

471 **(a)** Solve the inequality.

$$-7 < 2x - 1 \leq 5$$

..... **[3]**

(b) Represent your answer to part (a) on the number line.



[2]

472 **(a)** Solve the inequality.

$$-5 \leq 2x - 1 < 6$$

(a) Solve the inequality.


$$-5 \leq 2x - 1 < 6$$

..... **[3]**

(b) Write down all the integer values of x that satisfy this inequality.

..... **[2]**

473 The unshaded region R is defined by three inequalities.



(a) Write down these three inequalities.

.....

.....


..... **[3]**

(b) The point (a, b), where a and b are integers, lies in the region R.
It is given that $a = 3$.

Find all possible values of b.

..... **[3]**

474 The diagram shows a region R, which is unshaded.
The region is bounded by three lines.
One of the lines is broken.




Write down the three inequalities that define the region R.

.....

.....

..... **[3]**

475 The diagram shows a grid with three straight lines and an unshaded region labeled R.




(a) Write down the three inequalities that define the unshaded region R.

[3]

(b) Find the maximum value of $3x + 2y$ for integers x and y that lie in the region R.

..... **[2]**

476 A region R is defined by three inequalities.



The region R, unshaded, is defined by three inequalities.
Write down these three inequalities.

.....

.....

..... **[6]**

477 **(a)** Solve the inequality.


$$\frac{3x-2}{5} - \frac{2x-5}{3} \leq 2$$

..... **[4]**

(b) A region R is defined by the three inequalities $y > x$, $y < 4$, and $x + y > 2$.
Write down all the points in region R that have integer coordinates.

..... **[3]**

478 The diagram shows a region R unshaded.




Write down the three inequalities that define the region R.

.....

.....

..... **[3]**

479 The diagram shows three straight lines: $x = 1$, $y = 2x$, and $x + y = 6$.
The unshaded region R is bounded by these lines.



Write down the three inequalities that define the unshaded region R.

.....

.....

..... **[3]**

480 **(a)** Solve the double inequality.


$$-7 \leq 3x - 1 < 11$$

..... **[3]**

(b) Write down all the integer values of x that satisfy the inequality in part (a).

..... **[2]**

481 The diagram shows a region R which is unshaded.
The unwanted regions are shaded.



(a) Write down the three inequalities that define region R.

.....

.....

..... **[4]**

(b) Find the point (x, y), with integer coordinates, inside region R which maximizes the value of $2y - x$.

((..... , ...)) **[2]**

482 **(a)** Solve the inequality.

$$-5 < 2x + 1 \leq 11$$

..... **[3]**

(b) Write down all the integers that satisfy the inequality.

$$-2.5 \leq y < 1$$


..... **[2]**

(c) Use your graphic display calculator to solve the inequality.

$$2x^2 - 7x + 3 > 0$$

..... **[2]**

483 The diagram shows a grid with three straight lines.



(a) The unshaded region R is defined by three inequalities.
The boundaries are included in the region R.
Write down the three inequalities.

.....

.....

..... **[5]**

(b) A point (x, y) lies in the region R.
Both x and y are integers.
Find the maximum value of $2x + y$.

..... **[2]**

484 **(a)** Solve the inequality.

$$-5 < 2x - 1 \leq 7$$

..... **[2]**

(b) Write down all the integers that satisfy the inequality in part (a).

..... **[2]**

(c) A rectangular garden has length $(3x + 2)$ metres and width x metres.
The perimeter of the garden is strictly less than 40 metres.


(i) Write down an inequality in terms of x and show that it simplifies to $x < 4.5$.

[2]

(ii) Given that x is a positive integer, find the maximum possible area of the garden.

Area = ... m^2 **[2]**

485 The diagram shows a grid with three lines drawn.
The unwanted regions are shaded grey.



(a) Write down the three inequalities that define the unshaded region R.

.....

.....

..... **[3]**

(b) Find the coordinates of the point (x, y) in region R where x and y are integers, such that the value of $2x + 3y$ is a maximum.

((..... , ...)) **[2]**

486 A factory produces two types of bicycles: Mountain bikes and City bikes.
In a week, the factory produces x Mountain bikes and y City bikes.

- The factory produces at least 15 Mountain bikes.
- The factory can produce at most 40 bicycles in a week.
- The number of City bikes produced is at least half the number of Mountain bikes produced.

(a) Write down three inequalities, in terms of x and y, to show this information.

.....


.....

..... **[3]**

(b) The profit on each Mountain bike is \$400 and the profit on each City bike is \$150.
Work out the maximum possible profit the factory can make in a week.

\$ ..... **[3]**

487 The unshaded region R in the diagram is defined by three inequalities.



(a) Write down the three inequalities that define the unshaded region R.

.....

.....

..... **[3]**

(b) Use a graphic display calculator to solve the inequality.


$x^2 - 4x - 2 \leq 0$

Give your answer correct to 2 decimal places.

..... **[3]**

488 **(a)** On the grid, draw the lines $y = 2$, $y = x + 2$, and $2x + y = 8$.
Shade the unwanted regions to show the region R defined by the inequalities:

- $y \geq 2$
- $y \leq x + 2$
- $2x + y \leq 8$




[4]

(b) Find the largest value of $3x + 2y$ for points (x, y) in the region R where both x and y are integers.

..... **[2]**

489 The diagram shows three straight lines on a coordinate grid.



(a) The region R is unshaded.
Write down the three inequalities that define the region R.

.....

.....

..... **[3]**

(b) Find the maximum value of $2x + 3y$ given that x and y are integers that satisfy these three inequalities.

..... **[3]**

(c) A point is chosen at random from the points with integer coordinates in region R.
Work out the probability that $x = y$.

. . . . . **[2]**

490 **(a)** Solve the inequality.

$$-5 < 2x - 1 \leq 7$$

..... **[2]**

(b) Find the smallest integer x that satisfies the inequality.

$$\frac{2x-5}{3} - \frac{x-3}{4} > 1$$

$x =$ ..... **[3]**

(c) A taxi company, City Cabs, charges a fixed fare of \$3.20 plus \$1.50 per kilometre.
Another company, Metro Taxis, charges a fixed fare of \$5.00 plus \$1.10 per kilometre.
Construct and solve an inequality to find the minimum whole number of kilometres a passenger must travel for City Cabs to be more expensive than Metro Taxis.

..... kilometres **[3]**

491 Tickets for a community show cost \$5 for children and \$10 for adults.
The maximum number of tickets that can be sold is 500.
The organizers must sell at least 150 adult tickets.
The total amount of money raised from ticket sales must be at least \$3000.

Let x be the number of children's tickets sold and y be the number of adult tickets sold.


(a) Write down three inequalities in x and y to show this information.

.....

.....

..... **[3]**

(b) On the grid, draw three lines to show these inequalities.
Shade the unwanted regions to leave the region representing the possible number of tickets sold.



[4]

(c) The organizers want to maximize the number of children's tickets sold.
Find this maximum number of children's tickets and the corresponding number of adult tickets.

Children's tickets = ... , Adult tickets = ... **[2]**

492 A carpenter makes x tables and y chairs each week.
The total number of tables and chairs made each week must not exceed 60.
The number of tables made must be at least 10.
The number of chairs made must be at least half the number of tables made.


(a) Write down three inequalities, in terms of x and y, to show this information.

.....

.....

..... **[3]**

(b) On the grid, draw three straight lines and shade the unwanted regions to show the region which satisfies these inequalities.
Label the region R.



[4]

(c) The profit on each table is \$40 and the profit on each chair is \$25.
Work out the maximum possible weekly profit.

$\$$ ..... **[2]**

493 A clothing company manufactures two types of shirts: Standard and Premium.
In one day, they make x Standard shirts and y Premium shirts.
They can make a maximum of 60 shirts in total per day.
Each Standard shirt requires 1 metre of fabric and each Premium shirt requires 2 metres of fabric.
The company has a maximum of 80 metres of fabric available per day.
They must make at least 10 Premium shirts per day.


(a) Write down three inequalities in x and y, other than $x \geq 0$, to represent this information.

.....

.....

..... **[3]**

(b) On the grid provided, draw three lines to show these inequalities.
Shade the unwanted regions to leave the region that satisfies the inequalities unshaded.



[4]

(c) The profit on each Standard shirt is \$5 and the profit on each Premium shirt is \$8.
Find the maximum possible profit the company can make in one day.

$ ..... **[2]**

494 **(a)** Find the exact range of values of x for which

$$\frac{2x-3}{4} < \frac{x+5}{3} \leq \frac{3x-1}{2}$$

..... **[5]**

(b) A mobile phone company offers two plans.
Plan A costs \$12 per month plus \$0.07 per minute of calls.
Plan B costs \$25 per month for unlimited calls.

Let x be the number of minutes of calls in a month.

(i) Write down an inequality in terms of x to show the condition when Plan B is strictly cheaper than Plan A.

..... **[1]**

(ii) Solve your inequality to find the minimum number of whole minutes of calls required for Plan B to be cheaper than Plan A.

..... minutes **[3]**

495 **(a)** A factory makes x Mountain bicycles and y Road bicycles each day.

- The factory makes at most 60 bicycles in total each day.
- It makes at least 10 Road bicycles.
- It makes at least 15 Mountain bicycles.
- The number of Road bicycles is at most twice the number of Mountain bicycles.


Write down four inequalities, in terms of x and y, to show this information.

.....

.....

..... **[4]**

(b) On the grid, draw four lines to show these inequalities.
By shading the unwanted regions, indicate the region R which satisfies all four inequalities.



[5]

(c) The profit on each Mountain bicycle is \$30 and the profit on each Road bicycle is \$50.
Find the maximum possible profit the factory can make in one day, and the number of each type of bicycle it must make to achieve this profit.

Maximum profit = \$ .....

Mountain bicycles = ...

Road bicycles = ... **[3]**

496 Here are the first five terms of a number sequence.

7 11 15 19 23

(a) Write down the next term in this sequence.

..... **[1]**

(b) Find an expression, in terms of n, for the nth term of this sequence.

..... **[2]**

497 Here are the first four terms of a sequence.

5 11 17 23

(a) Find the next term of this sequence.

..... **[1]**

(b) Find an expression for the nth term of this sequence.

..... **[2]**

(c) Explain why 150 is not a term in this sequence.

..... **[1]**

498 Here are the first four terms of a sequence.

5 12 19 26

(a) Write down the next term in the sequence.

..... **[1]**

(b) Find an expression for the nth term of the sequence.

..... **[2]**

(c) Explain why 150 is not a term in this sequence.

..... **[1]**

499 Here are the first four terms of a sequence.

5 11 17 23

(a) Find the next term in this sequence.

..... **[1]**

(b) Find an expression for the nth term of this sequence.

..... **[2]**

(c) Explain why 150 is not a term in this sequence.

..... **[1]**

500 **(a)** Find the nth term of the sequence.

$$5 \quad 9 \quad 13 \quad 17 \quad 21$$

..... **[2]**

(b) Find the nth term of the sequence.

$$2 \quad 6 \quad 18 \quad 54 \quad 162$$

..... **[2]**

501 The first four terms of a sequence are

7 11 15 19

(a) Write down the next two terms of the sequence.

..... **[1]**

(b) Find an expression for the nth term of this sequence.

nth term = ... **[2]**

(c) Explain why 102 is not a term in this sequence.

..... **[2]**

502 **(a)** Sequence A is a linear sequence.

$$5 \quad 9 \quad 13 \quad 17 \quad \ldots$$

(i) Find the next two terms of sequence A.

..... **[1]**

(ii) Find the nth term of sequence A.

..... **[2]**


(b) Sequence B is a geometric sequence.

$$2 \quad 6 \quad 18 \quad 54 \quad \ldots$$

Find the nth term of sequence B.

..... **[2]**

503 The diagram shows a sequence of patterns made from small squares.



Pattern 1

Pattern 2

Pattern 3

(a) Find the number of squares in Pattern 4.

..... **[1]**

(b) Find an expression, in terms of n, for the number of squares in Pattern n.

..... **[2]**

(c) Which pattern has exactly 141 squares?

Pattern ..... **[2]**

504 The first four terms of a sequence are

$$-1 \quad 4 \quad 13 \quad 26$$

(a) Find an expression for the nth term of the sequence.

..... **[3]**

(b) Find the value of n when the nth term is 433.

$n =$ ..... **[3]**

505 The first five terms of a sequence, T_n, are given below.

$$4 \qquad 14 \qquad 34 \qquad 68 \qquad 120$$

(a) By finding the first, second and third differences, show that the nth term of the sequence is a cubic expression in n.

[2]

(b) Find an expression for T_n, the nth term of the sequence.

$T_n =$ ..... **[5]**

506 Here are the first five terms of a sequence.

3 9 17 27 39

Find an expression, in terms of n, for the nth term of this sequence.

..... **[3]**

507 Here are the first five terms of a sequence.

4 11 22 37 56

(a) Work out the next term of the sequence.

..... **[1]**

(b) Find an expression for the nth term of the sequence.

..... **[3]**

508 The nth term of a sequence is $u_n = an^2 + bn$.

(a) Given that $u_1 = 5$ and $u_2 = 14$, write down two simultaneous equations in a and b.

.....

..... **[2]**

(b) Solve these simultaneous equations to find the value of a and the value of b.

$a =$ ..... , $b =$ ..... **[3]**

509 **(a)** The nth term of a sequence is $n^2 - 3n + 5$.
Work out the first three terms of this sequence.

1st term = ... , 2nd term = ... , 3rd term = ...
[2]

(b) The first four terms of another sequence are

2 9 20 35

Find an expression for the nth term of this sequence.

..... **[3]**

510 The first four terms of a sequence, T, are

$$2 \quad 9 \quad 22 \quad 41$$

(a) Find an expression for the nth term, T_n.

$T_n =$ ..... **[3]**

(b) A second sequence is given by the formula $U_n = 2^n + 3$.
Work out the value of U_5.

$U_5 =$ ..... **[1]**

(c) Find the value of n for which $U_n = 131$.

$n =$ ..... **[2]**

511 The first four terms of a sequence are

$$-1 \quad 2 \quad 9 \quad 20$$

(a) Find the nth term of this sequence.

..... **[3]**

(b) The nth term of another sequence is given by $3^n - 2$.
Find the value of the term in this sequence that is closest to 1000.

..... **[3]**

512 The first five terms of a sequence are

$$2 \quad 9 \quad 20 \quad 35 \quad 54 \quad \ldots$$

(a) Find the next term.

..... **[1]**

(b) Find the nth term of the sequence.

..... **[3]**

(c) The kth term of the sequence is 464.
Find the value of k.

$k =$ ..... **[3]**

513 Consider the following three sequences, P, Q and R.
Sequence P:

$$2 \quad 9 \quad 28 \quad 65 \quad 126 \quad \ldots$$

Sequence Q:

$$3 \quad 8 \quad 15 \quad 24 \quad 35 \quad \ldots$$

Sequence R:

$$\frac{2}{3} \quad \frac{9}{8} \quad \frac{28}{15} \quad \frac{65}{24} \quad \frac{126}{35} \quad \ldots$$

(a) Find the nth term of sequence P.

..... **[2]**

(b) Find the nth term of sequence Q.

..... **[2]**

(c) Write down the nth term of sequence R.

..... **[1]**

(d) Find the 12th term of sequence R, giving your answer as a fraction in its simplest form.

..... **[2]**

514 Here are the first five terms of a sequence.

2 9 28 65 126

(a) Find the next term of this sequence.

..... **[1]**

(b) Find an expression for the nth term of this sequence.

..... **[3]**

(c) The kth term of the sequence is 8001.
Find the value of k.

$k =$ ..... **[3]**

515 The table shows the first five terms of sequences A, B and C.

n	1	2	3	4	5
Sequence A	2	5	10	17	26
Sequence B	1	8	27	64	125
Sequence C	-1	3	17	47	99

(a) Find the nth term of sequence A.

..... **[2]**

(b) Find the nth term of sequence B.

..... **[1]**

(c) Use your answers to parts (a) and (b) to find an expression, in terms of n, for the nth term of sequence C.
Give your answer in its simplest form.

..... **[2]**

(d) Determine which term in sequence A has a value of 10001.

$n =$ ..... **[2]**

516 **(a)** The first five terms of a sequence are

$$-1 \quad 8 \quad 33 \quad 80 \quad 155$$

Use the method of differences to find an expression for the nth term of this sequence.

..... **[4]**

(b) An exponential sequence has nth term $P_n = 5 \times 3^{n-1}$.
The kth term of this sequence is 3645.
Work out the value of k.

$k =$ ..... **[2]**

(c) The nth term of sequence T is $T_n = \dfrac{2n-1}{3n+2}$.
Write down the value that T_n approaches as n becomes very large.

..... **[1]**

517 **(a)** A sequence has the following first four terms:

1 15 53 127

Use the method of differences to find an expression for the nth term.

nth term = ... **[4]**

(b) Another sequence, V, has its nth term defined by

$$V_n = \frac{n^2 + 1}{2^n}$$

(i) Find the values of V_1, V_2 and V_3.
Give each fraction in its simplest form.

V_1 = ... , V_2 = ... , V_3 = ... **[2]**

(ii) Work out the exact value of $V_5 - V_4$.

..... **[3]**

518 **(a)** The first five terms of a sequence are

4 5 14 37 80

Use the method of differences to find an expression for the nth term.

..... **[4]**

(b) The nth term of a different sequence is given by $T_n = a \times 2^n + b$.
The first term of this sequence is 10 and the third term is 34.

Find the values of a and b.

$a =$ ..... , $b =$ ..... **[4]**

(c) Calculate the 12th term of the sequence described in part (b).

..... **[2]**

519 The first five terms of three sequences, A, B and C, are shown below.

Sequence A:	5	9	13	17	21
Sequence B:	2	6	20	50	102
Sequence C:	$\frac{5}{2}$	$\frac{3}{2}$	$\frac{13}{20}$	$\frac{17}{50}$	$\frac{7}{34}$

(a) Find the nth term of sequence A.

..... **[2]**

(b) **(i)** Work out the 6th term of sequence B.

..... **[1]**

(ii) Find an expression for the nth term of sequence B.

..... **[4]**

(c) **(i)** Work out the 6th term of sequence C, giving your answer as a fraction in its simplest form.

..... **[1]**

(ii) Find an expression for the nth term of sequence C.

..... **[2]**

520 y is directly proportional to the square of x.
When $x = 3$, $y = 36$.

(a) Find an equation connecting y and x.

$y =$ ..... **[2]**

(b) Find the value of y when $x = 5$.

$y =$ ..... **[1]**

521 y is directly proportional to $(x + 2)$.
When $x = 3$, $y = 20$.

Find the value of y when $x = 7$.

$y =$ ..... **[3]**

522 y is directly proportional to the square of x.
When $x = 3$, $y = 36$.

(a) Find an equation connecting y and x.

$y =$ ..... **[2]**

(b) Find the possible values of x when $y = 64$.

$x =$ ..... **[2]**

523 y is directly proportional to x.
When $x = 5$, $y = 15$.

(a) Find an equation connecting y and x.

..... **[2]**

(b) Work out the value of y when $x = 12$.

$y =$ ..... **[2]**

524 y is directly proportional to $(x+2)^3$.
When $x = 1$, $y = 135$.

(a) Find an equation for y in terms of x.

$y =$ ..... **[2]**

(b) Find the value of y when $x = 0$.

$y =$ ..... **[1]**

(c) Find the value of x when $y = 320$.

$x =$ ..... **[2]**

525 The variable y is directly proportional to the square of x.
When $x = 3$, $y = 36$.

(a) Find an equation connecting y and x.

$y =$ ..... **[2]**

(b) Find the value of y when $x = 5$.

$y =$ ..... **[1]**

(c) Find the values of x when $y = 196$.

$x =$ ..... **[2]**

526 y is directly proportional to the cube of x.
When $x = 2$, $y = 24$.

(a) Find an equation connecting y and x.

$y =$ ..... **[2]**

(b) Work out the value of y when $x = 5$.

$y =$ ..... **[1]**

(c) Work out the value of x when $y = 81$.

$x =$ ..... **[2]**

527 The table shows some corresponding positive values of p and q.

p	2	4	10
q	100	25	4

(a) Determine the algebraic equation connecting q and p.

$q =$ ..... **[2]**

(b) Find the value of q when $p = 5$.

$q =$ ..... **[1]**

(c) Find the value of p when $q = 1$.

$p =$ ..... **[2]**

528 P is directly proportional to the square root of Q.
Q is inversely proportional to R^3.
When $Q = 9$, $P = 12$.
When $R = 2$, $Q = 2$.

(a) Find an equation for P in terms of R.

$P =$ ..... **[5]**

(b) Calculate the value of P when $R = 4$.

$P =$ ..... **[2]**

529 A is directly proportional to the square of B.
B is inversely proportional to $(C + 2)$.
Assume that $B > 0$ and $(C + 2) > 0$.
When $B = 4$, $A = 80$.
When $C = 2$, $B = 3$.

(a) Find the value of A when $C = 4$.

$A =$ ..... **[4]**

(b) Find an expression for C in terms of A.

$C =$ ..... **[4]**

530 y is inversely proportional to the square of x.
When $x = 4$, $y = 3$.

(a) Find an equation connecting y and x.

..... **[3]**

(b) Work out the possible values of x when $y = 0.75$.

$x =$ ..... **[3]**

531 P is inversely proportional to the square of Q.
When $Q = 4$, $P = 1.5$.

(a) Find an equation connecting P and Q.

$P =$ ..... **[3]**

(b) Find the value of P when $Q = 10$.

$P =$ ..... **[1]**

(c) Find the positive value of Q when $P = 24$.

$Q =$ ..... **[2]**

532 The quantity P is inversely proportional to the square root of Q.
When $Q = 9$, $P = 12$.

(a) (i) Find the value of P when $Q = 144$.

$P =$ ..... **[3]**

(ii) Find the value of Q when $P = 1.5$.

$Q =$ ..... **[2]**

(b) Calculate the percentage decrease in P when the value of Q is multiplied by 4.

..... % **[2]**

533 M is inversely proportional to the square root of N.
When $N = 16$, $M = 15$.

(a) Find an equation connecting M and N.

$M =$ ..... **[2]**

(b) Find the value of M when $N = 100$.

$M =$ ..... **[2]**

(c) Find the value of N when $M = 1.2$.

$N =$ ..... **[3]**

534 The table shows some values of two variables, x and y.
The relationship between x and y can be modelled by a variation equation.

x	2	4	6
y	36	9	4

(a) Use the data to identify the correct variation model and find an equation connecting y and x.

$y =$ ..... **[3]**

(b) Find the positive value of x when $y = 2.25$.

$x =$ ..... **[2]**

(c) When x is multiplied by 3, y is multiplied by a constant p.
Find the value of p.

$p =$ ..... **[2]**

535 p is inversely proportional to the square root of q.
When $q = 144$, $p = 5$.

(a) Find an equation connecting p and q.

..... **[2]**

(b) Find the value of q when $p = 15$.

$q =$ ..... **[2]**

(c) The value of q is increased by 300%.
Find the percentage decrease in p.

..... % **[3]**

536 P is inversely proportional to the square root of Q.
When $Q = 36$, $P = 4$.

(a) Find an equation for P in terms of Q.

$P =$ ..... **[2]**

(b) Calculate the value of P when $Q = 144$.

$P =$ ..... **[1]**

(c) Calculate the value of Q when $P = 1.5$.

$Q =$ ..... **[2]**

(d) The value of Q is increased by 300%.
Calculate the percentage decrease in the value of P.

..... % **[3]**

537 **(a)** Some corresponding values of x and y are given below.

$$x = 2,\ y = 75 \qquad x = 4,\ y = 18.75 \qquad x = 10,\ y = 3$$

Identify which of the following models best fits the data:

$$y \propto x \qquad y \propto x^2 \qquad y \propto \frac{1}{x} \qquad y \propto \frac{1}{x^2}$$

Hence, find the equation for y in terms of x.

$y =$ ..... **[3]**

(b) P is directly proportional to the cube of Q.
When Q is increased by 50%, work out the percentage increase in P.

..... % **[3]**

538 **(a)** y is directly proportional to the cube of x.

When x increases by 20%, find the percentage increase in y.

..... % **[3]**

(b) The table shows some corresponding values of v and A.

Determine the best variation model connecting A and v from the following list:

$$A \propto v \qquad A \propto v^2 \qquad A \propto \sqrt{v} \qquad A \propto \frac{1}{v}$$

You must show working to justify your decision.

v	2	4	10
A	12	48	300

..... **[3]**

539 **(a)** The table shows some values of x and y.

x	2	4	5	8
y	32	8	5.12	2

(i) Identify the variation model and write down the equation for y in terms of x.

$y =$ ..... **[3]**

(ii) Calculate the value of y when $x = 10$.

$y =$ ..... **[1]**

(b) A is directly proportional to the cube of B.
When B is increased by $x\%$, the value of A increases by 72.8%.
Find the value of x.

$x =$ ..... **[3]**

540 **(a)** y is inversely proportional to the square root of x.

When x is increased by 44%, find the percentage change in y.

..... % **[4]**

(b) P is directly proportional to the cube of Q.
Q is inversely proportional to R.
When $R = 2$, $P = 10$.

Find the value of P when $R = 4$.

$P =$ ..... **[3]**

541 The table shows some corresponding values of x and y.

x	2	4	8
y	100	25	6.25

(a) Determine the relationship between x and y and find an equation connecting them.

..... **[3]**

(b) z is directly proportional to x.
When $x = 4$, $z = 10$.

(i) Find an equation for x in terms of z.

$x =$ ..... **[2]**

(ii) Hence find an equation for y in terms of z.
Give your answer in its simplest form.

$y =$ ..... **[3]**

542 The table shows some corresponding values of x and y.

x	2	4	10
y	40	5	0.32

(a) Determine the proportional relationship between y and x and find an equation connecting them.

$y =$ ..... **[3]**

(b) Find the value of x when $y = 2560$.


$x =$ ..... **[2]**

(c) The variable x is inversely proportional to w.
When $w = 5$, $x = 2$.

Find an equation connecting y and w.
Give your answer in its simplest form.

$y =$ ..... **[4]**

543 The diagram shows the graph of $y = a\sin(bx)$ for $0° \leq x \leq 360°$.




(a) Find the value of a.

$a =$ ..... **[1]**

(b) Find the value of b.

$b =$ ..... **[1]**

544 The diagram shows a sketch of the graph of $y = a\cos(bx)$ for $0° \leq x \leq 360°$.




(a) Write down the value of a.

$a =$ ..... **[1]**

(b) Find the value of b.

$b =$ ..... **[2]**

545 The diagram shows a sketch of the graph of the function $f(x) = ax^2 + c$.
The graph passes through the points $(0, -5)$ and $(2, 7)$.




(a) Write down the value of c.

$c =$ ..... **[1]**

(b) Find the value of a.

$a =$ ..... **[2]**

546 The diagram shows the sketch of the graph of $y = a\cos(bx)$ for $0° \leq x \leq 120°$.




(a) Write down the amplitude of the graph.

..... **[1]**

(b) Find the value of a and the value of b.

$a =$ ..... , $b =$ ..... **[2]**

547 The diagram shows the graph of the trigonometric function $f(x) = a\cos(bx)$ for $0° \leq x \leq 360°$.




(a) Write down the value of a and the value of b.

$a =$ ..... , $b =$ ..... **[2]**

(b) Write down the coordinates of the first minimum point where $x > 0$.

(..... , ...) **[1]**

548 The diagram shows the graph of $y = a\sin(bx)$ for $0° \leq x \leq 360°$.




(a) Find the value of a and the value of b.

$a =$ ..... , $b =$ ..... **[2]**

(b) Write down the period of the function.

..... ° **[1]**

549 Four graphs are shown below, labelled A, B, C, and D.



(a) Write down the letter of the graph that represents each function.

(i)

$$y = x^2 - 1$$

..... **[1]**

(ii)

$$y = 2^x$$

..... **[1]**

(iii)

$$y = \frac{1}{x}$$


..... **[1]**

(iv)

$$y = 2x - 1$$

..... **[1]**

550 The diagram shows the graph of $y = a\cos(bx)$ for $0° \leq x \leq 360°$.



(a) Write down the amplitude of the graph.

..... **[1]**


(b) Write down the period of the graph.

..... ° **[1]**

(c) Find the value of a and the value of b.

$a =$ ..... , $b =$ ..... **[2]**

551 The diagram shows the graph of $y = a\sin(bx)$ for $0° \leq x \leq 180°$.



(a) State the amplitude.

..... **[1]**


(b) State the period of the function.

..... ° **[1]**

(c) Write down the value of a and the value of b.

$a =$ ..... , $b =$ ..... **[2]**

552 The diagram shows the graph of $y = a\sin(bx)$ for $0° \leq x \leq 360°$.




(a) Find the value of a and the value of b.

$a =$ ..... , $b =$ ..... **[2]**

(b) The graph of $y = kx^3$ passes through the point $(2, -24)$.
Find the value of k.

$k =$ ..... **[2]**

553 The diagram shows the graph of $y = a\sin(bx)$ for $0° \leq x \leq 360°$.
The values of a and b are positive integers.



(a) Find the value of a and the value of b.

$a =$ ..... , $b =$ ..... **[2]**


(b) Write down the period of the function.

..... ° **[1]**

(c) State the number of solutions to the equation $a\sin(bx) = 2$ in the interval $0° \leq x \leq 360°$.

..... **[1]**

554 The diagram shows the graph of the function $y = x^2 + bx + c$.



(a) Write down the value of c.

$c =$ ..... **[1]**

(b) Find the value of b.

$b =$ ..... **[2]**


(c) Write down the coordinates of the minimum point of the graph.

(..... , ...) **[2]**

555 Six equations are shown below.

$$y = 2^x \qquad y = x^2 \qquad y = -x^2 \qquad y = x^3 \qquad y = \frac{1}{x} \qquad y = \cos x$$

The diagrams show four of these graphs, labelled A, B, C and D.




Write down the correct equation for each graph.

$A =$ ..... , $B =$ ..... , $C =$ ..... , $D =$ .....

[4]

556 The diagram shows a sketch of the graph of $y = x^2 + bx + c$.
The minimum point on the graph is $M(3, -4)$.




(a) Work out the value of b and the value of c.

$b =$ ..... , $c =$ ..... **[3]**

(b) Find the coordinates of the points where the graph crosses the x-axis.

..... **[2]**

557 The diagram shows a sketch of the graph of $f(x) = a\sin(bx)$ for $0° \leq x \leq 360°$.



(a) Write down the value of a.

$a =$ ..... **[1]**

(b) Write down the value of b.

$b =$ ..... **[1]**

(c) Solve the equation $f(x) = 1.5$ for $0° \leq x \leq 180°$.


$x =$ ..... **[3]**

558 The graph of the quadratic function $f(x) = ax^2 + bx + c$ has a line of symmetry at $x = 2$.
The graph passes through the points $(0, -5)$ and $(2, 3)$.

Find the values of a, b and c.

$a =$ ..... , $b =$ ..... , $c =$ ..... **[5]**

559 The diagram shows a sketch of the curve with equation $y = x^3 + ax^2 + bx$.
The curve passes through the points $Q(-1, 3)$ and $R(2, 0)$.



(a) Use the coordinates of Q and R to form two simultaneous equations in terms of a and b.

.....

..... **[2]**


(b) Solve your equations to find the value of a and the value of b.

$a =$ ..... , $b =$ ..... **[2]**

(c) Find the coordinates of the other two points where the curve crosses the x-axis.

..... **[3]**

560 The diagram shows the graph of $y = f(x)$ where $f(x) = ax^3 + bx^2$.
The graph passes through the origin $(0, 0)$, the point $(3, 0)$ and has a local minimum point at $(2, -4)$.



(a) Show that $3a + b = 0$.

[1]


(b) Find the value of a and the value of b.

$a =$ ..... , $b =$ ..... **[3]**

(c) Write down the range of values of k for which the equation $f(x) = k$ has exactly one solution.

..... **[2]**

561 The diagram shows a sketch of the graph of $y = f(x)$, where $f(x) = ax^2 + bx + c$.
The curve passes through the origin O and intersects the x-axis at the point A.
The point $M(3, 18)$ is the maximum point of the curve.



(a) Write down the value of c.

$c =$ ..... **[1]**

(b) Write down the coordinates of A.

(..... , ...) **[1]**


(c) Find the value of a and the value of b.

$a =$ ..... , $b =$ ..... **[3]**

(d) Find the range of values of k for which the equation $f(x) = k$ has two distinct positive real roots.

..... **[2]**

562 The diagram shows the graph of the cubic function $y = ax^3 + bx^2 + cx + d$.



(a) Write down the value of d.

$d =$ ..... **[1]**

(b) The graph intersects the x-axis at $x = -3$ and $x = 3$.
Show that $b = 0$ and $9a + c = 0$.

[3]

(c) The graph passes through the point $(1, -16)$.
Find the value of a and the value of c.


$a =$ ..... , $c =$ ..... **[3]**

(d) A different function $g(x) = px^2 + qx + r$ has a minimum point at $(0, -5)$ and passes through the point $(2, 7)$.

Find the values of p, q and r.

$p =$ ..... , $q =$ ..... , $r =$ ..... **[3]**

563 The diagram shows the graph of $y = a\sin(bx)$ for $0° \leq x \leq 360°$.



(a) Write down the value of a.

$a =$ ..... **[1]**


(b) Find the value of b.

$b =$ ..... **[1]**

(c) Calculate the value of y when $x = 30°$.
Give your answer correct to 3 significant figures.

$y =$ ..... **[2]**

564 The diagram shows the graph of $y = a\sin(bx)$ for $0° \leq x \leq 360°$.



(a) Find the value of a and the value of b.

$a =$ ..... , $b =$ ..... **[2]**


(b) The graph of $y = p^x$ passes through the point $(3, 125)$.
Find the value of p.

$p =$ ..... **[1]**

(c) The graph of $y = \dfrac{k}{x}$ passes through the point $(0.5, 8)$.
Find the value of k.

$k =$ ..... **[1]**

565 The diagram shows the graph of $y = \frac{a}{x} + b$.
The equation of the horizontal asymptote is $y = 2$.
The graph passes through the point $(1, 6)$.



(a) Write down the value of b.

$b =$ ..... **[1]**


(b) Find the value of a.

$a =$ ..... **[2]**

(c) The graph of $y = k^x$, where $k > 0$, also passes through the point $(2, 4)$, which lies on the curve $y = \frac{a}{x} + b$.
Find the value of k.

$k =$ ..... **[2]**

566 The diagram shows a sketch of the graph of $f(x) = \frac{a}{x} + b$, where $xe0$.
The dashed line is a horizontal asymptote.
The graph passes through the point $(2, 7)$.



(a) Write down the value of b.

$b =$ ..... **[1]**


(b) Find the value of a.

$a =$ ..... **[2]**

(c) Find the coordinates of the point where the graph crosses the x-axis.

(..... , ...) **[2]**

567 The diagram shows a sketch of the graph of $f(x) = a\cos(bx)$ for $0° \leq x \leq 180°$.



(a) Write down the value of a.

$a =$ ..... **[1]**


(b) Find the value of b.

$b =$ ..... **[2]**

(c) Determine the number of solutions to the equation $a\cos(bx) = 1.5$ in the interval $0° \leq x \leq 360°$.

..... **[2]**

568 The diagram shows the graph of a quadratic function $f(x) = ax^2 + bx + c$.
The graph crosses the y-axis at $(0, -4)$ and the x-axis at $(-1, 0)$ and $(4, 0)$.



(a) Write down the value of c.

$c =$ ..... **[1]**


(b) Determine the value of a and the value of b.

$a =$ ..... , $b =$ ..... **[3]**

(c) Calculate the exact coordinates of the minimum point of the graph.

(..... , ...) **[2]**

569 The diagram shows a sketch of the graphs of $y = a^x$ and $y = \frac{b}{x}$ for $x > 0$.
The graphs intersect at the point $P(3, 8)$.



(a) Find the value of a and the value of b.

$a =$ ..... , $b =$ ..... **[3]**


(b) The graph of $y = a^x$ intersects the y-axis at the point Q.
Write down the coordinates of Q.

(..... , ...) **[1]**

(c) Calculate the gradient of the straight line joining Q and P.


Gradient = ... **[2]**

570 Four graphs A, B, C and D are shown below.




Graph A

Graph B



Graph C



Graph D

(a) Write down the letter of the graph which could represent the function

(i) $f(x) = \dfrac{a}{x}$ where $a > 0$

..... **[1]**

(ii) $f(x) = a^x$ where $a > 1$

..... **[1]**

(iii) $f(x) = -ax^2$ where $a > 0$

..... **[1]**

(b) The equation of Graph B is $y = p^x$.
The graph passes through the point $(3, 125)$.
Find the value of p.

$p =$ ..... **[2]**


(c) The equation of Graph C is $y = \dfrac{q}{x}$.

The graph passes through the point $(0.5, 8)$.

Find the value of q.

$q =$ ..... **[2]**

571 The diagram shows the graph of $y = ax^2 + bx + c$.
The graph intersects the x-axis at $(-1, 0)$ and $(5, 0)$, and the y-axis at $(0, -5)$.



(a) Find the values of a, b and c.

$a =$ ..... , $b =$ ..... , $c =$ ..... **[3]**


(b) Write down the coordinates of the minimum point of the graph.

(..... , ...) **[2]**

(c) The graph of $y = p \times q^x$ passes through the points $(0, 2)$ and $(2, 18)$, where $q > 0$.
Find the value of p and the value of q.

$p =$ ..... , $q =$ ..... **[3]**

572 The diagram shows a sketch of the curve $f(x) = x^3 + ax^2 + bx + c$.
The curve crosses the y-axis at $(0, 4)$.
It crosses the x-axis at $(-1, 0)$ and touches the x-axis at $(2, 0)$.



(a) Write down the value of c.

$c =$ ..... **[1]**


(b) By using the x-intercepts, find the value of a and the value of b.

$a =$ ..... , $b =$ ..... **[4]**

(c) Use the graph to state the number of solutions to the equation $f(x) = 3$.

..... **[1]**

573 The diagram shows a sketch of the graph of $f(x) = x^3 + bx^2 + cx + d$.
The curve intersects the y-axis at 4 and the x-axis at -1 and 2.
The point $(2, 0)$ is a local minimum.



(a) Write down the value of d.

$d =$ ..... **[1]**


(b) Find the values of b and c.

$b =$ ..... , $c =$ ..... **[4]**

(c) Write down the number of solutions to the equation $f(x) = -3$.

Number of solutions = ... **[1]**

574 The diagram shows the graph of $y = x^3 + bx^2 + cx + d$.
The graph crosses the axes at $(-2, 0)$, $(2, 0)$, $(p, 0)$ and $(0, 12)$.



(a) Write down the value of d.

$d =$ ..... **[1]**


(b) Find the values of b and c.

$b =$ ..... , $c =$ ..... **[4]**

(c) Find the value of p.

$p =$ ..... **[2]**

575 The diagram shows a sketch of the graph of $f(x) = ax^3 + bx$.
The curve passes through the points $P(2, -16)$ and $Q(-3, 9)$.



(a) Find the value of a and the value of b.

$a =$ ..... , $b =$ ..... **[3]**


(b) Find the coordinates of the local maximum point of the graph.

(..... , ...) **[2]**

(c) The equation $f(x) = k$ has exactly one solution.
Write down the range of values for k.

..... **[2]**

576 The diagram shows a sketch of the curve $y = ax^3 + bx$.
The curve passes through the origin O, and the points $(2, 10)$ and $(-1, 4)$.
The curve crosses the x-axis at the points A, O, and B.




(a) Find the value of a and the value of b.

$a =$ ..... , $b =$ ..... **[4]**

(b) Calculate the exact length of the line segment AB.

..... **[3]**

577 The diagram shows the graph of the exponential function $f(x) = a \cdot b^x + c$.
The horizontal line $y = -3$ is an asymptote to the graph.
The graph passes through the points $(0, -1)$ and $(1, 3)$.



(a) State the value of c.

$c =$ ..... **[1]**

(b) Work out the value of a and the value of b.


$a =$ ..... , $b =$ ..... **[4]**

(c) The graph of $g(x) = \dfrac{k}{x}$ intersects the graph of $f(x)$ at the point where $x = 1$.

Find the value of k.

$k =$ ..... **[2]**

578 The diagram shows the graph of a cubic function $f(x) = ax^3 + bx^2 + cx + d$.
The graph crosses the x-axis at $(-3, 0)$, $(1, 0)$ and $(4, 0)$.
It crosses the y-axis at $(0, -24)$.



(a) Find the values of a, b, c and d.


$a =$ ..... , $b =$ ..... , $c =$ ..... , $d =$ ..... **[6]**

(b) The graph of $y = f(x)$ is translated by the vector $\begin{pmatrix} 0 \\ 10 \end{pmatrix}$ to give the graph of $y = g(x)$.

Find the coordinates of the y-intercept of the graph of $y = g(x)$.

(..... , ...) **[2]**

579 The diagram shows part of the graph of $y = \frac{a}{x} + b$ for $x > 0$.
The graph passes through the points $(1, 8)$ and $(4, 3.5)$.



(a) Find the value of a and the value of b.

$a =$ ..... , $b =$ ..... **[3]**

(b) Write down the equation of the horizontal asymptote to the graph.

..... **[1]**

(c) The line $y = x - 3$ intersects the graph of $y = \frac{6}{x} + 2$ at the point P.


(i) Show that the x-coordinate of P satisfies the equation $x^2 - 5x - 6 = 0$.

[2]

(ii) Find the coordinates of P.

(..... , ...) **[2]**

580 The diagram shows a sketch of the graph of $f(x) = x^3 + bx^2 + cx + 4$.
The graph crosses the x-axis at $(-1, 0)$ and touches the x-axis at $(2, 0)$.
The graph crosses the y-axis at $(0, 4)$.



(a) **(i)** Use the point $(-1, 0)$ to show that $b - c = -3$.

[2]


(ii) Find the values of b and c.

$b =$ ..... , $c =$ ..... **[4]**

(b) The graph of $g(x) = p \cdot q^x$ passes through the points $(0, 2)$ and $(3, 54)$.
Given that $q > 0$, work out the values of p and q.

$p =$ ..... , $q =$ ..... **[3]**

581 The diagram shows the graphs of $y = f(x)$ and $y = g(x)$ for $x \geq 0$.
It is given that $f(x) = px^2 + qx + r$.
The graph of $y = f(x)$ has a maximum point at $M(2, 9)$ and intersects the y-axis at $P(0, 5)$.



(a) Find the values of p, q, and r.

$p =$ ..... , $q =$ ..... , $r =$ ..... **[4]**

(b) The graph of $y = f(x)$ intersects the positive x-axis at the point R.
Find the coordinates of R.

(..... , ...) **[2]**


(c) It is given that $g(x) = a\sin(bx^\circ)$, where $a > 0$ and $b > 0$.
The graph of $y = g(x)$ has its first positive x-intercept at the point R.
The amplitude of $g(x)$ is equal to the y-coordinate of P.

Find the value of a and the value of b.

$a =$ ..... , $b =$ ..... **[3]**

582 Consider the function $f(x) = x^2 - 4x + 1$ for $-1 \leq x \leq 5$.

(a) Use your graphic display calculator to sketch the graph of $y = f(x)$ on the axes provided.



[2]

(b) Write down the coordinates of the vertex of the graph.


(..... , ...) **[1]**

(c) Find the zeros of $f(x)$.
Give your answers correct to 2 decimal places.

$x =$ ..... , $x =$ ..... **[2]**

583 Let $f(x) = -x^2 + 6x - 5$.

(a) On the axes provided, sketch the graph of $y = f(x)$ for $0 \leq x \leq 6$.



[2]

(b) Find the zeros of $f(x)$.


$x =$ ..... , $x =$ ..... **[2]**

(c) Find the coordinates of the vertex of the graph.

(..... , ...) **[2]**

584 Let $f(x) = -2x^2 + 5x + 3$.

(a) On the axes, sketch the graph of $y = f(x)$ for $-1.5 \leq x \leq 4$.



[2]

(b) Find the coordinates of the local maximum.


(..... , ...) **[2]**

(c) Find the zeros of $f(x)$.

$x =$ ..... , $x =$ ..... **[2]**

585 Consider the function $f(x) = x^3 - 3x^2 + 2$ for $-1 \leq x \leq 3$.

(a) On the grid, sketch the graph of $y = f(x)$.



[2]

(b) Find the zeros of $f(x)$.


$x =$ ..... **[2]**

(c) Find the coordinates of the local minimum point of the graph.

(..... , ...) **[2]**

586 A function is given by $f(x) = 2x^2 - 5x - 3$ for the domain $-2 \leq x \leq 4$.

(a) On the axes, sketch the graph of $y = f(x)$.



[2]

(b) Write down the zeros of $f(x)$.


$x =$ ..... , $x =$ ..... **[2]**

(c) Write down the coordinates of the local minimum.

(..... , ...) **[2]**

587 Consider the function $f(x) = \frac{x^3}{4} - 2x + 1$ for $-3.5 \leq x \leq 3.5$.

(a) On the grid, sketch the graph of $y = f(x)$.



[2]

(b) Find the coordinates of the local maximum point and the local minimum point.
Give your answers correct to 2 decimal places.


Local maximum = ... , Local minimum = ... **[2]**

(c) Write down the range of values of k for which the equation $f(x) = k$ has three distinct real solutions.

..... **[2]**

588 Let $f(x) = x^3 - 3x + 1$.

(a) On the axes provided, sketch the graph of $y = f(x)$ for $-2.5 \leq x \leq 2.5$.



[2]

(b) Find the coordinates of the local maximum point.

(..... , ...) **[2]**


(c) On the same axes, sketch the graph of $y = 2x - 1$. **[1]**

(d) Find the x-coordinates of the points of intersection of the two graphs.

$x =$ ..... , $x =$ ..... , $x =$ ..... **[2]**

589 Let $f(x) = \dfrac{2}{x+1}$ and $g(x) = x - 2$ for $-5 \leq x \leq 5$.

(a) On the grid, sketch the graphs of $y = f(x)$ and $y = g(x)$.



[3]

(b) Write down the equation of the vertical asymptote of the graph of $y = f(x)$.


..... **[1]**

(c) Find the coordinates of the points of intersection of the two graphs.

Point 1 = ... , Point 2 = ... **[3]**

590 Let $f(x) = x^3 - 4x^2 + x + 2$ for the domain $-1 \leq x \leq 4$.

(a) On the axes, sketch the graph of $y = f(x)$.



[2]

(b) Write down the x-coordinates of the points where $f(x) = 0$.

$x =$ ..... , $x =$ ..... , $x =$ ..... **[3]**

(c) Write down the coordinates of the local minimum.

(..... , ...) **[2]**


(d) On the same axes, sketch the graph of $g(x) = -2x + 1$. **[1]**

(e) Find the x-coordinates of the points of intersection of the graphs of $f(x)$ and $g(x)$.

$x =$ ..... , $x =$ ..... , $x =$ ..... **[3]**

591 Let $f(x) = x^3 - 3x + 2$.

(a) On the axes, sketch the graph of $y = f(x)$ for $-2.5 \leq x \leq 2.5$.



[3]

(b) Find the coordinates of the local maximum.

(..... , ...) **[2]**

(c) Find the zeros of $f(x)$.

$x =$ ..... , $x =$ ..... **[2]**


(d) On the same diagram, sketch the graph of $y = 2x + 1$. **[2]**

(e) Find the x-coordinates of the points of intersection of $y = f(x)$ and $y = 2x + 1$.

..... **[3]**

592 Let $f(x) = 3x \cdot 2^{-x}$ and $g(x) = 0.5x + 0.2$ for $0 \leq x \leq 5$.

(a) On the axes provided, sketch the graph of $y = f(x)$ and the graph of $y = g(x)$ for $0 \leq x \leq 5$.



[3]

(b) Find the coordinates of the local maximum of $y = f(x)$.
Give your answer correct to 2 decimal places.


(..... , ...) **[2]**

(c) Solve the inequality $f(x) > g(x)$ for $0 \leq x \leq 5$.
Give your answer correct to 2 decimal places.

..... **[2]**

593 The function $f(x)$ is defined as $f(x) = \dfrac{x^2}{2^x}$ for the domain $-1 \leq x \leq 5$.

(a) On the grid, sketch the graph of $y = f(x)$.



[3]

(b) Find the coordinates of the local maximum point.

(..... , ...) **[2]**

(c) Find the x-coordinates of the points where $f(x) = 0.5$.


$x =$ ..... **[2]**

(d) Write down the range of values of k for which the equation $f(x) = k$ has exactly 3 different solutions.

..... **[2]**

594 The function $f(x)$ is defined as $f(x) = 3x^4 - 8x^3 - 6x^2 + 24x$ for the domain $-2 \leq x \leq 3$.

(a) On the axes provided, sketch the graph of $y = f(x)$.



[3]

(b) Write down the coordinates of the two local minima.

.....

..... **[2]**

(c) Write down the coordinates of the local maximum.

(..... , ...) **[1]**

(d) The equation $f(x) = k$ has exactly four distinct solutions.
Write down the range of values for k.


..... **[2]**

(e) The graph of $y = f(x)$ is translated by the vector $\begin{pmatrix} 0 \\ 5 \end{pmatrix}$.
Write down the coordinates of the local maximum of the translated graph.

(..... , ...) **[1]**

595 Let $f(x) = \dfrac{x^2 - 2x + 4}{x - 2}$.

(a) On the axes provided, sketch the graph of $y = f(x)$ for $-2 \leq x \leq 6, x \neq 2$.



[3]

(b) Write down the equation of the vertical asymptote.

..... **[1]**

(c) Find the coordinates of the local maximum point.

(..... , ...) **[2]**

(d) Find the coordinates of the local minimum point.


(..... , ...) **[2]**

(e) The equation $f(x) = k$ has no real solutions.
Write down the range of values of k.

..... **[2]**

596 Let $f(x) = \dfrac{x^3 + 2}{x}$.

(a) On the axes, sketch the graph of $y = f(x)$ for $-3 \leq x \leq 3$.



[3]

(b) Find the coordinates of the local minimum.

(..... , ...) **[2]**

(c) Find the zero of $f(x)$.

$x =$ ..... **[1]**

(d) Write down the equation of the vertical asymptote.

..... **[1]**

(e) Solve the equation $f(x) = 4$.

..... **[3]**

597 Functions f and g are defined as follows.

$f(x) = 2x - 5$, $g(x) = x^2 + 1$


(a) Find the value of $f(g(3))$.

..... **[2]**

(b) Find $f^{-1}(x)$.

$f^{-1}(x) =$ ..... **[2]**

598 The mapping diagram shows the function $f(x) = 3x - 4$ for the domain $\{-1, 0, 2, 5\}$.



(a) Work out the value of k.

$k =$ ..... **[1]**

(b) Write down the range of f.

..... **[1]**

(c) Find $f^{-1}(x)$.

$f^{-1}(x) =$ ..... **[2]**

599 It is given that $f(x) = 3x - 2$ and $g(x) = \dfrac{10}{x}$.

(a) Find $f(4)$.

..... **[1]**

(b) Find x when $g(x) = 2$.

$x =$ ..... **[1]**

(c) Find $f^{-1}(x)$.

$f^{-1}(x) =$ ..... **[2]**

600 The functions f and g are defined as $f(x) = 3x - 2$ and $g(x) = x^2 + 4$.

(a) Work out

(i)

hspace*0.8cm $f(-4)$

..... **[1]**

(ii)

hspace*0.8cm $gf(2)$

..... **[2]**

(b) Find $f^{-1}(x)$.

$f^{-1}(x) =$ ..... **[2]**

601 $f(x) = 4x - 5$
$g(x) = \dfrac{x}{3} + 2$

(a) Work out $f(-2)$.

..... **[1]**

(b) Find $f^{-1}(x)$.

$f^{-1}(x) =$ ..... **[2]**

(c) Find $fg(x)$ in its simplest form.

$fg(x) =$ ..... **[2]**

602 The functions f and g are defined as $f(x) = 5x - 2$, $g(x) = x^2 + 3$.

(a) Find the value of $f(4)$.

..... **[1]**

(b) Find $f^{-1}(x)$.

$f^{-1}(x) =$ ..... **[2]**

(c) Find $fg(2)$.

..... **[2]**

603 The functions f and g are defined as $f(x) = 4x - 5$, $g(x) = \dfrac{12}{x}$.

(a) Work out the value of $f(3)$.

..... **[1]**

(b) Find $f^{-1}(x)$.

$f^{-1}(x) =$ ..... **[2]**

(c) Work out $fg(2)$.

..... **[2]**

604 Consider the functions $f(x) = 3x - 2$ and $g(x) = x^2 + 4$.

(a) Find $f(4)$.

..... **[1]**

(b) Find $f^{-1}(x)$.

$f^{-1}(x) =$ ..... **[2]**

(c) Find $gf(2)$.

..... **[2]**

605 Let $f(x) = 5 - 2x$, $g(x) = x^2 - 3$.

(a) Work out $f(-4)$.

..... **[1]**

(b) Work out $gf(2)$.

..... **[2]**

(c) Find $f^{-1}(x)$.

$f^{-1}(x) =$ ..... **[2]**

606 $f(x) = 4x - 3$, $g(x) = 2 - 5x$

(a) Work out $f(2)$.

..... **[1]**

(b) Find $f^{-1}(x)$.

$f^{-1}(x) =$ ..... **[2]**

(c) Work out $gf(1)$.

..... **[2]**

607 Let $f(x) = 4x - 3$ and $g(x) = 2x^2 + 1$.

(a) Work out $fg(3)$.

..... **[2]**

(b) Find $f^{-1}(x)$.

$f^{-1}(x) =$ ..... **[2]**

(c) Solve the equation

$$f(x) = 17$$

$x =$ ..... **[2]**

608 Two functions, f and g, are defined as follows:

$$f(x) = 3x - 2 \qquad g(x) = x^2 + 1$$

(a) Work out $f(4)$.

..... **[1]**

(b) Find $f^{-1}(x)$.

$f^{-1}(x) =$ ..... **[2]**

(c) Find $gf(x)$ in its simplest form.

$gf(x) =$ ..... **[3]**

609 The functions f and g are defined as $f(x) = \dfrac{3x+2}{x-1}$, $x \neq 1$, $g(x) = x - 2$.

(a) Find $f^{-1}(x)$.

$f^{-1}(x) =$ ..... **[3]**

(b) Solve the equation.

$f(x) = g(x)$

$x =$ ..... , $x =$ ..... **[4]**

610 Consider the functions $h(x) = \dfrac{12}{x-1}$ and $k(x) = 2x + 3$.

(a) State the value of x which must be excluded from the domain of $h(x)$.

$x =$ ..... **[1]**

(b) Find $k^{-1}(x)$.

$k^{-1}(x) =$ ..... **[2]**

(c) Find $hk(x)$, giving your answer in its simplest form.

$hk(x) =$ ..... **[2]**

(d) Solve the equation.

$h(x) = k(2)$

$x =$ ..... **[3]**

611 Let $f(x) = \dfrac{4}{2x-3}$ for $x \neq 1.5$, and $g(x) = x + 5$.

(a) Find $f^{-1}(x)$.

$f^{-1}(x) =$ ..... **[3]**

(b) Solve the equation $f(x) = g(-3)$.

$x =$ ..... **[3]**

(c) Express $gf(x)$ as a single fraction in its simplest form.

$gf(x) =$ ..... **[3]**

612 The functions f, g and h are defined as $f(x) = x^2 - 2$, $g(x) = \dfrac{x-3}{2}$, $h(x) = 5 - 3x$.

(a) Find $h^{-1}(x)$.

$h^{-1}(x) =$ ..... **[2]**

(b) Solve the equation

$f(x) = 34$

$x =$ ..... , $x =$ ..... **[2]**

(c) Find $hg(x)$ in its simplest form.

..... **[3]**

(d) Work out $fh(2)$.

..... **[2]**

613 The functions f and g are defined as $f(x) = \dfrac{x+3}{x-1}$, where $x \neq 1$, and $g(x) = 2x - 5$.

(a) Find $f^{-1}(x)$.

$f^{-1}(x) =$ ..... **[3]**

(b) Show that $ff(x) = x$.

[3]

(c) Solve the equation $g(x^2) = f(3)$.

$x =$ ..... **[4]**

614 $f(x) = \dfrac{3x+2}{x-1}$, $g(x) = \dfrac{2}{x+1}$

(a) Find $f^{-1}(x)$.

$f^{-1}(x) =$ ..... **[3]**

(b) Find $fg(x)$, giving your answer as a single fraction in its simplest form.

..... **[3]**

(c) Solve the equation.

$f(x) = f^{-1}(x)$

Give your answers in the form $a \pm \sqrt{b}$, where a and b are integers.

$x =$ ..... **[4]**

615 The functions f and g are defined as follows:

$f(x) = \dfrac{3x+5}{x-1}, x \neq 1$, and $g(x) = x^2 + 1$

(a) Find $f^{-1}(x)$.

$f^{-1}(x) =$ ..... **[4]**

(b) Solve the equation $f(x) = x$.

$x =$ ..... , $x =$ ..... **[4]**

(c) Find $gf(2)$.

$gf(2) =$ ..... **[2]**

616 $f(x) = \dfrac{4}{x-1}$ for $xe1$, $g(x) = 2x + 3$, $h(x) = x^2 - 4$.

(a) Find an expression, in its simplest form, for

(i) $fg(x)$,

$fg(x) =$ ..... **[2]**

(ii) $fh(x) - 1$ as a single fraction.

..... **[3]**

(b) Find $f^{-1}(x)$.

$f^{-1}(x) =$ ..... **[3]**

(c) Solve the equation $gf(x) = 5$.

$x =$ ..... **[3]**

617 The functions h and k are defined as $h(x) = \dfrac{2x+1}{3}$ and $k(x) = 5 - x$.

(a) Find $h^{-1}(x)$.

$h^{-1}(x) =$ ..... **[2]**

(b) Find an expression for $hk(x)$ in its simplest form.

$hk(x) =$ ..... **[2]**

(c) Solve the equation $h(x) = k(x)$.

$x =$ ..... **[3]**

618 Functions h and k are defined as follows.

$h(x) = \dfrac{4}{x-3}$ for $x \neq 3$

$k(x) = 2x + 1$

(a) Find $k(h(5))$.

..... **[2]**

(b) Find $h^{-1}(x)$.

$h^{-1}(x) =$ ..... **[2]**

(c) Solve the equation $h(k(x)) = 2$.

$x =$ ..... **[3]**

619 The functions f, g and h are defined as follows:

$f(x) = 2x + 3$, $g(x) = x^2 - 4$, $h(x) = 5^x$

(a) Find $h(-1)$ as a fraction in its simplest form.

..... **[1]**

(b) Find $gf(x)$ in its simplest form.

$gf(x) =$ ..... **[3]**

(c) Find $f^{-1}(x)$.

$f^{-1}(x) =$ ..... **[2]**

(d) Find the value of x when $h(x) = 125$.

$x =$ ..... **[1]**

620 Two functions, f and g, are defined as follows:

$$f(x) = \frac{5}{x-2} \qquad g(x) = x + 2$$

(a) State the value of x that must be excluded from the domain of $f(x)$.

$x =$ ..... **[1]**

(b) Find $f^{-1}(x)$.

$f^{-1}(x) =$ ..... **[3]**

(c) Solve the equation $f(x) = g(x)$.

$x =$ ..... , $x =$ ..... **[4]**

621 $f(x) = 2x - 3$, $g(x) = x^2 - x - 7$

(a) Work out $g(-2)$.

..... **[2]**

(b) Find $gf(x)$ in its simplest form.


..... **[3]**

(c) Solve the equation.

$f(x) = g(x)$

$x =$ ..... , $x =$ ..... **[3]**

622 The diagram shows the graph of $y = f(x)$ for $-2 \leq x \leq 4$, where $f(x) = x^2 - 2x - 3$.
The function g is defined by $g(x) = 2x + 1$.



(a) Use the graph to write down the range of f for $-2 \leq x \leq 4$.

..... **[2]**

(b) Find $fg(3)$.

$fg(3) =$ ..... **[2]**

(c) Find $gf(x)$, giving your answer in the form $ax^2 + bx + c$.

$gf(x) =$ ..... **[2]**

(d) Solve the equation $g^{-1}(x) = 6$.

$x =$ ..... **[2]**

623 Let $f(x) = \dfrac{2x+1}{3}$ and $g(x) = \dfrac{5}{x-2}$, where $x \neq 2$.

(a) Find $gf(x)$.
Give your answer as a single fraction in its simplest form.

..... **[3]**

(b) Find $g^{-1}(x)$.

$g^{-1}(x) =$ ..... **[3]**

(c) Solve the equation

$$f(x) = g(3)$$

$x =$ ..... **[3]**

624 $f(x) = 2x - 1$
$g(x) = x^2 - 4$
$h(x) = \dfrac{2x+1}{x-3}, \quad x \neq 3$

(a) Work out $g(-3)$.

..... **[1]**

(b) Find $h^{-1}(x)$.

$h^{-1}(x) =$ ..... **[3]**

(c) Find $gf(x)$ in the form $ax^2 + bx + c$.

$gf(x) =$ ..... **[3]**

(d) Solve the equation $f(x) = g(x)$.

$x =$ ..... , $x =$ ..... **[3]**

625 The functions f, g and h are defined as $f(x) = 4 - 3x$, $g(x) = \dfrac{5}{x-2}$, $x \neq 2$, $h(x) = x^2 - x - 7$.

(a) (i) Find the value of $f(0.5)$.

..... **[1]**

(ii) Find $gf(x)$ in its simplest form.

$gf(x) =$ ..... **[2]**

(b) Find the range of $h(x)$.

..... **[2]**

(c) Solve the equation.

$h(x) = 2x - 1$ Give your answers correct to 2 decimal places.

$x =$ ..... , $x =$ ..... **[4]**

626 Functions f, g and h are defined as follows.

$f(x) = 3x - 2$

$g(x) = x^2 - 4x$

$h(x) = \dfrac{2x+1}{x-1}$ for $x \neq 1$

(a) Find $h^{-1}(x)$.

$h^{-1}(x) =$ ..... **[3]**

(b) Find an expression for $g(f(x))$ in the form $ax^2 + bx + c$.

..... **[3]**

(c) Solve the equation $f(g(x)) = x$.
Give your answers correct to 2 decimal places.
You must show all your working.

$x =$ ..... , x =

..... **[4]**

627 Let $f(x) = 3x + 2$, $g(x) = x^2 - x$ and $h(x) = 5^x$.

(a) Find $fg(x)$.
Give your answer in the form $ax^2 + bx + c$.

..... **[2]**

(b) Find $gf(x)$.
Give your answer in the form $ax^2 + bx + c$.

..... **[3]**

(c) Solve the equation

$$gf(x) = 0$$

$x =$ ..... , $x =$ ..... **[3]**

(d) Find the value of x when $h(x) = \frac{1}{125}$.

$x =$ ..... **[2]**

628 $f(x) = \dfrac{3}{x+2}, \quad x \neq -2$
$g(x) = 5x - 4$

(a) Find $g^{-1}(x)$.

$g^{-1}(x) =$ ..... **[2]**

(b) Find $fg(x)$ as a single fraction in its simplest form.

$fg(x) =$ ..... **[2]**

(c) Find $ff(x)$ as a single fraction in its simplest form.

$ff(x) =$ ..... **[3]**

(d) Solve the equation $f(x) = g(x)$.
Show all your working.

$x =$ ..... , $x =$ ..... **[4]**

629 Let $f(x) = \dfrac{2x+5}{x-3}$ for $x \neq 3$, and $g(x) = 3x - 1$.

(a) Find $f^{-1}(x)$.

$f^{-1}(x) =$ ..... **[3]**

(b) Express $fg(x)$ as a single fraction in its simplest form.

$fg(x) =$ ..... **[3]**

(c) Solve the equation $gf(x) = x$.
Show all your working and give your answers correct to 2 decimal places.

$x =$ ..... , $x =$ ..... **[5]**

630 Consider the functions $f(x) = \dfrac{3}{x+2}$, $g(x) = (3x+5)^2$ and $h(x) = \dfrac{3x+5}{x-2}$.

(a) Find $h^{-1}(x)$.

$h^{-1}(x) =$ ..... **[3]**

(b) Find $fg(x)$.
Give your answer as a single fraction in its simplest form.

$fg(x) =$ ..... **[3]**

(c) Solve the equation.

$h(x) = x$

Show all your working and give your answers correct to 2 decimal places.

$x =$ ..... , or $x =$ ..... **[5]**

631 The functions f and g are defined as:

$$f(x) = \frac{2x+3}{x-1} \qquad g(x) = x^2 - 4x$$

(a) Find $f^{-1}(x)$.

$f^{-1}(x) =$ ..... **[4]**

(b) Find $ff(x)$, giving your answer as a single fraction in its simplest form.

$ff(x) =$ ..... **[4]**

(c) Solve the equation $g(x) = f(2)$.
Show all your working and give your answers correct to 2 decimal places.

$x =$ ..... , $x =$ ..... **[4]**

632 The functions f and g are defined as $f(x) = \dfrac{x+5}{x-2}$, $g(x) = x^2 + 3x$.

(a) State the value of x for which $f(x)$ is undefined.

$x =$ ..... **[1]**

(b) Find $f^{-1}(x)$.

$f^{-1}(x) =$ ..... **[4]**

(c) Solve the equation

$g(x) = 10$

$x =$ ..... , $x =$ ..... **[3]**

(d) Solve the equation

$f(x) = x$

Show all your working and give your answers correct to 2 decimal places.

$x =$ ..... , $x =$ ..... **[4]**

633 A quadratic function has x-intercepts at $x = -3$ and $x = 4$.
The coefficient of x^2 is 1.

Write down the equation of this function in the form $y = ax^2 + bx + c$.

$y =$ ..... **[2]**

634 A quadratic function has x-intercepts at $x = 4$ and $x = -2$.
The coefficient of x^2 is 1.

Find the equation of the quadratic function in the form $y = x^2 + bx + c$.

$y =$ ..... **[2]**

635 A quadratic function $f(x)$ has x-intercepts at $x = -4$ and $x = 3$.
The coefficient of x^2 is 1.

Find the equation of the function in the form $f(x) = x^2 + bx + c$.

$f(x) =$ ..... **[2]**

636 A quadratic function has x-intercepts at $x = -4$ and $x = 7$.
The coefficient of x^2 is 1.
Write down the equation of the function.
Give your answer in the form $y = x^2 + bx + c$.

$y =$ ..... **[2]**

637 A quadratic function has the equation $y = x^2 + bx + c$.
The vertex of the graph of this function is $(4, -3)$.

Find the value of b and the value of c.

$b =$ ..... , $c =$ ..... **[3]**

638 A quadratic function is given by

$$y = x^2 + bx + c$$

The vertex of the graph of this function is $(3, -4)$.

Find the value of b and the value of c.

$b =$ ..... , $c =$ ..... **[3]**

639 A quadratic function has a minimum point at $(3, -4)$ and passes through the point $(5, 0)$.
The coefficient of x^2 is 1.

Find the equation of the function in the form $y = x^2 + bx + c$.

$y =$ ..... **[3]**

640 The graph of a quadratic function, $y = x^2 + bx + c$, crosses the x-axis at $(-4, 0)$ and $(2, 0)$.

(a) Work out the value of b and the value of c.

$b =$ ..... , $c =$ ..... **[2]**

(b) Find the coordinates of the vertex of the graph.

(..... , ...) **[2]**

641 A quadratic function $y = f(x)$ has its vertex at $(3, -4)$.
The coefficient of x^2 is 1.

(a) Find the equation of the function in the form $y = x^2 + bx + c$.

$y =$ ..... **[2]**

(b) Find the coordinates of the points where the graph of $y = f(x)$ crosses the x-axis.

..... **[2]**

642 A quadratic function is given by the equation $y = (x - h)^2 + k$.
The vertex of the graph of this function is $(3, -2)$.


(a) Write down the value of h and the value of k.

$h =$ ..... , $k =$ ..... **[2]**

(b) The graph passes through the point $(5, p)$.
Work out the value of p.

$p =$ ..... **[2]**

643 The diagram shows a sketch of the quadratic curve $y = x^2 + bx + c$.
The vertex of the curve is at $(3, -4)$.



(a) Write down the equation of the curve in the form $y = (x - h)^2 + k$.

$y =$ ..... **[2]**


(b) Find the values of b and c.

$b =$ ..... , $c =$ ..... **[2]**

(c) Write down the coordinates of the point where the curve crosses the y-axis.

(..... , ...) **[1]**

644 The diagram shows the graph of a quadratic function $y = x^2 + px + q$.
The graph has a minimum point (vertex) at $M(2, -9)$.



(a) Find the value of p and the value of q.

$p =$ ..... , $q =$ ..... **[3]**

(b) The graph crosses the x-axis at the points A and B.
Find the coordinates of A and the coordinates of B.


$A =$ ..... , $B =$ ..... **[3]**

645 The graph of a quadratic function $y = f(x)$ intersects the x-axis at $(-2, 0)$ and $(5, 0)$.
The graph intersects the y-axis at $(0, -20)$.

Work out the function $f(x)$, giving your answer in the form $ax^2 + bx + c$.

$f(x) =$ ..... **[4]**


646 The diagram shows a sketch of the quadratic graph $y = f(x)$.
The graph crosses the x-axis at $(-2, 0)$ and $(6, 0)$.
It crosses the y-axis at $(0, -24)$.



Find the expression for $f(x)$ in the form $ax^2 + bx + c$.

$f(x) =$ ..... **[4]**

647 The diagram shows a sketch of the quadratic function $y = f(x)$.
The graph crosses the x-axis at $(-1, 0)$ and $(4, 0)$.
The graph crosses the y-axis at $(0, -8)$.




(a) Find the equation of the quadratic function in the form $y = ax^2 + bx + c$.

$y =$ ..... **[3]**

(b) Find the coordinates of the minimum point of the graph.

(..... , ...) **[3]**

648 The diagram shows the graph of a quadratic function $y = ax^2 + bx + c$.
The curve crosses the x-axis at $(-2, 0)$ and $(4, 0)$, and crosses the y-axis at $(0, 16)$.



(a) Find the equation of the curve in the form $y = ax^2 + bx + c$.

$y =$ ..... **[4]**

(b) Find the coordinates of the turning point of the curve.

(..... , ...) **[2]**

(c) Write down the equation of the line of symmetry of the curve.

..... **[1]**

649 A quadratic function is given by $f(x) = a(x-h)^2 + k$.
The graph of $y = f(x)$ has a maximum point at $(-1, 8)$ and passes through the point $(2, -10)$.

(a) Write down the value of h and the value of k.

$h =$ ..... , $k =$ ..... **[2]**


(b) Find the value of a.

$a =$ ..... **[3]**

(c) Write $f(x)$ in the form $ax^2 + bx + c$.

$f(x) =$ ..... **[3]**


650 The diagram shows the graph of a quadratic function $y = f(x)$.
The graph crosses the x-axis at $x = -1$ and $x = 4$.
The graph crosses the y-axis at $(0, -8)$.



Find the equation of the quadratic function in the form $f(x) = ax^2 + bx + c$.

$f(x) =$ ..... **[4]**

651 The diagram shows a sketch of the curve $y = f(x)$ where $f(x)$ is a quadratic function.
The maximum point on the curve is $M(-2, 8)$ and the curve crosses the y-axis at $(0, -4)$.



(a) Find the equation of the curve.
Give your answer in the form $f(x) = ax^2 + bx + c$.

$f(x) =$ ..... **[4]**

(b) Solve $f(x) = 0$ to find the exact x-coordinates of the points where the curve crosses the x-axis.

Give your answers in the form $\frac{p \pm q\sqrt{6}}{r}$, where p, q and r are integers.

$x =$ ..... **[3]**

652 A quadratic curve has its vertex at $(3, -5)$ and passes through the point $(1, 7)$.
Find the equation of the curve in the form $y = a(x - h)^2 + k$.

$y =$ ..... **[3]**

653 The graph of a quadratic function has its vertex at $(-2, 5)$ and passes through the point $(1, -13)$.
Find the equation of the function in the form $y = a(x - h)^2 + k$.

$y =$ ..... **[3]**

654 The graph of a quadratic function has its vertex at $(2, 7)$ and passes through the point $(4, -5)$.

(a) Find the equation of the function in the form $y = a(x - h)^2 + k$.

$y =$ ..... **[3]**

(b) Find the y-intercept of the graph.

..... **[1]**

655 A quadratic curve crosses the x-axis at $(-1, 0)$ and $(5, 0)$.

The curve also passes through the point $(2, -18)$.

Find the equation of the curve in the form $y = ax^2 + bx + c$.


$y =$ ..... **[4]**

656 A quadratic curve has a vertex at the point $(2, -5)$ and passes through the point $(4, 7)$.

Find the equation of the curve.
Give your answer in the form $y = ax^2 + bx + c$.

$y =$ ..... **[4]**

657 The diagram shows a sketch of the graph of a quadratic function $y = a(x - h)^2 + k$.
The vertex of the graph is $V(2, 5)$.
The graph passes through the point $P(4, -7)$.



(a) Write down the value of h and the value of k.

$h =$ ..... , $k =$ ..... **[2]**

(b) Work out the value of a.

$a =$ ..... **[2]**

(c) Find the coordinates of the y-intercept of the graph.

(..... , ...) **[2]**

658 The graph of a quadratic function $f(x)$ intersects the x-axis at $x = -2$ and $x = 5$.
The graph passes through the point $(3, 10)$.


(a) Find an expression for $f(x)$ in the form $ax^2 + bx + c$.

$f(x) =$ ..... **[4]**

(b) Find the coordinates of the vertex of the graph of $y = f(x)$.

(..... , ...) **[2]**


659 The diagram shows a sketch of the graph of $y = ax^2 + bx + c$.
The vertex of the graph is $V(-1, 8)$ and it passes through the point $P(2, -10)$.



Work out the values of a, b and c.

$a =$ ..... , $b =$ ..... , $c =$ ..... **[5]**

660 The diagram shows the sketch of a quadratic function.
The curve crosses the x-axis at -3 and 2, and intersects the y-axis at 12.




(a) Find the equation of the quadratic function.
Give your answer in the form $y = ax^2 + bx + c$.

$y =$ ..... **[4]**

(b) Calculate the coordinates of the vertex of this curve.

((..... , ...)) **[2]**

661 The diagram shows a sketch of the quadratic function $y = f(x)$.
The curve crosses the x-axis at $x = -1.5$ and $x = 4$.
It passes through the point $(2, -14)$.



(a) Find the equation of the quadratic function in the form $f(x) = ax^2 + bx + c$.

$f(x) =$ ..... **[4]**

(b) Find the exact coordinates of the turning point of the graph.

(..... , ...) **[3]**

662 A quadratic function is given by $f(x) = a(x-h)^2 + k$.
The maximum point of the graph of $y = f(x)$ is $(-4, 20)$.
The graph intersects the x-axis at $x = -2$ and $x = p$.

(a) **(i)** Write down the value of h and the value of k.

$h =$ ..... , $k =$ ..... **[2]**

(ii) Find the value of a.

$a =$ ..... **[2]**


(iii) Find the value of p.

$p =$ ..... **[1]**

(iv) Express $f(x)$ in the form $ax^2 + bx + c$.

$f(x) =$ ..... **[2]**

663 The graph of a quadratic function $y = f(x)$ intersects the x-axis at the points $A(-1, 0)$ and $B(5, 0)$. The graph intersects the y-axis at the point $C(0, 10)$.



(a) Find the equation of the function $f(x)$.
Give your answer in the form $f(x) = ax^2 + bx + c$.

$f(x) =$ ..... **[4]**


(b) Find the coordinates of the turning point (vertex) of the graph.

(..... , ...) **[2]**

(c) The graph of $y = f(x)$ is translated by the vector $\begin{pmatrix} 2 \\ -3 \end{pmatrix}$ to give the graph of $y = g(x)$.
Find the equation of $g(x)$ in the form $g(x) = a(x - h)^2 + k$.

$g(x) =$ ..... **[2]**

664 The diagram shows the graph of a quadratic function $y = f(x)$.
The vertex of the graph is at $V(-2, 8)$ and it intersects the y-axis at $(0, 5)$.



(a) Find the equation of the function $f(x)$ in the form $f(x) = a(x - h)^2 + k$.

$f(x) =$ ..... **[3]**


(b) Write $f(x)$ in the form $ax^2 + bx + c$.

$f(x) =$ ..... **[2]**

(c) Calculate the exact x-coordinates of the points where the graph intersects the x-axis.
Give your answers correct to 3 significant figures.

$x =$ ..... , $x =$ ..... **[3]**

665 The diagram shows a sketch of the quadratic curve $y = ax^2 + bx + c$.
The curve crosses the x-axis at $A(-2, 0)$ and $B(5, 0)$, and crosses the y-axis at $C(0, 15)$.



(a) Find the values of a, b, and c.

$a =$ ..... , $b =$ ..... , $c =$ ..... **[5]**


(b) Find the coordinates of the vertex of the curve.

(..... , ...) **[3]**

(c) The equation $ax^2 + bx + c = k$ has no real roots.
Write down the inequality for k.

..... **[1]**

666 The diagram shows a sketch of the graph of a quadratic function $y = f(x)$.
The maximum point of the graph is $V(-2, 18)$.
The graph crosses the y-axis at $(0, 10)$.



(a) Find the equation of the quadratic function in the form $y = ax^2 + bx + c$.

$y =$ ..... **[3]**

(b) The graph crosses the x-axis at A and B.
Find the length of the line segment AB.

$AB =$ ..... **[3]**

(c) The line $y = -2x + k$ is a tangent to the curve $y = f(x)$.

Find the value of k.

$k =$ ..... **[4]**

667 A quadratic function $f(x) = a(x-h)^2 + k$ has its vertex at $(4, -5)$ and passes through the point $(1, 8.5)$.

(a) Find the values of a, h, and k.

$a =$ ..... , $h =$ ..... , $k =$ ..... **[3]**

(b) Solve the equation $f(x) = 0$, giving your answers correct to 2 decimal places.

$x =$ ..... , $x =$ ..... **[3]**

(c) The line $y = 2x - 7$ intersects the curve $y = f(x)$ at two points, P and Q.
Find the x-coordinates of P and Q, giving your answers correct to 2 decimal places.

$x =$ ..... , $x =$ ..... **[3]**

(d) The graph of $y = f(x)$ is translated by the vector $\begin{pmatrix} -3 \\ 9 \end{pmatrix}$ to give the graph of $y = g(x)$.
Write down the expression for $g(x)$ in the form $p(x-q)^2 + r$.

$g(x) =$ ..... **[2]**

668 The quadratic function $f(x) = ax^2 + bx + c$ has a minimum turning point at $(1, -8)$.
The graph of $y = f(x)$ crosses the x-axis at $x = -1$ and $x = 3$.

(a) Show that $f(x) = 2x^2 - 4x - 6$.

[3]

(b) Solve the equation.

$f(x) = 10$

$x =$ ..... , $x =$ ..... **[3]**

(c) A straight line has equation $y = mx - 8$.
The line is a tangent to the graph of $y = f(x)$.

Find the possible values of m.

$m =$ ..... , $m =$ ..... **[4]**


(d) The graph of $y = f(x)$ is translated by the vector $\begin{pmatrix} 0 \\ k \end{pmatrix}$.

The new graph has no x-intercepts.

Write down the inequality for k.

..... **[2]**

669 The diagram shows the sketch of the graph of a function $y = f(x)$.
The dashed lines are the asymptotes to the curve.



(a) Write down the equation of the vertical asymptote.

..... **[1]**

(b) Write down the equation of the horizontal asymptote.

..... **[1]**

670 Let $f(x) = \dfrac{7}{x-4} + 3$.

(a) Write down the equation of the vertical asymptote of the graph of $y = f(x)$.

..... **[1]**

(b) Write down the equation of the horizontal asymptote of the graph of $y = f(x)$.

..... **[1]**

671 The function f is given by $f(x) = \dfrac{5}{x+3} - 2$.


(a) Write down the equation of the vertical asymptote of the graph of $y = f(x)$.

$x =$ ..... **[1]**

(b) Write down the equation of the horizontal asymptote of the graph of $y = f(x)$.

$y =$ ..... **[1]**

672 The diagram shows the graph of a function $y = f(x)$.
The graph has two asymptotes, each parallel to one of the coordinate axes.
The equations of the asymptotes involve only integer values.



(a) Write down the equation of the vertical asymptote of the graph of $y = f(x)$.

..... **[1]**


(b) Write down the equation of the horizontal asymptote of the graph of $y = f(x)$.

..... **[1]**

(c) The graph of $y = f(x)$ is reflected in the y-axis to form the graph of $y = g(x)$.
Write down the equation of the vertical asymptote of the graph of $y = g(x)$.

..... **[1]**

673 The diagram shows the graph of $y = f(x)$ and its two asymptotes, which are shown as dashed lines.



(a) Write down the equation of the vertical asymptote of the graph.

..... **[1]**

(b) Write down the equation of the horizontal asymptote of the graph.

..... **[1]**

(c) Use the graph to write down

(i) the value of $f(4)$,

$f(4) =$ ..... **[1]**

(ii) the value of x for which $f(x) = -2$.

$x =$ ..... **[1]**

674 A function is defined as $f(x) = \dfrac{4}{x-1} + 3$.

(a) Write down the equation of the

(i) vertical asymptote,

..... **[1]**

(ii) horizontal asymptote.

..... **[1]**


(b) Find the coordinates of the point where the graph of $y = f(x)$ crosses the x-axis.

((..... , ...)) **[2]**

(c) Find the coordinates of the point where the graph of $y = f(x)$ crosses the y-axis.

((..... , ...)) **[1]**

675 The diagram shows the graph of $f(x) = \tan x$ for $0° \leq x \leq 360°$.



(a) Write down the equations of the two asymptotes of the graph of $f(x) = \tan x$ for $0° \leq x \leq 360°$.

$x =$ ..... , $x =$ ..... **[2]**

(b) Explain how the graph shows that $\tan 90°$ is undefined.

..... **[1]**

(c) The graph of $y = \tan x$ is translated by the vector $\begin{pmatrix} 35° \\ 0 \end{pmatrix}$.
Write down the equations of the two asymptotes of the translated graph for $0° \leq x \leq 360°$.

$x =$ ..... , $x =$ ..... **[2]**

(d) Given that $\tan 45° = 1$, use the graph to write down the other solution to $\tan x = 1$ for $0° \leq x \leq 360°$.

$x =$ ..... ° **[1]**

676 Consider the functions $g(x) = \tan x$ and $h(x) = \dfrac{4x-1}{2x+5}$.

(a) Write down the equations of the two asymptotes of the graph of $y = g(x)$ for $0° \leq x \leq 360°$.

$x =$ ..... , $x =$ ..... **[2]**

(b) Write down the equations of the vertical and horizontal asymptotes of the graph of $y = h(x)$.

Vertical asymptote $x =$ ..... , Horizontal asymptote $y =$ ..... **[2]**

(c) Solve the equation

$h(x) = 0$

$x =$ ..... **[2]**

677 A function is defined as $f(x) = \dfrac{3}{x+2} + 1$.


(a) Write down the equations of the two asymptotes of the graph of $y = f(x)$.

Vertical asymptote: ..... , Horizontal asymptote: ..... **[2]**

(b) Find the coordinates of the points where the graph of $y = f(x)$ crosses the x-axis and the y-axis.

x-axis at ..... , y-axis at ..... **[2]**

(c) On the grid, sketch the graph of $y = f(x)$.



[3]

678 Let $f(x) = \dfrac{3}{x+2} - 4$.

(a) Write down the equation of the vertical asymptote of the graph of $y = f(x)$.

..... **[1]**

(b) Write down the equation of the horizontal asymptote of the graph of $y = f(x)$.

..... **[1]**

(c) Find the coordinates of the points where the graph of $y = f(x)$ intersects the coordinate axes.

x-intercept = ... , y-intercept = ... **[5]**

(d) The graph of $y = \tan x$ has an infinite number of vertical asymptotes.
Write down the equations of the two asymptotes in the interval $0° \leq x \leq 360°$.

$x =$ ..... , $x =$ ..... **[2]**

679 A function is given by $f(x) = \dfrac{a}{x-b} + c$, where a, b and c are constants.
The graph of $y = f(x)$ has a vertical asymptote at $x = 2$ and a horizontal asymptote at $y = -1$.
The graph intersects the x-axis at the point $(5, 0)$.

(a) Write down the value of b and the value of c.

$b =$ ..... , $c =$ ..... **[2]**

(b) Work out the value of a.

$a =$ ..... **[2]**


(c) The graph of $y = f(x)$ intersects the straight line $y = -x + 5$ at two points.
Find the x-coordinates of these two points.

$x =$ ..... , $x =$ ..... **[2]**

680 The diagram shows the graph of $y = f(x)$ where $f(x) = \dfrac{p}{x-q} + r$.

The graph has asymptotes at $x = 2$ and $y = -1$.

The graph passes through the point $(3, 3)$.



(a) Find the values of q and r.

$q =$ ..... , $r =$ ..... **[2]**

(b) Find the value of p.


$p =$ ..... **[2]**

(c) Find the coordinates of the point where the graph of $y = f(x)$ crosses the x-axis.

(..... , ...) **[3]**

681 The function $g(x) = \tan x$ is defined for $0° \leq x \leq 360°$.

(a) On the grid, sketch the graph of $y = g(x)$.



[2]


(b) Write down the equations of the asymptotes of the graph of $y = g(x)$ in this domain.

[2]

(c) Solve the equation $2\tan x + 3 = 0$ for $0° \leq x \leq 360°$.
Give your answers correct to 1 decimal place.

$x =$ ..... , $x =$ ..... **[3]**

682 The diagram shows the graph of $y = h(x)$, where $h(x) = \dfrac{5}{x - 1.5} + 2$ for $x \in 1.5$.
The dashed lines show the asymptotes of the curve.



(a) From the graph, write down the equation of

(i) the vertical asymptote,

..... **[1]**

(ii) the horizontal asymptote.

..... **[1]**

(b) Explain why the equation $h(x) = 2$ has no solution.

..... **[1]**

(c) The graph of $y = h(x)$ is reflected in the y-axis.
Write down the equations of the vertical and horizontal asymptotes of the reflected graph.

Vertical asymptote: ..... , Horizontal asymptote: ..... **[2]**

(d) Solve the equation $h(x) = 3.4$, giving your answer correct to 3 significant figures.

$x =$ ..... **[3]**

683 The function $f(x)$ is given by $f(x) = \dfrac{p}{x-q} + r$, where p, q, and r are constants.
The graph of $y = f(x)$ has a vertical asymptote at $x = 5$ and a horizontal asymptote at $y = -3$.
The graph passes through the point $(7, -2)$.

(a) Write down the value of q and the value of r.

$q =$ ..... , $r =$ ..... **[2]**

(b) Find the value of p.

$p =$ ..... **[3]**

(c) Find $f^{-1}(x)$.

$f^{-1}(x) =$ ..... **[4]**

(d) Write down the equations of the asymptotes of the graph of $y = f^{-1}(x)$.

$x =$ ..... , $y =$ ..... **[2]**

684 A function is defined as $y = f(x)$.

(a) Describe fully the single transformation that maps the graph of $y = f(x)$ onto the graph of $y = f(x - 4)$.

.....

..... **[2]**

(b) The graph of $y = f(x)$ is translated by the vector $\begin{pmatrix} 0 \\ 7 \end{pmatrix}$ to give the graph of $y = g(x)$.
Write down the expression for $g(x)$ in terms of $f(x)$.

$g(x) =$ ..... **[1]**

685 **(a)** The graph of $y = x^2$ is translated by the vector $\begin{pmatrix} 0 \\ -5 \end{pmatrix}$ to map onto the graph of $y = g(x)$.
Write down $g(x)$ in terms of x.

$g(x) =$ ..... **[1]**

(b) The graph of $y = x^2$ is translated by the vector $\begin{pmatrix} 3 \\ 0 \end{pmatrix}$ to map onto the graph of $y = h(x)$.
Write down $h(x)$ in terms of x.

$h(x) =$ ..... **[1]**

(c) Describe fully the single transformation that maps the graph of $y = x^2$ onto the graph of $y = (x + 4)^2 - 1$.

.....

.....

..... **[2]**

686 A function is defined as $f(x) = 2x - 5$.

(a) The graph of $y = f(x)$ is translated by the vector $\begin{pmatrix} 0 \\ 7 \end{pmatrix}$ to give the graph of $y = g(x)$.
Find $g(x)$ in terms of x, giving your answer in its simplest form.

$g(x) =$ ..... **[2]**


(b) Describe fully the single transformation that maps the graph of $y = f(x)$ onto the graph of $y = f(x - 4)$.

.....

.....

..... **[2]**

687 The diagram shows the graphs of $y = f(x)$ and $y = g(x)$ on a coordinate grid.



(a) Write down the column vector that describes the translation mapping the graph of $y = f(x)$ onto the graph of $y = g(x)$.

$\begin{pmatrix} \quad \\ \quad \end{pmatrix}$ **[2]**

(b) Given that $f(x) = x^2$, write down $g(x)$ in the form $(x - a)^2 + b$.

$g(x) =$ ..... **[2]**

(c) Write down the coordinates of the vertex of the graph of $y = f(x + 5) - 1$.

(..... , ...) **[2]**

688 The graph of $y = f(x)$ has a local maximum at the point $P(-2, 5)$ and a local minimum at the point $Q(3, -1)$.

(a) Write down the coordinates of the local maximum of the graph of $y = f(x) - 4$.

(..... , ...) **[1]**

(b) Write down the coordinates of the local minimum of the graph of $y = f(x - 2)$.

(..... , ...) **[1]**

(c) The graph of $y = f(x)$ is translated by the vector $\begin{pmatrix} a \\ b \end{pmatrix}$ such that the image of the local maximum P is at the origin $(0, 0)$.

(i) Find the value of a and the value of b.

$a =$ ..... , $b =$ ..... **[2]**

(ii) Write down the coordinates of the image of the local minimum Q under the same translation.

(..... , ...) **[1]**

689 Let $f(x) = 3x^2$.

(a) The graph of $y = f(x)$ is translated by the vector $\begin{pmatrix} -2 \\ 0 \end{pmatrix}$ to give the graph of $y = g(x)$.

Find $g(x)$ in terms of x.

Give your answer in the form $ax^2 + bx + c$.

$g(x) =$ ..... **[3]**

(b) The graph of $y = g(x)$ is translated by the vector $\begin{pmatrix} 0 \\ -8 \end{pmatrix}$ to give the graph of $y = h(x)$.


Write down the coordinates of the minimum point of the graph of $y = h(x)$.

(..... , ...) **[2]**


(c) Find the y-intercept of the graph of $y = h(x)$.

..... **[1]**

690 The diagram shows the graph of $y = f(x)$ on a coordinate grid.



(a) On the grid below, draw the graph of $y = f(x + 2) - 3$.



[2]

(b) The graph of $y = f(x)$ is mapped onto the graph of $y = h(x)$ by a single transformation.
The point $(1, 4)$ on $y = f(x)$ is mapped onto $(1, -1)$ on $y = h(x)$, and the point $(3, 1)$ is mapped onto $(3, -4)$.
Describe fully the single transformation that maps the graph of $y = f(x)$ onto the graph of $y = h(x)$.

.....

.....

..... **[2]**


(c) Write down $h(x)$ in terms of $f(x)$.

$h(x) =$ ..... **[1]**

(d) Find the coordinates of the image of the point $(4, 2)$ on $y = f(x)$ when it is mapped onto the graph of $y = f(x - 3) + 4$.

(..... , ...) **[2]**

691 The diagram shows a sketch of the graph of $y = f(x)$.
The curve intersects the x-axis at $(2, 0)$ and $(6, 0)$ and has a minimum point at $M(4, -5)$.



(a) A transformation maps the graph of $y = f(x)$ onto the graph of $y = f(x) + 4$.
Write down the coordinates of the image of point M.

((..... , ...)) **[1]**

(b) The graph of $y = f(x)$ is mapped onto the graph of $y = g(x)$ by a translation with vector $\begin{pmatrix} -3 \\ 2 \end{pmatrix}$.
Write down the expression for $g(x)$ in terms of $f(x)$.

$g(x) =$ ..... **[2]**

(c) The graph of $y = f(x - k)$ passes through the origin $(0, 0)$.
Find the two possible values of the constant k.

$k =$ ..... , $k =$ ..... **[2]**

(d) The function $h(x)$ is such that the graph of $y = h(x)$ has a minimum point at $(7, -5)$.
Find $h(x)$ in terms of $f(x)$.

$h(x) =$ ..... **[1]**

692 A function is given as $f(x) = x^2 - 2x - 3$.

(a) The graph of $y = f(x)$ is translated by the vector $\begin{pmatrix} -4 \\ 0 \end{pmatrix}$ to form the graph of $y = g(x)$.
Find $g(x)$ in the form $ax^2 + bx + c$.

$g(x) =$ ..... **[4]**

(b) The graph of $y = f(x)$ is mapped onto the graph of $y = h(x)$ by a translation with vector $\begin{pmatrix} 0 \\ -5 \end{pmatrix}$.
Solve the equation $h(x) = 0$.

$x =$ ..... , $x =$ ..... **[3]**

693 The functions $f(x)$ and $g(x)$ are defined as $f(x) = x^2 - 6x + 14$ and $g(x) = \dfrac{3}{x+2} - 5$.

(a) By writing $f(x)$ in the form $(x-a)^2 + b$, find the column vector of the translation that maps the graph of $y = x^2$ onto the graph of $y = f(x)$.

$\begin{pmatrix} \quad \\ \quad \end{pmatrix}$ **[3]**

(b) The graph of $y = g(x)$ is translated by the vector $\begin{pmatrix} 4 \\ 2 \end{pmatrix}$ to map onto the graph of $y = h(x)$. Find $h(x)$ in its simplest form.

$h(x) =$ ..... **[3]**

(c) Write down the equations of the vertical and horizontal asymptotes of the graph of $y = h(x)$.

$x =$ ..... , $y =$ ..... **[2]**

694 A function is defined as $f(x) = \dfrac{20}{x}$.

(a) The graph of $y = f(x)$ is mapped onto the graph of $y = g(x)$ by a translation with vector $\begin{pmatrix} 0 \\ -4 \end{pmatrix}$.

(i) Write down an expression for $g(x)$.

$g(x) =$ ..... **[1]**

(ii) Solve the equation $g(x) = 0$.

$x =$ ..... **[2]**

(b) The graph of $y = f(x)$ is translated by the vector $\begin{pmatrix} k \\ 0 \end{pmatrix}$ to give the graph of $y = h(x)$.
The graph of $y = h(x)$ passes through the point $(8, 5)$.
Work out the value of k.

$k =$ ..... **[3]**

(c) Let $p(x) = f(x - 3) + 7$.
Write down the equations of the vertical and horizontal asymptotes of the graph of $y = p(x)$.

Vertical asymptote: ..... , Horizontal asymptote: ..... **[2]**

695 The function f is defined by $f(x) = 2x^2 - 5x + 1$.

(a) The graph of $y = f(x)$ is translated by the vector $\begin{pmatrix} -2 \\ -6 \end{pmatrix}$ to give the graph of $y = g(x)$.

(i) Write down $g(x)$ in terms of $f(x)$.

$g(x) =$ ..... **[1]**

(ii) Find $g(x)$ in the form $ax^2 + bx + c$.

$g(x) =$ ..... **[3]**


(b) The graph of $y = f(x)$ is mapped onto the graph of $y = h(x)$ by a single translation, where $h(x) = 2x^2 - 13x + 18$.
Find the column vector of this translation.

$\begin{pmatrix} \quad \\ \quad \end{pmatrix}$ **[4]**

(c) Find the coordinates of the point of intersection of the graphs $y = f(x)$ and $y = g(x)$.

(..... , ...) **[3]**

696 The diagram shows a sketch of the function $f(x) = 10^x$.



(a) Write down the inverse function, $f^{-1}(x)$.

$f^{-1}(x) =$ ..... **[1]**

(b) Work out the exact value of $f^{-1}\left(\frac{1}{100}\right)$.

..... **[1]**

(c) Write down the exact solution to the equation $f(x) = 45$.

$x =$ ..... **[1]**

697 **(a)** Write $10^3 = 1000$ as a logarithmic equation.

..... **[1]**

(b) Find the exact value of x when $\log x = -2$.

$x =$ ..... **[1]**

(c) Solve $5^x = 12$, giving your answer in the form $\dfrac{\log a}{\log b}$.

$x =$ ..... **[1]**

698 The functions f and g are defined as $f(x) = 10^x$ and $g(x) = \log x$.

(a) Find the value of $g(100)$.

..... **[1]**

(b) Find the value of x when $f(x) = 0.001$.

$x =$ ..... **[1]**

(c) Write down the expression for $g(f(x))$ in its simplest form.

$g(f(x)) =$ ..... **[1]**

699 **(a)** Work out the value of

$\log 10000$

..... **[1]**

(b) Write the following equation in the form $\log_a b = c$.

$5^3 = 125$

..... **[1]**

(c) Solve the equation

$\log_2 x = 6$

$x =$ ..... **[1]**

700 **(a)** Find the value of x when $\log_2 x = 5$.

$x =$ ..... **[1]**

(b) Find the value of y when $10^y = 0.01$.

$y =$ ..... **[1]**

(c) Write the equation $P = k^t$ in logarithmic form.

..... **[1]**

701 **(a)** Write the following as a logarithmic equation.

$5^3 = 125$

..... **[1]**

(b) Find the value of $\log 1000$.

..... **[1]**

(c) Solve the equation.

$\log_2 x = 5$

$x =$ ..... **[2]**

702 **(a)** Write the equation $10^5 = 100000$ in the form $\log a = b$.

..... **[1]**

(b) Find the value of x when $\log x = 3$.

$x =$ ..... **[2]**

(c) The function $f(x)$ is defined as $f(x) = 10^x$.
Find $f^{-1}(x)$.

$f^{-1}(x) =$ ..... **[1]**

703 All logarithms in this question are in the base shown, or base 10
10 where no base is written.

(a) Write down the value of

(i) log
1000
log1000,

..... **[1]**

(ii) log
2
32
log
2
32.

..... **[1]**

(b) Given that
log
3
=
4
log
3
x=4, find the value of
x.

=
x= ..... **[1]**

(c) The function
f is defined by
(
)
=
10
f(x)=10
x
.
Write down
1
(

)
f
1
(x).

1
(
)
=
f
1
(x)= ..... **[1]**

704 **(a)** Find the value of x in each of the following equations.

(i) $10^x = 100000$

$x =$ ..... **[1]**

(ii) $\log x = -2$

$x =$ ..... **[1]**

(iii) $\log_2 64 = x$

$x =$ ..... **[1]**

(b) Given that $f(x) = 10^x$, write down an expression for $f^{-1}(x)$.

$f^{-1}(x) =$ ..... **[1]**

705 **(a)** Write down the value of

$\log 1000$

..... **[1]**

(b) Solve the equation.

$\log x = -2$

$x =$ ..... **[2]**

(c) Write the equation $5^x = 125$ in the form $x = \log_a b$.

$x =$ ..... **[1]**

706 Answer the following independent questions on logarithmic and exponential equations.

(a) Write the exponential equation $10^4 = 10000$ in logarithmic form.

..... **[1]**


(b) Find the exact value of x when $\log_5 x = 3$.

$x =$ ..... **[2]**

(c) The solution to the equation $7^x = 25$ can be written in the form $x = \dfrac{\log b}{\log a}$.
Write down the value of a and the value of b.

$a =$ ..... , $b =$ ..... **[2]**

707 The diagram shows a sketch of the graph of $y = f(x)$ for $x > 0$.
The function is defined as $f(x) = p\log x + q$, where $\log x$ denotes the logarithm to base 10.
The graph passes through the points $(10, 8)$ and $(100, 11)$.



(a) **(a)** Find the value of p and the value of q.

$p =$ ..... , $q =$ ..... **[3]**

(b) Solve the equation $f(x) = 23$.

$x =$ ..... **[2]**

(c) Find $f^{-1}(x)$.

$f^{-1}(x) =$ ..... **[2]**

708 The logarithmic function is the inverse of the exponential function.

(a) Write the equation $y = 6^x$ in logarithmic form.

(a) Write the equation $y = 6^x$ in logarithmic form.

$x =$ ..... **[1]**

(b) Solve the equation.

$3^x = 20$ Give your answer correct to 3 significant figures.

$x =$ ..... **[2]**

709 A population of insects is growing exponentially.
The number of insects, N, after t days is given by

$$N = 250 \times 1.08^t$$

(a) Work out the number of insects after 10 days.
Give your answer to the nearest integer.

..... **[2]**

(b) Calculate the number of days it takes for the population to reach 1000.
Give your answer correct to 1 decimal place.

..... days **[3]**

710 The value of a machine, V dollars, depreciates exponentially according to the formula

$$V = 15000 \times 0.8^t$$

where t is the number of years after it is purchased.

(a) Calculate the value of the machine after 4 years.

$\$ \ldots\ldots\ldots\ldots\ldots$ **[2]**

(b) Find the number of years it takes for the value of the machine to decrease to \$7500.
Give your answer correct to 3 significant figures.

$\ldots\ldots\ldots\ldots\ldots$ years **[3]**

711 A machine is purchased for \$25000. Its value depreciates exponentially by 12% each year.

(a) Calculate the value of the machine after 4 years.
Give your answer correct to the nearest dollar.

% ..... **[2]**

(b) Find the number of complete years it takes for the value of the machine to fall below \$10000.
You must show all your working.

..... years **[3]**

712 **(a)** Solve the equation.

$3^{2x} = 50$ Give your answer correct to 3 significant figures.

$x =$ ..... **[3]**

(b) Make x the subject of the formula.

$y = 10^{x-4}$

$x =$ ..... **[2]**

713 A population of bacteria is modelled by the formula

$P = 5000 \times 1.12^t$ where P is the population and t is the time in hours after the start of an experiment.

(a) **(a)** Write down the initial population of the bacteria.

..... **[1]**

(b) Find the time it takes for the population to reach 20 000.
Give your answer correct to 1 decimal place.

$t =$ ..... hours **[4]**

714 An investment of \$8000 earns compound interest at a rate of 3.5% per year. The value of the investment after n years is \$$V$.

(a) Write down an equation for V in terms of n.

$V =$ ..... **[1]**

(b) Calculate the minimum number of complete years it takes for the value of the investment to exceed \$12000.
Show all your working.

..... years **[3]**

(c) Solve the equation.

$2^x = 55$ Give your answer correct to 3 significant figures.

$x =$ ..... **[2]**

715 **(a)** Solve the equation $3^x = 20$.
Give your answer correct to 3 significant figures.

$x =$ ..... **[3]**

(b) The mass, M grams, of a radioactive substance after t days is given by the formula
$M = 500 \times 0.85^t$ Calculate the value of t when $M = 100$.
Give your answer correct to 3 significant figures.

$t =$ ..... **[3]**

716 The value, V dollars, of a painting t years after it was bought is given by the formula

$$V = 4500 \times 1.08^t$$

(a) Calculate the value of the painting 5 years after it was bought.

Give your answer correct to the nearest dollar.


..... **[2]**

(b) Calculate the time it takes for the value of the painting to reach \$10 000.

Give your answer in years, correct to 2 decimal places.

..... years **[4]**

717 The diagram shows the graph of
=
log
y=logx for
0.1
10
0.1x10.



(a) Use the graph to estimate the value of
log
6
log6.

..... **[1]**

(b) Use the graph to solve the equation
log
=
0.5
logx=0.5.

=
x= ..... **[1]**

(c) Solve the equation
3
+
1
=
20
3
x+1
=20, giving your answer correct to

3
3 significant figures.

=
x= ..... **[3]**

(d) Solve the equation
log
(
2
3
)
=
1.4
log(2x3)=1.4, giving your answer correct to
3
3 significant figures.

=
x= ..... **[2]**

718 The mass, M grams, of a radioactive substance after t days is given by the formula $M = 400 \times 0.88^t$.

(a) Calculate the mass of the substance after 5 days.
Give your answer correct to 1 decimal place.

..... grams **[2]**

(b) Find the number of days it takes for the mass of the substance to reduce to exactly half of its initial value.

..... days **[3]**

(c) Another radioactive substance has an initial mass of 800 grams.
After 10 days, its mass is 300 grams.
It decays according to the formula $M = 800 \times k^t$.
Find the value of k, giving your answer correct to 3 significant figures.

$k =$ ..... **[3]**

719 An amount of \$8000 is invested in an account that pays compound interest at a rate of 4.5% per year. The value of the investment, V dollars, after t years is given by the formula

$$V = 8000 \times 1.045^t$$

(a) Calculate the value of the investment after 5 years.

\$ ..... **[2]**

(b) Use logarithms to calculate the number of complete years it takes for the value of the investment to first exceed \$15000.

..... years **[3]**

(c) Solve the equation.

$7^{x-1} = 250$ Give your answer correct to 3 significant figures.

$x =$ ..... **[3]**

720 A population of insects grows exponentially.
The number of insects, N, after t weeks is given by the formula

$$N = 800 \times 2^t$$

(a) Work out the number of insects after 4 weeks.

..... **[1]**

(b) The number of insects reaches 102 400.
Work out the exact value of t.

$t =$ ..... **[2]**

(c) Another population of insects can be modelled by the formula $P = 1500 \times 3^t$.
Find the value of t when this population reaches 20 000.
Give your answer correct to 3 significant figures.

$t =$ ..... **[3]**

(d) A sum of \$2000 is invested at an annual compound interest rate of 5%. The total value, V, of the investment after n years is given by

$$V = 2000 \times 1.05^n$$

Calculate the minimum number of complete years required for the investment to double in value.

..... years **[3]**

721 A function f is defined as $f(x) = 4 \times 10^{2x} - 7$.

(a) Find $f^{-1}(x)$.


$f^{-1}(x) =$ ..... **[3]**

(b) Solve the equation $f(x) = 53$.
Give your answer correct to 3 significant figures.

$x =$ ..... **[3]**

722 A function is defined as $f(x) = 2^x$.

(a) On the grid, sketch the graph of $y = f(x)$ for $-2 \leq x \leq 3$.



[2]

(b) Write down the equation of the inverse function, $f^{-1}(x)$.

$f^{-1}(x) =$ ..... **[1]**

(c) On the same grid in part (a), sketch the graph of $y = f^{-1}(x)$ for $0 < x \leq 8$. **[2]**

(d) Solve the equation $2^x = 12$.
Give your answer correct to 3 significant figures.

$x =$ ..... **[2]**

723 **(a)** The mass, M grams, of a radioactive substance after t days is given by $M = 250 \times 0.92^t$.

(i) Write down the initial mass of the substance.

..... grams **[1]**

(ii) Find the number of days it takes for the mass to reduce to 50 grams.
Give your answer correct to 3 significant figures.

..... days **[3]**

(b) An amount of \$800 is invested at a rate of 6% per year compound interest. Calculate the number of complete years it takes for the investment to first exceed \$2000.


..... years **[3]**

724 The mass, m grams, of a radioactive isotope decays according to the formula

$m = 800 \times 10^{-kt}$ where t is the time in years and k is a constant.

The diagram shows a sketch of the decay curve.

After 15 years, the mass is 250 grams.



(a) Show that $k = 0.0337$ correct to 3 significant figures.

[3]

(b) Find the number of years it takes for the mass to become 50 grams.

..... years **[3]**

(c) Make t the subject of the formula $m = 800 \times 10^{-kt}$.

$t =$ ..... **[2]**

725 Let $f(x) = 5 \times 10^x$ and $g(x) = \log x$.

(a) Find the value of $f(g(100))$.

..... **[2]**

(b) Find $f^{-1}(x)$.

$f^{-1}(x) =$ ..... **[2]**

(c) Solve the equation.

$f(x) = 200$ Give your answer correct to 3 significant figures.

$x =$ ..... **[2]**

(d) Find $g(f(x))$ in the form $x + k$, where k is a constant.

..... **[2]**

726 A function is defined as $f(x) = 5 \times 3^x - 2$.

(a) Find the value of $f(-2)$, giving your answer as a fraction in its simplest form.

..... **[2]**

(b) Find $f^{-1}(x)$.

Give your answer in terms of base 10 logarithms.

$f^{-1}(x) =$ ..... **[3]**


(c) Solve the equation.

$f(x) = 500$

Give your answer correct to 3 significant figures.

$x =$ ..... **[3]**

727 The diagram shows the graph of $y = f(x)$.



(a) The graph of $y = f(x)$ passes through the point $(8, 3)$.
Given that $f(x) = \log_a x$, show that $a = 2$.

[2]

(b) **(i)** On the grid, sketch the graph of $y = f^{-1}(x)$. **[2]**

(ii) Write down the equation of the asymptote of the graph of $y = f^{-1}(x)$.

..... **[1]**

(c) By drawing a suitable straight line on the grid, find an approximate solution to the equation $\log_2 x + x = 8$.

$x =$ ..... **[3]**

728 The function $f(x)$ is defined as $f(x) = 4 \times 10^{0.5x}$.

(a) Find the value of $f(3)$.
Give your answer correct to 3 significant figures.

..... **[2]**

(b) Find $f^{-1}(x)$.


$f^{-1}(x) =$ ..... **[3]**

(c) Solve the equation

$$f(x) = 200$$

$x =$ ..... **[3]**

729 The diagram shows the graph of $y = f(x)$ where $f(x) = a \times b^x$.
The graph intersects the y-axis at the point $(0, 12)$ and passes through the point $P(3, 96)$.



(a) Find the value of a and the value of b.

$a =$ ..... , $b =$ ..... **[3]**

(b) Find $f^{-1}(x)$.
Give your answer in terms of base 10 logarithms.

$f^{-1}(x) =$ ..... **[3]**

(c) Solve the equation $f(x) = 500$.
Give your answer correct to 3 significant figures.

$x =$ ..... **[3]**

730 The value, \$V, *of a machine t years after it is purchased is modelled by the formula* $V = 24000 \times 0.85^t$.

(a) Calculate the value of the machine after 4 years, giving your answer correct to the nearest dollar.

\$ ..... **[2]**

(b) Find the number of years it takes for the value of the machine to fall to \$9000.
Give your answer correct to 3 significant figures.

$t =$ ..... years **[3]**

(c) A different asset increases in value by 6.8% each year.
Find the minimum whole number of years it takes for the value of this asset to double.

..... years **[3]**

(d) The function g is defined by $g(x) = 5 \times 2^x + 3$.
Find $g^{-1}(x)$.

$g^{-1}(x) =$ ..... **[3]**

731 The function $f(x) = A \times b^x$, where $b > 0$, passes through the points $(0, 7)$ and $(3, 56)$.

(a) Find the value of A and the value of b.

$A =$ ..... , $b =$ ..... **[3]**

(b) Find the inverse function, $f^{-1}(x)$.

$f^{-1}(x) =$ ..... **[3]**

(c) The equation $f(x) = 600$ is equivalent to $2^x = k$.

(i) Work out the exact value of k as a fraction.

$k =$ ..... **[1]**

(ii) Solve the equation $f(x) = 600$.
Give your answer correct to 3 decimal places.

$x =$ ..... **[2]**

(d) A different function $g(x)$ is defined as $g(x) = \log_c(x)$.
The graph of $y = g(x)$ passes through the point $(81, 4)$.


(i) Find the value of c.

$c =$ ..... **[2]**

(ii) Solve the equation $g(x) = -2$.

$x =$ ..... **[1]**

732 The diagram shows three vertices, $A(1, 2)$, $B(5, 2)$ and $C(7, 5)$, of a parallelogram $ABCD$.



(a) Find the coordinates of vertex D.

((..... , ...)) **[2]**

(b) Calculate the area of the parallelogram $ABCD$.

..... **[2]**

733 Points $P(-4, 2)$ and $Q(6, 2)$ are plotted on a Cartesian grid.

(a) Write down the coordinates of the midpoint of the line segment PQ.


(..... , ...) **[1]**

(b) Point R lies on the positive y-axis.
The area of triangle PQR is 35 square units.
Find the coordinates of R.

(..... , ...) **[3]**

734 The points $A(-1,3)$, $B(2,-1)$ and $C(6,1)$ are three vertices of a parallelogram $ABCD$.

(a) On the grid, plot the points A, B and C.



[1]


(b) Find the coordinates of the fourth vertex, D.

(..... , ...) **[2]**

(c) Write down the coordinates of the centre of the parallelogram.

(..... , ...) **[2]**

735 The diagram shows three points, A, B, and C, plotted on a Cartesian coordinate grid.



(a) Write down the coordinates of point A and point C.

$A =$ ..... , $C =$ ..... **[2]**

(b) Write down the coordinates of point D such that $ABCD$ is a parallelogram.


$D =$ (..... , ...) **[2]**

(c) Calculate the area of parallelogram $ABCD$.

..... square units **[2]**

736 In a Cartesian coordinate plane, a kite $PQRS$ has vertices $P(1,5)$, $Q(4,2)$, $R(1,-4)$, and $S(-2,2)$.

(a) Plot the points $P(1,5)$, $Q(4,2)$, $R(1,-4)$, and $S(-2,2)$ on the grid below and join them in order to form the kite $PQRS$.



[2]

(b) Write down the coordinates of the point where the diagonals of kite $PQRS$ intersect.

(..... , ...) **[1]**


(c) Calculate the area of kite $PQRS$.

..... square units **[2]**

(d) Point P is reflected in the x-axis to give point P', and point Q is reflected in the y-axis to give point Q'.
Write down the coordinates of P' and Q'.

$P' =$ ..... , $Q' =$ ..... **[2]**

737 The points $P(a, b)$ and $Q(b, a)$ are in the first quadrant, with $a > b$.
The origin is denoted by $O(0, 0)$.



(a) By drawing a suitable rectangle or otherwise, show that the area of triangle OPQ is $\frac{1}{2}(a^2 - b^2)$.

[3]

(b) The area of triangle OPQ is 24 and the length of PQ is $4\sqrt{2}$.
Find the values of a and b.

$a =$ ..... , $b =$ ..... **[5]**

738 A rectangle has its sides parallel to the coordinate axes.
Two opposite vertices of the rectangle are $A(-2, -4)$ and $C(x, y)$, where $x > 0$ and $y > 0$.
The area of the rectangle is 54 square units and its perimeter is 30 units.

(a) Show that $x + y = 9$.

[2]

(b) Form an equation in x and solve it to find the two possible pairs of coordinates for C.

[4]

(c) For the case where $x > y$, write down the coordinates of the other two vertices of the rectangle.

[2]

739 Points A and B have coordinates $(3k, -k)$ and $(k+4, -5k)$ respectively, where k is a constant.

(a) The midpoint of the line segment AB lies on the y-axis.
Find the value of k.

$k =$ ..... **[2]**


(b) Using your value of k from part (a), find the exact length of AB.
Give your answer in the form $a\sqrt{b}$ where b is a prime number.

..... **[3]**

(c) A point C lies on the x-axis.
The distance AC is equal to the distance BC.
Find the coordinates of C.

(..... , ...) **[4]**

740 A regular hexagon $ABCDEF$ has its center at the origin $O(0, 0)$.
The point A has coordinates $(6, 0)$.



(a) Write down the coordinates of point D.

(..... , ...) **[1]**

(b) Calculate the coordinates of point B.
Give your values correct to 2 decimal places.

(..... , ...) **[3]**

(c) Point P lies on the line segment AD such that $AP : PD = 1 : 3$.
Find the coordinates of P.

(..... , ...) **[2]**

741 A straight line has the equation $2y + 3x = 12$.
The line intersects the y-axis at A and the x-axis at B.
The point O is the origin $(0, 0)$.

(a) Find the coordinates of A and B.

$A =$ ..... , $B =$ ..... **[2]**

(b) Point C is such that $OACB$ is a rectangle.
Write down the coordinates of C.

((..... , ...)) **[1]**

(c) Point P lies on the line segment AB and has an x-coordinate of 2.
Find the coordinates of P.

((..... , ...)) **[2]**

(d) Find the area of triangle OPB.

..... **[2]**

742 A straight line L has the equation $3x + 2y = 18$.

(a) The line L crosses the x-axis at point A and the y-axis at point B.
Find the coordinates of A and B.

$A =$

$B =$ **[2]**

(b) Point P lies on the line segment AB such that the ratio $AP : PB = 2 : 1$.
Find the coordinates of P.

(..... , ...) **[3]**

(c) Does the point $(4, 3)$ lie on the line L?
Show how you decide.

[2]

743 A rectangle $ABCD$ has vertices $A(-3,-1)$, $B(5,-1)$, $C(5,k)$, and $D(h,k)$, where $k>0$.

(a) State the value of h and the quadrant in which point D lies.

$h =$ ..... , Quadrant: ..... **[2]**

(b) Given that the area of rectangle $ABCD$ is 52 square units, calculate the value of k.

$k =$ ..... **[3]**

(c) Point M is the centre of rectangle $ABCD$.
Find the coordinates of M.

$M =$ (..... , ...) **[2]**

(d) Point $E(p,q)$ lies on the y-axis such that triangle ABE has an area of 18 square units and $q>0$.
Find the coordinates of E.


$E =$ (..... , ...) **[3]**

744 Points A and B have coordinates $(2, -3)$ and $(6, 5)$ respectively.

Work out the gradient of the line AB.

..... **[2]**

745 The diagram shows a straight line L drawn on a coordinate grid.



(a) Find the gradient of line L.

..... **[2]**

(b) Write down the coordinates of the y-intercept of line L.


(..... , ...) **[1]**

746 The points A and B have coordinates $(-2, 8)$ and $(4, -4)$ respectively.

(a) Work out the gradient of the line AB.

..... **[2]**

(b) A straight line L is drawn on the grid.
Find the gradient of line L.



..... **[2]**

747 The points $A(-4, 7)$ and $B(2, -5)$ lie on a straight line.


(a) Calculate the gradient of the line AB.

..... **[2]**

(b) Point C has coordinates $(0, c)$.
The gradient of the line AC is equal to the gradient of the line AB.
Find the value of c.

$c =$ ..... **[2]**

748 The diagram shows two straight lines, L_1 and L_2, drawn on a coordinate grid.



(a) Find the gradient of line L_1.

..... **[2]**

(b) Find the gradient of line L_2.

..... **[2]**

(c) A straight line L_3 passes through the points $P(-3, 7)$ and $Q(5, -9)$.
Calculate the gradient of line L_3.

..... **[2]**

749 The points $A(-2, 5)$, $B(4, k)$, and $C(10, -1)$ lie in a coordinate plane.

(a) Given that the gradient of the line segment AB is $-\frac{1}{2}$, calculate the value of k.

$k =$ ..... **[3]**

(b) Using your value of k from part (a), show that the points A, B, and C lie on the same straight line.

[2]

(c) Point D has coordinates $(p, 3p - 1)$.
Find the value of p for which the line AD has a gradient of 2.

$p =$ ..... **[3]**

750 **(a)** Point P has coordinates $(2a, 3a)$ and point Q has coordinates $(5a, a + 4)$.

Find an expression for the gradient of the line PQ in terms of a.
Give your answer in its simplest form.

..... **[2]**

(b) The line PQ is perpendicular to a line with gradient 3.
Work out the value of a.

$a =$ ..... **[3]**

(c) The points $A(-1, 2)$, $B(3, 8)$ and $C(c, 23)$ lie on a straight line.
Work out the value of c.

$c =$ ..... **[3]**

751 The line L_1 passes through the points $P(-2, 3a)$ and $Q(4, a+6)$.

(a) Find an expression for the gradient of L_1 in terms of a.
Give your answer in its simplest form.

..... **[3]**

(b) The line L_2 has equation $y = 3x - 5$.
Given that L_1 is parallel to L_2, find the value of a.

$a =$ ..... **[2]**

(c) Using your value of a, find the coordinates of the midpoint of PQ.

(..... , ...) **[3]**

752 The points $A(1,2)$, $B(5,k)$ and $C(k,2k)$ are three distinct collinear points, where k is a constant.

(a) Write down an expression, in terms of k, for the gradient of the line segment AB.

..... **[1]**

(b) By finding an expression for the gradient of BC, show that $k^2 - 11k + 10 = 0$.

[3]

(c) Solve the equation to find the two possible values of k.

$k =$ ..... , k =

..... **[2]**

(d) Calculate the exact gradient of the line passing through point B and the origin $(0,0)$.

..... **[2]**

753 The points A, B and C have coordinates $(1, 3)$, $(4, 9)$ and $(8, y)$ respectively.

(a) The points A, B and C lie on the same straight line.
Work out the value of y.

$y =$ ..... **[2]**


(b) A different point D has coordinates $(p, p^2 - p - 3)$.
The gradient of the line AD is 2.
Work out the possible values of p.

$p =$ ..... , $p =$ ..... **[4]**

(c) A line L has gradient m.
The line L passes through the points $(3, m)$ and $(m + 1, 12)$.
Find the two possible values of m.

$m =$ ..... , $m =$ ..... **[4]**

754 **(a)** The diagram shows a straight line L passing through the points $P(-2, 1)$ and $Q(4, 5)$.
Find the gradient of line L.



..... **[2]**

(b) A different line passes through the points $(3, k)$ and $(8, -2)$.
The gradient of this line is -1.4.
Calculate the value of k.

$k =$ ..... **[3]**

755 Point A has coordinates $(-3, 5)$ and point B has coordinates $(4, -9)$.

(a) Calculate the gradient of the line AB.

..... **[2]**

(b) Point C has coordinates $(k, 1)$.
The gradient of the line AC is equal to the gradient of the line AB.
Find the value of k.

$k =$ ..... **[3]**

756 Point P has coordinates $(k-3, 2k)$ and point Q has coordinates $(2, 5)$.

(a) The gradient of the line PQ is -3.
Work out the value of k.

$k =$ ..... **[3]**

(b) Using your value of k from part (a), find the coordinates of the midpoint of PQ.

(..... , ...) **[2]**

757 A straight line passes through the points $P(3k, 4)$ and $Q(k-2, 8)$.
The gradient of this line is $-\frac{1}{2}$.

(a) Find the value of k.

$k =$ ..... **[3]**


(b) Write down the coordinates of P and Q.

$P =$ ..... , $Q =$ ..... **[1]**

(c) Another point R lies on the y-axis such that the gradient of the line passing through P and R is 1.
Find the coordinates of R.

(..... , ...) **[3]**

758 The diagram shows a sketch of the curve $y = x^2 - 3x + 2$ and two points, $P(1, 0)$ and $Q(t, t^2 - 3t + 2)$, on the curve, where $t > 1$.



(a) Show that the gradient of the chord PQ simplifies to $t - 2$.

[3]


(b) Find the coordinates of point Q when the gradient of the chord PQ is 2.8.

$Q = (\ldots\ldots\ ,\ \ldots\ldots)$ **[3]**

(c) Point R has coordinates $(3.2, -1.5)$.
Find the value of t, where $t > 1$, for which the gradient of QR is equal to the gradient of PQ.
Give your answer correct to 3 significant figures.

$t =$ ..... **[4]**

759 The points $A(-2, 5)$ and $B(4, -3)$ are shown on the coordinate grid.



(a) Find the coordinates of the midpoint of the line segment AB.

((..... , ...)) **[2]**

(b) Calculate the exact length of the line segment AB.

..... **[3]**

760 Point A has coordinates $(1, 4)$ and point B has coordinates $(7, -4)$.

(a) Calculate the coordinates of the midpoint of the line segment AB.

(..... , ...) **[2]**

(b) Calculate the length of the line segment AB.

..... **[3]**

761 The points A and B have coordinates $(2, -3)$ and $(8, 5)$ respectively.


(a) Find the coordinates of the midpoint of the line segment AB.

(..... , ...) **[2]**

(b) Calculate the exact length of the line segment AB.

..... **[3]**

762 The diagram shows a line segment AB plotted on a Cartesian coordinate grid.



(a) Write down the coordinates of point A and point B.

$A =$ ..... , $B =$ ..... **[2]**

(b) Find the coordinates of the midpoint M of the line segment AB.

$M =$ (..... , ...) **[2]**

(c) Calculate the length of the line segment AB.

..... units **[2]**

763 A circle has its centre at the origin, $O(0, 0)$.
The point $P(-3, 4)$ lies on the circumference of this circle.

(a) Work out the radius of the circle.

..... **[2]**

(b) The line segment PQ is a diameter of the circle.
Find the coordinates of Q.

(..... , ...) **[2]**

(c) The point R has coordinates $(k, \sqrt{11})$.
R also lies on the circumference of the circle, and k is a negative number.
Work out the exact value of k.

$k =$ ..... **[3]**

764 The line segment PQ has midpoint $M(3, -1)$.
The coordinates of point P are $(-2, 4)$.

(a) Find the coordinates of point Q.

$Q = (\ldots\ldots\ldots\ ,\ \ldots\ldots\ldots)$ **[2]**

(b) Calculate the exact length of PQ, giving your answer in the form $k\sqrt{2}$, where k is an integer.

$PQ = \ldots\ldots\ldots\ldots\ldots$ **[3]**

(c) Point R lies on the y-axis such that $PR = QR$.
Find the coordinates of R.

$R = (\ldots\ldots\ldots\ ,\ \ldots\ldots\ldots)$ **[4]**

765 Points A and B have coordinates $(3p, p-2)$ and $(p, 2p+1)$ respectively, where p is a constant.

(a) The midpoint of AB lies on the line $y = x + 1$.
Find the value of p.

$p =$ ..... **[3]**

(b) Given instead that the length of the line segment AB is 6, show that $5p^2 + 6p - 27 = 0$.

[3]

(c) Solve the equation $5p^2 + 6p - 27 = 0$ to find the two possible values of p.

$p =$ ..... , $p =$ ..... **[3]**

766 A circle has diameter AB, where A is the point $(-1, 2)$ and B is the point $(5, 8)$.

(a) Find the coordinates of the centre of the circle, C.

(..... , ...) **[2]**

(b) Find the exact radius of the circle.
Give your answer in the form $a\sqrt{b}$, where a and b are positive integers and b is as small as possible.

..... **[3]**

(c) Point D is such that the midpoint of CD is $M\left(\frac{5}{2}, \frac{3}{2}\right)$.

Find the exact length of AD, giving your answer in the form $c\sqrt{d}$, where c and d are integers and d is as small as possible.

..... **[5]**

767 The points A and B have coordinates $(-2, 5)$ and $(4, -3)$ respectively.

(a) Calculate the exact length of the line segment AB.

..... **[3]**

(b) Find the coordinates of the midpoint of AB.

(..... , ...) **[2]**

768 Point P has coordinates $(3, k)$ and point Q has coordinates $(7, 5)$.
The length of the line segment PQ is 5 units.

(a) Find the two possible values of k.

$k =$ ..... , $k =$ ..... **[3]**

(b) M is the midpoint of PQ.
For the case where $k < 5$, find the coordinates of M.

((..... , ...)) **[2]**

769 The midpoint of the line segment PQ is $M(1, -2)$.
The coordinates of P are $(-3, 5)$.

(a) Find the coordinates of Q.

(..... , ...) **[2]**

(b) Calculate the length of the line segment PQ.
Give your answer correct to 3 significant figures.

..... **[3]**

770 Point P has coordinates $(-3, 5)$ and point Q has coordinates $(k, -1)$.
The length of the line segment PQ is 10 units.


(a) Find the two possible values of k.

$k =$ ..... , $k =$ ..... **[4]**

(b) Given that $k > 0$, point M is the midpoint of PQ.
Find the coordinates of M.

(..... , ...) **[2]**

771 The points $A(-3, 2)$ and $B(5, 8)$ are the endpoints of a diameter of a circle.



(a) Find the coordinates of the centre of the circle, C.

((..... , ...)) **[2]**

(b) Calculate the area of the circle.
Give your answer correct to 1 decimal place.

..... **[3]**

(c) The point $D(k, 8)$ lies on the circumference of the circle.
Find the possible values of k.

$k =$ ..... , $k =$ ..... **[3]**

772 The points $A(2p, p)$ and $B(3p, 4p)$ are joined by a straight line, where $p > 0$.

(a) The length of the line segment AB is $5\sqrt{10}$.
Find the value of p.

$p =$ ..... **[4]**


(b) Point C lies on the line segment AB such that the ratio $AC : CB = 2 : 3$.
Using your value of p from part (a), find the coordinates of C.

(..... , ...) **[3]**

(c) The point $D(x, y)$ is such that C is the midpoint of BD.
Work out the coordinates of D.

(..... , ...) **[2]**

773 The diagram shows a sketch of triangle ABC with vertices $A(-3, 2)$, $B(5, 6)$, and $C(9, -2)$.



(a) Show that triangle ABC is isosceles.

[4]

(b) M is the midpoint of AC.

(i) Find the coordinates of M.

$M = (\ldots\ldots\ ,\ \ldots\ldots)$ **[2]**

(ii) Calculate the length of BM, giving your answer correct to 3 significant figures.

$BM = \ldots\ldots\ldots\ldots\ldots$ **[2]**

(c) Calculate the area of triangle ABC.

..... square units **[2]**

(d) Point D is such that $ABCD$ is a square.
Find the coordinates of D.


$D = ($ ..... , ... $)$ **[2]**

774 Find the gradient and the y-intercept of the line with equation

$3x + 5y = 20$

Gradient = ... , y-intercept = ... **[3]**

775 The diagram shows a straight line L drawn on a coordinate grid.



(a) Find the equation of line L.
Give your answer in the form $y = mx + c$.

$y =$ ..... **[3]**

(b) Write the equation of line L in the form $ax + by = c$, where a, b and c are integers.

..... **[1]**

776 **(a)** Find the gradient and the y-intercept of the line $3x + 5y = 15$.

Gradient = ... , y-intercept = ... **[3]**

(b) Find the equation of the line parallel to $3x + 5y = 15$ that passes through the point $(0, -4)$. Give your answer in the form $y = mx + c$.

$y =$ ..... **[2]**

777 A straight line L has the equation $3x - 4y = 12$.


(a) Find the gradient and the y-intercept of line L.

Gradient = ... , y-intercept = ... **[3]**

(b) Another line M is parallel to L and passes through the point $(0, 5)$.
Write down the equation of line M in the form $y = mx + c$.

$y =$ ..... **[2]**

778 The diagram shows a straight line L drawn on a coordinate grid.



(a) Find the equation of line L in the form $y = mx + c$.

$y =$ ..... **[3]**

(b) A vertical line M passes through the point $(-2, 4)$.
Write down the equation of line M.

..... **[1]**

(c) Find the coordinates of the point where line L and line M intersect.

(..... , ...) **[2]**

779 Line L_1 has equation $4x + 5y = 20$.

(a) Find the gradient and the y-intercept of line L_1.

Gradient = ... , y-intercept = ... **[3]**

(b) Find the coordinates of the point where line L_1 crosses the x-axis.

(..... , ...) **[2]**

(c) Line L_2 passes through the points $(-3, -5)$ and $(3, 7)$.
Find the equation of line L_2 in the form $ax + by = c$, where a, b, and c are integers.

..... **[3]**

780 A is the point $(p, -1)$ and B is the point $(5, 2p - 1)$.
The gradient of the line AB is 3.

(a) Find the value of p.

$p =$ ..... **[3]**

(b) Find the equation of the line AB.
Give your answer in the form $y = mx + c$.

..... **[2]**

(c) The line AB intersects the x-axis at the point C and the y-axis at the point D.
Work out the area of triangle COD, where O is the origin.

..... **[3]**

781 A straight line L passes through the points $A(-2, 3k)$ and $B(4, k)$, where k is a non-zero constant.

(a) Find the gradient of line L in terms of k.
Give your answer in its simplest form.

..... **[2]**

(b) Find the equation of line L in the form $ax + by = c$, where a, b, and c are expressions in terms of k.

..... **[3]**

(c) Line L also passes through the point $(1, 10)$.
Find the value of k.

$k =$ ..... **[2]**

(d) Hence, write down the coordinates of the x-intercept of line L.

(..... , ...) **[2]**

782 The line L passes through the points $A(2a, a+4)$ and $B(a-1, 3a+2)$.
The gradient of line L is -4.

(a) Find the value of a.

$a =$ ..... **[3]**

(b) Find the equation of line L.
Give your answer in the form $px + qy = r$, where p, q and r are integers and $p > 0$.

..... **[3]**

(c) Line L intersects the line $y = 3x - 2$ at the point M.
Find the coordinates of M.

(..... , ...) **[3]**

783 A is the point $(-3, 1)$ and B is the point $(5, 5)$.

(a) Find the equation of the perpendicular bisector of the line segment AB.
Give your answer in the form $y = mx + c$.

$y =$ ..... **[4]**


(b) The perpendicular bisector of AB intersects the line $x - 2y = 5$ at the point P.
Find the coordinates of P.

(..... , ...) **[3]**

(c) Find the exact length of AP.

..... **[3]**

784 The diagram shows a line L drawn on a coordinate grid.



(a) Work out the gradient of line L.

..... **[2]**

(b) Find the equation of line L.
Give your answer in the form $ax + by = c$, where a, b and c are integers.

..... **[3]**

785 Point A has coordinates $(-4, 3)$ and point B has coordinates $(6, -2)$.

(a) Find the equation of the line passing through A and B.
Give your answer in the form $y = mx + c$.

$y =$ ..... **[3]**


(b) The line $3x - 5y = 15$ intersects the x-axis at point C and the y-axis at point D.
Find the coordinates of C and D.

$C =$ ..... , $D =$ ..... **[2]**

(c) Write down the equation of the line that is parallel to the y-axis and passes through point A.

..... **[1]**

786 The diagram shows a straight line L.




(a) Find the equation of line L.
Give your answer in the form $ax + by = c$, where a, b, and c are integers.

..... **[3]**

(b) Line M is perpendicular to line L and passes through the point $(9, -1)$.
Find the equation of line M in the form $y = mx + c$.

$y =$ ..... **[3]**

787 The diagram shows a straight line L_1 drawn on a coordinate grid.




(a) Find the equation of line L_1 in the form $y = mx + c$.

$y =$ ..... **[3]**

(b) Line L_2 passes through the point $(2, 7)$ and the midpoint of the segment of L_1 that lies between the axes.
Find the equation of L_2 in the form $ax + by = c$, where a, b, and c are integers.

..... **[4]**

788 The diagram shows a sketch of two straight lines, L_1 and L_2, which intersect at the point P.
Line L_1 has equation $3x + 2y = 24$ and crosses the x-axis at A and the y-axis at B.
Line L_2 has equation $y = 1.5x - 3$ and crosses the x-axis at C.



(a) Write down the coordinates of A, B, and C.

$A =$ ..... , $B =$ ..... , $C =$ ..... **[3]**

(b) Find the coordinates of the point of intersection P.

$P =$ (..... , ...) **[3]**

(c) Calculate the area of quadrilateral $OCPB$.

..... square units **[3]**

(d) A third line L_3 has equation $kx - 4y = 14$ and also passes through point P.
Calculate the value of k.

$k =$ ..... **[2]**

789 Find the equation of the line that is parallel to $y = -2x + 7$ and passes through the point $(3, 4)$.
Give your answer in the form $y = mx + c$.

$y =$ ..... **[2]**

790 Line L has the equation $4x + 2y = 9$.
Find the equation of the line parallel to L that passes through the point $(3, -1)$.
Give your answer in the form $y = mx + c$.


$y =$ ..... **[3]**

791 Line L has the equation $y = 6 - 3x$.

Find the equation of the line parallel to L that passes through the point $(-2, 5)$.
Give your answer in the form $y = mx + c$.

$y =$ ..... **[3]**

792 The diagram shows two parallel straight lines, L_1 and L_2, on a coordinate grid.
Line L_1 has equation $y = 2x + 1$ and line L_2 passes through the point $(2, 1)$.



(a) Write down the gradient of line L_1.

..... **[1]**

(b) Find the equation of line L_2 in the form $y = mx + c$.

$y =$ ..... **[2]**

(c) A third line L_3 is parallel to L_1 and passes through the point $(-1, 5)$.
Find the equation of line L_3 in the form $y = mx + c$.

$y =$ ..... **[2]**

793 A straight line L_1 has the equation $2y - 5x = 8$.

(a) Find the gradient of line L_1.

..... **[2]**

(b) Line L_2 is parallel to L_1 and passes through the point $(0, -3)$.
Write down the equation of L_2.

$y =$ ..... **[2]**

(c) Line L_3 is parallel to L_1 and passes through the point $(4, 1)$.
Find the equation of L_3 in the form $ax + by = c$, where a, b, and c are integers.

..... **[3]**

794 The points A and B have coordinates $(-3, 5)$ and $(5, -1)$ respectively.
Line L is parallel to the line segment AB and passes through the point $(4, -2)$.

(a) Find the gradient of the line segment AB.

..... **[2]**

(b) Find the equation of line L.
Give your answer in the form $y = mx + c$.

$y =$ ..... **[3]**

795 Line L_1 has equation $3x + 4y = 12$.

(a) Find the gradient of line L_1.

..... **[2]**

(b) Line L_2 is parallel to line L_1 and passes through the point $(4, 3)$.
Find the equation of line L_2 in the form $ax + by = c$, where a, b, and c are integers.

..... **[3]**

(c) Line L_2 crosses the x-axis at A and the y-axis at B.
Find the coordinates of the midpoint of AB.

(..... , ...) **[3]**

796 Line P has equation $4x - ky = 8$.
Line Q has equation $y = (k-3)x + 2$.
The lines P and Q are parallel.

(a) Find the two possible values of the constant k.

$k =$ ..... **[4]**

(b) Given that line P has a positive gradient,

(i) find the equation of line P in the form $y = mx + c$.

$y =$ ..... **[2]**

(ii) A line R is parallel to line P and passes through the point $(-5, 7)$.
Find the equation of line R in the form $ax + by = c$, where a, b, and c are integers.

..... **[2]**

797 Line M has the equation $5x - 2y = 15$.
Point P has coordinates $(-4, 1)$.
Find the equation of the line parallel to M that passes through P.
Give your answer in the form $y = mx + c$.

$y =$ ..... **[3]**

798 A is the point $(-3, 2)$ and B is the point $(5, 6)$.


(a) Calculate the gradient of the line AB.

..... **[2]**

(b) Line L is parallel to AB and passes through the point $(4, -1)$.
Find the equation of line L, giving your answer in the form $ax + by = c$, where a, b and c are integers.

..... **[3]**

799 The diagram shows the points $A(-2, 1)$ and $B(4, 4)$ joined by a straight line.



(a) Calculate the gradient of the line AB.

..... **[2]**

(b) Line M passes through the point $(2, -3)$ and is parallel to the line AB.
Find the equation of line M in the form $y = mx + c$.

$y =$ ..... **[3]**

(c) Line M intersects the x-axis at point P and the y-axis at point Q.
Calculate the length of the line segment PQ.
Give your answer correct to 3 significant figures.

..... **[4]**

800 Points $A(-1, 5)$ and $B(3, -3)$ lie on line L_1.
Line L_2 is parallel to L_1 and passes through the point $C(2, 7)$.
Line L_2 crosses the x-axis at D and the y-axis at E.

(a) Find the equation of line L_2.
Give your answer in the form $y = mx + c$.

$y =$ ..... **[4]**

(b) Calculate the exact area of triangle ODE, where O is the origin $(0, 0)$.

..... $units^2$ **[3]**

801 Line L_1 has the equation $3x - 4y = 12$.
Line L_2 is parallel to L_1.


Line L_2 crosses the x-axis at point A and the y-axis at point B.
The length of the line segment AB is 15.

Find the two possible equations of L_2.
Give your answers in the form $y = mx + c$.

.....

..... **[7]**

802 The points $A(1,2)$, $B(7,4)$, and $C(10,9)$ are three vertices of a parallelogram $ABCD$.
Line CD is parallel to line AB and line AD is parallel to line BC.



Not to scale

(a) Find the equation of the line CD.

..... **[3]**


(b) Find the equation of the line AD.

..... **[3]**

(c) Work out the coordinates of the vertex D.

(..... , ...) **[3]**

803 The diagram shows a sketch of a parallelogram $ABCD$ on a coordinate grid.
Vertex A has coordinates $(-3, 2)$ and vertex B has coordinates $(5, 4)$.
The line passing through B and C has equation $5x - 2y = 17$.



(a) Find the equation of the line passing through A and D in the form $y = mx + c$.

$y =$ ..... **[3]**

(b) Vertex D lies on the y-axis.
Write down the coordinates of D and find the coordinates of vertex C.

$D =$ ..... , $C =$ ..... **[3]**

(c) Line L is parallel to AB and passes through the midpoint of BC.
Find the equation of line L in the form $ax + by = c$, where a, b, and c are integers.

..... [4]

(d) Calculate the length of diagonal AC, giving your answer correct to 3 significant figures.

$AC =$ ..... [2]

804 Line L has the equation $5y + 2x = 15$.
Find the gradient of a line that is perpendicular to L.

..... **[2]**

805 Find the equation of the line that is perpendicular to $y = 3x - 5$ and passes through the point $(6, 2)$. Give your answer in the form $y = mx + c$.

$y =$ ..... **[3]**

806 Line L_1 has the equation $y = 5 - 2x$.
Line L_2 is perpendicular to L_1 and passes through the point $(4, 1)$.


(a) Find the gradient of line L_1.

..... **[1]**

(b) Find the equation of line L_2.
Give your answer in the form $y = mx + c$.

$y =$ ..... **[3]**

807 The diagram shows a straight line L_1 drawn on a coordinate grid.



(a) Write down the gradient of line L_1.

..... **[1]**

(b) Find the gradient of a line that is perpendicular to L_1.

..... **[1]**

(c) Line L_2 is perpendicular to L_1 and passes through the point $(4, 3)$.
Find the equation of line L_2 in the form $y = mx + c$.

$y =$ ..... **[3]**

808 A straight line L has the equation $2y = 5x + 8$.

(a) Find the gradient of line L.

..... **[2]**

(b) Find the gradient of a line that is perpendicular to line L.

..... **[1]**

(c) A line M is perpendicular to line L and passes through the point $(10, -3)$.
Find the equation of line M in the form $y = mx + c$.

$y =$ ..... **[3]**

809 A is the point $(-4, 7)$ and B is the point $(2, -1)$.

Find the equation of the perpendicular bisector of the line segment AB.
Give your answer in the form $ax + by = c$, where a, b and c are integers.

..... **[5]**

810 Line L has equation $2x + 3y = 12$.
Point A has coordinates $(-3, 8)$ and point B has coordinates $(9, -2)$.

(a) Find the gradient of a line that is perpendicular to line L.

..... **[2]**

(b) For the line segment joining $A(-3, 8)$ and $B(9, -2)$,

(i) find the coordinates of the midpoint of AB,

(..... , ...) **[2]**

(ii) find the equation of the perpendicular bisector of AB, giving your answer in the form $ax + by = c$, where a, b, and c are integers.

..... **[4]**

811 A is the point $(-3, 8)$ and B is the point $(9, -2)$.


(a) Find the coordinates of the midpoint of AB.

(..... , ...) **[2]**

(b) Find the equation of the perpendicular bisector of AB.
Give your answer in the form $ax + by = c$, where a, b and c are integers.

..... **[4]**

812 Line L_1 has the equation $3x + 2y = 12$.
Line L_2 is perpendicular to L_1 and intersects L_1 at a point P on the y-axis.



Not to scale

(a) Find the coordinates of point P.

(..... , ...) **[1]**

(b) Find the equation of line L_2.
Give your answer in the form $y = mx + c$.

$y =$ ..... **[3]**

(c) Point Q lies on L_2 and has an x-coordinate of 9.
Point R lies on L_1 and has an x-coordinate of 4.
Work out the area of triangle PQR.

..... **[5]**

813 Line L_1 has equation $kx + 2y = 12$.
Line L_2 passes through the points $A(0, 5)$ and $B(2, k + 2)$.
The constant k is a non-zero integer.

(a) Find the gradient of line L_1 in terms of k.

..... **[1]**

(b) Find the gradient of line L_2 in terms of k.

..... **[2]**

(c) Given that L_1 and L_2 are perpendicular, find the two possible values of k.

$k =$ ..... **[4]**

(d) Given that the gradient of line L_1 is positive, find the equation of line L_2 in the form $ax + by = c$, where a, b, and c are integers.

..... **[3]**

814 The equation of line P is $3x - 4y = 12$.
Line Q is perpendicular to line P and passes through the point $(6, -2)$.

Find the equation of line Q.
Give your answer in the form $y = mx + c$.

$y =$ ..... **[4]**

815 The points A and B have coordinates $(-2, 4)$ and $(6, 0)$ respectively.

(a) Find the coordinates of the midpoint of AB.

(..... , ...) **[2]**

(b) Find the equation of the perpendicular bisector of the line segment AB.
Give your answer in the form $y = mx + c$.

$y =$ ..... **[4]**

816 The coordinates of point A are $(-4, 5)$ and the coordinates of point B are $(2, -7)$.

(a) Find the coordinates of the midpoint of the line segment AB.

(..... , ...) **[2]**

(b) Calculate the gradient of the line AB.

..... **[2]**

(c) Find the equation of the perpendicular bisector of the line segment AB.
Give your answer in the form $y = mx + c$.

$y =$ ..... **[3]**

(d) The perpendicular bisector intersects the x-axis at point P.
Find the coordinates of P.

(..... , ...) **[2]**

817 The line L_1 has equation $2y + 3x = 12$.
The line L_2 is perpendicular to L_1 and intersects L_1 at the point P on the y-axis.

(a) Find the coordinates of P.

(..... , ...) **[2]**


(b) Find the equation of line L_2.
Give your answer in the form $y = mx + c$.

$y =$ ..... **[3]**

(c) Line L_2 crosses the x-axis at the point Q.
Calculate the exact length of PQ.

..... **[4]**

818 The diagram shows a sketch of triangle ABC with vertices $A(-1, 2)$ and $B(5, 5)$.
Angle $ABC = 90°$ and vertex C lies on the x-axis.



(a) Find the equation of the line passing through B and C in the form $y = mx + c$.

$y =$ ..... **[3]**

(b) Find the coordinates of vertex C.

$C =$ (..... , ...) **[2]**

(c) Calculate the area of triangle ABC.


..... square units **[3]**

(d) The perpendicular bisector of AC intersects the y-axis at point P.

Find the coordinates of P, giving the y-coordinate correct to 3 significant figures.

$P = (\ldots\ldots\ldots\ ,\ \ldots\ldots\ldots)$ **[4]**

819 The diagram shows a regular polygon $ABCDEF$ with centre O.



(a) Write down the mathematical name of the polygon $ABCDEF$.

..... **[1]**


(b) Write down the mathematical name for the angle AFE.

..... **[1]**

(c) Write down the mathematical name for the triangle AOB.

..... **[1]**

820 The diagram shows a circle with centre O.
The straight line L touches the circle at the point P.



(a) Write down the mathematical name for the straight line CD.

..... **[1]**

(b) Write down the mathematical name for the straight line L.

..... **[1]**


(c) Write down the mathematical name for the shaded region OAB.

..... **[1]**

(d) The angle AOB is $60°$.
Write down the mathematical name for this type of angle.

..... **[1]**

821 The diagram shows a circle, centre O.
A and B are points on the circumference of the circle.
The line passing through T touches the circle at B.



(a) Write down the geometrical name of the line AB.

..... **[1]**

(b) Write down the geometrical name of the line BT.

..... **[1]**


(c) Write down the geometrical name of the shaded region.

..... **[1]**

(d) Write down the mathematical name of triangle OAB.

..... **[1]**

822 The diagram shows a circle with centre O.
The lines AB and CD intersect at E.
AB touches the circle at P, and CD passes through the centre O.



(a) Write down the mathematical name for the line segment OP.

..... **[1]**

(b) Write down the mathematical name for the line AB.

..... **[1]**

(c) Write down the mathematical name for the line segment CD.

..... **[1]**

(d) State the size of angle OPT.

..... ° **[1]**

823 **(a)** **(i)** A polygon has interior angles that sum to 1080°.

Work out the number of sides of this polygon.

..... **[2]**

(ii) Write down the mathematical name of this polygon.

..... **[1]**


(b) Write down the mathematical name of a quadrilateral with all four sides equal in length and no right angles.

..... **[1]**

(c) Write down the mathematical name of a 3D solid that has a uniform cross-section and two parallel circular faces.

..... **[1]**

824 The diagram shows a circle, centre O, with diameter AB.
The points A, B, C, and T lie on the circumference of the circle.
The straight line PQ touches the circle at T.



(a) Write down the mathematical name for

(i) the straight line PQ,

..... **[1]**

(ii) the straight line BC,

..... **[1]**


(iii) the shaded region bounded by BC and the minor arc BC.

..... **[1]**

(b) State the geometrical word that describes the relationship between radius OT and line PQ, and write down the size of angle OTQ.

Relationship: ..... , Angle $OTQ =$ ..... **[2]**

825 The diagram shows a circle with centre O.
The line AB touches the circle at the point C.
The line OD passes through C.



(a) Write down the mathematical name for the straight line AB.

..... **[1]**

(b) Write down the mathematical name for the straight line OC.

..... **[1]**


(c) Write down the size of angle OCB.

Angle OCB = ... ° **[1]**

(d) The length of OC is 5 cm.
Calculate the circumference of the circle.
Give your answer in terms of π.

..... cm **[2]**

826 Points A and B lie on a circle, centre O.
The lines TA and TB are tangents to the circle.
The region enclosed by the arc AB and the radii OA and OB is shaded.



(a) Write down the mathematical name for the quadrilateral $OATB$.

..... **[1]**

(b) Write down the mathematical name for the shaded region.

..... **[1]**


(c) Write down the mathematical name for the straight line AB.

..... **[1]**

(d) Triangle OAT and triangle OBT are exactly the same size and shape.
Write down the mathematical term for two identical shapes.

..... **[1]**

827 The diagram shows a quadrilateral $ABCD$.
$AB = AD$ and $CB = CD$.
The diagonals AC and BD intersect at M.



(a) Write down the mathematical name of the quadrilateral $ABCD$.

..... **[1]**

(b) Write down the mathematical term that describes the relationship between the line AC and the line BD.

..... **[1]**

(c) Triangle ABM and Triangle ADM are identical in shape and size.
Write down the mathematical word for this.

..... **[1]**

(d) The angle ABC is $110°$.
Write down the mathematical name for this type of angle.

..... **[1]**

(e) Write down the mathematical name for the line segment AC in relation to the line segment BD.

..... **[1]**

828 A regular polygon has n sides.
Each interior angle of this polygon is 156°.

(a) Work out the size of one exterior angle of this polygon.

..... ° **[1]**

(b) Find the value of n.

$n =$ ..... **[2]**


(c) Write down the mathematical name for a regular polygon with 10 sides.

..... **[1]**

(d) State whether a regular 15-sided polygon has an acute, obtuse, or reflex interior angle.

..... **[1]**

829 The diagram shows a circle, centre O.
P, Q and R are points on the circumference.
The straight line ST touches the circle at P.



(a) Write down the mathematical name of the line ST.

..... **[1]**

(b) Write down the mathematical name of the line segment QR.

..... **[1]**

(c) Write down the mathematical name of the shaded region bounded by the line QR and the minor arc QR.

..... **[1]**


(d) Write down the mathematical name of the shaded region bounded by the radii OP and OQ and the minor arc PQ.

..... **[1]**

(e) The angle marked w is greater than $180°$ and less than $360°$.
Write down the geometrical term for this type of angle.

..... **[1]**

830 In the diagram, $ABCD$ is a quadrilateral with AB parallel to DC.
The side DC is extended to E, and AC is a diagonal.
Angle $ADC = 65°$, angle $BAC = 34°$, and angle $ABC = 112°$.



(a) Write down the special mathematical name for quadrilateral $ABCD$.

..... **[1]**

(b) For vertex B,

(i) write down the geometrical term that describes the interior angle ABC,

..... **[1]**

(ii) calculate the size of the reflex angle at vertex B.

..... ° **[1]**

(c) Calculate the size of

(i) angle ACD,

Angle $ACD =$ ..... ° **[1]**

(ii) the exterior angle BCE.

Angle $BCE =$ ..... ° **[2]**

831 **(a)** The interior angle of a regular polygon is 144°.

(i) Work out the number of sides of this polygon.

..... **[2]**

(ii) Write down the geometrical name of this polygon.

..... **[1]**

(b) A hexagon has interior angles of $x°$, $2x°$, $(x + 30)°$, $(3x + 60)°$, 110° and 100°.

(i) Calculate the value of x.

$x =$ ..... **[3]**


(ii) Write down the geometrical name for an angle of $(x + 30)°$.

..... **[1]**

(iii) Write down the geometrical name for an angle of $(3x + 60)°$.

..... **[1]**

832 The diagram shows a regular pentagon $ABCDE$ and a regular hexagon $AEFGHI$ joined along the edge AE.



(a) Calculate the size of an interior angle of the regular pentagon $ABCDE$.

..... ° **[2]**


(b) Calculate the size of an interior angle of the regular hexagon $AEFGHI$.

..... ° **[2]**

(c) Calculate the size of the reflex angle BAI.

..... ° **[3]**

833 The diagram shows a large solid cone with a smaller solid cone removed from the top.
The cut is made parallel to the base of the large cone.
All dimensions are in centimetres.



(a) Write down the mathematical name of the solid with height 20.

..... **[1]**

(b) The small cone is geometrically related to the original large cone.

(i) Write down the mathematical word that describes the relationship between the small cone and the large cone.

..... **[1]**

(ii) Work out the value of the radius, r.

$r =$ ..... **[2]**

(iii) Calculate the slant edge of the solid with height 20.

..... *cm* **[3]**

834 A map shows three towns A, B and C.
Town B is 45 km from A on a bearing of 062°.
Town C is 45 km from A on a bearing of 138°.

(a) Write down the mathematical name of triangle ABC.

..... **[1]**

(b) Work out the bearing of A from C.

..... ° **[2]**

(c) **(i)** Point M is the midpoint of the straight line BC.
Write down the mathematical name of the line AM in relation to the line BC.

..... **[1]**


(ii) Calculate the bearing of M from A.

..... ° **[2]**

(d) A fourth town D is located such that the line AB is parallel to the line DC and the line AD is not parallel to the line BC.
Write down the mathematical name of the quadrilateral $ABCD$.

..... **[1]**

835 The diagram shows a solid frustum obtained by cutting a smaller right circular cone of height h cm from a larger right circular cone of height $(h + 12)$ cm.
The radius of the top circular face is r cm and the radius of the bottom circular base is $(r + 5)$ cm.
The two cones are mathematically similar.



(a) Given that the scale factor relating the linear dimensions of the smaller cone to the larger cone is $\frac{3}{5}$, show that $h = 18$.

[3]


(b) Find the value of r.

$r =$ ..... **[2]**

(c) The ratio of the volume of the smaller cone to the volume of the frustum is $27 : k$.
Find the value of k.

$k =$ ..... **[2]**

836 The diagram shows a frustum of a cone.
The radius of the circular base is 12 cm and the radius of the circular top is 4 cm.
The vertical height of the frustum is 15 cm.



(a) Calculate the volume of the frustum.
The volume, V, of a cone with radius r and height h is $V = \frac{1}{3}\pi r^2 h$.

..... cm^3 **[5]**

(b) A different shape is geometrically similar to the frustum.
The ratio of the surface area of this shape to the surface area of the frustum is 9 : 25.
Calculate the vertical height of this shape.

..... cm **[3]**

837 A regular polygon has n sides.

(a) The interior angle of this regular polygon is 4 times the size of its exterior angle.
Calculate the value of n and write down the mathematical name of this polygon.

$n =$ ..... , Name = ... **[4]**

(b) A solid uniform prism has a cross-section in the shape of this regular polygon.
Write down the number of faces, edges, and vertices of this prism.


Faces = ... , Edges = ... , Vertices = ... **[3]**

(c) Two mathematically similar cones have base areas in the ratio 9 : 16.
The volume of the smaller cone is 108π cm^3.

Calculate the volume of the larger cone.
Give your answer in terms of π.

..... cm^3 **[3]**

838 The diagram shows triangle ABC.
D is a point on AB and E is a point on AC such that DE is parallel to BC.
$AD = 6$ cm, $DB = 4$ cm, $AE = 7.5$ cm and $BC = 15$ cm.



(a) Explain why triangle ADE is similar to triangle ABC.

[2]

(b) Calculate the length of DE.

..... cm **[2]**

(c) Calculate the length of EC.

..... cm **[2]**


(d) State the geometrical name of the quadrilateral $DBCE$.

..... **[1]**

(e) The area of triangle ADE is 18 cm^2.
Calculate the area of the quadrilateral $DBCE$.

..... cm^2 **[3]**

839 The diagram shows a solid formed by cutting a smaller right cone of height 6 cm from the top of a larger right cone of height 15 cm and base radius 10 cm. The cut is made along a plane parallel to the circular base.



(a) Write down the mathematical name for

(i) the remaining solid after the top cone is removed,

..... **[1]**

(ii) a solid pyramid that has 4 triangular faces, 6 edges, and 4 vertices.

..... **[1]**

(b) Explain why the removed cone is similar to the original large cone, and write down the scale factor that maps the original large cone onto the removed cone.

Reason: ..... , Scale factor = ... **[2]**

(c) Calculate the radius of the upper circular face of the remaining solid.

..... cm **[2]**

(d) A regular polygon has an exterior angle of 36°.

(i) Write down the mathematical name of this regular polygon.

..... **[2]**

(ii) A right prism has this regular polygon as its cross-section.
State the total number of faces and the total number of edges of this prism.


Faces = ... , Edges = ... **[2]**

840 The bearing of point Y from point X is 048°.

Work out the bearing of point X from point Y.

..... ° **[2]**


841 The diagram shows the positions of a ship, S, and a lighthouse, L.



The bearing of the lighthouse from the ship is 115°.
Work out the bearing of the ship from the lighthouse.

..... ° **[2]**

842 The diagram shows the positions of three towns, A, B and C.
B is due north of A and the bearing of C from A is 058°.



(a) Find the bearing of A from C.

..... ° **[2]**

(b) Given that angle $ABC = 84°$, find the bearing of C from B.

..... ° **[1]**

843 The positions of three towns, A, B and C, are marked on a map.


(a) The bearing of town B from town A is 068°.
Work out the bearing of town A from town B.

..... ° **[2]**

(b) Town C is due west of town B.
Write down the bearing of town C from town B.

..... ° **[1]**

844 The bearing of a lighthouse L from a port P is 128°.
A boat B is positioned such that it is due South of P and due West of L.



(a) Find the bearing of P from L.

..... ° **[2]**

(b) Work out the size of angle BPL.

..... ° **[1]**

(c) Work out the size of angle PLB.

..... ° **[2]**

845 The bearing of a lighthouse, L, from a boat, B, is 148°.

(a) Work out the bearing of B from L.

..... ° **[2]**

(b) A port, P, is due north of L and due east of B.


(i) Write down the bearing of B from P.

..... ° **[1]**

(ii) Work out the size of angle LBP.

..... ° **[2]**

846 The diagram shows the positions of three points, P, Q and R.
Point Q is due south of point P.
Point R is due east of point Q.
The bearing of point R from point P is 124°.




(a) Find the bearing of point P from point R.

..... ° **[2]**

(b) Calculate the angle PRQ.

Angle PRQ = ... ° **[2]**

847 The diagram shows the positions of three points P, Q and R.
The bearing of Q from P is 115°.
The bearing of R from Q is 205°.




(a) Show that angle $PQR = 90°$.

[2]

(b) The distance PQ is 15 km and the distance QR is 8 km.
Calculate the distance PR.

..... km **[3]**

848 A ship sails from port P on a bearing of 125° for 45 km to point Q.
It then sails from Q on a bearing of 240° for 60 km to point R.




(a) Show that angle $PQR = 65°$.

[2]

(b) Calculate the distance PR.

$PR =$ ..... km **[3]**

849 The diagram shows the positions of three towns, A, B and C.
The bearing of B from A is 065°.
Angle $CAB = 85°$ and angle $ABC = 70°$.



(a) Calculate the bearing of C from A.

..... ° **[1]**

(b) Calculate the bearing of A from B.

..... ° **[2]**


(c) Calculate the bearing of C from B.

..... ° **[2]**

(d) Calculate angle ACB.

..... ° **[1]**

850 A ship sails 40 km from a port P to point Q on a bearing of 050°.
It then sails 60 km from Q to point R on a bearing of 140°.



(a) Show that the angle PQR is 90°.

[2]


(b) Calculate the distance PR.

..... km **[2]**

(c) Calculate the bearing of R from P.

..... ° **[3]**

851 A ship leaves a port A and travels 12 km on a bearing of 035° to point B.
From B, the ship travels 16 km on a bearing of 125° to point C.



(a) Show that angle $ABC = 90°$.

[2]


(b) Calculate the direct distance AC.

$AC =$ ..... km **[3]**

(c) Calculate the bearing of C from A.

..... ° **[3]**

852 Point B is on a bearing of 065° from point A.
Point C is on a bearing of 155° from point B.
The distance $AB = BC$.



(a) Show that angle $ABC = 90°$.

[2]


(b) Work out the bearing of A from C.

..... ° **[3]**

(c) Point D is such that the quadrilateral $ABCD$ is a square.
Find the bearing of D from A.

..... ° **[2]**

853 A helicopter flies from base A to base B, a distance of 120 km on a bearing of 060°.
It then flies from base B to base C, a distance of 180 km on a bearing of 170°.




(a) Calculate the distance AC.

..... km **[4]**

(b) Calculate the bearing of C from A.

..... ° **[3]**

854 Three towns are located at points P, Q, and R.
Town Q is 24 km from P on a bearing of 042°.
Town R is 35 km from P on a bearing of 118°.




(a) Calculate the distance QR.

$QR =$ ..... km **[3]**

(b) Calculate the bearing of R from Q.

..... ° **[4]**

855 Three lighthouses A, B and C are situated such that B is 12 km from A on a bearing of 042°. C is 18 km from A and the bearing of C from B is 145°.




(a) Calculate the bearing of C from A.

..... ° **[5]**

(b) A boat sails from A in a straight line such that it is always equidistant from B and C. Calculate the bearing on which the boat sails.

..... ° **[2]**

856 A ship leaves port P and sails 40 km on a bearing of 050° to reach point A.
It then sails 60 km on a bearing of 160° to reach point B.




(a) Calculate the distance PB.

..... km **[4]**

(b) Calculate the bearing of B from P.

..... ° **[4]**

857 A ship sails 45 km from Port P on a bearing of 050° to Point Q.
It then sails 60 km on a bearing of 120° to Point R.



(a) Calculate the distance PR.

..... km **[4]**


(b) Calculate the bearing of R from P.

..... ° **[3]**

(c) The ship sails directly back from R to P.
Calculate the bearing of P from R.

..... ° **[2]**

858 The diagram shows triangle PQR.
Point S lies on PQ and point T lies on PR.
The line ST is parallel to the line QR.




NOT TO SCALE

Given that $PS = 6$ cm, $SQ = 3$ cm and $ST = 8$ cm, calculate the length of QR.

$QR =$ ..... cm **[2]**

859 The diagram shows triangle ABC.
D is a point on AB and E is a point on AC such that DE is parallel to BC.
$AD = 4$ cm, $DB = 2$ cm and $DE = 5$ cm.




(a) Calculate the length of BC.

..... *cm* **[2]**

(b) The area of triangle ADE is 8 cm^2.
Work out the area of the trapezium $DBCE$.

..... *cm*^{2} **[2]**

860 Triangle ABC and triangle DEF are mathematically similar.



NOT TO SCALE

(a) Calculate the length DE.

$DE =$ ..... cm **[2]**

(b) The area of triangle DEF is 72 cm^2.
Calculate the area of triangle ABC.

..... cm^2 **[2]**

861 Two mathematically similar solid cylinders are shown as Cylinder A and Cylinder B.
Cylinder A has a height of 6 cm.
Cylinder B has a height of 15 cm.


(a) The volume of Cylinder A is 48 cm^3.
Calculate the volume of Cylinder B.

..... cm^3 **[2]**

(b) The curved surface area of Cylinder B is 200 cm^2.
Calculate the curved surface area of Cylinder A.

..... cm^2 **[2]**

862 Triangle ABC and triangle DEF are mathematically similar.
$AB = 3$ cm and $DE = 9$ cm.



NOT TO SCALE


(a) The length of BC is 5 cm.
Calculate the length of EF.

..... cm **[2]**

(b) The area of triangle DEF is 54 cm^2.
Calculate the area of triangle ABC.

..... cm^2 **[2]**

863 In the diagram, ADE is a triangle and DE is parallel to BC.
$AD = 4$ cm, $DB = 2$ cm and $DE = 5$ cm.




(a) Calculate the length of BC.

$BC =$ ..... cm **[2]**

(b) The area of triangle ADE is 8 cm^2.
Calculate the area of the trapezium $DBCE$.

Area = ... cm^2 **[3]**

864 In the diagram, ABC is a triangle and the line XY is parallel to BC.
$AX = 4$ cm, $XB = 2$ cm and $XY = 6$ cm.




(a) Calculate the length of BC.

$BC =$ ..... cm **[2]**

(b) The area of triangle AXY is 32 cm^2.
Calculate the area of the trapezium $XBCY$.

Area = ... cm^2 **[3]**

865 The diagram shows a triangle ABC.
D is a point on AB and E is a point on AC such that DE is parallel to BC.
$AD = 4$ cm, $DB = 2$ cm and $DE = 6$ cm.



(a) Calculate the length of BC.

..... cm **[2]**

(b) The area of triangle ADE is 12 cm^2.
Calculate the area of the trapezium $DBCE$.

..... cm^2 **[3]**

866 Two mathematically similar bottles have heights of 10 cm and 15 cm.


(a) The surface area of the smaller bottle is 120 cm^2.
Work out the surface area of the larger bottle.

..... cm^2 **[3]**

(b) The volume of the larger bottle is 540 cm^3.
Work out the volume of the smaller bottle.

..... cm^3 **[3]**

867 In the diagram, the line ST is parallel to QR.
$PS = 6$ cm, $SQ = 3$ cm and $ST = 8$ cm.
The length of PR is 12 cm.



(a) Calculate the length of QR.

..... cm **[2]**


(b) Calculate the length of PT.

..... cm **[2]**

(c) The area of triangle PST is 20 cm^2.
Calculate the area of the trapezium $SQRT$.

..... cm^2 **[3]**

868 Cone A and Cone B are mathematically similar.
The total surface area of Cone A is 20 cm^2 and the total surface area of Cone B is 45 cm^2.




(a) The volume of Cone A is 16 cm^3.
Calculate the volume of Cone B.

..... cm^3 **[3]**

(b) The height of Cone B is 12 cm.
Calculate the height of Cone A.

..... cm **[2]**

869 The diagram shows two similar cones, P and Q.
Cone P has a volume of 400 cm^3 and a height of 12 cm.
Cone Q has a volume of 1350 cm^3.




(a) Calculate the height of cone Q.

Height = ... cm **[3]**

(b) The curved surface area of cone Q is 360 cm^2.
Calculate the curved surface area of cone P.

Surface area = ... cm^2 **[2]**

870 The diagram shows a triangle ABC.
The point D lies on AB and the point E lies on AC.
The angle ADE is equal to the angle ACB.



(a) Explain why triangle ADE is mathematically similar to triangle ACB.

[3]


(b) It is given that $AD = x$ cm, $DB = (x + 1)$ cm, $AE = 4$ cm and $EC = 5$ cm.
Find the value of x.

$x =$ ..... **[4]**

(c) Find the ratio of the area of triangle ADE to the area of the quadrilateral $DBCE$.
Give your answer in its simplest form.

..... **[2]**

871 Triangles ABC and DEF are mathematically similar.
In the diagram, $AB = 4$ cm, $BC = 6$ cm and $DE = 14$ cm.




(a) Calculate the length of EF.

..... cm **[2]**

(b) The area of triangle ABC is 8.5 cm^2.
Calculate the area of triangle DEF.

..... cm^2 **[2]**

872 Two solid cones, A and B, are mathematically similar.



(a) Cone A has a volume of 150 cm^3 and a height of 12 cm.
Cone B has a volume of 1200 cm^3.
Calculate the height of cone B.

..... cm **[3]**

(b) The surface area of cone B is 440 cm^2.
Calculate the surface area of cone A.

..... cm^2 **[2]**

873 Two mathematically similar containers, A and B, have volumes of 320 cm^3 and 1080 cm^3 respectively.

(a) Container A has a height of 18 cm.

Calculate the height of container B.


Height = ... cm **[3]**

(b) Container B has a surface area of 630 cm^2.

Calculate the surface area of container A.

Surface area = ... cm^2 **[3]**

874 Two mathematically similar containers, A and B, are shown in the diagram.
Container A has a height of 12 cm and container B has a height of 30 cm.



NOT TO SCALE

(a) Container A has a volume of 320 cm^3.
Calculate the volume of container B.

Volume = ... cm^3 **[2]**

(b) Container B has a surface area of 1125 cm^2.
Calculate the surface area of container A.

Surface area = ... cm^2 **[2]**

(c) The containers are completely filled with liquid.
The cost of filling container A is \$4.80.
Assuming the cost is directly proportional to the volume, calculate the cost of filling container B.

..... **[2]**

875 Two mathematically similar solids, P and Q, have volumes of 320 cm^3 and 8640 cm^3 respectively.

(a) The height of solid P is 12 cm.
Calculate the height of solid Q.

..... cm **[3]**


(b) The surface area of solid Q is 2250 cm^2.
Calculate the surface area of solid P.

..... cm^2 **[2]**

(c) Solid P is made of a metal with a density of 8.5 g/cm^3.
Calculate the mass of solid P in kilograms.

..... kg **[2]**

876 The diagram shows triangle ADE with a line BC drawn parallel to DE.
$AB = 6$ cm, $BD = 4$ cm and $BC = 7.5$ cm.



NOT TO SCALE

(a) Explain why triangle ABC is mathematically similar to triangle ADE.

[2]


(b) Calculate the length of DE.

$DE =$ ..... cm **[2]**

(c) The area of triangle ABC is 18 cm^2.
Calculate the area of the quadrilateral $BCED$.

..... cm^2 **[3]**

877 In the diagram, DE is parallel to BC.
$AD = 6$ cm, $DB = 4$ cm and $DE = 9$ cm.



NOT TO SCALE

(a) Explain why triangle ADE is similar to triangle ABC.

.....

.....

..... **[2]**


(b) Calculate the length of BC.

$BC =$ ..... cm **[3]**

(c) The area of triangle ADE is 27 cm^2.
Calculate the area of the trapezium $DBCE$.

..... cm^2 **[3]**

878 The diagram shows two mathematically similar solid statues, P and Q.
Statue P has a height of 15 cm and a mass of 500 g.
Statue Q has a height of 45 cm.
Assume that both statues are made from the same uniform material.



(a) Calculate the mass of statue Q.
Give your answer in kilograms.

..... kg **[3]**


(b) A third similar statue, R, has a mass of 4 kg.
Calculate the height of statue R.

..... cm **[3]**

(c) The surface area of statue Q is 2430 cm^2.
Calculate the surface area of statue P.

..... cm^2 **[2]**

879 The diagram shows two geometrically similar solid metal cones, P and Q.
The volume of cone P is 320 cm^3 and the volume of cone Q is 1080 cm^3.



(a) The height of cone P is 12 cm.
Calculate the height of cone Q.

..... cm **[3]**


(b) The surface area of cone Q is 450 cm^2.
Calculate the surface area of cone P.

..... cm^2 **[2]**

(c) Both cones are melted down to make a single new solid cone R, which is geometrically similar to P and Q.
Calculate the height of cone R.
Give your answer correct to 3 significant figures.

..... cm **[3]**

880 The diagram shows a trapezium $ABCD$ with AB parallel to DC.
The diagonals AC and BD intersect at point X.
$AB = 9$ cm, $DC = 15$ cm and $BX = 6$ cm.
The area of triangle ABX is 27 cm^2.



NOT TO SCALE

(a) Explain why triangle ABX is similar to triangle CDX.

[2]

(b) Calculate the length of BD.

..... cm **[3]**


(c) Calculate the area of triangle CDX.

..... cm^2 **[2]**

(d) Calculate the area of triangle BCX.

..... cm^2 **[2]**

881 In the diagram, the line XY is parallel to BC.
$AX = 4$ cm, $XB = x$ cm, $AY = 5$ cm and $YC = (x + 2)$ cm.



(a) Calculate the value of x.

$x =$ ..... **[3]**


(b) $XY = 3.5$ cm.
Calculate the length of BC.

..... cm **[2]**

(c) The area of triangle AXY is A cm^2.
Find an expression for the area of the trapezium $XBCY$ in terms of A.
Give your answer in its simplest form.

..... **[2]**

882 Cone A and Cone B are mathematically similar solids.



NOT TO SCALE


(a) The volume of Cone A is 108 cm^3 and the volume of Cone B is 500 cm^3.
The sum of the surface areas of Cone A and Cone B is 340 cm^2.
Calculate the surface area of Cone B.

..... cm^2 **[4]**

(b) Cone B has a total vertical height of 20 cm.
A smaller cone, mathematically similar to Cone B, is removed from the top of Cone B to leave a frustum.
The volume of the frustum is $\frac{7}{8}$ of the volume of Cone B.
Calculate the vertical height of the removed small cone.

..... cm **[3]**

883 The diagram shows a circle with points A, B, C and D on its circumference.
The chords AC and BD intersect at X.



(a) Show that triangle AXB is similar to triangle DXC.
Give a geometric reason for each statement you make.

[3]


(b) $AX = 4$ cm, $BX = 5$ cm, $CX = 7.5$ cm and $AB = 6$ cm.
Calculate the length of DX and the length of DC.

$DX =$ ..... , $DC =$ ..... **[3]**

(c) The area of triangle AXB is 11.8 cm^2.
Calculate the area of triangle DXC.

..... cm^2 **[2]**

884 In the diagram, DE is parallel to BC.
$AD = (x + 2)$ cm, $DB = (x - 1)$ cm, $AE = 8$ cm and $EC = x$ cm.



NOT TO SCALE

(a) Show that $x^2 - 6x + 8 = 0$.

[3]


(b) Solve the equation $x^2 - 6x + 8 = 0$.

$x =$ ..... , $x =$ ..... **[2]**

(c) Given that the area of triangle ADE is 24 cm^2 and $x = 4$, calculate the area of the trapezium $DBCE$.

..... cm^2 **[3]**

885 In the diagram, D is a point on AB and E is a point on AC.
Angle AED = Angle ABC.
$AD = 6$ cm, $AE = 5$ cm and $DB = 9$ cm.



(a) Show that triangle ADE is similar to triangle ACB.
Give reasons for your statements.

[3]


(b) Calculate the length of EC.

$EC =$ ..... cm **[3]**

(c) Find the ratio of the area of triangle ADE to the area of quadrilateral $BCED$ in its simplest form.

..... **[2]**

886 The diagram shows a trapezium $ABCD$ with AB parallel to DC.
The diagonals AC and BD intersect at X.
$AB = 7$ cm and $DC = 10.5$ cm.



(a) Explain why triangle ABX is similar to triangle CDX.

[2]


(b) $AX = 4$ cm.
Calculate the length of AC.

$AC =$ ..... cm **[3]**

(c) The area of triangle ABX is 16 cm^2.
Calculate the area of the trapezium $ABCD$.

..... cm^2 **[4]**

887 The diagram shows triangle ABC.
D is a point on AC and E is a point on AB.
Angle ABC = Angle AED.



NOT TO SCALE

(a) Explain why triangle ABC is similar to triangle AED.

.....

.....

..... **[2]**

(b) Given that $AC = 15$ cm and $AD = 6$ cm, write down the scale factor of the enlargement that maps triangle AED onto triangle ABC.

..... **[1]**


(c) $AE = x$ cm and $AB = (2x + 2.5)$ cm.
Use similarity to set up an equation and find the value of x.

$x =$ ..... **[3]**

(d) The area of the quadrilateral $BCDE$ is 42 cm^2.
Calculate the area of triangle ADE.

..... cm^2 **[3]**

888 The diagram shows a trapezium $ABCD$ with AB parallel to DC.
The diagonals AC and BD intersect at X.



(a) Explain, giving geometric reasons, why triangle ABX is similar to triangle CDX.

[3]

(b) $AX = 4$ cm and $CX = 6$ cm.
$AB = x$ cm and $DC = (2x - 3)$ cm.

Work out the value of x.

$x =$ ..... **[3]**

(c) The area of triangle ABX is 24 cm^2.
Calculate the area of the trapezium $ABCD$.

Area = ... cm^2 **[4]**

889 Two solid statues, A and B, are mathematically similar.

(a) Statue A has a height of 12 cm and a volume of 800 cm^3.
Statue B has a volume of 2700 cm^3.
Calculate the height of Statue B.

..... cm **[3]**

(b) The cost to paint a statue is directly proportional to its surface area.
It costs \$18 to paint Statue A.
Calculate the cost to paint Statue B.

\$ ..... **[3]**

(c) A third mathematically similar statue, C, has a surface area that is 44% greater than the surface area of Statue A.
Calculate the volume of Statue C.

..... cm^3 **[4]**

890 Solid Cone A has a base radius of 10 cm and a perpendicular height of 24 cm.

(a) Calculate the total surface area of Cone A.
Give your answer in terms of π.

..... cm^2 **[3]**

(b) Cone B is mathematically similar to Cone A.
The volume of Cone B is 100π cm^3.
Calculate the total surface area of Cone B.
Give your answer in terms of π.

..... cm^2 **[4]**

(c) A frustum is made by cutting a smaller cone, which is geometrically identical to Cone B, from the top of Cone A.
This frustum is melted down and recast completely into a solid sphere.
Calculate the radius of this sphere.

..... cm **[4]**

891 Answer the following questions about the symmetry of two-dimensional shapes.

(a) Write down the number of lines of symmetry of a regular nonagon (a polygon with 9 sides).

..... **[1]**

(b) Write down the order of rotational symmetry of a parallelogram.

..... **[1]**

(c) Write down the mathematical name of a triangle that has exactly 1 line of symmetry and rotational symmetry of order 1.

..... **[1]**

892 Answer the following questions about 2D symmetry.

(a) Write down the order of rotational symmetry of a regular decagon.

..... **[1]**


(b) A quadrilateral has exactly two lines of symmetry and rotational symmetry of order 2.
State the mathematical name of this quadrilateral.

..... **[1]**

(c) A shape has rotational symmetry of order 5.
Calculate the smallest angle of rotation that maps the shape onto itself.

..... $^\circ$ **[1]**

893 The diagram shows a shaded shape on a coordinate grid.




(a) Write down the order of rotational symmetry of the shaded shape.

..... **[1]**

(b) Write down the number of lines of symmetry of the shaded shape.

..... **[1]**

(c) On the grid below, draw the image of the shape after a reflection in the line $x = 2$.



[2]

894 Write down the symmetry properties for the following geometric shapes.

(a) A regular polygon has 15 sides.

Write down:

(i) the number of lines of symmetry,

..... **[1]**

(ii) the order of rotational symmetry.

..... **[1]**

(b) A quadrilateral has exactly two lines of symmetry and rotational symmetry of order 2.


Write down the mathematical name of this quadrilateral.

..... **[1]**


(c) Write down the number of planes of symmetry of a solid right pyramid with a regular pentagonal base.

..... **[1]**


895 Three 2D shapes are shown below.



Parallelogram



Regular Pentagon




Isosceles Trapezium

Complete the table to show the number of lines of symmetry and the order of rotational symmetry for each shape.

[6]

896 The diagram shows a solid regular hexagonal prism.



(a) Write down the number of planes of symmetry of this regular hexagonal prism.

..... **[2]**

(b) Write down the order of rotational symmetry of the prism about the vertical axis passing through the centres of its hexagonal faces.

..... **[1]**

(c) A different solid is a right square-based pyramid.

Write down the number of planes of symmetry of this pyramid.

..... **[1]**

897 Consider the symmetry properties of the following 3D solid shapes.

(a) A right square-based pyramid has a base of side length 8 cm and perpendicular height 10 cm.
Write down the number of planes of symmetry for this pyramid.

..... **[1]**

(b) Write down the total number of planes of symmetry of a solid regular pentagonal prism.

..... **[2]**

(c) A right circular cone has base radius r and slant height l.
Describe fully its axis of symmetry.

..... **[2]**

898 The points $A(1, 6)$ and $B(7, 4)$ are two vertices of a quadrilateral $ABCD$.
The point $M(3, 2)$ is the centre of rotational symmetry of $ABCD$.
The quadrilateral $ABCD$ has rotational symmetry of order 4.

(a) Find the coordinates of C.

(..... , ...) **[2]**

(b) The line AC is a line of symmetry of the quadrilateral.


Find the equation of the line AC.
Give your answer in the form $y = mx + c$.

$y =$ ..... **[3]**

(c) Calculate the exact area of the quadrilateral $ABCD$.

..... **[3]**

899 The diagram shows a solid right prism.
The cross-section of the prism is an equilateral triangle.



(a) State the number of planes of symmetry of this solid prism.

..... **[2]**

(b) State the order of rotational symmetry of the prism about the axis passing through the centroids of the two equilateral triangular faces.

..... **[1]**

(c) A different solid is a right regular hexagonal prism.

State the number of planes of symmetry of this hexagonal prism.

..... **[2]**

900 **(a)** A regular polygon has an exterior angle of 15°.

(i) Calculate the number of sides of this polygon.

..... **[2]**

(ii) Write down the number of lines of symmetry of this polygon.

..... **[1]**

(b) Write down the mathematical name of a quadrilateral that has exactly 1 line of symmetry and rotational symmetry of order 1.


..... **[1]**

(c) A solid cone has a base radius of 5 cm and a perpendicular height of 12 cm.


State the number of planes of symmetry of this cone.

..... **[1]**

901 The diagram shows a solid square-based pyramid and a solid cylinder.



Square-based pyramid



Cylinder

(a) State the number of planes of symmetry for each solid.

(i) The solid square-based pyramid.

..... **[1]**

(ii) The solid cylinder.

..... **[1]**

(b) A solid square-based pyramid has a base side length of 6 cm.
The pyramid has exactly 4 planes of symmetry and its volume is 48 cm^3.
Calculate the length of a sloping edge of this pyramid.

..... cm **[5]**

902 A rhombus $ABCD$ has its centre at the origin $(0, 0)$.

(a) The line of symmetry passing through vertices A and C has the equation $y = 3x$.
Find the equation of the line of symmetry passing through vertices B and D.

$y =$ ..... **[2]**

(b) Write down the order of rotational symmetry of the rhombus $ABCD$.

..... **[1]**

(c) A different shape is a regular polygon with n sides.
The number of lines of symmetry of this polygon is equal to n.
The interior angle of this polygon is $165°$.

Calculate the value of n.

$n =$ ..... **[3]**

903 **(a)** A solid wooden block is in the shape of a right prism.
The cross-section of the prism is a regular octagon.

(i) Write down the order of rotational symmetry of the cross-section.

..... **[1]**

(ii) State the total number of planes of symmetry of this right prism.

..... **[2]**

(b) A different solid is a right pyramid with a regular polygon of n sides as its base.
The pyramid has exactly 12 planes of symmetry.


(i) Write down the value of n.

$n =$ ..... **[1]**

(ii) Calculate the sum of the interior angles of the base of this pyramid.


..... ° **[2]**

(c) A shape S is drawn on a coordinate grid.
The shape S has a line of symmetry with equation $y = x$.
One vertex of the shape is at the point $P(3, -2)$.
Write down the coordinates of the image of P after a reflection in this line of symmetry.



(..... , ...) **[2]**

904 Triangle ABC has vertices at $A(1, 2)$, $B(5, 4)$ and $C(3, 8)$.



(a) Show that triangle ABC is an isosceles triangle.

[3]

(b) Triangle ABC has exactly one line of symmetry.
Find the equation of this line of symmetry.
Give your answer in the form $y = mx + c$.

$y =$ ..... **[4]**

(c) A different shape is a regular polygon with n sides.
It has exactly 15 lines of symmetry.
Calculate the size of one interior angle of this polygon.

..... ° **[2]**

905 A right solid prism has a uniform cross-section which is a regular polygon with n sides.
The prism has exactly 7 planes of symmetry.
The length of the prism is not equal to its width.

(a) Explain why $n = 6$.

[2]

(b) The longest diagonal of the hexagonal cross-section is 12 cm.
The volume of the prism is 450 cm^3.


(i) Calculate the length of the prism.

..... cm **[4]**

(ii) Calculate the total surface area of the prism.

..... cm^2 **[4]**

906 In the diagram, the line AB is parallel to the line CD.
P is a point on AB.
Q and R are points on CD.
The line PQ and the line PR are drawn.
Angle $APQ = 48°$ and angle $BPR = 56°$.



NOT TO SCALE

(a) Write down the value of angle PQR.

Angle $PQR =$ ..... ° **[1]**


(b) Write down the value of angle PRQ.

Angle $PRQ =$ ..... ° **[1]**

(c) Work out the value of angle QPR.

Angle $QPR =$ ..... ° **[1]**

907 The diagram shows a straight line AB intersecting two parallel lines, CD and EF.



(a) Find the value of x.

$x =$ ..... **[1]**


(b) Give a geometric reason for your answer to part (a).

..... **[1]**

(c) Calculate the interior angle of a regular decagon (10-sided polygon).

..... ° **[1]**

908 In the diagram, the line AB is parallel to the line CD.
The point E lies between the two parallel lines.



NOT TO SCALE


(a) Calculate the size of angle AEC.

Angle AEC = ... ° **[2]**

(b) A regular polygon has an exterior angle of 24°.
Work out the number of sides of this polygon.

..... **[2]**

909 In the diagram, the line PQ is parallel to the line RS.
The point T lies on PQ, and the points U and V lie on RS.
The line RS is extended to a point W.
Angle $PTU = 52°$ and angle $QTV = 61°$.



(a) Calculate angle TUV.

Angle TUV = ... ° **[1]**


(b) Calculate angle UTV.

Angle UTV = ... ° **[2]**

(c) Calculate angle TVW.

Angle TVW = ... ° **[2]**

910 The diagram shows part of a regular polygon with n sides.
The exterior angle of this polygon is 24°.



(a) Calculate the value of n.

$n =$ ..... **[2]**


(b) Calculate the sum of the interior angles of this polygon.

..... ° **[2]**

(c) A different regular polygon has an interior angle of 165°.
Work out the number of sides of this polygon.

..... **[2]**

911 The diagram shows a triangle ABC with a line DE drawn parallel to BC.
D lies on AB and E lies on AC.
Angle $ADE = 55°$ and angle $AED = 70°$.



(a) Calculate angle DAE.

Angle $DAE =$ ..... ° **[2]**

(b) Calculate angle ABC.

Angle $ABC =$ ..... ° **[1]**

(c) Explain why triangle ADE is an isosceles triangle.

[2]

912 Answer the following questions about polygons and angles.

(a) An irregular polygon has n sides.
Three of the interior angles are each $140°$.
The remaining $(n-3)$ interior angles are each $165°$.

Work out the value of n.

$n =$ ..... **[4]**

(b) In triangle ABC, the angles are $\angle A = a°$, $\angle B = b°$ and $\angle C = c°$.
It is given that $c - b = a$.

Prove that ABC is a right-angled triangle.

[2]

(c) Polygon P is a regular polygon with 12 sides.
Polygon Q is a regular polygon with k sides.
The exterior angle of Polygon Q is half the size of the exterior angle of Polygon P.
Polygon P and Polygon Q share a common edge AB.


(i) Find the value of k.

$k =$ ..... **[2]**

(ii) Work out the size of the angle at vertex B that lies outside both polygons.

..... ° **[2]**

913 The diagram shows an irregular pentagon.
The interior angles of the pentagon are $x°$, $(x + 20)°$, $(2x - 10)°$, $(2x + 30)°$ and $140°$.



NOT TO SCALE


(a) Calculate the value of x.

$x =$ ..... **[3]**

(b) Find the size of the largest interior angle of the pentagon.

..... ° **[2]**

914 The diagram shows an irregular pentagon.



NOT TO SCALE


(a) Work out the value of x.

$x =$ ..... **[3]**

(b) Another regular polygon has an interior angle that is 4 times the size of its exterior angle.
Calculate the number of sides of this polygon.

..... **[3]**

915 In the diagram, the line AB is parallel to the line CD.
The transversal EF intersects AB at G and CD at H.
The point K lies on CD.



(a) Angle $EGB = (5x + 10)°$ and angle $GHD = (7x - 30)°$.
Find the value of x.
Give a geometrical reason for your working.

$x =$ ..... , Reason: ..... **[3]**

(b) Angle $HGK = 35°$.
Calculate angle GKH.

..... ° **[3]**

916 **(a)** This question is about the interior and exterior angles of polygons.

(i) The interior angles of a hexagon are $x°$, $2x°$, $(x + 30)°$, $(2x + 10)°$, $(3x - 20)°$, and $160°$.
Work out the value of x.

$x =$ ..... **[3]**

(ii) A regular polygon has an interior angle that is 6.5 times its exterior angle.
Calculate the number of sides of this polygon.


..... **[3]**

(iii) Two regular polygons, P and Q, have m sides and n sides respectively.
The sum of their exterior angles is $54°$.
The difference between their interior angles is $18°$.
Find the value of m and the value of n.

$m =$ ..... , $n =$ ..... **[5]**

917 **(a)** In the diagram, the line AB is parallel to the line CD.
X is a point between AB and CD.
The point P is on AB and the point Q is on CD.
Angle $BPX = (3x + 15)°$, angle $DQX = (2x + 10)°$ and angle $PXQ = 75°$.

Find the value of x.



NOT TO SCALE


$x =$ ..... **[3]**

(b) Polygon P is a regular polygon with n sides.
Polygon Q is a regular polygon with $3n$ sides.
The size of an interior angle of polygon Q is 16° greater than the size of an interior angle of polygon P.

Calculate the value of n.

$n =$ ..... **[4]**

918 The diagram shows a regular pentagon $ABCDE$ and a regular hexagon $EDFGHI$ sharing a common edge ED.



(a) Calculate angle CDE.

Angle CDE = ... ° **[2]**


(b) Calculate angle AEI.

Angle AEI = ... ° **[2]**

(c) Calculate angle EAI.

Angle EAI = ... ° **[3]**

919 The diagram shows three regular polygons meeting at the point P.
The polygons are a square, a regular pentagon, and a regular polygon with n sides.
The square shares the side PC with the pentagon and the side PA with the n-gon.
The pentagon shares the side PF with the n-gon.
They meet with no gaps and do not overlap.



(a) Calculate the interior angle of a regular pentagon.

..... ° **[2]**


(b) Find the value of n.

$n =$ ..... **[3]**

(c) The vertices A and F of the n-gon are joined by a straight line.
Calculate angle PFA.

..... ° **[2]**

920 The diagram shows two parallel lines, L_1 and L_2.
The point A lies on L_1 and the point C lies on L_2.
The point B lies between the two parallel lines.
The acute angle between L_1 and AB is $(3x + 10)°$.
The acute angle between L_2 and CB is $(2x + 20)°$.
The reflex angle ABC is $(300 - 2x)°$.




(a) Show that $x = 10$.

[4]

(b) The points A, B, and C form a triangle ABC.
Given that $AB = BC$, calculate the angle BCA.

Angle $BCA =$ ..... ° **[3]**

921 Points A, B and C lie on a circle, centre O.
Angle $AOB = 130°$.




(a) Find the size of angle ACB.

..... ° **[1]**

(b) Find the size of angle OAB.

..... ° **[2]**

922 A, B and C are points on a circle, center O.
Angle $ABC = 58°$.




(a) Work out the obtuse angle AOC.

Angle $AOC =$ ..... ° **[1]**

(b) Work out angle OAC.

Angle $OAC =$ ..... ° **[2]**

923 The diagram shows a circle, centre O.
A, B and P are points on the circumference.
AB is a diameter of the circle.
The line TC is a tangent to the circle at the point P.
Angle $BPC = 34°$.



NOT TO SCALE


(a) Write down the value of angle PAB.

Angle $PAB =$ ..... ° **[1]**

(b) Work out the value of angle POB.

Angle $POB =$ ..... ° **[2]**

924 A, B, C and D are points on a circle, centre O.
Angle $AOB = 110°$.



NOT TO SCALE

(a) Find the size of angle ACB.

Angle $ACB =$ ..... ° **[1]**


(b) Find the size of angle ADB.

Angle $ADB =$ ..... ° **[1]**

(c) State the geometrical reason for your answer to part (b).

..... **[1]**

925 A, B, C and D are points on a circle, center O.
Angle $ABC = 56°$.



(a) Calculate the obtuse angle AOC.

..... ° **[1]**


(b) Calculate angle ADC.

..... ° **[1]**

(c) State the geometric reason for your answer to part (b).

..... **[1]**

926 A, B, C and D are points on a circle, centre O.
Angle $ADB = 48°$.



(a) **(i)** Find angle ACB.

Angle $ACB =$ ..... ° **[1]**


(ii) Give a geometrical reason for your answer to part (a)(i).

..... **[1]**

(b) Find angle AOB.

Angle $AOB =$ ..... ° **[1]**

927 Points A, B, C and D lie on the circumference of a circle, center O.
Angle $AOB = 130°$.




(a) Calculate angle ACB.

..... ° **[1]**

(b) Calculate angle ADB.

..... ° **[2]**

928 A, B, C and D are points on the circumference of a circle, centre O.
Angle $BDC = 42°$ and angle $CBD = 65°$.



(a) Find angle BAC.

Angle $BAC =$ ..... ° **[1]**


(b) Find angle BCD.

Angle $BCD =$ ..... ° **[1]**

(c) Find angle BAD.

Angle $BAD =$ ..... ° **[2]**

929 The points A, B, and C lie on a circle, centre O.
The line SAT is a tangent to the circle at A.
Angle $SAB = 58°$.



(a) Find angle ACB and state a geometrical reason for your answer.

(i) Angle ACB

..... ° **[1]**


(ii) Reason

..... **[1]**

(b) Find angle OBA.

..... ° **[2]**

930 The diagram shows a circle with centre O.
The points A, B, C and D lie on the circumference of the circle.
Angle $AOB = 110°$ and angle $BDC = 30°$.



(a) Calculate angle ACB.

..... ° **[1]**

(b) State the geometrical reason for your answer to part (a).

..... **[1]**


(c) Calculate angle BAC.

..... ° **[1]**

(d) Calculate angle OAB.

..... ° **[2]**

931 A, B, C and D are points on the circumference of a circle, centre O.
Angle $AOC = 124°$.



(a) Calculate angle ABC.

..... ° **[2]**


(b) Calculate angle ADC.

..... ° **[2]**

(c) Calculate angle OAC.

..... ° **[2]**

932 A, B and C are points on a circle, centre O.
TA is a tangent to the circle at A.
C is a point on the major arc AB.
Angle $TAB = 42°$ and angle $BOC = 110°$.



(a) Write down the value of

(i) angle BCA,

Angle $BCA =$ ..... ° **[1]**

(ii) angle BAC.

Angle $BAC =$ ..... ° **[1]**


(b) Calculate angle OAB.

Angle $OAB =$ ..... ° **[2]**

(c) Calculate angle ABC.

Angle $ABC =$ ..... ° **[2]**

933 P, Q, R and S are points on a circle.
TQU is a tangent to the circle at Q.
Angle $RQU = 55°$ and Angle $QPS = 100°$.



NOT TO SCALE

(a) Find the size of angle QSR.

Angle $QSR =$ ..... ° **[1]**


(b) Find the size of angle QRS.

Angle $QRS =$ ..... ° **[2]**

(c) Calculate the size of angle SQR.

Angle $SQR =$ ..... ° **[2]**

934 Points P, Q, R and S lie on a circle.
The straight line XPT is a tangent to the circle at P.
Angle $QPT = 42°$, angle $PQR = 105°$ and angle $SPR = 48°$.



(a) Find the size of angle PRQ.

..... ° **[1]**

(b) Find the size of angle PSR.

..... ° **[1]**


(c) Find the size of angle PRS.

..... ° **[2]**

(d) Find the size of angle XPS.

..... ° **[1]**

935 In the diagram, AB is a diameter of the circle, center O.
The line XAY is a tangent to the circle at A.
C and D are points on the circle.
Angle $XAC = 54°$.



(a) **(i)** Write down the size of angle ACB.

..... ° **[1]**

(ii) Give a geometric reason for your answer.

..... **[1]**


(b) Find angle ABC.

..... ° **[1]**

(c) Find angle ADC.

..... ° **[2]**

936 P, Q and R are points on a circle, centre O.
ST is a tangent to the circle at R.
Angle $PQR = 118°$.



(a) Find reflex angle POR.

Reflex angle $POR =$ ..... ° **[2]**


(b) Find obtuse angle POR.

Obtuse angle $POR =$ ..... ° **[1]**

(c) Find angle PRS.

Angle $PRS =$ ..... ° **[2]**

937 A, B, C and D are points on a circle, centre O.
TA and TB are tangents to the circle at A and B.
Angle $ATB = 44°$.



(a) Calculate angle AOB.

Angle AOB = ... ° **[2]**


(b) Calculate angle ACB.

Angle ACB = ... ° **[1]**

(c) Calculate angle ADB.

Angle ADB = ... ° **[2]**

938 A, B, C and D are points on a circle.
AB is a diameter of the circle.
$CD = CB$ and angle $CAB = 28°$.



(a) (i) Find angle CBA.

Angle $CBA =$ ..... ° **[1]**

(ii) Give a geometrical reason for your answer.

..... **[1]**


(b) Find angle CAD.

Angle $CAD =$ ..... ° **[1]**

(c) Find angle ADC.

Angle $ADC =$ ..... ° **[2]**

939 A, B, C and D lie on a circle, centre O.
AB is a diameter of the circle.
The line XY is a tangent to the circle at B.
Angle $CBY = 58°$.



(a) (i) Find angle BAC.

..... ° **[1]**

(ii) State the circle theorem that gives this angle directly from angle CBY.

..... **[1]**


(b) Calculate angle ABC.

..... ° **[3]**

(c) Calculate angle ADC.

..... ° **[2]**

940 A, B, C and D are points on a circle, center O.
AD is a diameter of the circle.
PAT is a tangent to the circle at A.
Angle $PAB = 58°$ and angle $BDC = 22°$.



(a) Calculate angle ADB.

Angle $ADB =$ ..... ° **[2]**


(b) Calculate angle ABC.

Angle $ABC =$ ..... ° **[2]**

(c) Calculate angle OBC.

Angle $OBC =$ ..... ° **[3]**

941 A, B, C and D lie on a circle, centre O.
Angle $ABC = (x + 32)°$ and the obtuse angle $AOC = (4x - 20)°$.
The line AD is extended to the point E.




(a) Find the value of x.

$x =$ ..... **[3]**

(b) Work out the size of angle CDE.

Angle $CDE =$ ..... ° **[2]**

942 A, B and C are points on a circle, center O.
The lines TA and TC are tangents to the circle at A and C respectively.
Angle $ATC = 48°$.



(a) Calculate angle AOC.

..... ° **[2]**


(b) Calculate angle ABC.

..... ° **[1]**

(c) Calculate angle OAC.

..... ° **[2]**

943 P, Q, R and S are points on a circle, center O.
PR is a diameter of the circle.
TPV is a tangent to the circle at P.
Angle $QPR = 32°$ and angle $RPS = 41°$.



(a) Calculate angle PRQ.

Angle $PRQ =$ ..... ° **[2]**


(b) Calculate angle TPQ.

Angle $TPQ =$ ..... ° **[2]**

(c) Calculate angle PQS.

Angle $PQS =$ ..... ° **[2]**

944 AD is a diameter of the circle, centre O.
The points A, B, C, and D lie on the circumference.
The chords AC and BD intersect at X.
Angle $BAC = 32°$ and angle $CAD = 24°$.



(a) Find angle ADC.

..... ° **[2]**

(b) Find angle ABC.

..... ° **[1]**


(c) Find angle DBC.

..... ° **[1]**

(d) Find angle AXB.

..... ° **[2]**

945 The diagram shows a circle with centre O.
The line PAQ is a tangent to the circle at the point A.
The points B, C and D lie on the circle.
Angle $QAB = 62°$.



(a) Calculate angle ACB and give a geometrical reason for your answer.

$Angle ACB =$ ..... ° , Reason: ..... **[2]**


(b) Calculate angle ADB.

..... ° **[2]**

(c) Calculate angle OBA.

..... ° **[2]**

946 The points A, B, C and D lie on a circle.
XY is a tangent to the circle at A.
Angle $BCD = 115°$, angle $XAB = (2x + 15)°$ and angle $YAD = (x + 10)°$.




(a) Find the value of x.

$x =$ ..... **[4]**

(b) Calculate angle ABD.

Angle $ABD =$ ..... ° **[2]**

947 A, B, C and Q are points on a circle, center O.
The line PQR is a tangent to the circle at Q.
The chord AB is parallel to PR.
Angle $AQR = 58°$.



(a) (i) Find the size of angle ABQ.

..... ° **[1]**

(ii) Give a geometric reason for your answer.

..... **[1]**


(b) Find angle AOB.

..... ° **[2]**

(c) Find angle ACB.

..... ° **[2]**

948 A, B, C and D are points on a circle, centre O.
AB is a diameter of the circle.
PQ is a tangent to the circle at C.
Angle $CAB = (2x - 5)°$ and Angle $BCQ = (x + 17)°$.



NOT TO SCALE

(a) Set up an equation in terms of x and solve it to show that $x = 22$.
Give a geometrical reason to justify your equation.

[3]


(b) Work out the size of angle CBA.

Angle $CBA =$ ..... ° **[2]**

(c) Find the size of angle ADC.

Angle $ADC =$ ..... ° **[2]**

949 Points A, B, C and D lie on a circle.
The straight line PAQ is a tangent to the circle at A.
Angle $PAB = (2x + 10)°$, angle $ABC = (3x - 5)°$ and angle $CAQ = (x + 25)°$.
D is a point on the major arc BC.



(a) Find the value of x.
Show all your working.

$x =$ ..... **[3]**


(b) Find the size of angle BAC.

..... ° **[2]**

(c) Find the size of angle BDC.

..... ° **[2]**

950 TA and TB are tangents to a circle, centre O, at the points A and B.
C is a point on the major arc AB and D is a point on the minor arc AB.
Angle $ATB = (x + 10)°$
Angle $TAB = (2y - 10)°$
Angle $ACB = (3y - 40)°$



(a) Find the value of y.

$y =$ ..... **[3]**


(b) Find the value of x.

$x =$ ..... **[2]**

(c) Find angle ADB.

Angle $ADB =$ ..... ° **[2]**

951 A, B, C and D are points on a circle, centre O.
TA and TC are tangents to the circle at A and C.
Angle $ATC = 48°$ and angle $CAD = 31°$.



(a) **(i)** Find angle ABC.

Angle $ABC =$ ..... ° **[2]**

(ii) Find angle ADC.

Angle $ADC =$ ..... ° **[2]**


(iii) Find angle ACD.

Angle $ACD =$ ..... ° **[1]**

(iv) Calculate angle AOC.

Angle $AOC =$ ..... ° **[2]**

952 The points A, B, C, and D lie on a circle, centre O.
PA and PB are tangents to the circle at A and B.
Angle $APB = 2x°$.
C is a point on the major arc AB and D is a point on the minor arc AB.



(a) Find an expression, in terms of x, for each of the following angles.
Give each answer in its simplest form.

Angle AOB

..... ° **[1]**

(b) Angle ACB

..... ° **[1]**


(c) Angle ADB

..... ° **[2]**

(d) Given that angle $ADB = 3 \times$ angle ACB, find the value of x.

$x =$ ..... **[3]**

953 The diagram shows a circle containing the points A, B, C and D.
The straight line PAQ is a tangent to the circle at A.
The chords AC and BD intersect.
Angle $PAD = (2x + 3y)^\circ$ and angle $ABD = (x + 2y + 10)^\circ$.
Angle $BAC = (3x - y)^\circ$ and angle $BDC = (x + y + 8)^\circ$.



NOT TO SCALE

(a) **(i)** Use the circle theorems to write down two equations in x and y.

[2]


(ii) Solve your equations to find the value of x and the value of y.
Show all your working.

$x =$ ..... , $y =$ ..... **[3]**

(b) Given that angle $ABC = 95^\circ$, work out the size of angle QAB.

Angle $QAB =$ ..... ° **[3]**

954 A, B and C are points on a circle, center O.
The straight line PTQ is a tangent to the circle at the point T.
$\angle ATP = (4x - 5)^\circ$ and $\angle ABT = (2x + 25)^\circ$
$\angle BTQ = (3y)^\circ$ and $\angle BAT = (y + 12)^\circ$



(a) Use the alternate segment theorem to find the values of x and y.

$x =$ ..... , $y =$ ..... **[4]**


(b) Calculate angle ATB.

..... ° **[2]**

(c) C is a point on the minor arc AT.
Calculate angle ACT.

..... ° **[2]**

955 The diagram shows a circle with centre O.
The lines TA and TB are tangents to the circle at the points A and B.
C is a point on the major arc AB.
Angle $ATB = x°$ and angle $ACB = (2x - 30)°$.



(a) (i) Find the value of x.

$x =$ ..... **[4]**


(ii) Calculate angle OAB.

..... ° **[2]**

(b) D is a point on the minor arc AB.
Calculate angle ADB.

..... ° **[2]**

956 PA and PB are tangents to a circle, centre O, from a point P.
C is a point on the major arc AB.
Angle $APB = 4x°$.
Angle $ACB = (3x - 10)°$.



(a) Find an expression for angle PAB in terms of x.

..... ° **[3]**

(b) Use the alternate segment theorem to form an equation in x, and show that $x = 20$.

[3]


(c) Calculate angle AOB.

..... ° **[3]**

(d) D is a point on the minor arc AB.
Calculate angle ADB.

..... ° **[3]**

957 The diagram shows a circle with centre O.
P is an external point.
The lines PA and PB are tangents to the circle at the points A and B.
Angle $APB = 44°$.



NOT TO SCALE


(a) Calculate angle PAB.

Angle PAB = ... ° **[2]**

(b) Calculate angle AOB.

Angle AOB = ... ° **[2]**

958 P is a point outside a circle, centre O.
PA and PB are tangents to the circle at A and B.
Angle $APB = 40°$.



(a) Write down the geometrical reason why triangle PAB is isosceles.

..... **[1]**


(b) Calculate angle PAB.


Angle $PAB =$ ..... ° **[1]**

(c) Calculate angle AOB.

Angle $AOB =$ ..... ° **[2]**

959 P and Q are points on a circle, centre O.
TP and TQ are tangents to the circle from an external point T.
The line TO intersects the circle at S.
Angle $PTQ = 56°$.





(a) Calculate angle POT.

Angle $POT =$ ..... ° **[2]**


(b) Explain why triangle TPQ is isosceles.

..... **[1]**

(c) Calculate angle TPQ.

Angle $TPQ =$ ..... ° **[1]**

960 The diagram shows a circle, centre O.
TA and TB are tangents to the circle at A and B.
Angle $ATB = 44°$.



(a) Work out angle AOB.

Angle $AOB =$ ..... ° **[2]**


(b) Work out angle OAB.

Angle $OAB =$ ..... ° **[2]**

(c) Write down the geometrical reason why the length of TA is equal to the length of TB.

..... **[1]**

961 P is an external point.
PA and PB are tangents to a circle, centre O.
The line OP intersects the chord AB at right angles.
$OA = 7$ cm and $PA = 24$ cm.



(a) Write down the length of PB.

..... cm **[1]**


(b) Calculate the length of OP.

..... cm **[2]**

(c) Calculate the shortest distance from O to the chord AB.

..... cm **[3]**

962 The diagram shows a circle, centre O, with radius 25 cm.
AB and CD are two chords of the circle such that $AB = CD$.
The perpendicular distance from O to AB is $(x + 3)$ cm.
The perpendicular distance from O to CD is $(3x - 5)$ cm.




(a) Find the value of x.

$x =$ ..... **[2]**

(b) Calculate the length of the chord AB.

$AB =$ ..... cm **[3]**

963 A circle, centre O, has a radius of 8.5 cm.
A chord AB has length 14 cm.
M is the midpoint of AB.




(a) Calculate the distance OM.

$OM =$ ..... cm **[3]**

(b) Another chord CD is drawn such that it is parallel to AB and is at a distance of 3 cm from O.
Calculate the length of the chord CD.

$CD =$ ..... cm **[2]**

964 The diagram shows a circle, centre O, with radius 13 cm.
AB is a chord of the circle of length 24 cm.
The point M is the midpoint of the chord AB.
Another chord CD is drawn inside the circle.



NOT TO SCALE

(a) Calculate the perpendicular distance, OM, from the centre of the circle to the chord AB.

$OM =$ ..... cm **[3]**

(b) The perpendicular distance from the centre O to the chord CD is also 5 cm.
Write down the length of the chord CD.

$CD =$ ..... cm **[1]**


(c) Calculate the area of triangle OAB.

Area =

.....

cm^2 **[2]**

965 A, B, C and D are points on a circle, centre O and radius 25 cm.
The chords AB and CD each have a length of 14 cm.
M is the midpoint of AB and N is the midpoint of CD.
Angle $MON = 100°$.



(a) Show that $OM = 24$ cm.

[2]


(b) Calculate the length of MN.

..... cm **[3]**

(c) Calculate the area of triangle MON.

..... cm^2 **[2]**

966 The diagram shows a circle, centre O, with radius 13 cm.
AB is a chord of the circle of length 24 cm.



(a) Calculate the shortest distance from O to the chord AB.

..... *cm* **[3]**

(b) CD is another chord of the circle.
CD is parallel to AB and has length 24 cm.
Write down the shortest distance from O to CD.

..... *cm* **[1]**


(c) EF is a third chord of the circle.
The shortest distance from O to EF is 12 cm.
Calculate the length of the chord EF.

..... *cm* **[3]**

(d) State the geometrical property of a circle used to find the length of half the chord in part (a).

..... **[1]**

967 Two circles, C_1 and C_2, have a common centre O.
The radius of C_1 is r cm and the radius of C_2 is R cm, where $R > r$.
A chord AB of the larger circle C_2 is tangent to the smaller circle C_1 at the point T.
The length of the chord AB is 24 cm and the area of the annular region between the two circles is $k\pi$ cm^2.



(a) Show that $R^2 - r^2 = 144$.

[3]


(b) Find the value of k.

$k =$ ..... **[2]**

(c) A second chord CD of the larger circle C_2 has length 24 cm.
State, with a reason, the distance from O to the chord CD.

..... **[2]**

968 A circle has centre O and radius 10 cm.
MN is a chord of length 16 cm.
The perpendicular bisector of MN intersects MN at X and the minor arc MN at Y.
PQ is a second chord which is parallel to MN.
The perpendicular distance between MN and PQ is 11 cm.



(a) Explain why the line containing XY passes through the centre O.

..... **[1]**


(b) Calculate the length of XY.

$XY =$ ..... cm **[3]**

(c) Calculate the length of the chord PQ.

$PQ =$ ..... cm **[4]**

969 TA and TB are tangents to a circle, centre O and radius r, at the points A and B.
The line TO intersects the chord AB at M.
Angle ATO = Angle $BTO = x$.



(a) Show that the length of the chord AB is $2r\cos x$.

[2]


(b) A second chord BC is drawn such that $BC = AB$.
Write down the shortest distance from O to the chord BC, in terms of r and x.

..... **[1]**

(c) The area of triangle OAB is equal to the area of triangle ATB.
Find the exact value of x.

$x =$ ..... $^\circ$ **[5]**

970 A circle is inscribed in a quadrilateral $ABCD$.
The circle touches the sides AB, BC, CD, and DA at the points P, Q, R, and S respectively.



(a) Write down the geometrical reason why $AP = AS$.

..... **[1]**

(b) Show that $AB + CD = AD + BC$.

[3]


(c) Given that $AB = (2x + 5)$ cm, $BC = (3x - 2)$ cm, $CD = (x + 8)$ cm and $AD = (2x + 1)$ cm, work out the value of x.

$x =$ ..... **[3]**

(d) Calculate the perimeter of the quadrilateral $ABCD$.

Perimeter = ... cm **[2]**

971 The diagram shows a circle with centre O and radius 7 cm.
The lines TP and TQ are tangents to the circle at the points P and Q.
The distance from the centre O to the external point T is 25 cm.



NOT TO SCALE

(a) Show that the length of the tangent TP is 24 cm.

[2]

(b) Calculate angle PTQ.

Angle PTQ = ... ° **[3]**

(c) Calculate the area of the shaded region bounded by the tangents TP, TQ and the minor arc PQ.

Area =

.....

cm^2 **[4]**

972 **(a)** Change 52000 cm^2 into m^2.

..... m^2 **[1]**


(b) Change 4.7 m^3 into litres.

..... litres **[1]**

(c) Change 850 millilitres into cm^3.

..... cm^3 **[1]**

973 A rectangular piece of card and a water jug are being measured.



(a) Calculate the area of the rectangular piece of card in square metres (m^2).

..... m^2 **[2]**

(b) The jug holds 3.2 litres of water.
Write down this volume in cm^3.

..... cm^3 **[1]**

974 **(a)** Change 3.5 m^2 into cm^2.

..... cm^2 **[1]**

(b) A rectangular tank measures 60 cm by 45 cm by 30 cm.
Calculate the volume of the tank in litres.

..... litres **[3]**

975 A solid wooden block is in the shape of a cuboid.
The dimensions of the block are 40 cm, 250 mm and 0.2 m.

(a) Work out the volume of the block in cm^3.

..... cm^3 **[2]**


(b) Convert your answer to part (a) into litres.

..... litres **[1]**

(c) The density of the wood is 0.6 g/cm^3.
Calculate the mass of the block in kilograms.

..... kg **[2]**

976 A rectangular block of metal has length 2.5 m, width 1.4 m and thickness 8 mm.



NOT TO SCALE


(a) Work out the volume of the block of metal in cm^3.

..... cm^3 **[2]**

(b) The mass of 1 cm^3 of this metal is 7.8 g.
Work out the mass of the block of metal in kilograms.

..... kg **[2]**

977 A metal water tank is made in the shape of a cuboid.



(a) Calculate the capacity of the water tank in litres.

..... litres **[3]**

(b) A solid metal sphere is melted down to make the tank.
The metal sphere has a volume of 0.024 m^3.
The mass of 1 cm^3 of this metal is 7.8 g.

Calculate the total mass of the sphere, giving your answer in kilograms.

..... kg **[3]**

978 During a storm, 15 mm of rain falls on a flat rectangular roof measuring 15 m by 12 m.

(a) Calculate the volume of rain that falls on the roof.
Give your answer in litres.

..... litres **[3]**

(b) The water from the roof flows into a cylindrical tank with a base radius of 60 cm.
Calculate the increase in the depth of the water in the tank.
Give your answer in centimetres, correct to 1 decimal place.

..... cm **[4]**

979 A solid metal sphere has a radius of 15 cm.
The volume, V, of a sphere with radius r is $V = \frac{4}{3}\pi r^3$.

(a) Calculate the volume of the sphere in m^3.
Give your answer in standard form.

..... m^3 **[3]**

(b) The mass of the sphere is 125 kg.
Calculate the density of the metal in g/cm^3.

..... g/cm^3 **[3]**

(c) The sphere is melted down to make completely solid identical cones.
Each cone has a base radius of 24 mm and a height of 5.5 cm.
The volume, V, of a cone with radius r and height h is $V = \frac{1}{3}\pi r^2 h$.


(i) Work out the volume of one cone in cm^3.

..... cm^3 **[3]**

(ii) Work out the maximum number of completely solid cones that can be made.

..... **[2]**

980 Water flows through a cylindrical pipe into an empty rectangular pool.
The pool measures 4 m by 2.5 m.



(a) The cylindrical pipe has a radius of 15 mm.
Water flows through the pipe at a speed of 0.8 m/s.


Show that the volume of water flowing out of the pipe in one second is 0.565 litres, correct to 3 significant figures.

[3]

(b) Calculate the time taken for the depth of water in the pool to reach 1.2 m.
Give your answer in hours and minutes, correct to the nearest minute.

..... hours ..... minutes **[4]**

981 Water flows through a cylindrical pipe of internal radius 2.5 cm at a speed of 5.4 km/h.
The water flows into an empty rectangular tank of length 3 m and width 2 m.



NOT TO SCALE

(a) Show that the volume of water flowing into the tank in 1 minute is 176.7 litres, correct to 1 decimal place.

[3]

(b) Calculate the depth of the water in the tank after 45 minutes.
Give your answer in centimetres.

..... cm **[4]**

982 Water flows through a cylindrical pipe of radius 2.5 cm at a speed of 18 km/h.



(a) Calculate the volume of water, in litres, that flows through the pipe in 1 minute.

..... litres **[5]**

(b) The water from the pipe is used to fill an empty rectangular reservoir measuring 50 m by 30 m by 4 m.
Calculate the time it takes to fill the reservoir completely.
Give your answer in days and hours, correct to the nearest hour.

days ..... , hours ..... **[7]**

983 Water flows through a cylindrical pipe of radius 2.5 cm at a speed of 1.6 m/s.

(a) Calculate the volume of water, in litres, that flows through the pipe in one minute.
Give your answer correct to 3 significant figures.

..... litres **[4]**


(b) This water flows into an empty rectangular tank measuring 4.2 m by 3.5 m by 2.4 m.
Calculate the time taken to fill the tank completely.
Give your answer in hours and minutes, correct to the nearest minute.

hours ..... , minutes ..... **[5]**

(c) The outside vertical walls of the rectangular tank are to be painted.
1 litre of paint covers 45000 cm^2.
Paint is sold in 2.5-litre tins.
Work out the minimum number of tins of paint needed.

..... tins **[6]**

984 An isosceles triangle and a rectangle are shown below.
The area of the triangle is equal to the area of the rectangle.



NOT TO SCALE


(a) Calculate the value of x.

$x =$ ..... **[2]**

(b) Calculate the perimeter of the isosceles triangle.

..... cm **[3]**

985 The diagram shows a parallelogram $ABCD$.
The area of the parallelogram is 120 cm^2 and its perimeter is 54 cm.



(a) Work out the value of h.

$h =$ ..... cm **[2]**


(b) Work out the length of AD.

$AD =$ ..... cm **[2]**

(c) Calculate the area of triangle ACD.

Area = ... cm^2 **[1]**

986 The diagram shows a rectangle.
All measurements are in centimetres.




(a) The perimeter of the rectangle is 34 cm.
Find the value of x.

$x =$ ..... **[3]**

(b) Calculate the area of the rectangle.

..... cm^2 **[2]**

987 The diagram shows a right-angled trapezium.




(a) Calculate the area of the trapezium.

..... cm^2 **[2]**

(b) Calculate the perimeter of the trapezium.

..... cm **[3]**

988 $ABCD$ is a rectangle with length 14 cm and width 9 cm.
The point E lies on the edge CD such that $CE = 5$ cm.



NOT TO SCALE


(a) Calculate the area of the shaded triangle ABE.

..... cm^2 **[2]**

(b) Calculate the perimeter of the shaded triangle ABE.

..... cm **[4]**

989 The diagram shows a trapezium $ABCD$ with AB parallel to DC.
The perpendicular height of the trapezium is 8 cm.
All measurements are in centimetres.




(a) Calculate the area of the trapezium.

..... cm^2 **[2]**

(b) Calculate the perimeter of the trapezium.

..... cm **[4]**

990 The diagram shows a rectangle and a triangle.
The area of the rectangle is equal to the area of the triangle.




(a) Form an equation in x and solve it to find the value of x.

$x =$ ..... **[4]**

(b) Work out the perimeter of the rectangle.

..... cm **[2]**

991 The diagram shows a trapezium.
All measurements are in centimetres.




(a) The area of the trapezium is 120 cm^2.
Work out the value of x.

$x =$ ..... **[3]**

(b) The trapezium is an isosceles trapezium, with the two slanted sides being equal in length.
Calculate the perimeter of the trapezium.

Perimeter = ... cm **[4]**

992 A trapezium $ABCD$ has right angles at A and B.
The side AD is parallel to BC.
$AD = (x + 4)$ cm, $BC = (3x - 2)$ cm and $AB = (x - 1)$ cm.



NOT TO SCALE

(a) The area of the trapezium is 20 cm^2.


(i) Show that $2x^2 - x - 21 = 0$.

[3]

(ii) Calculate the perimeter of the trapezium.

..... cm **[5]**

993 The diagram shows a right-angled trapezium with parallel sides of length x cm and $(x + 4)$ cm.
The height of the trapezium is $(x - 1)$ cm.
The area of the trapezium is 40 cm^2.



(a) Show that $x^2 + x - 42 = 0$.

[3]


(b) Solve the equation $x^2 + x - 42 = 0$ to find the value of x.

$x =$ ..... **[2]**

(c) Calculate the perimeter of the trapezium.

..... cm **[4]**

994 A rectangle and a right-angled triangle are shown.
All measurements are in centimetres.



(a) The area of the rectangle is equal to the area of the right-angled triangle.
Show that $2x^2 - 13x - 8 = 0$.

[4]

(b) Solve the equation $2x^2 - 13x - 8 = 0$.
Show all your working and give your answers correct to 2 decimal places.

$x =$ ..... , $x =$ ..... **[4]**

(c) Calculate the perimeter of the rectangle.

Perimeter = ... cm **[2]**

995 A piece of wire of length 60 cm is cut into two pieces.
One piece is bent to form a square.
The other piece is bent to form a rectangle where the length is exactly twice the width.
Let the width of the rectangle be y cm.

(a) Write down an expression, in terms of y, for the side length of the square.

..... cm **[2]**

(b) The total area enclosed by the square and the rectangle is 113 cm^2.
Show that $17y^2 - 180y + 448 = 0$.

[3]

(c) Solve the equation.

$$17y^2 - 180y + 448 = 0$$


Show all your working.

$y =$ ..... , $y =$ ..... **[3]**

(d) Given that the side length of the square must be greater than the width of the rectangle, calculate the area of the square.

..... cm^2 **[2]**

996 The diagram shows a sector of a circle, center O, with radius 12 cm and sector angle 45°.




(a) Work out the exact length of the arc of the sector.
Give your answer in terms of π.

..... cm **[2]**

(b) Work out the exact area of the sector.
Give your answer in terms of π.

..... cm^2 **[2]**

997 The diagram shows a minor sector AOB of a circle with centre O and radius 12 cm.
The sector angle is 60°.



NOT TO SCALE


(a) Work out the length of the minor arc AB.
Give your answer in terms of π.

..... cm **[2]**

(b) Work out the area of the minor sector AOB.
Give your answer in terms of π.

..... cm^2 **[2]**

998 The diagram shows a sector of a circle with centre O and radius 12 cm.
The sector angle is 45°.




(a) Work out the exact area of the sector.
Give your answer in terms of π.

Area = ... cm^2 **[2]**

(b) Work out the exact perimeter of the sector.
Give your answer in terms of π.

Perimeter = ... cm **[3]**

999 The diagram shows a sector of a circle with centre O and radius 6 cm.
The sector angle is 150°.



(a) Calculate the area of the sector.
Give your answer in terms of π.

..... cm^2 **[2]**

(b) Calculate the perimeter of the sector.
Give your answer in terms of π.

..... cm **[3]**

1000 **(a)** A sector of a circle, center O, has radius r cm and sector angle $\theta°$.
The area of the sector is 10π cm^2 and the length of its arc is 2π cm.

Find the value of r and the value of θ.


$r =$ ..... , $\theta =$ ..... **[5]**

(b) Another sector of a circle has radius x cm and sector angle $(x + 20)°$.
The length of its arc is $\frac{16\pi}{5}$ cm.

Find the value of x.
You must show all your working.

$x =$ ..... **[4]**

1001 The diagram shows a sector of a circle with centre O and angle 72°.
The area of the sector is 85 cm^2.



NOT TO SCALE


(a) Calculate the radius of the sector.

..... cm **[3]**

(b) Calculate the perimeter of the sector.

..... cm **[3]**

1002 The diagram shows a major sector of a circle with centre O and radius r cm.
The reflex angle is 250°.
The length of the major arc is 35 cm.




(a) Calculate the value of r.

$r =$ ..... **[3]**

(b) Calculate the area of the major sector.

..... cm^2 **[3]**

1003 The diagram shows a shaded shape made from two concentric sectors of circles with centre O.
The inner radius is 5 cm and the outer radius is 9 cm.
The angle of both sectors is 120°.



(a) Calculate the area of the shaded shape.

Area = ... cm^2 **[3]**


(b) Calculate the perimeter of the shaded shape.

Perimeter = ... cm **[3]**

(c) A complete circle has the same area as the shaded shape.
Calculate the radius of this circle.

Radius = ... cm **[2]**

1004 The diagram shows a sector OAB of a circle with center O and radius 8.5 cm.
The sector angle is 110°.
A straight line connects A and B, forming a triangle OAB and a shaded segment.



(a) Calculate the area of the sector OAB.

..... cm^2 **[2]**


(b) Calculate the area of the shaded segment.

..... cm^2 **[3]**

(c) Calculate the perimeter of the shaded segment.

..... cm **[4]**

1005 The diagram shows a circle with centre O and radius 8 cm.
A and B are points on the circumference such that angle $AOB = 110°$.



(a) Calculate the area of the shaded segment.

..... cm^2 **[4]**

(b) Calculate the perimeter of the shaded segment.

..... cm **[3]**

1006 A sector of a circle has radius r cm and sector angle $x°$.

(a) The arc length of the sector is 6π cm and the area of the sector is 27π cm^2.

(i) Work out the value of r.

$r =$ ..... **[3]**


(ii) Work out the value of x.

$x =$ ..... **[2]**

(b) Another sector has radius R cm and sector angle $y°$.
The perimeter of this sector is equal to the circumference of a circle with radius R cm.
Find an expression for y in terms of π.

$y =$ ..... **[4]**

1007 The diagram shows a circle, centre O, with radius 15 cm.
AB is a chord of the circle with length 24 cm.
The region bounded by the chord AB and the minor arc AB is shaded.



NOT TO SCALE

(a) Show that angle $AOB = 106.3°$, correct to 1 decimal place.

[3]

(b) For the following calculations, use angle $AOB = 106.3°$.


(i) Calculate the perimeter of the shaded minor segment.

..... cm **[3]**

(ii) Calculate the area of the shaded minor segment.

..... cm^2 **[4]**

1008 The diagram shows a solid sphere with center O and radius 3 cm.




(a) Calculate the exact volume of the sphere.
Give your answer in terms of π.

..... cm^3 **[2]**

(b) Calculate the exact surface area of the sphere.
Give your answer in terms of π.

..... cm^2 **[2]**

1009 A solid is made by joining a solid hemisphere of radius 3 cm to a solid cylinder of radius 3 cm and height 10 cm.



NOT TO SCALE


(a) Work out the volume of the solid.
Give your answer in terms of π.

..... cm^3 **[3]**

(b) Work out the total surface area of the solid.
Give your answer in terms of π.

..... cm^2 **[3]**

1010 The diagram shows a solid cone.
The base radius of the cone is 6 cm and the slant height is l cm.
The total surface area of the cone is 96π cm^2.




(a) Show that the slant height, l, is 10 cm.

[3]

(b) Work out the volume of the cone.
Give your answer in terms of π.

Volume = ... cm^3 **[4]**

1011 The diagram shows a solid formed by joining a cone to a hemisphere with the same radius.
The radius of the hemisphere is 3 cm and the slant height of the cone is 5 cm.




(a) Calculate the exact volume of the solid.
Give your answer in terms of π.

..... cm^3 **[4]**

(b) Calculate the total surface area of the solid.
Give your answer in terms of π.

..... cm^2 **[3]**

1012 The diagram shows a solid triangular prism.
The cross-section of the prism is a right-angled triangle with sides 6 cm, 8 cm and 10 cm.
The length of the prism is 15 cm.




(a) Calculate the volume of the prism.

..... cm^3 **[2]**

(b) Calculate the total surface area of the prism.

..... cm^2 **[3]**

1013 The diagram shows a solid cylinder and a solid sphere.




(a) The cylinder has radius $r = 5$ cm and height $h = 12$ cm.
Work out the total surface area of the cylinder.

Surface area = ... cm^2 **[3]**

(b) The sphere has the same volume as the cylinder.
Calculate the radius, R, of the sphere.

$R =$ ..... cm **[3]**

1014 The diagram shows a solid pyramid with a square base.
The base has sides of length 10 cm.
The volume of the pyramid is 400 cm^3.



NOT TO SCALE


(a) Calculate the perpendicular height of the pyramid.

..... cm **[2]**

(b) Calculate the total surface area of the pyramid.

..... cm^2 **[4]**

1015 A solid is formed by attaching a right circular cone to the top of a solid cylinder.
The cylinder and the base of the cone both have a radius of 5 cm.
The height of the cylinder is 12 cm and the total height of the solid is 24 cm.



(a) Calculate the volume of the solid.

..... cm^3 **[4]**

(b) Calculate the total surface area of the solid.

..... cm^2 **[5]**

1016 A solid metal cylinder has radius x cm and height $3x$ cm.
The cylinder is melted down and completely recast into solid spheres.
Each sphere has a radius of $\frac{x}{2}$ cm.

(a) Find an expression, in terms of x and π, for the volume of the cylinder.
Give your answer in its simplest form.

..... cm^3 **[2]**

(b) Find an expression, in terms of x and π, for the volume of one sphere.
Give your answer in its simplest form.

..... cm^3 **[2]**


(c) Work out the number of complete spheres that can be made.

..... **[1]**

(d) The total surface area of the original cylinder is 140 cm^2.
Work out the value of x.

$x =$ ..... **[3]**

1017 The diagram shows a water trough in the shape of a prism of length 120 cm.
The cross-section of the prism is a trapezium $ABCD$.
AB is parallel to DC.
$AB = 25$ cm, $DC = 15$ cm and the perpendicular distance between AB and DC is 18 cm.




(a) Calculate the volume of the trough in litres.

Volume = ... litres **[4]**

(b) The trough is initially completely empty.
Water flows into the trough at a constant rate of 0.8 litres per second.
Calculate the depth of the water in the trough after 30 seconds.

Depth = ... cm **[5]**

1018 A solid is made by removing a small cone from a large cone to form a frustum.
The large cone has a base radius of 10 cm and a perpendicular height of 24 cm.
The small cone has a base radius of 5 cm and a perpendicular height of 12 cm.



NOT TO SCALE

(a) Calculate the volume of the frustum.

..... cm^3 **[3]**


(b) Calculate the total surface area of the solid frustum.

..... cm^2 **[4]**

(c) The solid frustum is melted down to form solid spheres of radius 3 cm.
Calculate the maximum number of complete spheres that can be made.

..... **[3]**

1019 A solid metal pyramid has a square base of side 12 cm and a perpendicular height of 10 cm.



(a) Calculate the volume of the pyramid.

..... cm^3 **[2]**

(b) Calculate the total surface area of the pyramid.

..... cm^2 **[4]**


(c) The metal pyramid is melted down and recast to make solid spheres of radius 2.5 cm.
Calculate the number of complete spheres that can be made.

..... **[3]**

(d) A mathematically similar pyramid has a volume of 1620 cm^3.
Calculate the perpendicular height of this similar pyramid.

..... cm **[3]**


1020 A shape is made from a rectangle and a semicircle.
The rectangle has a length of 10 cm and a width of 6 cm.
The semicircle is attached to one of the 6 cm sides.



Find the exact perimeter of the shape.
Give your answer in the form $a + b\pi$.

..... cm **[3]**

1021 The diagram shows a square $ABCD$ with side length 10 cm.
A quarter circle with center A and radius 10 cm is drawn inside the square.
The region inside the square but outside the quarter circle is shaded.




(a) Find the exact perimeter of the shaded region.
Give your answer in terms of π.

..... cm **[2]**

(b) Find the exact area of the shaded region.
Give your answer in terms of π.

..... cm^2 **[2]**

1022 The diagram shows a compound shape formed by joining a semi-circle to a rectangle.
The rectangle has a length of 8 cm and a width of 4 cm.
The diameter of the semi-circle is 4 cm.




(a) Calculate the perimeter of the shape.
Give your answer in its simplest form, in terms of π.

..... *cm* **[2]**

(b) Calculate the area of the shape.
Give your answer in its simplest form, in terms of π.

..... cm^2 **[2]**

1023 A compound shape is made from a rectangle and a semicircle.
The rectangle has width 8 cm and length 12 cm. The semicircle is attached to one of the sides of width 8 cm.




(a) Calculate the exact perimeter of the compound shape.
Give your answer in its simplest form, in terms of π.

..... cm **[2]**

(b) Calculate the exact area of the compound shape.
Give your answer in its simplest form, in terms of π.

..... cm^2 **[2]**

1024 The diagram shows a shape made from a rectangle and a semicircle.
The rectangle has length 12 cm and width 8 cm.
The semicircle has a diameter of 8 cm.




(a) Find the perimeter of the shape.
Give your answer in its simplest form in terms of π.

Perimeter = ... cm **[2]**

(b) Find the area of the shape.
Give your answer in its simplest form in terms of π.

Area = ... cm^2 **[3]**

1025 The diagram shows a compound shape made from a rectangle and a semicircle.
The rectangle has a length of 10 cm and a width of 8 cm.
The semicircle is attached to one of the 8 cm sides.




(a) Work out the exact area of the shape.
Give your answer in the form $a + b\pi$, where a and b are integers.

Area = ... cm^2 **[3]**

(b) Work out the exact perimeter of the shape.
Give your answer in the form $c + d\pi$, where c and d are integers.

Perimeter = ... cm **[3]**

1026 The diagram shows a compound shape made from a rectangle and a semicircle.
The rectangle has a length of 10 cm and a width of 6 cm.
A semicircle of diameter 6 cm has been removed from one end of the rectangle.



NOT TO SCALE


(a) Work out the perimeter of the shape.
Give your answer in the form $a + b\pi$, where a and b are integers.

..... cm **[3]**

(b) Work out the area of the shape.
Give your answer in the form $c - d\pi$, where c is an integer and d is a decimal.

..... cm^2 **[3]**

1027 The diagram shows a shape formed by a square of side 4 cm and a quarter of a circle with center C.




(a) Find the area of the shape.
Give your answer in terms of π.

Area = ... cm^2 **[3]**

(b) Find the perimeter of the shape.
Give your answer in terms of π.

Perimeter = ... cm **[3]**

1028 The diagram shows a compound shape made from a rectangle and a semicircle.
The rectangle has length 12 cm and width 10 cm.
The semicircle has a diameter of 10 cm.




(a) Calculate the perimeter of the shape.
Give your answer in the form $a + b\pi$.

..... cm **[3]**

(b) Calculate the area of the shape.
Give your answer in the form $c + d\pi$.

..... cm^2 **[3]**

1029 The diagram shows a shape made from a rectangle with dimensions 10 cm by 6 cm. A semicircle of diameter 6 cm has been removed from one end.



NOT TO SCALE


(a) Calculate the area of the shape.
Give your answer in terms of π.

..... cm^2 **[3]**

(b) Calculate the perimeter of the shape.
Give your answer in terms of π.

..... cm **[3]**

1030 A solid is made by attaching a hemisphere of radius 5 cm to one end of a solid cylinder.
The cylinder has a radius of 5 cm and a height of 12 cm.




(a) Calculate the total volume of the solid.
Give your answer correct to 3 significant figures.

..... cm^3 **[3]**

(b) Calculate the total surface area of the solid.
Give your answer correct to 3 significant figures.

..... cm^2 **[3]**

1031 The diagram shows a solid in the shape of a cylinder with a hemisphere attached to the top.
The radius of both the cylinder and the hemisphere is 5 cm.
The total height of the solid is 25 cm.



NOT TO SCALE


(a) Calculate the total volume of the solid.

..... cm^3 **[3]**

(b) Calculate the total surface area of the solid.

..... cm^2 **[4]**

1032 The diagram shows a solid compound shape made from a cylinder and a cone.
The cylinder and the cone both have a base radius of 5 cm.
The height of the cylinder is 12 cm and the height of the cone is 6 cm.




(a) Calculate the volume of the solid.

..... cm^3 **[3]**

(b) Calculate the total surface area of the solid.

..... cm^2 **[4]**

1033 A solid is made by placing a solid hemisphere on top of a solid cylinder.
The cylinder and the hemisphere both have radius 6 cm.
The total height of the solid is 20 cm.




(a) Calculate the volume of the solid.

..... cm^3 **[3]**

(b) Calculate the total surface area of the solid.

..... cm^2 **[4]**

1034 A solid is made from a cylinder and a cone.
The cylinder has radius 6 cm and height 15 cm.
The cone has radius 6 cm and height 8 cm.




(a) Calculate the total volume of the solid.

Volume = ... cm^3 **[3]**

(b) Calculate the total surface area of the solid.

Surface area = ... cm^2 **[4]**

1035 A solid is made by attaching a hemisphere of radius 3 cm to one end of a solid cylinder of radius 3 cm and height 10 cm.




(a) Calculate the volume of the solid.

Volume = ... cm^3 **[3]**

(b) Calculate the total surface area of the solid.

Total surface area = ... cm^2 **[4]**

1036 A solid object is formed by joining a hemisphere to one end of a solid cylinder.
The cylinder has a radius of 4 cm and a length of 12 cm.
The hemisphere has the same radius as the cylinder.




(a) Calculate the volume of the solid.

Volume = ... cm^3 **[3]**

(b) Calculate the total surface area of the solid.

Surface area = ... cm^2 **[4]**

1037 The diagram shows a solid frustum.
The frustum is made by removing a smaller cone from a larger cone.
The larger cone has a base radius of 8 cm and a vertical height of 24 cm.
The smaller cone has a base radius of 5 cm and a vertical height of 15 cm.




(a) Calculate the volume of the solid frustum.

..... cm^3 **[3]**

(b) Calculate the total surface area of the solid frustum.

..... cm^2 **[5]**

1038 A solid is formed by placing a hemisphere of radius 5 cm on top of a cylinder of radius 5 cm.
The total height of the solid is 18 cm.



NOT TO SCALE


(a) Calculate the volume of the solid.
Give your answer correct to 3 significant figures.

..... cm^3 **[4]**

(b) Calculate the total surface area of the solid.
Give your answer correct to 3 significant figures.

..... cm^2 **[4]**

1039 The diagram shows a solid made by joining a solid hemisphere to a solid cone.
The common radius of the hemisphere and the base of the cone is 5 cm.
The total height of the solid is 17 cm.



(a) Calculate the volume of the solid.

Volume = ... cm^3 **[3]**


(b) Calculate the total surface area of the solid.

Surface area = ... cm^2 **[3]**

(c) The solid is melted down and completely recast into a single solid cylinder of radius 4 cm.
Calculate the height of the cylinder.

Height = ... cm **[3]**

1040 A solid cone of height 24 cm and base radius 10 cm is cut parallel to its base to form a frustum and a smaller cone.
The radius of the top circular face of the frustum is 4 cm.



(a) Show that the height of the removed small cone is 9.6 cm.

[2]


(b) Calculate the volume of the frustum.

..... cm^3 **[3]**

(c) The solid frustum is melted down and recast entirely into a solid sphere.
Calculate the radius of this sphere.

..... cm **[3]**

1041 The diagram shows a solid metal frustum.
The frustum is formed by removing a small cone of height 10 cm from a larger cone.
The radius of the circular base of the frustum is 12 cm and the radius of the circular top is 4 cm.
The height of the frustum is 20 cm.




(a) Calculate the volume of the solid frustum.

..... cm^3 **[4]**

(b) Calculate the total surface area of the solid frustum.

..... cm^2 **[4]**

1042 A container is in the shape of a frustum of a cone.
The original cone has a base radius of 15 cm and a vertical height of 40 cm.
A smaller cone of vertical height 16 cm is removed from the top to form the frustum.



(a) Show that the radius r of the top circular face of the frustum is 6 cm.

[2]


(b) Calculate the volume of the frustum.

Volume = ... cm^3 **[3]**

(c) The solid frustum is melted down and recast entirely into a solid sphere.
Calculate the radius of the sphere.

Radius = ... cm **[3]**

1043 A solid frustum of a cone has top radius 4 cm, bottom radius 8 cm and vertical height 15 cm.
It is formed by removing a smaller cone from the top of a larger cone.



(a) Show that the original height of the cone before the top was removed was 30 cm.

[2]


(b) Calculate the volume of the frustum.

Volume = ... cm^3 **[3]**

(c) Calculate the total surface area of the solid frustum.

Surface area = ... cm^2 **[4]**

1044 The diagram shows three identical circles, each of radius r cm.
The circles touch each other externally.
The region enclosed between the three circles is shaded.



(a) Show that the area of the shaded region, A, is given by the formula

$$A = r^2\left(\sqrt{3} - \frac{\pi}{2}\right)$$

[4]

(b) The area of the shaded region is 15 cm^2.


Work out the value of r.

$r =$ ..... **[3]**

(c) Calculate the total perimeter of the shaded region.

..... cm **[3]**

1045 A frustum is made by removing a smaller cone from the top of a larger cone.
The larger cone has base radius 10 cm and vertical height 24 cm.
The frustum has vertical height 15 cm.



(a) Find the radius of the top of the frustum.

..... cm **[2]**


(b) Calculate the volume of the frustum.

..... cm^3 **[3]**

(c) Calculate the total surface area of the frustum.

..... cm^2 **[5]**

1046 A frustum is made by removing a small cone from the top of a large cone.
The large cone has a base radius of 8 cm and a total height of 24 cm.
The removed small cone has a base radius of 3 cm.



(a) Show that the height of the small cone removed is 9 cm.

[2]


(b) Calculate the volume of the frustum.

Volume = ... cm^3 **[3]**

(c) Calculate the total surface area of the frustum.

Surface area = ... cm^2 **[5]**

1047 The diagram shows a solid frustum of a cone.
The frustum is made by removing a small cone from the top of a larger solid cone.
The base radius of the frustum is 12 cm and the top radius is 4 cm.
The height of the frustum is 15 cm.



(a) Calculate the height of the small cone that was removed.

Height = ... cm **[3]**


(b) Calculate the volume of the solid frustum.

Volume = ... cm^3 **[3]**

(c) Calculate the total surface area of the solid frustum.

Surface area = ... cm^2 **[5]**

1048 The diagram shows a compound solid made from a frustum of a cone and a solid hemisphere.
The frustum is created by removing a small cone from a large cone.
The base radius of the frustum is 12 cm and its top radius is 6 cm.
The height of the frustum is 8 cm.
A solid hemisphere of radius 6 cm is attached to the top face of the frustum.



NOT TO SCALE

(a) Show that the perpendicular height of the large cone from which the frustum is made is 16 cm.

[3]


(b) Calculate the total volume of the compound solid.

..... cm^3 **[4]**

(c) Calculate the total surface area of the compound solid.

..... cm^2 **[4]**

1049 The diagram shows a closed container in the shape of a frustum of a cone.
The radius of the top circular face is 6 cm and the radius of the base is 4 cm.
The vertical height of the container is 10 cm.



NOT TO SCALE

(a) The frustum is made by removing a smaller cone from a larger cone.
Calculate the vertical height of the larger cone.

..... cm **[3]**


(b) Calculate the volume of the container.
Give your answer correct to 3 significant figures.

..... cm^3 **[3]**

(c) Calculate the total surface area of the closed container.
Give your answer correct to 3 significant figures.

..... cm^2 **[5]**

1050 A chord of a circle, centre O, has length 24 cm. The perpendicular distance from the centre of the circle to the chord is 5 cm.




Calculate the radius of the circle.

..... cm **[3]**

1051 The points $A(-4, 6)$ and $B(8, 1)$ lie on a coordinate grid.
Work out the exact length of the line segment AB.

..... **[3]**

1052 Points $A(-2, 1)$ and $B(4, 9)$ are plotted on a coordinate grid.



Calculate the exact length of the line segment AB.


$AB =$ ..... units **[3]**

1053 The point A has coordinates $(1, 4)$ and the point B has coordinates $(k, 19)$.
The exact distance between point A and point B is 17.

Find the two possible values of k.

$k =$ ..... , $k =$ ..... **[4]**

1054 The diagram shows a right-angled triangle.
All lengths are in centimetres.




(a) Show that $x^2 - 16x + 48 = 0$.

[3]

(b) Solve the equation to find the value of x, and hence calculate the perimeter of the triangle.

Perimeter = ... cm **[4]**

1055 The diagram shows two intersecting circles.
The circle with centre P has radius 10 cm.
The circle with centre Q has radius 17 cm.
The points A and B are the intersections of the two circles.
The length of the common chord AB is 16 cm.
M is the point of intersection of the chord AB and the line PQ.



NOT TO SCALE


(a) Show that the distance PM is 6 cm.

[2]

(b) Calculate the distance PQ.

..... cm **[3]**

1056 The diagram shows a right-angled triangle.
The lengths of the sides are $x - 1$, $x + 6$, and $2x + 1$, where x is an integer.
All lengths are in centimetres.



(a) Show that $x^2 - 3x - 18 = 0$.

[4]


(b) Solve the equation by factorising to find the value of x.
You must show your working.

$x =$ ..... **[3]**

(c) Work out the area of the triangle.

..... cm^2 **[3]**

1057 The diagram shows a circle, centre O, with radius 8.5 cm.
AB is a chord of the circle.
The perpendicular distance from O to the chord AB is 4 cm.




NOT TO SCALE

Calculate the length of the chord AB.

..... cm **[3]**

1058 The diagram shows a circle, centre O, with radius 8.5 cm.
AB is a chord of length 15 cm.
M is the midpoint of AB.



(a) Calculate the distance OM, labeled d in the diagram.


$d =$ ..... cm **[3]**

(b) A second chord, CD, is drawn in the same circle such that its perpendicular distance from the centre O is 7.5 cm.

Calculate the length of the chord CD.

Length = ... cm **[3]**

1059 The diagram shows a circle, centre O, with radius 12.4 cm.
AB is a chord of the circle.
The perpendicular distance from O to the chord AB is 5.2 cm.



(a) Calculate the length of the chord AB.


..... cm **[3]**

(b) Another chord CD is drawn parallel to AB and on the opposite side of the centre O.
The length of the chord CD is 18 cm.

Calculate the perpendicular distance between the two chords AB and CD.

..... cm **[4]**

1060 The diagram shows a coordinate grid.
Point A is $(-2, -1)$ and point B lies on the line $x = 6$.
The y-coordinate of B is positive.
The distance AB is 10 units.




(a) Find the y-coordinate of point B.

$y =$ ..... **[3]**

(b) Point C lies on the x-axis.
The triangle ABC is isosceles with $AC = BC$.
Calculate the x-coordinate of C.

$x =$ ..... **[4]**

1061 A right-angled triangle has sides of length x cm, $(x + 7)$ cm and a hypotenuse of length $(x + 8)$ cm.



(a) Use Pythagoras' theorem to write down an equation in x and show that it simplifies to $x^2 - 2x - 15 = 0$

[3]


(b) Solve the equation to find the value of x.

$x =$ ..... **[2]**

(c) Calculate the area of the triangle.


Area = ... cm^2 **[2]**

1062 A straight ladder of length 6.5 m leans against a vertical wall.
The base of the ladder is 2.1 m from the bottom of the wall on horizontal ground.



Calculate the angle the ladder makes with the horizontal ground.

..... ° **[3]**




NOT TO SCALE

1063 Triangle XYZ is right-angled at Y.
Angle $ZXY = 58°$ and $XY = 9.4$ cm.

Calculate the length of XZ.

..... cm **[3]**

1064 A surveyor stands at a point P, which is 45 m away from the base, B, of a vertical tower TB. The angle of elevation of the top of the tower, T, from P is 38°.



(a) Calculate the height of the tower, TB.

..... m **[2]**


(b) Calculate the straight-line distance PT.

..... m **[2]**

(c) A bird sits at a point X on the tower, where $BX = 20$ m.
Calculate the angle of elevation of the bird from P.

..... ° **[3]**

1065 A boat sails from a point A in a straight line on a bearing of 052°.
A lighthouse, L, is located 18 km due North of A.



NOT TO SCALE

(a) Calculate the shortest distance from the lighthouse L to the path of the boat.


..... km **[2]**

(b) The boat travels at a constant speed of 14 km/h.

Calculate the time it takes for the boat to reach the point on its path closest to the lighthouse.
Give your answer in minutes, correct to the nearest minute.

..... minutes **[3]**

1066 A ship sails from port P on a bearing of 040° for 80 km to point A.
It then changes course and sails on a bearing of 130° for 60 km to point B.



(a) Calculate the direct distance from P to B.

$PB =$ ..... km **[3]**


(b) Calculate the bearing of B from P.

Bearing = ... ° **[3]**

(c) Calculate the shortest distance from point A to the path PB.

..... km **[3]**

1067 A ship sails from port A to point B on a bearing of 052° for 15 km.
It then sails from point B to point C on a bearing of 142° for 20 km.



(a) Show that angle $ABC = 90°$.

[2]


(b) Calculate the distance AC.

..... km **[3]**

(c) Calculate the bearing of C from A.

..... ° **[4]**

1068 The diagram shows a quadrilateral $PQRS$ formed by two right-angled triangles, PQR and PRS.
Angle $PQR = 90°$ and angle $PRS = 90°$.
$PQ = 8.5$ cm, $QR = 11.2$ cm and angle $PSR = 41°$.



NOT TO SCALE

(a) Calculate the length of PR.

..... cm **[2]**


(b) Calculate the length of PS.

..... cm **[2]**

(c) Calculate the perpendicular distance from R to the line PS.

..... cm **[2]**

1069 Point A and point B are on horizontal ground, in a straight line with the base C of a vertical building CD.
The distance between A and B is 50 m.
The angle of elevation of the top of the building, D, from A is 25° and from B is 42°.




(a) Calculate the height of the building CD.

$CD =$ ..... m **[5]**

(b) A bird flies directly along the straight path from A to D.
Calculate the shortest distance from point B to the flight path AD.

..... m **[3]**

1070 The diagram shows a quadrilateral $ABCD$.
Angle $ABC = 90°$ and angle $ADC = 90°$.
$AB = 8$ cm, $BC = 15$ cm, and $CD = 12$ cm.



(a) Calculate the length of AC.

..... cm **[2]**

(b) Calculate the length of AD.

..... cm **[3]**

(c) Calculate the shortest distance from D to the line AC.

..... cm **[3]**

(d) Calculate the size of angle DAB.

..... ° **[4]**

1071 Without using a calculator, answer the following questions.

(a) Write down the exact value of $\cos 30°$.

..... **[1]**

(b) Show that

$$\frac{1-\sin 30°}{\tan^2 60°} = \frac{1}{6}$$

[2]

1072 Without using a calculator, write down the exact value of each of the following expressions.
You must show all your working.

(a) $\sin 90^\circ + \cos 0^\circ$

..... **[1]**

(b) $\tan 45^\circ \times \sin 30^\circ$

..... **[1]**

(c) $\dfrac{\tan 60^\circ}{\cos 30^\circ}$

..... **[2]**

1073 **(a)** Work out the exact value of

$$2\sin 60° \times \tan 30°$$


..... **[2]**

(b) Find the value of x, where $0° \leq x \leq 90°$, such that

$$\sqrt{2}\cos x = \tan 45°$$

$x =$ ..... ° **[2]**

1074 The diagram shows two right-angled triangles, PQR and PRS.
$PQ = 6$ cm, angle $PQR = 90°$ and angle $PRQ = 45°$.
Angle $PRS = 90°$ and angle $PSR = 60°$.




(a) Calculate the exact length of PR.

..... cm **[2]**

(b) Calculate the exact length of PS.
Give your answer in the form $a\sqrt{b}$ where a and b are integers.

..... cm **[3]**

1075 The diagram shows two right-angled triangles, PQR and PRS.
Angle $PQR = 90°$ and angle $PRS = 90°$.
Angle $PRQ = 45°$ and angle $SPR = 30°$.
$PR = 8\sqrt{2}$ cm.



NOT TO SCALE


(a) Without using a calculator, show that the exact length of PQ is 8 cm.

[2]

(b) Without using a calculator, work out the exact length of PS.
Give your answer in the form $\frac{a\sqrt{b}}{c}$, where a, b and c are integers.

$PS =$ ..... cm **[3]**

1076 The diagram shows a triangle ABC with a point D on the side AC.
BD is perpendicular to AC.
$AB = 8\sqrt{2}$ cm, $\angle BAC = 45°$ and $\angle BCA = 30°$.



Calculate the exact perimeter of triangle ABC.
Give your answer in the form $a + b\sqrt{2} + c\sqrt{3}$ where a, b, and c are integers.

Perimeter = ... cm **[6]**

1077 Do not use a calculator for this question.
All working must be clearly shown.

(a) Show that

$$\frac{1-\tan^2 30^\circ}{1+\tan^2 30^\circ} = \cos 60^\circ$$

[3]


(b) Solve the equation.

$$x \sin 60^\circ = \tan 60^\circ - x \cos 60^\circ$$

Give your answer in the form $a - \sqrt{b}$, where a and b are integers.

$x =$ ..... **[4]**

1078 The diagram shows a rectangle $ABCD$.
E is a point on the side CD.
$AE = 12$ cm, angle $DAE = 30°$ and angle $EAB = 60°$.



(a) Show that the exact length of AD is $6\sqrt{3}$ cm.

[2]


(b) The exact area of the rectangle $ABCD$ is $60\sqrt{3}$ cm^2.
Work out the exact length of CE.

..... cm **[3]**

(c) Find the exact value of $\tan(\angle EBC)$.
Give your answer in the form $\dfrac{a\sqrt{b}}{c}$ where a, b, and c are integers.

..... **[3]**

1079 The diagram shows a triangle PQR.
$PQ = x$ cm, $PR = (x + 3)$ cm and $QR = 7$ cm.
Angle $QPR = 60°$.



(a) Show that $x^2 + 3x - 40 = 0$.

[4]

(b) Solve the equation to find the value of x, and hence find the perimeter of triangle PQR.

Perimeter = ... cm **[3]**

(c) Find the exact area of triangle PQR.


Area = ... cm^2 **[3]**

1080 Solve the equation for $0° \leq x \leq 360°$.

$\sqrt{2}\sin x - 1 = 0$

$x =$ ..... **[3]**

1081 The diagram shows the graph of $y = \cos x$ for $0° \leq x \leq 360°$.




(a) Write down the coordinates of the lowest point on the graph.

(..... , ...) **[1]**

(b) Use the graph, or otherwise, to solve the equation $\cos x = 0.5$ for $0° \leq x \leq 360°$.

$x =$ ..... **[2]**

1082 **(a)** On the grid, sketch the graph of $y = \sin x$ for $0° \leq x \leq 360°$.



[2]

(b) Write down the exact solutions to the equation $\sin x = 0.5$ for $0° \leq x \leq 360°$.

$x =$ ..... , $x =$ ..... **[2]**

1083 Answer the following questions for $0° \leq x \leq 360°$.

(a) Solve the equation

$$2\cos x - 1 = 0$$

$x =$ ..... **[3]**


(b) Write down the number of solutions to the equation

$$3\sin x = 5$$

..... **[1]**

1084 The properties of trigonometric graphs can be used to solve equations.

(a) On the grid, sketch the graph of $y = \cos x$ for $0° \leq x \leq 360°$.



[2]

(b) Write down the coordinates of the minimum point on this graph.

(..... , ...) **[1]**


(c) Solve the equation for $0° \leq x \leq 360°$.

$2\cos x + 1 = 0$

$x =$ ..... **[3]**

1085 The functions $g(x) = \sin x$ and $h(x) = \cos x$ are defined for $0° \leq x \leq 360°$.

(a) On the grid provided, sketch the graphs of $y = g(x)$ and $y = h(x)$.



[3]

(b) Solve the equation $g(x) = h(x)$ for $0° \leq x \leq 360°$.

$x =$ ..... **[2]**

(c) Use your graphs to write down the range of values of x for which $g(x) > h(x)$ for $0° \leq x \leq 360°$.

..... **[2]**

1086 Solve the following equations for $0° \leq x \leq 360°$.

(a) $2\sin x + \sqrt{3} = 0$

$x =$ ..... **[3]**

(b) $4\cos^2 x - 3 = 0$

$x =$ ..... **[4]**

1087 Consider the trigonometric function $f(x) = 4\sin x + 3$ for $0° \leq x \leq 360°$.


(a) Solve the equation $f(x) = 0$.
Give your answers correct to 1 decimal place.

$x =$ ..... **[3]**

(b) Write down the exact coordinates of the minimum point on the graph of $y = f(x)$.

(..... , ...) **[2]**

1088 **(a)** On the axes provided, sketch the graph of $y = \tan x$ for $0° \leq x \leq 360°$.



[2]

(b) Solve the equation $3\tan x + 5 = 0$ for $0° \leq x \leq 360°$.
Give your answers correct to 1 decimal place.

$x =$ ..... **[3]**

1089 **(a)** Solve the equation for $0° \leq x \leq 360°$.

$$4\cos x + 1 = 0$$

Give your answers correct to 1 decimal place.

$x =$ ..... , $x =$ ..... **[3]**

(b) Solve the equation for $0° \leq x \leq 360°$.

$$2\tan x - 5 = 0$$

Give your answers correct to 1 decimal place.

$x =$ ..... , $x =$ ..... **[3]**

1090 **(a)** The equation $a\sin x - b = 0$ has a solution $x = 30°$.
Given that a and b are positive integers with no common factors, find the value of a and the value of b.

$a =$ ..... , $b =$ ..... **[2]**

(b) Write down the second solution to the equation $a\sin x - b = 0$ for $0° \leq x \leq 360°$.

$x =$ ..... ° **[1]**

(c) A horizontal line $y = k$ intersects the graph of $y = \cos x$ at exactly two points in the interval $0° \leq x \leq 360°$.
The x-coordinate of one of these points is $242°$.
Find the value of k correct to 3 significant figures and find the x-coordinate of the second point of intersection.

$k =$ ..... , $x =$ ..... **[3]**

1091 Algebraic techniques can be applied to solve trigonometric equations.

(a) Solve the equation for $0° \leq x \leq 360°$.

$$5\tan x + 14 = 0$$

Give your answers correct to 1 decimal place.


$x =$ ..... **[3]**

(b) Solve the equation for $0° \leq x \leq 360°$.

$$2\sin^2 x - \sin x = 0$$

$x =$ ..... **[4]**

1092 The diagram shows triangle PQR.
$PQ = 8$ cm, $PR = 5$ cm and angle $QPR = 60°$.



NOT TO SCALE


(a) Calculate the exact area of triangle PQR.
Give your answer in the form $a\sqrt{3}$.

..... cm^2 **[2]**

(b) Calculate the exact length of QR.

$QR =$ ..... cm **[3]**

1093 In triangle ABC, $AB = 5\,\text{cm}$, $BC = x\,\text{cm}$ and $AC = 7\,\text{cm}$.
Angle $ABC = 60°$.



NOT TO SCALE

(a) Use the cosine rule to show that $x^2 - 5x - 24 = 0$.

[3]


(b) Solve the equation $x^2 - 5x - 24 = 0$ to find the length of BC.

$BC =$ ..... cm **[2]**

(c) Calculate the exact area of triangle ABC.
Give your answer in the form $a\sqrt{b}$, where a and b are integers.

Area = ... cm^2 **[2]**

1094 The diagram shows triangle ABC.
$AB = 8\,\text{cm}$, $AC = 12\,\text{cm}$ and angle $BAC = 42°$.




(a) Calculate the area of triangle ABC.

..... cm^2 **[2]**

(b) Calculate the length of BC.

$BC =$ ..... cm **[3]**

1095 The diagram shows a triangle ABC.




(a) Calculate the area of triangle ABC.

..... cm^2 **[2]**

(b) Calculate the length of BC.

..... cm **[3]**

1096 The diagram shows triangle ABC.




(a) Calculate the area of triangle ABC.

..... cm^2 **[2]**

(b) Calculate the length of BC.

..... cm **[3]**

1097 A triangle ABC is shown below.



NOT TO SCALE


(a) Calculate the area of triangle ABC.

..... cm^2 **[2]**

(b) Calculate the length of BC.

..... cm **[3]**

1098 In triangle PQR, $PQ = 7.2\,\text{cm}$, $PR = 9.5\,\text{cm}$ and angle $QPR = 64°$.



NOT TO SCALE


(a) Calculate the area of triangle PQR.

Area = ... cm^2 **[2]**

(b) Calculate the length of QR.

QR = ... cm **[3]**

1099 The diagram shows a triangle ABC with $AB = 8\,\text{cm}$, $AC = 12\,\text{cm}$ and angle $BAC = 65°$.



NOT TO SCALE


(a) Calculate the length of BC.

$BC =$ ..... cm **[3]**

(b) Calculate the area of triangle ABC.

Area = ... cm^2 **[2]**

1100 The diagram shows a triangle ABC.
$AB = 8$ cm, $AC = 11$ cm and angle $BAC = 65°$.




(a) Calculate the area of triangle ABC.

Area = ... cm^2 **[2]**

(b) Calculate the length of BC.

BC = ... cm **[3]**

1101 In triangle ABC, $AC = 12$ cm, angle $BAC = 45°$ and angle $ABC = 60°$.




(a) Calculate the length of BC.

..... cm **[3]**

(b) Calculate the area of triangle ABC.

..... cm^2 **[3]**

1102 The diagram shows triangle ABC.
$AC = 12$ cm, angle $BAC = 40°$ and angle $ABC = 60°$.




(a) Calculate the length of BC.

$BC =$ ..... cm **[3]**

(b) Calculate the area of triangle ABC.

Area = ... cm^2 **[3]**

1103 The diagram shows triangle PQR.
$PQ = 7$ cm, $QR = 8$ cm and $PR = 12$ cm.




(a) Calculate the largest angle in triangle PQR.

..... ° **[3]**

(b) Calculate the area of triangle PQR.

Area = ... cm^2 **[2]**

1104 In triangle PQR, $PR = 8$ cm, $QR = 12$ cm and angle $PQR = 35°$.
Angle QPR is an obtuse angle.



NOT TO SCALE


(a) Calculate the size of angle QPR.

..... ° **[4]**

(b) Calculate the area of triangle PQR.

..... cm^2 **[2]**

1105 The diagram shows a triangle PQR.
The length of PQ is 15 cm and the length of PR is 12 cm.
Angle $PQR = 42°$.
Angle PRQ is obtuse.




(a) Calculate angle PRQ.

..... ° **[3]**

(b) Calculate the length of QR.

..... cm **[3]**

1106 The diagram shows triangle PQR.
$PQ = 14$ cm, $QR = 10$ cm and angle $QPR = 35°$.
Angle PRQ is obtuse.




(a) Calculate angle PRQ.

Angle $PRQ =$ ..... ° **[3]**

(b) Calculate the area of triangle PQR.

Area = ... cm^2 **[3]**

1107 The diagram shows triangle PQR.
The angle PQR is obtuse.




(a) Calculate the obtuse angle PQR.

..... ° **[4]**

(b) Calculate the area of triangle PQR.

..... cm^2 **[3]**

1108 The diagram shows triangle PQR.
$PQ = 15\,\text{cm}$, $QR = 12\,\text{cm}$ and $PR = 18\,\text{cm}$.



(a) Use the cosine rule to show that angle $PQR = 82.8°$, correct to 1 decimal place.

[3]


(b) Calculate the area of triangle PQR.

..... cm^2 **[2]**

(c) Calculate the shortest distance from R to the line PQ.

..... cm **[2]**

1109 In triangle ABC, $AB = 12\,\text{cm}$, $BC = 9\,\text{cm}$ and angle $BAC = 35°$.



NOT TO SCALE


(a) Calculate the two possible values of angle ACB.
Give your answers correct to 1 decimal place.

..... ° **[4]**

(b) Given that angle ACB is obtuse, calculate the area of triangle ABC.

Area = ... cm^2 **[3]**

1110 The diagram shows a triangle PQR.
$PQ = 15.2\,\text{cm}$, $PR = 18.5\,\text{cm}$ and angle $PQR = 72°$.



NOT TO SCALE

(a) Calculate the size of angle PRQ.

Angle $PRQ =$ ..... ° **[3]**


(b) Calculate the length of QR.

$QR =$ ..... cm **[3]**

(c) Calculate the shortest distance from P to the line QR.

Distance = ... cm **[2]**

1111 The diagram shows a quadrilateral $ABCD$.
$AB = 8$ cm, $AD = 10$ cm, $BC = 14$ cm and $CD = 15$ cm.
Angle $BAD = 110°$.



(a) Calculate the length of the diagonal BD.

..... cm **[3]**


(b) Calculate the angle BCD.

..... ° **[3]**

(c) Calculate the area of the quadrilateral $ABCD$.

..... cm^2 **[3]**

1112 The diagram shows a quadrilateral $ABCD$.
$AB = 12.4$ cm, $BC = 9.8$ cm and angle $ABC = 105°$.
Angle $CAD = 41°$ and angle $ADC = 68°$.



NOT TO SCALE

(a) Calculate the length of AC.
Give your answer correct to 3 significant figures.

$AC =$ ..... cm **[3]**

(b) Calculate the length of CD.
Give your answer correct to 3 significant figures.

$CD =$ ..... cm **[3]**

(c) Calculate the total area of the quadrilateral $ABCD$.

..... cm^2 **[3]**

1113 In triangle ABC, $AB = 15$ cm, $BC = 12$ cm and angle $BAC = 48°$.


(a) Calculate the two possible values of angle BCA.

and ..... **[3]**

(b) Hence, calculate the difference between the two possible areas of triangle ABC.

..... cm^2 **[4]**

1114 In triangle ABC, $AB = 10.5\,\text{cm}$, $BC = 8.2\,\text{cm}$ and angle $BAC = 35°$.
Angle BCA is obtuse.



(a) Calculate the size of angle BCA.

..... ° **[3]**


(b) Calculate the length of AC.

$AC =$ ..... cm **[3]**

(c) A point D lies outside the triangle ABC such that $CD = 6.4\,\text{cm}$ and angle ACD is acute.
The area of triangle ACD is $8.5\,\text{cm}^2$.
Calculate the size of angle ACD.

..... ° **[2]**

1115 The diagram shows triangle DEF.
$DE = 10.5$ cm, $DF = 8.2$ cm and angle $DEF = 42°$.
Angle DFE is an obtuse angle.



(a) Calculate angle DFE.

Angle $DFE =$ ..... ° **[3]**


(b) Calculate the length of EF.

$EF =$ ..... cm **[3]**

(c) Calculate the shortest distance from D to the line EF.

..... cm **[2]**

1116 The diagram shows a quadrilateral $ABCD$.
$AB = 6$ cm, $AD = 8$ cm, $DC = 5$ cm, and $BC = 9$ cm.
Angle $DAB = 110°$.



NOT TO SCALE

(a) Calculate the length of the diagonal BD.

..... cm **[3]**


(b) Calculate the size of angle BCD.

..... ° **[3]**

(c) Calculate the total area of the quadrilateral $ABCD$.

..... cm^2 **[3]**

1117 The diagram shows a quadrilateral $ABCD$.
$AB = 7$ cm, $AD = 9$ cm, $BC = 10$ cm and $CD = 14$ cm.
Angle $BAD = 110°$.



(a) Calculate the length of BD.

..... cm **[3]**

(b) Calculate angle BCD.

..... ° **[3]**

(c) Calculate the total area of the quadrilateral $ABCD$.

..... cm^2 **[3]**

1118 The diagram shows a quadrilateral $ABCD$.
$AB = 7.4$ cm, $BC = 9.1$ cm, and $CD = 10.5$ cm.
Angle $ABC = 112°$ and angle $CAD = 43°$.

(a) Calculate the length of AC.

..... cm **[3]**


(b) Calculate the acute angle ADC.

..... ° **[3]**

(c) Calculate the shortest distance from D to the line AC.

..... cm **[3]**

1119 The diagram shows a quadrilateral $ABCD$.
$AB = 7.5$ cm, $AD = 9.2$ cm and angle $BAD = 105°$.
$BC = 13$ cm and $CD = 10.5$ cm.



(a) Calculate the length of BD.

$BD =$ ..... cm **[3]**


(b) Calculate angle BCD.

Angle $BCD =$ ..... ° **[3]**

(c) Calculate the area of the quadrilateral $ABCD$.

Area = ... cm^2 **[3]**

1120 The diagram shows a quadrilateral $ABCD$.
$AB = 8.5\,\text{cm}$, $AD = 11.2\,\text{cm}$ and angle $BAD = 105°$.
$BC = 11.0\,\text{cm}$ and angle $BDC = 38°$.
Angle BCD is obtuse.



NOT TO SCALE

(a) Calculate the length of BD.

$BD =$ ..... cm **[3]**


(b) Calculate the size of angle BCD.

Angle $BCD =$ ..... ° **[3]**

(c) Calculate the area of the quadrilateral $ABCD$.

Area = ... cm^2 **[4]**

1121 Ship A and Ship B are at sea.
Ship B is 40 km from Ship A on a bearing of 050°.
A lighthouse L is 35 km from Ship B.
The bearing of L from A is 085°.
Given that the angle ALB is obtuse:



(a) Calculate the obtuse angle ALB.

..... ° **[4]**


(b) Calculate the distance AL.

..... km **[3]**

(c) Calculate the bearing of L from B.

..... ° **[3]**

1122 The diagram shows a cuboid $ABCDEFGH$.
$AB = 8$ cm, $BC = 6$ cm and $CG = 5$ cm.



(a) Calculate the length of the diagonal AC.

..... cm **[2]**


(b) Calculate the length of the space diagonal AG.

..... cm **[2]**

(c) Calculate the angle between the line AG and the base $ABCD$.

..... ° **[2]**

1123 The diagram shows a pyramid $VABCD$ with a square base $ABCD$ of side 10 cm.
The point M is the centre of the base and V is vertically above M.
The vertical height VM of the pyramid is 12 cm.




(a) Calculate the length of the edge VA.

..... cm **[3]**

(b) Calculate the angle between the edge VA and the base $ABCD$.

..... ° **[2]**

1124 The diagram shows a right triangular prism $ABCDEF$.
The base ABC is a horizontal right-angled triangle with angle $ABC = 90°$.
$AB = 12$ cm, $BC = 5$ cm and the vertical height $AD = 8$ cm.



(a) Calculate the length of AF.

..... cm **[3]**


(b) Calculate the angle between the line AF and the base ABC.

..... ° **[2]**

(c) Calculate the angle between the line AF and the vertical plane $ABED$.

..... ° **[3]**


1125 The grid shows triangle A.



(a) On the grid, draw the image of triangle A after a translation by the vector $\begin{pmatrix} -4 \\ 2 \end{pmatrix}$.
Label the image B. **[2]**

(b) On the grid, draw the image of triangle A after a reflection in the line $x = -1$.
Label the image C. **[2]**


1126 The diagram shows a triangle, Shape A, drawn on a coordinate grid.



(a) On the grid, draw the image of Shape A after a translation by the vector $\begin{pmatrix} -4 \\ 2 \end{pmatrix}$.
Label this image B. **[2]**

(b) On the grid, draw the image of Shape A after a reflection in the line $y = -1$.
Label this image C. **[2]**

1127 The diagram shows two triangles, A and B, plotted on a coordinate grid.



(a) On the grid, draw the image of triangle A after a translation by the vector $\begin{pmatrix} -4 \\ 2 \end{pmatrix}$. **[2]**


(b) Describe fully the single transformation that maps triangle A onto triangle B.

.....

.....

..... **[2]**

1128 The grid shows triangle A and triangle B.




(a) On the grid, draw the image of triangle A after a translation by the vector $\begin{pmatrix} -2 \\ -5 \end{pmatrix}$. **[2]**

(b) Describe fully the single transformation that maps triangle A onto triangle B.

.....
.....
.....

[2]

1129 The diagram shows triangle A on a coordinate grid.




(a) On the grid, translate shape A by the vector $\begin{pmatrix} -5 \\ 2 \end{pmatrix}$.
Label the image B. **[2]**

(b) On the grid, reflect shape A in the x-axis.
Label the image C. **[2]**

(c) On the grid, rotate shape A by 180° about the origin $(0, 0)$.
Label the image D. **[2]**


1130 The diagram shows a shape P plotted on a coordinate grid.



(a) On the grid, draw the image of shape P after an enlargement, scale factor -2, centre $(0, 0)$. **[2]**

(b) On the grid, draw the image of shape P after a rotation of $90°$ clockwise about the point $(1, -1)$. **[2]**

1131 The diagram shows two triangles, Shape P and Shape Q, drawn on a grid.



(a) Describe fully the single transformation that maps Shape P onto Shape Q.


.....

.....

..... **[3]**

(b) Shape P is enlarged by a scale factor of $\frac{1}{2}$ with centre of enlargement $(-2, 2)$.
On the grid, draw the image of Shape P and label it R. **[2]**

1132 The diagram shows shape A on a coordinate grid.



(a) Shape A is reflected in the line $y = x$ to give shape B.
On the grid, draw and label shape B. **[2]**


(b) Shape B is rotated by 90° clockwise about the point $(2, 2)$ to give shape C.
On the grid, draw and label shape C. **[2]**

(c) Describe fully the single transformation that maps shape A onto shape C.

.....

..... **[2]**

1133 The grid shows triangle T.



(a) On the grid, draw the image of triangle T after an enlargement with scale factor $-\frac{1}{2}$ and centre $(0, 2)$. **[2]**

(b) Triangle T is reflected in the line $x = 1$ to give triangle U.
Triangle U is then reflected in the line $y = 2$ to give triangle V.
Describe fully the single transformation that maps triangle T onto triangle V.


.....
.....
.....

[3]

(c) Triangle T is enlarged by a scale factor of k to give triangle W.
The area of triangle W is 48 square units.
Find the value of k, given that $k > 0$.

$k =$ ..... **[2]**

1134 The diagram shows two triangles, Shape T and Shape U.



(a) Describe fully the single transformation that maps Shape T onto Shape U.

.....

.....

..... **[3]**

(b) Describe fully the single transformation that maps Shape U onto Shape T.


.....

..... **[2]**

(c) Shape T is translated by the vector $\begin{pmatrix} -3 \\ 1 \end{pmatrix}$ to give Shape V.
Shape V is then reflected in the line $y = x$ to give Shape W.
Find the coordinates of the three vertices of Shape W.

..... **[3]**

1135 The grid shows shapes P and Q.



(a) Describe fully the single transformation that maps shape P onto shape Q.


.....

.....

..... **[3]**

(b) On the grid, enlarge shape P by scale factor $\frac{1}{2}$, centre $(6, 6)$.
Label the image R. **[2]**

1136 The grid shows shapes P and Q.




(a) On the grid, draw the image of shape P after an enlargement with scale factor 0.5 and centre $(0, 0)$. **[2]**

(b) Describe fully the single transformation that maps shape P onto shape Q.

.....
.....
.....

[3]

1137 The diagram shows triangles P and Q on a coordinate grid.



(a) Describe fully the single transformation that maps shape P onto shape Q.

.....

.....

..... **[3]**

(b) On the grid, enlarge shape P by scale factor $\frac{1}{2}$ with centre $(0, 4)$.
Label the image R. **[3]**


(c) Describe fully the single transformation that maps shape R onto shape Q.

.....

.....

..... **[3]**

1138 The grid shows triangles T and U.



(a) Describe fully the single transformation that maps triangle T onto triangle U.

.....

.....

..... **[3]**


(b) Triangle T is reflected in the line $y = x$ to give triangle V.
Triangle V is then rotated 90° clockwise about the origin to give triangle W.
Describe fully the single transformation that maps triangle T onto triangle W.

.....

.....

..... **[2]**

1139 The diagram shows triangle T plotted on a coordinate grid.



(a) Triangle T is mapped onto triangle U by an enlargement, scale factor $\frac{1}{2}$, centre $(0, 4)$.
On the grid, draw triangle U. **[2]**

(b) Triangle U is mapped onto triangle V by a reflection in the line $y = x$.
On the grid, draw triangle V. **[2]**

(c) Describe fully the single transformation that maps triangle U back onto triangle T.

.....

.....

..... **[3]**

1140 Vectors $\mathbf{p}$ and $\mathbf{q}$ are given as

$$\mathbf{p} = \begin{pmatrix} 4 \\ -3 \end{pmatrix} \qquad \mathbf{q} = \begin{pmatrix} -2 \\ 5 \end{pmatrix}$$

(a) Work out

$$2\mathbf{p} + \mathbf{q}$$

$\begin{pmatrix} \quad \\ \quad \end{pmatrix}$ **[2]**

(b) Point A has coordinates $(1, 4)$.
The translation from A to B is given by the vector $\mathbf{p}$.
Work out the coordinates of B.

(..... , ...) **[1]**

1141 The vectors **a** and **b** are given by

$\mathrm{a} = \begin{pmatrix} 4 \\ -3 \end{pmatrix}$ and $\mathrm{b} = \begin{pmatrix} -2 \\ 5 \end{pmatrix}$

(a) Work out $\mathbf{a} + 2\mathbf{b}$ as a single column vector.

$\begin{pmatrix} \quad \\ \quad \end{pmatrix}$ **[2]**

(b) Point P has coordinates $(3, -1)$.
Point Q is translated from P by the vector **a**.
Find the coordinates of point Q.

(..... , ...) **[2]**

1142 Given the vectors $\mathbf{p} = \begin{pmatrix} 4 \\ -3 \end{pmatrix}$ and $\mathbf{q} = \begin{pmatrix} -2 \\ 5 \end{pmatrix}$.

(a) Work out $3\mathbf{p}$.

$\begin{pmatrix} \quad \\ \quad \end{pmatrix}$ **[2]**


(b) Work out $\mathbf{p} - 2\mathbf{q}$.

$\begin{pmatrix} \quad \\ \quad \end{pmatrix}$ **[2]**

(c) Point M is translated to point N by the vector $\mathbf{q}$.
Given that the coordinates of N are $(6, -1)$, find the coordinates of M.

(..... , ...) **[2]**

1143 The diagram shows a parallelogram $OABC$, where O is the origin.
$\vec{OA} = \mathbf{a}$ and $\vec{OC} = \mathbf{c}$.
M is the midpoint of AB and N is the point on BC such that $BN : NC = 2 : 1$.



(a) Find each of the following vectors in terms of **a** and **c**.
Give each answer in its simplest form.

(i) $\vec{OM}$

$\vec{OM} =$ ..... **[2]**


(ii) $\vec{MN}$

$\vec{MN} =$ ..... **[2]**

(b) $\vec{ON} = k\mathbf{a} + \mathbf{c}$.
Write down the value of k.

$k =$ ..... **[2]**

1144 The diagram shows vectors **a**, **b** and **c** plotted on a Cartesian coordinate grid.



(a) Write **c** as a column vector.

$\begin{pmatrix} \quad \\ \quad \end{pmatrix}$ **[1]**

(b) Vector **c** can be written in the form $p\mathbf{a} + q\mathbf{b}$, where p and q are integers.
Find the value of p and the value of q.

$p =$ ..... , $q =$ ..... **[4]**

1145 $\mathbf{p} = \begin{pmatrix} 4 \\ -3 \end{pmatrix}$ $\mathbf{q} = \begin{pmatrix} -2 \\ 5 \end{pmatrix}$


(a) Work out $3\mathbf{p} - 2\mathbf{q}$.

$\begin{pmatrix} \quad \\ \quad \end{pmatrix}$ **[2]**

(b) Point A has coordinates $(3, -1)$.
$\vec{AB} = \mathbf{p} + \mathbf{q}$.
Find the coordinates of B.

(..... , ...) **[2]**

1146 In triangle OAB, $\vec{OA} = \mathbf{a}$ and $\vec{OB} = \mathbf{b}$.
P is the point on AB such that $AP : PB = 3 : 1$.



(a) Find $\vec{OP}$ in terms of $\mathbf{a}$ and $\mathbf{b}$.
Give your answer in its simplest form.

$\vec{OP} =$ ..... **[3]**

(b) M is the midpoint of OA.
Find $\vec{MP}$ in terms of $\mathbf{a}$ and $\mathbf{b}$.
Give your answer in its simplest form.

$\vec{MP} =$ ..... **[2]**

1147 **(a)** $\mathbf{u} = \begin{pmatrix} 4 \\ -3 \end{pmatrix}$, $\mathbf{v} = \begin{pmatrix} -1 \\ y \end{pmatrix}$ and $\mathbf{w} = \begin{pmatrix} 10 \\ 6 \end{pmatrix}$.

Given that $3\mathbf{u} + k\mathbf{v} = \mathbf{w}$, work out the value of k and the value of y.

$k =$ ..... , $y =$ ..... **[3]**

(b) Points P, Q and R have coordinates $(2, -1)$, $(5, 5)$ and $(11, 17)$ respectively.

(i) Write $\vec{PQ}$ as a column vector.

$\begin{pmatrix} \quad \\ \quad \end{pmatrix}$ **[1]**

(ii) Show that P, Q and R lie on a straight line.

[2]

1148 Points A, B and C have coordinates $(1, -2)$, $(7, 10)$ and $(4, 4)$ respectively.

(a) Write $\vec{AB}$ as a column vector.

$\begin{pmatrix} \quad \\ \quad \end{pmatrix}$ **[2]**

(b) Show that $\vec{AC} = \frac{1}{2}\vec{AB}$.

[2]


(c) Point C is translated by the vector $\begin{pmatrix} -5 \\ 2 \end{pmatrix}$ to point D.
Find the coordinates of D.

(..... , ...) **[2]**

(d) Work out the vector $\vec{BD}$ as a column vector.

$\begin{pmatrix} \quad \\ \quad \end{pmatrix}$ **[2]**

1149 $OABC$ is a parallelogram.
$\vec{OA} = \mathbf{a}$ and $\vec{OC} = \mathbf{c}$.
X is a point on AB such that $AX : XB = 1 : 2$.
Y is a point on the line CB extended such that $CB : BY = 1 : 2$.



(a) Find $\vec{OX}$ in terms of **a** and **c**.

$\vec{OX} =$ ..... **[2]**

(b) Find $\vec{OY}$ in terms of **a** and **c**.

$\vec{OY} =$ ..... **[2]**

(c) Show that the points O, X and Y lie on a straight line.

[2]

(d) Find the ratio $OX : XY$.

$OX : XY =$ ..... **[1]**

1150 The position vectors of points A, B, and C relative to an origin O are given by

$= \begin{pmatrix} 2 \\ -1 \end{pmatrix}$, $= \begin{pmatrix} 6 \\ 7 \end{pmatrix}$, and $= \begin{pmatrix} k+1 \\ 3k-2 \end{pmatrix}$

where k is a constant.

(a) Find $\vec{AB}$ as a column vector.

$\vec{AB} = \begin{pmatrix} \quad \\ \quad \end{pmatrix}$ **[2]**

(b) Given that points A, B, and C are collinear, find the value of k.

$k =$ ..... **[3]**

(c) Using your value of k from part (b), write down $\vec{AC}$ in terms of $\vec{AB}$.

$\vec{AC} =$ ..... **[2]**

1151 Given the vectors $\mathbf{u} = \begin{pmatrix} 3 \\ -2 \end{pmatrix}$ and $\mathbf{v} = \begin{pmatrix} -1 \\ 4 \end{pmatrix}$.

(a) Point P has coordinates $(5, -1)$.
The vector $\vec{PQ} = 2\mathbf{u} - \mathbf{v}$.
Find the coordinates of Q.

(..... , ...) **[3]**

(b) Point R has coordinates $(14, 3)$.
The vector $\vec{PR}$ can be written in the form $h\mathbf{u} + k\mathbf{v}$.
Work out the values of the constants h and k.

$h =$ ..... , $k =$ ..... **[5]**


1152 Vector $\mathbf{v}$ is given as

$$\mathbf{v} = \begin{pmatrix} 5 \\ -12 \end{pmatrix}$$

Work out $|\mathbf{v}|$, the magnitude of vector $\mathbf{v}$.

..... **[2]**

1153 The diagram shows vectors **a** and **b** drawn on a square grid.



(a) Work out $|\mathbf{a}|$.

$|\mathbf{a}| =$ ..... **[2]**

(b) Work out $|2\mathbf{a} + 3\mathbf{b}|$.

$|2\mathbf{a} + 3\mathbf{b}| =$ ..... **[3]**

1154 Given the vectors $\mathbf{p} = \begin{pmatrix} -8 \\ 15 \end{pmatrix}$ and $\mathbf{q} = \begin{pmatrix} 2 \\ -4 \end{pmatrix}$.

(a) Calculate $|\mathbf{p}|$.

..... **[2]**

(b) Calculate $|\mathbf{p} + 2\mathbf{q}|$.

..... **[3]**

1155 **(a)** The vector $\mathbf{a} = \begin{pmatrix} p \\ p \end{pmatrix}$, where $p > 0$.

Given that $|\mathbf{a}| = \sqrt{18}$, find the value of p.

$p =$ ..... **[2]**

(b) The vectors $\mathbf{u}$ and $\mathbf{v}$ are given by

$$\mathbf{u} = \begin{pmatrix} 4 \\ 1 \end{pmatrix}, \qquad \mathbf{v} = \begin{pmatrix} -1 \\ 2 \end{pmatrix}$$

A vector $\mathbf{w}$ is defined as $\mathbf{w} = \mathbf{u} + t\mathbf{v}$.

Given that $|\mathbf{w}| = \sqrt{29}$, find the two possible values of t.

$t =$ ..... , or $t =$ ..... **[5]**

1156 Vector $\mathbf{u} = \begin{pmatrix} x \\ x+2 \end{pmatrix}$.

(a) The magnitude of vector $\mathbf{u}$ is 10.
Show that $x^2 + 2x - 48 = 0$.

[3]

(b) Solve the equation

$$x^2 + 2x - 48 = 0$$

$x =$ ..... , $x =$ ..... **[2]**

(c) Hence write down the two possible column vectors for $\mathbf{u}$.

[2]

1157 Points A, B, and C have coordinates $(3, -2)$, $(-5, y)$, and $(7, 1)$ respectively.

(a) Calculate $|\vec{AC}|$.

..... **[2]**

(b) The magnitude of vector $\vec{AB}$ is 17.
Work out the two possible values of y.

$y =$ ..... , $y =$ ..... **[5]**

1158 The points P, Q and R have coordinates $(2, -3)$, $(k, 5)$ and $(4, 1)$ respectively.

(a) Given that $|\vec{PQ}| = 10$, find the two possible values of k.

$k =$ ..... , or $k =$ ..... **[4]**

(b) Using the positive value of k from part (a), calculate the magnitude of $\vec{QR}$.
Give your answer correct to 3 significant figures.

$|\vec{QR}| =$ ..... **[3]**

1159 **(a)** The points A and B have coordinates $(2, -3)$ and $(9, 4)$ respectively.
Work out the column vector $\vec{AB}$.

$\begin{pmatrix} \quad \\ \quad \end{pmatrix}$ **[2]**

(b) Calculate $|\vec{AB}|$.
Give your answer correct to 3 significant figures.

..... **[2]**

(c) Vector $\mathbf{p} = \begin{pmatrix} k \\ 4 \end{pmatrix}$.
Given that $|\mathbf{p}| = 5$, find the two possible values of k.

$k =$ ..... , $k =$ ..... **[3]**

1160 Given the vectors $\mathbf{u} = \begin{pmatrix} 2k \\ k-3 \end{pmatrix}$ and $\mathbf{v} = \begin{pmatrix} k+1 \\ 2 \end{pmatrix}$.

(a) Given that $|\mathbf{u} + \mathbf{v}| = \sqrt{50}$, show that $5k^2 + 2k - 24 = 0$.

[4]

(b) Solve the equation

$$5k^2 + 2k - 24 = 0$$

$k =$ ..... , $k =$ ..... **[4]**

(c) Hence, find the two possible column vectors for $\mathbf{u}$.

$\mathbf{u} = \begin{pmatrix} \quad \\ \quad \end{pmatrix}$ ..... , $\mathbf{u} = \begin{pmatrix} \quad \\ \quad \end{pmatrix}$ ..... **[2]**

1161 Event A is that a train arrives on time.

$P(A) = 0.83$

(a) Work out $P(A')$.

..... **[1]**

(b) State in words what the event A' represents.

..... **[1]**

1162 A biased spinner can land on 1, 2, 3, 4, or 5.
The table shows the probabilities of the spinner landing on each of the numbers.

Outcome	1	2	3	4	5
Probability	0.15	$2x$	0.31	x	0.24

(a) Find the value of x.

$x =$ ..... **[2]**

(b) The spinner is spun once.
Find the probability that it does not land on 2.

..... **[2]**

1163 A biased spinner can land on Red, Blue, Green, or Yellow.
The table shows the probabilities of the spinner landing on Red, Blue, and Green.

Colour	Red	Blue	Green	Yellow
Probability	0.15	0.3	0.45	

(a) Work out the probability that the spinner lands on Yellow.

..... **[2]**

(b) Work out the probability that the spinner lands on Red or Blue.

..... **[1]**

(c) Let G be the event that the spinner lands on Green.
Find $P(G')$.

$P(G') =$ ..... **[2]**

1164 A biased spinner can land on 1, 2, 3 or 4.
The table shows the probabilities of the spinner landing on each number.

Number	1	2	3	4
Probability	$\frac{1}{5}$	0.35	x	$2x$

(a) Work out the value of x.

$x =$ ..... **[2]**


(b) Find the probability that the spinner does not land on 4.

..... **[2]**

1165 In a group of 60 people, some like tea (T) and some like coffee (C).

- The total number of people who like tea is 38.
- The total number of people who like coffee is 24.
- The number of people who like neither is x.
- The number of people who like both is $2x$.

(a) Use this information to complete the Venn diagram in terms of x.



[3]

(b) Find the value of x.

$x =$ ..... **[2]**

(c) A person is chosen at random from the group.
Find $P(T \cap C')$.

$P(T \cap C') =$ ..... **[2]**

1166 A bag contains only red counters, blue counters, and green counters.
The number of red counters is x^2.
The number of blue counters is $3x$.
The number of green counters is 2.

(a) Write down an expression, in its simplest form, for the total number of counters in the bag.

..... **[1]**

(b) A counter is chosen at random.
The probability that it is red is $\frac{8}{15}$.
Show that $7x^2 - 24x - 16 = 0$.

[2]

(c) Solve the equation $7x^2 - 24x - 16 = 0$ to find the value of x.

$x =$ ..... **[3]**

(d) Find the probability that a randomly chosen counter is not blue.
Give your answer as a fraction in its simplest form.

..... **[2]**

1167 A bag contains only red marbles, blue marbles, and green marbles.
The probability of choosing a red marble at random is $\frac{2}{5}$.
The probability of choosing a blue marble at random is $\frac{1}{3}$.
There are 12 green marbles in the bag.

(a) Calculate the probability of choosing a green marble at random.

..... **[2]**

(b) Calculate the total number of marbles in the bag.


..... **[3]**

(c) Write down the probability that a marble chosen at random is not blue.

..... **[1]**

1168 A group of 120 people were asked about the pets they own.
$C = \{\text{people who own a cat}\}$
$D = \{\text{people who own a dog}\}$
$n(\mathsf{U}) = 120$
$n(C) = 55$
$n(D) = 68$
18 people own neither a cat nor a dog.

(a) Complete the Venn diagram to show the number of people in each region.



[3]

(b) A person is chosen at random.
Find $P(C \cap D)$.

..... **[1]**


(c) Find $P(D')$.

..... **[2]**

(d) Find the probability that a person chosen at random owns a cat or a dog, but not both.

..... **[2]**

1169 The Venn diagram shows information about the number of students in a class of 50 who study French (F) and German (G).



(a) Work out the value of x.

$x =$ ..... **[2]**

(b) A student is chosen at random.
Find $P(F')$.

..... **[2]**

(c) A student is chosen at random.
Find the probability that this student studies exactly one of these languages.

..... **[2]**

1170 A biased five-sided spinner is numbered 1, 2, 3, 4, and 5.
The table shows the probability of the spinner landing on each number.

Number	1	2	3	4	5
Probability	0.15	0.25	x	0.2	$3x$

(a) Work out the value of x.

$x =$ ..... **[2]**

(b) The spinner is spun 250 times.
Calculate the expected number of times the spinner lands on a number greater than 3.

..... **[2]**

1171 A biased 4-sided spinner, numbered 1, 2, 3, and 4, is spun 200 times.
The table shows the frequencies of each outcome.

Number	1	2	3	4
Frequency	45	35	70	50

(a) Write down the relative frequency of the spinner landing on 3.

..... **[1]**

(b) The spinner is spun 1000 times.
Estimate the number of times it will land on 1.

..... **[2]**

(c) Is the spinner fair?
Give a reason for your answer.

..... **[1]**

(d) Explain why spinning the spinner 1000 times gives a more reliable estimate of the probability than spinning it 200 times.

..... **[1]**

1172 A biased four-sided die is rolled 60 times.
The scores are 1, 2, 3, and 4.
The frequencies of each score are shown in the table.

Score	1	2	3	4
Frequency	x	$2x + 5$	$3x - 2$	$x + 8$

(a) Find the value of x.

$x =$ ..... **[2]**

(b) Find the relative frequency of rolling a 3.

..... **[1]**

(c) The die is rolled 1200 times.
Calculate the expected number of times it rolls a 2.

..... **[2]**

1173 A spinner has four colours: Red, Blue, Green, and Yellow.
A student spins it 50 times and records the results.
The table shows the relative frequencies for three of the colours.

Colour	Red	Blue	Green	Yellow
Relative frequency	0.15	0.45	0.10	x

(a) Find the value of x.

$x =$ ..... **[1]**

(b) The spinner is spun 300 times in total.
Estimate the number of times it lands on Blue.

..... **[2]**

(c) The spinner is meant to be fair.
Comment on whether the results show it is fair or biased.

..... **[1]**

1174 A bag contains a large number of coloured beads.
A bead is chosen at random, its colour recorded, and then returned to the bag.
This experiment is repeated 300 times.
The table shows the frequencies of the colours drawn.

Colour	Red	Blue	Green	Yellow
Frequency	84	105	x	y

(a) The relative frequency of choosing a Green bead is 0.22.
Work out the value of x and the value of y.

$x =$ ..... , $y =$ ..... **[3]**

(b) There are exactly 15 Yellow beads in the bag.
Estimate the total number of beads in the bag.

..... **[2]**

(c) A new experiment is done where a bead is chosen and replaced 800 times.
Calculate the expected number of times a Blue or Green bead is chosen.

..... **[2]**

1175 A researcher records the colours of cars passing a checkpoint.
Over one week, 800 cars pass the checkpoint.
The frequencies of the colours are shown in the table.

Colour	Silver	Black	White	Other
Frequency	260	180	w	$w + 40$

(a) Find the relative frequency of a car being Silver.
Give your answer as a fraction in its simplest form.

..... **[2]**

(b) Find the value of w.

$w =$ ..... **[2]**

(c) The researcher uses these results to estimate the probability that the next car passing the checkpoint is White.
Work out this estimate.

..... **[1]**

(d) In one month, 12000 cars are expected to pass the checkpoint.
Calculate the expected number of Black cars.

..... **[2]**

1176 A biased 5-sided spinner has sides numbered 1, 2, 3, 4, and 5.
The table shows the probability of the spinner landing on each number in terms of x.

Number	1	2	3	4	5
Probability	0.15	$2x$	$x + 0.05$	0.2	x

(a) Find the value of x.

$x =$ ..... **[2]**

(b) The spinner is spun 600 times.
Work out the expected number of times it lands on a prime number.

..... **[2]**

(c) A different spinner has 3 sections: Red, Blue, and Green.
The probability of landing on Red is p.
The probability of landing on Blue is $2p$.
The probability of landing on Green is $1 - 3p$.
This spinner is spun N times.

The expected number of times it lands on Red is 54 and the expected number of times it lands on Green is 36.
Work out the value of N and the value of p.

$N =$ ..... , $p =$ ..... **[4]**

1177 A biased four-sided spinner is numbered 1, 2, 3 and 4.
The table shows the probability of scoring each number.
The spinner is spun 400 times.
Let E_1 be the expected number of times the spinner lands on 1.
Let E_3 be the expected number of times the spinner lands on 3.
You are given that $E_1 \times E_3 = 5600$.
Also, the probability of scoring a 3 is greater than the probability of scoring a 1.

Score	1	2	3	4
Probability	x	$2x$	y	y

(a) Show that $300x^2 - 100x + 7 = 0$.

[3]

(b) Solve the equation to find the value of x and the value of y.

$x =$ ..... , $y =$ ..... **[4]**

(c) Calculate the expected number of times the spinner lands on an even number.

..... **[2]**

1178 Two biased coins, Coin A and Coin B, are tossed.
The probability of Coin A landing on Heads is p.
The probability of Coin B landing on Heads is $p + 0.15$.
Coin A is tossed 300 times and Coin B is tossed 200 times.
The total expected number of Heads from these 500 tosses is 255.

(a) Form an equation in p and show that $p = 0.45$.

[3]

(b) Both coins are tossed together 1000 times.
Work out the expected number of times they land on different outcomes.

..... **[3]**

(c) In a new experiment, Coin A is tossed N times.
The relative frequency of landing on Heads is 0.48 and it lands on Heads 72 times.

(i) Work out the value of N.


$N =$ ..... **[2]**

(ii) Calculate the expected number of Tails in these N tosses, using the theoretical probability $p = 0.45$.

..... **[2]**

1179 A bag contains 5 red counters and 3 blue counters.
A counter is taken at random from the bag, its colour is noted and it is replaced.
A second counter is then taken at random.

(a) Complete the tree diagram.




[2]

(b) Calculate the probability that both counters are red.

..... **[2]**

1180 A bag contains 4 red discs and 5 blue discs.
Jamil takes one disc at random, notes its colour, and does not replace it.
He then takes a second disc at random.

(a) Complete the tree diagram.




[2]

(b) Calculate the probability that Jamil takes one red disc and one blue disc.

..... **[3]**

1181 A basket contains 7 green apples and 3 red apples.
Two apples are chosen at random from the basket, without replacement.

(a) Complete the tree diagram.




[2]

(b) Calculate the probability that both apples chosen are the same colour.

..... **[3]**

1182 A class of 30 students were surveyed about the subjects they study.
17 study French (F), 14 study Spanish (S), and 5 study neither subject.

(a) Complete the Venn diagram to represent this information.



[2]

(b) A student is chosen at random from the class.
Work out the probability that this student:

(i) studies both French and Spanish,

..... **[1]**

(ii) studies French but not Spanish,


..... **[1]**

(iii) is a member of the set $F \cup S$.

..... **[1]**

1183 In a group of 30 students, 18 study Biology (B), 15 study Chemistry (C), and 4 study neither Biology nor Chemistry.

(a) Complete the Venn diagram to show the number of students in each region.



[2]

(b) A student is chosen at random from the group.

(i) Find the probability that this student studies Chemistry but not Biology.

..... **[1]**

(ii) Find $\mathrm{P}(B \cup C)$.


..... **[1]**

(iii) Given that the student chosen studies Biology, find the probability that this student also studies Chemistry.

..... **[1]**

1184 A bag contains 8 balls.
5 of the balls are Red and 3 of the balls are Blue.
A ball is taken at random from the bag, its colour is noted, and it is replaced.
A second ball is then taken at random from the bag.

(a) Complete the tree diagram.



[2]

(b) Calculate the probability that both balls are Red.


..... **[1]**

(c) Calculate the probability of picking one Red ball and one Blue ball in any order.

..... **[2]**

1185 A bag contains 5 red counters and 3 blue counters.
A counter is picked at random and its colour is noted.
It is then replaced in the bag.
A second counter is then picked at random.

(a) Complete the tree diagram.



[2]

(b) Work out the probability that both counters are red.


..... **[1]**

(c) Work out the probability that one counter is red and one counter is blue.

..... **[2]**

1186 A bag contains 7 red apples and 3 green apples.
Liam takes two apples at random, without replacement.

(a) Complete the tree diagram.




[2]

(b) Work out the probability that Liam takes two apples of the same colour.

..... **[3]**

1187 In a group of 30 students, 18 study Biology (B) and 15 study Chemistry (C).
5 students study neither Biology nor Chemistry.

(a) Complete the Venn diagram to show the number of students in each region.



[2]

(b) A student is chosen at random from the group.

(i) Find the probability that this student studies Biology but not Chemistry.

..... **[1]**

(ii) Find $\mathrm{P}(B \cup C)$.

..... **[1]**


(c) Find the probability that a student chosen at random from those who study Chemistry also studies Biology.

..... **[2]**

1188 There are 40 students in a group.

- 24 students play Football (F).
- 15 students play Cricket (C).
- 8 students play both Football and Cricket.

(a) Complete the Venn diagram to show this information.



[2]

(b) A student is chosen at random.
Calculate the probability that the student plays neither Football nor Cricket.


..... **[2]**

(c) Find $n(F \cup C)$.

..... **[2]**

1189 A bag contains 7 red counters and 5 blue counters.
A counter is taken at random, its colour is noted, and it is NOT replaced in the bag.
A second counter is then taken at random and its colour is noted.

(a) Complete the tree diagram by writing the missing probabilities on the branches.



[3]

(b) Calculate the probability that both counters are red.


..... **[2]**

(c) Calculate the probability that one counter is red and one counter is blue.

..... **[2]**

1190 A bag contains 8 balls.
There are 5 Red balls and 3 Blue balls.
A ball is taken at random from the bag, its colour is recorded, and it is replaced.
A second ball is then taken at random from the bag.

(a) Complete the tree diagram.



[2]

(b) Calculate the probability that both balls are Red.

..... **[2]**

(c) Calculate the probability that one ball is Red and one ball is Blue.

..... **[3]**

1191 A box contains x blue counters and 4 green counters.
Two counters are chosen at random from the box, one after another, without replacement.

(a) Show that the probability that both counters chosen are blue is $\dfrac{x(x-1)}{x^2+7x+12}$.

[2]

(b) The probability that both counters chosen are blue is $\dfrac{1}{3}$.

(i) Show that $x^2 - 5x - 6 = 0$.

[3]

(ii) Solve the equation $x^2 - 5x - 6 = 0$ to find the number of blue counters in the box.

$x =$ ..... **[3]**

1192 A bag contains x red counters and 5 blue counters.
Two counters are taken at random from the bag without replacement.
The probability that both counters are blue is $\frac{2}{15}$.

(a) Show that $x^2 + 9x - 130 = 0$.

[3]

(b) Find the value of x.

$x =$ ..... **[2]**

(c) Calculate the probability that the two counters chosen are different colours.

..... **[3]**

1193 A box contains x red balls and 4 blue balls.
Two balls are taken from the box at random, without replacement.
The probability that both balls are red is $\frac{1}{3}$.

(a) Show that $x^2 - 5x - 6 = 0$.

[3]

(b) Work out the number of red balls in the box.

..... **[2]**

(c) Calculate the probability that the two balls taken have different colours.

..... **[3]**

1194 A bag contains n counters.
3 of the counters are red and the rest are blue.
Two counters are chosen at random, without replacement.
The probability that both counters are red is $\frac{1}{15}$.

(a) Show that $n^2 - n - 90 = 0$.

[3]

(b) Find the number of blue counters in the bag.

..... **[2]**

(c) Find the probability that the two counters chosen have different colours.

..... **[3]**

1195 A bag contains n balls.
3 of the balls are red and the rest are blue.
Two balls are chosen at random, without replacement.
The probability that both balls are red is $\frac{1}{15}$.

(a) Show that $n^2 - n - 90 = 0$.

[3]

(b) Find the value of n.

$n =$ ..... **[2]**

(c) Find the probability that the two balls have different colours.

..... **[3]**

1196 A box contains n counters.
3 of the counters are Red and all the remaining counters are Blue.
Two counters are taken at random from the box, without replacement.
The probability that both counters are Red is $\frac{1}{15}$.

(a) Show that $n^2 - n - 90 = 0$.

[3]

(b) Find the value of n.


$n =$ ..... **[2]**

(c) Calculate the probability that exactly one of the two counters chosen is Red.

..... **[3]**

1197 A biased coin has a probability of showing Heads of $\frac{3}{5}$.
The coin is flipped twice.

(a) Complete the tree diagram.



[2]


(b) Calculate the probability that the coin shows exactly one Head.

..... **[2]**

1198 In a survey, 40 students are asked if they study History (H) or Geography (G).

- 23 students study History.
- 19 students study Geography.
- 7 students study neither subject.

(a) Complete the Venn diagram to show this information.



[2]

(b) A student is chosen at random from the 40 students.
Find the probability that the student studies History but not Geography.

..... **[1]**


(c) Two students are chosen at random from those who study History.
Calculate the probability that they both also study Geography.

..... **[2]**

1199 There are 30 students in a class.
U = {students in the class}
F = {students who play Football}
T = {students who play Tennis

18 students play Football. 14 students play Tennis. 4 students do not play Football and do not play Tennis.

(a) Complete the Venn diagram to show the number of students in each region.



[2]


(b) A student is chosen at random from the class.
Find $P(F \cap T)$.

$P(F \cap T) =$ ..... **[1]**

(c) Two students are chosen at random from the class.
Calculate the probability that both students play Football but not Tennis.

..... **[3]**

1200 A group of 40 students are asked if they play Football (F) or Tennis (T).
The Venn diagram shows the number of students in each region in terms of x.



(a) Find the value of x.

$x =$ ..... **[2]**

(b) Find $P(F \cup T)$.

..... **[1]**


(c) Two students are chosen at random from the group of 40.
Calculate the probability that both students play Football only.

..... **[3]**

1201 In a group of 40 students:

- 24 play Football (F)
- 19 play Tennis (T)
- 7 do not play either sport.

(a) Complete the Venn diagram to show this information.



[2]

(b) A student is chosen at random from the group.
Find the probability that the student plays both Football and Tennis.


..... **[1]**

(c) Two students who play Football are chosen at random.
Find the probability that exactly one of them also plays Tennis.

..... **[3]**

1202 In a group of 50 students, 32 study History (H), 21 study Geography (G), and 6 study neither subject.

(a) Complete the Venn diagram to show the number of students in each region.



[3]

(b) A student is chosen at random from the group of 50.

Find:

(i) $P(H \cap G)$

$P(H \cap G) =$ ..... **[1]**

(ii) $P(H \cup G')$


$P(H \cup G') =$ ..... **[1]**

(c) Two students who study History are chosen at random.
Calculate the probability that both students also study Geography.

..... **[2]**

1203 In a group of 40 students, 24 study Geography (G), 18 study History (H), and 6 study neither subject.

(a) Complete the Venn diagram to show the number of students in each region.



[2]

(b) A student is chosen at random from the group of 40 students.

(i) Find the probability that the student studies History but not Geography.

..... **[1]**

(ii) Find $P(G \cup H')$.


..... **[2]**

(c) Two students are chosen at random without replacement from the group of 40 students. Work out the probability that both students study Geography.

..... **[2]**

1204 In a group of 40 students, 24 play tennis (T) and 18 play golf (G).
5 students play neither tennis nor golf.

(a) Complete the Venn diagram to show this information.



[2]

(b) Two students are chosen at random from the group of 40.
Calculate the probability that both students play tennis but not golf.


. . . . . **[3]**

(c) A student who plays golf is chosen at random.
Find the probability that they also play tennis.

. . . . . **[2]**

1205 A box contains 7 red counters and 3 blue counters.
A counter is chosen at random from the box and its colour is recorded.
It is not replaced.
A second counter is then chosen at random from the box.

(a) Complete the tree diagram.



[3]

(b) Calculate the probability that both counters chosen are red.


..... **[2]**

(c) Calculate the probability that the two counters chosen are of different colours.

..... **[2]**

1206 A bag contains 5 red sweets and 4 yellow sweets.
Maya takes a sweet at random from the bag and eats it.
She then takes a second sweet at random from the bag and eats it.

(a) Complete the tree diagram.



[2]

(b) Calculate the probability that Maya eats two sweets of the same colour.


..... **[2]**

(c) Calculate the probability that Maya eats at least one yellow sweet.

..... **[3]**

1207 A bag contains 5 red counters and 3 blue counters.
Two counters are taken from the bag at random, without replacement.

(a) Complete the tree diagram by writing the missing probabilities on the branches.



[3]

(b) Calculate the probability that both counters taken are the same colour.


..... **[2]**

(c) Calculate the probability that at least one of the counters taken is red.

..... **[2]**

1208 In a group of 40 students, 24 like football (F) and 18 like cricket (C).
x students like both sports.
3 students do not like either sport.

(a) Complete the Venn diagram to show this information in terms of x.



[2]

(b) Work out the value of x.

$x =$ ..... **[2]**

(c) Two students are chosen at random from the group of 40.
Work out the probability that one student likes football only and the other student likes cricket only.


..... **[3]**

1209 In a class of 30 students, 18 study French (F) and 14 study Spanish (S).
3 students study neither French nor Spanish.
Let x be the number of students who study both French and Spanish.

(a) Find the value of x.

$x =$ ..... **[2]**

(b) Complete the Venn diagram to show the number of students in each region.




[2]

(c) Two students are chosen at random from the class.
Calculate the probability that both students study French but do not study Spanish.

..... **[3]**

1210 A box contains 7 red counters and 3 blue counters.
Two counters are taken at random from the box, without replacement.

(a) Complete the tree diagram.



[3]

(b) Calculate the probability that both counters are red.

..... **[2]**

(c) Calculate the probability that the two counters are different colours.

..... **[2]**

1211 A bag contains n sweets.
5 of the sweets are orange and the rest are lemon.
Two sweets are taken from the bag at random, without replacement.
The probability that both sweets are orange is $\frac{2}{21}$.

(a) Show that $n^2 - n - 210 = 0$.

[3]

(b) Solve the equation $n^2 - n - 210 = 0$ to find the number of sweets in the bag.

$n =$ ..... **[2]**

(c) Find the probability that the two sweets taken have different flavours.

..... **[3]**

1212 Box A contains x blue counters and 3 red counters.
Box B contains 4 blue counters and 2 red counters.
A counter is taken at random from Box A and placed into Box B.
A counter is then taken at random from Box B.

(a) Show that the probability of taking a blue counter from Box B is

$$\frac{5x+12}{7(x+3)}$$

[4]

(b) The probability of taking a blue counter from Box B is $\frac{3}{4}$.

(i) Work out the value of x.

$x =$ ..... **[3]**

(ii) Using your value of x, find the probability that the counter transferred from Box A to Box B was red, given that the counter taken from Box B is blue.

..... **[1]**

1213 A bag contains 4 red discs and some blue discs.
Let the total number of discs in the bag be x.
Two discs are taken at random from the bag, without replacement.
The probability that both discs are red is $\frac{1}{11}$.

(a) Show that $x^2 - x - 132 = 0$.

[3]

(b) Find the number of blue discs in the bag.

..... **[3]**

(c) Calculate the probability that one disc is red and one disc is blue.

..... **[3]**

1214 A box contains n chocolates.
There are 4 milk chocolates and the rest are dark chocolates.
Two chocolates are taken at random from the box, without replacement.
The probability that both chocolates are milk chocolate is $\frac{1}{14}$.

(a) Show that $n^2 - n - 168 = 0$.

[3]

(b) Solve the equation to find the number of dark chocolates in the box.

..... **[3]**

(c) Calculate the probability that the two chocolates taken are of different types.

..... **[3]**

1215 A box contains n sweets. 6 of the sweets are lemon flavour and the rest are orange flavour.

(a) Hannah takes two sweets at random from the box without replacement.
The probability that both sweets are lemon flavour is $\frac{1}{3}$.

(i) Show that $n^2 - n - 90 = 0$.

[3]

(ii) Solve the equation $n^2 - n - 90 = 0$ to find the total number of sweets in the box.

$n =$ ..... **[3]**

(b) Three sweets are now taken at random without replacement from the original box of 10 sweets.
Calculate the probability that at least one of the sweets taken is orange flavour.

..... **[3]**

1216 A box contains n chocolates, of which 4 are dark chocolates and the rest are milk chocolates.

(a) Two chocolates are taken from the box at random, without replacement.
The probability that both chocolates are dark is $\frac{2}{15}$.
Show that $n^2 - n - 90 = 0$.

[3]

(b) Solve the equation $n^2 - n - 90 = 0$ to find the total number of chocolates in the box.

$n =$ ..... **[2]**

(c) Three chocolates are now taken at random from the original box of chocolates, without replacement.
Work out the probability that exactly two of the three chocolates taken are dark chocolates.

..... **[4]**

1217 A bag contains x green apples and 4 red apples.
Two apples are taken at random, without replacement.

(a) The probability that both apples are green is $\frac{5}{18}$.

Show that $13x^2 - 53x - 60 = 0$.

[4]

(b) Solve the equation $13x^2 - 53x - 60 = 0$ to find the value of x.

$x =$ ..... **[3]**

(c) Work out the probability that exactly one of the two apples taken is green.

..... **[3]**

1218 70 students are asked to choose their favourite sport from Football, Tennis, or Basketball.
The incomplete two-way table shows some information about their choices.

(a) Complete the two-way table.

	Football	**Tennis**	**Basketball**	**Total**
Boys	15		12	
Girls		14		30
Total	23		20	70

[3]

(b) A student is chosen at random.
Find the probability that the student is a boy whose favourite sport is Tennis.

..... **[1]**

1219 50 students each choose one sport to play: Tennis, Football, or Swimming.
The students are classified as Boys or Girls.

(a) Complete the two-way table to show this information.

	Tennis	**Football**	**Swimming**	**Total**
Boys	7			28
Girls	8		6	
Total		20		50

[3]

(b) A student is chosen at random.
Find the probability that the student is a boy who chose Football.

..... **[1]**

1220 50 students were asked to choose their favourite sport from Football, Basketball, and Tennis.
The two-way table shows some information about their choices.

(a) Complete the two-way table.

	Football	**Basketball**	**Tennis**	**Total**
Boys	15		5	28
Girls		10	6	
Total	21			50

[3]

(b) A student who chose Tennis is chosen at random.
Work out the probability that the student is a boy.

..... **[2]**

1221 80 students were asked to choose their favourite sport from Football, Basketball, and Tennis.

- There are 35 girls in total.
- 38 students chose Football, of which 24 were boys.
- 12 girls chose Basketball.
- 20 students chose Tennis.

(a) Complete the two-way table.

	Football	**Basketball**	**Tennis**	**Total**
Boys				
Girls				
Total				

[3]

(b) A student is chosen at random.
Find the probability that the student is a boy whose favourite sport is Basketball.

..... **[1]**

(c) A girl is chosen at random.
Find the probability that her favourite sport is Tennis.

..... **[2]**

1222 A survey of adults recorded their preferred hot drink, either Tea or Coffee.
Let x be the number of males who prefer Coffee.
The number of males who prefer Tea is $2x + 10$.
The number of females who prefer Coffee is $x + 5$.
The number of females who prefer Tea is $3x - 11$.

(a) Write down an expression, in terms of x, for the total number of adults.
Give your answer in its simplest form.

..... **[2]**

(b) The survey included 130 adults.
Find the value of x.

$x =$ ..... **[2]**

(c) Complete the two-way table.

	Coffee	**Tea**	**Total**
Male			
Female			
Total			

[3]

(d) One of the adults who prefers Coffee is chosen at random.
Find the probability that this adult is female.

..... **[2]**

1223 A group of 120 adults were asked whether they prefer Tea or Coffee.

- 55 of the adults are women.
- 70 of the adults prefer Tea.
- 25 of the men prefer Coffee.

(a) Complete the two-way table to show this information.

	Tea	**Coffee**	**Total**
Men			
Women			
Total			

[3]

(b) Work out the number of women who prefer Tea as a percentage of the total number of adults.

..... % **[2]**

1224 A company has 120 employees. There are 75 men and 45 women.
They travel to work by bus, train, or car.

- 20% of the men travel by bus.
- $\frac{1}{3}$ of the women travel by bus.
- A total of 38 employees travel by train, of which 14 are women.
- The remaining employees travel by car.

(a) Construct and complete the two-way table to show this information.

	Bus	**Train**	**Car**	**Total**
Men				
Women				
Total				

[4]

(b) An employee who travels by car is chosen at random.
Work out the probability that this employee is a man.

..... **[2]**

1225 In a school of 400 students, the ratio of juniors to seniors is 3 : 2.
Each student travels to school by Bus, by Car, or Walks.

- 45% of the 400 students take the Bus.
- Of the seniors, 30 take the Car and 50 Walk.
- There are 80 juniors who take the Car.

(a) Complete the two-way table to classify this data.

	Bus	**Car**	**Walk**	**Total**
Juniors				
Seniors				
Total				400

[3]

(b) A student is chosen at random.
Find the probability that they are a junior who walks.

..... **[1]**

(c) Two seniors are chosen at random.
Find the probability that they both take the bus.

..... **[2]**

1226 80 students travel to school either by Bus or by Walking.
The two-way table shows the number of boys and girls using each method of transport, in terms of x.

	Bus	**Walk**	**Total**
Boys	x	$2x$	
Girls	$x + 5$	$3x - 2$	
Total			80

(a) Work out the value of x.

$x =$ ..... **[3]**

(b) Complete the two-way table with the number of students in each category.

	Bus	**Walk**	**Total**
Boys			
Girls			
Total			80

[2]

(c) A student is chosen at random from those who walk to school.
Find the probability that this student is a girl.

..... **[2]**

1227 A company has 300 employees. Each employee is either a Manager, an Office Worker, or a Technician.
Each employee works either Full-Time or Part-Time.

- The ratio of Managers : Office Workers : Technicians is 2 : 5 : 3.
- 20% of the Managers work Part-Time.
- 40% of the Office Workers work Part-Time.
- The number of Part-Time Technicians is equal to the number of Part-Time Managers.

(a) Complete the two-way table for this information.

	Full-Time	**Part-Time**	**Total**
Managers			
Office Workers			
Technicians			
Total			

[4]

(b) A Part-Time employee is chosen at random.
Find the probability that they are an Office Worker.

..... **[2]**

(c) Two Managers are chosen at random.
Find the probability that exactly one of them works Part-Time.

..... **[3]**

1228 A sports club has 200 members. Each member plays either Golf or Tennis.
There are x male members in the club.

- The number of male members who play Golf is $\frac{1}{4}x$.
- The number of female members who play Tennis is 30 more than the number of male members who play Golf.
- A total of 110 members play Tennis.

(a) Write down an expression, in terms of x, for the number of male members who play Tennis.
Give your answer in its simplest form.

..... **[1]**

(b) Use the information about the members who play Tennis to form an equation in x, and show that $x = 80$.

[2]

(c) Complete the two-way table to show the number of members in each category.

	Golf	**Tennis**	**Total**
Male			
Female			
Total			

[3]

(d) Two members are chosen at random from those who play Golf.
Calculate the probability that they are both female.

..... **[3]**

1229 A group of 120 students are classified by gender (Boys, Girls) and handedness (Left-handed, Right-handed).
The two-way table shows the number of students in each category in terms of x and y.
Two students are chosen at random from the 120 students.
The probability that both students are left-handed is $\frac{3}{34}$.

	Left-handed	**Right-handed**
Boys	x	y
Girls	$2x$	$y - 6$

(a) Show that $3x^2 - x - 420 = 0$.

[4]

(b) Find the value of x and the value of y.

$x =$ ..... , $y =$ ..... **[4]**

(c) A right-handed student is chosen at random.
Find the probability that the student is a girl.

..... **[2]**

1230 The table shows information about the marks scored by two different classes, Class A and Class B, in a mathematics examination.

Statistic	Class A	Class B
Mean	62	70
Median	65	72
Range	40	25
Interquartile Range	15	10

(a) Which class had the higher average mark?
Give a reason for your answer.

..... **[1]**

(b) Which class had marks that were more consistent?
Give a reason for your answer.

..... **[1]**

(c) A student says, "Every student in Class B scored a higher mark than every student in Class A."
Explain why this statement cannot be confirmed using only the information in the table.

.....

.....

..... **[2]**

1231 The number of goals scored by Team A in each of their last 6 matches is recorded below.

3 1 0 4 2 2

Team B also played 6 matches.
For Team B, the median number of goals scored was 1.5 and the range was 5.

(a) Work out the median and the range of the number of goals scored by Team A.

Median = ... , Range = ... **[2]**


(b) Make two comments comparing the performance of Team A with the performance of Team B.

.....

.....

..... **[2]**

1232 The dual bar chart shows the grades achieved by a group of boys and a group of girls in an examination.



(a) Write down the modal grade for the boys.

..... **[1]**

(b) Find the number of girls who scored Grade C.

..... **[1]**

(c) Calculate the percentage of all students who scored Grade A.

..... % **[2]**

1233 The stem-and-leaf diagram shows the ages, in years, of 15 members of a chess club.

1	2 5 8
2	0 1 1 4 7 9
3	2 5 6 9
4	1 3

Key: 2 | 1 represents 21 years

(a) Find the median age.

..... years **[1]**

(b) Find the range of the ages.

..... years **[1]**

(c) Find the interquartile range.

..... years **[2]**

(d) A member is chosen at random.
Find the probability that they are older than 30 years.

..... **[1]**

1234 A set of 5 integers has the following statistical properties:

- Mode = 3
- Median = 7
- Mean = 8
- Range = 11

(a) Find the 5 integers.

..... **[4]**

(b) A sixth integer, x, is added to the set.
The new mean is 9.
Work out the value of x.

$x =$ ..... **[2]**


(c) A student claims: "If another completely different set of 5 integers has the same range of 11, it must also have the same standard deviation."

State whether the student is correct.
Give a statistical reason for your answer.

.....

..... **[1]**

1235 The box-and-whisker plots show information about the times, in minutes, taken by two groups of people (Group X and Group Y) to complete a puzzle.



(a) Find the median time for Group Y.

..... minutes **[1]**

(b) Find the interquartile range for Group X.

..... minutes **[1]**

(c) Make two different comments comparing the distributions of the times taken by Group X and Group Y.

.....

.....

.....

..... **[2]**


(d) Which group had a larger proportion of people completing the puzzle in less than 30 minutes? Explain your reasoning.

.....

.....

..... **[2]**

1236 The box-and-whisker plots show information about the times, in minutes, taken by two groups of students, Group A and Group B, to complete a mathematics test.



(a) Write down the median time for Group A.

..... minutes **[1]**

(b) Work out the interquartile range for Group B.

..... minutes **[1]**

(c) Make two comments comparing the distributions of the times taken by Group A and Group B.


Comment 1:

Comment 2: **[2]**

(d) Group A has 80 students.
Estimate the number of students in Group A who took longer than 55 minutes.

..... students **[2]**

1237 The box-and-whisker plots show the distribution of test scores for two classes, Class X and Class Y.



(a) Write down the median score for Class X.

..... **[1]**

(b) Work out the interquartile range for Class Y.

..... **[2]**

(c) Make two different comments comparing the distribution of test scores in Class X with Class Y.

[2]

(d) A student states: "Because the median score for Class X is higher, every student in Class X scored more than every student in Class Y."

Is this statement correct?
Give a statistical reason for your answer.

.....

..... **[1]**

1238 The table shows information about the mass, m grams, of 60 mangoes collected by a farmer.

Mass (m grams)	$100 < m \leq 120$	$120 < m \leq 140$	$140 < m \leq 160$	$160 < m \leq 180$	$180 < m \leq 200$
Frequency	10	20	15	10	5

(a) State the modal class.

..... **[1]**

(b) Calculate an estimate of the mean mass of the mangoes.

..... g **[4]**


(c) The farmer states that exactly 15 mangoes have a mass of 150 grams because 150 is the midpoint of the class $140 < m \leq 160$.
Explain why the farmer's reasoning is incorrect.

.....

.....

..... **[2]**

1239 The box-and-whisker plots show information about the times, in minutes, taken by a group of Adults and a group of Students to complete a logic puzzle.



(a) Use the diagram to find the interquartile range of the times taken by the Adults.

..... minutes **[2]**

(b) Make two different statements comparing the distribution of the times taken by the Adults with the times taken by the Students.

.....

.....

..... **[2]**

(c) A student claims: "Because the range of the Students' times is smaller, this proves that every student completed the puzzle faster than the slowest Adult."

Comment on the validity of this reasoning.

.....


.....

..... **[2]**

1240 The pie chart shows the distribution of grades (A, B, C, and D) achieved by 120 students in Class P.

The frequency table shows the distribution of the same grades achieved by the 80 students in Class Q.

The ratio of the number of students in Class Q achieving Grade A : Grade B : Grade C : Grade D is 1 : 3 : 2 : 2.



Class P

(a) Find the number of students who achieved Grade A in Class P.

..... **[3]**

(b) A student claims: "A higher percentage of students in Class Q achieved Grade B than in Class P."

Show that this claim is correct.

[3]

(c) The school principal states that Class P performed better overall because more students achieved Grade C in Class P than in Class Q.

Give one statistical reason why comparing the absolute frequencies (number of students) for


each grade is not an appropriate way to determine which class performed better.

.....

.....

..... **[1]**

1241 The cumulative frequency curve shows information about the masses, m grams, of 120 apples from Farm X.



(a) Use the graph to find an estimate for the median mass.

..... g **[1]**

(b) Use the graph to find an estimate for the interquartile range.

..... g **[2]**

(c) Apples with a mass greater than 128 g are classed as Large.
Find the percentage of these apples that are Large.

..... % **[2]**

(d) 120 apples from Farm Y have a median mass of 105 g and an interquartile range of 22 g.
Compare the distribution of the masses of the apples from Farm X and Farm Y.

.....

..... [2]

1242 A scientist records different types of weather data.
State whether each of the following types of data is discrete or continuous.

(a) The daily maximum temperature.

..... **[1]**

(b) The number of rainy days in a month.

..... **[1]**

(c) The amount of rainfall in one hour.

..... **[1]**

1243 Statistical data can be classified as either discrete or continuous.

(a) State whether each of the following variables is discrete or continuous data.

(i) The number of cars in a car park.

..... **[1]**

(ii) The temperature of a cup of coffee.

..... **[1]**

(iii) The time it takes to complete a jigsaw puzzle.

..... **[1]**

(iv) The number of keys on a piano.

..... **[1]**

1244 A sports analyst collects data during a football match.

(a) State whether the number of goals scored in the match is discrete or continuous.

..... **[1]**

(b) State whether the mass of the football is discrete or continuous.

..... **[1]**

(c) State whether the time taken to complete the match is discrete or continuous.

..... **[1]**

(d) State whether the temperature on the pitch is discrete or continuous.

..... **[1]**

1245 A student collects data for a statistics project.
For each of the variables below, state whether the data collected is discrete or continuous.

(a) The number of goals scored in a football match.

..... **[1]**

(b) The mass of a football.

..... **[1]**

(c) The volume of water in a swimming pool.

..... **[1]**

(d) The number of people swimming in a pool.

..... **[1]**

1246 Two sets of data are collected during a fitness test.
Data Set A is the exact time, t seconds, taken by athletes to run a race.
Data Set B is the number of push-ups, n, completed by the athletes.

(a) Data Set A is grouped into classes such as $10 \leq t < 12$.
Data Set B is grouped into classes such as $10 - 19$.
Explain why the classes for Data Set A use inequalities while the classes for Data Set B use a hyphen.

.....

..... **[2]**

(b) A student decides to draw a histogram to represent Data Set B. One of the classes is $20 - 29$.

(i) State the lower and upper class boundaries that must be used for this class on the horizontal axis of the histogram.

Lower = ... , Upper = ... **[1]**

(ii) Calculate the class width used for the frequency density calculation.

..... **[1]**

(c) Continuous data can theoretically take any value within a range.
Give an example of a continuous variable that is practically recorded as discrete data in everyday life.

.....

..... **[1]**

1247 A botanist studies a new species of plant in a greenhouse.
She records the height, h cm, of each plant and the total number of leaves on each plant.

(a) State whether the height of a plant is discrete or continuous data.

..... **[1]**

(b) The botanist wants to group the heights of the plants into a frequency table.
She suggests using the following class intervals:

$$10 - 19 \qquad 20 - 29 \qquad 30 - 39$$

Explain why these class intervals are not appropriate for the height of the plants.

.....

..... **[2]**

(c) State whether the number of leaves on a plant is discrete or continuous data.

..... **[1]**

1248 A student collects data about a local park for a science project.

(a) Explain the mathematical difference between discrete and continuous data.

.....

.....

..... **[2]**

(b) The student measures the lengths of 50 leaves to the nearest millimetre.

(i) State whether the true length of a leaf is discrete or continuous.

..... **[1]**

(ii) Explain why measuring the lengths to the nearest millimetre does not make the variable discrete.

.....

..... **[1]**

1249 A farmer weighs 60 apples.
The masses, m grams, are recorded in the table.

Mass (m grams)	**Frequency**
$80 < m \leq 100$	12
$100 < m \leq 120$	26
$120 < m \leq 140$	15
$140 < m \leq 160$	7

(a) State whether the mass of an apple is discrete or continuous data.

..... **[1]**

(b) Calculate an estimate of the mean mass.

..... g **[3]**

(c) Explain why your answer to part (b) is an estimate.

..... **[1]**

1250 A scientist measures the lengths of 50 worms and counts the number of spots on each worm.
The frequency table shows the data collected for the number of spots.

Number of spots	$0-4$	$5-9$	$10-14$	$15-19$
Frequency	12	18	14	6

(a) State which of the two variables, length or number of spots, is continuous.

..... **[1]**

(b) The length of one worm is recorded as 14.5 cm, correct to the nearest millimetre.

(i) Write down the lower bound of this length.

..... *cm* **[1]**

(ii) Write down the upper bound of this length.

..... *cm* **[1]**

(c) Using the frequency table, write down the class width for the $5-9$ class.

..... **[2]**

1251 Data is collected about the members of an athletics club.

(a) State whether each of the following variables is discrete or continuous.

(i) The number of members in the club.

..... **[1]**

(ii) The exact distance a member jumps in the long jump.

..... **[1]**

(b) The shoe sizes of the members are recorded.
Shoe sizes are available in sizes such as $7, 7.5, 8, 8.5, 9$.
A student claims that because these shoe sizes include decimal values, the data is continuous.
Explain why the student is incorrect.

.....

.....

..... **[2]**

1252 Measurements and quantities in statistics are defined by their ability to take certain values.

(a) Explain the mathematical difference between continuous data and discrete data.

.....

..... **[2]**

(b) A person's exact foot length is measured in centimetres.
This measurement is then used to determine their shoe size, which can take values such as 7, 7.5, 8, or 8.5.

(i) State whether exact foot length is discrete or continuous data.

..... **[1]**

(ii) A student argues that shoe size must be continuous because it includes decimal values like 7.5.
Explain why the student is incorrect.

.....

.....
[2]

1253 A factory packs identical books into boxes.
Each box contains exactly 24 books.

(a) State whether each of the following variables is discrete or continuous.

(i) The number of books in a box.

..... **[1]**

(ii) The mass of a book.

..... **[1]**

(b) The mass of one book is 850 g, correct to the nearest 10 g.
Find the upper bound for the total mass of the 24 books.
Give your answer in kilograms.

..... kg **[2]**

(c) An empty box has a mass of 1.2 kg, correct to 1 decimal place.
Calculate the lower bound for the total mass of a box containing 24 books.

..... kg **[3]**

1254 The number of goals scored by a football team in each of their last 10 matches is shown below.

2 0 1 4 1 0 3 1 2 1

(a) Find the mode.

..... **[1]**

(b) Find the median.

..... **[2]**

(c) Calculate the mean.

..... **[2]**

1255 A student records the number of points scored by a basketball team in 11 matches.

14 20 8 24 16 5 26 13 22 10 18

(a) Find the median.

..... **[1]**

(b) Find the range.

..... **[1]**

(c) Find the interquartile range.

..... **[2]**

(d) Calculate the mean.

..... **[2]**

1256 The list shows the number of points scored by a basketball player in 7 games.

$$7 \quad 12 \quad 8 \quad 7 \quad 12 \quad x \quad 10$$

The mode of the number of points is 7.

(a) Write down the value of x.

$x =$ ..... **[1]**

(b) Find the range of the number of points.

..... **[1]**

(c) Find the median.

..... **[1]**

(d) Calculate the mean.

..... **[2]**

(e) In the 8th game, the player scores y points.
The new mean number of points for all 8 games is 9.5.
Find the value of y.

$y =$ ..... **[2]**

1257 The table shows the heights, h cm, of 80 plants.

Height (h cm)	**Frequency**
$10 < h \leq 20$	12
$20 < h \leq 30$	25
$30 < h \leq 40$	28
$40 < h \leq 50$	11
$50 < h \leq 60$	4

(a) Write down the modal class.

..... **[1]**

(b) Calculate an estimate of the mean height.

..... cm **[4]**

1258 A survey was conducted on the masses, m grams, of 80 apples. The results are shown in the frequency table.

Mass (m grams)	$80 < m \leq 90$	$90 < m \leq 100$	$100 < m \leq 110$	$110 < m \leq 120$	$120 < m \leq 130$
Frequency	8	22	31	14	5

(a) Write down the modal class.

..... **[1]**

(b) Calculate an estimate of the mean mass of the apples.

..... g **[4]**

(c) Explain why your answer to part (b) is an estimate.

.....

..... **[2]**

1259 The table shows information about the time, t seconds, taken by 100 students to complete a puzzle.

Time (t seconds)	Frequency
$10 < t \leq 20$	14
$20 < t \leq 30$	28
$30 < t \leq 40$	32
$40 < t \leq 50$	18
$50 < t \leq 60$	8

(a) Write down the modal time interval.

..... **[1]**

(b) Calculate an estimate of the mean time.

..... seconds **[4]**

(c) Calculate the percentage of students who took strictly more than 40 seconds to complete the puzzle.

..... % **[2]**

(d) Write down the interval that contains the upper quartile.

..... **[1]**

1260 A factory produces 60 items.
The masses, m kg, of these items are recorded in the table.
The estimate of the mean mass is 48 kg.

Mass (m kg)	**Frequency**
$0 < m \leq 20$	p
$20 < m \leq 40$	14
$40 < m \leq 60$	16
$60 < m \leq 80$	q
$80 < m \leq 100$	5

(a) Find the value of p and the value of q.

$p =$ ..... , $q =$ ..... **[5]**

(b) State the class interval which contains the median mass.
You must show your working.

..... **[2]**

1261 The mean of five numbers is 12.
The median of the five numbers is 11.
The mode of the five numbers is 8.
The range of the five numbers is 15.

(a) Find the five numbers.

..... **[5]**

(b) A sixth number, k, is added to the list.
The new mean is 13.
Work out the value of k.

$k =$ ..... **[2]**

1262 The table shows information about the masses, m kg, of 50 parcels.

Mass (m kg)	Frequency
$0 < m \leq 2$	15
$2 < m \leq 4$	x
$4 < m \leq 6$	y
$6 < m \leq 8$	8
$8 < m \leq 10$	2

The estimate of the mean mass of the parcels is 3.8 kg.

(a) Use the total number of parcels to write down an equation in x and y, and show that it simplifies to $x + y = 25$.

[1]

(b) Use the estimate of the mean to show that $3x + 5y = 101$.

[4]

(c) Solve the simultaneous equations to find the value of x and the value of y.
You must show all your working.

$x =$ ..... , $y =$ ..... **[3]**

(d) Write down the interval that contains the median mass.

..... **[1]**

1263 The number of text messages received by 15 students in one hour is shown below.

12	23	15	18	21	14	17	15	22	19	15	24	11	16	20

(a) Find the mean.

..... **[1]**

(b) Find the median.

..... **[1]**

(c) Find the interquartile range.

..... **[2]**

1264 A survey was conducted to find the number of books read by 21 students during a month.
The results are shown in the table.

Number of books	1	2	3	4	5
Frequency	2	5	7	4	3

(a) Use a graphic display calculator to find the following statistics for this data.

(i) Calculate the mean number of books read.
Give your answer correct to 3 significant figures.

Mean = ... **[2]**

(ii) Find the median.

Median = ... **[1]**

(iii) Find the interquartile range.

Interquartile range = ... **[2]**

1265 The marks scored by 15 students in a mathematics test are shown below.

15	19	14	28	11	16	9	20	15	13	17	21	10	18	14

(a) Find the mean.

..... **[2]**

(b) Find the median.

..... **[1]**

(c) Find the interquartile range.

..... **[2]**

1266 The table shows information about the time, t minutes, taken by 100 people to travel to work.

Time (t minutes)	Frequency
$0 < t \leq 10$	14
$10 < t \leq 20$	28
$20 < t \leq 30$	35
$30 < t \leq 40$	18
$40 < t \leq 60$	5

(a) Write down the modal class.

..... **[1]**

(b) Use your graphic display calculator to calculate an estimate of the mean time.

..... minutes **[3]**

(c) Explain why your answer to part (b) is an estimate.

..... **[1]**

1267 The table shows the time, t minutes, taken by 80 people to complete a crossword puzzle.

Time (t minutes)	$0 < t \leq 10$	$10 < t \leq 20$	$20 < t \leq 30$	$30 < t \leq 40$	$40 < t \leq 60$
Frequency	12	25	28	11	4

(a) Calculate an estimate of the mean time.

..... minutes **[3]**

(b) Explain why your answer to part (a) is an estimate.

..... **[1]**

(c) An 81st person completes the puzzle in 12 minutes.
Without calculating the new mean, explain how this affects the mean time.

..... **[1]**

1268 A survey was conducted on the time, t minutes, taken by 80 students to travel to school. The results are shown in the frequency table.

Time (t minutes)	$0 < t \leq 10$	$10 < t \leq 20$	$20 < t \leq 30$	$30 < t \leq 40$	$40 < t \leq 50$
Frequency	12	25	28	11	4

(a) Calculate an estimate of the mean time.

..... minutes **[2]**

(b) Explain why your answer to part (a) is an estimate.

.....

..... **[1]**

(c) The mean time for a different group of 120 students to travel to school is 24.5 minutes. Calculate the overall mean time for the 200 students.

..... minutes **[3]**

1269 The table shows the number of goals scored by a football team in 35 matches.

Number of goals	0	1	2	3	4	5
Frequency	3	x	y	6	4	1

(a) The mean number of goals scored is 2.2.
Find the value of x and the value of y.

$x =$ ..... , $y =$ ..... **[4]**

(b) Find the median number of goals.

..... **[1]**

1270 The table shows information about the number of goals scored by a football team in a number of matches.

Number of goals	0	1	2	3	4
Frequency	6	x	8	4	2

The mean number of goals scored is 1.5.

(a) Show that $x = 12$.

[3]

(b) Find the median.

..... **[1]**

(c) Find the interquartile range.

..... **[2]**

1271 A game is played where players can score $1, 2, 3, 4$, or 5 points.
The table shows the frequencies of the scores obtained by a group of players.
The mean score is 2.8.

Score	1	2	3	4	5
Frequency	10	k	$2k+3$	7	5

(a) Show algebraically that $k = 5$.

[4]

(b) Use your graphic display calculator to find the following statistics for the scores.

(i) Find the median.

Median = ... **[1]**

(ii) Find the lower quartile.

Lower quartile = ... **[1]**

(iii) Find the upper quartile.

Upper quartile = ... **[1]**

1272 The stem-and-leaf diagram shows the masses, in kg, of 15 parcels.

1	2	4	7		
2	0	1	1	1	5
3	0	2	6	8	
4	1	4	5		

Key: 1 | 2 represents 12 kg

(a) Find the mode of the masses.

..... kg **[1]**

(b) Find the median mass.

..... kg **[1]**

(c) Work out the range of the masses.

..... kg **[1]**

(d) A parcel is chosen at random.
Find the probability that its mass is strictly greater than 30 kg.

..... **[2]**

1273 The masses, in kilograms, of 15 parcels are recorded below.

27	15	31	42	23	12	53	38	49	27	45	40	34	51	42

(a) Complete the ordered stem-and-leaf diagram to show this information.

1	
2	
3	
4	
5	

Key: 1 | 2 represents 12 kg

[3]


(b) Find the median mass.

..... kg **[1]**

(c) Work out the range.

..... kg **[1]**

1274 The pie chart shows information about the favourite subjects of 180 students.



(a) Find the value of x.

$x =$ ..... **[1]**

(b) Find the number of students whose favourite subject is Science.

..... **[2]**

(c) Write down the fraction of students whose favourite subject is Maths.
Give your fraction in its simplest form.

..... **[2]**

1275 The masses, m kilograms, of 15 parcels are recorded.

3.2	1.5	4.6	2.1	1.8	3.5	2.9	4.1	3.2	1.9	2.4	3.7	4.0	2.6	1.7

(a) Complete the ordered stem-and-leaf diagram to show this information.

1	
2	
3	
4	

Key: 1 | 5 represents 1.5 kg

[3]


(b) Find the median mass.

..... kg **[1]**

(c) A parcel is chosen at random.
Find the probability that its mass is greater than 3.5 kg.

..... **[1]**

1276 The pie chart shows information about how 120 students travel to school.



(a) Calculate the number of students who travel by Bus.

..... **[2]**

(b) There are 25 students who travel by Car.
Calculate the sector angle for Car.

..... ° **[2]**

(c) Calculate the percentage of students who travel by Bicycle.

..... % **[3]**

1277 A survey asked 120 students about their favourite sport.
The results are shown in the table.

Sport	**Frequency**
Football	45
Basketball	35
Tennis	25
Swimming	15

(a) A pie chart is drawn to show this information.
Calculate the sector angle for Basketball.

..... ° **[2]**

(b) The pie chart has a radius of 6 cm.

Calculate the area of the sector representing Tennis.

..... cm^2 **[3]**

(c) In a different school, the pie chart for favourite sports has a sector angle of 108° for Football.
This sector represents 54 students.
Calculate the total number of students in this school.

..... **[2]**

1278 A pie chart is drawn to show the favourite sport of a group of students.
The angles of the sectors for each sport are given in terms of x.

Football: $(2x + 30)^\circ$ Basketball: x° Tennis: $(x + 50)^\circ$ Swimming: $(2x - 20)^\circ$

(a) Find the value of x.

$x =$ ..... **[2]**


(b) 10 students chose Tennis as their favourite sport.
Calculate the total number of students in the group.

..... **[2]**

(c) Two students are chosen at random from the group.
Calculate the probability that one student chose Football and the other student chose Basketball.

..... **[3]**

1279 A pie chart shows the grades (A, B, C, and D) achieved by students in a mathematics test.
The angles of the sectors are given in terms of x.



(a) Show that $x = 45$.

[3]


(b) There are 24 students who achieved Grade A.
Calculate the total number of students who took the test.

..... **[2]**

(c) Calculate the percentage of students who achieved Grade C.

..... % **[3]**

1280 A company records the number of Laptops and Tablets sold over three months.
The dual bar chart shows the sales for January, February, and March.
The bar for Tablets sold in March is incomplete and not shown.



(a) The total number of Laptops sold over the three months is 20% more than the total number of Tablets sold.
Calculate the number of Tablets sold in March.

..... **[4]**


(b) In April, the total number of items sold is $\frac{1}{3}$ of the total number of items sold in January, February, and March combined.
The ratio of Laptops to Tablets sold in April is 6 : 5.
Work out the number of Laptops sold in April.

..... **[4]**

(c) The price of a Laptop is \$450 and the price of a Tablet is \$200.
Calculate the percentage of the total revenue in January that came from Tablet sales.

..... % **[3]**

1281 The scatter diagram shows the relationship between the maximum daily temperature, x°C, and the number of ice creams sold, y, on 10 different days.



(a) State the type of correlation shown in the scatter diagram.

..... **[1]**

(b) On the scatter diagram, draw a line of best fit. **[1]**


(c) Use your line of best fit to estimate the number of ice creams sold on a day when the maximum temperature is 22°C.

..... **[1]**

1282 The table shows the outside temperature, T°C, and the number of hot drinks, N, sold by a cafe on 10 different days.

Temperature (T°C)	5	8	12	15	18	20	22	25	28	30
Number of hot drinks (N)	140	125	100	95	70	65	50	40	25	15

(a) Complete the scatter diagram.
The first four points have been plotted for you.



[2]

(b) What type of correlation is shown by the scatter diagram?


..... **[1]**

(c) Draw a line of best fit on the scatter diagram. **[1]**

1283 A teacher records the time, x minutes, spent studying and the score, y marks, for 8 students.
The table shows the results.

Time (x min)	10	20	30	40	50	60	70	80
Score (y marks)	15	25	20	40	50	70	60	80

(a) On the grid, complete the scatter diagram to show this information.



[2]

(b) What type of correlation is shown by the scatter diagram?

..... **[1]**

(c) On the scatter diagram, draw a line of best fit. **[1]**

(d) Another student spent 45 minutes studying.
Use your line of best fit to estimate their score.

..... **[1]**

1284 The table shows the time spent studying, x hours, and the test score, y marks, for a group of 8 students.

Study time (x hours)	2	3.5	4	5	6.5	8	9	10
Test score (y marks)	45	52	60	68	75	82	85	95

(a) Use your graphic display calculator to find the equation of the linear regression line, giving y in terms of x.
Give your answer in the form $y = mx + c$.

$y =$ ..... **[2]**

(b) Use your equation to estimate the test score for a student who studies for 7 hours.

..... **[2]**


(c) Explain why it may not be reliable to use this equation to estimate the test score for a student who studies for 15 hours.

..... **[1]**

1285 A car dealer records the age, x years, and the price, y thousand dollars, of 12 used cars of the same model.
The table shows the data.

Age (x years)	1	2	2	3	3	4	5	5	6	7	7	9
Price (y thousand \$)	19	18	16	15	14	12	10	9	7	6	5	1

(a) The scatter diagram shows the first 10 points from the table.
Plot the remaining 2 points on the scatter diagram.



[1]

(b) Calculate the mean age and the mean price.

Mean age = ... , Mean price = ... **[2]**

(c) On the scatter diagram, plot the mean point and draw a line of best fit. **[2]**

(d) State the type of correlation shown.

..... **[1]**

(e) Use your line of best fit to estimate the price of a car that is 8 years old.


..... thousand dollars **[1]**

1286 A teacher records the number of hours spent studying, x, and the test score, y, for 7 students.

Study time (x hours)	2	3	4	5	6	7	8
Test score (y)	42	48	63	68	81	88	100

The scatter diagram for this data is partially completed.

(a) Plot the remaining three points on the scatter diagram.



[2]

(b) Calculate the mean study time, $\bar{x}$, and the mean test score, $\bar{y}$.

$\bar{x} =$ ..... , $\bar{y} =$ ..... **[2]**

(c) On the scatter diagram, plot the mean point and draw a line of best fit. **[2]**

(d) Another student studied for 5.5 hours.
Use your line of best fit to estimate their test score.

..... **[1]**

1287 A researcher measures the height, x cm, and the mass, y kg, of 12 adult males.
The results are summarised as follows:

$$\sum x = 2100 \qquad \sum y = 924$$

The equation of the linear regression line is $y = 0.85x - c$, where c is a constant.

(a) Calculate the mean height, $\bar{x}$, and the mean mass, $\bar{y}$.

$\bar{x} =$ ..... , $\bar{y} =$ ..... **[2]**

(b) The point $(\bar{x}, \bar{y})$ lies on the linear regression line.
Find the value of c.

$c =$ ..... **[2]**

(c) A new adult male is added to the study.
His height is 180 cm.
His mass is exactly equal to the value predicted by the regression line.
Calculate the new mean mass for all 13 men.

..... kg **[2]**

1288 An automotive engineer tests 8 different cars to investigate the relationship between engine size, x litres, and fuel consumption, y km/l.
The results are shown in the table.

Engine size (x litres)	1.0	1.2	1.5	1.8	2.0	2.4	2.5	3.0
Fuel consumption (y km/l)	21.5	20.0	18.2	16.5	15.0	13.0	12.5	9.8

(a) **(i)** Use a graphic display calculator to find the equation of the linear regression line. Give your answer in the form $y = mx + c$.

$y =$ ..... **[2]**

(ii) Explain what the value of the gradient represents in this context.

..... **[1]**

(b) Use your equation from part (a)(i) to estimate the fuel consumption of a car with an engine size of 1.6 litres.

..... km/l **[2]**

(c) Explain why it may not be reliable to use this equation to estimate the fuel consumption of a car with an engine size of 5.0 litres.

..... **[1]**

1289 The table shows the temperature, x°C, and the number of ice creams sold, y, at a shop over 8 days.

Temperature (x°C)	12	15	18	20	22	25	28	30
Ice creams sold (y)	40	52	68	75	84	98	110	115

(a) Calculate the mean temperature and the mean number of ice creams sold.

Mean temperature = ... , Mean ice creams sold = ... **[2]**

(b) Use a graphic display calculator to find the equation of the linear regression line for this data. Give your answer in the form $y = mx + c$, with values of m and c correct to 3 significant figures.

$y =$ ..... **[2]**


(c) The shop owner predicts the number of ice creams sold on a day when the temperature is 24°C. Use your equation to estimate this number.

..... **[2]**

(d) Comment on the reliability of using your equation to predict the number of ice creams sold on a day when the temperature is 42°C.

..... **[1]**

1290 The cumulative frequency diagram shows the times taken, t minutes, by 120 students to complete a test.



(a) Use the diagram to estimate the median time.

..... minutes **[1]**


(b) Use the diagram to estimate the interquartile range.

..... minutes **[2]**

(c) Students who take longer than 45 minutes are given a penalty.
Estimate the number of students who are given a penalty.

..... **[2]**

1291 The cumulative frequency diagram shows information about the distance, d km, travelled by 60 people to work.



(a) Use the diagram to find an estimate for:

(i) the median,

..... km **[1]**

(ii) the interquartile range.

..... km **[2]**


(b) Find the number of people who travelled more than 45 km to work.

..... **[3]**

1292 The table shows the cumulative frequencies for the times, t minutes, taken by 80 students to complete a puzzle.

Time (t minutes)	$t \leq 10$	$t \leq 20$	$t \leq 30$	$t \leq 40$	$t \leq 50$	$t \leq 60$
Cumulative frequency	6	22	48	68	78	80

(a) On the grid, draw a cumulative frequency diagram to show this information.



[3]

(b) Use your diagram to estimate the median time.

..... minutes **[2]**

1293 A company records the salaries, s thousand dollars, of its 200 employees.
The frequency table shows the results.


Salary (s thousand dollars)	$20 < s \leq 30$	$30 < s \leq 40$	$40 < s \leq 50$	$50 < s \leq 60$	$60 < s \leq 70$
Frequency	24	66	70	28	12

(a) Complete the cumulative frequency table.

Salary (s thousand dollars)	$s \leq 30$	$s \leq 40$	$s \leq 50$	$s \leq 60$	$s \leq 70$
Cumulative frequency	24				

[2]

(b) On the grid, draw a cumulative frequency diagram to show this information.




[3]

(c) Use your diagram to estimate the 70th percentile of the salaries.

..... thousand **[2]**

1294 The cumulative frequency diagram shows information about the distances, d km, travelled by 120 people to work.



(a) Use the diagram to estimate the median distance.

..... km **[2]**

(b) Use the diagram to estimate the interquartile range.

..... km **[3]**

(c) Estimate the number of people who travelled more than 35 km to work.

..... **[3]**

1295 A table shows information about the masses of 120 apples.


Mass (m grams)	$50 < m \leq 100$	$100 < m \leq 150$	$150 < m \leq 200$	$200 < m \leq 250$	$250 < m \leq 300$
Frequency	12	34	46	20	8

(a) Complete the cumulative frequency table.

Mass (m grams)	$m \leq 100$	$m \leq 150$	$m \leq 200$	$m \leq 250$	$m \leq 300$
Cumulative frequency	12				120

[2]

(b) On the grid, draw a cumulative frequency diagram to show this information.



[3]


(c) Use your diagram to estimate the 60th percentile.

..... g **[2]**

(d) Apples with a mass greater than 220 grams are classified as large.
Estimate the number of large apples.

..... **[3]**

1296 The cumulative frequency diagram shows information about the masses, m grams, of 200 parcels.



(a) Find an estimate for the 80th percentile.

..... grams **[2]**

(b) Use the cumulative frequency diagram to complete the frequency table.


Mass (m grams)	$0 < m \leq 10$	$10 < m \leq 20$	$20 < m \leq 30$	$30 < m \leq 40$
Frequency				

[2]

(c) Calculate an estimate of the mean mass.

..... grams [4]

1297 The cumulative frequency curve shows information about the masses, m grams, of 100 potatoes.



(a) Use the curve to estimate the number of potatoes with a mass greater than 220 grams.

..... **[3]**

(b) The curve was drawn using grouped data.
Find the values of p, q, and r.


Mass (m grams)	$50 < m \leq 100$	$100 < m \leq 150$	$150 < m \leq 200$	$200 < m \leq 250$	$250 < m \leq 300$
Frequency	p	q	40	r	6

$p =$ ..... , $q =$ ..... , $r =$ ..... **[3]**

(c) Calculate an estimate of the mean mass of the potatoes.

..... g **[3]**

1298 The cumulative frequency diagram shows the distribution of marks scored by 200 candidates in an examination.



(a) Use your graph to estimate the interquartile range.

..... **[2]**

(b) 15% of the candidates score less than m marks.
Find the value of m.

$m =$ ..... **[2]**


(c) Candidates scoring more than P marks are awarded a Distinction.
Given that 30 candidates are awarded a Distinction, find the value of P.

$P =$ ..... **[2]**

(d) Candidates who score less than 45 marks have to take a supplementary test.
Calculate the percentage of candidates who do NOT have to take this test.

..... % **[4]**

1299 The diagram shows the positions of three lighthouses,
A,
B, and
C. Lighthouse
C is due east of lighthouse
A, and the bearing of
B from
A is
054
054
.



(a) Write down the three-figure bearing of
C from
A.

.....

[1]

(b) Calculate the three-figure bearing of
A from
B.

.....


[2]

(c) Given that
=
AB=AC, calculate the three-figure bearing of
C from
B.

.....

[2]

1300 The scale drawing shows the positions of two towns,
P and
Q, on a map.
The scale is
1
1 cm represents
5
5 km.



(a) Measure the three-figure bearing of
Q from
P.

.....


[1]

(b) Work out the actual distance, in kilometres, from town
P to town
Q.

..... km **[2]**

(c) A third town,
R, is
25
25 km from
Q on a bearing of
140
140
.
Using a ruler and protractor, mark and label the position of town

R on the scale drawing below.




[2]

(d) Without measuring, calculate the three-figure bearing of P from Q.

.....

[2]

1301 The diagram shows the positions of four radar stations,
A,
B,
C, and
D, on flat ground.
Station
C is due east of station
D. The bearing of
A from
D is
018.6
018.6
, the bearing of
B from
A is
048.4
048.4
, and the bearing of
B from
C is
322.8
322.8
.



(a) Write down the three-figure bearing of
D from
C.

.....

[1]

(b) Calculate the three-figure bearing of
D from
A.

. . . . . .

[2]

(c) Calculate the size of

(i) angle DAB,

Angle = DAB= . . . . . .

[2]

(ii) angle ABC.

Angle = ABC= . . . . . .

[3]

(d) Calculate the three-figure bearing of C from B.

. . . . . .

[2]

1 **(a)** $2^3 \times 3^2 \times 7$ **[B2]**

M1 for a correct partial factorisation (e.g., 8×63 or 2×252) or a correct prime factor tree/ladder.

(b) 14 **[B2]**

M1 for identifying the needed powers 2×7 or writing $2^4 \times 3^2 \times 7^2$.

[Back to Question] *[Total Marks: 4]*

2 **(a)** $2^2 \times 3 \times 7$ **[B2]**

M1 for $2, 2, 3, 7$ correctly identified in a factor tree or division ladder

(b) 252 **[B2]**

M1 for writing $126 = 2 \times 3^2 \times 7$ or writing multiples of 84 and 126

[Back to Question] *[Total Marks: 4]*

3 **(a)** **(i)** 12 or 36 **[B1]**

(ii) 64 **[B1]**

(iii) 23 **[B1]**

(b) Prime factors: $36 = 2^2 \times 3^2$ and $64 = 2^6$ **[M1]**

LCM $= 2^6 \times 3^2 = 576$ **[A1]**

[Back to Question] *[Total Marks: 5]*

4 **(a)** 13 **[B1]**

(b) $\sqrt{7}$ **[B1]**

(c) -5 **[B1]**

(d) 0.25 **[B1]**

(e) 64 **[B1]**

[Back to Question] *[Total Marks: 5]*

5 **(a)** **(i)** 23 **[B1]**

(ii) 8 or 27 or 64 **[B1]**

(iii) 64 **[B1]**

(b) $2^2 \times 3^2$ (or $2 \times 2 \times 3 \times 3$) **[B2]**

Partial working: Correct factor tree or repeated division showing 2 and 3 as prime factors **[M1]**

[Back to Question] *[Total Marks: 5]*

6 **(a)** **(i)** 2 or 23 **[B1]**

(ii) 27 or 64 **[B1]**

(iii) 64 **[B1]**

(iv) 15 **[B1]**

(b) $2^3 \times 3^2$ **[B2]**

If 0 scored, M1 for $2 \times 2 \times 2 \times 3 \times 3$ or a correct factor tree/division **[M1]**

[Back to Question] *[Total Marks: 6]*

7 **(a)** **(i)** 13 **[B1]**

(ii) 8 or -1 **[B1]**

(iii) $\sqrt{30}$ or π **[B1]**

(iv) -5 or -1 **[B1]**

(b) $\frac{5}{4}$ or $1\frac{1}{4}$ **[B2]**

M1 for $\frac{1}{0.8}$ or $\frac{10}{8}$ or $\frac{8}{10}$ seen.

[Back to Question] *[Total Marks: 6]*

8 **(a)** 36 (or $2^2 \times 3^2$) **[B2]**

Partial working: Identifies 2^2 and 3^2 as common prime factors **[M1]**

(b) $2^3 \times 3^4 \times 5 \times 7$ **[B2]**

Partial working: One incorrect power or missing prime factor **[B1]**

(c) 66 **[B3]**

Partial working: Identifies that C must contain the factor 11 from the LCM **[M1]**

Partial working: Identifies that C must contain exactly $2^1 \times 3^1$ to satisfy the HCF condition **[M1]**

[Back to Question] *[Total Marks: 7]*

9 **(a)** Irrational **[B1]**

(b) Any valid example, e.g. $\sqrt{2}$, $\sqrt{3}$, π (if $y^2 = \pi^2$ which is irrational, so this is wrong)

Only square roots of non-square rational numbers, e.g. $\sqrt{2}$ **[B1]**

(c) **(i)** 53 and Yes **[B2]**

M1 for substituting $n = 3$: $3^2 + 3 + 41$

(ii) Substituting $n = 41$ yields $41^2 + 41 + 41$ **[M1]**

This factorises to $41(41 + 1 + 1) = 41 \times 43$, which has factors other than 1 and itself, therefore it is not prime. **[A1]**

[Back to Question] *[Total Marks: 6]*

10 **(a)** 2 **[B3]**

M2 for $540 = 2^2 \times 3^3 \times 5$ and $180 = 2^2 \times 3^2 \times 5$ written as prime products

or M1 for either 540 or 180 correctly written as a product of prime factors.

(b) $y = 4$ **[B1]**

$z = 3$ **[B1]**

(c) 15 **[B2]**

M1 for identifying the needed powers to reach multiples of 3, e.g. $3^1 \times 5^1$.

[Back to Question] *[Total Marks: 7]*

11 **(a)** 40 **[B2]**

If 0 scored, allow [M1] for setting up the equation $12x = 4 \times 120$ or finding prime factors of 120 indicating $2^3 \times 5$.

(b) **(i)** $2^4 \times 3^2 \times 7$ **[B2]**

If 0 scored, allow [M1] for a correct factor tree or continuous division with at least 2 correct prime division steps.

(ii) 216 **[B3]**

If 0 scored, allow [M1] for writing $72 = 2^3 \times 3^2$.

Allow a further [M1] for deducing that N must contain $2^3 \times 3^3$ to be the smallest perfect cube with the correct HCF (power of 2 is exactly 3, power of 3 is at least 3).

[Back to Question] *[Total Marks: 7]*

12 **(a)** **(i)** pq **[B1]**

(ii) p^3q^3 **[B1]**

(b) $pq = 15$, so the prime numbers are 3 and 5 **[M1]**

$m > n \implies p^3q > pq^3 \implies p^2 > q^2 \implies p > q$ **[M1]**

$p = 5$ and $q = 3$ **[A1]**

(c) $m \times n = (p^3q)(pq^3) = p^4q^4$ **[M1]**

$= (p^2q^2)^2$ which is the square of an integer, hence a square number **[A1]**

(d) Substitutes values: $m = 5^3 \times 3^1$ **[M1]**

Recognises the power of 3 needs to be a multiple of 3, so $k = 3^2 = 9$ **[A1]**

[Back to Question] *[Total Marks: 9]*

13 $x = 2$ **[B2]**

$y = 3$ **[B2]**

If 0 scored, award M1 for expressing $180 = 2^2 \times 3^2 \times 5$ or $12600 = 2^3 \times 3^2 \times 5^2 \times 7$

[Back to Question] *[Total Marks: 4]*

14 **(a)** 36 (or $2^2 \times 3^2$) **[B1]**

(b) $2^3 \times 3^3 \times 5 \times 7$ **[B2]**

If 0 scored, allow [M1] for a product with at least three correct prime bases and powers, or for the correct evaluation 7560.

(c) 10 **[B2]**

If 0 scored, allow [M1] for identifying required missing powers 2×5 or $2^1 \times 5^1$.

[Back to Question] *[Total Marks: 5]*

15 **(a)** Identifies HCF as $2^1 \times 3^1$ **[M1]**

$x = 1$ **[A1]**

(b) Prime factorises $8100 = 81 \times 100 = 3^4 \times 2^2 \times 5^2$ **[M1]**

Compares highest powers of 3 between A (3^3) and C (3^y) to the LCM (3^4) **[M1]**

$y = 4$ **[A1]**

(c) $\sqrt{11}$ **[B1]**

4π **[B1]**

[Back to Question] *[Total Marks: 7]*

16 **(a)** 12 **[B2]**

If 0 scored, M1 for $84 = 2^2 \times 3 \times 7$ and $120 = 2^3 \times 3 \times 5$ OR for a common factor of 4 or 6 identified **[M1]**

(b) 840 **[B2]**

If 0 scored, M1 for $2^3 \times 3 \times 5 \times 7$ **[M1]**

(c) $\frac{5}{4}$ or $1\frac{1}{4}$ **[B2]**

If 0 scored, M1 for $\frac{1}{0.8}$ or $\frac{10}{8}$ seen **[M1]**

(d) Irrational, because 120 is not a perfect square (or $\sqrt{120}$ cannot be written as a simple fraction) **[B2]**

If 0 scored, B1 for Irrational with no/incorrect reason **[B1]**

[Back to Question] *[Total Marks: 8]*

17 **(a)** **(i)** $x = 2$, $y = 8$, $z = 10$ **[B2]**

Partial working: Lists correctly prime numbers $x < 10$ and cube numbers $y > 1$ **[M1]**

(ii) $x = 7$, $y = 8$, $z = 15$ OR $x = 2$, $y = 64$, $z = 66$ **[B2]**

Partial working: Identifies another valid combination of a prime and cube summing to a triangle number **[M1]**

(b) (i) e.g. $p = \sqrt{2}$, $q = -\sqrt{2}$ OR $p = 1 + \sqrt{3}$, $q = 1 - \sqrt{3}$ **[B1]**

(ii) e.g. $p = \sqrt{2}$, $q = \sqrt{8}$ OR $p = \sqrt{3}$, $q = 2\sqrt{3}$ **[B1]**

[Back to Question] *[Total Marks: 6]*

18 (a) $x = 2$ **[B1]**

$y = 2$ **[B1]**

M1 for $84 = 2^2 \times 3 \times 7$ OR $3528 = 2^3 \times 3^2 \times 7^2$ seen **[M1]**

(b) 1764 **[B1]**

(c) 10:21:00 **[B3]**

M2 for LCM of 45 and 72 = 360 seconds (or 6 minutes) **[M2]**

If M2 not scored, M1 for listing multiples of 45 and 72 with at least one common multiple, OR $45 = 3^2 \times 5$ and $72 = 2^3 \times 3^2$ **[M1]**

[Back to Question] *[Total Marks: 7]*

19 (a) Correct elements in all four regions: 2 in the intersection, $4, 6, 8, 10$ in A only, $3, 5, 7$ in B only, and $1, 9$ outside both circles **[B2]**

If 0 scored, allow [B1] for 2 or 3 correct regions

(b) $\{2\}$ or 2 **[B1]**

(c) 2 **[B1]**

[Back to Question] *[Total Marks: 4]*

20 (a) 2, 3, 5, 7 **[B1]**

(b) 1, 2, 3, 4, 6, 12 **[B1]**

(c) 8 **[B1]**

(d) 2, 3 **[B1]**

(e) 8 or 9 or 10 **[B1]**

[Back to Question] *[Total Marks: 5]*

21 (a) $\{3, 5, 7\}$ **[B1]**

(b) 4 **[B2]**

M1 for correctly identifying $(A \cup B)' = \{4, 6, 8, 10\}$ or $A \cup B = \{1, 2, 3, 5, 7, 9\}$.

(c) $\in$ **[B1]**

(d) $\subset$ or $\subseteq$ **[B1]**

[Back to Question] *[Total Marks: 5]*

22 **(a)** Intersection: $3, 5, 7$ **[B1]**

Remainder of diagram correct: A only region has 2; B only region has $1, 9$; Outer region has $4, 6, 8, 10$ **[B1]**

(b) 2 **[B1]**

(c) 6 **[B1]**

(d) $A \cap B'$ **[B1]**

[Back to Question] *[Total Marks: 5]*

23 **(a)** $1, 2, 3, 6, 9$ **[B1]**

(b) $2, 3$ **[B1]**

(c) 3 **[B2]**

Partial working: Correctly identifies elements in the union $\{1, 2, 3, 5, 6, 7, 9\}$ OR elements outside $\{4, 8, 10\}$ **[M1]**

(d) $\notin$ **[B1]**

[Back to Question] *[Total Marks: 5]*

24 **(a)** 4 **[B2]**

M1 for $6 + 3 + 7 + 1 + 2 + 4 + 8 + x = 35$ or $31 + x = 35$.

(b) 9 **[B1]**

(c) $A' \cap B \cap C$ or $B \cap C \cap A'$ **[B1]**

(d) 16 **[B2]**

M1 for correctly identifying $n(B \cap C) = 6$ or summing $12 + 4$.

[Back to Question] *[Total Marks: 6]*

25 **(a)** 7 **[B3]**

Partial working: Correct simplified equation $7x + 3 = 52$ **[M2]**

Partial working: Sets up unsimplified sum $(2x + 1) + x + (3x - 2) + (x + 4) = 52$ **[M1]**

(b) 33 **[B2]**

Partial working: Evaluates correct sub-regions for $x = 7$ (e.g. $15 + 7 + 11$) OR identifies correct algebraic expression $4x + 5$ **[M1]**

(c) Region strictly inside A but outside B is shaded (the left crescent) **[B1]**

[Back to Question] *[Total Marks: 6]*

26 **(a)** Intersection: x and Outer region: 4 **[B1]**

P only region: $28 - x$ AND C only region: $22 - x$ **[B1]**

(b) $(28 - x) + x + (22 - x) + 4 = 45$ oe **[M1]**

$x = 9$ **[A1]**

(c) 19 **[B1]**

[Back to Question] *[Total Marks: 5]*

27 **(a)** Summing all regions: $(x + 5) + 2x + x + 3 + 4 + 2 + 5 + (x + 1) = 45$ (or equivalent) **[M1]**

$5x + 20 = 45 \implies 5x = 25$ **[M1]**

$x = 5$ **[A1]**

(b) 34 **[B1]**

(c) Shading exactly the region representing P only (the top part of circle P, excluding the intersections with Q and R) **[B1]**

[Back to Question] *[Total Marks: 5]*

28 **(a)** 5 in outer region **[B1]**

8 in intersection **[B1]**

17 in F only AND 10 in S only **[B1]**

(b) 17 **[B1]**

(c) $\frac{27}{40}$ **[B2]**

M1 for $17 + 10$ or 27 seen **[M1]**

(d) $\frac{8}{25} \times \frac{7}{24}$ **[M1]**

$\frac{7}{75}$ **[A1]**

[Back to Question] *[Total Marks: 7]*

29 **(a)** Intersection of all three: 8 AND Outer region: 3 **[B1]**

Two-set only intersections: 12, 7, and 10 in correct regions **[B1]**

Single set regions: x (in C), $2x$ (in D), and x (in B) **[B1]**

(b) $x + 2x + x + 8 + 12 + 7 + 10 + 3 = 100$ oe **[M1]**

$x = 15$ **[A1]**

(c) Identifies the regions required: Only C + Only D + ($C \cap D$ only) OR uses $x + 2x + 12$ **[M1]**

57 **[A1]**

[Back to Question] *[Total Marks: 7]*

30 **(a)** 4 in center intersection **[B1]**

8, 5, 7 in the two-sport intersections **[B1]**

13, 9, 9 in the one-sport regions AND 5 in outer region **[B1]**

(b) 5 **[B1]**

(c) 17 **[B1]**

(d) $B \cap T' \cap S'$ or $B \cap (T \cup S)'$ **[B2]**

M1 for $B \cap \ldots$ **[M1]**

[Back to Question] *[Total Marks: 7]*

31 **(a)** $2x$ in the T only region **[B1]**

x in the intersection and $3x$ in the C only region **[B1]**

2 in the region outside both circles **[B1]**

(b) 8 **[B2]**

M1 for $2x + x + 3x + 2 = 50$ or $6x = 48$.

(c) 16 **[B1]**

(d) Region outside T shaded, as well as the entirety of C (the only unshaded region is 'T only') **[B2]**

B1 for shading T' correctly OR shading C correctly.

[Back to Question] *[Total Marks: 8]*

32 **(a)** Places 15 in the central intersection **[B1]**

Places 15 in $B \cap T$ only, 13 in $T \cap C$ only, and 7 in $B \cap C$ only **[B1]**

Places x, $2x$, $x + 5$, and 8 in the respective remaining regions (B only, T only, C only, and outside the sets) **[B1]**

(b) $x + 2x + (x + 5) + 15 + 15 + 13 + 7 + 8 = 103 \implies 4x + 63 = 103$ (or equivalent valid method) **[M1]**

$x = 10$ **[A1]**

(c) 85 **[B1]**

(d) $\dfrac{13}{63}$ **[B2]**

If 0 scored, allow [M1] for correctly identifying the denominator $n(T) = 63$ or the numerator $n(T \cap C \cap B') = 13$

[Back to Question] *[Total Marks: 8]*

33 **(a)** All 8 regions correctly populated **[B3]**

Partial working: 5 to 7 regions correctly populated **[B2]**

Partial working: 3 to 4 regions correctly populated (e.g. places $12, 15, x$ correctly) **[B1]**

(Correct values are: Center = 12. $M \cap D$ only = 13. $D \cap A$ only = 20. $M \cap A$ only = 16. M only = 9. D only = 20. A only = x. Outside = 15)

(b) 15 **[B2]**

Partial working: Forms correct sum $9 + 20 + x + 13 + 16 + 20 + 12 + 15 = 120$ OR $105 + x = 120$ **[M1]**

(c) 70 **[B2]**

Partial working: Identifies the 20 students in only Drama and Art OR correctly calculates $50 + 20$ **[M1]**

(d) Region strictly inside M but outside both D and A is shaded (the top crescent / "only M" region) **[B1]**

[Back to Question] *[Total Marks: 8]*

34 (a) 11 **[B1]**

(b) 61 **[B2]**

M1 for 64 or 3 seen.

[Back to Question] *[Total Marks: 3]*

35 (a) 21 **[B2]**

M1 for 13 or 8

(b) 80 **[B2]**

M1 for 16 or 5

[Back to Question] *[Total Marks: 4]*

36 (a) Calculates $3^3 = 27$ and $\sqrt{144} = 12$ **[M1]**

15 **[A1]**

(b) Calculates $2^4 = 16$ and $\sqrt[3]{125} = 5$ **[M1]**

80 **[A1]**

[Back to Question] *[Total Marks: 4]*

37 (a) 14 **[B1]**

(b) 59 **[B2]**

If 0 scored, M1 for 64 or 5 seen **[M1]**

(c) 5 **[B2]**

If 0 scored, M1 for $169 - 144$ or 25 seen **[M1]**

[Back to Question] *[Total Marks: 5]*

38 **(a)** 14 **[B1]**

(b) 48 **[B2]**

If 0 scored, allow [M1] for correctly evaluating either $\sqrt[3]{27} = 3$ or $2^4 = 16$

(c) 8.8 or $\frac{44}{5}$ **[B2]**

If 0 scored, allow [M1] for a correct numerator of 44 or a correct denominator of 5

[Back to Question] *[Total Marks: 5]*

39 **(a)** $\frac{37}{7}$ or $5\frac{2}{7}$ **[B2]**

M1 for $25 + 12$ or $8 - 1$ seen.

(b) 4 **[B2]**

M1 for 9×9 or $3^2 \times 3^2$ or 81 seen.

[Back to Question] *[Total Marks: 4]*

40 **(a)** Numerator correctly evaluated to $64 + 15 = 79$ **[B1]**

Denominator correctly evaluated to 4 **[B1]**

19.75 **[B1]**

(b) Substitution leading to $25 - (-8)$ **[M1]**

33 **[A1]**

[Back to Question] *[Total Marks: 5]*

41 **(a)** 5 **[B1]**

(b) 150 **[B2]**

If 0 scored, allow [M1] for 6×5^2 or 6×25

(c) 121 **[B2]**

If 0 scored, allow [M1] for evaluating $3^3 - 4^2 = 11$

[Back to Question] *[Total Marks: 5]*

42 **(a)** $3x + 4 = 64$ **[M1]**

$3x = 60$ **[M1]**

20 **[A1]**

(b) $x^2 = 64 + 36 = 100$ **[M1]**

$10, -10$ **[A1]**

(c) $3\sqrt{x} = 21$ or $\sqrt{x} = 7$ **[M1]**

49 **[A1]**

[Back to Question] *[Total Marks: 7]*

43 **(a)** 125 **[B2]**

If 0 scored, M1 for evaluating $16 - 9 - 2$ to get 5 **[M1]**

(b) -2 **[B3]**

M1 for $\sqrt{64}$ or 8 **[M1]**

M1 for $5 - 9$ or -4 **[M1]**

(c) 729 **[B3]**

M1 for $2\sqrt[3]{y} = 6$ or $\sqrt[3]{y} = 3$ **[M1]**

M1 for $y = 27$ **[M1]**

[Back to Question] *[Total Marks: 8]*

44 **(a)** 5.3990299... **[B2]**

M1 for numerator evaluated to 163.91... or denominator evaluated to 30.359...

(b) 5.399 **[B1]**

(c) 6 **[B2]**

M1 for evaluating $6^4 = 1296$ and $7^4 = 2401$

[Back to Question] *[Total Marks: 5]*

45 **(a)** **(i)** 1.49315... **[B2]**

If 0 scored, M1 for 3.329 seen or 72.25 and 68.921 **[M1]**

(ii) 1.49 **[B1]**

(b) 96 **[B2]**

If 0 scored, M1 for 23.04 or 0.24 seen **[M1]**

[Back to Question] *[Total Marks: 5]*

46 **(a)** 1.59 **[B2]**

M1 for 23.02... or 14.44 seen.

(b) 1734 **[B3]**

M1 for $\sqrt[3]{4913}$ or 17 seen.

M1 for 6×17^2.

[Back to Question] *[Total Marks: 5]*

47 **(a)** 1.5 **[B1]**

(b) Rearranges to $2x^4 = 1250$ leading to $x^4 = 625$ **[M1]**

$x = 5$ **[B1]**

$x = -5$ **[B1]**

(c) Correct substitution: $\sqrt{\dfrac{3(4.2)^2 + (-5.7)}{1.8}}$ **[M1]**

Evaluates numerator to 47.22 OR fraction to 26.2333... **[M1]**

5.12 **[A1]**

[Back to Question] *[Total Marks: 7]*

48 **(a)** 0.239 **[B2]**

If 0 scored, allow [M1] for evaluating the numerator as 19.679... or the denominator as 82.464...

(b) 64 **[B3]**

If 0 scored, allow [M1] for isolating x^2 correctly, e.g., $x^2 = 16^3$ or $x^2 = 4096$

Allow a further [M1] for $x = \sqrt{4096}$

(c) 7.44 or 7.443... **[B3]**

If 0 scored, allow [M1] for finding the volume of the cube as 12^3 or 1728

Allow a further [M1] for correctly rearranging the sphere volume formula to solve for r, e.g., $r^3 = \dfrac{1728 \times 3}{4\pi}$ or $r = \sqrt[3]{\dfrac{1296}{\pi}}$

[Back to Question] *[Total Marks: 8]*

49 **(a)** $\dfrac{5}{8}$ 64% 0.65 $\dfrac{2}{3}$ **[B2]**

M1 for correct conversions to decimals seen: 0.666..., 0.64, 0.625

(b) $\dfrac{17}{20}$ **[B1]**

[Back to Question] *[Total Marks: 3]*

50 **(a)** $\dfrac{7}{9}$ **[B1]**

(b) $\dfrac{36}{99}$ **[M1]**

$\dfrac{4}{11}$ **[A1]**

[Back to Question] *[Total Marks: 3]*

51 **(a)** $\dfrac{9}{20}$ **[B1]**

(b) 0.36 **[B1]**

(c) $\frac{204}{100}$ or $2\frac{4}{100}$ **[M1]**

$2\frac{1}{25}$ **[A1]**

[Back to Question] *[Total Marks: 4]*

52 **(a)** $\frac{9}{20}$ **[B1]**

(b) 0.375 **[B1]**

(c) 0.4 $\frac{5}{12}$ 42% $\frac{3}{7}$ **[B2]**

If 0 scored, allow [M1] for converting at least two of the fractions or percentages to decimals correctly, e.g., $0.416\ldots$, 0.42, $0.428\ldots$

[Back to Question] *[Total Marks: 4]*

53 **(a)** $\frac{75}{1000}$ or equivalent unsimplified fraction **[M1]**

$\frac{3}{40}$ **[A1]**

(b) 3.4 **[B1]**

(c) M1 for converting at least two of the other forms to decimals or percentages correctly (e.g. $\frac{5}{8} = 0.625$, $\frac{7}{11} = 0.636\ldots$, $62\% = 0.62$) **[M1]**

0.6 62% $\frac{5}{8}$ $\frac{7}{11}$ **[A1]**

[Back to Question] *[Total Marks: 5]*

54 Length multiplier $= \frac{4}{3}$ **[B1]**

Area multiplier $= \frac{7}{6}$ **[B1]**

$\frac{4}{3} \times$ Width multiplier $= \frac{7}{6}$ oe **[M1]**

Width multiplier $= \frac{7}{6} \times \frac{3}{4} = \frac{7}{8}$ **[M1]**

$x = 12.5$ **[A1]**

[Back to Question] *[Total Marks: 5]*

55 **(a)** 60 **[B3]**

M1 for $60\% = \frac{3}{5}$ or $\frac{1}{5} = 20\%$ seen

M1 for $\frac{3}{5} - \frac{1}{5} = \frac{2}{5}$ or $60\% - 20\% = 40\%$

M1 for $\frac{24}{2} \times 5$ or $\frac{24}{40} \times 100$

(b) 400 **[B3]**

M1 for $100\% - 30\% = 70\%$ or $0.7x$ seen

M1 for $\frac{3}{4} \times 0.7x = 210$ or $0.525x = 210$ or equivalent equation

[Back to Question] *[Total Marks: 6]*

56 **(a)** Total number of counters $= x + (2x + 5) + (3x - 1) = 6x + 4$ **[M1]**

$\frac{2x+5}{6x+4} = \frac{35}{100}$ or $\frac{7}{20}$ **[M1]**

$20(2x + 5) = 7(6x + 4)$ leading to $40x + 100 = 42x + 28$ **[M1]**

36 **[A1]**

(b) Substituting $x = 36$ to find green counters $= 3(36) - 1 = 107$ and total counters $= 6(36) + 4 = 220$ **[M1]**

$\frac{107}{220} \times 100$ **[M1]**

$48\frac{7}{11}$ **[A1]**

[Back to Question] *[Total Marks: 7]*

57 **(a)** $1000x = 245.4545\ldots$ and $10x = 2.4545\ldots$ or $100x = 24.5454\ldots$ and $x = 0.24545\ldots$ **[M1]**

$990x = 243$ or $99x = 24.3$ **[M1]**

$\frac{27}{110}$ **[A1]**

(b) Identifies that the denominator $42 = 2 \times 3 \times 7$ (or states that 42 has prime factors other than 2 and 5) **[B1]**

Terminating decimals can only have prime factors of 2 and/or 5 in their denominator. Since the fraction is in its simplest form, the factors of 3 and 7 do not cancel out, so it must recur. **[B1]**

(c) $\frac{10x+y}{99} = \frac{k}{11}$ or $\frac{9k}{99}$ (or states that the two-digit number xy is a multiple of 9) **[M1]**

9 **[A1]**

[Back to Question] *[Total Marks: 7]*

58 **(a)** States appropriate pair of equations to eliminate the recurring part, e.g., $100x = 27.77\ldots$ and $10x = 2.77\ldots$ **[M1]**

Subtracts to get $90x = 25$ leading to $x = \frac{25}{90}$, and correctly simplifies this to $\frac{5}{18}$ **[A1]**

(b) **(i)** 36 **[B3]**

If 0 scored, allow [M1] for calculating the fraction or percentage of the whole school for either boys or girls, e.g., $\frac{2}{5} \times 45\%$ or $\frac{3}{5} \times 30\%$

Allow [M1] for evaluating either as 18%

(ii) 400 **[B2]**

If 0 scored, allow [M1] for a correct algebraic setup or division, e.g., $\frac{144}{0.36}$ or $36\% = 144$

[Back to Question] *[Total Marks: 7]*

59 **(a)** 315 **[B1]**

(b) 27.5 **[B2]**

M1 for $\frac{3}{8} \times 100 = 37.5\%$ or for $100 - (35 + 37.5)$

(c) $\frac{9}{160}$ **[B2]**

M1 for $\frac{15}{100} \times \frac{3}{8}$ or $\frac{47.25}{840}$

[Back to Question] *[Total Marks: 5]*

60 **(a)** $100\% - 45\% - 37.5\% = 17.5\%$ oe **[M1]**

$\frac{140}{0.175}$ oe **[M1]**

800 **[A1]**

(b) Junior students $= \frac{3}{8} \times 800 = 300$ **[M1]**

$1.5G + G = 300$ oe **[M1]**

120 **[A1]**

[Back to Question] *[Total Marks: 6]*

61 **(a)** Converts at least two of the non-decimals to decimals correctly (e.g. 0.71 and 0.75) **[M1]**

$71\%, 0.714, \frac{5}{7}, \frac{3}{4}$ **[A1]**

(b) $\frac{1}{2}\left(\frac{2}{5} + \frac{5}{8}\right)$ or equivalent **[M1]**

$\frac{41}{80}$ **[A1]**

(c) $9x - 4x = 35$ or states the multiplier is $35 \div 5 = 7$ **[M1]**

$\frac{28}{63}$ **[A1]**

[Back to Question] *[Total Marks: 6]*

62 **(a)** $\frac{2}{5} = 40\%$ or 0.4, and total percentage of Sales and Production $= 40\% + 28\% = 68\%$ **[M1]**

$100\% - 68\% = 32\%$ and equating $0.32 \times$ Total $= 24$ or equivalent **[M1]**

75 **[A1]**

(b) Number of women in Sales $= 0.60 \times 30 = 18$ **[M1]**

Number of women in Production $= \frac{3}{7} \times 21 = 9$ (or men $= 12$, so women $= 9$) **[M1]**

Total women $= 18 + 9 + 12 = 39$ **[M1]**

52 **[A1]**

[Back to Question] *[Total Marks: 7]*

63 **(a)** 369 **[B2]**

If 0 scored, allow [M1] for 450×0.82 or 450×0.18

(b) 740 **[B3]**

If 0 scored, allow [M2] for $\frac{629}{1 - 0.15}$ or $\frac{629}{0.85}$

If 0 scored, allow [M1] for identifying that $85\% = \$629$

(c) 4 **[B2]**

If 0 scored, allow [M1] for a correct calculation showing the compound effect, e.g., $1.2 \times 0.8 = 0.96$ or choosing a starting value of 100 and reaching 96

[Back to Question] *[Total Marks: 7]*

64 (i) $>$ **[B1]**

(ii) $>$ **[B1]**

[Back to Question] *[Total Marks: 2]*

65 Converts to a common format (e.g. decimals: $0.625, 0.62, 0.615, 0.6$) **[M1]**

$\frac{3}{5}, 61.5\%, 0.62, \frac{5}{8}$ **[A1]**

[Back to Question] *[Total Marks: 2]*

66 **(a)** $>$ **[B1]**

(b) $=$ **[B1]**

(c) $<$ **[B1]**

[Back to Question] *[Total Marks: 3]*

67 **(a)** **(i)** $<$ **[B1]**

(ii) $=$ **[B1]**

(b) M1 for converting at least three values into a comparable form (e.g. percentages: 55%, 54.54%..., 54%, 55.55%...) **[M1]**

$54\%, \frac{6}{11}, 0.55, \frac{5}{9}$ **[A1]**

[Back to Question] *[Total Marks: 4]*

68 **(a)** **(i)** $=$ **[B1]**

(ii) $>$ **[B1]**

(iii) $>$ **[B1]**

(b) $0.3 \quad 33\% \quad \frac{1}{3} \quad \frac{3}{8}$ **[B2]**

If 0 scored, allow [M1] for converting at least two of the values correctly to decimals, e.g. 0.33, $0.333\ldots$, 0.375

[Back to Question] *[Total Marks: 5]*

69 **(a)** $p^2 \quad p \quad \sqrt{p} \quad \frac{1}{p}$ **[B2]**

M1 for evaluating the expressions with a chosen test value for p (e.g., $p = 0.25$) OR for any 3 expressions in the correct relative order.

(b) **(i)** $>$ **[B1]**

(ii) $<$ **[B1]**

[Back to Question] *[Total Marks: 4]*

70 **(a)** $x^2, x, \sqrt{x}, \frac{1}{x}$ **[B2]**

(B1 for 3 in correct relative order or if reversed)

(b) $x^{-2}, \frac{1}{x}, x, x^3$ **[B2]**

(B1 for 3 in correct relative order or if reversed)

[Back to Question] *[Total Marks: 4]*

71 **(a)** M1 for correctly calculating at least two values: $x^2 = 0.16$, $x^3 = -0.064$, $\frac{1}{x} = -2.5$ **[M1]**

$\frac{1}{x}, x, x^3, x^2$ **[A1]**

(b) M1 for squaring both terms to get 75 and 80, OR expressing as $\sqrt{75}$ and $\sqrt{80}$ **[M1]**

$<$ **[A1]**

(c) M1 for identifying a common denominator, e.g. cross multiplying to get $9m < 35$ **[M1]**

M1 for $35 < 9(m + 1)$ leading to $26 < 9m$ **[M1]**

3 **[A1]**

[Back to Question] *[Total Marks: 7]*

72 **(a)** $p^2 \quad p \quad \sqrt{p} \quad \frac{1}{p}$ **[B2]**

If 0 scored, allow [M1] for a correct order of any three of the terms or for successfully substituting a test value such as $p = \frac{1}{4}$ to evaluate all four expressions.

(b) **(i)** $<$ **[B1]**

(ii) $<$ **[B1]**

(c) c, b, a or $5^{20}, 3^{30}, 2^{50}$ **[A1]**

Shows index manipulation to obtain a common power, e.g. $(2^5)^{10}$, $(3^3)^{10}$ and $(5^2)^{10}$ **[M1]**

Evaluates the bases correctly as 32, 27 and 25 **[M1]**

[Back to Question] *[Total Marks: 7]*

73 $0.12 \quad 12.1\% \quad \frac{4}{33} \quad \sqrt{0.015}$ **[B2]**

M1 for correct decimal equivalents seen for at least two of the non-decimal values (e.g., 0.121, 0.1212..., 0.1224...).

[Back to Question] *[Total Marks: 2]*

74 Converts all to a common unit, e.g. grams: 0.52 g, 0.04 g, 4.5 g, 0.6 g **[M2]**

(M1 for 2 correct conversions to a common unit)

4.5×10^3 mg, 6×10^{-7} tonnes, 5.2×10^{-4} kg, 0.04 g **[A1]**

[Back to Question] *[Total Marks: 3]*

75 **(a)** B1 for at least two correct evaluated values ($a = 3100$, $b = 3130.49\ldots$, $c = 3125$, $d = 3136$) **[B1]**

B1 for all four evaluated correctly **[B1]**

d, b, c, a **[A1]**

(b) 15 **[B1]**

[Back to Question] *[Total Marks: 4]*

76 **(a)** $5^{-3} \quad 0.00805 \quad \frac{1}{120} \quad 8.5 \times 10^{-3}$ **[B2]**

If 0 scored, allow [M1] for evaluating at least two correctly as standard decimals: 0.008, 0.0085, 0.00833...

(b) $11.5 \leq m < 12.5$ **[B2]**

If 0 scored, allow [M1] for identifying the bounds 11.5 and 12.5

[Back to Question] *[Total Marks: 4]*

77 **(a)** Converts at least two values to decimals or percentages correctly (e.g. 0.875, 0.86, 0.833...) **[M1]**

$\frac{5}{6}$, 86%, 0.865, $\frac{7}{8}$ **[A1]**

(b) Finds a common denominator (e.g. 6, 12, etc.) **[M1]**

Any correct fraction (e.g. $\frac{5}{12}$, $\frac{2}{5}$) **[A1]**

[Back to Question] *[Total Marks: 4]*

78 **(a)** $<$ **[B1]**

(b) Evaluates at least two expressions correctly (0.25, 0.707..., 2) **[M1]**

a^2, a, $\sqrt{a}$, $\frac{1}{a}$ **[A1]**

(c) False **[B1]**

Provides a valid counterexample, e.g. let $p = 2$ and $q = -3$. Then $2 > -3$, but $\frac{1}{2} > -\frac{1}{3}$. **[B1]**

[Back to Question] *[Total Marks: 5]*

79 **(a)** 9 **[B1]**

(b) 25 **[B1]**

(c) 27 **[B2]**

If 0 scored, M1 for -4 or -3 seen **[M1]**

[Back to Question] *[Total Marks: 4]*

80 **(a)** 34 **[B2]**

M1 for $-2 \times -3 = 6$ or $12 - (-6) + 16$ or $18 + 16$ seen

(b) 1 **[B2]**

M1 for $\frac{1}{6} \times \frac{3}{2} = \frac{3}{12}$ or $\frac{1}{4}$ seen

[Back to Question] *[Total Marks: 4]*

81 **(a)** Evaluates multiplication and division correctly to get $+12$ and -5 **[M1]**

19 **[A1]**

(b) Correct division seen (e.g. $\frac{2}{5} \times \frac{7}{4}$ or $\frac{14}{20}$ or $\frac{7}{10}$) **[M1]**

$\frac{3}{5}$ **[A1]**

[Back to Question] *[Total Marks: 4]*

82 **(a)** $15 - 3 \times 6$ or $15 - 18$ **[M1]**

-3 **[A1]**

(b) $\frac{9}{4} \times \frac{8}{3}$ **[M1]**

6 **[A1]**

$4\frac{1}{2}$ or $\frac{9}{2}$ **[A1]**

[Back to Question] *[Total Marks: 5]*

83 **(a)** 28 **[B2]**

If 0 scored, allow [M1] for correctly evaluating the bracket to -4 or for showing $12 - (-16)$

(b) 30 **[B2]**

If 0 scored, allow [M1] for an equivalent fraction such as $\frac{120}{4}$ or $\frac{12}{0.4}$

(c) 3.5 **[B1]**

[Back to Question] *[Total Marks: 5]*

84 **(a)** $1\frac{13}{30}$ **[B3]**

If 0 scored, allow [M1] for correctly evaluating the multiplication as $\frac{18}{30}$ or $\frac{3}{5}$

Allow a further [M1] for finding a correct common denominator to add the fractions, e.g., $\frac{25}{30} + \frac{18}{30}$ or giving the improper fraction $\frac{43}{30}$

(b) 6 **[B3]**

If 0 scored, allow [M1] for correctly evaluating the bracket as $\frac{3}{8}$

Allow a further [M1] for converting to an improper fraction and correctly multiplying, e.g., $\frac{9}{4} \times \frac{8}{3}$

[Back to Question] *[Total Marks: 6]*

85 **(a)** $\frac{8}{6} - \frac{5}{6}$ or $\frac{1}{2}$ **[M1]**

$\frac{12}{5} \div$ their $\frac{1}{2}$ **[M1]**

$\frac{12}{5} \times \frac{2}{1}$ or equivalent multiplication **[M1]**

$4\frac{4}{5}$ **[A1]**

(b) 0.54 or $\frac{45 \times 12}{600}$ or equivalent **[M1]**

$54 \div 6$ or $0.54 \div 0.06$ **[M1]**

9 **[A1]**

[Back to Question] *[Total Marks: 7]*

86 Denominator: $2 - 0.4 = 1.6$ or $\frac{8}{5}$ **[B1]**

Right hand side: $-1\frac{1}{4} = -\frac{5}{4}$ **[B1]**

$x + \frac{1}{3} = -\frac{5}{4} \times \frac{8}{5}$ **[M1]**

$x + \frac{1}{3} = -2$ **[A1]**

$x = -\frac{7}{3}$ or $-2\frac{1}{3}$ **[A1]**

[Back to Question] *[Total Marks: 5]*

87 Converts mixed numbers to improper fractions and multiplies correctly: $\frac{9}{4} \times \frac{4}{3} = 3$ (or $\frac{12}{4}$ or $\frac{36}{12}$) **[M1]**

Evaluates the numerator correctly: $3\frac{2}{3} + 3 = 6\frac{2}{3}$ (or $\frac{20}{3}$) **[M1]**

Evaluates the denominator correctly using a common denominator: $\frac{27}{6} - \frac{11}{6} = \frac{16}{6} = \frac{8}{3}$ **[M1]**

Sets up the correct division for the final step: $\frac{20}{3} \div \frac{8}{3} = \frac{20}{3} \times \frac{3}{8}$ (or $\frac{20}{8}$) **[M1]**

$\frac{5}{2}$ or $2\frac{1}{2}$ **[A1]**

[Back to Question] *[Total Marks: 5]*

88 (a) $6\frac{2}{5}$ **[B4]**

M1 for changing mixed numbers to improper fractions, e.g., $\frac{5}{4}$ or $\frac{8}{3}$

M1 for writing fractions with a common denominator to subtract, e.g., $\frac{15}{12} - \frac{10}{12}$

M1 for $\frac{8}{3} \times \frac{12}{5}$ or $\frac{32}{5}$

(b) $38\frac{1}{3}$ **[B4]**

M1 for $83\frac{1}{3} \div 12\frac{1}{2}$

M1 for $\frac{250}{3} \times \frac{2}{25}$ or width $= \frac{20}{3}$ $(6\frac{2}{3})$

M1 for $2 \times \left(12\frac{1}{2} + \frac{20}{3}\right)$ or $2 \times \left(\frac{75}{6} + \frac{40}{6}\right)$

[Back to Question] *[Total Marks: 8]*

89 **(a)** Finds the temperature at 14:00 ($-4.5 + 11.2 = 6.7°C$) or sets up the full expression $(-4.5 + 11.2 - 8.1)$ **[M1]**

-1.4 **[A1]**

(b) Evaluates numerator to 19.63... or denominator to 5.115 **[M1]**

3.84 **[A1]**

[Back to Question] *[Total Marks: 4]*

90 Net rate of fill $= 0.75 - 0.45 = 0.3$ litres/min oe **[M1]**

Total time $= \frac{85.5}{0.3} = 285$ minutes **[M1]**

$285 \div 60 = 4$ with remainder 45 **[M1]**

4 hours 45 minutes **[A1]**

[Back to Question] *[Total Marks: 4]*

91 **(a)** 37.5 **[B2]**

M1 for 15×2.8 or 42 seen

(b) 23 **[B3]**

M1 for $37.5 - 1.2m = 9.9$ or $37.5 - 9.9$ or 27.6 seen

M1 for $27.6 \div 1.2$

[Back to Question] *[Total Marks: 5]*

92 **(a)** $\frac{2}{5} \times 1000$ or 400 **[M1]**

18.5×20 or 370 **[M1]**

770 **[A1]**

(b) Target volume $= 500$ or volume to lose $= 770 - 500 = 270$ **[M1]**

$270 \div 4.5$ **[M1]**

60 **[A1]**

(c) Volume to fill $= 1000 - 500 = 500$ **[M1]**

1 hour 20 minutes = 80 minutes **[M1]**

$500 \div 80$ leading to 6.25 **[A1]**

[Back to Question] *[Total Marks: 9]*

93 **(a)** Number of bottles = 360 **[B1]**

Volume left over = 0.5 **[B2]**

If 0 scored for the volume left over, allow [M1] for $450.5 - (360 \times 1.25)$

(b) 4 hours 54 minutes **[B4]**

If 0 scored, allow [M1] for calculating the volume of the pool: $24.5 \times 12 \times 1.8 = 529.2$ m^3

Allow a further [M1] for converting the volume to litres: $529.2 \times 1000 = 529200$ litres

Allow a further [M1] for calculating total minutes: $\frac{529200}{1800} = 294$ minutes

(c) -8.5 **[B3]**

If 0 scored, allow [M1] for correctly interpreting 6 hours 20 minutes as a fraction or decimal of hours, e.g., $6\frac{1}{3}$ hours or $\frac{19}{3}$ hours

Allow a further [M1] for calculating the total temperature rise: $1.5 \times \frac{19}{3} = 9.5$

[Back to Question] *[Total Marks: 10]*

94 **(a)** 1 **[B1]**

(b) 9 **[B1]**

(c) x^3 **[B1]**

[Back to Question] *[Total Marks: 3]*

95 **(a)** 5 **[B1]**

(b) $\frac{1}{36}$ **[B1]**

(c) t^8 **[B1]**

[Back to Question] *[Total Marks: 3]*

96 **(a)** m^7 **[B1]**

(b) $\frac{1}{25}$ or 0.04 **[B1]**

(c) 7 **[B1]**

[Back to Question] *[Total Marks: 3]*

97 **(a)** $\frac{1}{9}$ **[B1]**

(b) $\frac{2}{3}$ **[B1]**

(c) x^2 **[B1]**

(d) y^{12} **[B1]**

[Back to Question] *[Total Marks: 4]*

98 **(a)** $\frac{1}{8}$ **[B2]**

M1 for $(\sqrt[4]{16})^{-3}$ or $(2)^{-3}$ or $\dfrac{1}{16^{\frac{3}{4}}}$

(b) $4w^8y^{-2}$ or $\dfrac{4w^8}{y^2}$ **[B3]**

B1 for 4

B1 for w^8

B1 for y^{-2} or $\dfrac{1}{y^2}$

[Back to Question] *[Total Marks: 5]*

99 **(a)** $16x^6y^2$ **[B2]**

If 0 scored, allow [M1] for a product containing two of 16, x^6, y^2

(b) $\dfrac{3a^4}{b^3}$ or $3a^4b^{-3}$ **[B2]**

If 0 scored, allow [M1] for a term containing two of 3, a^4, b^{-3}

(c) $\dfrac{1}{9}$ **[B2]**

If 0 scored, allow [M1] for evaluating the cube root to 3 or for showing $\dfrac{1}{27^{\frac{2}{3}}}$ or 9

[Back to Question] *[Total Marks: 6]*

100 **(a)** **(i)** 16 **[B2]**

If 0 scored, M1 for $(\sqrt[3]{64})^2$ or 4^2 or $4096^{\frac{1}{3}}$ seen **[M1]**

(ii) $\dfrac{1}{16}$ or 0.0625 **[B1]**

(b) **(i)** $\dfrac{5a^4}{b^4}$ or $5a^4b^{-4}$ **[B2]**

If 0 scored, M1 for $\dfrac{15}{3} = 5$ or a^{6-2} or b^{3-7} **[M1]**

(ii) $2m^3n^{-1}$ or $\dfrac{2m^3}{n}$ **[B2]**

If 0 scored, M1 for $4^{\frac{1}{2}} = 2$ or $m^{6\times\frac{1}{2}}$ or $n^{-2\times\frac{1}{2}}$ **[M1]**

[Back to Question] *[Total Marks: 7]*

101 **(a)** $16^{\frac{1}{4}} = 2$ or equivalent correct first step **[M1]**

$\dfrac{1}{8}$ **[A1]**

(b) 25 **[B1]**

x^6y^{-4} or $\dfrac{x^6}{y^4}$ **[B1]**

(c) Expresses terms with a common base, e.g. $4^x = 2^{2x}$ or $32 = 2^5$ **[M1]**

Sets up a correct linear equation from indices, e.g. $3x - 1 + 2x = 5$ **[M1]**

$x = 1.2$ or $x = \frac{6}{5}$ **[A1]**

[Back to Question] *[Total Marks: 7]*

102 (a) $\frac{4y^4}{5x^2}$ **[B3]**

B1 for reciprocal $\left(\frac{16y^8}{25x^4}\right)^{\frac{1}{2}}$ seen

B1 for $\sqrt{16} = 4$ and $\sqrt{25} = 5$

B1 for $(y^8)^{\frac{1}{2}} = y^4$ and $(x^4)^{\frac{1}{2}} = x^2$

(b) -6 **[B4]**

M1 for changing to base 2: $4 = 2^2$ and $8 = 2^3$

M1 for $2^{3x-1} \times 2^{2(x+2)} = 2^{3(x-3)}$

M1 for equating powers: $3x - 1 + 2x + 4 = 3x - 9$

[Back to Question] *[Total Marks: 7]*

103 (a) Inverts fraction to remove negative power or correctly applies negative power to all terms, e.g. $\left(\frac{y^4}{16x^8}\right)^{\frac{1}{2}}$ or $\frac{16^{-\frac{1}{2}}x^{-4}}{y^{-2}}$ **[M1]**

Correctly evaluates the coefficient $16^{-\frac{1}{2}} = \frac{1}{4}$ or correctly halves powers **[M1]**

$\frac{y^2}{4x^4}$ or $0.25x^{-4}y^2$ **[A1]**

(b) Recognises that $4^x = 2^{2x}$ or substitutes $u = 4^x$ **[M1]**

Expresses 2^{2x+1} as 2×2^{2x} or $2(4^x)$ **[M1]**

Simplifies to a single exponential term, e.g. $-2^{2x} = -8$ or $u - 2u = -8$ **[M1]**

$x = 1.5$ or $x = \frac{3}{2}$ **[A1]**

[Back to Question] *[Total Marks: 7]*

104 (a) $\frac{27y^6}{8x^9}$ or $3.375y^6x^{-9}$ **[B3]**

If 0 scored, allow [M1] for taking the reciprocal to remove the negative power: $\left(\frac{81y^8}{16x^{12}}\right)^{\frac{3}{4}}$

Allow a further [M1] for finding the 4th root of the expression: $\left(\frac{3y^2}{2x^3}\right)^3$

(b) 6 **[B3]**

If 0 scored, allow [M1] for expressing both sides with a common base 2, e.g. $(2^3)^x = (2^2)^{x+3}$

Allow a further [M1] for equating the powers to set up the linear equation $3x = 2(x + 3)$

(c) 1 **[B3]**

If 0 scored, allow [M1] for expressing all terms as powers of 5, e.g. $5^1 \times (5^2)^k = 5^3$

Allow a further [M1] for correctly forming the equation for the indices $1 + 2k = 3$

[Back to Question] *[Total Marks: 9]*

105 **(a)** -1.5 or $-\frac{3}{2}$ **[B3]**

If 0 scored, M2 for $5^k = (5^3)^{-\frac{1}{2}}$ or $5^k = 5^{-\frac{3}{2}}$ **[M2]**

If 0 or M2 not scored, M1 for $\sqrt{125} = 125^{\frac{1}{2}}$ or $\frac{1}{125^{0.5}}$ or $125 = 5^3$ **[M1]**

(b) $\frac{27y^6}{8x^9}$ or $3.375y^6x^{-9}$ **[B3]**

If 0 scored, M2 for $\left(\frac{81y^8}{16x^{12}}\right)^{\frac{3}{4}}$ or $\frac{16^{-\frac{3}{4}}x^{-9}}{81^{-\frac{3}{4}}y^{-6}}$ **[M2]**

If 0 or M2 not scored, M1 for finding the 4th root: $\frac{2}{3}$, x^3, or y^2 correctly applied to at least one term **[M1]**

(c) 1.5 or $\frac{3}{2}$ **[B3]**

If 0 scored, M2 for $2x - 1 + 2x = 5$ or $4x - 1 = 5$ **[M2]**

If 0 or M2 not scored, M1 for writing with common base 2: $4^x = 2^{2x}$ or $32 = 2^5$ **[M1]**

[Back to Question] *[Total Marks: 9]*

106 **(a)** 4.5×10^5 **[B1]**

(b) 0.000308 **[B1]**

(c) 8×10^8 **[B1]**

[Back to Question] *[Total Marks: 3]*

107 **(a)** 4.52×10^5 **[B1]**

(b) 0.00607 **[B1]**

(c) 1.2×10^{10} **[B2]**

If 0 scored, M1 for 12×10^9 **[M1]**

[Back to Question] *[Total Marks: 4]*

108 **(a)** 7.14×10^{-3} **[B1]**

(b) 50200 **[B1]**

(c) 1.8×10^{10} **[B2]**

M1 for 18×10^9 or 18000000000 seen

[Back to Question] *[Total Marks: 4]*

109 **(a)** **(i)** 3.4×10^{-4} **[B1]**

(ii) 52000 **[B1]**

(b) 8×10^8 **[B2]**

If B0, M1 for $2 \times 4 = 8$ or $10^3 \times 10^5 = 10^8$ seen.

[Back to Question] *[Total Marks: 4]*

110 **(a)** 4.52×10^{-4} **[B1]**

(b) 371000 **[B1]**

(c) 8×10^7 **[B2]**

If 0 scored, allow [M1] for 80000000 or 8×10^k where $k \neq 7$

[Back to Question] *[Total Marks: 4]*

111 **(a)** $597 \times 10^{22} - 7.35 \times 10^{22}$ or $5.97 \times 10^{24} - 0.0735 \times 10^{24}$ **[M1]**

5.8965×10^{24} **[A1]**

(b) Sets up the correct division: $\dfrac{1.99 \times 10^{30}}{5.97 \times 10^{24}}$ **[M1]**

Evaluates correctly, e.g., $0.3333... \times 10^6$ **[M1]**

3.33×10^5 **[A1]**

[Back to Question] *[Total Marks: 5]*

112 **(a)** 4.5×10^2 **[B3]**

If 0 scored, allow [M1] for $\dfrac{3.6}{8} \times 10^3$ or 0.45×10^3

Allow a further [M1] for correctly converting to standard form, or for seeing 450

(b) 5.7×10^8 **[B2]**

If 0 scored, allow [M1] for $(50 \times 10^7) + (7 \times 10^7)$ or 570000000 or digits 57

(c) $3.2 \times 10^{2m+n+1}$ **[B2]**

If 0 scored, allow [M1] for $(4 \times 10^{2m}) \times (8 \times 10^n)$ or $32 \times 10^{2m+n}$

[Back to Question] *[Total Marks: 7]*

113 **(a)** 3×10^8 **[B3]**

If 0 scored, M1 for $5.4 \times 10^5 + 0.6 \times 10^5$ or $54 \times 10^4 + 6 \times 10^4$ or 6.0×10^5 seen **[M1]**

M1 for $\dfrac{\text{their } 6.0 \times 10^5}{2 \times 10^{-3}}$ **[M1]**

(b) 5×10^4 **[B2]**

If 0 scored, M1 for $\sqrt{25 \times 10^8}$ or 50000 seen **[M1]**

(c) $5 \times 10^{2x-y-1}$ **[B3]**

If 0 scored, M1 for $p^2 = 4 \times 10^{2x}$ **[M1]**

M1 for $\dfrac{4 \times 10^{2x}}{8 \times 10^y}$ or $0.5 \times 10^{2x-y}$ **[M1]**

[Back to Question] *[Total Marks: 8]*

114 **(a)** $4 \times 10^{m+n+1}$ or $4.0 \times 10^{m+n+1}$ **[B2]**

M1 for $40 \times 10^{m+n}$ seen

(b) $3.125 \times 10^{2m-n}$ **[B3]**

M1 for $(5 \times 10^m)^2$ or 25×10^{2m} seen

M1 for $\dfrac{25}{8} \times 10^{2m-n}$

(c) 2 **[B3]**

M1 for $8 \times 10^n = 5.08 \times 10^m - 5 \times 10^m$

M1 for $8 \times 10^n = 0.08 \times 10^m$ or $8 \times 10^n = 8 \times 10^{m-2}$

[Back to Question] *[Total Marks: 8]*

115 **(a)** 500 **[B2]**

M1 for $1.5 \times 10^8 \div 3.0 \times 10^5$

(b) 6.0×10^3 or 6×10^3 **[B2]**

M1 for $1.5 \times 10^8 \div 2.5 \times 10^4$ or 6000 seen

[Back to Question] *[Total Marks: 4]*

116 **(a)** $\dfrac{1.99 \times 10^{30}}{1.90 \times 10^{27}}$ **[M1]**

$1047.36\ldots$ or $1.047\cdots \times 10^3$ **[A1]**

1.05×10^3 **[A1]**

(b) $(4.5 \times 10^{13}) \times 1000$ or $(4.5 \times 10^{13}) \times 10^3$ **[M1]**

4.5×10^{16} **[A1]**

[Back to Question] *[Total Marks: 5]*

117 **(a)** 3.61×10^7 **[B2]**

If B0, M1 for $3.4 \times 10^7 + 0.21 \times 10^7$ or $34 \times 10^6 + 2.1 \times 10^6$ or 36100000 **[M1]**

(b) Converts minutes to seconds: $5 \times 60 = 300$ **[M1]**

Sets up division: $\dfrac{300}{4 \times 10^{-9}}$ **[M1]**

7.5×10^{10} **[A1]**

[Back to Question] *[Total Marks: 5]*

118 **(a)** 6.04×10^{24} **[B2]**

If 0 scored, allow [M1] for 6.0435×10^{24} or 604.35×10^{22} or digits 604

(b) 81 **[B2]**

If 0 scored, allow [M1] for $\dfrac{5.97 \times 10^{24}}{7.35 \times 10^{22}}$ or 81.2...

(c) 1.79 **[B2]**

If 0 scored, allow [M1] for $300 \times 5.97 \times 10^{24}$ or 1.791×10^{27} or digits 179

[Back to Question] *[Total Marks: 6]*

119 **(a)** 5.8965×10^{24} **[B2]**

If 0 scored, M1 for $5.97 \times 10^{24} - 0.0735 \times 10^{24}$ or $597 \times 10^{22} - 7.35 \times 10^{22}$ seen **[M1]**

(b) 5×10^2 or 5.00×10^2 **[B2]**

If 0 scored, M1 for $\dfrac{1.50 \times 10^8}{3.00 \times 10^5}$ or 500 **[M1]**

(c) 5.10×10^{14} **[B3]**

If 0 scored, M2 for $4 \times \pi \times (6.37 \times 10^6)^2$ **[M2]**

If 0 or M2 not scored, M1 for $(6.37 \times 10^6)^2$ or $4.057... \times 10^{13}$ **[M1]**

[Back to Question] *[Total Marks: 7]*

120 **(a)** $\pi \times (4.5 \times 10^2)^2 \times (1.2 \times 10^3)$ **[M1]**

$7.63407\cdots \times 10^8$ **[A1]**

7.634×10^8 **[A1]**

(b) Volume in $m^3 = \dfrac{7.634 \times 10^8}{10^6} = 763.4\ m^3$ (or equivalent rate conversion $1.5 \times 10^2\ cm^3/\text{s}$) **[M1]**

Time in seconds $= \dfrac{763.4}{1.5 \times 10^{-4}} = 5.089\cdots \times 10^6$ **[M1]**

Time in hours $= \dfrac{5.089\cdots \times 10^6}{3600} = 1413.7\ldots$ **[M1]**

1.41×10^3 **[A1]**

[Back to Question] *[Total Marks: 7]*

121 **(a)** **(i)** 47.54 **[B1]**

(ii) 48 **[B1]**

(iii) 48 **[B1]**

[Back to Question] *[Total Marks: 3]*

122 **(a)** 780 **[B1]**

(b) 0.036 **[B1]**

(c) 4.10 **[B1]**

[Back to Question] *[Total Marks: 3]*

123 **(a)** 48.98 **[B1]**

(b) 0.0031 **[B1]**

(c) 600 **[A1]**

M1 for $\frac{8 \times 30}{0.4}$

[Back to Question] *[Total Marks: 4]*

124 **(a)** **(i)** 34.57 **[B1]**

(ii) 0.070 **[B1]**

(b) $\frac{60 \times 4}{0.5}$ **[M1]**

480 **[B1]**

[Back to Question] *[Total Marks: 4]*

125 **(a)** 0.048 **[B1]**

(b) 380 **[B1]**

(c) $\frac{60 \times 4}{20 - 3}$ or $\frac{240}{17}$ **[M1]**

14 or 14.1 (from $\frac{240}{17}$) or 15 (if 17 is estimated as 20) **[A1]**

[Back to Question] *[Total Marks: 4]*

126 **(a)** **(i)** 203.50 **[B1]**

(ii) 203 **[B1]**

(iii) 200 **[B1]**

(b) 560 **[A1]**

If 0 scored, M1 for $\frac{7 \times 40}{0.5}$ **[M1]**

[Back to Question] *[Total Marks: 5]*

127 $\frac{60 \times 3}{0.2}$ **[M1]**

$\frac{180}{0.2}$ **[M1]**

900 **[A1]**

[Back to Question] *[Total Marks: 3]*

128 **(a)** $\frac{8 \times 30}{0.4}$ **[M1]**

600 **[A1]**

(b) $\sqrt{100}$ or 10, and 20, and 0.5 **[M1]**

$\frac{10 + 20}{0.5^2}$ or $\frac{30}{0.25}$ **[M1]**

120 **[A1]**

[Back to Question] *[Total Marks: 5]*

129 **(a)** $\frac{\sqrt{100} \times 4^2}{0.2}$ **[M1]**

$\frac{10 \times 16}{0.2}$ or $\frac{160}{0.2}$ **[M1]**

800 **[B1]**

(b) 500×200 **[M1]**

100000 **[B1]**

[Back to Question] *[Total Marks: 5]*

130 **(a)** 2000 **[A3]**

If 0 scored, M1 for rounding the numbers to 4×10^6, 0.05 and 20 **[M1]**

M1 for evaluating the numerator to 2000 and the denominator to 1 **[M1]**

(b) Greater (than) **[B1]**

Valid reason, e.g. the numerator was rounded up and the denominator was rounded down. **[B1]**

[Back to Question] *[Total Marks: 5]*

131 **(a)** 1600 **[A1]**

M1 for $\frac{\sqrt[3]{8} \times 20^2}{0.5}$

M1 for 2×400 or 800 seen in numerator

(b) 3.2×10^6 **[A1]**

M1 for $4 \times 10^5 \times 2^3$

M1 for 32×10^5 or 3200000 seen

[Back to Question] *[Total Marks: 6]*

132 **(a)** $\sqrt{100} \times 4^3$ or 10×64 $(= 640)$ **[M1]**

0.2×50 $(= 10)$ **[M1]**

$\frac{10 \times 64}{10} = 64$ **[A1]**

(b) Uses $a = 4$, $x = 0.05$, $b = 1$, $c = 0.8$ **[M1]**

$\frac{4 \times 0.05}{1 - 0.8}$ or $\frac{0.2}{0.2}$ **[M1]**

1 **[A1]**

[Back to Question] *[Total Marks: 6]*

133 **(a)** 40, 5, and 1000 seen **[M1]**

$\frac{40 \times 5^2}{\sqrt[3]{1000}}$ or $\frac{40 \times 25}{10}$ **[M1]**

100 **[A1]**

(b) Radius ≈ 4 and Height ≈ 10 **[B1]**

Volume $= 3 \times 4^2 \times 10$ **[M1]**

480 **[A1]**

(c) Distance ≈ 4000 and Speed ≈ 600 **[B1]**

$\frac{4000}{600}$ or $\frac{20}{3}$ or 6.66... hours **[M1]**

6 hours and 40 minutes **[A1]**

[Back to Question] *[Total Marks: 9]*

134 **(a)** $125 \times 74 = 9250$ **[M1]**

9300 **[A1]**

(b) $x^2 = 9250$ **[M1]**

$x = \sqrt{9250}$ (96.176...) **[M1]**

96.2 **[A1]**

[Back to Question] *[Total Marks: 5]*

135 **(a)** 404 **[A2]**

If 0 scored, M1 for $\frac{1}{3} \times \pi \times 5.82^2 \times 11.4$ **[M1]**

(b) 360 **[A2]**

If 0 scored, M1 for $\frac{1}{3} \times 3 \times 6^2 \times 10$ **[M1]**

(c) 70 **[A2]**

If 0 scored, M1 for $850 \div 12$ or 70.83... seen **[M1]**

[Back to Question] *[Total Marks: 6]*

136 **(a)** 87.1 **[B2]**

M1 for 87.098... seen or $\frac{78.38...}{0.9}$

(b) **(i)** 27 **[A1]**

M1 for $\frac{4}{3} \times \pi \times 1.85^3$ or 26.5... seen

(ii) 35 hours 22 minutes **[A1]**

M1 for their exact volume (or 27) $\times 1000$ (e.g. 26522 litres)

M1 for $\frac{\text{their } 26522}{12.5}$ $(= 2121.76$ mins$)$

[Back to Question] *[Total Marks: 7]*

137 **(a)** **(i)** 70.35465... **[B1]**

Accept an answer containing at least 5 decimal places.

(ii) 70.4 **[B1]**

(b) $\frac{30^2 - 10^2}{\sqrt{100}}$ or $\frac{900 - 100}{10}$ **[M1]**

80 **[B1]**

[Back to Question] *[Total Marks: 4]*

138 **(a)** $r \approx 4$, $h \approx 20$, density ≈ 8 **[M1]**

Estimated Mass $= 3 \times 4^2 \times 20 \times 8$ **[M1]**

7680 **[A1]**

(b) Volume $= \pi \times (4.15)^2 \times 19.8 = 1071.3\ldots$ **[M1]**

Mass $= 1071.3\cdots \times 8.13 = 8710.19\ldots$ **[M1]**

8710 **[A1]**

[Back to Question] *[Total Marks: 6]*

139 **(a)** $3 : 5 : 7$ **[B1]**

(b) 80 **[B2]**

If 0 scored, M1 for $\frac{240}{8+10+12} \times 10$ **[M1]**

[Back to Question] *[Total Marks: 3]*

140 **(a)** M1 for 1.2 kg = 1200 g or 450 g = 0.45 kg

$3 : 8$ **[A1]**

(b) M1 for $3 + 4 + 5$ or 12 seen

140 **[A1]**

[Back to Question] *[Total Marks: 4]*

141 **(a)** Total parts $= 2 + 3 + 5 = 10$ **[M1]**

Ben's share $= \frac{3}{10} \times 840 = 252$ **[A1]**

(b) $\frac{5}{10}$ **[M1]**

$\frac{1}{2}$ **[A1]**

[Back to Question] *[Total Marks: 4]*

142 **(a)** $3 + 4 + 5 = 12$ **[M1]**

$\frac{5}{12} \times 120 = 50$ **[A1]**

(b) $\frac{3}{12} \times 120$ or 3×10 **[M1]**

30 **[A1]**

(c) $4 : 5$ **[B1]**

[Back to Question] *[Total Marks: 5]*

143 **(a)** $45 : 120$ **[M1]**

$3 : 8$ **[A1]**

(b) $3.6 \times 100 = 360$ cm **[M1]**

$\frac{360}{2+3+7} \times 7$ **[M1]**

210 **[A1]**

[Back to Question] *[Total Marks: 5]*

144 **(a)** $\dfrac{m+30}{f} = 3$ or $m + 30 = 3f$ **[B1]**

$\dfrac{m}{f+20} = 1$ or $m = f + 20$ **[B1]**

(b) M1 for substituting to eliminate one variable, e.g., $f + 20 + 30 = 3f$ or $m + 30 = 3(m - 20)$

M1 for correctly simplifying their equation to $2f = 50$ or $2m = 90$

$m = 45$, $f = 25$ **[A1]**

(c) 9 : 5 **[B1]**

[Back to Question] *[Total Marks: 6]*

145 **(a)** 15 : 20 : 8 **[B2]**

If 0 scored, M1 for $3 \times 5 : 4 \times 5$ or $5 \times 4 : 2 \times 4$ **[M1]**

(b) 24 **[B2]**

If 0 scored, M1 for $\dfrac{129}{15 + 20 + 8} \times 8$ **[M1]**

(c) 6 **[B3]**

If 0 scored, M1 for $\dfrac{60 - y}{24 - y} = 3$ or $\dfrac{60 - y}{24 - y} = \dfrac{3}{1}$ **[M1]**

M1 for $60 - y = 72 - 3y$ (or equivalent linear equation) **[M1]**

[Back to Question] *[Total Marks: 7]*

146 **(a)** Actual distance $= 8.4 \times 250\,000 = 2\,100\,000$ cm **[M1]**

Distance in m $= 21\,000$ or distance in km $= \dfrac{2\,100\,000}{100\,000}$ **[M1]**

21 **[A1]**

(b) Area scale factor $= 1^2 : (250\,000)^2 = 1 : 6.25 \times 10^{10}$ **[M1]**

35 km^2 $= 35 \times 10^{10}$ cm^2 **[M1]**

Map area $= \dfrac{35 \times 10^{10}}{6.25 \times 10^{10}} = 5.6$ **[A1]**

[Back to Question] *[Total Marks: 6]*

147 **(a)** 16.4×250000 or 4100000 **[M1]**

41 **[A1]**

(b) Scale factor for area is $(250000)^2$ or 1 km $= 10^5$ cm so 1 km^2 $= 10^{10}$ cm^2 **[M1]**

$\dfrac{35 \times 10^{10}}{250000^2}$ or $\dfrac{35}{2.5^2}$ **[M1]**

5.6 **[A1]**

(c) 400000 **[B1]**

[Back to Question] *[Total Marks: 6]*

148 **(a)** 29 **[B2]**

If 0 scored, M1 for $14.5 \times 1000 \times 100 \div 50000$ or equivalent **[M1]**

(b) 3.1 **[B2]**

If 0 scored, M1 for $12.4 \times \left(\frac{50000}{100000}\right)^2$ or 12.4×0.25 **[M1]**

(c) Calculates requires 360 g of flour and 225 g of butter **[B2]**

If B0 scored, M1 for $\frac{240}{8} \times 12$ or $\frac{150}{8} \times 12$ **[M1]**

No, because he needs 225 g of butter but only has 200 g **[B1]**

[Back to Question] *[Total Marks: 7]*

149 **(a)** $16.8 \times 250\,000 = 4\,200\,000$ cm **[M1]**

42 **[A1]**

(b) Scale of area $= 1^2 : (2.5)^2$ OR 1 cm^2 represents 6.25 km^2 **[M1]**

8.96 **[A1]**

(c) 14 cm^2 represents 56 km^2 $\implies$ 1 cm^2 represents 4 km^2 **[M1]**

Linear ratio $= \sqrt{1}$ cm represents $\sqrt{4}$ km $\implies$ 1 cm to 2 km **[M1]**

200 000 **[A1]**

[Back to Question] *[Total Marks: 7]*

150 **(a)** M1 for $5 - 3 = 2$ parts

M1 for $84 \div 2$ or 42 seen

336 **[A1]**

(b) **(i)** M1 for 8.5×25000 or 212500 seen

2.125 **[A1]**

(ii) M1 for 1 cm represents 0.25 km or 1 km represents 4 cm

M1 for area scale factor 1 cm^2 represents 0.0625 km^2 or $3.5 \times 10^{10} \div 25000^2$

56 **[A1]**

[Back to Question] *[Total Marks: 8]*

151 **(a)** Let total workers be $8x$. Males $= 5x$, Females $= 3x$. **[M1]**

Full-time males $= \frac{3}{5} \times 5x = 3x$. Part-time males $= \frac{2}{5} \times 5x = 2x$. **[M1]**

Full-time females $= \frac{4}{5} \times 3x = 2.4x$. Part-time females $= \frac{1}{5} \times 3x = 0.6x$. **[M1]**

Total full-time $= 3x + 2.4x = 5.4x$. Total part-time $= 2x + 0.6x = 2.6x$. Ratio is $5.4 : 2.6 = 27 : 13$. **[A1]**

(b) Difference $= 3x - 0.6x = 2.4x$ **[M1]**

$2.4x = 42 \implies x = 17.5$ **[M1]**

Total workers $= 8x = 8 \times 17.5 = 140$ **[A1]**

[Back to Question] *[Total Marks: 7]*

152 **(a)** $5 - 3 = 2$ parts represent 42 workers **[M1]**

1 part is $\dfrac{42}{2} = 21$ **[M1]**

$8 \times 21 = 168$ **[A1]**

(b) Day shift workers: $5 \times 21 = 105$. Night shift workers: $3 \times 21 = 63$ **[M1]**

Day shift males: $\dfrac{3}{7} \times 105 = 45$. Day shift females: $105 - 45 = 60$ **[M1]**

Night shift males: $\dfrac{2}{9} \times 63 = 14$. Night shift females: $63 - 14 = 49$ **[M1]**

Total males $= 45 + 14 = 59$. Total females $= 60 + 49 = 109$. Ratio is $59 : 109$ **[A1]**

[Back to Question] *[Total Marks: 7]*

153 **(a)** Alex : Ben $= 15 : 20$ and Ben : Chris $= 20 : 24$ **[M1]**

$15 : 20 : 24$ **[A1]**

(b) Difference in parts $= 24 - 15 = 9$ parts $\implies$ 9 parts $= 16\,200$ **[M1]**

Total parts $= 15 + 20 + 24 = 59$ **[M1]**

$P = \dfrac{16\,200}{9} \times 59 = 106\,200$ **[A1]**

(c) Alex's share $= \dfrac{15}{59} \times 106\,200 = 27\,000$ **[B1]**

$27\,000 \times \left(1 + \dfrac{4.2}{100}\right)^5$ **[M1]**

33 167 **[A1]**

[Back to Question] *[Total Marks: 8]*

154 Time taken $= 1.5$ hours (or 1 hour 30 minutes) **[B1]**

$120 \div 1.5 = 80$ **[B1]**

[Back to Question] *[Total Marks: 2]*

155 Time $= 1.5$ hours **[B1]**

Average Speed $= \dfrac{120}{1.5}$ **[M1]**

80 **[A1]**

[Back to Question] *[Total Marks: 3]*

156 **(a)** M1 for $108 \div 1.5$ or $\frac{108}{90} \times 60$

72 **[A1]**

(b) M1 for $72 \times \frac{1000}{3600}$ or $72 \div 3.6$

20 **[A1]**

[Back to Question] *[Total Marks: 4]*

157 **(a)** $\frac{120}{1.5}$ or $\frac{120}{90} \times 60$ **[M1]**

80 **[A1]**

(b) $\frac{8}{100} \times 350$ or 8×3.5 **[M1]**

28 **[A1]**

[Back to Question] *[Total Marks: 4]*

158 **(a)** Time $= 2.25$ or $\frac{9}{4}$ hours **[M1]**

$180 \div 2.25 = 80$ **[A1]**

(b) Time in seconds $= 450 \div 1.5 = 300$ **[M1]**

$300 \div 60 = 5$ **[A1]**

[Back to Question] *[Total Marks: 4]*

159 **(a)** M1 for $\frac{30}{v} + \frac{40}{v+5} = 4$

M1 for $30(v+5) + 40v = 4v(v+5)$ (removing fractions)

A1 for obtaining $2v^2 - 25v - 75 = 0$ with no errors or omitted steps in working

(b) M2 for $(2v+5)(v-15) = 0$ or correct substitution into the quadratic formula
(M1 for $(2v+a)(v+b) = 0$ where $ab = -75$ or $2b + a = -25$)

$v = 15$ and $v = -2.5$ **[A1]**

(c) 2 **[B1]**

[Back to Question] *[Total Marks: 7]*

160 Time $= 4\frac{25}{60}$ hours or 4.416... hours **[M1]**

Average Speed $= \frac{385}{4.416...}$ **[M1]**

87.2 (correct to 1 decimal place) **[A1]**

[Back to Question] *[Total Marks: 3]*

161 **(a)** 45×1000 or $\frac{45}{60}$ **[M1]**

$\frac{45000}{60} = 750$ **[A1]**

(b) Base area $= 150 \times 80 = 12000$ cm^2 **[M1]**

Rate of height increase $= \frac{\text{Volume per second}}{\text{Base area}} = \frac{750}{12000}$ **[M1]**

0.0625 **[A1]**

[Back to Question] *[Total Marks: 5]*

162 **(a)** $\frac{350}{100} \times 6.4$ **[M1]**

22.4 **[A1]**

(b) Cost in euros $= 22.4 \times 1.35 = 30.24$ **[M1]**

Cost in dollars $= 30.24 \div 0.85$ **[M1]**

35.58 **[A1]**

[Back to Question] *[Total Marks: 5]*

163 **(a)** M1 for 52×1000 or 52000 seen

M1 for their $52000 \div 1.6$ (implies 32500 seconds)

9 hours 2 minutes **[A1]**

(b) M1 for 52×2.15 or 111.8 seen

M1 for their 111.8×0.84

93.91 **[A1]**

[Back to Question] *[Total Marks: 6]*

164 **(a)** Fuel used $= \frac{325}{100} \times 8.4$ or 27.3 **[M1]**

27.3×1.35 **[M1]**

36.855 or 36.86 **[A1]**

(b) 24 minutes $= 0.4$ hours or $\frac{2}{5}$ hours **[B1]**

Distance $= 75 \times 4.4$ **[M1]**

330 **[A1]**

[Back to Question] *[Total Marks: 6]*

165 **(a)** Volume in $m^3 = 1.8 \times 1.5 \times 1.2 = 3.24$ **[M1]**

$3.24 \times 1000 = 3240$ litres **[A1]**

(b) Time in seconds $= \dfrac{3240}{2.5} = 1296$ **[M1]**

Time in minutes $= \dfrac{1296}{60} = 21.6$ **[A1]**

(c) Rate $= \dfrac{\text{Volume}}{\text{Time}} = \dfrac{3240}{45}$ **[M1]**

$x = 72$ **[A1]**

[Back to Question] *[Total Marks: 6]*

166 **(a)** Volume $= 2.5 \times 1.4 \times 1.2$ or 4.2 (m^3) **[M1]**

Volume in litres $= 4.2 \times 1000$ or 4200 **[M1]**

Time in seconds $= \dfrac{4200}{0.35}$ or 12000 **[M1]**

3 h 20 min **[A1]**

(b) $\dfrac{780}{0.84}$ or 928.57... **[M1]**

$928.57 - 850$ **[M1]**

78.57 **[A1]**

[Back to Question] *[Total Marks: 7]*

167 **(a)** Time A to B $= \dfrac{d}{v}$ **[M1]**

Time B to C $= \dfrac{2d}{v+15}$ **[M1]**

Total Time $T = \dfrac{d}{v} + \dfrac{2d}{v+15} = \dfrac{d(v+15)+2dv}{v(v+15)} = \dfrac{dv+15d+2dv}{v(v+15)} = \dfrac{3dv+15d}{v(v+15)}$ **[A1]**

(b) Substitute $d = 120$ and $T = 4.5$ into the equation from part (a): $4.5 = \dfrac{3(120)v + 15(120)}{v(v+15)}$ **[M1]**

$4.5(v^2 + 15v) = 360v + 1800$ **[M1]**

$4.5v^2 + 67.5v - 360v - 1800 = 0$ **[M1]**

$4.5v^2 - 292.5v - 1800 = 0 \implies divideby1.5toget3\text{v}^2 - 195v - 1200 = 0$ **[A1]**

[Back to Question] *[Total Marks: 7]*

168 **(a)** $\dfrac{2400}{v}$ **[B1]**

(b) $\dfrac{2400}{v} + \dfrac{2400}{v+40} = 10.5$ **[M1]**

$2400(v+40) + 2400v = 10.5v(v+40)$ **[M1]**

$2400v + 96000 + 2400v = 10.5v^2 + 420v$ **[M1]**

$10.5v^2 - 4380v - 96000 = 0$ **[A1]**

Multiply by $\frac{2}{3}$ to get $7v^2 - 2920v - 192000 = 0$ **[A1]**

(c) $v = \frac{-(-2920) \pm \sqrt{(-2920)^2 - 4(7)(-192000)}}{2(7)}$ **[M1]**

$v = 475.25...$ or $v = -57.68...$ (discard negative) **[A1]**

475 (correct to nearest integer) **[A1]**

[Back to Question] *[Total Marks: 7]*

169 **(a)** $\frac{30}{x} + \frac{18}{x-4} = 2.5$ **[M1]**

$\frac{30(x-4) + 18x}{x(x-4)} = 2.5$ **[M1]**

$48x - 120 = 2.5(x^2 - 4x)$ **[M1]**

$48x - 120 = 2.5x^2 - 10x \implies 2.5x^2 - 58x + 120 = 0 \implies 5x^2 - 116x + 240 = 0$ **[A1]**

(b) $x = \frac{-(-116) \pm \sqrt{(-116)^2 - 4(5)(240)}}{2(5)}$ **[M1]**

$x = \frac{116 \pm \sqrt{13456 - 4800}}{10} = \frac{116 \pm \sqrt{8656}}{10}$ **[M1]**

$x = 20.90$ or $x = 2.30$ **[A1]**

[Back to Question] *[Total Marks: 7]*

170 **(a)** $\frac{15}{x} + \frac{12}{x-4} = 3.5$ **[M1]**

Multiply by $2x(x-4)$ to clear denominators (or equivalent): $30(x-4) + 24x = 7x(x-4)$ **[M1]**

$54x - 120 = 7x^2 - 28x \implies 7x^2 - 82x + 120 = 0$ **[A1]**

(b) **(i)** $(7x - 12)(x - 10) = 0$ OR $\frac{82 \pm \sqrt{(-82)^2 - 4(7)(120)}}{14}$ **[M2]**

If M0, allow M1 for $(7x \pm a)(x \pm b) = 0$ where $ab = 120$ or $7b \pm a = -82$

$x = 10$, $x = \frac{12}{7}$ (or 1.71) **[A1]**

(ii) Identifies valid speed $x = 10$ (since $x - 4 > 0 \implies x > 4$) **[B1]**

Time $= \frac{15}{10} = 1.5$ **[B1]**

[Back to Question] *[Total Marks: 8]*

171 **(a)** $\frac{150}{x}$ **[B1]**

(b) $\frac{150}{x} + \frac{90}{x+15} = 3$ **[M1]**

$\frac{150(x+15) + 90x}{x(x+15)} = 3$ or $150(x+15) + 90x = 3x(x+15)$ **[M1]**

$150x + 2250 + 90x = 3x^2 + 45x$ **[M1]**

$3x^2 - 195x - 2250 = 0$ leading to $x^2 - 65x - 750 = 0$ **[A1]**

(c) $(x - 75)(x + 10) = 0$ or $\dfrac{65 \pm \sqrt{(-65)^2 - 4(1)(-750)}}{2}$ **[M2]**

If M0 scored, allow [M1] for $(x \pm 75)(x \pm 10) = 0$ or for correct substitution into the quadratic formula.

75 **[A1]**

[Back to Question] *[Total Marks: 8]*

172 Profit $= \$400 - \$320 = \$80$ or $\dfrac{400}{320} \times 100$

[M1]

25 **[A1]**

[Back to Question] *[Total Marks: 2]*

173 (a) $\dfrac{15}{100} \times 80$ or 12 **[M1]**

68 **[A1]**

(b) $\dfrac{4000 - 3200}{4000} \times 100$ or $\dfrac{800}{4000} \times 100$ **[M1]**

20 **[A1]**

[Back to Question] *[Total Marks: 4]*

174 (a) $\dfrac{24}{60} \times 100$ **[M1]**

40 **[A1]**

(b) $\dfrac{15}{100} \times 80 = 12$ OR 80×0.85 **[M1]**

68 **[A1]**

[Back to Question] *[Total Marks: 4]*

175 (a) $\dfrac{15}{100} \times 320$ or $10\% = 32, 5\% = 16$ **[M1]**

48 **[A1]**

(b) $\dfrac{20}{100} \times 45 = 9$ or $\dfrac{80}{100} \times 45$ **[M1]**

$45 - 9$ (if first method used) **[M1]**

36 **[A1]**

[Back to Question] *[Total Marks: 5]*

176 **(a)** Discount $= 0.15 \times 120 = 18$ OR 0.85×120 **[M1]**

102 **[A1]**

(b) Let cost price be C. $1.2C = 120$ **[M1]**

$C = \dfrac{120}{1.2}$ **[M1]**

100 **[A1]**

[Back to Question] *[Total Marks: 5]*

177 **(a)** M1 for $15 \times (1 - 0.20)$ or $15 - 3$

12 **[A1]**

(b) M2 for $56 \div 0.8$ or $\dfrac{56}{80} \times 100$
(M1 for $80\% = 56$ or equivalent)

70 **[A1]**

(c) M1 for $\dfrac{42 - 30}{30} \times 100$ or $\dfrac{12}{30} \times 100$

40 **[A1]**

[Back to Question] *[Total Marks: 7]*

178 **(a)** $4500 \times \left(1 + \dfrac{3.2}{100}\right)^5$ **[M2]**

OR 4500×1.032^5 **[M2]**

(M1 for 4500×1.032^k where $k > 1$)

5267.57... **[A1]**

(b) $\dfrac{408 - 320}{320} \times 100$ or $\dfrac{88}{320} \times 100$ **[M1]**

27.5 **[A1]**

[Back to Question] *[Total Marks: 5]*

179 **(a)** $45\,000 \times 1.042$ OR $45\,000 + 45\,000 \times 0.042$ **[M1]**

46 890 **[A1]**

(b) $100\% - 24\% = 76\%$ **[B1]**

$\dfrac{342}{0.76}$ OR $\dfrac{342}{76} \times 100$ **[M1]**

450 **[A1]**

[Back to Question] *[Total Marks: 5]*

180 **(a)** $4500 \times (1 + \dfrac{2.4}{100})^5$ or 4500×1.024^5 **[M2]**

If M0, M1 for 4500×1.024 or 4608 seen.

5066.57 or 5066.56... **[A1]**

(b) Interest earned $= 5066.57 - 4500 = 566.57$ (accept unrounded) **[M1]**

$\frac{4500 \times r \times 5}{100} = 566.57$ OR $225r = 566.57$ **[M1]**

2.52 or 2.518... **[A1]**

[Back to Question] *[Total Marks: 6]*

181 **(a)** $8000 \times \left(1 + \frac{3.5}{100}\right)^4$ or 8000×1.035^4 **[M2]**

If M0 scored, allow [M1] for 8000×1.035^k where $k > 1$

9180.18 **[A1]**

(b) $x \times \left(1 - \frac{4}{100}\right)^3 = 45000$ or $x \times 0.96^3 = 45000$ **[M1]**

$\frac{45000}{0.96^3}$ **[M1]**

50863 **[A1]**

[Back to Question] *[Total Marks: 6]*

182 **(a)** $4500 \times \left(1 + \frac{2.4}{100}\right)^5$ or 4500×1.024^5 **[M2]**

[M1] for 4500×1.024^n where $n > 1$

5066.55 **[A1]**

(b) $100\% - 18\% = 82\%$ or 0.82 **[B1]**

$\frac{557.60}{0.82}$ or $x \times 0.82 = 557.60$ **[M1]**

680 **[A1]**

[Back to Question] *[Total Marks: 6]*

183 **(a)** M2 for $8000 \times \left(1 + \frac{3.5}{100}\right)^4$

(M1 for 8000×1.035^k where $k > 1$)

9180.19 **[A1]**

(b) M2 for $24000 \times 0.88^n < 10000$ or evaluating at least two consecutive trials (e.g. $n = 6$ and $n = 7$)

(M1 for 24000×0.88^n seen)

A1 for $n = 7$ (Value at year 6 is \$11145.69, Value at year 7 is \$9808.21)

(c) M2 for $150000 \div 1.25$

(M1 for $125\% = 150000$ or equivalent)

120000 **[A1]**

[Back to Question] *[Total Marks: 9]*

184 **(a)** $(1.25 \times 10^6) \times 1.045^{10}$ **[M1]**

1941214.7... **[A1]**

1.94×10^6 **[A1]**

(b) $299.25 = 86\%$ or $\dfrac{299.25}{1 - 0.14}$ **[M1]**

348 **[A1]**

(c) Let V be the original value. $V \times 0.85 \times 0.90^2 = 17212.50$ **[M1]**

$V = \dfrac{17212.50}{0.85 \times 0.81} = 25000$ **[A1]**

[Back to Question] *[Total Marks: 7]*

185 **(a)** $1\,250\,000 \times (1 + \dfrac{x}{100})^3 = 1\,367\,631$ **[M1]**

$(1 + \dfrac{x}{100})^3 = 1.0941048$ **[M1]**

$1 + \dfrac{x}{100} = \sqrt[3]{1.0941048} = 1.0304\cdots \implies x = 3.04\cdots \approx 3$ **[A1]**

(b) Target population $= 2 \times 1\,250\,000 = 2\,500\,000$ **[B1]**

$1\,250\,000 \times (1.03)^n > 2\,500\,000 \implies 1.03^n > 2$ **[M1]**

Using logs: $n \log(1.03) > \log(2)$ OR trial and improvement yielding n between 23 and 24 **[M1]**

$n > 23.449\ldots$, so it takes 24 years. Year $= 2015 + 24 = 2039$ **[A1]**

[Back to Question] *[Total Marks: 7]*

186 **(a)** $p \times 1.2 = 960$ or $\dfrac{960}{1.2}$ **[M1]**

800 **[A1]**

(b) Selling price $= 1.15q$ **[B1]**

Marked Price $\times 0.9 = 1.15q$ or $\dfrac{1.15q}{0.9}$ **[M1]**

$\dfrac{23q}{18}$ or $\dfrac{23}{18}q$ **[A1]**

(c) $500 \times \left(1 + \dfrac{r}{100}\right)^{10} = 740.12$ or $\left(1 + \dfrac{r}{100}\right)^{10} = \dfrac{740.12}{500}$ **[M1]**

$1 + \dfrac{r}{100} = \sqrt[10]{1.48024}$ or 1.04 **[M1]**

4 **[A1]**

[Back to Question] *[Total Marks: 8]*

187 **(a)** $5000 \times \left(1 + \frac{2.5}{100}\right)^3$ **[M1]**

5384.45 **[A1]**

(b) $\left(1 + \frac{r}{100}\right)^5 = 1.30$ **[M1]**

$1 + \frac{r}{100} = \sqrt[5]{1.30}$ **[M1]**

5.39 (or 5.387...) **[A1]**

(c) $V \times (1 - 0.12)^3 = 17\,036.80$ **[M1]**

$V = \frac{17\,036.80}{0.88^3}$ **[M1]**

25 000 **[A1]**

[Back to Question] *[Total Marks: 8]*

188 **(a)** Expression for 3 years depreciation: e.g. $V \times (1 - 0.15) \times (1 - 0.08)^2$ or $V \times 0.85 \times 0.92^2$ **[M2]**

[M1] for $V \times 0.85$ or $V \times 0.92^2$ seen

$V \times 0.71944 = 17986$ or $V = \frac{17986}{0.85 \times 0.92^2}$ **[M1]**

25 000 **[A1]**

(b) $\frac{25000 - 17986}{25000} \times 100$ or $100\% - (0.85 \times 0.92^2 \times 100)\%$ **[M1]**

28.1 **[A1]**

(c) $P \times \left(1 + \frac{r}{100}\right)^{12} = 2P$ or $\left(1 + \frac{r}{100}\right)^{12} = 2$ **[M1]**

$1 + \frac{r}{100} = \sqrt[12]{2}$ or 1.059... **[M1]**

5.95 (correct to 3 significant figures) **[A1]**

[Back to Question] *[Total Marks: 9]*

189 **(a)** M2 for $\left(1 + \frac{r}{100}\right)^6 = 2$ or equivalent

(M1 for $\left(1 + \frac{r}{100}\right)^6$ seen)

12.2 **[A1]**

(b) M2 for $X \times 1.40 \times 0.85 = 142.80$ or $1.19X = 142.80$

(M1 for $X \times 1.40$ or Selling Price $\times 0.85 = 142.80$)

120 **[A1]**

(c) M1 for $P \times \frac{x}{100} \times 5 = 250$ (implies $Px = 5000$)

M2 for $P\left(1 + \frac{x}{100}\right)^2 - P = 102.50$

(M1 for $P\left(1 + \frac{x}{100}\right)^2 = P + 102.50$)

M1 for correctly substituting to form an equation in one variable, e.g. $\frac{5000}{x}\left(\frac{2x}{100}+\frac{x^2}{10000}\right)=102.50$

$x=5$ **[A1]**

$P=1000$ **[A1]**

[Back to Question] *[Total Marks: 11]*

190 Numerator $\approx 1.907878...$, Denominator $= 1.025$ **[M1]**

1.861344784 **[A1]**

[Back to Question] *[Total Marks: 2]*

191 (a) 3.068822... **[B2]**

If 0 scored, allow [B1] for seeing 15.36 or 2.485... or 0.81

(b) 3.069 [B1] (strict FT from their (a) if given to at least 4 d.p.)

[Back to Question] *[Total Marks: 3]*

192 (a) Numerator $\sqrt{22.01}=4.69148\ldots$ or Denominator $12.167-1.4=10.767$ seen **[M1]**

0.435727839 **[A1]**

(b) 0.436 **[B1]**

[Back to Question] *[Total Marks: 3]*

193 (a) M1 for $\sqrt{158.79}$ or $12.60119\ldots$ or 4.48 seen

2.812765825 (accept to at least 4 decimal places) **[A1]**

(b) 2.81 **[B1ft]**

(c) 3.75×10^8 **[B1]**

[Back to Question] *[Total Marks: 4]*

194 (a) 4.423... or $\frac{15.98...}{3.612}$ **[M1]**

4.42 or 4.424 **[A1]**

(b) 1.4862... − 0.8191... or 0.6671... **[M1]**

0.667 **[A1]**

[Back to Question] *[Total Marks: 4]*

195 (a) 4.8 **[M1]**

4.80 **[A1]**

(b) $2 + \frac{18}{60}$ or $\frac{138}{60}$ **[M1]**

2.3 **[A1]**

[Back to Question] *[Total Marks: 4]*

196 **(a)** $150.0625 + 4.4412...$ or $154.5037...$ **[M1]**

154.50 **[A1]**

(b) $\frac{1.3639...}{0.9510...}$ or $1.434...$ **[M1]**

1.43 **[A1]**

[Back to Question] *[Total Marks: 4]*

197 **(a)** 8.40 **[B1]**

(b) 5 hours 9 minutes **[B1]**

(c) $280 \div 85$ **[M1]**

3.294... **[A1]**

3 hours 18 minutes **[A1]**

[Back to Question] *[Total Marks: 5]*

198 **(a)** **(i)** Numerator 4.94 or denominator 1.14... seen **[M1]**

4.332666... **[A1]**

(ii) 4.33 **[B1]**

(b) 0.35×60 **[M1]**

5 h 21 min **[A1]**

[Back to Question] *[Total Marks: 5]*

199 **(a)** $\pi \times 4.2^2$ or $55.417...$ **[M1]**

55 **[A1]**

(b) $23 \div 60$ or $0.3833...$ seen **[M1]**

Time $= 4 + \frac{23}{60}$ or $4.3833...$ hours **[M1]**

Distance $= 78 \times 4.3833...$ **[M1]**

341.9 **[A1]**

[Back to Question] *[Total Marks: 6]*

200 **(a)** Time $= 3 + \frac{40}{60}$ or $\frac{11}{3}$ hours **[M1]**

Cost $= 45 + 32.50 \times \frac{11}{3}$ **[M1]**

164.17 **[A1]**

(b) $230.25 - 45 = 185.25$ **[M1]**

$\frac{185.25}{32.50} = 5.7$ hours **[M1]**

5 hours 42 minutes **[A1]**

[Back to Question] *[Total Marks: 6]*

201 **(a)** M1 for $\frac{4}{3} \times \pi \times 4.85^3$

477.8 **[A1]**

(b) M1 for $\frac{1}{3} \times \pi \times 1.2^2 \times 3.5$ (implies $5.2778\ldots$)

M1 for $\frac{\text{their } 477.8}{\text{their } 5.2778\ldots}$

A1 for $90.5\ldots$

90 **[B1]**

(c) 0.584 (accept 0.582 to 0.585) **[B1ft]**

[Back to Question] *[Total Marks: 7]*

202 **(a)** $\frac{4}{3}\pi(1.4)^3$ **[M1]**

11.49404... **[A1]**

(b) their 11.494... $\div$ 0.035 **[M1]**

328.4... or 328 (minutes) **[A1]**

their 328.4... $\div$ 60 (converting to hours) **[M1]**

5 hours 28 minutes **[A1]**

[Back to Question] *[Total Marks: 6]*

203 **(a)** 14.4×10^3 or 14400 **[M1]**

1.44×10^4 **[A1]**

(b) $\frac{1}{2.5066...} \times 0.7788...$ or $0.398942... \times 0.778800...$ or $0.310696...$ **[M1]**

0.3107 **[A1]**

(c) $1.5 = 2\pi\sqrt{\frac{l}{9.81}} \implies \frac{1.5}{2\pi} = \sqrt{\frac{l}{9.81}} \implies 0.2387... = \sqrt{\frac{l}{9.81}}$ **[M1]**

$l = 9.81 \times (0.2387...)^2 \implies l = 0.55909...$ **[M1]**

0.56 **[A1]**

[Back to Question] *[Total Marks: 6]*

204 **(a)** Numerator $= 340000 + 27000 = 367000$ or 3.67×10^5 **[M1]**

$3.67 \times 10^5 \div (8.1 \times 10^{-3}) = 45308641.98...$ **[M1]**

4.53×10^7 or 4.531×10^7 **[A1]**

(b) Upper bound for $r = 5.25$ and Upper bound for $h = 12.45$ **[M1]**

$V = \frac{1}{3} \times \pi \times 5.25^2 \times 12.45$ **[M1]**

359.34... or 359 **[A1]**

[Back to Question] *[Total Marks: 6]*

205 **(a)** M1 for $2\pi\sqrt{\frac{0.85}{9.81}}$

1.850 **[A1]**

(b) M1 for $\left(\frac{T}{2\pi}\right)^2 = \frac{l}{g}$ or $\frac{T^2}{4\pi^2} = \frac{l}{g}$

$\frac{gT^2}{4\pi^2}$ **[A1]**

(c) M1 for their $l = \frac{9.81 \times 2.5^2}{4\pi^2}$

A1 for $l = 1.5532\ldots$ metres

155.3 **[A1]**

[Back to Question] *[Total Marks: 7]*

206 **(a)** $\frac{1}{3} \times \pi \times 3.2^2 \times (3 \times 8.45 - 3.2)$ **[M1]**

237.5 **[A1]**

(b) $\frac{3V}{\pi h^2} = 3R - h$ **[M1]**

$\frac{3V}{\pi h^2} + h = 3R$ **[M1]**

$R = \frac{V}{\pi h^2} + \frac{h}{3}$ or $R = \frac{3V + \pi h^3}{3\pi h^2}$ **[A1]**

(c) $\frac{500}{\pi \times 5.6^2} + \frac{5.6}{3}$ **[M1]**

6.94 **[A1]**

[Back to Question] *[Total Marks: 7]*

207 **(a)** $\frac{1}{3} \times \pi \times 2.31^2 \times (3 \times 8.74 - 2.31)$ **[M1]**

133.607... **[A1]**

133.6 **[A1]**

(b) Rounding initial values: $R = 8.7$ and $h = 2.3$ **[B1]**

$\frac{1}{3} \times \pi \times 2.3^2 \times (3 \times 8.7 - 2.3)$ or 131.84... **[M1]**

Difference = 133.607... − 131.844... = 1.76... **[M1]**

1.8 **[A1]**

[Back to Question] *[Total Marks: 7]*

208 **(a)** 2 hours 35 minutes **[B1]**

(b) M1 for 15 35 or 17 70 seen

18 10 **[A1]**

[Back to Question] *[Total Marks: 3]*

209 **(a)** 2 hours 15 minutes **[B1]**

(b) Adds 2 hours 35 minutes to 16 45 (e.g., 18 80) **[M1]**

19 20 **[A1]**

[Back to Question] *[Total Marks: 3]*

210 Arrival time in London local time = 21 : 45 + 6 hours 50 minutes **[M1]**

04 : 35 (next day) **[M1]**

Dubai local time = 04 : 35 + 4 hours = 08 : 35 **[A1]**

[Back to Question] *[Total Marks: 3]*

211 **(a)** London departure time = 10 30 or Dubai arrival time = 23 15 **[M1]**

8 hours 45 minutes **[A1]**

(b) London departure time = 18 45 or Dubai arrival time = 06 05 (Saturday) **[M1]**

02 05 Saturday **[A1]**

[Back to Question] *[Total Marks: 4]*

212 **(a)** 22 50 + 14 h 35 min **[M1]**

13 25 **[A1]**

(b) 0.4×60 **[M1]**

3 h 24 min **[A1]**

[Back to Question] *[Total Marks: 4]*

213 **(a)** 115 minutes = 1 hour 55 minutes **[M1]**

21 30 (or 21 : 30) **[A1]**

(b) Arrival time in New York = 21 40 + 7 h 15 m = 04 55 (next day) **[M1]**

Arrival time in Paris = 04 55 + 6 hours **[M1]**

10 55 (or 10 : 55) **[A1]**

[Back to Question] *[Total Marks: 5]*

214 Flight arrival time in Dubai local time: 14 30 + 14 h 15 min = 28 45 (or 04 45 next day) **[M1]**

Adds time zone difference: 28 45 + 7 hours (or 04 45 + 7 hours) **[M1]**

11 45 **[A1]**

[Back to Question] *[Total Marks: 3]*

215 **(a)** $3 \times 10^5 \times 60 \times 60 \times 24 \times 365$ **[M1]**

9.46×10^{12} or 9.4608×10^{12} **[A1]**

(b) $\dfrac{4.5 \times 10^{13}}{9.4608 \times 10^{12}}$ **[M1]**

4.76 **[A1]**

[Back to Question] *[Total Marks: 4]*

216 **(a)** 11 : 03 − 08 : 15 **[M1]**

2 hours 48 minutes **[A1]**

(b) 48 minutes $= \dfrac{48}{60}$ hours = 0.8 hours **[M1]**

Average speed $= \dfrac{420}{2.8}$ **[M1]**

150 **[A1]**

[Back to Question] *[Total Marks: 5]*

217 **(a)** M1 for arrival time in Dubai (10 00 Tuesday) OR departure time in London (22 15 Monday) seen

M1 for a valid time zone adjustment and flight duration addition

06 00, Tuesday **[A1]**

(b) M1 for $1.5 \times 365 \times 24 \times 60$

A1 for 788400

7.884×10^5 **[B1]**

[Back to Question] *[Total Marks: 6]*

218 **(a)** London departure time = 16 30 − 14 h 15 min or 02 15 **[M1]**

Tokyo departure time = 02 15 + 9 h **[M1]**

11 15 **[A1]**

(b) 1 min 45 s = 105 seconds or 8 hours = 480 minutes **[B1]**

$(8 \times 3600) \div 105$ or $480 \div 1.75$ **[M1]**

274 **[A1]**

[Back to Question] *[Total Marks: 6]*

219 **(a)** Time = $\frac{42.2}{12.5}$ or 3.376 hours **[M1]**

$0.376 \times 60 = 22.56$ minutes **[M1]**

$0.56 \times 60 = 33.6$ seconds **[M1]**

3 hours, 22 minutes, 34 seconds **[A1]**

(b) Correct difference calculation, e.g. 12 hours 64 minutes −08 hours 35 minutes **[M1]**

4 hours 29 minutes **[A1]**

[Back to Question] *[Total Marks: 6]*

220 **(a)** M1 for $30 \times 24 \times 60$ or 43200 minutes seen

M1 for $\frac{43200}{94.5}$ or 457.142...

457 **[A1]**

(b) M1 for 100×94.5 or 9450 minutes

M1 for converting minutes into hours and days (e.g., 157.5 hours OR 6 days 13 hours 30 minutes)

M1 for adding their 6 days and 13 hours 30 minutes to 05 50 on 1st March

19 20, 7th March **[A1]**

[Back to Question] *[Total Marks: 7]*

221 **(a)** $920 \div 24$ **[M1]**

38.333... hours **[A1]**

38 hours 20 minutes **[A1]**

(b) Adds their time from part (a) to Wednesday 17 50 **[M1]**

Wednesday 17 50 + 20 minutes = 18 10 **[M1]**

Wednesday 18 10 + 38 hours = Thursday 18 10 + 14 hours **[M1]**

Day: Friday **[A1]**

Time: 08 10 **[A1]**

[Back to Question] *[Total Marks: 7]*

222 **(a)** $\frac{6600}{800}$ or 8.25 **[M1]**

8 hours 15 minutes **[A1]**

(b) Arrival time in City A time = 05 55 **[M1]**

00 55 Wednesday **[A1]**

(c) Departure from City B = 15 15 (local time) or 20 15 (City A time) **[M1]**

Arrival in City B time = 23 00 **[M1]**

04 00 Thursday **[A1]**

[Back to Question] *[Total Marks: 7]*

223 **(a)** $\frac{35}{x} + \frac{42}{x-5}$ **[B1]**

(b) 4 hours 33 minutes $= 4 + \frac{33}{60}$ or $\frac{91}{20}$ hours **[B1]**

$\frac{35}{x} + \frac{42}{x-5} = \frac{91}{20}$ **[M1]**

Multiplies by common denominator: $20 \times 35(x-5) + 20 \times 42x = 91x(x-5)$ **[M1]**

$700x - 3500 + 840x = 91x^2 - 455x$ **[M1]**

$1540x - 3500 = 91x^2 - 455x \implies 91x^2 - 1995x + 3500 = 0$ **[M1]**

Divides by 7 to reach $13x^2 - 285x + 500 = 0$ without errors **[A1]**

(c) Uses quadratic formula or factors: $(13x - 25)(x - 20) = 0$ **[M1]**

$x = 20$ or $x = \frac{25}{13}$ (or 1.92...) **[A1]**

$x = 20$ only (rejects $\frac{25}{13}$ as speed $x - 5$ must be positive) **[A1]**

[Back to Question] *[Total Marks: 8]*

224 **(a)** Minutes in a year $= 365 \times 24 \times 60$ **[M1]**

525600 minutes **[M1]**

Number of orbits $= \frac{525600}{92} = 5713.043\ldots$ **[M1]**

5713 **[A1]**

(b) **(i)** $\frac{11440}{880} = 13$ **[B1]**

(ii) Time difference $= 9 - (-4) = 13$ hours (Q is 13 hours ahead) **[M1]**

Departure time in City Q's local time = 22 : 45 (Monday) +13 hours = 11 : 45 (Tuesday)

[M1]

Arrival time in City Q's local time = 11 : 45 (Tuesday) +13 hours (flight time) = 00 : 45 (Wednesday) [A1]

[Back to Question] *[Total Marks: 8]*

225 **(a)** 15×92.75 or 1380 min 675 s [M1]

23 h 11 min 15 s [A1]

(b) Arrival time in City A local time = 09 45 + 8 h 20 min or 18 05 [M1]

26 05 − 18 05 [M1]

8 [A1]

(c) Distance travelled by Train P at 08:30 = $90 \times 0.5 = 45$ km [M1]

Relative speed of approaching trains = $90 + 120 = 210$ km/h [M1]

Time taken to meet from 08:30 = $(360 - 45) \div 210 = 1.5$ hours [M1]

10 00 [A1]

[Back to Question] *[Total Marks: 9]*

226 **(a)** 1680 [B1]

(b) M1 for 1680 − 1050 or 630 seen

140 [A1]

[Back to Question] *[Total Marks: 3]*

227 $255 \div 0.85$ [M1]

\$300 [A1]

\$100 [A1]

[Back to Question] *[Total Marks: 3]*

228 $(3 \times 1.20) + (4 \times 2.50)$ or $3.60 + 10.00$ or 13.60 [M1]

$20 - 13.60$ [M1]

6.40 [A1]

[Back to Question] *[Total Marks: 3]*

229 $300 \times 1.50 = 450$ [M1]

$450 - 210 = 240$ [M1]

$240 \div 1.50 = 160$ [A1]

[Back to Question] *[Total Marks: 3]*

230 **(a)** Cost of pens $= 4 \times 2.35 = 9.40$ **[M1]**

Cost of notebooks $= 3 \times 4.50 = 13.50$ **[M1]**

Change $= 50 - (9.40 + 13.50) = 27.10$ **[A1]**

(b) 54 **[B1]**

[Back to Question] *[Total Marks: 4]*

231 **(a)** 850×148 **[M1]**

125800 **[A1]**

(b) $125800 - (45000 + 31500)$ or 49300 **[M1]**

$\frac{49300}{152}$ or 324.342... **[M1]**

324.34 **[A1]**

[Back to Question] *[Total Marks: 5]*

232 **(a)** $1\,250\,000 \div 145$ **[M1]**

8620.69 **[A1]**

(b) Total cost $= 8620.69 + 600 = 9220.69$ **[M1]**

Profit $= 12\,500 - 9220.69 = 3279.31$ (or $\frac{12500 - 9220.69}{9220.69}$) **[M1]**

35.6 (accept $35.56\ldots$) **[A1]**

[Back to Question] *[Total Marks: 5]*

233 **(a)** 450×1.14 **[M1]**

513 **[A1]**

(b) $1.32 \div 1.14$ **[M1]**

1.1578... **[A1]**

1.16 **[A1]**

[Back to Question] *[Total Marks: 5]*

234 **(a)** M1 for 680×1.32 (implies 897.60)

M1 for their $897.60 - 850$
OR
M1 for $850 \div 1.32$ (implies $643.939\ldots$)
M1 for (680 − their $643.939\ldots$) $\times 1.32$

47.60 (accept 47.6) **[A1]**

(b) M1 for 850×1.04 or $850 \times 0.04 + 850$ (implies 884)

M1 for their 884×0.85

751.40 (accept 751.4) **[A1]**

[Back to Question] *[Total Marks: 6]*

235 **(a)** Amount in euros $= 450 \times 1.18 = 531$ **[M1]**

Remaining euros $= 531 - 285 = 246$ **[M1]**

Amount back in pounds $= 246 \div 1.22$ **[M1]**

201.639... rounding to 201.64 **[A1]**

(b) Sale price $= 850 \times 0.85$ $(= 722.5)$ **[M1]**

Final price $= 722.5 \times 0.95 = 686.375$ which rounds to 686.38 **[A1]**

[Back to Question] *[Total Marks: 6]*

236 **(a)** 850×0.98 or $850 - (850 \times 0.02)$ **[M1]**

833 **[A1]**

149940 **[A1]**

(b) $149940 - 115000$ $(= 34940)$ **[M1]**

their $34940 - 500$ $(= 34440)$ **[M1]**

their $34440 \div 184$ **[M1]**

187.17 (accept 187.17...) **[A1]**

[Back to Question] *[Total Marks: 7]*

237 **(a)** $\frac{17400 + 1200}{1.24}$ or $\frac{18600}{1.24}$ **[M1]**

15000 **[A1]**

(b) $17100 - 15000$ or 2100 **[M1]**

$\frac{2100}{15000} \times 100$ **[M1]**

14 **[A1]**

(c) $2100 \times (1 - 0.02)$ or 2100×0.98 or 2058 **[M1]**

2058×1.16 **[M1]**

2387.28 **[A1]**

[Back to Question] *[Total Marks: 8]*

238 Total revenue in USD $= 1000 \times 4.20 = 4200$ **[B1]**

Convert revenue to EUR: $4200 \div 1.20$ $(= 3500)$ **[M1]**

Equate to cost with profit multiplier: Total Cost $\times 1.40 = 3500$ or $3500 \div 1.40$ $(= 2500)$ **[M1]**

Form equation for cost: $1000y + 250 = 2500$ **[M1]**

$y = 2.25$ **[A1]**

[Back to Question] *[Total Marks: 5]*

239 **(a)** 800×0.85 or 680 **[M1]**

$680 - 420$ or 260 **[M1]**

$260 \div 0.82$ **[M1]**

317.07 **[A1]**

(b) $61.20 \div (1 - 0.15)$ or $61.20 \div 0.85$ **[M1]**

72 **[A1]**

[Back to Question] *[Total Marks: 6]*

240 **(a)** Cost of Brand A per 100g $= 12.50 \div 5 = 2.50$ (or per gram $= 0.025$) **[M1]**

Cost of Brand B per 100g $= 8.40 \div 3 = 2.80$ (or per gram $= 0.028$) **[M1]**

Brand A with correct supporting working **[A1]**

(b) $x + y = 2000$ **[M1]**

Cost equation: $0.025x + 0.028y = 54.10$ (or equivalent, e.g. $\frac{12.5}{500}x + \frac{8.4}{300}y = 54.1$) **[M1]**

Substitute $y = 2000 - x$: $0.025x + 0.028(2000 - x) = 54.10$ **[M1]**

$0.025x + 56 - 0.028x = 54.10 \implies -0.003x = -1.9 \implies x = 633.3...$ and $y = 1366.6...$

Accept $x = 633$ and $y = 1367$ (or exact fractions $\frac{1900}{3}$ and $\frac{4100}{3}$) **[A1]**

[Back to Question] *[Total Marks: 7]*

241 **(a)** $\frac{400}{x}$ **[B1]**

(b) $\frac{450}{x - 0.05} - \frac{400}{x} = 100$ **[M1]**

$450x - 400(x - 0.05) = 100x(x - 0.05)$ **[M1]**

$450x - 400x + 20 = 100x^2 - 5x$ **[M1]**

$50x + 20 = 100x^2 - 5x \implies 100x^2 - 55x - 20 = 0$ **[A1]**

Dividing by 5 yields $20x^2 - 11x - 4 = 0$ (must show working to reach final equation) **[A1]**

(c) $(5x - 4)(4x + 1) = 0$ or use of quadratic formula $\frac{11 \pm \sqrt{(-11)^2 - 4(20)(-4)}}{2(20)}$ **[M2]**

If M2 not scored, allow [M1] for $(5x \pm a)(4x \pm b) = 0$ where $ab = 4$ or $a(4) \pm b(5) = -11$, or correct substitution into formula but with one error.

0.8 (with -0.25 rejected or ignored) **[A1]**

[Back to Question] *[Total Marks: 8]*

242 **(a)** M1 for $21500 - 24000x = 740$

M1 for $24000x = 20760$

0.865 **[A1]**

(b) M1 for 780×1.28 (implies 998.4)

M1 for their $998.4 \div (1 - 0.025)$ or their $998.4 \div 0.975$

1024 **[A1]**

(c) M1 for setting up a correct equation, e.g. $R \times (1 - 0.08) \times k = R \times 0.828$ or $0.92k = 0.828$

M1 for $k = \dfrac{0.828}{0.92}$

0.9 **[A1]**

[Back to Question] *[Total Marks: 9]*

243 **(a)** 840×1.15 or 966 **[M1]**

$\dfrac{821.10}{966}$ or 0.85 **[M1]**

15 **[A1]**

(b) $\dfrac{2430000}{450}$ or 5400 **[M1]**

$\dfrac{5400}{7.2}$ or 750 **[M1]**

90 **[A1]**

(c) 840×1.025 or 861 **[M1]**

$\dfrac{861}{1.32}$ or 652.272... **[M1]**

652.27 **[A1]**

[Back to Question] *[Total Marks: 9]*

244 **(a)** **(a)** $8000(1 + \dfrac{p}{100})^3 = 27000$ **[M1]**

$1 + \dfrac{p}{100} = \sqrt[3]{\dfrac{27000}{8000}} = \dfrac{3}{2}$ **[M1]**

50 **[A1]**

(b) $27000 \times (1.5)^2$ or $8000 \times (1.5)^5$ **[M1]**

60 750 **[A1]**

[Back to Question] *[Total Marks: 5]*

245 **(a)** $\left(1 + \dfrac{r}{100}\right)^2 = 1.44$ or $x^2 = 1.44$ **[M1]**

$1 + \dfrac{r}{100} = 1.2$ or $x = 1.2$ **[M1]**

20 **[A1]**

(b) Number of doubling periods $= 45 \div 15 = 3$ or 80×2^3 **[M1]**

640 **[A1]**

[Back to Question] *[Total Marks: 5]*

246 M1 for $24000 \times (1 - 0.08)^5$ or 24000×0.92^5

A1 for 15817.96...

15818 **[B1]**

[Back to Question] *[Total Marks: 3]*

247 $125\,000 \times (1 + 0.018)^8$ or $125\,000 \times 1.018^8$ [M2]
(If not M2, award M1 for 1.018^8 or $125\,000 \times 1.018^k$ where $k > 1$)

144 183.6... **[A1]**

144 184 **[B1]**

[Back to Question] *[Total Marks: 3]*

248 $26000 \times (1 - 0.15)^4$ or 26000×0.85^4 **[M2]**

If M0, allow [M1] for 26000×0.85^k where $k > 1$

13572 **[A1]**

[Back to Question] *[Total Marks: 3]*

249 $450 \times \left(1 + \frac{6.5}{100}\right)^{12}$ **[M1]**

450×1.065^{12} **[M1]**

958.01 (accept 958) **[A1]**

[Back to Question] *[Total Marks: 3]*

250 $32000 \times (1 - 0.12)^5$ oe **[M2]**

For 32000×0.88^n where $n > 1$, award **[M1]**

16887 **[A1]**

[Back to Question] *[Total Marks: 3]*

251 **(a)** $65000 \times \left(1 + \frac{2.8}{100}\right)^5$ or 65000×1.028^5 **[M1]**

74624 **[A1]**

(b) $\frac{74624 - 65000}{65000} \times 100$ or $(1.028^5 - 1) \times 100$ **[M1]**

14.8 **[A1]**

[Back to Question] *[Total Marks: 4]*

252 **(a)** 1500×1.08^{12} **[M1]**

3777 **[A1]**

(b) $1500 \times 1.08^n > 3000$ or $1.08^n = 2$ **[M1]**

Trial with values (e.g., $1.08^9 = 1.999...$ and $1.08^{10} = 2.158...$) or uses logarithms (e.g., $n = \dfrac{\log(2)}{\log(1.08)} = 9.006...$) **[M1]**

10 **[A1]**

[Back to Question] *[Total Marks: 5]*

253 **(a)** **(a)** $85000(1 + \dfrac{r}{100})^{12} = 112000$ **[M1]**

$1 + \dfrac{r}{100} = \left(\dfrac{112}{85}\right)^{\frac{1}{12}}$ **[M1]**

2.33 (from 2.327...) **[A1]**

(b) $85000 \times \left(1 + \dfrac{2.327...}{100}\right)^{20}$ or $112000 \times \left(1 + \dfrac{2.327...}{100}\right)^{8}$ **[M1]**

135 000 **[A1]**

[Back to Question] *[Total Marks: 5]*

254 **(a)** $28\,000 \times \left(1 - \dfrac{r}{100}\right)^4 = 16\,150$ **[M1]**

$\left(1 - \dfrac{r}{100}\right) = \sqrt[4]{\dfrac{16\,150}{28\,000}}$ or $\sqrt[4]{0.57678...}$ **[M1]**

12.9 (or 12.88 to 12.89) **[A1]**

(b) $16\,150 \times (1 - 0.1289...)^3$ or $28\,000 \times (1 - 0.1289...)^7$ **[M1]**

10 677 **[A1]**

[Back to Question] *[Total Marks: 5]*

255 **(a)** $45000 \times \left(1 - \dfrac{x}{100}\right)^4 = 26000$ **[M1]**

$1 - \dfrac{x}{100} = \sqrt[4]{\dfrac{26000}{45000}}$ or $1 - \dfrac{x}{100} = 0.8711...$ **[M1]**

12.9 **[A1]**

(b) $45000 \times (0.8711...)^n < 10000$ or $(0.8711...)^n < \dfrac{2}{9}$ **[M1]**

Testing values or using logarithms: $n = 10 \implies$ \$11342 and $n = 11 \implies$ \$9881 **[M1]**

11 **[A1]**

[Back to Question] *[Total Marks: 6]*

256 **(a)** M1 for $125000 \times \left(1 + \frac{2.4}{100}\right)^{10}$ or 125000×1.024^{10}

A1 for $158461.5\ldots$

158500 **[B1]**

(b) M1 for $85000 \times \left(1 - \frac{r}{100}\right)^{8} = 68000$

M1 for $1 - \frac{r}{100} = \sqrt[8]{\frac{68000}{85000}}$ or $0.8^{\frac{1}{8}}$

2.75 (accept 2.750 to 2.751) **[A1]**

[Back to Question] *[Total Marks: 6]*

257 **(a)** $28\,500 \times \left(1 - \frac{12}{100}\right)^{5}$ **[M1]**

$28\,500 \times 0.88^{5}$ **[M1]**

15 040 **[A1]**

(b) $35\,000 \times \left(1 - \frac{r}{100}\right)^{4} = 15\,200$ **[M1]**

$\left(1 - \frac{r}{100}\right)^{4} = \frac{152}{350}$ **[M1]**

$1 - \frac{r}{100} = \sqrt[4]{\frac{152}{350}} = 0.8123\ldots$ **[M1]**

18.8 (accept 18.76 to 18.77) **[A1]**

[Back to Question] *[Total Marks: 7]*

258 **(a)** Let $k = 1 - \frac{r}{100}$. Then $M_0 k^2 = 80$ and $M_0 k^5 = 40.96$ **[M1]**

$k^3 = \frac{40.96}{80} = 0.512 \implies k = \sqrt[3]{0.512} = 0.8$ **[M1]**

$1 - \frac{r}{100} = 0.8 \implies r = 20$ **[A1]**

(b) $M_0 \times 0.8^2 = 80$ or $M_0 \times 0.64 = 80$ **[M1]**

125 **[A1]**

(c) $125 \times 0.8^n < 10$ or $0.8^n < 0.08$ **[M1]**

Testing values: $n = 11 \implies 0.8^{11} = 0.0858...$ and $n = 12 \implies 0.8^{12} = 0.0687...$ **[M1]**

12 **[A1]**

[Back to Question] *[Total Marks: 8]*

259 **(a)** M1 for $500 \times (1 - 0.12)^7$ or 500×0.88^7

204.3 (accept 204 or 204.33 . . .) **[A1]**

(b) M1 for $500 \times 0.88^d < 50$ or $0.88^d = 0.1$

M1 for evaluating trials (e.g. $0.88^{18} = 0.1001\ldots$ and $0.88^{19} = 0.0881\ldots$) OR using logarithms $d > \dfrac{\log(0.1)}{\log(0.88)}$

19 **[A1]**

(c) M1 for $\left(1 - \dfrac{r}{100}\right)^{15} = 0.5$

M1 for $\sqrt[15]{0.5}$ or $0.5^{\frac{1}{15}}$

4.52 (accept 4.515 to 4.516) **[A1]**

[Back to Question] *[Total Marks: 8]*

260 **(a)** $4000 \times \left(1 + \dfrac{4.5}{100}\right)^{10}$ or 4000×1.045^{10} **[M1]**

6211.88 (or 6211.87...) **[A1]**

(b) $6211.88 + 500$ or 6711.88 **[B1]**

$4000 \times \left(1 + \dfrac{r}{100}\right)^{10} = 6711.88...$ **[M1]**

$1 + \dfrac{r}{100} = \sqrt[10]{\dfrac{6711.88...}{4000}}$ **[M1]**

5.31 (or 5.31...) **[A1]**

(c) $2500 \times 2^{\frac{20}{8}}$ or $2500 \times 2^{2.5}$ **[M1]**

14 142 (or 14 142.13...) **[A1]**

[Back to Question] *[Total Marks: 8]*

261 **(a)** $2500 \times \left(1 + \dfrac{r}{100}\right)^6 = 9100$ **[M1]**

$\left(1 + \dfrac{r}{100}\right)^6 = 3.64$ **[M1]**

$1 + \dfrac{r}{100} = 3.64^{\frac{1}{6}} = 1.2403\ldots$ **[M1]**

24.0 (accept 24.03 . . .) **[A1]**

(b) 2500×1.2403^{10} or 9100×1.2403^4 **[M1]**

21 500 **[A1]**

(c) $2500 \times 1.2403^n > 50\,000 \implies 1.2403^n > 20$ **[M1]**

$n \times \log(1.2403) > \log(20) \implies n > 13.91\ldots$ (or equivalent trial and error) **[M1]**

14 **[A1]**

[Back to Question] *[Total Marks: 9]*

262 $5\sqrt{3} - 3\sqrt{3} + 2\sqrt{3}$ **[M1]**

$4\sqrt{3}$ **[A1]**

[Back to Question] *[Total Marks: 2]*

263 $5\sqrt{3}$ or $4\sqrt{3}$ or $3\sqrt{3}$ seen **[M1]**

$6\sqrt{3}$ **[A1]**

[Back to Question] *[Total Marks: 2]*

264 $6\sqrt{2}$ or $5\sqrt{2}$ or $3\sqrt{2}$ seen **[M1]**

$8\sqrt{2}$ **[A1]**

[Back to Question] *[Total Marks: 2]*

265 $5\sqrt{3}$ or $3\sqrt{3}$ or $2\sqrt{3}$ seen **[M1]**

$4\sqrt{3}$ **[A1]**

[Back to Question] *[Total Marks: 2]*

266 $\sqrt{25 \times 3} + \sqrt{16 \times 3} - \sqrt{9 \times 3} = 5\sqrt{3} + 4\sqrt{3} - 3\sqrt{3}$ **[M1]**

$6\sqrt{3}$ **[A1]**

[Back to Question] *[Total Marks: 2]*

267 **(a)** $3\sqrt{5}$ **[B1]**

(b) $2\sqrt{5}$ or $4\sqrt{5}$ seen **[M1]**

$6\sqrt{5}$ **[A1]**

[Back to Question] *[Total Marks: 3]*

268 $\sqrt{75} = 5\sqrt{3}$ **[B1]**

$\sqrt{27} = 3\sqrt{3}$ or $\sqrt{12} = 2\sqrt{3}$ **[B1]**

$6\sqrt{3}$ **[A1]**

[Back to Question] *[Total Marks: 3]*

269 $\sqrt{48} = 4\sqrt{3}$ **[B1]**

$\sqrt{12} = 2\sqrt{3}$ **[B1]**

$4\sqrt{3} - 2\sqrt{3} = 2\sqrt{3}$ **[B1]**

[Back to Question] *[Total Marks: 3]*

270 **(a)** M1 for $5\sqrt{3}$ or $2\sqrt{3}$

$7\sqrt{3}$ **[A1]**

(b) M1 for $\dfrac{14 \times \sqrt{7}}{\sqrt{7} \times \sqrt{7}}$ or $\dfrac{14\sqrt{7}}{7}$

$2\sqrt{7}$ **[A1]**

[Back to Question] *[Total Marks: 4]*

271 **(a)** $\sqrt{25 \times 3}$ or $\sqrt{25}\sqrt{3}$ **[M1]**

$5\sqrt{3}$ **[A1]**

(b) $6\sqrt{2} - 2\sqrt{2}$ **[M1]**

$4\sqrt{2}$ **[A1]**

[Back to Question] *[Total Marks: 4]*

272 **(a)** **(i)** $3\sqrt{5} + 2\sqrt{5}$ **[M1]**

$5\sqrt{5}$ **[A1]**

(ii) Multiplies numerator and denominator by $\sqrt{3}$: $\dfrac{18\sqrt{3}}{3}$ **[M1]**

$6\sqrt{3}$ **[A1]**

[Back to Question] *[Total Marks: 4]*

273 **(a)** $3\sqrt{5}$ **[B1]**

(b) $6\sqrt{2} + 5\sqrt{2} - 3\sqrt{2}$ **[M1]**

$8\sqrt{2}$ **[A1]**

(c) $\dfrac{21\sqrt{7}}{\sqrt{7}\sqrt{7}}$ or $\dfrac{21\sqrt{7}}{7}$ **[M1]**

$3\sqrt{7}$ **[A1]**

[Back to Question] *[Total Marks: 5]*

274 **(a)** $8 + 4\sqrt{3} - 2\sqrt{3} - 3$ **[M1]**

$5 + 2\sqrt{3}$ **[A1]**

(b) $\dfrac{21\sqrt{7}}{7}$ **[M1]**

$3\sqrt{7}$ **[A1]**

[Back to Question] *[Total Marks: 4]*

275 **(a)** $9 - 3\sqrt{5} - 3\sqrt{5} + 5$ (or $9 - 6\sqrt{5} + 5$) **[M1]**

$14 - 6\sqrt{5}$ **[A1]**

(b) $\dfrac{14 \times \sqrt{7}}{\sqrt{7} \times \sqrt{7}}$ **[M1]**

$2\sqrt{7}$ **[A1]**

[Back to Question] *[Total Marks: 4]*

276 **(a)** $12 + 3\sqrt{5} - 8\sqrt{5} - 10$ **[M1]**

$2 - 5\sqrt{5}$ **[A1]**

(b) Multiplies numerator and denominator by $\sqrt{7}$ to get $\dfrac{14\sqrt{7}}{7}$ **[M1]**

$2\sqrt{7}$ **[A1]**

[Back to Question] *[Total Marks: 4]*

277 **(a)** **(a)** $12 + 3\sqrt{5} - 8\sqrt{5} - 2(5)$ or $12 - 5\sqrt{5} - 10$ **[M1]**

$2 - 5\sqrt{5}$ **[A1]**

(b) $\dfrac{15 \times \sqrt{3}}{\sqrt{3} \times \sqrt{3}}$ or $\dfrac{15\sqrt{3}}{3}$ **[M1]**

$5\sqrt{3}$ **[A1]**

[Back to Question] *[Total Marks: 4]*

278 **(a)** $12 - 3\sqrt{2} + 4\sqrt{2} - 2$ **[M1]**

$10 + \sqrt{2}$ **[A1]**

(b) Multiply numerator and denominator by $(5 + \sqrt{3})$ **[M1]**

Denominator evaluates to $25 - 3 = 22$ **[M1]**

$5 + \sqrt{3}$ **[A1]**

[Back to Question] *[Total Marks: 5]*

279 **(a)** M1 for $9 + 3\sqrt{2} + 3\sqrt{2} + 2$ (allow one numerical error in expansion)

$11 + 6\sqrt{2}$ **[A1]**

(b) M1 for multiplying numerator and denominator by $(3 + \sqrt{5})$

M1 for denominator evaluated to 4 (from $3^2 - (\sqrt{5})^2$)

$3 + \sqrt{5}$ **[A1]**

[Back to Question] *[Total Marks: 5]*

280 **(a)** $20 - 9$ or $4(5) - 9$ **[M1]**

11 **[A1]**

(b) Multiplies numerator and denominator by $(3+\sqrt{2})$ **[M1]**

Denominator of $9-2$ or 7 seen **[M1]**

$6+2\sqrt{2}$ **[A1]**

[Back to Question] *[Total Marks: 5]*

281 **(a)** $12+15\sqrt{2}-2\sqrt{2}-5$ **[M1]**

$7+13\sqrt{2}$ **[A1]**

(b) Multiplies numerator and denominator by $(4+\sqrt{3})$ **[M1]**

Denominator of $16-3$ or 13 **[M1]**

$8+2\sqrt{3}$ or $2(4+\sqrt{3})$ **[A1]**

[Back to Question] *[Total Marks: 5]*

282 **(a)** $\frac{15\sqrt{5}}{5}$ **[M1]**

$3\sqrt{5}$ **[A1]**

(b) $10-2\sqrt{3}+5\sqrt{3}-3$ **[M1]**

$7+3\sqrt{3}$ **[A2]**

[Back to Question] *[Total Marks: 5]*

283 **(a)** $6-3\sqrt{5}+2\sqrt{5}-5$ **[M1]**

$1-\sqrt{5}$ **[A1]**

(b) $\frac{14\times\sqrt{7}}{\sqrt{7}\times\sqrt{7}}$ **[M1]**

$\frac{14\sqrt{7}}{7}$ **[M1]**

$2\sqrt{7}$ **[A1]**

[Back to Question] *[Total Marks: 5]*

284 **(a)** $15+5\sqrt{7}-6\sqrt{7}-2(7)$ **[M1]**

$1-\sqrt{7}$ **[A1]**

(b) Multiplies numerator and denominator by $(5+\sqrt{3})$: $\frac{22(5+\sqrt{3})}{(5-\sqrt{3})(5+\sqrt{3})}$ **[M1]**

Denominator becomes $25-3=22$ **[M1]**

$5+\sqrt{3}$ **[A1]**

[Back to Question] *[Total Marks: 5]*

285 **(a)** $20 + 8\sqrt{2} - 15\sqrt{2} - 6(2)$ **[M1]**

$8 - 7\sqrt{2}$ **[A1]**

(b) $\dfrac{11(4+\sqrt{5})}{(4-\sqrt{5})(4+\sqrt{5})}$ **[M1]**

$\dfrac{11(4+\sqrt{5})}{16-5}$ or $\dfrac{11(4+\sqrt{5})}{11}$ **[M1]**

$4 + \sqrt{5}$ **[A1]**

(c) $3x\sqrt{3} - x\sqrt{3} = 18$ **[M1]**

$2x\sqrt{3} = 18$ or $x\sqrt{3} = 9$ **[M1]**

$x = \dfrac{9}{\sqrt{3}} = 3\sqrt{3}$ **[A1]**

[Back to Question] *[Total Marks: 8]*

286 $\times \dfrac{3+\sqrt{7}}{3+\sqrt{7}}$ **[M1]**

Numerator $15 + 5\sqrt{7} + 3\sqrt{7} + 7$ or $22 + 8\sqrt{7}$ **[B1]**

Denominator $9 - 7$ or 2 **[B1]**

$11 + 4\sqrt{7}$ **[A1]**

[Back to Question] *[Total Marks: 4]*

287 M1 for rearranging to group x terms, e.g., $x\sqrt{3} - x = 2 + \sqrt{12}$

M1 for factorising x, e.g., $x(\sqrt{3} - 1) = 2 + 2\sqrt{3}$

M1 for multiplying by the conjugate to rationalise: $\dfrac{2+2\sqrt{3}}{\sqrt{3}-1} \times \dfrac{\sqrt{3}+1}{\sqrt{3}+1}$

M1 for correctly expanding the numerator to $8 + 4\sqrt{3}$ or evaluating the denominator to 2

$4 + 2\sqrt{3}$ **[A1]**

[Back to Question] *[Total Marks: 5]*

288 Forms an equation for the area: $(a + b\sqrt{3})(2 - \sqrt{3}) = 1 + 4\sqrt{3}$ **[M1]**

Expands the brackets: $2a - a\sqrt{3} + 2b\sqrt{3} - 3b = 1 + 4\sqrt{3}$ **[M1]**

Equates rational and irrational parts to form simultaneous equations: $2a - 3b = 1$ and $-a + 2b = 4$ **[M1]**

Attempts to solve the simultaneous equations (e.g. multiplies second equation by 2: $-2a + 4b = 8$ and adds to first to get $b = 9$) **[M1]**

$a = 14$, $b = 9$ **[A1]**

[Back to Question] *[Total Marks: 5]*

289 **(a)** Multiplies numerator and denominator by $(2+\sqrt{3})$ **[M1]**

Numerator expands to $10+5\sqrt{3}+2\sqrt{3}+3$ **[M1]**

Denominator is $4-3=1$ leading to final answer $13+7\sqrt{3}$ **[A1]**

(b) $x=\dfrac{14}{3-\sqrt{2}}$ **[M1]**

Multiplies numerator and denominator by $(3+\sqrt{2})$ to get $\dfrac{14(3+\sqrt{2})}{9-2}$ **[M1]**

$x=6+2\sqrt{2}$ **[A1]**

[Back to Question] *[Total Marks: 6]*

290 **(a)** Multiply numerator and denominator by $2+\sqrt{3}$ **[M1]**

Numerator becomes $(5+\sqrt{3})(2+\sqrt{3})=10+5\sqrt{3}+2\sqrt{3}+3$ (giving $13+7\sqrt{3}$) **[M1]**

$13+7\sqrt{3}$ **[A1]**

(b) Length $=\dfrac{12+4\sqrt{5}}{3+\sqrt{5}}$ **[M1]**

Multiply numerator and denominator by $3-\sqrt{5}$ **[M1]**

Numerator becomes $(12+4\sqrt{5})(3-\sqrt{5})=36-12\sqrt{5}+12\sqrt{5}-20=16$ **[M1]**

4 **[A1]**

[Back to Question] *[Total Marks: 7]*

291 **(a)** Multiplies numerator and denominator by $(5+\sqrt{3})$ **[M1]**

Denominator evaluates to $25-3$ or 22 **[M1]**

$5+\sqrt{3}$ **[A1]**

(b) $x(\sqrt{2}-1)=\sqrt{8}$ or $x(\sqrt{2}-1)=2\sqrt{2}$ **[M1]**

$x=\dfrac{2\sqrt{2}}{\sqrt{2}-1}$ **[M1]**

Multiplies numerator and denominator by $(\sqrt{2}+1)$ **[M1]**

$4+2\sqrt{2}$ **[A1]**

[Back to Question] *[Total Marks: 7]*

292 **(a)** $\dfrac{1}{2}\times(2+\sqrt{3})\times h=5+3\sqrt{3}$ **[M1]**

$h=\dfrac{10+6\sqrt{3}}{2+\sqrt{3}}$ **[M1]**

Multiply numerator and denominator by $(2-\sqrt{3})$ **[M1]**

$h=\dfrac{20-10\sqrt{3}+12\sqrt{3}-18}{4-3}=2+2\sqrt{3}$ **[A1]**

(b) $y^2=(2+\sqrt{3})^2+(2+2\sqrt{3})^2$ **[M1]**

$y^2 = (4 + 4\sqrt{3} + 3) + (4 + 8\sqrt{3} + 12)$ **[M1]**

$23 + 12\sqrt{3}$ **[A1]**

[Back to Question] *[Total Marks: 7]*

293 **(a)** $\dfrac{5 + 2\sqrt{3}}{2 + \sqrt{3}}$ **[M1]**

Multiplies numerator and denominator by $(2 - \sqrt{3})$ **[M1]**

$4 - \sqrt{3}$ **[A1]**

(b) $x\sqrt{5} - 2x = 2 + \sqrt{20}$ or $x(\sqrt{5} - 2) = \ldots$ **[M1]**

$x = \dfrac{2 + 2\sqrt{5}}{\sqrt{5} - 2}$ **[M1]**

Multiplies numerator and denominator by $(\sqrt{5} + 2)$ **[M1]**

$14 + 6\sqrt{5}$ **[A1]**

[Back to Question] *[Total Marks: 7]*

294 **(a)** $\dfrac{14 - 2\sqrt{5}}{3 + \sqrt{5}}$ **[M1]**

Multiplies numerator and denominator by $(3 - \sqrt{5})$ **[M1]**

Numerator expands to $42 - 14\sqrt{5} - 6\sqrt{5} + 10$ (or $52 - 20\sqrt{5}$) **[M1]**

$13 - 5\sqrt{5}$ **[A1]**

(b) Uses Pythagoras' theorem: $(3 + \sqrt{5})^2 + (13 - 5\sqrt{5})^2$ **[M1]**

$14 + 6\sqrt{5}$ or $294 - 130\sqrt{5}$ seen **[M1]**

$308 - 124\sqrt{5}$ with no errors seen **[A1]**

[Back to Question] *[Total Marks: 7]*

295 **(a)** Multiplies numerator and denominator by $(3 + \sqrt{3})$: $\dfrac{6(3 + \sqrt{3})}{(3 - \sqrt{3})(3 + \sqrt{3})}$ **[M1]**

Expands denominator correctly: $9 - 3$ or 6 **[M1]**

$\dfrac{6(3 + \sqrt{3})}{6}$ **[M1]**

$3 + \sqrt{3}$ **[A1]**

(b) $\dfrac{2\sqrt{6} + 3\sqrt{6}}{\sqrt{3}}$ or $\dfrac{5\sqrt{6}}{\sqrt{3}}$ or $5\sqrt{2}$ **[M1]**

$\dfrac{10\sqrt{2}}{2}$ or $5\sqrt{2}$ **[M1]**

0 (or $0\sqrt{2} + 0\sqrt{3}$) **[A1]**

[Back to Question] *[Total Marks: 7]*

296 **(a)** Area $= \frac{1}{2} \times$ base $\times$ height $\implies 5 + 2\sqrt{2} = \frac{1}{2} \times (3 + \sqrt{2}) \times h$ **[M1]**

$h = \frac{2(5 + 2\sqrt{2})}{3 + \sqrt{2}}$ **[A1]**

(b) Multiply numerator and denominator by $(3 - \sqrt{2})$: $\frac{2(5 + 2\sqrt{2})(3 - \sqrt{2})}{(3 + \sqrt{2})(3 - \sqrt{2})}$ **[M1]**

Denominator: $3^2 - (\sqrt{2})^2 = 9 - 2 = 7$ **[B1]**

Numerator: $2(15 - 5\sqrt{2} + 6\sqrt{2} - 4) = 2(11 + \sqrt{2}) = 22 + 2\sqrt{2}$ **[M1]**

Height $= \frac{22 + 2\sqrt{2}}{7}$ **[A1]**

(c) $\frac{1(\sqrt{7} + 2)}{(\sqrt{7} - 2)(\sqrt{7} + 2)}$ **[M1]**

$\frac{\sqrt{7} + 2}{7 - 4} = \frac{\sqrt{7} + 2}{3}$ **[A1]**

[Back to Question] *[Total Marks: 8]*

297 **(a)** $\frac{1}{2} \times (\sqrt{6} - \sqrt{2})(\sqrt{6} + \sqrt{2})$ **[M1]**

$\frac{1}{2}(6 - 2)$ **[M1]**

2 **[A1]**

(b) $(\text{Hypotenuse})^2 = (\sqrt{6} - \sqrt{2})^2 + (\sqrt{6} + \sqrt{2})^2$ **[M1]**

$(6 - 2\sqrt{12} + 2) + (6 + 2\sqrt{12} + 2)$ or $8 - 2\sqrt{12} + 8 + 2\sqrt{12}$ **[M1]**

$(\text{Hypotenuse})^2 = 16$ **[A1]**

Hypotenuse $= \sqrt{16} = 4$ **[A1]**

(c) Area scale factor $= \frac{18}{2} = 9$ **[M1]**

Length scale factor $= \sqrt{9} = 3$ **[M1]**

$4 \times 3 = 12$ **[A1]**

[Back to Question] *[Total Marks: 10]*

298 M1 for $3(-4)^2$ or $-5(-4)(2)$ or 48 or +40

88 **[A1]**

[Back to Question] *[Total Marks: 2]*

299 Correct substitution: $5(-2)^2 - 3(4)$ or $5(4) - 12$ **[M1]**

8 **[A1]**

[Back to Question] *[Total Marks: 2]*

300 $3(4) - 2(-5)$ or $12 - -10$ or $12 + 10$ **[M1]**

22 **[A1]**

[Back to Question] *[Total Marks: 2]*

301 Substitutes values correctly, e.g. $3(-3)^2 - 4(-5)$ **[M1]**

47 **[A1]**

[Back to Question] *[Total Marks: 2]*

302 Substitution of all three values correctly e.g. $\dfrac{3(5) - (-4)^2}{2(0.5)}$ **[M1]**

-1 **[A1]**

[Back to Question] *[Total Marks: 2]*

303 Substitute the given values into the formula: $v = 5 + (-2)(3)$ **[M1]**

$v = 5 - 6$ **[M1]**

-1 **[A1]**

[Back to Question] *[Total Marks: 3]*

304 **(a)** $4(-2) - 3(5)$ or $-8 - 15$ **[M1]**

-23 **[A1]**

(b) $\dfrac{5^2 - 7}{2(-3)}$ or $\dfrac{25 - 7}{-6}$ **[M1]**

-3 **[A1]**

[Back to Question] *[Total Marks: 4]*

305 **(a)** -6 or -20 seen **[M1]**

-26 **[A1]**

(b) $(-3)^2 = 9$ or $4(-3) = -12$ seen **[M1]**

-3 **[A1]**

[Back to Question] *[Total Marks: 4]*

306 **(a)** Substitutes values correctly: $5(3) - 2(-4)$ **[M1]**

$15 + 8 = 23$ **[A1]**

(b) Substitutes values: $\dfrac{(-4)^2 - (2)^3}{2(3)}$ **[M1]**

Evaluates numerator: $16 - 8 = 8$

$\frac{8}{6} = \frac{4}{3}$ or $1\frac{1}{3}$ **[A1]**

[Back to Question] *[Total Marks: 4]*

307 **(a)** **(a)** $40 + 25(3.5)$ or $40 + 87.5$ **[M1]**

127.50 **[A1]**

(b) $-12 + 9.8 \times 3$ **[M1]**

$-12 + 29.4$ **[M1]**

17.4 **[A1]**

[Back to Question] *[Total Marks: 5]*

308 **(a)** $40 + 25h$ **[B1]**

(b) $(-2)^3 = -8$ and $(-3)^2 = 9$ **[M1]**

$\sqrt{\frac{1}{16}} = \frac{1}{4}$ **[B1]**

-68 **[A1]**

[Back to Question] *[Total Marks: 4]*

309 Evaluates $p^{-2} = \left(\frac{1}{2}\right)^{-2} = 4$ **[B1]**

Evaluates $q^3 = (-2)^3 = -8$ **[B1]**

Evaluates $\sqrt[3]{r} = \sqrt[3]{-\frac{1}{8}} = -\frac{1}{2}$ **[B1]**

8 **[A1]**

[Back to Question] *[Total Marks: 4]*

310 **(a)** $d = \sqrt{(-8-4)^2 + (2--3)^2}$ or $\sqrt{144+25}$ **[M1]**

13 **[A1]**

(b) $17 = \sqrt{(3-k)^2 + (-7-8)^2}$ **[M1]**

$289 = (3-k)^2 + 225$ **[M1]**

$(3-k)^2 = 64$ or $k^2 - 6k - 55 = 0$ **[M1]**

$k = -5$ and $k = 11$ **[A1]**

[Back to Question] *[Total Marks: 6]*

311 **(a)** States $A =$ Curved Surface Area $+$ Base Area $= 2\pi rh + \pi r^2$ **[M1]**

Factorises πr out to obtain $\pi r(2h + r)$ **[A1]**

(b) **(i)** Substitutes $r = 4$ and $h = 6$: $A = \pi(4)(2(6) + 4)$ **[M1]**

$A = 4\pi(12 + 4) = 64\pi$ **[A1]**

(ii) Rearranges volume formula: $h = \dfrac{V}{\pi r^2}$ **[M1]**

Substitutes h into area formula: $A = 2\pi r\left(\dfrac{V}{\pi r^2}\right) + \pi r^2$ **[M1]**

Simplifies first term: $\dfrac{2\pi rV}{\pi r^2} = \dfrac{2V}{r}$, resulting in $A = \dfrac{2V}{r} + \pi r^2$ **[A1]**

[Back to Question] *[Total Marks: 7]*

312 **(a)** Substitute $x = -2$ and $y = 4$: $P = \dfrac{2(-2)^2 - 3(4)}{-2 - 4}$ **[M1]**

$P = \dfrac{2(4) - 12}{-6} = \dfrac{8 - 12}{-6}$ **[M1]**

$\dfrac{2}{3}$ (or equivalent) **[A1]**

(b) Multiply both sides by $(x - y)$: $P(x - y) = 2x^2 - 3y$ **[M1]**

Expand: $Px - Py = 2x^2 - 3y$ **[M1]**

Collect terms with y on one side: $3y - Py = 2x^2 - Px$ **[M1]**

Factorise and divide: $y(3 - P) = 2x^2 - Px \implies y = \dfrac{2x^2 - Px}{3 - P}$ (or equivalent) **[A1]**

[Back to Question] *[Total Marks: 7]*

313 **(a)** **(i)** $\dfrac{6^2 - (-2)^2}{6(-2)}$ **[M1]**

$-\dfrac{8}{3}$ **[A1]**

(ii) $\dfrac{0.5^2 - 4^2}{0.5(4)}$ **[M1]**

$\dfrac{-15.75}{2}$ **[M1]**

$-\dfrac{63}{8}$ **[A1]**

(b) $\dfrac{x^2 - 2^2}{2x} = \dfrac{3}{2}$ **[M1]**

$2(x^2 - 4) = 6x$ (or $x^2 - 3x - 4 = 0$) **[M1]**

4 **[A1]**

[Back to Question] *[Total Marks: 8]*

314 **(a)** $C + Rh$ or $C + hR$ **[B1]**

(b) M1 for $45 + 32.50 \times 3.5$ or $45 + 113.75$

158.75 **[A1]**

[Back to Question] *[Total Marks: 3]*

315 Substitutes values correctly: $\dfrac{\sqrt{12.5-3.84}}{(-1.2)^3}$ **[M1]**

Evaluates numerator to 2.94... or denominator to -1.728 **[M1]**

-1.70 (accept $-1.703...$ rounded strictly to 3 sig figs) **[A1]**

[Back to Question] *[Total Marks: 3]*

316 $\pi \times (4.5)^2 \times 8.2 + \dfrac{2}{3} \times \pi \times (4.5)^3$ **[M1]**

$521.6... + 190.8...$ or $712.5...$ **[M1]**

713 **[A1]**

[Back to Question] *[Total Marks: 3]*

317 (a) $45 + 32h$ (or equivalent) **[B1]**

(b) 157 **[B1]**

(c) Correct substitution: $(-3)^3 - 4(-3)(2.5)$ or $-27 - (-30)$ **[M1]**

3 **[A1]**

[Back to Question] *[Total Marks: 4]*

318 (a) $V = \dfrac{1}{3}\pi(6.4)^2(12.6)$ **[M1]**

540.5 **[A1]**

(b) $800 = \dfrac{1}{3}\pi r^2(14)$ **[M1]**

$r^2 = \dfrac{2400}{14\pi}$ or $54.567\ldots$ **[M1]**

7.39 **[A1]**

[Back to Question] *[Total Marks: 5]*

319 (a) $85(7) + 0.24(450)$ **[M1]**

703 **[A1]**

(b) $1475.60 = 85(14) + 0.24m$ **[M1]**

$1475.60 - 1190 = 0.24m$ (or $285.60 = 0.24m$) **[M1]**

1190 **[A1]**

[Back to Question] *[Total Marks: 5]*

320 **(a)** Substitute $h = 3.5$: $C = 45 + 35(3.5)$ **[M1]**

167.5 **[A1]**

(b) Substitute $C = 325$: $325 = 45 + 35h$ **[M1]**

$280 = 35h$ **[M1]**

8 **[A1]**

[Back to Question] *[Total Marks: 5]*

321 **(a)** Substitutes $h = 4.2$ and $a = 6.5$ into formula: $V = \frac{1}{6}\pi(4.2)(3(6.5)^2 + 4.2^2)$ **[M1]**

$V = 317.53... = 317.5$ **[A1]**

(b) Multiplies by 6 and divides by πh: $\frac{6V}{\pi h} = 3a^2 + h^2$ **[M1]**

Subtracts h^2: $\frac{6V}{\pi h} - h^2 = 3a^2$ **[M1]**

$a^2 = \frac{2V}{\pi h} - \frac{h^2}{3}$ or $a^2 = \frac{1}{3}\left(\frac{6V}{\pi h} - h^2\right)$ **[A1]**

(c) $a = \sqrt{\frac{2(150)}{\pi(3)} - \frac{3^2}{3}} = \sqrt{31.83... - 3} = \sqrt{28.83...} = 5.37$ **[B1]**

[Back to Question] *[Total Marks: 6]*

322 Numerator correctly evaluated: $-4.5 - (-3.2)^2 = -4.5 - 10.24 = -14.74$ **[B1]**

Denominator correctly evaluated: $1.8 - \sqrt{0.64} = 1.8 - 0.8 = 1$ **[B1]**

-14.74 **[B1]**

[Back to Question] *[Total Marks: 3]*

323 **(a)** **(a)** $5.2(4.5) + 0.5(-9.8)(4.5)^2$ **[M1]**

-75.825 **[A1]**

(b) $-9.8 = 4.9t + \frac{1}{2}(-9.8)t^2$ **[M1]**

$4.9t^2 - 4.9t - 9.8 = 0$ or $t^2 - t - 2 = 0$ **[M1]**

$t = 2$ (rejects -1 as time must be positive) **[A1]**

[Back to Question] *[Total Marks: 5]*

324 **(a)** M1 for correctly substituting the values into the formula: $\frac{1}{3}\pi(12.5)(4.2^2 + 4.2 \times 7.8 + 7.8^2)$

M1 for 111.24 or 463.5π or 1456.1...

1460 **[A1]**

(b) M1 for correctly substituting the values into the square root part: $\sqrt{(7.8 - 4.2)^2 + 12.5^2}$

M1 for $\sqrt{169.21}$ or 13.008...

736.9 **[A1]**

[Back to Question] *[Total Marks: 6]*

325 At least 3 terms correct from $6x^2 + 15x - 4x - 10$ **[M1]**

$6x^2 + 11x - 10$ **[A1]**

[Back to Question] *[Total Marks: 2]*

326 (a) Expansion to $6x - 3 - 2x - 8$ **[M1]**

$4x - 11$ **[A1]**

(b) Extracting a partial common factor, e.g. $5xy(3x - 4y)$ or $x(15xy - 20y^2)$ or $5(3x^2y - 4xy^2)$ **[M1]**

$5xy(3x - 4y)$ **[A1]**

[Back to Question] *[Total Marks: 4]*

327 (a) M1 for $8a - 12b$ or $-3a + 15b$ seen

$5a + 3b$ **[A1]**

(b) M1 for a correct partial factorisation, e.g., $6x(2xy - 3y^2)$ or $2xy(6x - 9y)$

$6xy(2x - 3y)$ **[A1]**

[Back to Question] *[Total Marks: 4]*

328 (a) $8x - 12y - 3x + 15y$ **[M1]**

$5x + 3y$ **[A1]**

(b) $7ab$ extracted as a common factor **[M1]**

$7ab(2a - 3b)$ **[A1]**

[Back to Question] *[Total Marks: 4]*

329 (a) $12x - 8 - 2x + 10$ **[M1]**

$10x + 2$ **[A1]**

(b) $5ab(3a - 2b)$ **[B2]**

If not B2, then B1 for $5a(3ab - 2b^2)$ or $5b(3a^2 - 2ab)$ or $ab(15a - 10b)$ **[B1]**

[Back to Question] *[Total Marks: 4]*

330 (a) $6x - 15 - 2x - 8$ **[M1]**

$4x - 23$ **[A1]**

(b) $4xy(2x-3y)$ or $4x(2xy-3y^2)$ or $4y(2x^2-3xy)$ or $xy(8x-12y)$ **[M1]**

$4xy(2x-3y)$ **[A1]**

[Back to Question] *[Total Marks: 4]*

331 **(a)** Correct expansion: $6x-15-4x+8$ **[M1]**

$2x-7$ **[A1]**

(b) $6xy(2x-3y)$ **[B2]**

Correct partial factorisation e.g., $6x(2xy-3y^2)$ or $2xy(6x-9y)$ **[M1]**

[Back to Question] *[Total Marks: 4]*

332 **(a)** $5a$ or $+3b$ **[B1]**

$5a+3b$ **[A1]**

(b) $6xy(2x-3y)$ **[B2]**

If B0, allow [M1] for a correct partial factorisation, e.g. $6x(2xy-3y^2)$ or $3xy(4x-6y)$

[Back to Question] *[Total Marks: 4]*

333 **(a)** Expansion to give $6x-15$ and $-4x+12$ **[M1]**

$2x-3$ **[A1]**

(b) $7xy(2x+3y)$ **[B2]**

Partial factorisation, e.g. $7x(2xy+3y^2)$ or $xy(14x+21y)$ **[B1]**

[Back to Question] *[Total Marks: 4]*

334 **(a)** $3(4x^2-9y^2)$ or $(6x-9y)(2x+3y)$ or $(2x-3y)(6x+9y)$ **[M1]**

$3(2x-3y)(2x+3y)$ **[A1]**

(b) $2a(x-3y)+b(x-3y)$ or $x(2a+b)-3y(2a+b)$ **[M1]**

$(2a+b)(x-3y)$ **[A1]**

[Back to Question] *[Total Marks: 4]*

335 **(a)** $2x^2+10x-3x-15$ **[M1]**

$2x^2+7x-15$ **[A1]**

(b) $(y-4x)(y+4x)$ **[B1]**

(c) $(p+q)(p-5q)$ **[B2]**

If B0, allow [M1] for $(p+aq)(p+bq)$ where $ab=-5$ or $a+b=-4$

[Back to Question] *[Total Marks: 5]*

336 **(a)** $2x^2 + 10x - 3x - 15$ **[M1]**

$2x^2 + 7x - 15$ **[A1]**

(b) $c(a - 3b) + 2d(a - 3b)$ or $a(c + 2d) - 3b(c + 2d)$ **[M1]**

$(a - 3b)(c + 2d)$ **[A1]**

(c) $(4x - 5y)(4x + 5y)$ **[B1]**

[Back to Question] *[Total Marks: 5]*

337 **(a)** M1 for 3 correct terms out of $2x^2 + 8x - 3x - 12$

$2x^2 + 5x - 12$ **[A1]**

(b) M1 for $x(a - b)$ or $-3y(a - b)$ seen

$(x - 3y)(a - b)$ **[A1]**

(c) $(7x - 4y)(7x + 4y)$ **[B2]**

If 0 scored, allow M1 for $(7x - 4y)(7x - 4y)$ or $(7x + 4y)(7x + 4y)$

[Back to Question] *[Total Marks: 6]*

338 **(a)** $(x + a)(x + b)$ where $ab = 15$ or $a + b = -8$ **[M1]**

$(x - 3)(x - 5)$ **[A1]**

(b) $(2x + a)(x + b)$ where $ab = -12$ or $2b + a = -5$ **[M1]**

$(2x + 3)(x - 4)$ **[A1]**

(c) $a(x - 3) + 2b(x - 3)$ or $x(a + 2b) - 3(a + 2b)$ **[M1]**

$(a + 2b)(x - 3)$ **[A1]**

[Back to Question] *[Total Marks: 6]*

339 **(a)** $6x^2 + 2x - 15x - 5$ (at least 3 correct terms) **[M1]**

$6x^2 - 13x - 5$ **[A1]**

(b) $(x + a)(x + b)$ where $ab = -18$ or $a + b = -7$ **[M1]**

$(x - 9)(x + 2)$ **[A1]**

(c) $2(9x^2 - 25y^2)$ **[M1]**

$(3x - 5y)(3x + 5y)$ seen **[M1]**

$2(3x - 5y)(3x + 5y)$ **[A1]**

[Back to Question] *[Total Marks: 7]*

340 **(a)** Correct expansion of any two brackets, e.g. $2x^2 - x - 6$ or $2x^2 - 7x - 15$ or $x^2 - 7x + 10$ **[M1]**

Correctly multiplies their 3-term quadratic by the remaining bracket to get at least 4

correct terms **[M1]**

$2x^3 - 11x^2 - x + 30$ **[A1]**

(b) Extracts common factor x to give $x(12x^2 + 17x - 5)$ **[M1]**

Correctly factorises the quadratic expression to $(4x - 1)(3x + 5)$ **[M1]**

$x(4x - 1)(3x + 5)$ **[A1]**

[Back to Question] *[Total Marks: 6]*

341 **(a)** Taking out common factor to reach $2(9x^2 - 25y^2)$ **[M1]**

Factorising difference of squares to reach $(3x - 5y)(3x + 5y)$ **[M1]**

$2(3x - 5y)(3x + 5y)$ **[A1]**

(b) Taking out common factor x to reach $x(3x^2 + 10x - 8)$ **[M1]**

Factorising quadratic to reach $(3x - 2)(x + 4)$ **[M1]**

$x(3x - 2)(x + 4)$ **[A1]**

[Back to Question] *[Total Marks: 6]*

342 **(a)** Correct expansion of two brackets, e.g. $(x^2 + x - 6)(2x - 1)$ or $(x - 2)(2x^2 + 5x - 3)$ **[M1]**

Multiplication of quadratic by third bracket yielding 6 terms, e.g.
$2x^3 - x^2 + 2x^2 - x - 12x + 6$ **[M1]**

$2x^3 + x^2 - 13x + 6$ **[A1]**

(b) **(i)** $(3x + a)(2x + b)$ where $ab = -10$ or $2a + 3b = -11$ **[M1]**

$(3x + 2)(2x - 5)$ **[A1]**

(ii) $a(x - y) + 3c(x - y)$ or $x(a + 3c) - y(a + 3c)$ **[M1]**

$(a + 3c)(x - y)$ **[A1]**

[Back to Question] *[Total Marks: 7]*

343 **(a)** $x(c - 5d) + 3y(c - 5d)$ or $c(x + 3y) - 5d(x + 3y)$ **[M1]**

$(x + 3y)(c - 5d)$ **[A1]**

(b) $2x(x^2 - 9)$ **[M1]**

$2x(x - 3)(x + 3)$ **[A1]**

(c) $2x(6x^2 - 7x - 5)$ **[M1]**

$2x(3x - 5)(2x + 1)$ **[A2]**

If not A2, then A1 for $(3x - 5)(2x + 1)$ or $2x(3x + 5)(2x - 1)$ **[A1]**

[Back to Question] *[Total Marks: 7]*

344 **(a)** Expands any two brackets, e.g. $(2x^2 - 7x - 4)(x + 3)$ **[M1]**

Multiplies their quadratic by the third bracket, e.g. $2x^3 - 7x^2 - 4x + 6x^2 - 21x - 12$ **[M1]**

$2x^3 - x^2 - 25x - 12$ **[A1]**

(b) $2a(x + y) - 3b(y + x)$ or $x(2a - 3b) + y(2a - 3b)$ **[M1]**

$(x + y)(2a - 3b)$ **[A1]**

(c) $(2x + 5)(3x - 4)$ **[B2]**

If B0, allow [M1] for $(2x + a)(3x + b)$ where $ab = -20$ or $3a + 2b = 7$

[Back to Question] *[Total Marks: 7]*

345 **(a)** $(2a - b)(3x - y)$ **[B2]**

Correct partial factorisation e.g., $2a(3x - y) - b(3x - y)$ or $3x(2a - b) - y(2a - b)$ **[M1]**

(b) **(i)** Extracts common factor of 2: $2(25x^2 - 4y^2)$ **[M1]**

Factorises difference of two squares: $(5x - 2y)(5x + 2y)$ **[M1]**

$2(5x - 2y)(5x + 2y)$ **[A1]**

(ii) Extracts common factor of x: $x(2x^2 + 5x - 12)$ **[M1]**

Factorises quadratic: $(2x - 3)(x + 4)$ **[M1]**

$x(2x - 3)(x + 4)$ **[A1]**

[Back to Question] *[Total Marks: 8]*

346 **(a)** Correct expansion of any two brackets, e.g., $(2x^2 - 5x - 3)$ or $(2x^2 + 9x + 4)$ or $(x^2 + x - 12)$ **[M1]**

Correct expansion of their quadratic with the third bracket, producing at least 5 terms, e.g., $2x^3 + 8x^2 - 5x^2 - 20x - 3x - 12$ **[M1]**

$2x^3 + 3x^2 - 23x - 12$ **[A1]**

(b) $(2x + 3)(x - 5)$ **[B2]**

Accept $(2x + a)(x + b)$ where $ab = -15$ or $2b + a = -7$ **[M1]**

[Back to Question] *[Total Marks: 5]*

347 **(a)** Four terms with at least three correct from $2x^2 + 10x - 3x - 15$ **[M1]**

$2x^2 + 7x - 15$ **[A1]**

(b) $(x + a)(x + b)$ where $ab = -18$ or $a + b = -7$ **[M1]**

$(x - 9)(x + 2)$ **[A1]**

(c) $(2x + a)(2x - a)$ or $(cx + 5y)(cx - 5y)$ **[M1]**

$(2x - 5y)(2x + 5y)$ **[A1]**

[Back to Question] *[Total Marks: 6]*

348 **(a)** $(x-4)(2x^2+7x+6)$ or $(2x^2-5x-12)(x+2)$ or $(x^2-2x-8)(2x+3)$ **[M1]**

$2x^3+7x^2+6x-8x^2-28x-24$ (allow one error in expansion) **[M1]**

$2x^3-x^2-22x-24$ **[A1]**

(b) $(4x-7y)(4x+7y)$ **[B2]**

If not B2, then B1 for $(4x-ky)(4x+ky)$ or $(kx-7y)(kx+7y)$ where $k \neq 0$ **[B1]**

(c) $(3x+2)(x-4)$ **[B2]**

If not B2, then B1 for $(3x+a)(x+b)$ where $ab=-8$ or $3b+a=-10$ **[B1]**

[Back to Question] *[Total Marks: 7]*

349 **(a)** Expands any two brackets correctly, e.g., $(2x^2-5x-12)(x+1)$ or $(x-4)(2x^2+5x+3)$ or $(x^2-3x-4)(2x+3)$ **[M1]**

Multiplies their resulting quadratic by the third bracket, allowing one error **[M1]**

$2x^3-3x^2-17x-12$ **[A1]**

(b) $(3x+a)(2x+b)$ or $(6x+c)(x+d)$ where $ab=-10$ or $cd=-10$ **[M1]**

$(3x+2)(2x-5)$ **[A1]**

(c) $x^2(x+3)-4(x+3)$ or $x(x^2-4)+3(x^2-4)$ **[M1]**

$(x^2-4)(x+3)$ **[M1]**

$(x-2)(x+2)(x+3)$ **[A1]**

[Back to Question] *[Total Marks: 8]*

350 **(a)** M1 for correctly expanding any pair of brackets, e.g., $2x^2-3x-2$ or $2x^2+7x+3$ or x^2+x-6

M1 for multiplying their quadratic expression by the third bracket, allowing one error in terms

$2x^3+3x^2-11x-6$ **[A1]**

(b) M1 for $(3x+a)(2x+b)$ where $ab=-10$ or $3b+2a=-11$

$(3x+2)(2x-5)$ **[A1]**

(c) M1 for partial factorisation extracting x: $x(3x^2+5x-2)$

M1 for factorising the quadratic factor: $(3x-1)(x+2)$

$x(3x-1)(x+2)$ **[A1]**

[Back to Question] *[Total Marks: 8]*

351 **(a)** $(x^2+x-6)(2x-1)$ or $(x-2)(2x^2+5x-3)$ or $(x+3)(2x^2-5x+2)$ **[M1]**

$2x^3-x^2+2x^2-x-12x+6$ (allow one error in expansion) **[M1]**

$2x^3+x^2-13x+6$ **[A1]**

(b) $(3x \pm a)(2x \pm b)$ where $ab=10$ or $6x^2-15x+4x-10$ **[M1]**

$(3x+2)(2x-5)$ **[A1]**

(c) $2x(x^2-9)$ **[M1]**

$(x-3)(x+3)$ seen **[M1]**

$2x(x-3)(x+3)$ **[A1]**

[Back to Question] *[Total Marks: 8]*

352 Correct common denominator of 15 with unsimplified numerator $3(2x)-5(x-1)$ **[M1]**

$\frac{x+5}{15}$ **[A1]**

[Back to Question] *[Total Marks: 2]*

353 M1 for a correct common denominator with at least one correct numerator, e.g.,
$\frac{5(2x-1)+3(x+4)}{15}$ or $\frac{10x-5}{15}+\frac{3x+12}{15}$

$\frac{13x+7}{15}$ **[A1]**

[Back to Question] *[Total Marks: 2]*

354 $\frac{5(3x)-4(x-2)}{20}$ or $\frac{15x}{20}-\frac{4x-8}{20}$ **[M1]**

$\frac{11x+8}{20}$ **[A1]**

[Back to Question] *[Total Marks: 2]*

355 Uses common denominator $(x+2)(3x-1)$ **[B1]**

Correct unsimplified numerator $5(3x-1)+2(x+2)$ **[M1]**

$\frac{17x-1}{(x+2)(3x-1)}$ or $\frac{17x-1}{3x^2+5x-2}$ **[A1]**

[Back to Question] *[Total Marks: 3]*

356 Writes fractions over a common denominator of 20 **[M1]**

Correct unsimplified numerator: $5(x+3)-4(2x-1)$ or $5x+15-8x+4$ **[M1]**

$\frac{19-3x}{20}$ or $\frac{-3x+19}{20}$ **[A1]**

[Back to Question] *[Total Marks: 3]*

357 Common denominator of 12 and correct numerator formulation: $\frac{3(3x)-4(x-2)}{12}$ **[M1]**

Numerator expanded correctly to $9x-4x+8$ **[M1]**

$\frac{5x+8}{12}$ **[A1]**

[Back to Question] *[Total Marks: 3]*

358 $\frac{5(2x)-3(x-4)}{15}$ or $\frac{10x}{15}-\frac{3(x-4)}{15}$ **[M1]**

$10x-3x+12$ or better as numerator (allow one sign error for M1, e.g., $10x-3x-12$) **[M1]**

$\frac{7x+12}{15}$ **[A1]**

[Back to Question] *[Total Marks: 3]*

359 **(a)** Uses a common denominator of 12, e.g. $\frac{4(2x)+3(x-1)}{12}$ **[M1]**

$\frac{11x-3}{12}$ **[A1]**

(b) $3x^2$ or $\frac{1}{y^2}$ seen **[M1]**

$\frac{3x^2}{y^2}$ or $3x^2y^{-2}$ **[A1]**

[Back to Question] *[Total Marks: 4]*

360 **(a)** $\frac{15x+8x}{20}$ **[M1]**

$\frac{23x}{20}$ **[A1]**

(b) $\frac{5a^2b}{12}\times\frac{9}{10ab^2}$ **[M1]**

$\frac{3a}{8b}$ **[A1]**

[Back to Question] *[Total Marks: 4]*

361 $(x-4)(x+4)$ seen **[B1]**

$(x-4)(x-1)$ seen **[B1]**

$\frac{x+4}{x-1}$ **[A1]**

[Back to Question] *[Total Marks: 3]*

362 $3(x^2-4)$ or $3(x-2)(x+2)$ **[B1]**

$(x+5)(x-2)$ [B2] (or [M1] for $(x+a)(x+b)$ where $ab=-10$ or $a+b=3$)

$\frac{3(x+2)}{x+5}$ or $\frac{3x+6}{x+5}$ **[A1]**

[Back to Question] *[Total Marks: 4]*

363 **(a)** $\dfrac{3(x+5)-2(x-2)}{(x-2)(x+5)}$ **[M1]**

$3x+15-2x+4$ correctly expanded on numerator **[M1]**

$\dfrac{x+19}{(x-2)(x+5)}$ or $\dfrac{x+19}{x^2+3x-10}$ **[A1]**

(b) $(x-4)(x+4)$ seen for numerator **[M1]**

$(x-4)(x+2)$ seen for denominator **[M1]**

$\dfrac{x+4}{x+2}$ **[A1]**

[Back to Question] *[Total Marks: 6]*

364 **(a)** M1 for factorising the numerator correctly: $(x-3)(x+3)$

M1 for factorising the denominator correctly: $(2x-1)(x+3)$

$\dfrac{x-3}{2x-1}$ **[A1]**

(b) M1 for the correct common denominator $(x-2)(x+1)$

M1 for the correct unsimplified numerator $4(x+1)-3(x-2)$

$\dfrac{x+10}{(x-2)(x+1)}$ or $\dfrac{x+10}{x^2-x-2}$ **[A1]**

[Back to Question] *[Total Marks: 6]*

365 Factorises denominator to $(2x-3)(2x+3)$ **[B1]**

Attempts to factorise numerator to $(2x+a)(x+b)$ where $ab=-12$ or $2b+a=5$ **[M1]**

Correctly factorises numerator to $(2x-3)(x+4)$ **[A1]**

$\dfrac{x+4}{2x+3}$ **[A1]**

[Back to Question] *[Total Marks: 4]*

366 $(3x+1)(x-2)$ **[B2]**

If not B2, then B1 for $(3x+a)(x+b)$ where $ab=-2$ or $3b+a=-5$ **[B1]**

$(x-2)(x+2)$ **[B1]**

$(x-2)(x+1)$ **[B1]**

$\dfrac{3x^2-5x-2}{x^2-4}\times\dfrac{x^2-x-2}{3x+1}$ or equivalent correct inverted product **[M1]**

$\dfrac{(x-2)(x+1)}{x+2}$ or $\dfrac{x^2-x-2}{x+2}$ **[A1]**

[Back to Question] *[Total Marks: 6]*

367 **(a)** Factorising numerator: $(2x+1)(x-3)$ **[B1]**

Factorising denominator: $(2x-3)(2x+3)$ **[B1]**

Note: no common factor cancels; the expression is already in its simplest form

Final answer: $\dfrac{2x^2-5x-3}{4x^2-9}$ or $\dfrac{(2x+1)(x-3)}{(2x-3)(2x+3)}$ **[A1]**

(b) Recognising $x^2-x-2=(x-2)(x+1)$ **[M1]**

Writing all over a common denominator of $(x-2)(x+1)$: $\dfrac{3(x+1)+4(x-2)-2x}{(x-2)(x+1)}$ **[M2]**

Simplifying numerator: $3x+3+4x-8-2x=5x-5$ **[M1]**

$\dfrac{5(x-1)}{(x-2)(x+1)}$ **[A1]**

[Back to Question] *[Total Marks: 7]*

368 Factorising numerator to $(x-3)(x+3)$ **[B1]**

Factorising denominator to $(2x-1)(x+3)$ **[B1]**

$\dfrac{x-3}{2x-1}$ **[A1]**

[Back to Question] *[Total Marks: 3]*

369 Denominator of $(x-2)(2x+1)$ **[B1]**

Correct unsimplified numerator: $5(2x+1)-3(x-2)$ **[M1]**

$\dfrac{7x+11}{(x-2)(2x+1)}$ **[A1]**

[Back to Question] *[Total Marks: 3]*

370 Numerator factorised as $(2x-3)(x-3)$ **[B2]**

Allow [M1] for $(2x+a)(x+b)$ where $ab=9$ or $2b+a=-9$

Denominator factorised as $2(x-3)(x+3)$ or $2(x^2-9)$ or $(2x-6)(x+3)$ **[B1]**

$\dfrac{2x-3}{2(x+3)}$ or $\dfrac{2x-3}{2x+6}$ **[A1]**

[Back to Question] *[Total Marks: 4]*

371 $2(x-2)(x+2)$ **[B2]**

If not B2, then B1 for $2(x^2-4)$ or $(2x-4)(x+2)$ **[B1]**

$(x-2)(x+3)$ **[B2]**

If not B2, then B1 for $(x+a)(x+b)$ where $ab=-6$ or $a+b=1$ **[B1]**

$\dfrac{2(x+2)}{x+3}$ or $\dfrac{2x+4}{x+3}$ **[A1]**

[Back to Question] *[Total Marks: 5]*

372 **(a)** Common denominator $(2x-1)(x+4)$ seen **[M1]**

Numerator: $(x+3)(x+4)-(x-2)(2x-1)$ **[M1]**

$(x^2+7x+12)-(2x^2-5x+2)$ correctly simplified to $-x^2+12x+10$ **[A1]**

(b) Factorising numerator to $(x-4)(x+4)$ or denominator to $x(x+4)$ **[M1]**

$\dfrac{x-4}{x}$ **[A1]**

[Back to Question] *[Total Marks: 5]*

373 Factorising x^2-25 to $(x-5)(x+5)$ and x^2-3x to $x(x-3)$ **[B1]**

Factorising x^2+6x+5 to $(x+1)(x+5)$ and $2x-6$ to $2(x-3)$ **[B1]**

Showing multiplication by the reciprocal, e.g. $\dfrac{(x-5)(x+5)}{x(x-3)}\times\dfrac{2(x-3)}{(x+1)(x+5)}$ **[M1]**

$\dfrac{2(x-5)}{x(x+1)}$ (or equivalent expanded form $\dfrac{2x-10}{x^2+x}$) **[A1]**

[Back to Question] *[Total Marks: 4]*

374 **(a)** M1 for factorising $2x^2-8x$ to $2x(x-4)$

M1 for factorising x^2-16 to $(x-4)(x+4)$

M1 for inverting the second fraction and factorising, i.e., multiplying by $\dfrac{3(x+4)}{6x}$

1 **[A1]**

(b) M1 for writing the denominator x^2-9 as $(x-3)(x+3)$

M1 for common denominator $(x-3)(x+3)$ and correct numerator $5(x+3)+2x+1$

$\dfrac{7x+16}{(x-3)(x+3)}$ or $\dfrac{7x+16}{x^2-9}$ **[A1]**

[Back to Question] *[Total Marks: 7]*

375 **(a)** $\dfrac{4(2x-1)-3(x-3)}{(x-3)(2x-1)}$ **[M1]**

Numerator expanded correctly to $8x-4-3x+9$ **[M1]**

$\dfrac{5x+5}{(x-3)(2x-1)}$ or $\dfrac{5(x+1)}{(x-3)(2x-1)}$ **[A1]**

(b) Multiplies by reciprocal: $\times\dfrac{4x+2}{x+5}$ **[M1]**

Factorises x^2-25 to $(x-5)(x+5)$ or $4x+2$ to $2(2x+1)$ **[M1]**

Factorises denominator $2x^2-9x-5$ to $(2x+1)(x-5)$ **[M1]**

2 **[A1]**

[Back to Question] *[Total Marks: 7]*

376 **(a)** Uses common denominator $(x-2)(x+2)$ or x^2-4 **[B1]**

Correct unsimplified numerator $3(x+2)-(x+5)$ **[M1]**

$\dfrac{2x+1}{(x-2)(x+2)}$ or $\dfrac{2x+1}{x^2-4}$ **[A1]**

(b) Factorises $4x^2-25$ as $(2x-5)(2x+5)$ **[B1]**

Factorises denominators as $x(2x-5)$ and $2x(2x+5)$ **[B2]**

Allow [B1] if only one denominator is factorised correctly

Correct cancellation leading to final answer $3x$ **[A1]**

[Back to Question] *[Total Marks: 7]*

377 **(a)** $\dfrac{5(2x+3)+4(x-2)}{(x-2)(2x+3)}$ **[M1]**

$10x+15+4x-8$ **[M1]**

$\dfrac{14x+7}{(x-2)(2x+3)}$ or $\dfrac{7(2x+1)}{(x-2)(2x+3)}$ **[A1]**

(b) $\dfrac{2x^2-5x-12}{4x^2-9}\times\dfrac{2x-3}{x-4}$ **[M1]**

$(2x+3)(x-4)$ **[B1]**

$(2x-3)(2x+3)$ **[B1]**

1 **[A1]**

[Back to Question] *[Total Marks: 7]*

378 **(a)** $(2x-1)(x+3)$ seen for numerator **[M1]**

$(2x-1)(2x+1)$ seen for denominator **[M1]**

Identifies and cancels the common factor $(2x-1)$ **[M1]**

$\dfrac{x+3}{2x+1}$ **[A1]**

(b) Factorises denominator: $x^2-4x+3=(x-1)(x-3)$ **[M1]**

Writes with a common denominator: $\dfrac{5(x-1)+(x+2)}{(x-1)(x-3)}$ **[M1]**

Simplifies numerator: $5x-5+x+2=6x-3$ **[M1]**

$\dfrac{3(2x-1)}{(x-1)(x-3)}$ or $\dfrac{6x-3}{x^2-4x+3}$ **[A1]**

[Back to Question] *[Total Marks: 8]*

379 If incorrect, award [B1] for any two of 12, x^8, or y^4 correct in a product

$12x^8y^4$ **[B2]**

[Back to Question] *[Total Marks: 2]*

380 **(a)** $12x^{-3}$ or $\frac{12}{x^3}$ **[B1]**

(b) $\frac{2x^3}{3}$ or $\frac{2}{3}x^3$ **[B2]**

Correct cube root of coefficient $\frac{2}{3}$ OR correct index x^3 seen **[M1]**

[Back to Question] *[Total Marks: 3]*

381 **(a)** $20x$ **[B1]**

(b) Numerator $4x^8$ or denominator y^6 seen **[M1]**

$\frac{4x^8}{y^6}$ **[A1]**

(c) -3 **[B1]**

[Back to Question] *[Total Marks: 4]*

382 **(a)** M1 for any two of 12, x^8 or y^4 seen in a product

$12x^8y^4$ **[A1]**

(b) M1 for $\frac{1}{16^{\frac{1}{4}}}$ or 2^{-1} or $\frac{1}{\sqrt[4]{16}}$

$\frac{1}{2}$ **[A1]**

[Back to Question] *[Total Marks: 4]*

383 **(a)** 15 or x^2 in a product **[M1]**

$15x^2$ **[A1]**

(b) $\sqrt[4]{16}$ or 2^3 or $4096^{\frac{1}{4}}$ **[M1]**

8 **[A1]**

[Back to Question] *[Total Marks: 4]*

384 **(a)** y^3 **[B1]**

(b) $\frac{1}{25}$ or 0.04 **[B1]**

(c) $8x^{12}$ **[B2]**

If B0, allow [M1] for $8x^k$ or kx^{12} where $k \neq 0$

[Back to Question] *[Total Marks: 4]*

385 **(a)** -3 **[B1]**

(b) $2y^3$ **[B2]**

If not B2, then B1 for 2 or y^3 seen **[B1]**

(c) $\frac{1}{8}$ **[B1]**

[Back to Question] *[Total Marks: 4]*

386 **(a)** w^{11} **[B1]**

(b) 3^2 or 9 seen, or x^6 seen **[M1]**

$9x^6$ **[A1]**

(c) $\frac{15}{5} = 3$ or $y^{8-(-2)}$ **[M1]**

$3y^{10}$ **[A1]**

[Back to Question] *[Total Marks: 5]*

387 **(a)** p^{11} **[B1]**

(b) $9q^6$ **[B2]**

Partial correct answer $9q^k$ or cq^6 **[B1]**

(c) $(\sqrt[3]{27})^2$ or 3^2 or $\sqrt[3]{729}$ **[M1]**

9 **[A1]**

[Back to Question] *[Total Marks: 5]*

388 **(a)** $4x^3y^4$ **[B2]**

If not B2, then B1 for two of 4, x^3, y^4 correct **[B1]**

(b) $(3^3)^{x-1} = (3^2)^{2x+1}$ or $3^{3(x-1)} = 3^{2(2x+1)}$ **[M1]**

$3x - 3 = 4x + 2$ **[M1]**

-5 **[A1]**

[Back to Question] *[Total Marks: 5]*

389 **(a)** $\frac{1}{4}$ or 0.25 **[B2]**

Correct intermediate step e.g., 2^{-2} or $\left(\frac{1}{32}\right)^{\frac{2}{5}}$ or $32^{\frac{1}{5}} = 2$ **[M1]**

(b) Writes 125 as 5^3 **[M1]**

Equates indices: $x + 2 = 3x$ **[M1]**

1 **[A1]**

[Back to Question] *[Total Marks: 5]*

390 **(a)** $(\sqrt[3]{125})^2$ or 5^2 or $125^2 = 15625$ and $\sqrt[3]{15625}$ **[M1]**

25 **[A1]**

(b) 3 **[B1]**

x^6 **[B1]**

y^{-8} or $\frac{1}{y^8}$ **[B1]**

Note: Award [B3] for a fully correct final answer of $\frac{3x^6}{y^8}$ or $3x^6y^{-8}$

[Back to Question] *[Total Marks: 5]*

391 **(a)** $\frac{1}{4}$ or 0.25 **[B1]**

(b) 25 or $\frac{1}{25}$ seen **[M1]**

1 **[A1]**

(c) $125 = 5^3$ seen or implied **[B1]**

$2x - 1 = 3x$ **[M1]**

$x = -1$ **[A1]**

[Back to Question] *[Total Marks: 6]*

392 **(a)** $4 \times \frac{3}{2} = 6$ or $x^{-3+\frac{1}{2}}$ **[M1]**

$6x^{-\frac{5}{2}}$ or $\frac{6}{x^{\frac{5}{2}}}$ **[A1]**

(b) $3^{2x+1} = 3^3$ or $2x + 1 = 3$ **[M1]**

1 **[A1]**

(c) $8^{\frac{1}{3}} = 2$ or $(a^6)^{\frac{1}{3}} = a^2$ or $(b^3)^{\frac{1}{3}} = b$ **[M1]**

$\frac{2a^2}{b}$ **[A1]**

[Back to Question] *[Total Marks: 6]*

393 **(a)** Numerator $8x^{15}$ **[B1]**

Denominator $27y^6$ **[B1]**

$\frac{8x^{15}}{27y^6}$ **[A1]**

(b) Writing both sides with a common base of 3, e.g. $3^{2x} = 3^{3(x+1)}$ **[M1]**

$2x = 3x + 3$ **[M1]**

-3 **[A1]**

[Back to Question] *[Total Marks: 6]*

394 **(a)** Takes the reciprocal to deal with the negative index: $\left(\frac{64y^{12}}{27x^9}\right)^{\frac{1}{3}}$ or evaluates cube root of constants correctly **[M1]**

Applies the power of $\frac{1}{3}$ to variables: $y^{12\times\frac{1}{3}}$ and $x^{9\times\frac{1}{3}}$ **[M1]**

$\frac{4y^4}{3x^3}$ **[A1]**

(b) Writes 32 as 2^5 **[B1]**

Equates indices: $3x-1=5x$ **[M1]**

-0.5 or $-\frac{1}{2}$ **[A1]**

[Back to Question] *[Total Marks: 6]*

395 **(a)** 8 **[B1]**

x^9 **[B1]**

y^6 as denominator or y^{-6} **[B1]**

(b) $8=2^3$ seen or implied by $8^x=2^{3x}$ **[M1]**

$x+3=3x$ **[M1]**

1.5 or $\frac{3}{2}$ **[A1]**

[Back to Question] *[Total Marks: 6]*

396 **(a)** $(3^2)^{2x}=(3^3)^{x+1}$ or $3^{4x}=3^{3x+3}$ **[M1]**

$4x=3x+3$ **[M1]**

3 **[A1]**

(b) Inverts fraction to remove negative power: $\left(\frac{81b^{12}}{16a^8}\right)^{\frac{3}{4}}$ **[M1]**

Applies power $\frac{1}{4}$ (fourth root) to all terms: $\frac{3b^3}{2a^2}$ **[M1]**

Cubes the result to get final answer: $\frac{27b^9}{8a^6}$ or $3.375b^9a^{-6}$ **[A1]**

[Back to Question] *[Total Marks: 6]*

397 **(a)** Expressing both sides with base 2: $(2^4)^{2x-1}=(2^3)^{x+2}$ **[M1]**

Equating powers: $4(2x-1)=3(x+2)$ or $8x-4=3x+6$ **[M1]**

2 **[A1]**

(b) Taking the reciprocal to remove negative index: $\left(\frac{16y^8}{81x^{12}}\right)^{\frac{3}{4}}$ **[M1]**

Evaluating the fractional power of the numbers: $16^{\frac{3}{4}} = 8$ and $81^{\frac{3}{4}} = 27$ **[M1]**

$\frac{8y^6}{27x^9}$ **[A1]**

[Back to Question] *[Total Marks: 6]*

398 **(a)** Simplifies the first fraction to $8x^2y^{-5}$ or $\frac{8x^2}{y^5}$ **[M1]**

Multiplies by the reciprocal of the second fraction: $\times\frac{y^4}{4x^3}$ **[M1]**

$\frac{2}{xy}$ or $2x^{-1}y^{-1}$ **[A1]**

(b) Writes 25 as 5^2 and 125 as 5^3 **[M1]**

Applies exponent rules correctly to get a single base on the left: $5^{2x-1+2(x+2)}$ **[M1]**

Equates indices to form a linear equation: $4x + 3 = 3x$ **[M1]**

-3 **[A1]**

[Back to Question] *[Total Marks: 7]*

399 **(a)** $12x^{\frac{3}{2}}$ seen as numerator **[M1]**

$2x^k$ for any k or $cx^{\frac{7}{6}}$ for any c **[M1]**

$2x^{\frac{7}{6}}$ **[A1]**

(b) Converting all terms to base 3: $3^{2(2x-1)} \times 3^{x+1} = 3^{3x}$ **[M1]**

Applying exponent multiplication law: $3^{4x-2+x+1}$ or 3^{5x-1} **[M1]**

Equating indices: $5x - 1 = 3x$ **[M1]**

0.5 or $\frac{1}{2}$ **[A1]**

[Back to Question] *[Total Marks: 7]*

400 **(a)** M1 for simplifying the numerator: $30x^{\frac{1}{2}}$

M1 for correctly handling the division: $6x^{\frac{1}{2}-\frac{3}{2}}$

$6x^{-1}$ or $\frac{6}{x}$ **[A1]**

(b) M1 for reciprocating to eliminate the negative index: $x^{\frac{2}{3}} = \frac{9}{4}$

M1 for isolating x: $x = \left(\frac{9}{4}\right)^{\frac{3}{2}}$

$\frac{27}{8}$ or $3\frac{3}{8}$ **[A1]**

(c) M1 for converting to base 3: $3^{2x} \times 3^{2x-1} = 3^{3(x+1)}$

M1 for applying index laws on the LHS to get 3^{4x-1}

M1 for equating powers correctly: $4x - 1 = 3x + 3$

4 **[A1]**

[Back to Question] *[Total Marks: 10]*

401 **(a)** M1 for $16^{\frac{3}{4}} = 8$ seen

M1 for $x^{12\times\frac{3}{4}} = x^9$ or $y^{8\times\frac{3}{4}} = y^6$ seen

$\frac{8x^9}{y^6}$ or $8x^9y^{-6}$ **[A1]**

(b) M1 for expressing both sides with a common base of 2: $2^{2(2x-1)}$ and $2^{3(x+2)}$

M1 for equating the exponents: $4x - 2 = 3x + 6$

8 **[A1]**

[Back to Question] *[Total Marks: 6]*

402 **(a)** $4x^2$ or y^{-3} or $\frac{1}{y^3}$ seen **[M1]**

$\frac{4x^2}{y^3}$ or $4x^2y^{-3}$ **[A1]**

(b) 3^{2x} or $3^{3(x-1)}$ seen **[B1]**

$3\sqrt{3} = 3^{1.5}$ or $3^{\frac{3}{2}}$ seen **[B1]**

$2x + 3x - 3 = 1.5$ or equivalent equation in x **[M1]**

$x = 0.9$ or $\frac{9}{10}$ **[A1]**

[Back to Question] *[Total Marks: 6]*

403 **(a)** Expressing all terms with a base of 2, e.g. $2^{2x} \times 2^{3(2x-1)} = 2^4$ **[M1]**

Correct linear equation for the exponents: $2x + 6x - 3 = 4$ or $8x = 7$ **[M1]**

$\frac{7}{8}$ or 0.875 **[A1]**

(b) Simplifying the numerator to $24x^1$ (or $24x$) **[M1]**

Correct manipulation of the coefficients $\frac{24}{8} = 3$ or indices $x^{1-(-3)}$ **[M1]**

$3x^4$ **[A1]**

[Back to Question] *[Total Marks: 6]*

404 **(a)** $\left(\frac{27y^6}{8x^9}\right)^{\frac{2}{3}}$ or $\frac{4x^6}{9y^4}$ (inverted or squared/cubed root correctly applied to some terms) **[M1]**

$\left(\frac{3y^2}{2x^3}\right)^2$ **[M1]**

$\frac{9y^4}{4x^6}$ or $2.25y^4x^{-6}$ **[A1]**

(b) $\frac{5^{a-2}\times 5^{2b}}{5^{3(a+1)}} = 5^4$ (expressing all terms as powers of 5) **[M1]**

$a-2+2b-3(a+1)=4$ **[M1]**

$-2a+2b-5=4$ or $-2a+2b=9$ **[A1]**

Accept any pair of values satisfying the equation, e.g. $a=0, b=4.5$ or $a=-4.5, b=0$ or $b=a+4.5$ **[A1]**

[Back to Question] *[Total Marks: 7]*

405 **(a)** $3x^2y^{-6}$ or $\frac{3x^2}{y^6}$ **[B3]**

Coefficient 3 seen **[M1]**

Correct subtraction of indices $x^{\frac{3}{2}-\left(-\frac{1}{2}\right)}$ leading to x^2 OR y^{-2-4} leading to y^{-6} **[M1]**

(b) Expresses all terms as powers of 2: $(2^2)^{2x-1}\times(2^3)^{x+1}=(2^5)^x$ **[M1]**

Correctly adds powers on the LHS: $2(2x-1)+3(x+1)$ or $4x-2+3x+3$ **[M1]**

Forms linear equation equating LHS and RHS powers: $7x+1=5x$ **[M1]**

$-\frac{1}{2}$ or -0.5 **[A1]**

[Back to Question] *[Total Marks: 7]*

406 $3(3x-1)=4(x+5)$ or $9x-3=4x+20$ **[M1]**

$5x=23$ **[M1]**

4.6 or $\frac{23}{5}$ **[A1]**

[Back to Question] *[Total Marks: 3]*

407 Multiplies all terms by 12 (or forms a single fraction): $3(2x-3)+4(x+1)=24$ **[M1]**

Expands brackets correctly: $6x-9+4x+4=24$ **[M1]**

2.9 or $\frac{29}{10}$ **[A1]**

[Back to Question] *[Total Marks: 3]*

408 Valid method to eliminate one variable, e.g. multiplying the first equation by 2 to get $4x-2y=14$ and adding the equations **[M1]**

$x = 4$ **[A1]**

$y = 1$ **[A1]**

[Back to Question] *[Total Marks: 3]*

409 Multiplies one or both equations to equate coefficients of x or y (e.g. $4x - 10y = 44$) **[M1]**

Adds or subtracts equations to eliminate one variable (e.g. $13y = -39$) **[M1]**

$x = 3.5$, $y = -3$ **[A1]**

[Back to Question] *[Total Marks: 3]*

410 **(a)** $4x - 12 = 2x + 10$ **[M1]**

$x = 11$ **[A1]**

(b) $5y = 2x - 1$ **[M1]**

$\frac{5y + 1}{2}$ **[A1]**

[Back to Question] *[Total Marks: 4]*

411 **(a)** $6x - 3 = 5x + 8$ **[M1]**

11 **[A1]**

(b) $y + 7 = 4x$ or $\frac{y}{4} = x - \frac{7}{4}$ **[M1]**

$\frac{y + 7}{4}$ or $\frac{y}{4} + \frac{7}{4}$ **[A1]**

[Back to Question] *[Total Marks: 4]*

412 **(a)** M1 for expanding brackets correctly: $6x - 15 = 4x + 4$

M1 for correctly isolating x terms, e.g., $6x - 4x = 4 + 15$ or $2x = 19$

9.5 or $\frac{19}{2}$ **[A1]**

(b) M1 for multiplying by 3 to get $3V = \pi r^2 h$

$\frac{3V}{\pi r^2}$ **[A1]**

[Back to Question] *[Total Marks: 5]*

413 **(a)** $6x - 3 = 5x + 8$ **[M1]**

11 **[A1]**

(b) $5y = 2x + 3$ **[M1]**

$2x = 5y - 3$ **[M1]**

$\frac{5y-3}{2}$ **[A1]**

[Back to Question] *[Total Marks: 5]*

414 **(a)** Expands brackets correctly to $6x - 3 = 4x + 11$ **[M1]**

7 **[A1]**

(b) Multiplies second equation by 2 to get $2x - 2y = 4$ OR substitutes $x = y + 2$ into first equation **[M1]**

Correctly finds $x = 4$ **[A1]**

Correctly finds $y = 2$ **[A1]**

[Back to Question] *[Total Marks: 5]*

415 **(a)** $5x - 3 = 2x + 12$ **[M1]**

5 **[A1]**

(b) Multiply by 12 to get $4(x + 2) + 3(2x - 1) = 36$ (or equivalent common denominator method) **[M1]**

$4x + 8 + 6x - 3 = 36$ **[M1]**

$\frac{31}{10}$ or 3.1 **[A1]**

[Back to Question] *[Total Marks: 5]*

416 **(a)** Expands bracket to obtain $6x - 15$ **[M1]**

Rearranges terms to isolate x, e.g., $6x - 4x = 7 + 15$ or $2x = 22$ **[M1]**

11 **[A1]**

(b) Multiplies by 5 to obtain $5y = 2x + 3$ **[M1]**

$\frac{5y-3}{2}$ **[A1]**

[Back to Question] *[Total Marks: 5]*

417 **(a)** $8x - 4 = 3x + 11$ **[M1]**

$5x = 15$ **[M1]**

3 **[A1]**

(b) $2y = 3x - 5$ **[M1]**

$2y + 5 = 3x$ **[M1]**

$\frac{2y+5}{3}$ or $\frac{2y}{3} + \frac{5}{3}$ **[A1]**

[Back to Question] *[Total Marks: 6]*

418 **(a)** $5x - 10 = 3x + 14$ **[M1]**

$2x = 24$ **[M1]**

12 **[A1]**

(b) $2A = h(a + b)$ or $\frac{A}{h} = \frac{a+b}{2}$ **[M1]**

$\frac{2A}{h} = a + b$ or $2A = ha + hb$ **[M1]**

$\frac{2A}{h} - a$ or $\frac{2A - ah}{h}$ **[A1]**

[Back to Question] *[Total Marks: 6]*

419 **(a)** $6x - 15 = 4x + 7$ **[M1]**

$2x = 22$ **[M1]**

11 **[A1]**

(b) Equates coefficients, e.g. multiplies first equation by 2 and second equation by 3 to get $4x + 6y = 26$ and $15x - 6y = 12$ **[M1]**

Adds equations to eliminate y, e.g. $19x = 38$, so $x = 2$ **[A1]**

$y = 3$ **[A1]**

[Back to Question] *[Total Marks: 6]*

420 **(a)** Expands brackets: $4x - 12 = 2x + 7$ **[M1]**

Rearranges to collect terms: $2x = 19$ **[M1]**

9.5 or $\frac{19}{2}$ **[A1]**

(b) Multiplies first equation by 2 to equate coefficients of y: $10x - 4y = 32$ (or equivalent valid method) **[M1]**

Adds equations to eliminate y: $13x = 39 \implies x = 3$ **[A1]**

Substitutes $x = 3$ into either equation to find y: $15 - 2y = 16 \implies y = -0.5$ **[A1]**

[Back to Question] *[Total Marks: 6]*

421 $y(x + 4) = 3x - 2$ **[M1]**

$xy - 3x = -4y - 2$ or $3x - xy = 4y + 2$ **[M1]**

$x(y - 3) = -4y - 2$ or $x(3 - y) = 4y + 2$ **[M1]**

$\frac{4y+2}{3-y}$ or $\frac{-4y-2}{y-3}$ **[A1]**

[Back to Question] *[Total Marks: 4]*

422 $y(x - 5) = 2x + 3$ or $xy - 5y = 2x + 3$ **[M1]**

$xy - 2x = 5y + 3$ or $2x - xy = -5y - 3$ **[M1]**

$x(y-2) = 5y+3$ or $x(2-y) = -5y-3$ **[M1]**

$\frac{5y+3}{y-2}$ or $\frac{-5y-3}{2-y}$ **[A1]**

[Back to Question] *[Total Marks: 4]*

423 Multiplies by the denominator: $y(2x+1) = 3x-5$ **[M1]**

Expands the bracket: $2xy + y = 3x - 5$ **[M1]**

Rearranges to collect all terms involving x on one side: $y+5 = 3x-2xy$ (or $2xy - 3x = -y-5$) **[M1]**

$\frac{y+5}{3-2y}$ or $\frac{-y-5}{2y-3}$ **[A1]**

[Back to Question] *[Total Marks: 4]*

424 (a) Multiplies equations to equate coefficients of x or y, e.g., $9x+6y=36$ and $10x-6y=2$ **[M1]**

Adds or subtracts equations to eliminate a variable **[M1]**

$x=2$ and $y=3$ **[A1]**

(b) $(2x \pm a)(x \pm b) = 0$ where $ab = 12$ **[M1]**

$(2x+3)(x-4) = 0$ **[M1]**

$x = 4$ and $x = -1.5$ (or $-\frac{3}{2}$) **[A1]**

[Back to Question] *[Total Marks: 6]*

425 (a) Correct method to equate coefficients (e.g., $10x+15y=65$ and $10x-4y=8$, leading to $19y=57$) or correctly substitute **[M1]**

$y=3$ **[A1]**

$x=2$ **[A1]**

(b) $y(x+2) = 3x-1$ **[M1]**

$xy + 2y = 3x - 1$ **[M1]**

$xy - 3x = -1 - 2y$ or $2y+1 = 3x-xy$ **[M1]**

$\frac{-1-2y}{y-3}$ or $\frac{2y+1}{3-y}$ **[A1]**

[Back to Question] *[Total Marks: 7]*

426 (a) $y(x+4) = 3x-2$ **[M1]**

$xy + 4y = 3x - 2$ **[M1]**

$xy - 3x = -4y - 2$ or $x(y-3) = -(4y+2)$ **[M1]**

$\frac{-4y-2}{y-3}$ or $\frac{4y+2}{3-y}$ **[A1]**

(b) $5(x+1) = 2(x-2)$ **[M1]**

$5x + 5 = 2x - 4$ **[M1]**

-3 **[A1]**

[Back to Question] *[Total Marks: 7]*

427 **(a)** Multiplies equations to equate coefficients, e.g., $9x + 6y = 36$ and $10x - 6y = 2$ **[M1]**

Adds or subtracts equations to eliminate one variable, e.g., $19x = 38$ **[M1]**

$x = 2$ **[A1]**

$y = 3$ **[A1]**

(b) Finds correct product and sum components, e.g., $-10x$ and $3x$ **[M1]**

Correctly factorises to $(2x+3)(x-5) = 0$ **[M1]**

$x = -1.5, x = 5$ **[A1]**

[Back to Question] *[Total Marks: 7]*

428 **(a)** Multiply the second equation by 2 to get $8x + 2y = 36$ (or correct substitution method) **[M1]**

$x = 5$ **[A1]**

$y = -2$ **[A1]**

(b) $T(p-2) = 3p + 4$ **[M1]**

$Tp - 2T = 3p + 4$ **[M1]**

$Tp - 3p = 2T + 4$ or $p(T-3) = 2T + 4$ **[M1]**

$\frac{2T+4}{T-3}$ or $\frac{-2T-4}{3-T}$ **[A1]**

[Back to Question] *[Total Marks: 7]*

429 **(a)** Multiplies the first equation by 2 to give $4x - 2y = 18$ (or rearranges to $y = 2x - 9$) **[M1]**

Adds the equations to eliminate y, yielding $7x = 28$ (or correctly substitutes to form an equation in one variable) **[M1]**

$x = 4$ **[A1]**

$y = -1$ **[A1]**

(b) Multiplies throughout by a common denominator (e.g., 12) to obtain $4(2x) - 3(x-4) = 30$ **[M1]**

Expands brackets correctly: $8x - 3x + 12 = 30$ **[M1]**

$5x = 18$ **[M1]**

3.6 or $\frac{18}{5}$ **[A1]**

[Back to Question] *[Total Marks: 8]*

430 **(a)** Multiply second equation by 2: $4x - 10y = 18$ **[M1]**

Subtract equations to eliminate x: $13y = -13$ **[M1]**

$y = -1$ **[A1]**

$x = 2$ **[A1]**

(b) $(3x + 2)(x - 4) = 0$ **[M2]**

If M0, M1 for $(3x \pm a)(x \pm b)$ where $ab = 8$ or $a \pm 3b = -10$

$x = -\frac{2}{3}$ **[A1]**

$x = 4$ **[A1]**

[Back to Question] *[Total Marks: 8]*

431 Forms common denominator or multiplies through: $2(x - 2) + 1(x - 1) = (x - 1)(x - 2)$ **[M1]**

Expands brackets correctly: $2x - 4 + x - 1 = x^2 - 3x + 2$ **[M1]**

Rearranges to a 3-term quadratic: $x^2 - 6x + 7 = 0$ **[A1]**

Applies quadratic formula: $\frac{-(-6) \pm \sqrt{(-6)^2 - 4(1)(7)}}{2(1)}$ or completes the square:

$(x - 3)^2 - 9 + 7 = 0$ **[M1]**

Correct unsimplified surd form: $\frac{6 \pm \sqrt{8}}{2}$ or $(x - 3)^2 = 2$ **[M1]**

$3 \pm \sqrt{2}$ **[A1]**

[Back to Question] *[Total Marks: 6]*

432 **(a)** B1 for $x + y = 150$

B1 for $12x + 8y = 1480$

M1 for a valid method to solve the equations (e.g., multiplying the first equation by 8 to get $8x + 8y = 1200$ and subtracting)

$x = 70$ and $y = 80$ **[A1]**

(b) M1 for isolating the square root: $\frac{T}{2\pi} = \sqrt{\frac{m}{k}}$

M1 for squaring both sides: $\frac{T^2}{4\pi^2} = \frac{m}{k}$

$\frac{kT^2}{4\pi^2}$ **[A1]**

[Back to Question] *[Total Marks: 7]*

433 **(a)** $p(x + 4) = 3x - 1$ **[M1]**

$px + 4p = 3x - 1$ **[M1]**

$3x - px = 4p + 1$ (or equivalent step isolating x terms on one side) **[M1]**

$\frac{4p+1}{3-p}$ (or $\frac{-4p-1}{p-3}$) **[A1]**

(b) $(2x \pm a)(x \pm b) = 0$ where $ab = -12$ or $2b \pm a = -5$ **[M1]**

$(2x+3)(x-4) = 0$ **[M1]**

$x = -1.5$, $x = 4$ **[A1]**

[Back to Question] *[Total Marks: 7]*

434 **(a)** $2(3x+2+x-1) = 42$ or $3x+2+x-1 = 21$ **[M1]**

$4x+1 = 21$ **[M1]**

5 **[A1]**

(b) $p^2 = \frac{x-3}{x}$ **[M1]**

$p^2x = x-3$ **[M1]**

$p^2x - x = -3$ or $x - p^2x = 3$ or $x(p^2-1) = -3$ **[M1]**

$\frac{3}{1-p^2}$ or $\frac{-3}{p^2-1}$ **[A1]**

[Back to Question] *[Total Marks: 7]*

435 **(a)** Multiplies by denominator: $y(x-3) = 2x+5$ **[M1]**

Expands and collects x terms on one side: $xy - 2x = 3y + 5$ **[M1]**

$\frac{3y+5}{y-2}$ or $\frac{-3y-5}{2-y}$ **[A1]**

(b) Multiplies throughout by common denominator $3(x-2)$ to reach $(x+1)(x-2) - 6 = 3(x-2)$ **[M1]**

Expands brackets correctly: $x^2 - 2x + x - 2 - 6 = 3x - 6$ **[M1]**

Rearranges to form a correct 3-term quadratic equation: $x^2 - 4x - 2 = 0$ **[A1]**

Uses the quadratic formula $\frac{-(-4) \pm \sqrt{(-4)^2 - 4(1)(-2)}}{2(1)}$ **[M1]**

4.45 **[A1]**

-0.45 **[A1]**

[Back to Question] *[Total Marks: 9]*

436 **(a)** Sets up equation for area: $(2x+3)(x-1) = 45$ **[M1]**

Expands and simplifies to given result: $2x^2 - 2x + 3x - 3 = 45 \implies 2x^2 + x - 48 = 0$ **[A1]**

(b) Uses quadratic formula with correct values: $\frac{-1 \pm \sqrt{1^2 - 4(2)(-48)}}{2(2)}$ **[M1]**

Correct discriminant: $\frac{-1 \pm \sqrt{385}}{4}$ **[M1]**

4.655... and $-5.155...$ **[A1]**

4.66 and -5.16 **[A1]**

(c) Multiplies by denominator: $y(x-3) = ax+5$ **[M1]**

Expands brackets: $xy - 3y = ax + 5$ **[M1]**

Collects terms in x on one side: $xy - ax = 3y + 5$ **[M1]**

Factorises x and divides: $\dfrac{3y+5}{y-a}$ (or equivalent) **[A1]**

[Back to Question] *[Total Marks: 10]*

437 $\dfrac{3(x-1)+1(x+2)}{(x+2)(x-1)} = 2$ or $3(x-1)+1(x+2) = 2(x+2)(x-1)$ **[M1]**

$3x - 3 + x + 2$ or $4x - 1$ **[M1]**

$2x^2 - 2x - 3 = 0$ or $-2x^2 + 2x + 3 = 0$ **[A2]**

$\dfrac{-(-2) \pm \sqrt{(-2)^2 - 4(2)(-3)}}{2(2)}$ or GDC equivalent to solve the 3-term quadratic **[M1]**

1.82 and -0.82 **[A1]**

[Back to Question] *[Total Marks: 6]*

438 Multiplies by the common denominator to get $3(2x+1)+4(x-2) = 2(x-2)(2x+1)$ [M2]
(allow M1 for $\dfrac{3(2x+1)+4(x-2)}{(x-2)(2x+1)} = 2$)

Expands brackets to get $10x - 5 = 2(2x^2 - 3x - 2)$ or $10x - 5 = 4x^2 - 6x - 4$ **[M1]**

Rearranges to form a correct 3-term quadratic equation equal to zero: $4x^2 - 16x + 1 = 0$ **[A1]**

Substitutes correctly into the quadratic formula: $x = \dfrac{-(-16) \pm \sqrt{(-16)^2 - 4(4)(1)}}{2(4)}$ **[M1]**

$x = 3.94$ and $x = 0.06$ **[A1]**

[Back to Question] *[Total Marks: 6]*

439 Multiplies by the common denominator to get $5(2x+1) - 3(x-2) = 2(x-2)(2x+1)$ **[M2]**

If not M2, M1 for a correct single fraction $\dfrac{5(2x+1)-3(x-2)}{(x-2)(2x+1)}$ seen

$4x^2 - 13x - 15 = 0$ **[A2]**

If not A2, A1 for correctly expanding sides to $7x + 11$ or $4x^2 - 6x - 4$

$x = \dfrac{-(-13) \pm \sqrt{(-13)^2 - 4(4)(-15)}}{2(4)}$ **[M1]**

4.15 **[A1]**

-0.90 **[A1]**

[Back to Question] *[Total Marks: 7]*

440 **(a)** $\dfrac{3(x+3)-2(x-1)}{(x-1)(x+3)} = 1$ **[M1]**

$3x + 9 - 2x + 2 = (x-1)(x+3)$ **[M1]**

$x + 11 = x^2 + 2x - 3$ **[M1]**

$x^2 + x - 14 = 0$ established with no errors seen **[A1]**

(b) $\dfrac{-1 \pm \sqrt{1^2 - 4(1)(-14)}}{2(1)}$ **[M1]**

$\dfrac{-1 \pm \sqrt{57}}{2}$ **[M1]**

$3.27, -4.27$ **[A1]**

[Back to Question] *[Total Marks: 7]*

441 **(a)** **(i)** Forms a correct initial equation involving time: $\dfrac{120}{x} - \dfrac{120}{x+10} = \dfrac{36}{60}$ **[M1]**

Simplifies the time difference to $\dfrac{3}{5}$ and multiplies by a common denominator: $600(x+10) - 600x = 3x(x+10)$ **[M1]**

$6000 = 3x^2 + 30x$ leading correctly to $x^2 + 10x - 2000 = 0$ with no errors **[A1]**

(ii) $(x+50)(x-40) = 0$ **[M1]**

$x = 40, x = -50$ **[A1]**

(b) Uses the quadratic formula with correct substitution: $x = \dfrac{-(-7) \pm \sqrt{(-7)^2 - 4(3)(-5)}}{2(3)}$ **[M1]**

Simplifies to $\dfrac{7 \pm \sqrt{109}}{6}$ **[M1]**

$x = 2.91$ or $x = -0.57$ **[A1]**

[Back to Question] *[Total Marks: 8]*

442 **(a)** Constructs initial time equation: $\dfrac{120}{x} + \dfrac{150}{x+10} = 4$ **[M1]**

Multiplies by common denominator $x(x+10)$ to reach $120(x+10) + 150x = 4x(x+10)$ **[M1]**

Expands brackets correctly: $120x + 1200 + 150x = 4x^2 + 40x$ **[M1]**

Rearranges to $4x^2 - 230x - 1200 = 0$ and divides by 2 to yield $2x^2 - 115x - 600 = 0$ with no errors seen **[A1]**

(b) Substitutes values into the quadratic formula $\dfrac{115 \pm \sqrt{(-115)^2 - 4(2)(-600)}}{4}$ **[M1]**

62.3 **[A1]**

(c) -1.84 **[B1]**

2.36 **[B1]**

[Back to Question] *[Total Marks: 8]*

443 **(a)** $\frac{40}{x}$ **[B1]**

(b) Converts 45 minutes to hours: $\frac{45}{60}$ or $\frac{3}{4}$ (or 0.75) **[B1]**

Forms a correct fractional equation: $\frac{40}{x} - \frac{40}{x+20} = \frac{3}{4}$ **[M1]**

Finds a common denominator or multiplies through to clear fractions:
$40(x+20) - 40x = \frac{3}{4}x(x+20)$ **[M1]**

Expands correctly and simplifies to the given quadratic with no errors: $800 = \frac{3}{4}(x^2 + 20x)$

$3200 = 3x^2 + 60x$

$3x^2 + 60x - 3200 = 0$ [A1]

(c) Correct substitution into the quadratic formula: $x = \frac{-60 \pm \sqrt{60^2 - 4(3)(-3200)}}{2(3)}$ **[M1]**

Correct simplification of the discriminant: $\frac{-60 \pm \sqrt{42000}}{6}$ **[M1]**

24.16 and -44.16 (both correct) **[A1]**

[Back to Question] *[Total Marks: 8]*

444 **(a)** M1 for multiplying by the common denominator to obtain
$5(x-3) + 2(x+2) = (x+2)(x-3)$

M1 for expanding the brackets correctly: $5x - 15 + 2x + 4 = x^2 - x - 6$

M1 for rearranging into a correct three-term quadratic equation, e.g., $x^2 - 8x + 5 = 0$

M1 for correct use of the quadratic formula, e.g., $\frac{-(-8) \pm \sqrt{(-8)^2 - 4(1)(5)}}{2(1)}$

7.32 and 0.68 **[A1]**

(b) M1 for multiplying by the denominator: $y(5x+1) = 2x - 3$

M1 for expanding the bracket and collecting x terms on one side: $5xy - 2x = -y - 3$ or $2x - 5xy = y + 3$

M1 for factorising x: $x(5y-2) = -y - 3$ or $x(2-5y) = y + 3$

$\frac{-y-3}{5y-2}$ or $\frac{y+3}{2-5y}$ **[A1]**

[Back to Question] *[Total Marks: 9]*

445 **(a)** $x(x-6) = 3(x+2)$ **[M1]**

$x^2 - 6x = 3x + 6$ **[M1]**

$x^2 - 9x - 6 = 0$ **[A1]**

Uses the quadratic formula, e.g., $x = \frac{-(-9) \pm \sqrt{(-9)^2 - 4(1)(-6)}}{2(1)}$ **[M1]**

$\frac{9 \pm \sqrt{105}}{2}$ **[A1]**

(b) $y(3x+1) = 2x - 5$ **[M1]**

$3xy + y = 2x - 5$ **[M1]**

Isolates terms containing x on one side, e.g., $3xy - 2x = -y - 5$ or $2x - 3xy = y + 5$ **[M1]**

$\frac{-y-5}{3y-2}$ or $\frac{y+5}{2-3y}$ **[A1]**

[Back to Question] *[Total Marks: 9]*

446 **(a)** $3(2x+1) + 2(x-1) = 2(x-1)(2x+1)$ **[M1]**

$6x + 3 + 2x - 2 = 4x^2 - 2x - 2$ **[M1]**

$4x^2 - 10x - 3 = 0$ **[A1]**

$\frac{10 \pm \sqrt{(-10)^2 - 4(4)(-3)}}{2(4)}$ **[M1]**

$2.77, -0.27$ **[A1]**

(b) $x^2 + (x+2)^2 = 340$ (or equivalent setup for consecutive even numbers) **[M1]**

$2x^2 + 4x - 336 = 0$ or $x^2 + 2x - 168 = 0$ **[A1]**

$(x+14)(x-12) = 0$ or correct application of quadratic formula **[M1]**

12 and 14 (both correct) **[A1]**

[Back to Question] *[Total Marks: 9]*

447 **(a)** $\frac{100}{x} - \frac{100}{x+15} = \frac{1}{2}$ (or equivalent e.g., $\frac{100}{x} = \frac{100}{x+15} + 0.5$) **[M1]**

$200(x+15) - 200x = x(x+15)$ **[M1]**

$200x + 3000 - 200x = x^2 + 15x$ leading to $x^2 + 15x - 3000 = 0$ **[A1]**

(b) $\frac{-15 \pm \sqrt{15^2 - 4(1)(-3000)}}{2(1)}$ **[M2]**

Allow [M1] for correctly substituting values into the quadratic formula or evaluating the discriminant $\sqrt{12225}$

47.78 **[A1]**

-62.78 **[A1]**

(c) Speed of car $= 47.78 + 15 = 62.78$ km/h **[M1]**

Time $= \frac{100}{62.78} = 1.592...$ hours **[M1]**

1 hour 36 minutes **[A1]**

[Back to Question] *[Total Marks: 10]*

448 **(a)** Multiplies by the denominator to obtain $y(x+4) = 3x - 2$ **[M1]**

Expands the bracket to obtain $xy + 4y = 3x - 2$ **[M1]**

Collects terms in x on one side, e.g., $xy - 3x = -4y - 2$ or $3x - xy = 4y + 2$ **[M1]**

Factorises x out and divides: $\dfrac{-4y-2}{y-3}$ or $\dfrac{4y+2}{3-y}$ **[A1]**

(b) Writes with common denominator or multiplies through to eliminate fractions:
$5(2x+1)-3(x-2)=2(x-2)(2x+1)$ **[M1]**

Correctly expands LHS: $10x+5-3x+6$ **[M1]**

Correctly expands RHS: $2(2x^2-3x-2)$ or $4x^2-6x-4$ **[M1]**

Rearranges to form a correct 3-term quadratic equation, e.g., $4x^2-13x-15=0$ **[A1]**

Correctly substitutes into the quadratic formula: $\dfrac{-(-13)\pm\sqrt{(-13)^2-4(4)(-15)}}{2(4)}$ **[M1]**

$x=4.15$ and $x=-0.90$ **[A1]**

[Back to Question] *[Total Marks: 10]*

449 **(a)** $y(x+2)=3x-1$ **[M1]**

$yx+2y=3x-1$ **[M1]**

$2y+1=3x-yx$ or $yx-3x=-2y-1$ **[M1]**

$x=\dfrac{2y+1}{3-y}$ or $x=\dfrac{-2y-1}{y-3}$ **[A1]**

(b) Multiply throughout by $(x-1)(x+2)$: $2(x+2)+3(x-1)=(x-1)(x+2)$ **[M1]**

Expand terms: $2x+4+3x-3=x^2+x-2$ **[M1]**

Collect to form a quadratic equation: $x^2-4x-3=0$ **[A1]**

Use the quadratic formula: $x=\dfrac{-(-4)\pm\sqrt{(-4)^2-4(1)(-3)}}{2(1)}$ **[M1]**

$x=\dfrac{4\pm\sqrt{28}}{2}$ **[M1]**

$x=4.65$ and $x=-0.65$ **[A1]**

[Back to Question] *[Total Marks: 10]*

450 **(a)** $\dfrac{150}{x}$ **[B1]**

(b) $\dfrac{150}{x-15}-\dfrac{150}{x}=0.5$ (or $\dfrac{30}{60}$ or $\dfrac{1}{2}$) **[M1]**

$150x-150(x-15)=0.5x(x-15)$ **[M1]**

$150x-150x+2250=0.5x^2-7.5x$ **[M1]**

$4500=x^2-15x \implies x^2-15x-4500=0$ with no errors seen **[A1]**

(c) $(x-75)(x+60)=0$ or $\dfrac{-(-15)\pm\sqrt{(-15)^2-4(1)(-4500)}}{2(1)}$ **[M2]**

(or $M1$ for correct factors $(x+a)(x+b)$ where $ab=-4500$ or $a+b=-15$, or for correct substitution into quadratic formula)

75 (rejects negative solution -60) **[A1]**

(d) $P(q-1)=2q+3$ **[M1]**

$Pq - P = 2q + 3$ **[M1]**

$Pq - 2q = P + 3$ **[M1]**

$\frac{P+3}{P-2}$ or $\frac{-P-3}{2-P}$ **[A1]**

[Back to Question] *[Total Marks: 12]*

451 Rearranges to collect x terms on one side and constants on the other, e.g., $-8 < 4x$ or $-4x < 8$ **[M1]**

$x > -2$ or $-2 < x$ **[A1]**

[Back to Question] *[Total Marks: 2]*

452 (a) $3x - 6 > 5x + 4$ **[M1]**

$x < -5$ **[A1]**

(b) -6 **[B1]**

[Back to Question] *[Total Marks: 3]*

453 Expands brackets correctly: $12x - 4 - 5x \geq 2x + 16$ **[M1]**

Rearranges to collect x terms: $7x - 2x \geq 16 + 4$ or $5x \geq 20$ **[M1]**

$x \geq 4$ **[A1]**

[Back to Question] *[Total Marks: 3]*

454 (a) $3x - 12 < 5x - 2$

[M1]

$-10 < 2x \implies x > -5$ (or $-5 < x$)

[A1]

(b) -4

[B1]

[Back to Question] *[Total Marks: 3]*

455 (a) $6x - 3 > 4x + 7$ **[M1]**

$2x > 10$ leading to $x > 5$ **[A1]**

(b) Open circle placed at 5 with an arrow pointing to the right **[B1]**

(c) 6 **[B1]**

[Back to Question] *[Total Marks: 4]*

456 **(a)** $-6 < 3x \leq 12$ **[M1]**

$-2 < x \leq 4$ **[A1]**

(b) $-1, 0, 1, 2, 3, 4$ **[B2]**

(or [B1] for 5 correct with no incorrect, or all 6 correct with one incorrect)

[Back to Question] *[Total Marks: 4]*

457 **(a)** $-4 < 2x \leq 8$ **[M1]**

$-2 < x \leq 4$ **[A1]**

(b) Open circle at -2 and closed circle at 4 **[B1]**

Thick line joining the two circles **[B1]**

[Back to Question] *[Total Marks: 4]*

458 **(a)** $4x \leq 20$ **[M1]**

$x \leq 5$ **[A1]**

(b) $3x + 6 < 20$ or $x + 2 < 6.66\ldots$ **[M1]**

4 **[A1]**

[Back to Question] *[Total Marks: 4]*

459 **(a)** $2x < 12$ **[M1]**

$x < 6$ **[A1]**

(b) Recognises open circle means strict inequality ($>$ or $<$) and arrow to right means greater than **[M1]**

$x > -2$ **[A1]**

[Back to Question] *[Total Marks: 4]*

460 **(a)** M1 for a correct first step, e.g., adding 1 to all parts to get $-4 < 2x \leq 8$

M1 for a correct second step, e.g., dividing all parts by 2

$-2 < x \leq 4$ **[A1]**

(b) M1 for an open circle at -2 and a closed solid circle at 4

A1 for a continuous thick line drawn between the two circles

[Back to Question] *[Total Marks: 5]*

461 **(a)** Expands brackets correctly: $3x - 6 \leq 5x + 8$ **[M1]**

Rearranges to isolate x, e.g. $-14 \leq 2x$ or $-2x \leq 14$ **[M1]**

$x \geq -7$ **[A1]**

(b) Places a closed, solid circle at -7 **[B1]**

Draws a line with an arrow pointing to the right from the circle **[B1]**

[Back to Question] *[Total Marks: 5]*

462 **(a)** $-9 > 2x$ or $-2x > 9$ **[M1]**

$x < -4.5$ **[A1]**

(b) $-4 < 2n \leq 4$ **[M1]**

$-2 < n \leq 2$ **[M1]**

$-1, 0, 1, 2$ **[A1]**

(c) $x > -1$ (accept any valid variable) **[B1]**

[Back to Question] *[Total Marks: 6]*

463 **(a)** $-15 \leq 3x$ or $-3x \leq 15$ **[M1]**

$x \geq -5$ or $-5 \leq x$ **[A1]**

(b) $-7 < 2x \leq 8$ **[M1]**

$-3.5 < x \leq 4$ **[A1]**

(c) Largest $= 4$ **[B1]**

Smallest $= -3$ **[B1]**

[Back to Question] *[Total Marks: 6]*

464 **(a)** $-3x < 9$ or $2 - 11 < 3x$ **[M1]**

$x > -3$ **[A1]**

(b) $8x - 4 \leq 5x + 17$ **[M1]**

$3x \leq 21$ **[M1]**

$x \leq 7$ **[A1]**

(c) $-3 < n \leq 7$ **[M1]**

$-2, -1, 0, 1, 2, 3, 4, 5, 6, 7$ **[A1]**

[Back to Question] *[Total Marks: 7]*

465 **(a)** Subtracts $3x$ and 3 to reach $\pm 4x \leq \pm 8$ or similar **[M1]**

$x \geq -2$ **[A1]**

(b) Solid circle (or equivalent clear inclusive mark) placed at -2 **[B1]**

Thick line or arrow pointing to the right starting from -2 **[B1]**

(c) Adds 5 to all parts to get $3 < 3x \leq 9$ **[M1]**

Divides by 3 to get $1 < x \leq 3$ **[A1]**

$2, 3$ **[A1]**

[Back to Question] *[Total Marks: 7]*

466 $y \geq 1$ **[B1]**

$y \leq x$ **[B1]**

$x + y \leq 6$ **[B1]**

[Back to Question] *[Total Marks: 3]*

467 $y \leq 5x - 4$ or equivalent **[B1]**

$2y \geq x + 1$ or equivalent **[B1]**

$x + y \leq 8$ or equivalent **[B1]**

[Back to Question] *[Total Marks: 3]*

468 $x \geq 2$ **[B1]**

$y > x$ **[B1]**

$x + y \leq 8$ (or $y \leq -x + 8$) **[B1]**

[Back to Question] *[Total Marks: 3]*

469 $x \geq 1$

[B1]

$y \geq 2$

[B1]

$x + y < 6$ (or $y < -x + 6$)

[B1]

[Back to Question] *[Total Marks: 3]*

470 **(a)** $-6 < 3x \leq 12$ **[M1]**

$-2 < x \leq 4$ **[A1]**

(b) Open circle at -2 **[B1]**

Closed circle at 4 and a line joining the two circles **[B1]**

[Back to Question] *[Total Marks: 4]*

471 **(a)** $-6 < 2x \leq 6$ (Allow M1 for one correct side e.g. $-6 < 2x$ or $2x \leq 6$) **[M2]**

$-3 < x \leq 3$ **[A1]**

(b) Open circle at -3 and closed circle at 3 **[B1]**

A single continuous line drawn connecting the two circles **[B1]**

[Back to Question] *[Total Marks: 5]*

472 **(a)** **(a)** Adds 1 to all three parts: $-4 \leq 2x < 7$ **[M1]**

Divides all three parts by 2 **[M1]**

$-2 \leq x < 3.5$ or $-2 \leq x < \frac{7}{2}$ **[A1]**

(b) Award [B1] for at least three correct integers and no incorrect integers, or all correct integers with only one incorrect integer

$-2, -1, 0, 1, 2, 3$ **[B2]**

[Back to Question] *[Total Marks: 5]*

473 **(a)** $x < 4$ **[B1]**

$y \geq -x + 3$ **[B1]**

$y \leq \frac{1}{2}x + 3$ (or equivalent, e.g. $2y \leq x + 6$) **[B1]**

(b) Substitutes $x = 3$ into the inequalities: $b \geq -3 + 3 = 0$ and $b \leq 0.5(3) + 3 = 4.5$ **[M1]**

Deduces the range $0 \leq b \leq 4.5$ **[A1]**

$0, 1, 2, 3, 4$ **[A1]**

[Back to Question] *[Total Marks: 6]*

474 $x \geq 1$ **[B1]**

$y \geq 2$ **[B1]**

$x + y < 6$ or $y < -x + 6$ **[B1]**

[Back to Question] *[Total Marks: 3]*

475 **(a)** $y \geq 1$ **[B1]**

$y \leq x$ **[B1]**

$x + y \leq 6$ or $y \leq -x + 6$ **[B1]**

(b) Tests a valid integer coordinate close to the boundary, e.g., $(5, 1)$, $(4, 2)$, or $(3, 3)$ **[M1]**

17 (from $x = 5, y = 1$) **[A1]**

[Back to Question] *[Total Marks: 5]*

476 Identifies horizontal line $y = 2$ **[M1]**

States $y \geq 2$ (accept $y > 2$) **[A1]**

Identifies dashed line passing through $(0, 6)$ and $(4, 2)$ **[M1]**

Finds equation $y = -x + 6$ and states $y < -x + 6$ **[A1]**

Identifies solid line passing through $(0, 2)$ and $(1, 5)$ **[M1]**

Finds equation $y = 3x + 2$ and states $y \leq 3x + 2$ **[A1]**

[Back to Question] *[Total Marks: 6]*

477 **(a)** Multiplies by 15 to clear denominators: $3(3x - 2) - 5(2x - 5) \leq 30$ **[M1]**

$9x - 6 - 10x + 25 \leq 30$ **[M1]**

$-x + 19 \leq 30$ **[M1]**

$x \geq -11$ **[A1]**

(b) Identifies that y can only take integer values of 2 or 3 (since $2y > x + y > 2 \implies y > 1$) **[M1]**

Finds correct points for one of the valid y values: e.g. $(1, 2)$ for $y = 2$ OR $(0, 3), (1, 3), (2, 3)$ for $y = 3$ **[B1]**

States all four correct points: $(1, 2), (0, 3), (1, 3), (2, 3)$ with no extras **[A1]**

[Back to Question] *[Total Marks: 7]*

478 $y \geq 2$ **[B1]**

$y \leq 2x$ **[B1]**

$x + y \leq 6$ **[B1]**

[Back to Question] *[Total Marks: 3]*

479 $x \geq 1$ **[B1]**

$y \leq 2x$ **[B1]**

$x + y \leq 6$ **[B1]**

(Note: Accept strict inequalities $<, >$)

[Back to Question] *[Total Marks: 3]*

480 **(a)** Adding 1 to all parts: $-6 \leq 3x < 12$ **[M2, M1 for one side correct]**

Dividing by 3: $-2 \leq x < 4$ **[A1]**

(b) Identifies at least 3 correct integers or includes 4 but excludes -2 **[M1]**

$-2, -1, 0, 1, 2, 3$ **[A1]**

[Back to Question] *[Total Marks: 5]*

481 **(a)** $y \geq 1$ **[B1]**

$y \leq x + 1$ **[B1]**

$y < 5 - x$ (or $x + y < 5$) **[B1]**

All three inequality signs correct (with strict inequality for the dashed line) **[B1]**

(b) Identifies $(1, 2)$ as a valid integer coordinate point inside R **[M1]**

$(1, 2)$ **[A1]**

[Back to Question] *[Total Marks: 6]*

482 **(a)** Subtracts 1 from all parts: $-6 < 2x \leq 10$ **[M1]**

Divides by 2 across all parts **[M1]**

$-3 < x \leq 5$ **[A1]**

(b) Identifies at least two correct integers from the set **[M1]**

$-2, -1, 0$ **[A1]**

(c) Finds the correct roots of the quadratic $2x^2 - 7x + 3 = 0$ as 0.5 and 3 **[B1]**

$x < 0.5$ or $x > 3$ **[B1]**

[Back to Question] *[Total Marks: 7]*

483 **(a)** $y \geq 1$ **[B1]**

$y \leq 2x - 2$ **[B2]**

(or $M1$ for finding the boundary equation $y = 2x - 2$)

$x + y \leq 7$ (or $y \leq -x + 7$) **[B2]**

(or $M1$ for finding the boundary equation $x + y = 7$)

(b) Identifies a valid integer coordinate inside or on the boundary of R that maximizes the expression, e.g., $(6, 1)$, $(5, 2)$, or $(4, 3)$ **[M1]**

13 (from $2(6) + 1 = 13$) **[A1]**

[Back to Question] *[Total Marks: 7]*

484 **(a)** $-4 < 2x \leq 8$ **[M1]**

$-2 < x \leq 4$ **[A1]**

(b) $-1, 0, 1, 2, 3, 4$ **[B2]**

1 omission or 1 extra integer **[B1]**

(c) **(i)** $2(3x + 2) + 2x < 40$ or $(3x + 2) + x < 20$ **[M1]**

$8x + 4 < 40$

8x ¡ 36

x ¡ $4.5 with outer errors [A1]$

(ii) Identifies maximum integer $x = 4$ **[M1]**

Length = 14, Width = 4, Area = 56 **[A1]**

[Back to Question] *[Total Marks: 8]*

485 **(a)** $y \geq 2$ **[B1]**

$y \leq x$ **[B1]**

$x + y < 8$ **[B1]**

(b) Identifies $(4, 3)$ or $(5, 2)$ as integer coordinates within the region R. Note that $(4, 4)$ is excluded due to the strict inequality $x + y < 8$. **[M1]**

Evaluates $2x + 3y$ and correctly identifies $(4, 3)$ as yielding the maximum value (17). **[A1]**

[Back to Question] *[Total Marks: 5]*

486 **(a)** $x \geq 15$ **[B1]**

$x + y \leq 40$ **[B1]**

$y \geq \frac{1}{2}x$ or $2y \geq x$ **[B1]**

(b) Identifies the objective function for profit: $P = 400x + 150y$ **[M1]**

Tests the valid integer vertex $(26, 14)$ or evaluates the profit equation showing profit is maximised at the highest possible integer value for x **[M1]**

12500 **[A1]**

[Back to Question] *[Total Marks: 6]*

487 **(a)** $x \geq 2$ **[B1]**

$y \geq 1$ **[B1]**

$x + y < 7$ or $y < -x + 7$ **[B1]**

(b) Finding critical values -0.45 and 4.45 (allow ≈ -0.449 and ≈ 4.449) **[B2]**

(Allow B1 for setting $x^2 - 4x - 2 = 0$ and finding one correct root or correct quadratic formula setup)

$-0.45 \leq x \leq 4.45$ **[B1]**

[Back to Question] *[Total Marks: 6]*

488 **(a)** Draw correct solid line $y = 2$

[B1]

Draw correct solid line $y = x + 2$

[B1]

Draw correct solid line $2x + y = 8$

[B1]

Correct shading of unwanted regions, leaving the triangle with vertices $(0, 2)$, $(3, 2)$, and $(2, 4)$ unshaded and clearly representing R.

[B1]

(b) Identify integer coordinates $(2, 4)$ or $(3, 2)$ as yielding the maximum value

[M1]

14

[A1]

[Back to Question] *[Total Marks: 6]*

489 **(a)** $y \geq 2$ **[B1]**

$y \leq x$ **[B1]**

$x + y \leq 8$ **[B1]**

(b) Identifies at least two valid integer coordinate pairs within the region R (e.g., $(6, 2)$, $(5, 3)$, $(4, 4)$) **[M1]**

Evaluates $2x + 3y$ for at least one critical integer point, e.g., $2(4) + 3(4) = 20$ or $2(5) + 3(3) = 19$ **[M1]**

20 **[A1]**

(c) Identifies the correct total number of integer points in R is 9, or identifies the 3 correct points where $x = y$: $(2, 2), (3, 3), (4, 4)$ **[M1]**

$\frac{3}{9}$ or $\frac{1}{3}$ **[A1]**

[Back to Question] *[Total Marks: 8]*

490 **(a)** $-4 < 2x \leq 8$ **[M1]**

$-2 < x \leq 4$ **[A1]**

(b) Multiplies throughout by 12 (or uses common denominator): $4(2x - 5) - 3(x - 3) > 12$ **[M1]**

Expands and simplifies: $8x - 20 - 3x + 9 > 12 \implies 5x - 11 > 12 \implies 5x > 23$ **[M1]**

5 **[A1]**

(c) Constructs correct inequality: $3.20 + 1.50k > 5.00 + 1.10k$ **[M1]**

Solves for k: $0.40k > 1.80 \implies k > 4.5$ **[M1]**

5 **[A1]**

[Back to Question] *[Total Marks: 8]*

491 **(a)** $x + y \leq 500$ **[B1]**

$y \geq 150$ **[B1]**

$5x + 10y \geq 3000$ (or simplified to $x + 2y \geq 600$) **[B1]**

(b) Line drawn correctly for $x + y = 500$ (passing through $(0, 500)$ and $(500, 0)$) **[B1]**

Line drawn correctly for $y = 150$ (horizontal line) **[B1]**

Line drawn correctly for $x + 2y = 600$ (passing through $(0, 300)$ and $(600, 0)$) **[B1]**

Correct region identified and all unwanted regions properly shaded (leaving quadrilateral with vertices $(0, 300), (0, 500), (350, 150), (300, 150)$ unshaded) **[B1]**

(c) Identifies the vertex $(350, 150)$ as giving the maximum x value **[M1]**

Children's tickets = 350, Adult tickets = 150 **[A1]**

[Back to Question] *[Total Marks: 9]*

492 **(a)** $x + y \leq 60$ **[B1]**

$x \geq 10$ **[B1]**

$y \geq 0.5x$ (or $2y \geq x$ or $y \geq \frac{1}{2}x$) **[B1]**

(b) Draws the line $x + y = 60$ correctly (from $(0, 60)$ to $(60, 0)$) **[B1]**

Draws the line $x = 10$ correctly (vertical line at 10) **[B1]**

Draws the line $y = 0.5x$ correctly (from $(0, 0)$ passing through $(20, 10)$, $(40, 20)$, etc.) **[B1]**

Shades the unwanted regions correctly, leaving the correct triangular region R unshaded **[B1]**

(c) Identifies $(40, 20)$ as the vertex for maximum profit (or tests multiple vertices) **[M1]**

2100 **[A1]**

[Back to Question] *[Total Marks: 9]*

493 **(a)** $x + y \leq 60$ **[B1]**

$x + 2y \leq 80$ **[B1]**

$y \geq 10$ **[B1]**

(b) B1 for correctly drawing the line $x + y = 60$

B1 for correctly drawing the line $x + 2y = 80$

B1 for correctly drawing the line $y = 10$

B1 for correctly shading the unwanted regions (above the two diagonal lines, and below the horizontal line) to leave the correct valid region unshaded

(c) M1 for a valid method to find the maximum profit, e.g., testing the vertex $(40, 20)$ in the objective function $5x + 8y$

360 **[A1]**

[Back to Question] *[Total Marks: 9]*

494 **(a)** Forms left inequality: $3(2x - 3) < 4(x + 5)$ or equivalent **[M1]**

Simplifies to $6x - 9 < 4x + 20$

x ¡ $\frac{29}{2}$ (or 14.5) [A1]

Forms right inequality: $2(x + 5) \leq 3(3x - 1)$ or equivalent **[M1]**

Simplifies to $2x + 10 \leq 9x - 3$

x $\geq \frac{13}{7}$ [A1]

$\frac{13}{7} \leq x < \frac{29}{2}$ (accept $1.86 \leq x < 14.5$ or equivalent double inequality) **[A1]**

(b) (i) $25 < 12 + 0.07x$ (or $12 + 0.07x > 25$) **[B1]**

(ii) $13 < 0.07x$ **[M1]**

$x > 185.71\ldots$ **[A1]**

186 **[A1]**

[Back to Question] *[Total Marks: 9]*

495 (a) $x + y \leq 60$ **[B1]**

$y \geq 10$ **[B1]**

$x \geq 15$ **[B1]**

$y \leq 2x$ **[B1]**

(b) Line $x + y = 60$ drawn correctly **[B1]**

Line $y = 10$ drawn correctly **[B1]**

Line $x = 15$ drawn correctly **[B1]**

Line $y = 2x$ drawn correctly **[B1]**

Correct region R indicated with all four unwanted regions shaded **[B1]**

(c) Identifies $(20, 40)$ as the vertex for maximum profit **[M1]**

Mountain bicycles = 20 and Road bicycles = 40 **[A1]**

Maximum profit = 2600 **[A1]**

[Back to Question] *[Total Marks: 12]*

496 (a) 27 **[B1]**

(b) Identifies the common difference of 4, e.g. writes $4n$ **[M1]**

$4n + 3$ **[A1]**

[Back to Question] *[Total Marks: 3]*

497 (a) 29 **[B1]**

(b) M1 for $6n + k$ where k is an integer

$6n - 1$ **[A1]**

(c) States that 150 is even but all terms in the sequence are odd OR solves $6n - 1 = 150$ to get $n = 25.1...$ and states n must be an integer **[B1]**

[Back to Question] *[Total Marks: 4]*

498 **(a)** 33 **[B1]**

(b) $7n + k$ for any integer k **[M1]**

$7n - 2$ **[A1]**

(c) $7n - 2 = 150$

$7\text{n} = 152 whichdoesnotgiveanintegervaluefor\text{n}.[B1]$

[Back to Question] *[Total Marks: 4]*

499 **(a)** 29 **[B1]**

(b) $6n + c$ or finding common difference of 6 **[M1]**

$6n - 1$ **[A1]**

(c) Sets $6n - 1 = 150 \implies 6n = 151$ and states that 151 is not divisible by 6 (or n is not an integer) **[B1]**

[Back to Question] *[Total Marks: 4]*

500 **(a)** Common difference is 4 **[M1]**

$4n + 1$ **[A1]**

(b) Identifies common ratio is 3 **[M1]**

$2 \times 3^{n-1}$ **[A1]**

[Back to Question] *[Total Marks: 4]*

501 **(a)** $23, 27$ **[B1]**

(b) $4n + k$ (for any k) **[M1]**

$4n + 3$ **[A1]**

(c) Equates $4n + 3 = 102$ and attempts to solve for n **[M1]**

$n = 24.75$, which is not a whole number/integer **[A1]**

[Back to Question] *[Total Marks: 5]*

502 **(a)** **(i)** 21 and 25 **[B1]**

(ii) Identifies common difference of 4 leading to $4n$ **[M1]**

$4n + 1$ **[A1]**

(b) Identifies common ratio of 3 **[M1]**

$2 \times 3^{n-1}$ **[A1]**

[Back to Question] *[Total Marks: 5]*

503 **(a)** 17 **[B1]**

(b) $4n + c$ where c is a constant **[M1]**

$4n + 1$ **[A1]**

(c) $4n + 1 = 141$ or $141 - 1 = 140$ **[M1]**

35 **[A1]**

[Back to Question] *[Total Marks: 5]*

504 **(a)** Identifying the second differences are constant 4 **[M1]**

Deducing the n^2 coefficient is 2, or setting up $2n^2 + bn + c$ **[M1]**

$2n^2 - n - 2$ **[A1]**

(b) $2n^2 - n - 2 = 433$ **[M1]**

$2n^2 - n - 435 = 0$ leading to correct factorisation $(2n + 29)(n - 15) = 0$ or correct substitution into the quadratic formula **[M1]**

15 (must reject negative solution if given) **[A1]**

[Back to Question] *[Total Marks: 6]*

505 **(a)** First differences: $10, 20, 34, 52$ **[M1]**

Second differences: $10, 14, 18$
Third differences: $4, 4$
Since the third difference is a non-zero constant, the sequence is cubic. **[A1]**

(b) Uses $6a = 4 \implies a = \frac{2}{3}$ **[M1]**

Uses $12a + 2b = 10 \implies 12(\frac{2}{3}) + 2b = 10 \implies 8 + 2b = 10 \implies b = 1$ **[M1]**

Uses $7a + 3b + c = 10 \implies 7(\frac{2}{3}) + 3(1) + c = 10 \implies c = 10 - 3 - \frac{14}{3} = \frac{7}{3}$ **[M1]**

Uses $a + b + c + d = 4 \implies \frac{2}{3} + 1 + \frac{7}{3} + d = 4 \implies 4 + d = 4 \implies d = 0$ **[M1]**

$\frac{2}{3}n^3 + n^2 + \frac{7}{3}n$ **[A1]**

[Back to Question] *[Total Marks: 7]*

506 Finds the first differences $6, 8, 10, 12$ and a constant second difference of 2 **[M1]**

Deduces that n^2 is part of the nth term expression **[M1]**

$n^2 + 3n - 1$ **[A1]**

[Back to Question] *[Total Marks: 3]*

507 **(a)** 79 **[B1]**

(b) Calculates second differences to be 4, or equates $2a = 4$ **[M1]**

Deduces that the sequence contains $2n^2$ **[M1]**

$2n^2 + n + 1$ **[A1]**

[Back to Question] *[Total Marks: 4]*

508 **(a)** $a + b = 5$ **[B1]**

$4a + 2b = 14$ **[B1]**

(b) M1 for a correct method to eliminate one variable, e.g. multiplying the first equation by 2 to give $2a + 2b = 10$ and subtracting from the second equation

$2a = 4$ leading to $a = 2$ **[A1]**

$b = 3$ **[A1]**

[Back to Question] *[Total Marks: 5]*

509 **(a)** One correct term evaluated **[M1]**

$3, 3, 5$ **[A1]**

(b) Second differences are 4 **[M1]**

Identifies $2n^2$ as part of the expression **[M1]**

$2n^2 + n - 1$ **[A1]**

[Back to Question] *[Total Marks: 5]*

510 **(a)** Finds first differences of $7, 13, 19$ and constant second differences of 6 **[M1]**

Deduces the quadratic term is $3n^2$ **[M1]**

$3n^2 - 2n + 1$ **[A1]**

(b) 35 **[B1]**

(c) $2^n + 3 = 131$ leading to $2^n = 128$ **[M1]**

7 **[A1]**

[Back to Question] *[Total Marks: 6]*

511 **(a)** Finds first differences: $3, 7, 11$ and second differences: $4, 4$ **[M1]**

Deduces the term involves $2n^2$ **[M1]**

$2n^2 - 3$ **[A1]**

(b) $3^n - 2 \approx 1000 \implies 3^n \approx 1002$ **[M1]**

Uses logarithms or trial and improvement: $n = 6 \implies 3^6 - 2 = 729 - 2 = 727$; $n = 7 \implies 3^7 - 2 = 2187 - 2 = 2185$ **[M1]**

727 **[A1]**

[Back to Question] *[Total Marks: 6]*

512 **(a)** 77 **[B1]**

(b) Finds first differences $(7, 11, 15, 19)$ and constant second difference of 4 **[M1]**

Deduces the coefficient of n^2 is 2 **[A1]**

Subtracts $2n^2$ from the sequence terms to identify the linear sequence $n - 1$ **[M1]**

$2n^2 + n - 1$ **[A1]**

(c) Sets up the equation $2k^2 + k - 1 = 464$ **[M1]**

Rearranges to form a quadratic equal to zero: $2k^2 + k - 465 = 0$ **[M1]**

15 **[A1]**

[Back to Question] *[Total Marks: 7]*

513 **(a)** Recognises the relationship to cube numbers $(1, 8, 27, 64, 125)$ **[M1]**

$n^3 + 1$ **[A1]**

(b) Identifies second difference is 2, or relates to square numbers **[M1]**

$n^2 + 2n$ or $(n + 1)^2 - 1$ **[A1]**

(c) $\dfrac{n^3 + 1}{n^2 + 2n}$ **[B1]**

(d) Substitutes $n = 12$ into their expression for sequence R, yielding $\dfrac{1728 + 1}{144 + 24}$ or $\dfrac{1729}{168}$ **[M1]**

Simplifies the fraction correctly to $\dfrac{247}{24}$ **[A1]**

[Back to Question] *[Total Marks: 7]*

514 **(a)** 217 **[B1]**

(b) M1 for finding the first differences $(7, 19, 37, 61)$ and second differences $(12, 18, 24)$

M1 for finding third differences are constant at 6, implying an n^3 term (or recognizing the sequence is $1^3 + 1, 2^3 + 1, 3^3 + 1, ...$)

$n^3 + 1$ **[A1]**

(c) M1 for setting up the equation $k^3 + 1 = 8001$

M1 for $k^3 = 8000$

$k = 20$ **[A1]**

[Back to Question] *[Total Marks: 7]*

515 **(a)** Identifies second difference of 2 or relates to square numbers **[M1]**

$n^2 + 1$ **[A1]**

(b) n^3 **[B1]**

(c) Recognises the relationship that Sequence C is Sequence B minus Sequence A **[M1]**

$n^3 - n^2 - 1$ **[A1]**

(d) Sets up the correct equation: $n^2 + 1 = 10001$ or $n^2 = 10000$ **[M1]**

100 **[A1]**

[Back to Question] *[Total Marks: 7]*

516 **(a)** Finds third differences are 6 (first diffs: $9, 25, 47, 75$; second diffs: $16, 22, 28$; third diffs: 6) **[M1]**

Identifies n^3 as the cubic term **[B1]**

Subtracts n^3 from the sequence to get a quadratic sequence $(-2, 0, 6, 16, 30)$ and finds its second differences are 4 to give $2n^2$ **[M1]**

$n^3 + 2n^2 - 4n$ **[A1]**

(b) $5 \times 3^{k-1} = 3645$

$3^{k-1} = 729$ [M1]

7 **[A1]**

(c) $\frac{2}{3}$ or 0.667 **[B1]**

[Back to Question] *[Total Marks: 7]*

517 **(a)** Finds first differences $14, 38, 74$ and second differences $24, 36$ **[M1]**

Finds third difference of 12 leading to a cubic term of $2n^3$ **[M1]**

Evaluates the sequence of residuals $T_n - 2n^3$ to find a constant difference of -1 **[M1]**

$2n^3 - 1$ **[A1]**

(b) **(i)** $V_1 = 1$, $V_2 = \frac{5}{4}$, $V_3 = \frac{5}{4}$ **[B2]**

Award B1 if any two are correct **[B1]**

(ii) Substitutes $n = 5$ to find $V_5 = \frac{25+1}{32} = \frac{13}{16}$ **[B1]**

Substitutes $n = 4$ to find $V_4 = \frac{16+1}{16} = \frac{17}{16}$ **[B1]**

Evaluates $\frac{13}{16} - \frac{17}{16} = -\frac{1}{4}$ (accept -0.25) **[A1]**

[Back to Question] *[Total Marks: 9]*

518 **(a)** Correct first differences $1, 9, 23, 43$ and second differences $8, 14, 20$ and third differences 6 **[M1]**

Deducing the cubic term coefficient is $a = 6 \div 6 = 1$ **[B1]**

Setting up equations to solve for other coefficients, e.g. $12a + 2b = 8$ **[M1]**

$n^3 - 2n^2 + 5$ **[A1]**

(b) $2a + b = 10$ **[B1]**

$8a + b = 34$ **[B1]**

Correct method to solve simultaneous equations, e.g. subtracting to get $6a = 24$ **[M1]**

$a = 4, b = 2$ **[A1]**

(c) Substituting $n = 12$ into their $4 \times 2^n + 2$ **[M1]**

16386 **[A1]**

[Back to Question] *[Total Marks: 10]*

519 **(a)** $4n + 1$ **[B2]**

If $4n + k$ seen, award **[M1]**

(b) **(i)** 182 **[B1]**

(ii) Identifies third differences are constant (6), so term contains n^3 **[M1]**

Subtracts n^3 terms $(1, 8, 27, 64, 125)$ from the sequence **[M1]**

Remainder sequence is $1, -2, -7, -14, -23$ **[M1]**

Deduces the remainder is $-n^2 + 2$ **[M1]**

$n^3 - n^2 + 2$ **[A1]**

(c) **(i)** $\frac{25}{182}$ **[B1]**

(ii) Recognises the pattern is Sequence A divided by Sequence B **[M1]**

$\frac{4n + 1}{n^3 - n^2 + 2}$ **[A1]**

[Back to Question] *[Total Marks: 10]*

520 **(a)** $y = kx^2$ or $36 = k(3^2)$ **[M1]**

$4x^2$ **[A1]**

(b) 100 **[B1]**

[Back to Question] *[Total Marks: 3]*

521 $y = k(x + 2)$ **[M1]**

$20 = k(3 + 2)$ leading to $k = 4$ **[M1]**

36 **[A1]**

[Back to Question] *[Total Marks: 3]*

522 **(a)** M1 for $y = kx^2$ or $36 = k(3)^2$

$4x^2$ **[A1]**

(b) M1 for $64 = 4x^2$ leading to $x^2 = 16$

4 and -4 (or ± 4) **[A1]**

[Back to Question] *[Total Marks: 4]*

523 **(a)** Sets up equation $y = kx$ and substitutes $x = 5, y = 15$ to get $15 = 5k$ **[M1]**

$y = 3x$ **[A1]**

(b) Substitutes $x = 12$ into their equation from part (a), e.g. $y = 3 \times 12$ **[M1]**

36 **[A1]**

[Back to Question] *[Total Marks: 4]*

524 **(a)** Substitutes into the proportion equation to find the constant, e.g. $135 = k(1+2)^3$ **[M1]**

$5(x+2)^3$ **[A1]**

(b) 40 **[B1]**

(c) Sets up the equation $320 = 5(x+2)^3$ and simplifies to $(x+2)^3 = 64$ **[M1]**

2 **[A1]**

[Back to Question] *[Total Marks: 5]*

525 **(a)** $y = kx^2$ leading to $36 = k(3)^2$ **[M1]**

$4x^2$ **[A1]**

(b) 100 **[B1]**

(c) $196 = 4x^2$ leading to $x^2 = 49$ **[M1]**

7 and -7 **[A1]**

[Back to Question] *[Total Marks: 5]*

526 **(a)** $y = kx^3$ **[M1]**

$24 = k(2)^3 \implies k = 3$ **[M1]**

$3x^3$ **[A1]**

(b) $y = 3(5)^3 = 3(125)$ **[M1]**

375 **[A1]**

(c) $81 = 3x^3$ **[M1]**

$x^3 = 27$ **[M1]**

3 **[A1]**

[Back to Question] *[Total Marks: 5]*

527 **(a)** Recognises inverse square proportion, e.g. tests p^2q **[M1]**

$2^2 \times 100 = 400$, $4^2 \times 25 = 400 \implies k = 400$ **[M1]**

$\dfrac{400}{p^2}$ **[A1]**

(b) $q = \dfrac{400}{5^2} = \dfrac{400}{25}$ **[M1]**

16 **[A1]**

(c) $1 = \dfrac{400}{p^2}$ **[M1]**

$p^2 = 400$ **[M1]**

20 (Accept ± 20) **[A1]**

[Back to Question] *[Total Marks: 5]*

528 **(a)** Sets up $P = k_1\sqrt{Q}$ and substitutes $Q = 9, P = 12 \implies 12 = k_1\sqrt{9} \implies 12 = 3k_1 \implies k_1 = 4$, so $P = 4\sqrt{Q}$ **[M1]**

Sets up $Q = \dfrac{k_2}{R^3}$ and substitutes $R = 2, Q = 2 \implies 2 = \dfrac{k_2}{2^3} \implies 2 = \dfrac{k_2}{8} \implies k_2 = 16$, so $Q = \dfrac{16}{R^3}$ **[M1]**

Substitutes Q into the P equation: $P = 4\sqrt{\dfrac{16}{R^3}}$ **[M1]**

Simplifies the square root: $P = 4 \times \dfrac{4}{\sqrt{R^3}}$ **[M1]**

$P = \dfrac{16}{R^{1.5}}$ or $P = 16R^{-1.5}$ or $P = \dfrac{16}{R\sqrt{R}}$ **[A1]**

(b) Substitutes $R = 4$ into their formula: $P = \dfrac{16}{4\sqrt{4}}$ or $P = \dfrac{16}{8}$ **[M1]**

2 **[A1]**

[Back to Question] *[Total Marks: 7]*

529 **(a)** M1 for setting up $A = kB^2$ and finding $k = 5$ (i.e., $A = 5B^2$)

M1 for setting up $B = \dfrac{m}{C+2}$ and finding $m = 12$ (i.e., $B = \dfrac{12}{C+2}$)

M1 for substituting $C = 4$ into their equation for B to get $B = 2$

20 **[A1]**

(b) M1 for substituting their expression for B into their equation for A: $A = 5\left(\dfrac{12}{C+2}\right)^2$

M1 for expanding or rearranging to isolate $(C+2)^2$: $(C+2)^2 = \dfrac{720}{A}$

M1 for taking the positive square root: $C + 2 = \sqrt{\dfrac{720}{A}}$

$\sqrt{\dfrac{720}{A}} - 2$ (or equivalent valid form e.g. $\dfrac{12\sqrt{5}}{\sqrt{A}} - 2$) **[A1]**

[Back to Question] *[Total Marks: 8]*

530 **(a)** Sets up equation $y = \dfrac{k}{x^2}$ **[M1]**

Substitutes $x = 4, y = 3$ to get $3 = \frac{k}{4^2} \implies k = 3 \times 16 = 48$ **[M1]**

$y = \frac{48}{x^2}$ **[A1]**

(b) Sets $0.75 = \frac{48}{x^2}$ **[M1]**

Rearranges to $x^2 = \frac{48}{0.75} = 64$ **[M1]**

$x = 8, x = -8$ (both required for final mark) **[A1]**

[Back to Question] *[Total Marks: 6]*

531 **(a)** $P = \frac{k}{Q^2}$ **[M1]**

$1.5 = \frac{k}{4^2}$ **[M1]**

$\frac{24}{Q^2}$ **[A1]**

(b) 0.24 or $\frac{6}{25}$ **[B1]**

(c) $24 = \frac{24}{Q^2}$ or $Q^2 = 1$ **[M1]**

1 **[A1]**

[Back to Question] *[Total Marks: 6]*

532 **(a)** **(i)** $P = \frac{k}{\sqrt{Q}}$ leading to $12 = \frac{k}{\sqrt{9}}$ **[M1]**

$k = 36$ **[B1]**

3 **[A1]**

(ii) $1.5 = \frac{36}{\sqrt{Q}}$ leading to $\sqrt{Q} = 24$ **[M1]**

576 **[A1]**

(b) Identifying that the new value of P involves dividing by $\sqrt{4}$ or 2 **[M1]**

50 **[A1]**

[Back to Question] *[Total Marks: 7]*

533 **(a)** $M = \frac{k}{\sqrt{N}}$ or $15 = \frac{k}{\sqrt{16}}$ **[M1]**

$\frac{60}{\sqrt{N}}$ **[A1]**

(b) $\frac{60}{\sqrt{100}}$ or $\frac{60}{10}$ **[M1]**

6 **[A1]**

(c) $1.2 = \dfrac{60}{\sqrt{N}}$ **[M1]**

$\sqrt{N} = \dfrac{60}{1.2}$ or $\sqrt{N} = 50$ **[M1]**

2500 **[A1]**

[Back to Question] *[Total Marks: 7]*

534 **(a)** M1 for recognising that y is inversely proportional to x^2 (e.g. testing $x^2y = c$)

M1 for a correct substitution to find k, e.g., $k = 4^2 \times 9 = 144$

$\dfrac{144}{x^2}$ **[A1]**

(b) M1 for $2.25 = \dfrac{144}{x^2}$ or $x^2 = 64$

8 **[A1]**

(c) M1 for substituting $3x$ into their equation: $y_{new} = \dfrac{144}{(3x)^2} = \dfrac{144}{9x^2}$

$\dfrac{1}{9}$ (or 0.111...) **[A1]**

[Back to Question] *[Total Marks: 7]*

535 **(a)** Sets up the correct proportional relationship, e.g. $p = \dfrac{k}{\sqrt{q}}$ or $5 = \dfrac{k}{\sqrt{144}}$ **[M1]**

$p = \dfrac{60}{\sqrt{q}}$ **[A1]**

(b) Substitutes $p = 15$ into their equation, e.g. $15 = \dfrac{60}{\sqrt{q}}$ or $\sqrt{q} = 4$ **[M1]**

16 **[A1]**

(c) Recognises that increasing by 300% means the new value of q is $4q$ **[B1]**

Substitutes $4q$ into the equation to find the new p: $p_{new} = \dfrac{60}{\sqrt{4q}} = \dfrac{60}{2\sqrt{q}} = \dfrac{1}{2}p$ **[M1]**

50 **[A1]**

[Back to Question] *[Total Marks: 7]*

536 **(a)** Substitutes into the proportion equation to find the constant, e.g. $4 = \dfrac{k}{\sqrt{36}}$ **[M1]**

$\dfrac{24}{\sqrt{Q}}$ **[A1]**

(b) 2 **[B1]**

(c) Sets up the equation $1.5 = \dfrac{24}{\sqrt{Q}}$ leading to $\sqrt{Q} = 16$ **[M1]**

256 **[A1]**

(d) Identifies the new value of Q is $4Q$, or uses numerical values (e.g. Q increases from 36 to 144) **[M1]**

Evaluates the new value of P as $\frac{1}{2}P$, or calculates the new P is 2 when original was 4 **[M1]**

50 **[A1]**

[Back to Question] *[Total Marks: 8]*

537 **(a)** Tests products or ratios, e.g., shows $x^2y = 300$ for all pairs or identifies $k = 300$ **[M1]**

Identifies $y \propto \frac{1}{x^2}$ as the correct model **[A1]**

$\frac{300}{x^2}$ **[A1]**

(b) Uses 1.5 multiplier and finds $(1.5)^3$ or 3.375 **[M1]**

Identifies increase factor as $3.375 - 1$ or 2.375 **[M1]**

237.5 **[A1]**

[Back to Question] *[Total Marks: 6]*

538 **(a)** Uses new $x = 1.2x$ to write $y_{new} = k(1.2x)^3$ **[M1]**

Calculates $1.2^3 = 1.728$ **[M1]**

72.8 **[A1]**

(b) Tests one incorrect model and shows the constant is not equal, e.g. for $A \propto v$, calculates $\frac{12}{2} = 6$ but $\frac{48}{4} = 12$ **[M1]**

Tests $A \propto v^2$ and shows $\frac{A}{v^2}$ is constant by calculating $\frac{12}{4} = 3$, $\frac{48}{16} = 3$, and $\frac{300}{100} = 3$ **[M1]**

$A \propto v^2$ **[A1]**

[Back to Question] *[Total Marks: 6]*

539 **(a)** **(i)** Recognises that x^2y is constant (128), or identifies the model as $y \propto \frac{1}{x^2}$ **[M1]**

Sets up the equation $y = \frac{k}{x^2}$ and correctly calculates $k = 128$ **[M1]**

$\frac{128}{x^2}$ **[A1]**

(ii) 1.28 **[B1]**

(b) Sets up the correct scale factor equation for B, e.g. $m^3 = 1.728$ or $\left(1 + \frac{x}{100}\right)^3 = 1.728$ **[M1]**

Calculates the multiplier $m = \sqrt[3]{1.728} = 1.2$ **[M1]**

20 **[A1]**

[Back to Question] *[Total Marks: 7]*

540 **(a)** Forms equation: $y = \dfrac{k}{\sqrt{x}}$ **[M1]**

Uses multiplier 1.44 for the increase in x: New $y = \dfrac{k}{\sqrt{1.44x}}$ **[M1]**

Simplifies denominator: $\sqrt{1.44} = 1.2$, so New $y = \dfrac{1}{1.2} \times \dfrac{k}{\sqrt{x}}$ (or $\dfrac{5}{6}y$) **[M1]**

Calculates percentage change: $\left(\dfrac{5}{6} - 1\right) \times 100$ **[M1]**

-16.7 (or 16.7 decrease) **[A1]**

(b) Deduces relationship between P and R: $P = k_1\left(\dfrac{k_2}{R}\right)^3 = \dfrac{K}{R^3}$ (or $P \propto \dfrac{1}{R^3}$) **[M1]**

Substitutes given values to find constant: $10 = \dfrac{K}{2^3} \implies K = 80$ **[M1]**

Substitutes $R = 4$ into their equation: $P = \dfrac{80}{4^3} = \dfrac{80}{64}$ **[M1]**

1.25 (or $\dfrac{5}{4}$) **[A1]**

[Back to Question] *[Total Marks: 7]*

541 **(a)** Tests a non-linear variation model and sets up an equation, e.g. $y = \dfrac{k}{x^2}$ **[M1]**

Correctly identifies the constant k by substituting a coordinate pair, e.g. $100 = \dfrac{k}{2^2} \implies k = 400$ **[M1]**

$y = \dfrac{400}{x^2}$ **[A1]**

(b) **(i)** Sets up $z = kx$ and finds k: $10 = 4k \implies k = 2.5$ **[M1]**

$\dfrac{z}{2.5}$ or $\dfrac{2z}{5}$ or $0.4z$ **[A1]**

(ii) Substitutes their expression for x into the equation from part (a): $y = \dfrac{400}{(\frac{2z}{5})^2}$ or $y = \dfrac{400}{(0.4z)^2}$ **[M1]**

Squares the fraction or decimal correctly: $y = \dfrac{400}{\frac{4z^2}{25}}$ or $y = \dfrac{400}{0.16z^2}$ **[M1]**

$\dfrac{2500}{z^2}$ **[A1]**

[Back to Question] *[Total Marks: 8]*

542 **(a)** Demonstrating that $y \times x^3$ is constant for at least two pairs (e.g., $40 \times 2^3 = 320$ and $5 \times 4^3 = 320$) **[M1]**

$y \propto \dfrac{1}{x^3}$ **[B1]**

$\dfrac{320}{x^3}$ **[A1]**

(b) $2560 = \frac{320}{x^3}$ leading to $x^3 = \frac{1}{8}$ **[M1]**

$\frac{1}{2}$ or 0.5 **[A1]**

(c) $x = \frac{k_2}{w}$ leading to $2 = \frac{k_2}{5}$ **[M1]**

$x = \frac{10}{w}$ **[A1]**

Substituting into their equation for y: $y = \frac{320}{(\frac{10}{w})^3}$ **[M1]**

$0.32w^3$ or $\frac{8}{25}w^3$ **[A1]**

[Back to Question] *[Total Marks: 9]*

543 **(a)** 4 **[B1]**

(b) 2 **[B1]**

[Back to Question] *[Total Marks: 2]*

544 **(a)** 4 **[B1]**

(b) Identifies period is 180° OR uses $360 \div 180$ **[M1]**

2 **[A1]**

[Back to Question] *[Total Marks: 3]*

545 **(a)** -5 **[B1]**

(b) M1 for substituting $(2, 7)$ and their c into the equation, e.g. $7 = a(2)^2 - 5$

3 **[A1]**

[Back to Question] *[Total Marks: 3]*

546 **(a)** 4 **[B1]**

(b) $a = 4$ **[B1]**

$b = 3$ **[B1]**

[Back to Question] *[Total Marks: 3]*

547 **(a)** $a = 3$ **[B1]**

$b = 2$ **[B1]**

(b) $(90, -3)$ **[B1]**

[Back to Question] *[Total Marks: 3]*

548 **(a)** $a = 4$ **[B1]**

$b = 2$ **[B1]**

(b) 180 **[B1]**

[Back to Question] *[Total Marks: 3]*

549 **(a)** **(i)** B **[B1]**

(ii) A **[B1]**

(iii) D **[B1]**

(iv) C **[B1]**

[Back to Question] *[Total Marks: 4]*

550 **(a)** 5 **[B1]**

(b) 180 **[B1]**

(c) $a = 5$ **[B1]**

$b = 2$ **[B1]**

[Back to Question] *[Total Marks: 4]*

551 **(a)** 4 **[B1]**

(b) 180 **[B1]**

(c) $a = 4$ **[B1]**

$b = 2$ **[B1]**

[Back to Question] *[Total Marks: 4]*

552 **(a)** $a = 5$ **[B1]**

$b = 3$ **[B1]**

(b) $-24 = k(2^3)$ or $8k = -24$ **[M1]**

-3 **[A1]**

[Back to Question] *[Total Marks: 4]*

553 **(a)** $a = 4$ **[B1]**

$b = 3$ **[B1]**

(b) 120 **[B1]**

(c) 6 **[B1]**

[Back to Question] *[Total Marks: 4]*

554 **(a)** -8 **[B1]**

(b) Substitutes an x-intercept from the graph, e.g. $4^2 + 4b - 8 = 0$ **[M1]**

-2 **[A1]**

(c) $(1, -9)$ [B2]
(If incorrect, B1 for $x = 1$ or $y = -9$)

[Back to Question] *[Total Marks: 5]*

555 A: $y = 2^x$ **[B1]**

B: $y = \dfrac{1}{x}$ **[B1]**

C: $y = -x^2$ **[B1]**

D: $y = x^3$ **[B1]**

[Back to Question] *[Total Marks: 4]*

556 **(a)** x-coordinate of minimum is $3 \implies -\dfrac{b}{2} = 3$ **[M1]**

$b = -6$ **[A1]**

Substitute $x = 3, y = -4, b = -6$ to find c: $-4 = 3^2 - 6(3) + c \implies c = 5$ **[A1]**

(b) $x^2 - 6x + 5 = 0$ **[M1]**

$(1, 0)$ and $(5, 0)$ **[A1]**

[Back to Question] *[Total Marks: 5]*

557 **(a)** 3 **[B1]**

(b) 2 **[B1]**

(c) M1 for setting up $3\sin(2x) = 1.5$ leading to $\sin(2x) = 0.5$

B1 for one correct value, e.g. $15°$ or $75°$

$15°$ and $75°$ **[A1]**

[Back to Question] *[Total Marks: 5]*

558 $c = -5$ **[B1]**

Uses line of symmetry $x = 2$ to form equation $-\dfrac{b}{2a} = 2 \implies b = -4a$ **[M1]**

Substitutes $(2, 3)$ and $c = -5$ into equation: $4a + 2b - 5 = 3$ **[M1]**

Solves simultaneous equations for a: $4a + 2(-4a) = 8 \implies -4a = 8 \implies a = -2$ **[M1]**

$b = -4(-2) = 8$ **[A1]**

[Back to Question] *[Total Marks: 5]*

559 **(a)** Substitute $(-1, 3)$: $(-1)^3 + a(-1)^2 + b(-1) = 3$

a - b = 4[$B1$]

Substitute $(2, 0)$: $(2)^3 + a(2)^2 + b(2) = 0$

2a + b = -4[$B1$]

(b) Valid method to solve simultaneous equations (e.g. addition: $3a = 0$) **[M1]**

$a = 0$ and $b = -4$ **[A1]**

(c) $x^3 - 4x = 0$ **[M1]**

$x(x^2 - 4) = 0$

x(x-2)(x+2) = 0[$M1$]

$(0, 0)$ and $(-2, 0)$ **[A1]**

[Back to Question] *[Total Marks: 7]*

560 **(a)** Substitutes $(3, 0)$ into $f(x)$ to get $a(3)^3 + b(3)^2 = 0 \implies 27a + 9b = 0$, leading to $3a + b = 0$. **[B1]**

(b) Substitutes $(2, -4)$ to form a second equation: $8a + 4b = -4$ (or $2a + b = -1$) **[M1]**

Solves simultaneous equations (e.g. substitutes $b = -3a$ into $2a + b = -1$) **[M1]**

$a = 1, b = -3$ **[A1]**

(c) Identifies the turning points at $y = 0$ and $y = -4$ as critical values **[M1]**

$k < -4$ or $k > 0$ **[A1]**

[Back to Question] *[Total Marks: 6]*

561 **(a)** 0 **[B1]**

(b) $(6, 0)$ **[B1]**

(c) Uses the vertex form $y = a(x - 3)^2 + 18$ OR substitutes $(3, 18)$ into $18 = 9a + 3b$ (knowing $c = 0$) **[M1]**

Uses a second point e.g., $(6, 0)$ to set up $0 = 36a + 6b$ OR substitutes $(0, 0)$ into vertex form: $0 = a(-3)^2 + 18$ **[M1]**

$a = -2, b = 12$ **[A1]**

(d) Identifies that $y = k$ must intersect the curve strictly between $y = 0$ and $y = 18$ on the positive x-axis **[M1]**

$0 < k < 18$ **[A1]**

[Back to Question] *[Total Marks: 7]*

562 **(a)** 0 **[B1]**

(b) Sets up equation for $x = 3$: $27a + 9b + 3c = 0$ (or $9a + 3b + c = 0$) **[M1]**

Sets up equation for $x = -3$: $-27a + 9b - 3c = 0$ (or $-9a + 3b - c = 0$) **[M1]**

Adds or subtracts equations correctly to deduce $18b = 0 \implies b = 0$, and substitutes to conclude $9a + c = 0$ **[A1]**

(c) Substitutes $x = 1, y = -16$ into the equation to get $a + c = -16$ **[M1]**

Solves $9a + c = 0$ and $a + c = -16$ simultaneously **[M1]**

$a = 2$, $c = -18$ **[A1]**

(d) $r = -5$ **[B1]**

$q = 0$ **[B1]**

Sets up $4p - 5 = 7$ to find $p = 3$ **[B1]**

[Back to Question] *[Total Marks: 10]*

563 **(a)** 4 **[B1]**

(b) 2 **[B1]**

(c) $4\sin(2 \times 30°)$ or $4\sin(60°)$ **[M1]**

3.46 **[A1]**

[Back to Question] *[Total Marks: 4]*

564 **(a)** $a = 4$ **[B1]**

$b = 2$ **[B1]**

(b) 5 **[B1]**

(c) 4 **[B1]**

[Back to Question] *[Total Marks: 4]*

565 **(a)** 2 **[B1]**

(b) Substitutes $(1, 6)$ into equation: $6 = \frac{a}{1} + 2$ **[M1]**

4 **[A1]**

(c) Substitutes $(2, 4)$ into $y = k^x$: $4 = k^2$ **[M1]**

2 **[A1]**

[Back to Question] *[Total Marks: 5]*

566 **(a)** 3 **[B1]**

(b) $7 = \frac{a}{2} + 3$ **[M1]**

$a = 8$ **[A1]**

(c) $\frac{8}{x} + 3 = 0$ **[M1]**

$\left(-\frac{8}{3}, 0\right)$ or $(-2.67, 0)$ **[A1]**

[Back to Question] *[Total Marks: 5]*

567 **(a)** 4 **[B1]**

(b) Recognises the period is $120°$ or sets $b \times 120 = 360$ **[M1]**

3 **[A1]**

(c) Recognises there are 2 solutions per period (or sketches horizontal line) **[M1]**

6 **[A1]**

[Back to Question] *[Total Marks: 5]*

568 **(a)** -4 **[B1]**

(b) Uses x-intercepts to set up equation: $a(-1)^2 + b(-1) - 4 = 0$ or $a(x+1)(x-4)$ **[M1]**

Substitute y-intercept $(0, -4)$ into factored form: $a(1)(-4) = -4 \implies a = 1$ **[M1]**

$b = -3$ **[A1]**

(c) Identifies x-coordinate of minimum as $x = \frac{-1+4}{2} = 1.5$ (or uses $x = \frac{-b}{2a}$) **[M1]**

$(1.5, -6.25)$ **[A1]**

[Back to Question] *[Total Marks: 6]*

569 **(a)** Uses $P(3, 8)$ in $y = a^x$: $a^3 = 8$ **[M1]**

$a = 2$ **[A1]**

Uses $P(3, 8)$ in $y = \frac{b}{x}$: $8 = \frac{b}{3} \implies b = 24$ **[B1]**

(b) $(0, 1)$ **[B1]**

(c) Gradient $= \frac{8-1}{3-0}$ **[M1]**

$\frac{7}{3}$ (or 2.33 or $2\frac{1}{3}$) **[A1]**

[Back to Question] *[Total Marks: 6]*

570 **(a)** **(i)** Graph C (or C) **[B1]**

(ii) Graph B (or B) **[B1]**

(iii) Graph D (or D) **[B1]**

(b) $p^3 = 125$ **[M1]**

$p = 5$ **[A1]**

(c) $\frac{q}{0.5} = 8$ **[M1]**

$q = 4$ **[A1]**

[Back to Question] *[Total Marks: 7]*

571 **(a)** $c = -5$ **[B1]**

Uses the roots to write $y = k(x+1)(x-5)$ **[M1]**

Expands to $x^2 - 4x - 5$ to find $a = 1$ and $b = -4$ **[A1]**

(b) $x = 2$ **[B1]**

$(2, -9)$ **[B1]**

(c) $2 = p \times q^0 \implies p = 2$ **[B1]**

$18 = 2 \times q^2 \implies q^2 = 9$ **[M1]**

$q = 3$ **[A1]**

[Back to Question] *[Total Marks: 8]*

572 **(a)** 4 **[B1]**

(b) Sets up equation using $(-1, 0)$: $(-1)^3 + a(-1)^2 + b(-1) + 4 = 0 \implies a - b = -3$ **[M1]**

Sets up equation using $(2, 0)$: $(2)^3 + a(2)^2 + b(2) + 4 = 0 \implies 4a + 2b = -12$ OR $2a + b = -6$ **[M1]**

Correct method to solve simultaneous equations (e.g. adding the equations) **[M1]**

$a = -3$ and $b = 0$ **[A1]**

(c) 3 **[B1]**

[Back to Question] *[Total Marks: 6]*

573 **(a)** 4 **[B1]**

(b) M1 for substituting $(-1, 0)$ to form an equation: $-1 + b - c + 4 = 0$ (or $b - c = -3$)

M1 for substituting $(2, 0)$ to form a second equation: $8 + 4b + 2c + 4 = 0$ (or $2b + c = -6$)

M1 for a valid method to solve their simultaneous equations

OR M2 for expanding $(x+1)(x-2)^2$

$b = -3$ **[A1]**

$c = 0$ **[A1]**

(c) 1 **[B1]**

[Back to Question] *[Total Marks: 6]*

574 **(a)** 12 **[B1]**

(b) Substitution of $(-2, 0)$ or $(2, 0)$ into the equation: $0 = -8 + 4b - 2c + 12$ or $0 = 8 + 4b + 2c + 12$ **[M1]**

Forming two correct simultaneous equations, e.g., $2b - c = -2$ and $2b + c = -10$ **[M1]**

$b = -3$ **[A1]**

$c = -4$ **[A1]**

(c) Setting up an equation for the roots, e.g., $(x-2)(x+2)(x-p) = x^3 - 3x^2 - 4x + 12$ OR $p^3 - 3p^2 - 4p + 12 = 0$ **[M1]**

$p = 3$ **[A1]**

[Back to Question] *[Total Marks: 7]*

575 **(a)** $8a + 2b = -16$ or $4a + b = -8$ **[M1]**

$-27a - 3b = 9$ or $-9a - b = 3$ **[M1]**

$a = 1, b = -12$ **[A1]**

(b) -2 **[B1]**

16 **[B1]**

(c) $k > 16$ **[B1]**

$k < -16$ **[B1]**

[Back to Question] *[Total Marks: 7]*

576 **(a)** $10 = a(2)^3 + b(2) \implies 8a + 2b = 10$ or $4a + b = 5$ **[M1]**

$4 = a(-1)^3 + b(-1) \implies -a - b = 4$ **[M1]**

Attempt to solve simultaneous equations (e.g., adding to eliminate b) **[M1]**

$a = 3$, $b = -7$ **[A1]**

(b) $3x^3 - 7x = 0$ **[M1]**

$x(3x^2 - 7) = 0 \implies x = \pm\sqrt{\frac{7}{3}}$ **[M1]**

Length $AB = 2 \times \sqrt{\frac{7}{3}} = \frac{2\sqrt{21}}{3}$ (or equivalent exact form) **[A1]**

[Back to Question] *[Total Marks: 7]*

577 **(a)** -3 **[B1]**

(b) Substitutes $(0, -1)$ into the equation: $a \cdot b^0 - 3 = -1$ **[M1]**

$a = 2$ **[A1]**

Substitutes $(1, 3)$ into the updated equation: $2 \cdot b^1 - 3 = 3$ **[M1]**

$b = 3$ **[A1]**

(c) Identifies that $g(1) = f(1)$, so $g(1) = 3$ **[M1]**

$\frac{k}{1} = 3 \implies k = 3$ **[A1]**

[Back to Question] *[Total Marks: 7]*

578 **(a)** Forms equation using roots: $f(x) = k(x+3)(x-1)(x-4)$ **[M1]**

Substitutes $(0, -24)$ to find k: $k(3)(-1)(-4) = -24$ **[M1]**

$12k = -24 \implies k = -2$ **[A1]**

Expands brackets: $-2(x^2 + 2x - 3)(x - 4) = -2(x^3 - 4x^2 + 2x^2 - 8x - 3x + 12)$ **[M1]**

Simplifies to: $-2(x^3 - 2x^2 - 11x + 12) = -2x^3 + 4x^2 + 22x - 24$ **[M1]**

$a = -2, b = 4, c = 22, d = -24$ **[A1]**

(b) Recognises translation moves the y-intercept up by 10 (calculates $-24 + 10$) **[M1]**

$(0, -14)$ **[A1]**

[Back to Question] *[Total Marks: 8]*

579 **(a)** Sets up simultaneous equations: $8 = a + b$ and $3.5 = \frac{a}{4} + b$ **[M1]**

Subtracts to eliminate b: $4.5 = \frac{3a}{4} \implies a = 6$ **[A1]**

$b = 2$ **[A1]**

(b) $y = 2$ **[B1]**

(c) **(i)** $x - 3 = \frac{6}{x} + 2$ **[M1]**

$x - 5 = \frac{6}{x} \implies x^2 - 5x = 6 \implies x^2 - 5x - 6 = 0$ **[A1]**

(ii) Solves $(x - 6)(x + 1) = 0$ to get $x = 6$ (rejects $x = -1$ as $x > 0$) **[M1]**

$(6, 3)$ **[A1]**

[Back to Question] *[Total Marks: 8]*

580 **(a)** **(i)** $f(-1) = (-1)^3 + b(-1)^2 + c(-1) + 4 = 0$ **[M1]**

$-1 + b - c + 4 = 0$ leading to $b - c = -3$ **[A1]**

(ii) Use point $(2, 0)$ to form equation: $2^3 + b(2)^2 + c(2) + 4 = 0$ **[M1]**

$8 + 4b + 2c + 4 = 0$ leading to $2b + c = -6$ **[A1]**

Attempt to solve simultaneous equations $b - c = -3$ and $2b + c = -6$ **[M1]**

$b = -3$, $c = 0$ **[A1]**

(b) $p \cdot q^0 = 2 \implies p = 2$ **[B1]**

$2 \cdot q^3 = 54 \implies q^3 = 27$ **[M1]**

$q = 3$ **[A1]**

[Back to Question] *[Total Marks: 9]*

581 **(a)** $r = 5$ **[B1]**

Uses vertex $(2, 9)$ to form an equation, e.g. $9 = p(2)^2 + q(2) + 5$ or uses vertex form $y = p(x - 2)^2 + 9$ **[M1]**

Forms a second equation using symmetry, e.g.

$$\frac{-q}{2p} = 2$$

q = -4p, *or substitutes* (0,5) *into vertex form* 5 = p(0-2)^2 + 9 [M1]

$p = -1$ and $q = 4$ **[A1]**

(b) Equating $f(x)$ to zero: $-x^2 + 4x + 5 = 0$

$x^2 - 4x - 5 = 0$ [M1]

$(5, 0)$ **[A1]**

(c) $a = 5$ **[B1]**

First positive intercept occurs when $5b = 180$ **[M1]**

$b = 36$ **[A1]**

[Back to Question] *[Total Marks: 9]*

582 **(a)** B1 for a smooth U-shaped parabola passing through $(0, 1)$, $(2, -3)$, and $(4, 1)$

B1 for correct endpoints plotted or stopped cleanly at $(-1, 6)$ and $(5, 6)$

(b) $(2, -3)$ **[B1]**

(c) 0.27 **[B1]**

3.73 **[B1]**

[Back to Question] *[Total Marks: 5]*

583 **(a)** Correct inverted U-shape parabola **[B1]**

Maximum approximately at $(3, 4)$ and crosses the x-axis at 1 and 5 **[B1]**

(b) $x = 1$ **[B1]**

$x = 5$ **[B1]**

(c) $(3, 4)$ **[B2]**

If B0, M1 for $x = 3$ or $y = 4$ identified

[Back to Question] *[Total Marks: 6]*

584 **(a)** Correct shape (inverted parabola) **[B1]**

Graph passes approximately through $(-0.5, 0)$, $(3, 0)$, $(0, 3)$ and vertex roughly at $(1.25, 6.1)$ **[B1]**

(b) $(1.25, 6.125)$ (Accept 6.13 or 6.1) **[B2]**

If B0, M1 for identifying correct x-coordinate 1.25 or correct derivative $-4x + 5 = 0$

(c) $x = -0.5$ **[B1]**

$x = 3$ **[B1]**

[Back to Question] *[Total Marks: 6]*

585 **(a)** Correct basic cubic shape with a local maximum and local minimum **[B1]**

Graph crosses y-axis at 2, has a maximum at $(0, 2)$ and a minimum at $(2, -2)$ **[B1]**

(b) Uses GDC to find roots **[M1]**

$-0.732, 1, 2.73$ (accept to 3 s.f.) **[A1]**

(c) Identifies correct turning point from GDC **[M1]**

$(2, -2)$ **[A1]**

[Back to Question] *[Total Marks: 6]*

586 **(a)** Correct U-shaped parabola passing through the y-axis at -3 **[B1]**

Vertices and endpoints approximately in the correct positions **[B1]**

(b) -0.5 **[B1]**

3 **[B1]**

(c) $x = 1.25$ **[B1]**

$y = -6.125$ **[B1]**

[Back to Question] *[Total Marks: 6]*

587 **(a)** B1 for a smooth cubic curve with y-intercept at $(0, 1)$ and correct general shape

B1 for correct turning points in the second and fourth quadrants and correct x-intercepts near -3.05, 0.52, and 2.53

(b) Local maximum $= (-1.63, 3.18)$ **[B1]**

Local minimum $= (1.63, -1.18)$ **[B1]**

(c) $-1.18 < k < 3.18$ **[B2]**

If 0 scored, allow B1 for identifying the boundary values -1.18 and 3.18

[Back to Question] *[Total Marks: 6]*

588 **(a)** Correct cubic shape with a local maximum and a local minimum **[B1]**

Local maximum approximately at $(-1, 3)$ and local minimum approximately at $(1, -1)$ **[B1]**

(b) $(-1, 3)$ **[B2]**

If B0, M1 for $x = -1$ or $y = 3$

(c) Correct straight line passing through $(0, -1)$ and $(2, 3)$ **[B1]**

(d) $x = 2$ **[B1]**

$x = 0.414$ and $x = -2.41$ **[B1]**

[Back to Question] *[Total Marks: 7]*

589 **(a)** Straight line for $g(x)$ passing through $(0, -2)$ and $(2, 0)$ **[B1]**

Two correct hyperbolic branches for $f(x)$ **[B1]**

Correct asymptotic behaviour near $x = -1$ **[B1]**

(b) $x = -1$ **[B1]**

(c) Sets up $\dfrac{2}{x+1} = x - 2$ or uses intersection feature on GDC **[M1]**

One correct coordinate pair: $(-1.56, -3.56)$ or $(2.56, 0.562)$ (to 3 s.f.) **[A1]**

Second correct coordinate pair **[A1]**

[Back to Question] *[Total Marks: 7]*

590 **(a)** Correct basic cubic shape **[B1]**

Local maximum near $x = 0$, local minimum near $x = 2.5$, and passing through $(0, 2)$ **[B1]**

(b) 1 **[B1]**

-0.562 (or $-0.561...$) **[B1]**

3.56 (or 3.561...) **[B1]**

(c) $x = 2.54$ (or 2.535...) **[B1]**

$y = -4.88$ (or $-4.879...$) **[B1]**

(d) Straight line passing through $(0, 1)$ with a negative gradient **[B1]**

(e) -0.468 (allow -0.47) **[B1]**

0.743 (allow 0.74) **[B1]**

3.73 **[B1]**

[Back to Question] *[Total Marks: 11]*

591 **(a)** Correct general cubic shape starting bottom left and ending top right **[B1]**

Local maximum correctly positioned at approximately $(-1, 4)$ **[B1]**

Local minimum correctly positioned at $(1, 0)$ **[B1]**

(b) $(-1, 4)$ **[B2]**

If B0, M1 for attempting to find stationary points e.g. evaluating max on calculator or $3x^2 - 3 = 0$

(c) $x = -2$ **[B1]**

$x = 1$ **[B1]**

(d) Straight line with positive gradient **[B1]**

Line passing through $(0, 1)$ and $(-0.5, 0)$ **[B1]**

(e) Uses calculator to find intersections or equates $x^3 - 3x + 2 = 2x + 1$ **[M1]**

-2.33 (or $-2.330\ldots$) **[B1]**

0.202 (or $0.2016\ldots$) and 2.13 (or $2.128\ldots$) **[B2, or B1 for 1 correct]**

[Back to Question] *[Total Marks: 12]*

592 **(a)** B1 for $y = f(x)$ starting at $(0, 0)$, rising to a single peak just above 1.5, and decaying towards the x-axis

B1 for $y = g(x)$ drawn as a straight line from $(0, 0.2)$ to $(5, 2.7)$

B1 for clearly showing two points of intersection in the first quadrant

(b) $(1.44, 1.59)$ **[B2]**

If 0 scored, allow B1 for $x = 1.44$ or $y = 1.59$

(c) M1 for finding the x-coordinates of the points of intersection: 0.08 (or $0.082\ldots$) and 2.46 (or $2.456\ldots$)

$0.08 < x < 2.46$ **[A1]**

[Back to Question] *[Total Marks: 7]*

593 **(a)** Graph touches the origin $(0, 0)$ **[B1]**

Graph reaches a maximum in the first quadrant and then approaches the x-axis asymptotically **[B1]**

Curve sketched with reasonable accuracy from $x = -1$ to $x = 5$, with $f(-1) = 2$ **[B1]**

(b) Uses maximum feature on GDC **[M1]**

$(2.89, 1.13)$ (accept $x = 2.88$ to 2.89 and $y = 1.12$ to 1.13) **[A1]**

(c) Draws line $y = 0.5$ on GDC or sets up equation **[M1]**

$-0.589, 0.835, 5.38$ (all three required, accept to 3 s.f.) **[A1]**

(d) Recognises the horizontal line $y = k$ must lie between the local minimum (0) and local maximum (1.13) **[M1]**

$0 < k < 1.13$ **[A1]**

[Back to Question] *[Total Marks: 9]*

594 **(a)** Correct W-shaped curve passing through the origin **[B1]**

Two minima (one in the third quadrant, one in the first quadrant) and one maximum roughly in correct places **[B1]**

Graph drawn between $x = -2$ and $x = 3$ with correct relative heights at the endpoints **[B1]**

(b) $(-1, -19)$ **[B1]**

$(2, 8)$ **[B1]**

(c) $(1, 13)$ **[B1]**

(d) Uses the y-values of the local extremum in the first quadrant to set bounds **[M1]**

$8 < k < 13$ **[A1]**

(e) $(1, 18)$ **[B1]**

[Back to Question] *[Total Marks: 9]*

595 **(a)** Correct shape with two distinct branches **[B1]**

Left branch passing through $(0, -2)$ and tending downwards as x

2^- [B1]

Right branch passing through $(4, 6)$ and tending upwards as x

2^+ [B1]

(b) $x = 2$ **[B1]**

(c) $(0, -2)$ **[B2]**

If B0, M1 for either coordinate correct

(d) $(4, 6)$ **[B2]**

If B0, M1 for either coordinate correct

(e) Identifies the critical values -2 and 6 from the local extrema **[M1]**

$-2 < k < 6$ **[A1]**

[Back to Question] *[Total Marks: 10]*

596 **(a)** Left branch in the 2nd and 3rd quadrants, passing through x-axis and asymptotic to y-axis downwards **[B1]**

Right branch in 1st quadrant with a clear minimum, asymptotic to y-axis upwards **[B1]**

Both branches curving upwards as x

$\pm\infty$ [B1]

(b) $(1, 3)$ **[B2]**

If B0, M1 for evaluating turning point on calculator or setting derivative to 0

(c) -1.26 (or $-1.259\ldots$) **[B1]**

(d) $x = 0$ **[B1]**

(e) Recognises need to find intersections of graph with $y = 4$ or rearranging to $x^3 - 4x + 2 = 0$ **[M1]**

-2.21 (or $-2.214\ldots$) **[B1]**

0.539 (or $0.5391\ldots$) and 1.68 (or $1.675\ldots$) **[B1]**

[Back to Question] *[Total Marks: 10]*

597 **(a)** $g(3) = 3^2 + 1 = 10$ **[M1]**

$f(10) = 2(10) - 5 = 15$ **[A1]**

(b) $y = 2x - 5$ OR $x = 2y - 5$ **[M1]**

$\dfrac{x+5}{2}$ **[A1]**

[Back to Question] *[Total Marks: 4]*

598 **(a)** 11 **[B1]**

(b) $\{-7, -4, 2, 11\}$ **[B1]**

(c) $x = 3y - 4$ or $y + 4 = 3x$ **[M1]**

$\frac{x+4}{3}$ **[A1]**

[Back to Question] *[Total Marks: 4]*

599 **(a)** 10 **[B1]**

(b) 5 **[B1]**

(c) $y = 3x - 2 \implies y + 2 = 3x$ **[M1]**

$\frac{x+2}{3}$ **[A1]**

[Back to Question] *[Total Marks: 4]*

600 **(a)** **(i)** -14 **[B1]**

(ii) $f(2) = 3(2) - 2 = 4$ **[M1]**

20 **[A1]**

(b) $y = 3x - 2 \implies y + 2 = 3x$ **[M1]**

$\frac{x+2}{3}$ **[A1]**

[Back to Question] *[Total Marks: 5]*

601 **(a)** -13 **[B1]**

(b) $y = 4x - 5$ or $x = 4y - 5$ **[M1]**

$\frac{x+5}{4}$ **[A1]**

(c) $4\left(\frac{x}{3} + 2\right) - 5$ **[M1]**

$\frac{4x}{3} + 3$ or $\frac{4x+9}{3}$ **[A1]**

[Back to Question] *[Total Marks: 5]*

602 **(a)** 18 **[B1]**

(b) $y = 5x - 2$ or $x = \frac{y+2}{5}$ **[M1]**

$\frac{x+2}{5}$ **[A1]**

(c) $g(2) = 2^2 + 3 = 7$ **[M1]**

$f(7) = 5(7) - 2 = 33$ **[A1]**

[Back to Question] *[Total Marks: 5]*

603 **(a)** 7 **[B1]**

(b) Let $y = 4x - 5$ leading to $y + 5 = 4x$ or equivalent **[M1]**

$\dfrac{x+5}{4}$ **[A1]**

(c) $g(2) = 6$ **[M1]**

19 **[A1]**

[Back to Question] *[Total Marks: 5]*

604 **(a)** 10 **[B1]**

(b) $y = 3x - 2 \implies 3x = y + 2$ **[M1]**

$\dfrac{x+2}{3}$ **[A1]**

(c) $f(2) = 3(2) - 2 = 4$ **[M1]**

$g(4) = 4^2 + 4 = 20$ **[A1]**

[Back to Question] *[Total Marks: 5]*

605 **(a)** 13 **[B1]**

(b) Finds $f(2) = 1$ **[M1]**

-2 **[A1]**

(c) Sets $y = 5 - 2x$ and rearranges to $2x = 5 - y$ **[M1]**

$\dfrac{5-x}{2}$ **[A1]**

[Back to Question] *[Total Marks: 5]*

606 **(a)** 5 **[B1]**

(b) $y = 4x - 3 \implies x = \dfrac{y+3}{4}$ or $x = 4y - 3$ **[M1]**

$\dfrac{x+3}{4}$ **[A1]**

(c) $f(1) = 4(1) - 3 = 1$ **[M1]**

$g(1) = 2 - 5(1) = -3$ **[A1]**

[Back to Question] *[Total Marks: 5]*

607 **(a)** Calculates $g(3) = 2(3)^2 + 1 = 19$ **[B1]**

Substitutes their 19 into $f(x)$: $4(19) - 3 = 73$ **[B1]**

(b) Let $y = 4x - 3$ and attempts to rearrange for x (e.g., $y + 3 = 4x$) **[M1]**

$\frac{x+3}{4}$ **[A1]**

(c) Sets $4x - 3 = 17$ **[M1]**

$x = 5$ **[A1]**

[Back to Question] *[Total Marks: 6]*

608 **(a)** 10 **[B1]**

(b) Sets $y = 3x - 2$ or $x = 3y - 2$ OR clearly shows inverse operations $(x + 2) \div 3$ **[M1]**

$\frac{x+2}{3}$ **[A1]**

(c) Substitutes $f(x)$ into $g(x)$: $(3x - 2)^2 + 1$ **[M1]**

Correct expansion of $(3x - 2)^2$: $9x^2 - 6x - 6x + 4$ OR $9x^2 - 12x + 4$ **[M1]**

$9x^2 - 12x + 5$ **[A1]**

[Back to Question] *[Total Marks: 6]*

609 **(a)** $y(x - 1) = 3x + 2$ **[M1]**

$xy - 3x = y + 2$ or $x(y - 3) = y + 2$ **[M1]**

$\frac{x+2}{x-3}$ **[A1]**

(b) $\frac{3x+2}{x-1} = x - 2$ **[M1]**

$3x + 2 = (x - 1)(x - 2)$ leading to $3x + 2 = x^2 - 3x + 2$ **[M1]**

$x^2 - 6x = 0$ or $x(x - 6) = 0$ **[M1]**

0 and 6 **[A1]**

[Back to Question] *[Total Marks: 7]*

610 **(a)** 1 **[B1]**

(b) $y = 2x + 3 \implies 2x = y - 3$ **[M1]**

$\frac{x-3}{2}$ **[A1]**

(c) $hk(x) = \frac{12}{(2x+3)-1}$ **[M1]**

$\frac{12}{2x+2} = \frac{6}{x+1}$ **[A1]**

(d) $k(2) = 2(2) + 3 = 7$ **[B1]**

$\frac{12}{x-1} = 7 \implies 12 = 7(x-1)$ **[M1]**

$12 = 7x - 7 \implies 19 = 7x \implies x = \frac{19}{7}$ (or $2\frac{5}{7}$) **[A1]**

[Back to Question] *[Total Marks: 8]*

611 **(a)** Sets $y = \frac{4}{2x-3}$ and multiplies by denominator to obtain $y(2x-3) = 4$ **[M1]**

Expands and rearranges to collect terms in x: $2xy = 4 + 3y$ **[M1]**

$\frac{4+3x}{2x}$ **[A1]**

(b) Finds $g(-3) = 2$ **[B1]**

Equates and attempts to solve: $\frac{4}{2x-3} = 2 \implies 4x - 6 = 4$ **[M1]**

2.5 (or $\frac{5}{2}$) **[A1]**

(c) Substitutes $f(x)$ into $g(x)$: $\frac{4}{2x-3} + 5$ **[M1]**

Expresses over a common denominator: $\frac{4+5(2x-3)}{2x-3}$ **[M1]**

$\frac{10x-11}{2x-3}$ **[A1]**

[Back to Question] *[Total Marks: 9]*

612 **(a)** Let $y = 5 - 3x$ leading to $3x = 5 - y$ or equivalent **[M1]**

$\frac{5-x}{3}$ **[A1]**

(b) $x^2 - 2 = 34$

$x^2 = 36$ [M1]

$x = 6, x = -6$ **[A1]**

(c) $5 - 3\left(\frac{x-3}{2}\right)$ **[M1]**

$\frac{10-3(x-3)}{2}$ or equivalent common denominator step **[M1]**

$\frac{19-3x}{2}$ **[A1]**

(d) $h(2) = 5 - 3(2) = -1$ **[M1]**

-1 **[A1]**

[Back to Question] *[Total Marks: 9]*

613 **(a)** $y = \frac{x+3}{x-1} \implies xy - y = x + 3$ **[M1]**

$xy - x = y + 3 \implies x(y-1) = y + 3$ **[M1]**

$\dfrac{x+3}{x-1}$ **[A1]**

(b) $\dfrac{\dfrac{x+3}{x-1}+3}{\dfrac{x+3}{x-1}-1}$ **[M1]**

$\dfrac{\dfrac{x+3+3(x-1)}{x-1}}{\dfrac{x+3-(x-1)}{x-1}}$ (or equivalent correct manipulation using a common denominator) **[M1]**

$\dfrac{x+3+3x-3}{x+3-x+1} = \dfrac{4x}{4} = x$ with no mathematical errors seen **[A1]**

(c) $f(3) = \dfrac{3+3}{3-1} = 3$ **[B1]**

$2x^2 - 5 = 3$ **[M1]**

$2x^2 = 8 \implies x^2 = 4$ **[M1]**

$2, -2$ **[A1]**

[Back to Question] *[Total Marks: 10]*

614 **(a)** $y(x-1) = 3x + 2$ or $x(y-1) = 3y + 2$ **[M1]**

$xy - 3x = y + 2$ or $x(y-3) = y + 2$ **[M1]**

$\dfrac{x+2}{x-3}$ **[A1]**

(b) $fg(x) = \dfrac{3\left(\dfrac{2}{x+1}\right)+2}{\dfrac{2}{x+1}-1}$ **[M1]**

Numerator becomes $\dfrac{6+2(x+1)}{x+1}$ or Denominator becomes $\dfrac{2-(x+1)}{x+1}$ **[M1]**

$\dfrac{2x+8}{1-x}$ **[A1]**

(c) $\dfrac{3x+2}{x-1} = \dfrac{x+2}{x-3} \implies (3x+2)(x-3) = (x+2)(x-1)$ **[M1]**

$3x^2 - 7x - 6 = x^2 + x - 2$ **[M1]**

$2x^2 - 8x - 4 = 0$ or $x^2 - 4x - 2 = 0$ **[M1]**

$2 \pm \sqrt{6}$ **[A1]**

[Back to Question] *[Total Marks: 10]*

615 **(a)** $y(x-1) = 3x + 5$ or $xy - y = 3x + 5$ **[M1]**

$xy - 3x = y + 5$ **[M1]**

$x(y-3) = y + 5$ **[M1]**

$\dfrac{x+5}{x-3}$ **[A1]**

(b) $\dfrac{3x+5}{x-1} = x$ **[M1]**

$3x + 5 = x^2 - x$ **[M1]**

$x^2 - 4x - 5 = 0$ **[M1]**

5 and -1 **[A1]**

(c) $f(2) = \dfrac{3(2)+5}{2-1} = 11$ **[M1]**

$11^2 + 1 = 122$ **[A1]**

[Back to Question] *[Total Marks: 10]*

616 **(a)** **(i)** $\dfrac{4}{(2x+3)-1}$ **[M1]**

$\dfrac{2}{x+1}$ **[A1]**

(ii) $\dfrac{4}{(x^2-4)-1} - 1$ **[M1]**

$\dfrac{4-(x^2-5)}{x^2-5}$ **[M1]**

$\dfrac{9-x^2}{x^2-5}$ **[A1]**

(b) $y(x-1) = 4$ or $x - 1 = \dfrac{4}{y}$ **[M1]**

$xy = 4 + y$ or $x = \dfrac{4}{y} + 1$ **[M1]**

$\dfrac{x+4}{x}$ **[A1]**

(c) $2\left(\dfrac{4}{x-1}\right) + 3 = 5$ **[M1]**

$\dfrac{8}{x-1} = 2$ **[M1]**

5 **[A1]**

[Back to Question] *[Total Marks: 11]*

617 **(a)** $y = \dfrac{2x+1}{3} \implies 3y = 2x + 1$ **[M1]**

$\dfrac{3x-1}{2}$ **[A1]**

(b) $hk(x) = \dfrac{2(5-x)+1}{3}$ **[M1]**

$\dfrac{11-2x}{3}$ **[A1]**

(c) $\dfrac{2x+1}{3} = 5 - x$ **[M1]**

$2x + 1 = 15 - 3x \implies 5x = 14$ **[M1]**

2.8 (or $\frac{14}{5}$) **[A1]**

[Back to Question] *[Total Marks: 7]*

618 **(a)** $h(5) = \frac{4}{5-3} = 2$ **[M1]**

$k(2) = 2(2) + 1 = 5$ **[A1]**

(b) $y(x-3) = 4$ OR $x(y-3) = 4$ **[M1]**

$\frac{4}{x} + 3$ OR $\frac{4+3x}{x}$ **[A1]**

(c) $h(2x+1) = \frac{4}{(2x+1)-3}$ **[M1]**

$\frac{4}{2x-2} = 2 \implies 4 = 4x - 4$ **[M1]**

$x = 2$ **[A1]**

[Back to Question] *[Total Marks: 7]*

619 **(a)** $\frac{1}{5}$ **[B1]**

(b) $(2x+3)^2 - 4$ **[M1]**

$4x^2 + 6x + 6x + 9 - 4$ **[M1]**

$4x^2 + 12x + 5$ **[A1]**

(c) $y = 2x + 3 \implies y - 3 = 2x$ **[M1]**

$\frac{x-3}{2}$ **[A1]**

(d) 3 **[B1]**

[Back to Question] *[Total Marks: 7]*

620 **(a)** 2 **[B1]**

(b) Sets $y = \frac{5}{x-2}$ and multiplies by $(x-2)$: $y(x-2) = 5$ **[M1]**

Rearranges to isolate x: $x - 2 = \frac{5}{y} \implies x = \frac{5}{y} + 2$ OR $xy - 2y = 5 \implies x = \frac{5+2y}{y}$ **[M1]**

$\frac{5}{x} + 2$ OR $\frac{5+2x}{x}$ **[A1]**

(c) Equates functions: $\frac{5}{x-2} = x + 2$ **[M1]**

Multiplies by $(x-2)$ to form a quadratic: $5 = (x+2)(x-2)$ **[M1]**

Expands and simplifies to $x^2 = 9$ or $x^2 - 9 = 0$ **[A1]**

3 and -3 **[A1]**

[Back to Question] *[Total Marks: 8]*

621 **(a)** $(-2)^2 - (-2) - 7$ **[M1]**

-1 **[A1]**

(b) $(2x-3)^2 - (2x-3) - 7$ **[M1]**

$4x^2 - 12x + 9 - 2x + 3 - 7$ **[M1]**

$4x^2 - 14x + 5$ **[A1]**

(c) $2x - 3 = x^2 - x - 7$ **[M1]**

$x^2 - 3x - 4 = 0$ or $(x-4)(x+1) = 0$ **[M1]**

$x = 4$ and $x = -1$ **[A1]**

[Back to Question] *[Total Marks: 8]*

622 **(a)** -4 and 5 seen **[B1]**

$-4 \leq f(x) \leq 5$ **[B1]**

(b) $g(3) = 7$ **[B1]**

$f(7) = 32$ **[B1]**

(c) $2(x^2 - 2x - 3) + 1$ **[M1]**

$2x^2 - 4x - 5$ **[A1]**

(d) $x = g(6)$ or $\dfrac{x-1}{2} = 6$ **[M1]**

13 **[A1]**

[Back to Question] *[Total Marks: 8]*

623 **(a)** Substitution of $f(x)$ into $g(x)$: $\dfrac{5}{\dfrac{2x+1}{3} - 2}$ **[M1]**

Writing the denominator as a single fraction: $\dfrac{2x+1-6}{3}$ or $\dfrac{2x-5}{3}$ **[M1]**

$\dfrac{15}{2x-5}$ **[A1]**

(b) Let $y = \dfrac{5}{x-2}$ and attempting to multiply by $(x-2)$ or swapping x and y **[M1]**

$x - 2 = \dfrac{5}{y}$ or $xy - 2y = 5$ **[M1]**

$\dfrac{5}{x} + 2$ or $\dfrac{5+2x}{x}$ **[A1]**

(c) Evaluates $g(3) = 5$ **[B1]**

Equating $f(x)$ to their $g(3)$: $\dfrac{2x+1}{3} = 5$ **[M1]**

$x = 7$ **[A1]**

[Back to Question] *[Total Marks: 9]*

624 **(a)** 5 **[B1]**

(b) $y(x-3) = 2x+1$ **[M1]**

$xy - 3y = 2x + 1$
$xy - 2x = 3y + 1$
$x(y-2) = 3y+1$ **[M1]**

$\dfrac{3x+1}{x-2}$ **[A1]**

(c) $(2x-1)^2 - 4$ **[M1]**

$4x^2 - 2x - 2x + 1 - 4$ **[M1]**

$4x^2 - 4x - 3$ **[A1]**

(d) $2x - 1 = x^2 - 4$
$x^2 - 2x - 3 = 0$ **[M1]**

$(x-3)(x+1) = 0$ **[M1]**

$x = 3, x = -1$ **[A1]**

[Back to Question] *[Total Marks: 10]*

625 **(a)** **(i)** 2.5 **[B1]**

(ii) $g(4-3x) = \dfrac{5}{(4-3x)-2}$ **[M1]**

$\dfrac{5}{2-3x}$ **[A1]**

(b) Minimum x-coordinate is $-\dfrac{-1}{2(1)} = 0.5$ or via completing square $(x-0.5)^2 - 7.25$ **[M1]**

$h(x) \geq -7.25$ **[A1]**

(c) $x^2 - x - 7 = 2x - 1$ leading to $x^2 - 3x - 6 = 0$ **[M1]**

$\dfrac{-(-3) \pm \sqrt{(-3)^2 - 4(1)(-6)}}{2(1)}$ **[M2]**

If M2 not scored, allow M1 for $\dfrac{3 \pm \sqrt{33}}{2}$ or correct use of formula with one error **[M1]**

-1.37 and 4.37 **[A1]**

[Back to Question] *[Total Marks: 9]*

626 **(a)** $y(x-1) = 2x+1 \implies xy - y = 2x + 1$ **[M1]**

$xy - 2x = y + 1 \implies x(y-2) = y+1$ **[M1]**

$\dfrac{x+1}{x-2}$ **[A1]**

(b) $g(3x-2) = (3x-2)^2 - 4(3x-2)$ **[M1]**

$9x^2 - 12x + 4 - 12x + 8$ **[M1]**

$9x^2 - 24x + 12$ **[A1]**

(c) $3(x^2 - 4x) - 2 = x$ **[M1]**

$3x^2 - 13x - 2 = 0$ **[A1]**

$x = \dfrac{13 \pm \sqrt{(-13)^2 - 4(3)(-2)}}{2(3)}$ **[M1]**

$x = 4.48$ or $x = -0.15$ **[A1]**

[Back to Question] *[Total Marks: 10]*

627 **(a)** Substitution of $g(x)$ into $f(x)$: $3(x^2 - x) + 2$ **[M1]**

$3x^2 - 3x + 2$ **[A1]**

(b) Substitution of $f(x)$ into $g(x)$: $(3x + 2)^2 - (3x + 2)$ **[M1]**

Correct expansion of $(3x + 2)^2$ to $9x^2 + 12x + 4$ **[M1]**

$9x^2 + 9x + 2$ **[A1]**

(c) Sets their three-term quadratic $gf(x)$ equal to 0: $9x^2 + 9x + 2 = 0$ **[M1]**

Valid method to solve quadratic (e.g., $(3x + 1)(3x + 2) = 0$ or correct use of formula) **[M1]**

$x = -\dfrac{1}{3}$ and $x = -\dfrac{2}{3}$ **[A1]**

(d) $5^x = 5^{-3}$ **[M1]**

$x = -3$ **[A1]**

[Back to Question] *[Total Marks: 10]*

628 **(a)** $x = 5y - 4$ or $y = 5x - 4$ **[M1]**

$\dfrac{x + 4}{5}$ **[A1]**

(b) $\dfrac{3}{(5x - 4) + 2}$ **[M1]**

$\dfrac{3}{5x - 2}$ **[A1]**

(c) $\dfrac{3}{\dfrac{3}{x + 2} + 2}$ **[M1]**

$\dfrac{3}{\dfrac{3 + 2(x + 2)}{x + 2}}$ **[M1]**

$\dfrac{3(x + 2)}{2x + 7}$ **[A1]**

(d) $\dfrac{3}{x + 2} = 5x - 4$

$3 = (5x - 4)(x + 2)$ **[M1]**

$3 = 5x^2 + 10x - 4x - 8$

$5x^2 + 6x - 11 = 0$ **[M1]**

$(5x + 11)(x - 1) = 0$ or correct use of quadratic formula **[M1]**

$x = 1, x = -2.2$ (or $-\frac{11}{5}$) **[A1]**

[Back to Question] *[Total Marks: 11]*

629 **(a)** Sets $y = \frac{2x+5}{x-3}$ and rearranges to get $y(x-3) = 2x+5$ **[M1]**

Expands and factorises to collect terms in x: $xy - 2x = 3y + 5 \implies x(y-2) = 3y+5$ **[M1]**

$\frac{3x+5}{x-2}$ **[A1]**

(b) Substitutes $g(x)$ into $f(x)$: $\frac{2(3x-1)+5}{(3x-1)-3}$ **[M1]**

Expands and simplifies the numerator to $6x+3$ **[M1]**

$\frac{6x+3}{3x-4}$ **[A1]**

(c) Sets up the equation $3\left(\frac{2x+5}{x-3}\right) - 1 = x$ **[M1]**

Eliminates the denominator to obtain $6x + 15 - (x-3) = x(x-3)$ (or equivalent) **[M1]**

Simplifies to form the correct 3-term quadratic equation: $x^2 - 8x - 18 = 0$ **[A1]**

Correct use of the quadratic formula $x = \frac{-(-8) \pm \sqrt{(-8)^2 - 4(1)(-18)}}{2(1)}$ **[M1]**

9.83 and -1.83 **[A1]**

[Back to Question] *[Total Marks: 11]*

630 **(a)** $y = \frac{3x+5}{x-2} \implies y(x-2) = 3x+5$ **[M1]**

$xy - 2y = 3x + 5 \implies xy - 3x = 2y + 5 \implies x(y-3) = 2y+5$ **[M1]**

$\frac{2x+5}{x-3}$ **[A1]**

(b) $\frac{3}{(3x+5)^2+2}$ **[M1]**

$\frac{3}{9x^2+30x+25+2} = \frac{3}{9x^2+30x+27}$ **[M1]**

$\frac{3}{3(3x^2+10x+9)} = \frac{1}{3x^2+10x+9}$ **[A1]**

(c) $\frac{3x+5}{x-2} = x \implies 3x+5 = x(x-2)$ **[M1]**

$3x + 5 = x^2 - 2x \implies x^2 - 5x - 5 = 0$ **[A1]**

$\frac{-(-5) \pm \sqrt{(-5)^2 - 4(1)(-5)}}{2(1)}$ **[M1]**

$\frac{5 \pm \sqrt{25+20}}{2} = \frac{5 \pm \sqrt{45}}{2}$ **[M1]**

5.85 and -0.85 **[A1]**

[Back to Question] *[Total Marks: 11]*

631 **(a)** Sets $y = \dfrac{2x+3}{x-1}$ and multiplies: $y(x-1) = 2x+3$ **[M1]**

Expands bracket and collects x terms on one side: $xy - 2x = y + 3$ **[M1]**

Factorises x: $x(y-2) = y+3$ **[M1]**

$\dfrac{x+3}{x-2}$ **[A1]**

(b) Substitutes $f(x)$ into itself: $\dfrac{2\left(\dfrac{2x+3}{x-1}\right)+3}{\left(\dfrac{2x+3}{x-1}\right)-1}$ **[M1]**

Forms common denominators for numerator and denominator: $\dfrac{\dfrac{2(2x+3)+3(x-1)}{x-1}}{\dfrac{(2x+3)-(x-1)}{x-1}}$ **[M1]**

Correctly simplified numerator: $4x+6+3x-3 = 7x+3$ OR correctly simplified denominator: $2x+3-x+1 = x+4$ **[B1]**

$\dfrac{7x+3}{x+4}$ **[A1]**

(c) Calculates $f(2) = \dfrac{2(2)+3}{2-1} = 7$ **[B1]**

Forms correct quadratic equation: $x^2 - 4x - 7 = 0$ **[B1]**

Correct substitution into quadratic formula: $\dfrac{4 \pm \sqrt{(-4)^2 - 4(1)(-7)}}{2}$ OR $\dfrac{4 \pm \sqrt{44}}{2}$ **[M1]**

5.32 and -1.32 **[A1]**

[Back to Question] *[Total Marks: 12]*

632 **(a)** 2 **[B1]**

(b) Let $y = \dfrac{x+5}{x-2}$ **[M1]**

$y(x-2) = x+5$

xy - 2y = x + 5[M1]

$xy - x = 2y + 5$

x(y - 1) = 2y + 5[M1]

$\dfrac{2x+5}{x-1}$ or equivalent **[A1]**

(c) $x^2 + 3x - 10 = 0$ **[M1]**

$(x+5)(x-2) = 0$ or correct substitution into quadratic formula **[M1]**

$x = -5, x = 2$ **[A1]**

(d)

$$\frac{x+5}{x-2} = x$$

$x^2 - 2x = x + 5$ [M1]

$x^2 - 3x - 5 = 0$ **[M1]**

$x = \dfrac{-(-3) \pm \sqrt{(-3)^2 - 4(1)(-5)}}{2}$ **[M1]**

$x = 4.19, x = -1.19$ **[A1]**

[Back to Question] *[Total Marks: 12]*

633 $y = (x + 3)(x - 4)$ **[M1]**

$x^2 - x - 12$ **[A1]**

[Back to Question] *[Total Marks: 2]*

634 $(x - 4)(x + 2)$ **[M1]**

$x^2 - 2x - 8$ **[A1]**

[Back to Question] *[Total Marks: 2]*

635 Uses $(x - p)(x - q)$ with $p = -4$ and $q = 3$: $(x + 4)(x - 3)$ **[M1]**

$x^2 + x - 12$ **[A1]**

[Back to Question] *[Total Marks: 2]*

636 Uses the intercepts to form $y = (x + 4)(x - 7)$ **[M1]**

$x^2 - 3x - 28$ **[A1]**

[Back to Question] *[Total Marks: 2]*

637 Uses the vertex form $y = (x - 4)^2 - 3$ **[M1]**

Expands to $x^2 - 8x + 16 - 3$ **[M1]**

$b = -8$ and $c = 13$ **[A1]**

[Back to Question] *[Total Marks: 3]*

638 Uses the vertex form $y = (x - 3)^2 - 4$ OR uses $-\dfrac{b}{2(1)} = 3$ **[M1]**

$b = -6$ **[A1]**

$c = 5$ **[A1]**

[Back to Question] *[Total Marks: 3]*

639 $y = (x - 3)^2 - 4$ **[M1]**

$y = x^2 - 6x + 9 - 4$ **[M1]**

$x^2 - 6x + 5$ **[A1]**

[Back to Question] *[Total Marks: 3]*

640 **(a)** $x^2 + bx + c = (x + 4)(x - 2)$ **[M1]**

$b = 2$, $c = -8$ **[A1]**

(b) Finds x-coordinate of vertex: $x = \dfrac{-4 + 2}{2} = -1$ **[M1]**

$(-1, -9)$ **[A1]**

[Back to Question] *[Total Marks: 4]*

641 **(a)** Uses vertex form: $y = (x - 3)^2 - 4$ **[M1]**

$x^2 - 6x + 5$ **[A1]**

(b) Sets $x^2 - 6x + 5 = 0 \implies (x - 1)(x - 5) = 0$ **[M1]**

$(1, 0)$ and $(5, 0)$ **[A1]**

[Back to Question] *[Total Marks: 4]*

642 **(a)** $h = 3$ **[B1]**

$k = -2$ **[B1]**

(b) $p = (5 - 3)^2 - 2$ **[M1]**

$p = 2$ **[A1]**

[Back to Question] *[Total Marks: 4]*

643 **(a)** $(x - 3)^2$ seen **[B1]**

$(x - 3)^2 - 4$ **[B1]**

(b) $y = x^2 - 6x + 9 - 4$ **[M1]**

$b = -6$ and $c = 5$ **[A1]**

(c) $(0, 5)$ **[B1]**

[Back to Question] *[Total Marks: 5]*

644 **(a)** Uses vertex form: $y = (x - 2)^2 - 9$ or x-coordinate of vertex $\dfrac{-p}{2} = 2$ **[M1]**

Expands to find p and q, or solves $2^2 + 2p + q = -9$ **[M1]**

$p = -4$ and $q = -5$ **[A1]**

(b) $x^2 - 4x - 5 = 0$ **[M1]**

$(x - 5)(x + 1) = 0$ **[M1]**

$A = (-1, 0)$ and $B = (5, 0)$ **[A1]**

[Back to Question] *[Total Marks: 6]*

645 $f(x) = a(x+2)(x-5)$ **[M1]**

$a(0+2)(0-5) = -20$ or $-10a = -20$ **[M1]**

$a = 2$ **[A1]**

$2x^2 - 6x - 20$ **[A1]**

[Back to Question] *[Total Marks: 4]*

646 $f(x) = a(x+2)(x-6)$ **[M1]**

$-24 = a(0+2)(0-6)$ **[M1]**

$a = 2$ **[A1]**

$2x^2 - 8x - 24$ **[A1]**

[Back to Question] *[Total Marks: 4]*

647 **(a)** Uses intercept form: $y = a(x+1)(x-4)$ **[M1]**

Substitutes $(0, -8)$ to find a: $-8 = a(0+1)(0-4) \implies -4a = -8 \implies a = 2$ **[M1]**

$2x^2 - 6x - 8$ **[A1]**

(b) Finds the x-coordinate of the vertex by symmetry: $x = \dfrac{-1+4}{2} = 1.5$ (or uses $x = -\dfrac{b}{2a}$) **[M1]**

Substitutes $x = 1.5$ into the equation: $y = 2(1.5)^2 - 6(1.5) - 8$ **[M1]**

$(1.5, -12.5)$ (or $(\dfrac{3}{2}, -\dfrac{25}{2})$) **[A1]**

[Back to Question] *[Total Marks: 6]*

648 **(a)** $y = a(x+2)(x-4)$ **[M1]**

$16 = a(0+2)(0-4)$ **[M1]**

$a = -2$ **[A1]**

$y = -2x^2 + 4x + 16$ **[A1]**

(b) $x = \dfrac{-2+4}{2} = 1$ **[M1]**

$(1, 18)$ **[A1]**

(c) $x = 1$ **[B1]**

[Back to Question] *[Total Marks: 7]*

649 **(a)** $h = -1$ **[B1]**

$k = 8$ **[B1]**

(b) $-10 = a(2-(-1))^2 + 8$ **[M1]**

$9a = -18$ **[M1]**

-2 **[A1]**

(c) $-2(x+1)^2+8$ **[M1]**

$-2(x^2+2x+1)+8$ **[M1]**

$-2x^2-4x+6$ **[A1]**

[Back to Question] *[Total Marks: 8]*

650 $f(x)=a(x+1)(x-4)$ **[M1]**

$-8=a(0+1)(0-4)$ **[M1]**

$a=2$ **[A1]**

$2x^2-6x-8$ **[A1]**

[Back to Question] *[Total Marks: 4]*

651 **(a)** Forms the equation $f(x)=a(x+2)^2+8$ **[M1]**

Substitutes $(0,-4)$ into their equation:

$-4=a(0+2)^2+8$ **[M1]**

$4a=-12 \implies a=-3$ **[A1]**

$-3x^2-12x-4$ **[A1]**

(b) Sets $-3(x+2)^2+8=0$ or $-3x^2-12x-4=0$ **[M1]**

Rearranges to $(x+2)^2=\frac{8}{3}$ OR correctly substitutes into the quadratic formula:

$x=\frac{-(-12)\pm\sqrt{(-12)^2-4(-3)(-4)}}{2(-3)}$ **[M1]**

$\frac{-6\pm2\sqrt{6}}{3}$ **[A1]**

[Back to Question] *[Total Marks: 7]*

652 $y=a(x-3)^2-5$ **[M1]**

$7=a(1-3)^2-5 \implies 12=4a \implies a=3$ **[M1]**

$3(x-3)^2-5$ **[A1]**

[Back to Question] *[Total Marks: 3]*

653 $y=a(x+2)^2+5$ **[M1]**

$-13=a(1+2)^2+5$ **[M1]**

$-18=9a$

a = -2$[M1]$

$-2(x+2)^2+5$ **[A1]**

[Back to Question] *[Total Marks: 3]*

654 **(a)** Uses vertex form: $y = a(x-2)^2 + 7$ **[M1]**

Substitutes $(4, -5)$ into their equation: $-5 = a(4-2)^2 + 7$ **[M1]**

$-3(x-2)^2 + 7$ **[A1]**

(b) -5 **[B1]**

[Back to Question] *[Total Marks: 4]*

655 Writes the function in the factored form $y = a(x+1)(x-5)$ **[M1]**

Substitutes $(2, -18)$ into their equation:

$-18 = a(2+1)(2-5)$ **[M1]**

$a = 2$ **[A1]**

$2x^2 - 8x - 10$ **[A1]**

[Back to Question] *[Total Marks: 4]*

656 Uses vertex form $y = a(x-2)^2 - 5$ **[M1]**

Substitutes $(4, 7)$ into their equation: $7 = a(4-2)^2 - 5$ **[M1]**

$4a = 12$

a = 3[$A1$]

$3x^2 - 12x + 7$ **[A1]**

[Back to Question] *[Total Marks: 4]*

657 **(a)** $h = 2$ **[B1]**

$k = 5$ **[B1]**

(b) Substitutes $(4, -7)$ into equation: $-7 = a(4-2)^2 + 5$ **[M1]**

$a = -3$ **[A1]**

(c) Substitutes $x = 0$ into $y = -3(x-2)^2 + 5$ **[M1]**

$(0, -7)$ **[A1]**

[Back to Question] *[Total Marks: 6]*

658 **(a)** Sets up the factored form $f(x) = a(x+2)(x-5)$ **[M1]**

Substitutes the point $(3, 10)$: $10 = a(3+2)(3-5)$ **[M1]**

Solves for a: $10 = a(5)(-2) \implies a = -1$ **[A1]**

$f(x) = -x^2 + 3x + 10$ **[A1]**

(b) Finds the x-coordinate of the vertex (e.g., midpoint of intercepts $\frac{-2+5}{2} = 1.5$ or $x = \frac{-b}{2a}$) **[M1]**

$(1.5, 12.25)$ **[A1]**

[Back to Question] *[Total Marks: 6]*

659 $y = a(x+1)^2 + 8$ **[M1]**

$-10 = a(2+1)^2 + 8$ **[M1]**

$9a = -18 \implies a = -2$ **[A1]**

$-2(x^2 + 2x + 1) + 8$ **[M1]**

$a = -2,\ b = -4,\ c = 6$ **[A1]**

[Back to Question] *[Total Marks: 5]*

660 (a) $y = a(x+3)(x-2)$ **[M1]**

$12 = a(0+3)(0-2)$ **[M1]**

$12 = -6a$

a = -2[$M1$]

$-2x^2 - 2x + 12$ **[A1]**

(b) $x = \dfrac{-3+2}{2}$ or $x = \dfrac{-(-2)}{2(-2)} = -0.5$ **[M1]**

$y = -2(-0.5)^2 - 2(-0.5) + 12 = 12.5$ **[M1]**

$(-0.5, 12.5)$ **[A1]**

[Back to Question] *[Total Marks: 6]*

661 (a) Sets up equation using x-intercepts: $f(x) = a(x+1.5)(x-4)$ OR $f(x) = a(2x+3)(x-4)$ **[M1]**

Substitutes $(2, -14)$ to find a: $-14 = a(2+1.5)(2-4)$ **[M1]**

Solves for a: $a = 2$ **[A1]**

$2x^2 - 5x - 12$ **[A1]**

(b) Identifies x-coordinate of vertex as midpoint of roots: $x = \dfrac{-1.5+4}{2} = 1.25$ OR uses $x = \dfrac{-b}{2a}$ **[M1]**

Substitutes $x = 1.25$ into their function: $y = 2(1.25)^2 - 5(1.25) - 12$ **[M1]**

$(1.25, -15.125)$ OR $\left(\dfrac{5}{4}, -\dfrac{121}{8}\right)$ **[A1]**

[Back to Question] *[Total Marks: 7]*

662 (a) (i) $h = -4$ **[B1]**

$k = 20$ **[B1]**

(ii) $0 = a(-2+4)^2 + 20$ **[M1]**

$a = -5$ **[A1]**

(iii) -6 **[B1]**

(iv) $f(x) = -5(x+4)^2 + 20 = -5(x^2 + 8x + 16) + 20$ **[M1]**

$-5x^2 - 40x - 60$ **[A1]**

[Back to Question] *[Total Marks: 7]*

663 **(a)** Sets up the equation in factored form: $f(x) = a(x+1)(x-5)$ **[M1]**

Substitutes $(0, 10)$ to find a: $10 = a(0+1)(0-5)$ **[M1]**

$a = -2$ **[A1]**

$-2x^2 + 8x + 10$ **[A1]**

(b) Finds the x-coordinate of the vertex using symmetry $x = \dfrac{-1+5}{2} = 2$ OR uses $x = -\dfrac{b}{2a}$ **[M1]**

$(2, 18)$ **[A1]**

(c) Translates the vertex $(2, 18)$

$(4,\ 15) OR applies transformation$ -2((x-2)-2)^2 + 18 − 3 [M1]

$-2(x-4)^2 + 15$ **[A1]**

[Back to Question] *[Total Marks: 8]*

664 **(a)** Uses the vertex coordinates to write $f(x) = a(x+2)^2 + 8$ **[M1]**

Substitutes the y-intercept $(0, 5)$: $5 = a(0+2)^2 + 8$ **[M1]**

$f(x) = -0.75(x+2)^2 + 8$ or equivalent using fractions like $-\dfrac{3}{4}$ **[A1]**

(b) Expands $(x+2)^2$ as $x^2 + 4x + 4$ **[M1]**

$f(x) = -0.75x^2 - 3x + 5$ **[A1]**

(c) Equates function to zero: $-0.75(x+2)^2 + 8 = 0$ or $-0.75x^2 - 3x + 5 = 0$ **[M1]**

Rearranges to $(x+2)^2 = \dfrac{32}{3}$ or correctly applies the quadratic formula **[M1]**

$x = 1.27$ and $x = -5.27$ **[A1]**

[Back to Question] *[Total Marks: 8]*

665 **(a)** $c = 15$ **[B1]**

$y = a(x+2)(x-5)$ **[M1]**

$15 = a(0+2)(0-5)$ **[M1]**

$a = -1.5$ **[A1]**

$b = 4.5$ **[A1]**

(b) $x = \dfrac{-2+5}{2} = 1.5$ **[M1]**

$y = -1.5(1.5)^2 + 4.5(1.5) + 15$ **[M1]**

$(1.5, 18.375)$ or $\left(\frac{3}{2}, \frac{147}{8}\right)$ **[A1]**

(c) $k > 18.375$ **[B1]**

[Back to Question] *[Total Marks: 9]*

666 **(a)** Uses vertex form: $y = a(x+2)^2 + 18$ **[M1]**

Substitutes $(0, 10)$ to find a: $10 = a(0+2)^2 + 18 \implies 4a = -8 \implies a = -2$ **[M1]**

$-2x^2 - 8x + 10$ **[A1]**

(b) Sets $f(x) = 0$: $-2x^2 - 8x + 10 = 0$ **[M1]**

$x^2 + 4x - 5 = 0 \implies (x+5)(x-1) = 0 \implies x = -5$ or $x = 1$ **[M1]**

6 **[A1]**

(c) Equates line and curve: $-2x^2 - 8x + 10 = -2x + k$ **[M1]**

Rearranges to form a quadratic equation: $2x^2 + 6x + (k-10) = 0$ (or equivalent) **[M1]**

Sets discriminant $\Delta = 0$ for a tangent: $6^2 - 4(2)(k-10) = 0$ **[M1]**

14.5 (or $\frac{29}{2}$) **[A1]**

[Back to Question] *[Total Marks: 10]*

667 **(a)** $h = 4$ and $k = -5$ **[B1]**

$8.5 = a(1-4)^2 - 5$ **[M1]**

$a = 1.5$ **[A1]**

(b) $1.5(x-4)^2 - 5 = 0$ or $(x-4)^2 = \frac{10}{3}$ **[M1]**

$x = 4 \pm \sqrt{\frac{10}{3}}$ **[M1]**

$x = 2.17$ and $x = 5.83$ **[A1]**

(c) $1.5(x-4)^2 - 5 = 2x - 7$ **[M1]**

$3x^2 - 28x + 52 = 0$ **[A1]**

$x = 2.56$ and $x = 6.77$ **[A1]**

(d) New vertex at $(1, 4)$ **[M1]**

$1.5(x-1)^2 + 4$ **[A1]**

[Back to Question] *[Total Marks: 11]*

668 **(a)** Forms equation using x-intercepts: $f(x) = a(x+1)(x-3)$ OR using vertex $f(x) = a(x-1)^2 - 8$ **[M1]**

Substitutes appropriate point to find a: $a(1+1)(1-3) = -8 \implies -4a = -8 \implies a = 2$ **[M1]**

Expands to show final form: $2(x^2 - 2x - 3) = 2x^2 - 4x - 6$ **[A1]**

(b) $2x^2 - 4x - 6 = 10 \implies 2x^2 - 4x - 16 = 0$ **[M1]**

Factors correctly to $(x-4)(x+2) = 0$ OR uses quadratic formula correctly **[M1]**

$x = 4$ and $x = -2$ **[A1]**

(c) Sets equations equal: $2x^2 - 4x - 6 = mx - 8$ leading to $2x^2 - (4+m)x + 2 = 0$ **[M1]**

States condition for tangent (discriminant $= 0$): $(-(4+m))^2 - 4(2)(2) = 0$ **[M1]**

Simplifies to $(4+m)^2 = 16$ or $m^2 + 8m = 0$ **[M1]**

$m = 0$, $m = -8$ **[A1]**

(d) Recognises minimum y-value must be strictly greater than 0: $-8 + k > 0$ **[M1]**

$k > 8$ **[A1]**

[Back to Question] *[Total Marks: 12]*

669 **(a)** $x = 2$ **[B1]**

(b) $y = 1$ **[B1]**

[Back to Question] *[Total Marks: 2]*

670 **(a)** $x = 4$ **[B1]**

(b) $y = 3$ **[B1]**

[Back to Question] *[Total Marks: 2]*

671 **(a)** -3 **[B1]**

(b) -2 **[B1]**

[Back to Question] *[Total Marks: 2]*

672 **(a)** $x = 2$ **[B1]**

(b) $y = 1$ **[B1]**

(c) $x = -2$ **[B1]**

[Back to Question] *[Total Marks: 3]*

673 **(a)** $x = 3$ **[B1]**

(b) $y = 1$ **[B1]**

(c) **(i)** 4 **[B1]**

(ii) 2 **[B1]**

[Back to Question] *[Total Marks: 4]*

674 **(a)** **(i)** $x = 1$ **[B1]**

(ii) $y = 3$ **[B1]**

(b) $\dfrac{4}{x-1} + 3 = 0$ **[M1]**

$(-\dfrac{1}{3}, 0)$ **[A1]**

(c) $(0, -1)$ **[B1]**

[Back to Question] *[Total Marks: 5]*

675 **(a)** $x = 90°$ **[B1]**

$x = 270°$ **[B1]**

(b) There is a vertical asymptote at $x = 90°$ (or the curve never touches $x = 90°$) **[B1]**

(c) $x = 125°$ **[B1]**

$x = 305°$ **[B1]**

(d) 225 **[B1]**

[Back to Question] *[Total Marks: 6]*

676 **(a)** $90°$ **[B1]**

$270°$ **[B1]**

(b) Vertical asymptote: $x = -2.5$ (or $-\dfrac{5}{2}$) **[B1]**

Horizontal asymptote: $y = 2$ **[B1]**

(c) $4x - 1 = 0$ **[M1]**

0.25 (or $\dfrac{1}{4}$) **[A1]**

[Back to Question] *[Total Marks: 6]*

677 **(a)** Vertical asymptote: $x = -2$ **[B1]**

Horizontal asymptote: $y = 1$ **[B1]**

(b) Substitute $y = 0$ to find x-intercept: $0 = \dfrac{3}{x+2} + 1 \implies x = -5$ **[M1]**

x-axis at $(-5, 0)$ and y-axis at $(0, 2.5)$ **[A1]**

(c) Two correct branches of the hyperbola sketched **[B1]**

Graph approaches $x = -2$ and $y = 1$ correctly without touching or crossing them **[B1]**

Graph passes through $(-5, 0)$ and $(0, 2.5)$ approximately **[B1]**

[Back to Question] *[Total Marks: 7]*

678 **(a)** $x = -2$ **[B1]**

(b) $y = -4$ **[B1]**

(c) Sets $x = 0$ to find the y-intercept: $y = \frac{3}{0+2} - 4$ **[M1]**

y-intercept is $(0, -2.5)$ **[A1]**

Sets $y = 0$ to find the x-intercept: $0 = \frac{3}{x+2} - 4$ **[M1]**

$4 = \frac{3}{x+2} \implies 4(x+2) = 3 \implies 4x + 8 = 3$ **[M1]**

x-intercept is $(-1.25, 0)$ **[A1]**

(d) $x = 90°$ **[B1]**

$x = 270°$ **[B1]**

[Back to Question] *[Total Marks: 9]*

679 **(a)** $b = 2$ **[B1]**

$c = -1$ **[B1]**

(b) Substitutes $(5, 0)$ into the equation: $\frac{a}{5-2} - 1 = 0$ **[M1]**

$a = 3$ **[A1]**

(c) Sets up the equation $\frac{3}{x-2} - 1 = -x + 5$ or uses GDC to find intersection of $y = f(x)$ and $y = -x + 5$ **[M1]**

$x = 3$ and $x = 5$ **[A1]**

[Back to Question] *[Total Marks: 6]*

680 **(a)** $q = 2$ **[B1]**

$r = -1$ **[B1]**

(b) Substitutes $(3, 3)$, $q = 2$ and $r = -1$ into $f(x)$: $3 = \frac{p}{3-2} - 1$ **[M1]**

$p = 4$ **[A1]**

(c) Sets $f(x) = 0$

$\frac{4}{x-2} - 1 = 0$ [M1]

$$\frac{4}{x-2} = 1$$

x - 2 = 4[M1]

$(6, 0)$ **[A1]**

[Back to Question] *[Total Marks: 7]*

681 **(a)** Two correct vertical asymptotes at $x = 90°$ and $x = 270°$ **[B1]**

Three correct branches passing through $(0, 0)$, $(180, 0)$, and $(360, 0)$ with correct general curvature **[B1]**

(b) $x = 90°$ **[B1]**

$x = 270°$ **[B1]**

(c) $\tan x = -1.5$ **[M1]**

$x = 123.7°$ **[A1]**

$x = 303.7°$ **[A1]**

[Back to Question] *[Total Marks: 7]*

682 **(a)** **(i)** $x = 1.5$ **[B1]**

(ii) $y = 2$ **[B1]**

(b) $y = 2$ is a horizontal asymptote so the curve never reaches $y = 2$ **[B1]**

(c) Vertical asymptote: $x = -1.5$ **[B1]**

Horizontal asymptote: $y = 2$ **[B1]**

(d) $\dfrac{5}{x - 1.5} + 2 = 3.4$ or $\dfrac{5}{x - 1.5} = 1.4$ **[M1]**

$x - 1.5 = \dfrac{5}{1.4}$ **[M1]**

5.07 **[A1]**

[Back to Question] *[Total Marks: 8]*

683 **(a)** $q = 5$ **[B1]**

$r = -3$ **[B1]**

(b) Substitutes $x = 7, y = -2$ and their values of q and r into the equation: $-2 = \dfrac{p}{7 - 5} - 3$ **[M1]**

$-2 = \dfrac{p}{2} - 3 \implies \dfrac{p}{2} = 1$ **[M1]**

$p = 2$ **[A1]**

(c) Sets $y = \dfrac{2}{x - 5} - 3$ and attempts to make x the subject, or swaps x and y **[M1]**

$y + 3 = \dfrac{2}{x - 5} \implies x - 5 = \dfrac{2}{y + 3}$ **[M1]**

$x = \dfrac{2}{y + 3} + 5$ **[M1]**

$f^{-1}(x) = \dfrac{2}{x + 3} + 5$ or equivalent **[A1]**

(d) $x = -3$ **[B1]**

$y = 5$ **[B1]**

(Strict follow through from their answers in part (a): $x = r, y = q$)

[Back to Question] *[Total Marks: 11]*

684 **(a)** Translation **[B1]**

by vector $\begin{pmatrix} 4 \\ 0 \end{pmatrix}$ **[B1]**

(b) $f(x) + 7$ **[B1]**

[Back to Question] *[Total Marks: 3]*

685 **(a)** $x^2 - 5$ **[B1]**

(b) $(x - 3)^2$ **[B1]**

(c) Translation **[B1]**

by vector $\begin{pmatrix} -4 \\ -1 \end{pmatrix}$ **[B1]**

[Back to Question] *[Total Marks: 4]*

686 **(a)** $g(x) = f(x) + 7$ **[M1]**

$g(x) = 2x + 2$ **[A1]**

(b) Translation **[B1]**

by vector $\begin{pmatrix} 4 \\ 0 \end{pmatrix}$ **[B1]**

[Back to Question] *[Total Marks: 4]*

687 **(a)** Horizontal shift 3 seen **[B1]**

$\begin{pmatrix} 3 \\ -2 \end{pmatrix}$ **[B1]**

(b) $(x - 3)^2$ seen **[B1]**

$(x - 3)^2 - 2$ **[B1]**

(c) $x = -5$ **[B1]**

$(-5, -1)$ **[B1]**

[Back to Question] *[Total Marks: 6]*

688 **(a)** $(-2, 1)$ **[B1]**

(b) $(5, -1)$ **[B1]**

(c) **(i)** $a = 2$ **[B1]**

$b = -5$ **[B1]**

(ii) $(5, -6)$ **[B1]**

(Strict follow through from their vector in (c)(i) applied to $(3, -1)$)

[Back to Question] *[Total Marks: 5]*

689 **(a)** $g(x) = f(x+2) = 3(x+2)^2$ **[M1]**

$3(x^2 + 4x + 4)$ **[M1]**

$3x^2 + 12x + 12$ **[A1]**

(b) $(-2, -8)$ **[B2]**

(Allow B1 for each correct coordinate)

(c) Substitutes $x = 0$ into $h(x) = 3(x+2)^2 - 8$ (or similar valid method) **[M1]**

4 **[A1]**

[Back to Question] *[Total Marks: 6]*

690 **(a)** Translation by 2 units left or 3 units down **[M1]**

Correct line segments joining $(-3, -2)$, $(-1, 1)$, $(1, -2)$, and $(2, -1)$ **[A1]**

(b) Translation **[B1]**

Vector $\begin{pmatrix} 0 \\ -5 \end{pmatrix}$ **[B1]**

(c) $f(x) - 5$ **[B1]**

(d) $x = 4 + 3 = 7$ or $y = 2 + 4 = 6$ **[M1]**

$(7, 6)$ **[A1]**

[Back to Question] *[Total Marks: 7]*

691 **(a)** $(4, -1)$ **[B1]**

(b) $f(x+3) + 2$ **[B2]**

(B1 for $f(x+3)$ and B1 for $+2$)

(c) $f(-k) = 0$, so $-k = 2$ or $-k = 6$ **[M1]**

$k = -2$ and $k = -6$ **[A1]**

(d) $f(x-3)$ **[B1]**

[Back to Question] *[Total Marks: 6]*

692 **(a)** $g(x) = f(x+4)$ **[M1]**

$(x+4)^2 - 2(x+4) - 3$ **[M1]**

$x^2 + 8x + 16 - 2x - 8 - 3$ **[M1]**

$x^2 + 6x + 5$ **[A1]**

(b) $h(x) = x^2 - 2x - 3 - 5 = x^2 - 2x - 8$ **[M1]**

$(x - 4)(x + 2) = 0$ **[M1]**

$x = 4$ and $x = -2$ **[A1]**

[Back to Question] *[Total Marks: 7]*

693 **(a)** Finds $(x - 3)^2$ as part of completing the square: $f(x) = (x - 3)^2 - 9 + 14$ **[M1]**

$f(x) = (x - 3)^2 + 5$ **[M1]**

Translation vector $\begin{pmatrix} 3 \\ 5 \end{pmatrix}$ **[A1]**

(b) Substitutes $x - 4$ into $g(x)$: $g(x - 4) = \dfrac{3}{(x - 4) + 2} - 5$ **[M1]**

Adds 2 to the function: $h(x) = \dfrac{3}{x - 2} - 5 + 2$ **[M1]**

$h(x) = \dfrac{3}{x - 2} - 3$ or equivalent single fraction $\dfrac{9 - 3x}{x - 2}$ **[A1]**

(c) $x = 2$ **[B1]**

$y = -3$ **[B1]**

(Strict follow through from their $h(x)$ in part (b))

[Back to Question] *[Total Marks: 8]*

694 **(a)** **(i)** $\dfrac{20}{x} - 4$ **[B1]**

(ii) $\dfrac{20}{x} - 4 = 0$ **[M1]**

$x = 5$ **[A1]**

(b) $h(x) = f(x - k) = \dfrac{20}{x - k}$ **[M1]**

$\dfrac{20}{8 - k} = 5$ **[M1]**

$k = 4$ **[A1]**

(c) Vertical asymptote: $x = 3$ **[B1]**

Horizontal asymptote: $y = 7$ **[B1]**

[Back to Question] *[Total Marks: 8]*

695 **(a)** **(i)** $f(x + 2) - 6$ **[B1]**

(ii) $2(x + 2)^2 - 5(x + 2) + 1 - 6$ **[M1]**

$2(x^2 + 4x + 4) - 5x - 15$ **[M1]**

$2x^2 + 3x - 7$ **[A1]**

(b) $2(x-p)^2 - 5(x-p) + 1 + q = 2x^2 - 13x + 18$ **[M1]**

$-4p - 5 = -13$ giving $p = 2$ **[A1]**

$2(2)^2 + 5(2) + 1 + q = 18$ giving $q = -1$ **[M1]**

$\begin{pmatrix} 2 \\ -1 \end{pmatrix}$ **[A1]**

(c) $2x^2 - 5x + 1 = 2x^2 + 3x - 7$ **[M1]**

$8x = 8$ giving $x = 1$ **[A1]**

$(1, -2)$ **[A1]**

[Back to Question] *[Total Marks: 11]*

696 **(a)** $\log_{10} x$ OR $\log x$ **[B1]**

(b) -2 **[B1]**

(c) $\log_{10} 45$ OR $\log 45$ **[B1]**

[Back to Question] *[Total Marks: 3]*

697 **(a)** $\log 1000 = 3$ **[B1]**

(b) 0.01 or $\frac{1}{100}$ **[B1]**

(c) $\frac{\log 12}{\log 5}$ **[B1]**

[Back to Question] *[Total Marks: 3]*

698 **(a)** 2 **[B1]**

(b) -3 **[B1]**

(c) x **[B1]**

[Back to Question] *[Total Marks: 3]*

699 **(a)** 4 **[B1]**

(b) $\log_5 125 = 3$ **[B1]**

(c) 64 **[B1]**

[Back to Question] *[Total Marks: 3]*

700 **(a)** 32 **[B1]**

(b) -2 **[B1]**

(c) $t = \log_k P$ **[B1]**

[Back to Question] *[Total Marks: 3]*

701 **(a)** $\log_5 125 = 3$ **[B1]**

(b) 3 **[B1]**

(c) $x = 2^5$ **[M1]**

32 **[A1]**

[Back to Question] *[Total Marks: 4]*

702 **(a)** $\log 100000 = 5$ **[B1]**

(b) 10^3 **[M1]**

1000 **[A1]**

(c) $\log x$ **[B1]**

[Back to Question] *[Total Marks: 4]*

703 **(a)** **(i)** 3
3 **[B1]**

(ii) 5
5 **[B1]**

(b)
=
3
4
=
81
x=3
4
=81 **[B1]**

(c) log
logx (or
log
10
log
10
x) **[B1]**

[Back to Question] *[Total Marks: 4]*

704 **(a)** **(i)** 5 **[B1]**

(ii) 0.01 or $\frac{1}{100}$ **[B1]**

(iii) 6 **[B1]**

(b) $\log x$ or $\log_{10} x$ **[B1]**

[Back to Question] *[Total Marks: 4]*

705 **(a)** 3 **[B1]**

(b) 10^{-2} or $\dfrac{1}{100}$ **[M1]**

0.01 **[A1]**

(c) $\log_5 125$ **[B1]**

[Back to Question] *[Total Marks: 4]*

706 **(a)** $\log_{10} 10000 = 4$ (or $\log 10000 = 4$) **[B1]**

(b) Uses index form $x = 5^3$ **[M1]**

125 **[A1]**

(c) $a = 7$ **[B1]**

$b = 25$ **[B1]**

[Back to Question] *[Total Marks: 5]*

707 **(a)** **(a)** $p(1) + q = 8$ or $p(2) + q = 11$ **[B1]**

A valid method to solve simultaneous equations e.g. $(2p + q) - (p + q) = 11 - 8$ **[M1]**

$p = 3$, $q = 5$ **[A1]**

(b) $3 \log x + 5 = 23 \implies 3 \log x = 18 \implies \log x = 6$ **[M1]**

1 000 000 or 10^6 **[A1]**

(c) $x = 3 \log y + 5 \implies \log y = \dfrac{x - 5}{3}$ **[M1]**

$10^{\frac{x-5}{3}}$ **[A1]**

[Back to Question] *[Total Marks: 7]*

708 **(a)** **(a)** $\log_6 y$ **[B1]**

(b) $x = \dfrac{\log 20}{\log 3}$ or $x \log 3 = \log 20$ **[M1]**

2.73 **[A1]**

[Back to Question] *[Total Marks: 3]*

709 **(a)** 250×1.08^{10} **[M1]**

540 **[A1]**

(b) $250 \times 1.08^t = 1000$ **[M1]**

$1.08^t = 4 \implies t = \dfrac{\log 4}{\log 1.08}$ **[M1]**

18.0 **[A1]**

[Back to Question] *[Total Marks: 5]*

710 **(a)** 15000×0.8^4 **[M1]**

6144 **[A1]**

(b) $7500 = 15000 \times 0.8^t$ **[M1]**

$0.5 = 0.8^t \implies t = \dfrac{\log 0.5}{\log 0.8}$ **[M1]**

3.11 **[A1]**

[Back to Question] *[Total Marks: 5]*

711 **(a)** $25000 \times (1 - 0.12)^4$ or 25000×0.88^4 **[M1]**

14992 **[A1]**

(b) $25000 \times 0.88^n = 10000$ or $0.88^n = 0.4$ **[M1]**

$n = \dfrac{\log 0.4}{\log 0.88}$ **[M1]**

$n = 7.16...$ **[A1]**

8 **[B1]**

[Back to Question] *[Total Marks: 5]*

712 **(a)** $2x = \dfrac{\log 50}{\log 3}$ **[M1]**

$2x = 3.56\ldots$ **[A1]**

1.78 **[A1]**

(b) $\log y = x - 4$ **[M1]**

$\log y + 4$ **[A1]**

[Back to Question] *[Total Marks: 5]*

713 **(a)** **(a)** 5000 **[B1]**

(b) $5000 \times 1.12^t = 20000$ **[M1]**

$1.12^t = 4$ **[M1]**

$t = \dfrac{\log 4}{\log 1.12}$ **[M1]**

12.2 **[A1]**

[Back to Question] *[Total Marks: 5]*

714 **(a)** 8000×1.035^n **[B1]**

(b) Sets up the inequality or equation: $8000 \times 1.035^n > 12000$ OR $1.035^n = 1.5$ **[M1]**

Uses logarithms to solve for n: $n = \dfrac{\log_{10} 1.5}{\log_{10} 1.035}$ **[M1]**

12 **[A1]**

(c) $x = \dfrac{\log_{10} 55}{\log_{10} 2}$ **[M1]**

5.78 **[A1]**

[Back to Question] *[Total Marks: 6]*

715 **(a)** $x \log 3 = \log 20$ **[M1]**

$x = \dfrac{\log 20}{\log 3}$ **[M1]**

2.73 **[A1]**

(b) $100 = 500 \times 0.85^t \implies 0.85^t = 0.2$ **[M1]**

$t \log 0.85 = \log 0.2 \implies t = \dfrac{\log 0.2}{\log 0.85}$ **[M1]**

9.90 **[A1]**

[Back to Question] *[Total Marks: 6]*

716 **(a)** 4500×1.08^5 **[M1]**

6612 **[A1]**

(b) $4500 \times 1.08^t = 10000$ **[M1]**

$1.08^t = \dfrac{10000}{4500}$ oe **[M1]**

$t = \dfrac{\log(20/9)}{\log 1.08}$ **[M1]**

10.38 **[A1]**

[Back to Question] *[Total Marks: 6]*

717 **(a)** 0.78
0.78 (accept
0.75
0.75 to
0.80
0.80) **[B1]**

(b) 3.2
3.2 (accept
3.1
3.1 to
3.3

3.3) **[B1]**

(c) (
+
1
)
log
3
=
log
20
(x+1)log3=log20 or
+
1
=
log
20
log
3
x+1=
log3
log20
[M1]

+
1
=
2.7268
...
x+1=2.7268... **[M1]**

1.73
1.73 **[A1]**

(d) 2
3
=
10
1.4
2x3=10
1.4
(or
2
3
=
25.118
...
2x3=25.118...) **[M1]**

14.1
14.1 **[A1]**

[Back to Question] *[Total Marks: 7]*

718 **(a)** Substitutes $t = 5$ into the formula: 400×0.88^5 **[M1]**

211.1 **[A1]**

(b) Sets up the equation $400 \times 0.88^t = 200$ and simplifies to $0.88^t = 0.5$ **[M1]**

$t = \dfrac{\log 0.5}{\log 0.88}$ **[M1]**

5.42 **[A1]**

(c) Sets up the equation $800 \times k^{10} = 300$ **[M1]**

$k^{10} = 0.375$ **[M1]**

0.907 **[A1]**

[Back to Question] *[Total Marks: 8]*

719 **(a)** 8000×1.045^5 **[M1]**

9969.46 (accept 9969 or 9970) **[A1]**

(b) $1.045^t = \dfrac{15000}{8000}$ or 1.875 **[M1]**

$t = \dfrac{\log 1.875}{\log 1.045}$ or 14.28... **[M1]**

15 **[A1]**

(c) $\log(7^{x-1}) = \log 250$ or $x - 1 = \log_7 250$ **[M1]**

$x - 1 = \dfrac{\log 250}{\log 7}$ or $x - 1 = 2.837...$ **[M1]**

3.84 **[A1]**

[Back to Question] *[Total Marks: 8]*

720 **(a)** 12800 **[B1]**

(b) $800 \times 2^t = 102400 \implies 2^t = 128$ **[M1]**

$t = 7$ **[A1]**

(c) $1500 \times 3^t = 20000 \implies 3^t = \dfrac{40}{3}$ (or 13.33...) **[M1]**

Uses logarithms to solve $3^t = \dfrac{40}{3}$, e.g. $t = \dfrac{\log(40/3)}{\log 3}$ **[M1]**

2.36 (from 2.358...) **[A1]**

(d) $2000 \times 1.05^n = 4000 \implies 1.05^n = 2$ **[M1]**

$n = \dfrac{\log 2}{\log 1.05}$ (or 14.2...) **[M1]**

15 **[A1]**

[Back to Question] *[Total Marks: 9]*

721 **(a)** $y = 4 \times 10^{2x} - 7 \implies y + 7 = 4 \times 10^{2x}$ **[M1]**

$\dfrac{y+7}{4} = 10^{2x} \implies 2x = \log\left(\dfrac{y+7}{4}\right)$ **[M1]**

$\dfrac{1}{2}\log\left(\dfrac{x+7}{4}\right)$ **[A1]**

(b) $4 \times 10^{2x} - 7 = 53 \implies 4 \times 10^{2x} = 60$ **[M1]**

$10^{2x} = 15 \implies 2x = \log 15$ **[M1]**

0.588 **[A1]**

[Back to Question] *[Total Marks: 6]*

722 **(a)** Exponential curve passing through $(0, 1)$ and $(1, 2)$ **[B1]**

Curve passing through $(3, 8)$ and approaching the negative x-axis as an asymptote **[B1]**

(b) $\dfrac{\log x}{\log 2}$ (or $\log_2 x$) **[B1]**

(c) Logarithmic curve passing through $(1, 0)$ and $(2, 1)$ **[B1]**

Curve passing through $(8, 3)$ and approaching the negative y-axis as an asymptote **[B1]**

(d) $x = \dfrac{\log 12}{\log 2}$ **[M1]**

3.58 **[A1]**

[Back to Question] *[Total Marks: 7]*

723 **(a)** **(i)** 250 **[B1]**

(ii) $250 \times 0.92^t = 50$ or $0.92^t = 0.2$ **[M1]**

$t = \dfrac{\log 0.2}{\log 0.92}$ **[M1]**

19.3 **[A1]**

(b) $800 \times 1.06^n > 2000$ or $1.06^n = 2.5$ **[M1]**

$n = \dfrac{\log 2.5}{\log 1.06}$ $(= 15.7\ldots)$ **[M1]**

16 **[A1]**

[Back to Question] *[Total Marks: 7]*

724 **(a)** Substitutes given values into the formula: $250 = 800 \times 10^{-15k}$ **[M1]**

Rearranges to isolate the exponential: $10^{-15k} = 0.3125$ **[M1]**

Takes logarithms to solve for k: $-15k = \log_{10}(0.3125) \implies k = \dfrac{\log_{10}(0.3125)}{-15} = 0.03367...$ **[A1]**

(b) Substitutes $m = 50$ and $k = 0.03367...$ into the formula: $50 = 800 \times 10^{-0.03367t}$ **[M1]**

Isolates the exponential and takes logarithms: $-0.03367t = \log_{10}\left(\dfrac{50}{800}\right)$ **[M1]**

35.8 (accept 35.7 to 35.8) **[A1]**

(c) Rearranges to $10^{-kt} = \dfrac{m}{800}$ and converts to logarithmic form: $-kt = \log_{10}\left(\dfrac{m}{800}\right)$ **[M1]**

$t = -\dfrac{1}{k}\log_{10}\left(\dfrac{m}{800}\right)$ OR $t = \dfrac{1}{k}\log_{10}\left(\dfrac{800}{m}\right)$ **[A1]**

[Back to Question] *[Total Marks: 8]*

725 **(a)** $g(100) = \log 100 = 2$ **[M1]**

$f(2) = 5 \times 10^2 = 500$ **[A1]**

(b) $y = 5 \times 10^x$ leads to $10^x = \dfrac{y}{5}$ **[M1]**

$\log\left(\dfrac{x}{5}\right)$ or $\log x - \log 5$ **[A1]**

(c) $5 \times 10^x = 200$ leads to $10^x = 40$ **[M1]**

$x = \log 40 = 1.602...$ **[A1]**

1.60 **[B1]**

(d) $g(f(x)) = \log(5 \times 10^x)$ **[M1]**

$= \log 5 + \log 10^x = \log 5 + x$ **[M1]**

$x + \log 5$ **[A1]**

[Back to Question] *[Total Marks: 8]*

726 **(a)** $5 \times 3^{-2} - 2$ **[M1]**

$-\dfrac{13}{9}$ **[A1]**

(b) $y = 5 \times 3^x - 2 \implies y + 2 = 5 \times 3^x$ **[M1]**

$3^x = \dfrac{y+2}{5}$ **[M1]**

$\dfrac{\log\left(\frac{x+2}{5}\right)}{\log 3}$ **[A1]**

(c) $5 \times 3^x - 2 = 500 \implies 3^x = 100.4$ **[M1]**

$x = \dfrac{\log 100.4}{\log 3}$ **[M1]**

4.20 **[A1]**

[Back to Question] *[Total Marks: 8]*

727 **(a)** $3 = \log_a 8$ **[M1]**

$a^3 = 8 \implies a = 2$ **[A1]**

(b) **(i)** Smooth exponential curve passing through $(0, 1)$ **[B1]**

Correct shape passing through $(1, 2)$, $(2, 4)$ and $(3, 8)$ **[B1]**

(ii) $y = 0$ (accept the x-axis) **[B1]**

(c) Rearranges to $\log_2 x = 8 - x$ **[M1]**

Draws the line $y = 8 - x$ correctly on the grid (passing through $(0, 8)$ and $(8, 0)$) **[M1]**

$x = 5.5$ (accept 5.4 to 5.6) **[A1]**

[Back to Question] *[Total Marks: 8]*

728 **(a)** Substitutes $x = 3$: $4 \times 10^{1.5}$ **[M1]**

126 **[A1]**

(b) Sets $y = 4 \times 10^{0.5x}$ and rearranges to $\frac{y}{4} = 10^{0.5x}$ **[M1]**

$\log\left(\frac{y}{4}\right) = 0.5x$ **[M1]**

$2\log\left(\frac{x}{4}\right)$ **[A1]**

(c) $4 \times 10^{0.5x} = 200$

$10^{0.5x} = 50$ [M1]

$0.5x = \log 50$ **[M1]**

3.40 **[A1]**

[Back to Question] *[Total Marks: 8]*

729 **(a)** $a \times b^0 = 12 \implies a = 12$ **[B1]**

$12 \times b^3 = 96 \implies b^3 = 8$ **[M1]**

$b = 2$ **[A1]**

(b) Sets up $y = 12 \times 2^x \implies 2^x = \frac{y}{12}$ **[M1]**

$x \log 2 = \log\left(\frac{y}{12}\right)$ **[M1]**

$f^{-1}(x) = \frac{\log(x/12)}{\log 2}$ **[A1]**

(c) $12 \times 2^x = 500 \implies 2^x = \frac{125}{3}$ (41.666...) **[M1]**

$x = \frac{\log(125/3)}{\log 2}$ **[M1]**

5.38 **[A1]**

[Back to Question] *[Total Marks: 9]*

730 **(a)** 24000
×
0.85
4
24000×0.85

4
(
=
12528.15
=12528.15) **[M1]**

12528
12528 **[A1]**

(b) $24000 \times 0.85^t = 9000$ or $0.85^t = 0.375$ **[M1]**

$t = \dfrac{\log 0.375}{\log 0.85}$ **[M1]**

6.04 **[A1]**

(c) $1.068^n = 2$ (or $1.068^n > 2$) **[M1]**

$n = \dfrac{\log 2}{\log 1.068} = 10.536\ldots$ **[M1]**

11 **[A1]**

(d) $2^y = \dfrac{x-3}{5}$ **[M1]**

$y = \log_2\left(\dfrac{x-3}{5}\right)$ or $y \log 2 = \log\left(\dfrac{x-3}{5}\right)$ **[M1]**

$log_2(\dfrac{x-3}{5}$ or
log
(
3
5
)
log
2
log2
log(
5
x3
)

[A1]

[Back to Question] *[Total Marks: 11]*

731 **(a)** Substitutes $(0, 7)$ to find A: $7 = A \times b^0 \implies A = 7$ **[B1]**

Substitutes $(3, 56)$ with $A = 7$: $7 \times b^3 = 56 \implies b^3 = 8$ **[M1]**

$b = 2$ **[A1]**

(b) Sets $y = 7 \times 2^x$ **[M1]**

Rearranges to make 2^x the subject: $2^x = \dfrac{y}{7}$ **[M1]**

Converts to logarithmic form: $\log_2\left(\dfrac{x}{7}\right)$ (or equivalent correct form) **[A1]**

(c) **(i)** $\dfrac{600}{7}$ **[B1]**

(ii) $x = \dfrac{\log(600/7)}{\log 2}$ **[M1]**

6.421 (from 6.4214...) **[A1]**

(d) **(i)** Sets $4 = \log_c 81 \implies c^4 = 81$ **[M1]**

$c = 3$ **[A1]**

(ii) $\dfrac{1}{9}$ (or 3^{-2}) **[B1]**

[Back to Question] *[Total Marks: 12]*

732 **(a)** x-coordinate 3 **[B1]**

y-coordinate 5 **[B1]**

(b) Base $AB = 4$ and height $= 3$ **[M1]**

12 **[A1]**

[Back to Question] *[Total Marks: 4]*

733 **(a)** $(1, 2)$ **[B1]**

(b) Identifies the base length $PQ = 10$ **[M1]**

Sets up the area equation:

$$\frac{1}{2} \times 10 \times h = 35$$

h = 7[$M1$]

$(0, 9)$ **[A1]**

[Back to Question] *[Total Marks: 4]*

734 **(a)** All three points A, B, and C plotted accurately on the grid **[B1]**

(b) Identifies the horizontal and vertical translation, e.g. $\vec{BA} = \begin{pmatrix} -3 \\ 4 \end{pmatrix}$ OR correctly plots D on the grid **[M1]**

$(3, 5)$ **[A1]**

(c) Calculates the midpoint of AC or BD: $\left(\dfrac{-1+6}{2}, \dfrac{3+1}{2}\right)$ **[M1]**

$(2.5, 2)$ **[A1]**

[Back to Question] *[Total Marks: 5]*

735 **(a)** $A = (-2, 3)$ **[B1]**

$C = (1, -2)$ **[B1]**

(b) Horizontal shift of 5 units or vertical shift of 5 units identified **[M1]**

$(-4, -2)$ **[A1]**

(c) Base $= 5$ and perpendicular height $= 5$ **[M1]**

25 **[A1]**

[Back to Question] *[Total Marks: 6]*

736 **(a)** All four points P, Q, R, and S plotted accurately **[B1]**

Points joined with straight line segments to form kite $PQRS$ **[B1]**

(b) $(1, 2)$ **[B1]**

(c) Diagonal lengths $PR = 9$ and $QS = 6$ **[M1]**

$\frac{1}{2} \times 9 \times 6 = 27$ **[A1]**

(d) $P' = (1, -5)$ **[B1]**

$Q' = (-4, 2)$ **[B1]**

[Back to Question] *[Total Marks: 7]*

737 **(a)** Identifies the area of the enclosing square as a^2 and the three outer right-angled triangles as $\frac{1}{2}ab$, $\frac{1}{2}ab$, and $\frac{1}{2}(a-b)^2$ **[M1]**

Sets up the subtraction for the area of OPQ: Area $= a^2 - \frac{1}{2}ab - \frac{1}{2}ab - \frac{1}{2}(a-b)^2$ **[M1]**

Correctly expands brackets and simplifies algebraically to $\frac{1}{2}(a^2 - b^2)$ without errors **[A1]**

(b) Sets up the distance squared equation: $(a-b)^2 + (b-a)^2 = (4\sqrt{2})^2$

$2(a-b)^2 = 32$ [M1]

Solves to find $a - b = 4$ (accepting $a > b$) **[A1]**

Uses the area equation:

$$\frac{1}{2}(a^2 - b^2) = 24$$

$a^2 - b^2 = 48$ [B1]

Factorises to $(a-b)(a+b) = 48$, substitutes $a - b = 4$ to obtain $a + b = 12$ **[M1]**

$a = 8$ and $b = 4$ **[A1]**

[Back to Question] *[Total Marks: 8]*

738 **(a)** Width $= x - (-2) = x + 2$ and Height $= y - (-4) = y + 4$ **[M1]**

$2(x+2) + 2(y+4) = 30 \implies 2x + 2y + 12 = 30 \implies x + y = 9$ **[A1]**

(b) Area $= (x+2)(y+4) = 54$ **[M1]**

Substitute $y = 9 - x$ to give $(x+2)(13-x) = 54$ **[M1]**

$x^2 - 11x + 28 = 0$ or equivalent three-term quadratic **[M1]**

$(4, 5)$ and $(7, 2)$ **[A1]**

(c) $(-2, 2)$ **[B1]**

$(7, -4)$ **[B1]**

[Back to Question] *[Total Marks: 8]*

739 **(a)** Sets the x-coordinate of the midpoint to zero: $\frac{3k + k + 4}{2} = 0$ **[M1]**

-1 **[A1]**

(b) Substitutes $k = -1$ to find $A(-3, 1)$ and $B(3, 5)$ **[M1]**

$AB^2 = (3 - (-3))^2 + (5 - 1)^2 = 6^2 + 4^2 = 52$ **[M1]**

$2\sqrt{13}$ **[A1]**

(c) Lets C be $(x, 0)$ and sets up equation $AC^2 = BC^2$:
$(x + 3)^2 + (0 - 1)^2 = (x - 3)^2 + (0 - 5)^2$ **[M1]**

Expands both sides correctly: $x^2 + 6x + 9 + 1 = x^2 - 6x + 9 + 25$ **[M1]**

Rearranges to solve for x: $12x = 24$

x = 2[$M1$]

$(2, 0)$ **[A1]**

[Back to Question] *[Total Marks: 9]*

740 **(a)** $(-6, 0)$ **[B1]**

(b) Identifies radius $OB = 6$ and angle from positive x-axis as $60°$ **[M1]**

$x = 6\cos 60° = 3$ and $y = 6\sin 60°$ $(= 3\sqrt{3})$ **[M1]**

$(3, 5.20)$ **[A1]**

(c) Calculates length $AD = 12$, and finds $AP = \frac{1}{4} \times 12 = 3$ **[M1]**

$(3, 0)$ **[A1]**

[Back to Question] *[Total Marks: 6]*

741 **(a)** $A(0, 6)$ **[B1]**

$B(4, 0)$ **[B1]**

(b) $(4, 6)$ **[B1]**

(c) $2y + 3(2) = 12$ **[M1]**

$(2, 3)$ **[A1]**

(d) Base $OB = 4$ and height of $\triangle OPB = 3$ (the y-coordinate of P) **[M1]**

6 **[A1]**

[Back to Question] *[Total Marks: 7]*

742 **(a)** Substitutes $y = 0$ to find $A(6, 0)$ **[B1]**

Substitutes $x = 0$ to find $B(0, 9)$ **[B1]**

(b) Recognises the change in x is -6 and change in y is 9 from A to B **[M1]**

Forms a valid ratio equation for either coordinate, e.g. x-coordinate $= 6 + \frac{2}{3}(-6)$ OR y-coordinate $= 0 + \frac{2}{3}(9)$ **[M1]**

$(2, 6)$ **[A1]**

(c) Substitutes $x = 4$ and $y = 3$ into the equation: $3(4) + 2(3)$ **[M1]**

Shows $12 + 6 = 18$ and clearly concludes 'Yes' **[A1]**

[Back to Question] *[Total Marks: 7]*

743 **(a)** $h = -3$ **[B1]**

Second quadrant (accept 2nd or II) **[B1]**

(b) Width $AB = 5 - (-3) = 8$ **[M1]**

$8(k + 1) = 52$ or $k + 1 = 6.5$ **[M1]**

5.5 **[A1]**

(c) $x = \frac{-3 + 5}{2} = 1$ or $y = \frac{-1 + 5.5}{2} = 2.25$ **[M1]**

$(1, 2.25)$ **[A1]**

(d) $p = 0$ **[B1]**

$\frac{1}{2} \times 8 \times (q + 1) = 18$ or $q + 1 = 4.5$ **[M1]**

$(0, 3.5)$ **[A1]**

[Back to Question] *[Total Marks: 10]*

744 $\frac{5 - (-3)}{6 - 2}$ or equivalent **[M1]**

2 **[A1]**

[Back to Question] *[Total Marks: 2]*

745 **(a)** Evidence of using $\frac{\text{change in } y}{\text{change in } x}$ with two correct points from the graph, e.g., $\frac{4 - (-2)}{3 - 0}$ **[M1]**

2 **[A1]**

(b) $(0, -2)$ **[B1]**

[Back to Question] *[Total Marks: 3]*

746 **(a)** Uses gradient formula: $\frac{-4 - 8}{4 - (-2)}$ or $\frac{8 - (-4)}{-2 - 4}$ **[M1]**

-2 **[A1]**

(b) Identifies coordinates of two clear points on the line, e.g., $(0, 1)$ and $(4, 4)$ **[M1]**

$\frac{3}{4}$ or 0.75 **[A1]**

[Back to Question] *[Total Marks: 4]*

747 **(a)** Uses the gradient formula: $\frac{-5-7}{2-(-4)}$ **[M1]**

-2 **[A1]**

(b) Sets up the equation: $\frac{c-7}{0-(-4)} = -2$ **[M1]**

-1 **[A1]**

[Back to Question] *[Total Marks: 4]*

748 **(a)** Correct vertical and horizontal change seen, e.g. $\frac{6}{4}$ or $\frac{3}{2}$ **[M1]**

1.5 (or $\frac{3}{2}$) **[A1]**

(b) Correct vertical and horizontal change seen, e.g. $\frac{-3}{6}$ **[M1]**

-0.5 (or $-\frac{1}{2}$) **[A1]**

(c) $\frac{-9-7}{5-(-3)}$ **[M1]**

-2 **[A1]**

[Back to Question] *[Total Marks: 6]*

749 **(a)** $\frac{k-5}{4-(-2)} = -\frac{1}{2}$ **[M1]**

$k - 5 = -3$ **[M1]**

2 **[A1]**

(b) Gradient of $BC = \frac{-1-2}{10-4} = -\frac{1}{2}$ **[M1]**

Gradient of AB = gradient of BC and point B is common to both segments **[A1]**

(c) $\frac{(3p-1)-5}{p-(-2)} = 2$ **[M1]**

$3p - 6 = 2(p + 2)$ **[M1]**

10 **[A1]**

[Back to Question] *[Total Marks: 8]*

750 **(a)** $\dfrac{(a+4)-3a}{5a-2a}$ or equivalent **[M1]**

$\dfrac{4-2a}{3a}$ **[A1]**

(b) States or implies gradient of PQ is $-\dfrac{1}{3}$ **[B1]**

$\dfrac{4-2a}{3a} = -\dfrac{1}{3}$ or $3(4-2a) = -3a$ **[M1]**

4 **[A1]**

(c) Gradient $AB = \dfrac{8-2}{3-(-1)}$ or $\dfrac{6}{4}$ or $\dfrac{3}{2}$ **[M1]**

$\dfrac{23-8}{c-3} = \dfrac{3}{2}$ or $\dfrac{23-2}{c-(-1)} = \dfrac{3}{2}$ **[M1]**

13 **[A1]**

[Back to Question] *[Total Marks: 8]*

751 **(a)** $\dfrac{(a+6)-3a}{4-(-2)}$ **[M1]**

$\dfrac{6-2a}{6}$ **[M1]**

$\dfrac{3-a}{3}$ or $1-\dfrac{a}{3}$ **[A1]**

(b) $\dfrac{3-a}{3} = 3$ oe **[M1]**

$a = -6$ **[A1]**

(c) $P = (-2,-18)$ and $Q = (4,0)$ **[M1]**

$\left(\dfrac{-2+4}{2}, \dfrac{-18+0}{2}\right)$ **[M1]**

$(1,-9)$ **[A1]**

[Back to Question] *[Total Marks: 8]*

752 **(a)** $\dfrac{k-2}{4}$ **[B1]**

(b) Finds an expression for the gradient of BC (e.g., $\dfrac{k}{k-5}$) and equates to the gradient of AB: $\dfrac{k-2}{4} = \dfrac{k}{k-5}$ **[M1]**

Expands brackets correctly: $(k-2)(k-5) = 4k$

$\mathrm{k}^2 - 7k + 10 = 4k$ [M1]

Rearranges to the given equation $k^2 - 11k + 10 = 0$ with no mathematical errors seen **[A1]**

(c) Factorises $(k-1)(k-10) = 0$ or uses the quadratic formula **[M1]**

1 and 10 **[A1]**

(d) Deduces that $k = 10$ must be used, as $k = 1$ would mean points A and C are the same,

contradicting that they are distinct **[B1]**

Calculates the gradient of OB using $(5, 10)$: $\dfrac{10-0}{5-0} = 2$ **[B1]**

[Back to Question] *[Total Marks: 8]*

753 **(a)** Calculates the gradient of AB as $\dfrac{9-3}{4-1} = 2$ and equates to gradient of BC or AC, e.g., $\dfrac{y-9}{8-4} = 2$ **[M1]**

$y = 17$ **[A1]**

(b) Sets up the gradient equation: $\dfrac{(p^2-p-3)-3}{p-1} = 2$ **[M1]**

Simplifies to form a 3-term quadratic equation: $p^2 - p - 6 = 2(p-1)$

$\mathrm{p}^2 - 3p - 4 = 0$ [M1]

Factorises $(p-4)(p+1) = 0$ or uses the quadratic formula correctly **[M1]**

$p = 4$ and $p = -1$ **[A1]**

(c) Sets up the gradient equation: $\dfrac{12-m}{(m+1)-3} = m$ **[M1]**

Multiplies by the denominator: $12 - m = m(m-2)$ **[M1]**

Rearranges into a 3-term quadratic: $m^2 - m - 12 = 0$ **[M1]**

$m = 4$ and $m = -3$ **[A1]**

[Back to Question] *[Total Marks: 10]*

754 **(a)** $\dfrac{5-1}{4-(-2)}$ or equivalent **[M1]**

$\dfrac{2}{3}$ **[A1]**

(b) $\dfrac{-2-k}{8-3} = -1.4$ or equivalent **[M1]**

$-2 - k = -7$ **[M1]**

5 **[A1]**

[Back to Question] *[Total Marks: 5]*

755 **(a)** $\dfrac{-9-5}{4-(-3)}$ or $\dfrac{5-(-9)}{-3-4}$ **[M1]**

-2 **[A1]**

(b) $\dfrac{1-5}{k-(-3)} = -2$ oe **[M1]**

$-4 = -2(k+3)$ or $-2k - 6 = -4$ **[M1]**

$k = -1$ **[A1]**

[Back to Question] *[Total Marks: 5]*

756 **(a)** Forms a correct equation for the gradient: $\frac{5-2k}{2-(k-3)} = -3$ OR $\frac{2k-5}{k-3-2} = -3$ **[M1]**

Correctly simplifies the denominator and equates: $5-2k = -3(5-k)$ **[M1]**

$k = 4$ **[A1]**

(b) Evaluates coordinates of P as $(1, 8)$ and substitutes into midpoint formula: $\left(\frac{1+2}{2}, \frac{8+5}{2}\right)$ **[M1]**

$(1.5, 6.5)$ **[A1]**

[Back to Question] *[Total Marks: 5]*

757 **(a)** Finds a correct algebraic expression for the gradient: $\frac{8-4}{k-2-3k}$ or $\frac{4}{-2k-2}$ **[M1]**

Equates to $-\frac{1}{2}$ and eliminates fractions: $8 = 2k+2$ or $-8 = -2k-2$ **[M1]**

3 **[A1]**

(b) $P(9, 4)$ and $Q(1, 8)$ **[B1]**

(c) Identifies the coordinates of R must be of the form $(0, y)$ **[B1]**

Sets up the gradient equation: $\frac{y-4}{0-9} = 1$ **[M1]**

$(0, -5)$ **[A1]**

[Back to Question] *[Total Marks: 7]*

758 **(a)** Gradient $= \frac{(t^2-3t+2)-0}{t-1}$ **[M1]**

Factorising numerator: $\frac{(t-1)(t-2)}{t-1}$ **[M1]**

Cancelling $(t-1)$ correctly to obtain $t-2$ **[A1]**

(b) $t-2 = 2.8$ giving $t = 4.8$ **[M1]**

$y = 4.8^2 - 3(4.8) + 2$ **[M1]**

$(4.8, 10.64)$ **[A1]**

(c) Gradient of $QR = \frac{(t^2-3t+2)-(-1.5)}{t-3.2}$ **[M1]**

$\frac{t^2-3t+3.5}{t-3.2} = t-2$ **[M1]**

$t^2-3t+3.5 = t^2-5.2t+6.4$ leading to $2.2t = 2.9$ **[M1]**

1.32 **[A1]**

[Back to Question] *[Total Marks: 10]*

759 **(a)** $\left(\frac{-2+4}{2}, \frac{5+(-3)}{2}\right)$ **[M1]**

$(1, 1)$ **[A1]**

(b) $(4-(-2))^2+(-3-5)^2$ or $6^2+(-8)^2$ **[M1]**

$\sqrt{36+64}$ or $\sqrt{100}$ **[M1]**

10 **[A1]**

[Back to Question] *[Total Marks: 5]*

760 **(a)** $\left(\frac{1+7}{2}, \frac{4+(-4)}{2}\right)$ **[M1]**

$(4, 0)$ **[A1]**

(b) $(7-1)^2+(-4-4)^2$ oe **[M1]**

$\sqrt{36+64}$ or $\sqrt{100}$ **[M1]**

10 **[A1]**

[Back to Question] *[Total Marks: 5]*

761 **(a)** Uses the midpoint formula: $\left(\frac{2+8}{2}, \frac{-3+5}{2}\right)$ **[M1]**

$(5, 1)$ **[A1]**

(b) Uses the distance formula: $\sqrt{(8-2)^2+(5-(-3))^2}$ **[M1]**

$\sqrt{6^2+8^2}$ or $\sqrt{36+64}$ **[M1]**

10 **[A1]**

[Back to Question] *[Total Marks: 5]*

762 **(a)** $A = (-2, 1)$ **[B1]**

$B = (4, 9)$ **[B1]**

(b) $\left(\frac{-2+4}{2}, \frac{1+9}{2}\right)$ **[M1]**

$(1, 5)$ **[A1]**

(c) $\sqrt{(4-(-2))^2+(9-1)^2}$ or $\sqrt{6^2+8^2}$ **[M1]**

10 **[A1]**

[Back to Question] *[Total Marks: 6]*

763 **(a)** Calculates length OP: $\sqrt{(-3)^2+4^2}$ or $\sqrt{9+16}$ **[M1]**

5 **[A1]**

(b) Recognises $O(0, 0)$ is the midpoint of PQ, so sets up $\frac{-3+x}{2}=0$ and $\frac{4+y}{2}=0$ OR uses translation from P to O **[M1]**

$(3, -4)$ **[A1]**

(c) Sets up equation for length OR: $k^2 + (\sqrt{11})^2 = 5^2$ **[M1]**

$k^2 + 11 = 25$

$k^2 = 14$ [M1]

$-\sqrt{14}$ **[A1]**

[Back to Question] *[Total Marks: 7]*

764 **(a)** $\frac{-2+x}{2} = 3$ or $\frac{4+y}{2} = -1$ **[M1]**

$(8, -6)$ **[A1]**

(b) $PQ = \sqrt{(8-(-2))^2 + (-6-4)^2}$ **[M1]**

$\sqrt{100+100}$ or $\sqrt{200}$ **[M1]**

$10\sqrt{2}$ **[A1]**

(c) Let $R = (0, y)$ and form $(0-(-2))^2 + (y-4)^2 = (0-8)^2 + (y-(-6))^2$ **[M1]**

$4 + y^2 - 8y + 16 = 64 + y^2 + 12y + 36$ **[M1]**

$-20y = 80$ giving $y = -4$ **[A1]**

$(0, -4)$ **[A1]**

[Back to Question] *[Total Marks: 9]*

765 **(a)** Finds coordinates of the midpoint in terms of p: $\left(\frac{4p}{2}, \frac{3p-1}{2}\right)$ or $\left(2p, \frac{3p-1}{2}\right)$ **[M1]**

Substitutes into the equation of the line: $\frac{3p-1}{2} = 2p + 1$ **[M1]**

-3 **[A1]**

(b) Sets up the distance squared equation: $(3p-p)^2 + (p-2-(2p+1))^2 = 6^2$ **[M1]**

Simplifies to $(2p)^2 + (-p-3)^2 = 36$ **[M1]**

Expands to $4p^2 + p^2 + 6p + 9 = 36$ and rearranges to $5p^2 + 6p - 27 = 0$ with no errors seen **[A1]**

(c) Factorises to $(5p-9)(p+3) = 0$ or uses correct quadratic formula **[M2]**

Award [M1] for a partial factorisation or one error in the quadratic formula

1.8 (or $\frac{9}{5}$) and -3 **[A1]**

[Back to Question] *[Total Marks: 9]*

766 **(a)** $\left(\frac{-1+5}{2}, \frac{2+8}{2}\right)$ **[M1]**

$(2, 5)$ **[A1]**

(b) $(2-(-1))^2 + (5-2)^2$ or $\frac{1}{2}\sqrt{(5-(-1))^2 + (8-2)^2}$ oe **[M1]**

$\sqrt{9+9}$ or $\sqrt{18}$ **[M1]**

$3\sqrt{2}$ **[A1]**

(c) $\frac{2+x}{2} = \frac{5}{2}$ and $\frac{5+y}{2} = \frac{3}{2}$ oe **[M1]**

$x = 3$ and $y = -2$ to give $D(3, -2)$ **[A1]**

Length $AD = \sqrt{(3-(-1))^2 + (-2-2)^2}$ oe **[M1]**

$\sqrt{16+16}$ or $\sqrt{32}$ **[M1]**

$4\sqrt{2}$ **[A1]**

[Back to Question] *[Total Marks: 10]*

767 **(a)** Uses distance formula: $\sqrt{(4-(-2))^2 + (-3-5)^2}$ **[M1]**

Simplifies to $\sqrt{6^2 + (-8)^2}$ or $\sqrt{36+64}$ **[M1]**

10 **[A1]**

(b) Uses midpoint formula: $\left(\frac{-2+4}{2}, \frac{5+(-3)}{2}\right)$ **[M1]**

$(1, 1)$ **[A1]**

[Back to Question] *[Total Marks: 5]*

768 **(a)** $(7-3)^2 + (5-k)^2 = 5^2$ or equivalent **[M1]**

$16 + (5-k)^2 = 25$ or $(5-k)^2 = 9$ **[M1]**

$k = 2$ and $k = 8$ **[A1]**

(b) Identifies $P(3, 2)$ to use in midpoint calculation: $\left(\frac{3+7}{2}, \frac{2+5}{2}\right)$ **[M1]**

$(5, 3.5)$ **[A1]**

[Back to Question] *[Total Marks: 5]*

769 **(a)** Sets up an equation for at least one coordinate, e.g., $\frac{-3+x}{2} = 1$ or $\frac{5+y}{2} = -2$ **[M1]**

$(5, -9)$ **[A1]**

(b) Uses distance formula with $P(-3, 5)$ and their $Q(5, -9)$: $\sqrt{(5-(-3))^2 + (-9-5)^2}$ **[M1]**

$\sqrt{8^2 + (-14)^2}$ or $\sqrt{64+196}$ or $\sqrt{260}$ **[M1]**

16.1 **[A1]**

[Back to Question] *[Total Marks: 5]*

770 **(a)** $(k-(-3))^2 + (-1-5)^2 = 10^2$ oe **[M1]**

$(k+3)^2 + 36 = 100$ or $k^2 + 6k + 9 + 36 = 100$ **[M1]**

$(k+3)^2 = 64$ or $k^2 + 6k - 55 = 0$ **[M1]**

$k = 5$ or $k = -11$ **[A1]**

(b) Chooses $k = 5$ to establish coordinates of Q as $(5, -1)$ **[M1]**

$(1, 2)$ **[A1]**

[Back to Question] *[Total Marks: 6]*

771 **(a)** $\left(\frac{-3+5}{2}, \frac{2+8}{2}\right)$ **[M1]**

$(1, 5)$ **[A1]**

(b) Radius $r = \sqrt{(1-(-3))^2 + (5-2)^2}$ or $\sqrt{4^2+3^2} = 5$ **[M1]**

Area $= \pi \times 5^2$ **[M1]**

78.5 **[A1]**

(c) Distance $CD = 5 \implies (k-1)^2 + (8-5)^2 = 5^2$ **[M1]**

$(k-1)^2 + 9 = 25 \implies (k-1)^2 = 16$ **[M1]**

$k = 5$ and $k = -3$ **[A1]**

[Back to Question] *[Total Marks: 8]*

772 **(a)** Sets up the distance formula: $\sqrt{(3p-2p)^2 + (4p-p)^2} = 5\sqrt{10}$ **[M1]**

Simplifies the terms inside the root: $\sqrt{p^2 + (3p)^2} = 5\sqrt{10}$ **[M1]**

Squares both sides and simplifies: $10p^2 = 250$ **[M1]**

$p = 5$ (rejects -5 as $p > 0$) **[A1]**

(b) Evaluates coordinates of $A(10, 5)$ and $B(15, 20)$ **[B1]**

Sets up a valid expression for the coordinates of C, e.g., x-coordinate $= 10 + \frac{2}{5}(15-10)$ or y-coordinate $= 5 + \frac{2}{5}(20-5)$ **[M1]**

$(12, 11)$ **[A1]**

(c) Uses the midpoint formula, e.g. $\frac{15+x}{2} = 12$ and $\frac{20+y}{2} = 11$, or $D = 2C - B$ **[M1]**

$(9, 2)$ **[A1]**

[Back to Question] *[Total Marks: 9]*

773 **(a)** $AB = \sqrt{(5-(-3))^2 + (6-2)^2} = \sqrt{80}$ $(= 8.94\ldots)$ **[M1]**

$BC = \sqrt{(9-5)^2 + (-2-6)^2} = \sqrt{80}$ $(= 8.94\ldots)$ **[M1]**

$AC = \sqrt{(9-(-3))^2 + (-2-2)^2} = \sqrt{160}$ $(= 12.6\ldots)$ **[M1]**

$AB = BC$ (and $ACeAB$), therefore triangle ABC is isosceles **[A1]**

(b) **(i)** $\left(\frac{-3+9}{2}, \frac{2+(-2)}{2}\right)$ **[M1]**

$(3, 0)$ **[A1]**

(ii) $BM = \sqrt{(5-3)^2 + (6-0)^2}$ $(= \sqrt{40})$ **[M1]**

6.32 **[A1]**

(c) $\frac{1}{2} \times \sqrt{160} \times \sqrt{40}$ or $\frac{1}{2} \times \sqrt{80} \times \sqrt{80}$ **[M1]**

40 **[A1]**

(d) $\frac{5+x}{2} = 3$ and $\frac{6+y}{2} = 0$ (or vector translation $B + \overrightarrow{AD}$) **[M1]**

$(1, -6)$ **[A1]**

[Back to Question] *[Total Marks: 12]*

774 Rearranging to $5y = -3x + 20$ or $y = -\frac{3}{5}x + 4$ **[M1]**

Gradient $= -\frac{3}{5}$ oe **[A1]**

y-intercept $= 4$ **[A1]**

[Back to Question] *[Total Marks: 3]*

775 **(a)** Gradient $= -2$ **[B1]**

y-intercept $= 4$ **[B1]**

$-2x + 4$ **[B1]**

(b) $2x + y = 4$ oe **[B1]**

[Back to Question] *[Total Marks: 4]*

776 **(a)** Rearranges equation to $y = mx + c$ form: $5y = -3x + 15$ or $y = -\frac{3}{5}x + 3$ **[M1]**

Gradient $= -\frac{3}{5}$ **[A1]**

y-intercept $= 3$ **[A1]**

(b) Identifies parallel gradient as $-\frac{3}{5}$ **[M1]**

$-\frac{3}{5}x - 4$ **[A1]**

[Back to Question] *[Total Marks: 5]*

777 **(a)** Rearranges the equation to make y the subject: $4y = 3x - 12$ or $y = \frac{3}{4}x - 3$ **[M1]**

Gradient $= \frac{3}{4}$ **[A1]**

y-intercept $= -3$ **[A1]**

(b) Identifies the gradient of M is $\frac{3}{4}$ **[M1]**

$\frac{3}{4}x + 5$ **[A1]**

[Back to Question] *[Total Marks: 5]*

778 **(a)** Gradient $= \dfrac{3-(-3)}{3-0} = 2$ **[M1]**

y-intercept $= -3$ **[B1]**

$2x - 3$ **[A1]**

(b) $x = -2$ **[B1]**

(c) $y = 2(-2) - 3 = -7$ **[M1]**

$(-2, -7)$ **[A1]**

[Back to Question] *[Total Marks: 6]*

779 **(a)** $5y = -4x + 20$ or $y = -\dfrac{4}{5}x + 4$ **[M1]**

Gradient $= -\dfrac{4}{5}$ (or -0.8) **[A1]**

y-intercept $= 4$ **[A1]**

(b) Substitute $y = 0$: $4x = 20$ **[M1]**

$(5, 0)$ **[A1]**

(c) Gradient $= \dfrac{7-(-5)}{3-(-3)} = 2$ **[M1]**

$y = 2x + 1$ **[M1]**

$2x - y = -1$ (or $-2x + y = 1$) **[A1]**

[Back to Question] *[Total Marks: 8]*

780 **(a)** $\dfrac{(2p-1)-(-1)}{5-p} = 3$ **[M1]**

$2p = 3(5 - p)$ **[M1]**

$p = 3$ **[A1]**

(b) Substituting $p = 3$ to find coordinates $A(3, -1)$ or $B(5, 5)$ and using gradient 3 to find equation **[M1]**

$y = 3x - 10$ **[A1]**

(c) Identifies x-intercept as $\dfrac{10}{3}$ or y-intercept as -10 **[M1]**

$\dfrac{1}{2} \times \dfrac{10}{3} \times 10$ **[M1]**

$\dfrac{50}{3}$ or 16.7 $(16.66\ldots)$ **[A1]**

[Back to Question] *[Total Marks: 8]*

781 **(a)** Sets up the gradient expression: $\frac{k-3k}{4-(-2)}$ **[M1]**

$-\frac{k}{3}$ **[A1]**

(b) Substitutes a point and gradient into the linear equation: $y-k=-\frac{k}{3}(x-4)$ or $y-3k=-\frac{k}{3}(x-(-2))$ **[M1]**

Multiplies by 3 to eliminate the fraction: $3y-3k=-kx+4k$ **[M1]**

$kx+3y=7k$ **[A1]**

(c) Substitutes $x=1$ and $y=10$ into their equation from part (b): $k(1)+3(10)=7k$ **[M1]**

5 **[A1]**

(d) Substitutes $k=5$ to find the line equation $5x+3y=35$ and sets $y=0$: $5x=35$ **[M1]**

$(7,0)$ **[A1]**

[Back to Question] *[Total Marks: 9]*

782 **(a)** $\frac{(3a+2)-(a+4)}{(a-1)-2a}=-4$ **[M1]**

$\frac{2a-2}{-a-1}=-4$ **[M1]**

$2a-2=4a+4$

-6 = 2a

a = -3[A1]

(b) Identifies coordinates of A as $(-6,1)$ or B as $(-4,-7)$ **[M1]**

$y-1=-4(x+6)$ or $y=-4x-23$ **[M1]**

$4x+y=-23$ **[A1]**

(c) $4x+(3x-2)=-23$ or $-4x-23=3x-2$ oe **[M1]**

$7x=-21$

x = -3[A1]

$y=-11$ **[A1]**

[Back to Question] *[Total Marks: 9]*

783 **(a)** Calculates the midpoint of $AB=\left(\frac{-3+5}{2},\frac{1+5}{2}\right)=(1,3)$ **[B1]**

Calculates the gradient of $AB=\frac{5-1}{5-(-3)}=\frac{4}{8}=\frac{1}{2}$ **[B1]**

Finds the perpendicular gradient $=-2$ **[M1]**

Uses perpendicular gradient -2 and midpoint $(1,3)$ to find equation, e.g., $3=-2(1)+c$ **[M1]**

$-2x+5$ **[A1]**

(b) Sets up simultaneous equations, e.g., substituting $y = -2x + 5$ into $x - 2y = 5$ **[M1]**

$x - 2(-2x + 5) = 5 \implies 5x - 10 = 5 \implies x = 3$ **[M1]**

$(3, -1)$ **[A1]**

(c) Uses distance formula for $A(-3, 1)$ and $P(3, -1)$ **[M1]**

$\sqrt{(3 - (-3))^2 + (-1 - 1)^2}$ **[M1]**

$\sqrt{40}$ or $2\sqrt{10}$ **[A1]**

[Back to Question] *[Total Marks: 10]*

784 **(a)** Evidence of using two integer coordinates on the line, e.g. $\dfrac{5 - (-1)}{-4 - 4}$ **[M1]**

$-\dfrac{3}{4}$ or -0.75 **[A1]**

(b) $y = -\dfrac{3}{4}x + 2$ or $y - 5 = -\dfrac{3}{4}(x + 4)$ **[M1]**

$4y = -3x + 8$ or $4y - 20 = -3x - 12$ **[M1]**

$3x + 4y = 8$ or equivalent integer form **[A1]**

[Back to Question] *[Total Marks: 5]*

785 **(a)** Gradient $= \dfrac{-2 - 3}{6 - (-4)}$ or $\dfrac{-5}{10} = -\dfrac{1}{2}$ **[M1]**

Substitution of $(-4, 3)$ or $(6, -2)$ into $y = -\dfrac{1}{2}x + c$, e.g. $3 = -\dfrac{1}{2}(-4) + c$ **[M1]**

$-\dfrac{1}{2}x + 1$ **[A1]**

(b) $C = (5, 0)$ **[B1]**

$D = (0, -3)$ **[B1]**

(c) $x = -4$ **[B1]**

[Back to Question] *[Total Marks: 6]*

786 **(a)** Identifies two correct points on the line, e.g., $(0, 4)$ and $(2, 7)$ **[M1]**

Calculates the gradient $m = \dfrac{7 - 4}{2 - 0} = \dfrac{3}{2}$ **[M1]**

States correct equation in required integer form, e.g., $3x - 2y = -8$ or $2y - 3x = 8$ **[A1]**

(b) Uses $m_1 \times m_2 = -1$ to find gradient of M: $-\dfrac{2}{3}$ **[M1]**

Substitutes $(9, -1)$ into $y = -\dfrac{2}{3}x + c$ to find c: $-1 = -\dfrac{2}{3}(9) + c$ **[M1]**

$-\dfrac{2}{3}x + 5$ **[A1]**

[Back to Question] *[Total Marks: 6]*

787 **(a)** Identifies the y-intercept is 4 or uses coordinates $(0, 4)$ **[B1]**

Calculates the gradient: $\frac{0-4}{6-0}$ or $\frac{-4}{6}$ **[M1]**

$-\frac{2}{3}x + 4$ **[A1]**

(b) Finds the coordinates of the midpoint of the segment $(0, 4)$ to $(6, 0)$: $\left(\frac{0+6}{2}, \frac{4+0}{2}\right)$ or $(3, 2)$ **[B1]**

Calculates the gradient of L_2: $\frac{7-2}{2-3}$ or $\frac{5}{-1} = -5$ **[M1]**

Uses point-gradient form for L_2: $y - 7 = -5(x - 2)$ or $y - 2 = -5(x - 3)$ **[M1]**

$5x + y = 17$ **[A1]**

[Back to Question] *[Total Marks: 7]*

788 **(a)** $A = (8, 0)$ **[B1]**

$B = (0, 12)$ **[B1]**

$C = (2, 0)$ **[B1]**

(b) $3x + 2(1.5x - 3) = 24$ **[M1]**

$6x = 30$ giving $x = 5$ **[A1]**

$(5, 4.5)$ **[A1]**

(c) Area of $\triangle OAB = \frac{1}{2} \times 8 \times 12 = 48$ **[M1]**

Area of $\triangle CAP = \frac{1}{2} \times (8 - 2) \times 4.5 = 13.5$ **[M1]**

34.5 **[A1]**

(d) $5k - 4(4.5) = 14$ **[M1]**

6.4 **[A1]**

[Back to Question] *[Total Marks: 11]*

789 Identifies the gradient $m = -2$ and sets up $4 = -2(3) + c$ or $y - 4 = -2(x - 3)$ **[M1]**

$-2x + 10$ **[A1]**

[Back to Question] *[Total Marks: 2]*

790 Rearranges equation to find gradient, e.g., $2y = -4x + 9$ implies $m = -2$ **[M1]**

Substitutes $(3, -1)$ into $y = -2x + c$ or writes $y - (-1) = -2(x - 3)$ **[M1]**

$-2x + 5$ **[A1]**

[Back to Question] *[Total Marks: 3]*

791 Gradient $m = -3$ **[B1]**

$5 = -3(-2) + c$ or $y - 5 = -3(x - (-2))$ **[M1]**

$y = -3x - 1$ **[A1]**

[Back to Question] *[Total Marks: 3]*

792 **(a)** 2 **[B1]**

(b) $1 = 2(2) + c$ or $y - 1 = 2(x - 2)$ **[M1]**

$2x - 3$ **[A1]**

(c) $5 = 2(-1) + c$ or $y - 5 = 2(x + 1)$ **[M1]**

$2x + 7$ **[A1]**

[Back to Question] *[Total Marks: 5]*

793 **(a)** Rearranges the equation to make y the subject: $2y = 5x + 8$ or $y = \frac{5}{2}x + 4$ **[M1]**

$\frac{5}{2}$ or 2.5 **[A1]**

(b) Identifies the gradient of L_2 is $\frac{5}{2}$ and the y-intercept is -3 **[M1]**

$\frac{5}{2}x - 3$ (or $2.5x - 3$) **[A1]**

(c) Substitutes $(4, 1)$ and $m = \frac{5}{2}$ into linear equation: $y - 1 = \frac{5}{2}(x - 4)$ or $1 = \frac{5}{2}(4) + c$ **[M1]**

Eliminates the fraction: $2y - 2 = 5(x - 4)$ or $2y - 2 = 5x - 20$ **[M1]**

$5x - 2y = 18$ (or $-5x + 2y = -18$) **[A1]**

[Back to Question] *[Total Marks: 7]*

794 **(a)** $\frac{-1 - 5}{5 - (-3)}$ or $\frac{5 - (-1)}{-3 - 5}$ **[M1]**

$-\frac{3}{4}$ **[A1]**

(b) $m = -\frac{3}{4}$ **[M1]**

$-2 = -\frac{3}{4}(4) + c$ or $y - (-2) = -\frac{3}{4}(x - 4)$ **[M1]**

$y = -\frac{3}{4}x + 1$ **[A1]**

[Back to Question] *[Total Marks: 5]*

795 **(a)** $4y = -3x + 12$ or $y = -\frac{3}{4}x + 3$ **[M1]**

$-\frac{3}{4}$ (or -0.75) **[A1]**

(b) Gradient of $L_2 = -\frac{3}{4}$ **[B1]**

$y - 3 = -\frac{3}{4}(x - 4)$ or $3(4) + 4(3) = c$ **[M1]**

$3x + 4y = 24$ **[A1]**

(c) $A = (8, 0)$ **[B1]**

$B = (0, 6)$ **[B1]**

$(4, 3)$ **[A1]**

[Back to Question] *[Total Marks: 8]*

796 **(a)** Finds the gradient of line P in terms of k: $m_P = \frac{4}{k}$ **[B1]**

Sets gradients equal: $\frac{4}{k} = k - 3$ **[M1]**

Forms a 3-term quadratic equation: $k^2 - 3k - 4 = 0$ **[M1]**

4 and -1 **[A1]**

(b) **(i)** Identifies that $k = 4$ produces a positive gradient ($m = 1$) while $k = -1$ produces a negative gradient ($m = -4$) **[M1]**

$x - 2$ **[A1]**

(ii) Substitutes $(-5, 7)$ and $m = 1$ into equation: $y - 7 = 1(x - (-5))$ **[M1]**

$x - y = -12$ or $y - x = 12$ (or equivalent integer form) **[A1]**

[Back to Question] *[Total Marks: 8]*

797 Rearranges $5x - 2y = 15$ to find the gradient, e.g. $2y = 5x - 15$

y $= \frac{5}{2}x - 7.5$, so $m = \frac{5}{2}$ [M1]

Substitutes $(-4, 1)$ and $m = \frac{5}{2}$ into $y = mx + c$: $1 = \frac{5}{2}(-4) + c$ **[M1]**

$\frac{5}{2}x + 11$ or $2.5x + 11$ **[A1]**

[Back to Question] *[Total Marks: 3]*

798 **(a)** $\frac{6 - 2}{5 - (-3)}$ oe **[M1]**

$\frac{1}{2}$ or 0.5 **[A1]**

(b) $y - (-1) = \frac{1}{2}(x - 4)$ oe **[M1]**

$2(y + 1) = x - 4$ or $2y + 2 = x - 4$ **[M1]**

$x - 2y = 6$ or $-x + 2y = -6$ **[A1]**

[Back to Question] *[Total Marks: 5]*

799 **(a)** Uses the gradient formula: $\frac{4-1}{4-(-2)}$ **[M1]**

$\frac{3}{6} = \frac{1}{2}$ or 0.5 **[A1]**

(b) States or implies gradient $m = 0.5$ **[B1]**

Substitutes $(2, -3)$ into $y = 0.5x + c$: $-3 = 0.5(2) + c$ **[M1]**

$0.5x - 4$ (or $\frac{1}{2}x - 4$) **[A1]**

(c) Finds coordinates of P by setting $y = 0$: $0 = 0.5x - 4 \implies x = 8$, so $P(8, 0)$ **[B1]**

Finds coordinates of Q by setting $x = 0$: $y = -4$, so $Q(0, -4)$ **[B1]**

Applies distance formula: $\sqrt{(8-0)^2 + (0-(-4))^2}$ or $\sqrt{8^2 + (-4)^2}$ **[M1]**

8.94 (from $\sqrt{80}$) **[A1]**

[Back to Question] *[Total Marks: 9]*

800 **(a)** Calculates the gradient of L_1: $m = \frac{-3-5}{3-(-1)}$ **[M1]**

$m = \frac{-8}{4} = -2$ **[A1]**

Uses parallel gradient -2 and point $C(2, 7)$ to find c: $7 = -2(2) + c$ **[M1]**

$-2x + 11$ **[A1]**

(b) Finds y-intercept $E(0, 11)$ and sets $y = 0$ to find x-intercept D: $0 = -2x + 11$

x = 5.5[$M1$]

Sets up area calculation for right-angled triangle: $\frac{1}{2} \times 5.5 \times 11$ **[M1]**

30.25 or $\frac{121}{4}$ **[A1]**

[Back to Question] *[Total Marks: 7]*

801 Gradient of $L_1 = \frac{3}{4}$ oe **[B1]**

Equation of L_2 is $y = \frac{3}{4}x + c$ or $3x - 4y = k$ **[M1]**

x-intercept of L_2 is $-\frac{4c}{3}$ (or $\frac{k}{3}$) AND y-intercept is c (or $-\frac{k}{4}$) oe **[M1]**

$(-\frac{4c}{3})^2 + c^2 = 15^2$ or $(\frac{k}{3})^2 + (-\frac{k}{4})^2 = 15^2$ oe **[M1]**

$\frac{25c^2}{9} = 225$ or $\frac{25k^2}{144} = 225$ **[M1]**

$c = \pm 9$ or $k = \pm 36$ **[A1]**

$y = \frac{3}{4}x + 9$ and $y = \frac{3}{4}x - 9$ **[A1]**

[Back to Question] *[Total Marks: 7]*

802 **(a)** Gradient of $AB = \dfrac{4-2}{7-1} = \dfrac{1}{3}$ **[M1]**

Substitute $(10, 9)$ into $y - y_1 = m(x - x_1)$ with their gradient: $y - 9 = \dfrac{1}{3}(x - 10)$ **[M1]**

$y = \dfrac{1}{3}x + \dfrac{17}{3}$ or $3y = x + 17$ **[A1]**

(b) Gradient of $BC = \dfrac{9-4}{10-7} = \dfrac{5}{3}$ **[M1]**

Substitute $(1, 2)$ into $y - y_1 = m(x - x_1)$ with their gradient: $y - 2 = \dfrac{5}{3}(x - 1)$ **[M1]**

$y = \dfrac{5}{3}x + \dfrac{1}{3}$ or $3y = 5x + 1$ **[A1]**

(c) Equate equations from (a) and (b): $\dfrac{1}{3}x + \dfrac{17}{3} = \dfrac{5}{3}x + \dfrac{1}{3}$ **[M1]**

$4x = 16 \implies x = 4$ **[A1]**

$y = 7$ **[A1]**

[Back to Question] *[Total Marks: 9]*

803 **(a)** Gradient of $BC = 2.5$ (or $\dfrac{5}{2}$) **[B1]**

$2 = 2.5(-3) + c$ or $y - 2 = 2.5(x + 3)$ **[M1]**

$2.5x + 9.5$ (or $\dfrac{5}{2}x + \dfrac{19}{2}$) **[A1]**

(b) $D = (0, 9.5)$ **[B1]**

Translation $\overrightarrow{AB} = \begin{pmatrix} 8 \\ 2 \end{pmatrix}$ applied to D **[M1]**

$C = (8, 11.5)$ **[A1]**

(c) Gradient of $AB = \dfrac{4-2}{5-(-3)} = \dfrac{1}{4}$ **[B1]**

Midpoint of $BC = \left(\dfrac{5+8}{2}, \dfrac{4+11.5}{2}\right) = (6.5, 7.75)$ **[B1]**

$y - 7.75 = \dfrac{1}{4}(x - 6.5)$ **[M1]**

$2x - 8y = -49$ (or $-2x + 8y = 49$) **[A1]**

(d) $\sqrt{(8-(-3))^2 + (11.5-2)^2}$ $(= \sqrt{211.25})$ **[M1]**

14.5 **[A1]**

[Back to Question] *[Total Marks: 12]*

804 Rearranges to $5y = -2x + 15$ to find gradient of L is $-\dfrac{2}{5}$ **[M1]**

$\dfrac{5}{2}$ or 2.5 **[A1]**

[Back to Question] *[Total Marks: 2]*

805 Identifies the gradient of the given line as 3 **[B1]**

Uses the perpendicular gradient $-\frac{1}{3}$ and substitutes $(6, 2)$ into $y = mx + c$ or $y - y_1 = m(x - x_1)$ **[M1]**

$-\frac{1}{3}x + 4$ **[A1]**

[Back to Question] *[Total Marks: 3]*

806 **(a)** -2 **[B1]**

(b) Gradient of $L_2 = \frac{1}{2}$ **[M1]**

$1 = \frac{1}{2}(4) + c$ or $y - 1 = \frac{1}{2}(x - 4)$ **[M1]**

$y = \frac{1}{2}x - 1$ **[A1]**

[Back to Question] *[Total Marks: 4]*

807 **(a)** 2 **[B1]**

(b) $-\frac{1}{2}$ (or -0.5) **[B1]**

(c) $3 = -\frac{1}{2}(4) + c$ or $y - 3 = -\frac{1}{2}(x - 4)$ **[M1]**

$c = 5$ **[A1]**

$-\frac{1}{2}x + 5$ **[A1]**

[Back to Question] *[Total Marks: 5]*

808 **(a)** Rearranges the equation to make y the subject: $y = \frac{5}{2}x + 4$ **[M1]**

$\frac{5}{2}$ or 2.5 **[A1]**

(b) $-\frac{2}{5}$ or -0.4 **[B1]**

(c) Substitutes $(10, -3)$ and $m = -\frac{2}{5}$ into linear equation: $-3 = -\frac{2}{5}(10) + c$ or $y - (-3) = -\frac{2}{5}(x - 10)$ **[M1]**

Solves for c: $-3 = -4 + c \implies c = 1$ **[M1]**

$-\frac{2}{5}x + 1$ (or $-0.4x + 1$) **[A1]**

[Back to Question] *[Total Marks: 6]*

809 Calculates the midpoint of $AB = \left(\frac{-4+2}{2}, \frac{7+(-1)}{2}\right) = (-1, 3)$ **[B1]**

Calculates the gradient of $AB = \dfrac{-1-7}{2-(-4)} = -\dfrac{8}{6} = -\dfrac{4}{3}$ **[B1]**

Finds the perpendicular gradient $= \dfrac{3}{4}$ **[M1]**

Substitutes the midpoint $(-1, 3)$ and gradient $\dfrac{3}{4}$ to find the equation, e.g.,
$y - 3 = \dfrac{3}{4}(x - (-1))$ **[M1]**

$3x - 4y = -15$ or $4y - 3x = 15$ or equivalent integer form **[A1]**

[Back to Question] *[Total Marks: 5]*

810 **(a)** Gradient of $L = -\dfrac{2}{3}$ **[M1]**

$\dfrac{3}{2}$ (or 1.5) **[A1]**

(b) **(i)** $\left(\dfrac{-3+9}{2}, \dfrac{8+(-2)}{2}\right)$ **[M1]**

$(3, 3)$ **[A1]**

(ii) Gradient of $AB = \dfrac{-2-8}{9-(-3)} = -\dfrac{5}{6}$ **[M1]**

Gradient of perpendicular bisector $= \dfrac{6}{5}$ **[A1]**

$y - 3 = \dfrac{6}{5}(x - 3)$ **[M1]**

$6x - 5y = 3$ (or $-6x + 5y = -3$) **[A1]**

[Back to Question] *[Total Marks: 8]*

811 **(a)** $\left(\dfrac{-3+9}{2}, \dfrac{8+(-2)}{2}\right)$ **[M1]**

$(3, 3)$ **[A1]**

(b) Gradient of $AB = \dfrac{-2-8}{9-(-3)}$ or $-\dfrac{10}{12}$ or $-\dfrac{5}{6}$ **[M1]**

Perpendicular gradient is $\dfrac{6}{5}$ **[M1]**

Substitute $(3, 3)$ into $y - 3 = \dfrac{6}{5}(x - 3)$ or $3 = \dfrac{6}{5}(3) + c$ **[M1]**

$6x - 5y = 3$ or $-6x + 5y = -3$ **[A1]**

[Back to Question] *[Total Marks: 6]*

812 **(a)** $(0, 6)$ **[B1]**

(b) Gradient of $L_1 = -\dfrac{3}{2}$ **[M1]**

Gradient of $L_2 = \dfrac{2}{3}$ **[M1]**

$y = \frac{2}{3}x + 6$ **[A1]**

(c) Coordinates of Q: $y = \frac{2}{3}(9) + 6 = 12 \implies Q(9, 12)$ **[B1]**

Coordinates of R: $3(4) + 2y = 12 \implies y = 0 \implies R(4, 0)$ **[B1]**

Identify triangle PQR is right-angled at P OR setup valid box/shoelace method **[M1]**

Length $PQ = \sqrt{9^2 + 6^2} = \sqrt{117}$ and Length $PR = \sqrt{4^2 + (-6)^2} = \sqrt{52}$ **[M1]**

Area $= \frac{1}{2} \times \sqrt{117} \times \sqrt{52} = 39$ **[A1]**

[Back to Question] *[Total Marks: 9]*

813 **(a)** Rearranges to $2y = -kx + 12$ to identify gradient $-\frac{k}{2}$ **[B1]**

(b) Uses the gradient formula: $\frac{(k+2) - 5}{2 - 0}$ **[M1]**

$\frac{k-3}{2}$ **[A1]**

(c) Applies the perpendicularity condition $m_1 \times m_2 = -1$ **[M1]**

Forms the equation: $\left(-\frac{k}{2}\right)\left(\frac{k-3}{2}\right) = -1$ **[M1]**

Expands and rearranges to form a quadratic equation:
$-k(k-3) = -4 \implies k^2 - 3k - 4 = 0$ **[M1]**

4 and -1 **[A1]**

(d) Deduces that $k = -1$ must be chosen because $m_1 = -\frac{k}{2}$ must be positive (yielding $m_1 = \frac{1}{2}$) **[M1]**

Calculates the gradient of L_2 when $k = -1$: $m_2 = \frac{-1-3}{2} = -2$, and uses the y-intercept from point $A(0, 5)$ to write $y = -2x + 5$ **[M1]**

$2x + y = 5$ **[A1]**

[Back to Question] *[Total Marks: 10]*

814 $4y = 3x - 12$ implies gradient of P is $\frac{3}{4}$ **[M1]**

Gradient of Q is $-\frac{4}{3}$ **[M1]**

Substitute $(6, -2)$ into $y = -\frac{4}{3}x + c$ or $y - (-2) = -\frac{4}{3}(x - 6)$ **[M1]**

$-\frac{4}{3}x + 6$ **[A1]**

[Back to Question] *[Total Marks: 4]*

815 **(a)** $\left(\frac{-2+6}{2}, \frac{4+0}{2}\right)$ **[M1]**

$(2, 2)$ **[A1]**

(b) Gradient of $AB = \frac{0-4}{6-(-2)} = -\frac{1}{2}$ **[M1]**

Gradient of perpendicular bisector $= 2$ **[M1]**

Substitute midpoint $(2, 2)$ into $y - y_1 = m(x - x_1)$: $y - 2 = 2(x - 2)$ **[M1]**

$y = 2x - 2$ **[A1]**

[Back to Question] *[Total Marks: 6]*

816 **(a)** Uses the midpoint formula: $\left(\frac{-4+2}{2}, \frac{5+(-7)}{2}\right)$ **[M1]**

$(-1, -1)$ **[A1]**

(b) Uses the gradient formula: $\frac{-7-5}{2-(-4)}$ or $\frac{5-(-7)}{-4-2}$ **[M1]**

$\frac{-12}{6} = -2$ **[A1]**

(c) Identifies the perpendicular gradient is $\frac{1}{2}$ (or 0.5) **[B1]**

Substitutes their midpoint $(-1, -1)$ and perpendicular gradient into $y = mx + c$:
$-1 = \frac{1}{2}(-1) + c$ **[M1]**

$\frac{1}{2}x - \frac{1}{2}$ (or $0.5x - 0.5$) **[A1]**

(d) Sets $y = 0$ in their equation from part (c): $0 = \frac{1}{2}x - \frac{1}{2}$ **[M1]**

$(1, 0)$ **[A1]**

[Back to Question] *[Total Marks: 9]*

817 **(a)** Substitutes $x = 0$ into $2y + 3x = 12$ because P is on the y-axis **[M1]**

$(0, 6)$ **[A1]**

(b) Rearranges L_1 equation to find gradient: $2y = -3x + 12 \implies y = -\frac{3}{2}x + 6$, so $m = -\frac{3}{2}$ **[B1]**

States perpendicular gradient for L_2 is $\frac{2}{3}$ **[M1]**

$\frac{2}{3}x + 6$ **[A1]**

(c) Substitutes $y = 0$ into the equation of L_2 to find the coordinates of Q **[M1]**

$0 = \frac{2}{3}x + 6 \implies x = -9$ **[A1]**

Uses the distance formula for $P(0, 6)$ and $Q(-9, 0)$: $\sqrt{(-9-0)^2 + (0-6)^2}$ **[M1]**

$\sqrt{117}$ or $3\sqrt{13}$ **[A1]**

[Back to Question] *[Total Marks: 9]*

818 **(a)** Gradient of $AB = \dfrac{5-2}{5-(-1)} = \dfrac{1}{2}$ **[M1]**

Gradient of $BC = -2$ **[A1]**

$-2x + 15$ **[A1]**

(b) $0 = -2x + 15$ **[M1]**

$(7.5, 0)$ **[A1]**

(c) $AB = \sqrt{(5-(-1))^2 + (5-2)^2} = \sqrt{45}$ and $BC = \sqrt{(7.5-5)^2 + (0-5)^2} = \sqrt{31.25}$ **[M1]**

Area $= \dfrac{1}{2} \times \sqrt{45} \times \sqrt{31.25}$ **[M1]**

18.75 **[A1]**

(d) Midpoint of $AC = \left(\dfrac{-1+7.5}{2}, \dfrac{2+0}{2}\right) = (3.25, 1)$ **[B1]**

Gradient of $AC = \dfrac{0-2}{7.5-(-1)} = -\dfrac{4}{17}$, so perpendicular gradient $= \dfrac{17}{4}$ $(= 4.25)$ **[M1]**

$y - 1 = 4.25(0 - 3.25)$ **[M1]**

$(0, -12.8)$ **[A1]**

[Back to Question] *[Total Marks: 12]*

819 **(a)** Hexagon **[B1]**

(b) Obtuse **[B1]**

(c) Equilateral **[B1]**

[Back to Question] *[Total Marks: 3]*

820 **(a)** Chord **[B1]**

(b) Tangent **[B1]**

(c) Sector **[B1]**

(d) Acute **[B1]**

[Back to Question] *[Total Marks: 4]*

821 **(a)** Chord **[B1]**

(b) Tangent **[B1]**

(c) Minor segment (Accept segment) **[B1]**

(d) Isosceles **[B1]**

[Back to Question] *[Total Marks: 4]*

822 **(a)** Radius **[B1]**

(b) Tangent **[B1]**

(c) Diameter **[B1]**

(d) 90 **[B1]**

[Back to Question] *[Total Marks: 4]*

823 **(a)** **(i)** $(n-2) \times 180 = 1080$ or $1080 \div 180$ **[M1]**

8 **[A1]**

(ii) Octagon **[B1]**

(b) Rhombus **[B1]**

(c) Cylinder **[B1]**

[Back to Question] *[Total Marks: 5]*

824 **(a)** **(i)** tangent **[B1]**

(ii) chord **[B1]**

(iii) minor segment (accept segment) **[B1]**

(b) Relationship: perpendicular **[B1]**

Angle $OTQ = 90°$ **[B1]**

[Back to Question] *[Total Marks: 5]*

825 **(a)** Tangent **[B1]**

(b) Radius **[B1]**

(c) 90 **[B1]**

(d) Uses $2 \times \pi \times 5$ **[M1]**

10π **[A1]**

[Back to Question] *[Total Marks: 5]*

826 **(a)** Kite **[B1]**

(b) Sector (or minor sector) **[B1]**

(c) Chord **[B1]**

(d) Congruent **[B1]**

[Back to Question] *[Total Marks: 4]*

827 **(a)** Kite **[B1]**

(b) Perpendicular **[B1]**

(c) Congruent **[B1]**

(d) Obtuse **[B1]**

(e) Perpendicular bisector **[B1]**

[Back to Question] *[Total Marks: 5]*

828 **(a)** $180 - 156 = 24$ **[B1]**

(b) $\frac{360}{24}$ **[M1]**

15 **[A1]**

(c) Decagon **[B1]**

(d) Obtuse **[B1]**

[Back to Question] *[Total Marks: 5]*

829 **(a)** Tangent **[B1]**

(b) Chord **[B1]**

(c) Segment **[B1]**

(d) Sector **[B1]**

(e) Reflex **[B1]**

[Back to Question] *[Total Marks: 5]*

830 **(a)** trapezium **[B1]**

(b) **(i)** obtuse **[B1]**

(ii) $360 - 112 = 248$ **[B1]**

(c) **(i)** 34 **[B1]**

(ii) Angle $BCA = 180° - 112° - 34° = 34°$ (or interior angle $BCD = 68°$) **[M1]**

112 **[A1]**

[Back to Question] *[Total Marks: 6]*

831 **(a)** **(i)** Exterior angle $= 180° - 144° = 36°$ **[M1]**

$\frac{360}{36} = 10$ **[A1]**

(ii) Decagon **[B1]**

(b) **(i)** Sum of interior angles $= (6 - 2) \times 180° = 720°$ **[B1]**

$x + 2x + (x + 30) + (3x + 60) + 110 + 100 = 720$ OR $7x + 300 = 720$ **[M1]**

$x = 60$ **[A1]**

(ii) Right angle (since $60 + 30 = 90$) **[B1]**

(iii) Reflex angle (since $3(60) + 60 = 240$) **[B1]**

[Back to Question] *[Total Marks: 8]*

832 **(a)** $\frac{(5-2)\times 180}{5}$ or $180 - \frac{360}{5}$ **[M1]**

108 **[A1]**

(b) $\frac{(6-2)\times 180}{6}$ or $180 - \frac{360}{6}$ **[M1]**

120 **[A1]**

(c) Identifies angles around point A: $108°$ and $120°$ **[M1]**

Calculates reflex angle $= 360 - 108 - 120$ **[M1]**

132 **[A1]**

[Back to Question] *[Total Marks: 7]*

833 **(a)** Frustum **[B1]**

(b) **(i)** Similar **[B1]**

(ii) Scale factor $= \frac{20+10}{10} = 3$ or $\frac{10}{30} = \frac{1}{3}$ **[M1]**

$r = \frac{12}{3} = 4$ **[A1]**

(iii) Radius difference $= 12 - 4 = 8$ **[M1]**

$\sqrt{20^2 + 8^2}$ **[M1]**

21.5 or 21.54... **[A1]**

[Back to Question] *[Total Marks: 7]*

834 **(a)** Isosceles **[B1]**

(b) $138 + 180$ or $360 - (180 - 138)$ **[M1]**

318 **[A1]**

(c) **(i)** Perpendicular bisector **[B1]**

(ii) Angle $BAC = 138 - 62 = 76°$ **[M1]**

Bearing $= 062 + (76 \div 2) = 100°$ **[A1]**

(d) Trapezium **[B1]**

[Back to Question] *[Total Marks: 7]*

835 **(a)** $\dfrac{h}{h+12} = \dfrac{3}{5}$ **[M1]**

$5h = 3(h+12)$ **[M1]**

$5h = 3h + 36 \implies 2h = 36 \implies h = 18$ **[A1]**

(b) $\dfrac{r}{r+5} = \dfrac{3}{5}$ or $\dfrac{r}{r+5} = \dfrac{18}{30}$ **[M1]**

$5r = 3r + 15 \implies 2r = 15 \implies r = 7.5$ **[A1]**

(c) Volume ratio (small cone : large cone) $= \left(\dfrac{3}{5}\right)^3 = \dfrac{27}{125}$ **[M1]**

Volume of frustum $= 125 - 27 = 98$, so $k = 98$ **[A1]**

[Back to Question] *[Total Marks: 7]*

836 **(a)** Uses similar triangles to find the height H of the full cone: $\dfrac{H}{12} = \dfrac{H-15}{4}$ **[M1]**

$4H = 12H - 180 \implies 8H = 180 \implies H = 22.5$ cm **[A1]**

Volume of full cone $= \dfrac{1}{3} \times \pi \times 12^2 \times 22.5 = 1080\pi$ **[M1]**

Volume of small cone removed $= \dfrac{1}{3} \times \pi \times 4^2 \times (22.5 - 15) = \dfrac{1}{3} \times \pi \times 16 \times 7.5 = 40\pi$ **[M1]**

Volume of frustum $= 1080\pi - 40\pi = 1040\pi$ or 3267 or 3270 (3 sf) **[A1]**

(b) Identifies the area scale factor is $\dfrac{9}{25}$ **[B1]**

Calculates the length scale factor: $\sqrt{\dfrac{9}{25}} = \dfrac{3}{5}$ **[M1]**

New height $= 15 \times \dfrac{3}{5} = 9$ **[A1]**

[Back to Question] *[Total Marks: 8]*

837 **(a)** Setting up an equation connecting interior and exterior angles, e.g., $4x + x = 180$ **[M1]**

Exterior angle $= 36°$ or Interior angle $= 144°$ **[A1]**

$n = 10$ **[A1]**

Decagon **[B1]**

(b) Faces $= 12$ **[B1]**

Edges $= 30$ **[B1]**

Vertices $= 20$ **[B1]**

(c) Finding the linear scale factor $\sqrt{\dfrac{16}{9}}$ or $\dfrac{4}{3}$ **[M1]**

Finding the volume scale factor $\left(\dfrac{4}{3}\right)^3$ or $\dfrac{64}{27}$ **[M1]**

256π **[A1]**

[Back to Question] *[Total Marks: 10]*

838 **(a)** Angle ADE = Angle ABC because they are corresponding angles **[B1]**

Angle A is common to both triangles OR Angle AED = Angle ACB (corresponding angles) **[B1]**

(b) Length scale factor $= \frac{6}{10}$ or 0.6 **[M1]**

$DE = 15 \times 0.6 = 9$ **[A1]**

(c) $AC = \frac{7.5}{0.6} = 12.5$ **[M1]**

$EC = 12.5 - 7.5 = 5$ **[A1]**

(d) Trapezium **[B1]**

(e) Area scale factor $= 0.6^2 = 0.36$ **[M1]**

Area of triangle $ABC = \frac{18}{0.36} = 50$ **[M1]**

Area of $DBCE = 50 - 18 = 32$ **[A1]**

[Back to Question] *[Total Marks: 10]*

839 **(a)** **(i)** frustum **[B1]**

(ii) tetrahedron **[B1]**

(b) Reason: the cut is parallel to the base so corresponding angles are equal **[B1]**

Scale factor = 0.4 (or $\frac{2}{5}$) **[B1]**

(c) $10 \times \frac{6}{15}$ **[M1]**

4 **[A1]**

(d) **(i)** Number of sides $= \frac{360}{36} = 10$ **[M1]**

decagon **[A1]**

(ii) Faces = 12 **[B1]**

Edges = 30 **[B1]**

[Back to Question] *[Total Marks: 10]*

840 $048 + 180$ **[M1]**

228 **[A1]**

[Back to Question] *[Total Marks: 2]*

841 Identifies the relationship between alternate angles or uses $180° + 115°$ **[M1]**

295 **[A1]**

[Back to Question] *[Total Marks: 2]*

842 **(a)** $180 + 58$ **[M1]**

238 **[A1]**

(b) $180 - 84 = 096$ or 096 **[B1]**

[Back to Question] *[Total Marks: 3]*

843 **(a)** $068 + 180$ **[M1]**

248 **[A1]**

(b) 270 **[B1]**

[Back to Question] *[Total Marks: 3]*

844 **(a)** $128 + 180$ **[M1]**

308 **[A1]**

(b) 52 **[B1]**

(c) $180 - 90 - 52$ **[M1]**

38 **[A1]**

[Back to Question] *[Total Marks: 5]*

845 **(a)** $180 + 148$ OR $360 - (180 - 148)$ **[M1]**

328 **[A1]**

(b) **(i)** 270 **[B1]**

(ii) Identifies the bearing of P from B is $090°$ **[B1]**

$148 - 90 = 58$ **[A1]**

[Back to Question] *[Total Marks: 5]*

846 **(a)** $124 + 180$ **[M1]**

304 **[A1]**

(b) $\angle PQR = 90°$ or $\angle QPR = 180° - 124° = 56°$ **[M1]**

$\angle PRQ = 180° - 90° - 56° = 34°$ **[A1]**

[Back to Question] *[Total Marks: 4]*

847 **(a)** Bearing of P from $Q = 115 + 180 = 295°$ **[M1]**

Angle $PQR = 295° - 205° = 90°$ **[A1]**

(b) $15^2 + 8^2$ **[M1]**

$225 + 64 = 289$ **[M1]**

17 **[A1]**

[Back to Question] *[Total Marks: 5]*

848 **(a)** Interior angle at Q on North line $= 180° - 125° = 55°$ **[M1]**

Angle $PQR = 360° - 240° - 55° = 65°$ **[A1]**

(b) $PR^2 = 45° + 60° - 2(45)(60)\cos(65°)$ **[M2]**

$PR = 57.8$ or $57.82...$ **[A1]**

[Back to Question] *[Total Marks: 5]*

849 **(a)** 150 **[B1]**

(b) $180 + 65$ OR $360 - (180 - 65)$ **[M1]**

245 **[A1]**

(c) Identifies bearing of A from B is 245° (can follow through from (b)) **[M1]**

$245 - 70 = 175$ **[A1]**

(d) 25 **[B1]**

[Back to Question] *[Total Marks: 6]*

850 **(a)** Calculates the interior angle at Q on the North side: $180° - 50° = 130°$ OR bearing of P from $Q = 180° + 50° = 230°$ **[M1]**

Calculates angle PQR: $360° - 130° - 140° = 90°$ OR $230° - 140° = 90°$ **[A1]**

(b) Uses Pythagoras' theorem: $PR^2 = 40^2 + 60^2$ **[M1]**

$72.11\cdots = 72.1$ **[A1]**

(c) Uses trigonometry to find angle QPR: $\tan(QPR) = \dfrac{60}{40}$ **[M1]**

Angle $QPR = 56.3\ldots°$ **[M1]**

Bearing $= 050° + 56.3° = 106.3°$ **[A1]**

[Back to Question] *[Total Marks: 7]*

851 **(a)** Interior angle at B (South to BA) $= 35°$ OR interior angle from North line $= 180 - 35 = 145°$ **[M1]**

Angle $ABC = 360 - 145 - 125 = 90°$ OR $125 - 35 = 90°$ **[A1]**

(b) $AC^2 = 12^2 + 16^2$ **[M1]**

$AC^2 = 144 + 256 = 400$ **[M1]**

$AC = 20$ **[A1]**

(c) $\tan(\angle BAC) = \dfrac{16}{12}$ or $\sin(\angle BAC) = \dfrac{16}{20}$ **[M1]**

$\angle BAC = 53.13°$ **[A1]**

Bearing $= 035° + 53.13° = 088.1°$ **[A1]**

[Back to Question] *[Total Marks: 8]*

852 **(a)** Bearing of A from B is $065° + 180° = 245°$ or Interior angle between North and AB is $180° - 65° = 115°$ **[M1]**

$245° - 155° = 90°$ or $360° - 115° - 155° = 90°$ **[A1]**

(b) Triangle ABC is right-angled and isosceles, so $\angle BCA = (180° - 90°) \div 2 = 45°$ **[M1]**

Bearing of B from C $= 155° + 180° = 335°$ **[M1]**

Bearing of A from C $= 335° - 45° = 290°$ **[A1]**

(c) $\angle DAB = 90°$ **[M1]**

Bearing of D from A $= 065° + 90° = 155°$ **[A1]**

[Back to Question] *[Total Marks: 7]*

853 **(a)** Calculates angle $ABC = 240° - 170° = 70°$ (or equivalent geometric reasoning) **[B1]**

Uses the cosine rule: $AC^2 = 120° + 180^2 - 2(120)(180)\cos(70°)$ **[M1]**

$AC^2 = 14400 + 32400 - 43200(0.3420\ldots) = 32024.7\ldots$ **[M1]**

$178.9\cdots = 179$ (to 3 sf) **[A1]**

(b) Uses the sine rule to find angle BAC: $\dfrac{\sin(BAC)}{180} = \dfrac{\sin(70°)}{178.95}$ **[M1]**

$\sin(BAC) = 0.9452\cdots \implies$ Angle $BAC = 70.9°$ **[M1]**

Bearing $= 060° + 70.9° = 130.9°$ **[A1]**

[Back to Question] *[Total Marks: 7]*

854 **(a)** $\angle QPR = 118° - 42° = 76°$ **[M1]**

$QR^2 = 24^2 + 35^2 - 2(24)(35)\cos(76°)$ **[M1]**

$QR = 37.3$ or $37.34\ldots$ **[A1]**

(b) $\dfrac{\sin(\angle PQR)}{35} = \dfrac{\sin(76°)}{37.34}$ OR $\cos(\angle PQR) = \dfrac{24^2 + 37.34° - 35^2}{2(24)(37.34)}$ **[M1]**

$\sin(\angle PQR) = 0.909\ldots$ OR $\cos(\angle PQR) = 0.415\ldots$ **[M1]**

$\angle PQR = 65.4°$ or $65.42\ldots°$ **[A1]**

Bearing $= 180 + 42 - 65.4 = 156.6°$ (or $157°$) **[A1]**

[Back to Question] *[Total Marks: 7]*

855 **(a)** Angle $ABC = 360° - 145° - (180° - 42°) = 77°$ **[M1]**

$\dfrac{\sin(BAC)}{BC} = \dfrac{\sin(77°)}{18}$ **[M1]**

$BC^2 = 12^2 + 18^2 - 2(12)(18)\cos(BAC)$ (Implicitly requires finding BC or BAC first. Correct approach: $\frac{\sin(BCA)}{12} = \frac{\sin(77°)}{18}$) **[M1]**

$\sin(BCA) = \frac{12\sin(77°)}{18} = 0.6495...$ **[A1]**

Angle $BCA = 40.5°$ **[A1]**

Angle $BAC = 180° - 77° - 40.5° = 62.5°$ **[M1]**

Bearing $= 042° + 62.5° = 104.5°$ **[A1]**

(b) Boat sails on perpendicular bisector of BC. Angle $BAC = 62.5°$. The bisector of angle BAC is not necessarily the perpendicular bisector of BC.

Line is perpendicular bisector of BC.

Let midpoint of BC be M. In $\triangle ABM$, AM is the path.

Use cosine rule to find BC: $BC^2 = 12^2 + 18^2 - 2(12)(18)\cos(62.5°)$

BC = 16.39...

$\sin(ABM)/AM = \sin(BAM)/8.19...$ (This approach is overly complex).

Simpler: Path is perpendicular bisector. Angle of BC with North is 145°.

Perpendicular to BC has bearing $145° - 90° = 055°$.

Recognise path is perpendicular to BC **[M1]**

Bearing $= 145 - 90 = 055$ (or 055°) **[A1]**

[Back to Question] *[Total Marks: 7]*

856 **(a)** Bearing of P from $A = 050 + 180 = 230°$ **[M1]**

Angle $PAB = 230° - 160° = 70°$ **[M1]**

$PB^2 = 40^2 + 60^2 - 2(40)(60)\cos(70°)$ **[M1]**

59.6 or 59.65... **[A1]**

(b) Uses sine rule or cosine rule to find angle APB **[M1]**

$\frac{\sin(APB)}{60} = \frac{\sin(70°)}{59.65}$ or $\cos(APB) = \frac{40^2 + 59.65^2 - 60^2}{2(40)(59.65)}$ **[M1]**

Angle $APB = 70.9°$ $(70.93...°)$ **[A1]**

Bearing $= 050 + 70.9 = 120.9°$ **[A1]**

[Back to Question] *[Total Marks: 8]*

857 **(a)** Identifies angle $PQR = 230° - 120° = 110°$ (or uses parallel lines to find $180 - 50 = 130$ and $360 - 130 - 120 = 110$) **[B1]**

Uses Cosine rule: $PR^2 = 45^2 + 60^2 - 2(45)(60)\cos(110°)$ [M2] (M1 for correct substitution into formula)

$PR = 86.4$ or 86.44... **[A1]**

(b) Uses Sine rule: $\frac{\sin(QPR)}{60} = \frac{\sin(110°)}{86.44}$ **[M1]**

Angle $QPR = 40.7°$ $(40.73\ldots)$ **[A1]**

Bearing $= 050° + 40.7° = 090.7°$ **[A1]**

(c) $090.7 + 180$ **[M1]**

270.7 **[A1]**

[Back to Question] *[Total Marks: 9]*

858 Scale factor $= \frac{6+3}{6} = 1.5$ (or ratio $\frac{QR}{8} = \frac{9}{6}$) **[M1]**

12 **[A1]**

[Back to Question] *[Total Marks: 2]*

859 (a) Linear scale factor $= \frac{4+2}{4} = 1.5$ (or equivalent ratio $\frac{6}{4}$) **[M1]**

7.5 **[A1]**

(b) Area of $ABC = 8 \times 1.5^2 = 18$ **[M1]**

10 **[A1]**

[Back to Question] *[Total Marks: 4]*

860 (a) Scale factor $= \frac{18}{6} = 3$ **[M1]**

15 **[A1]**

(b) Area scale factor $= 3^2 = 9$ **[M1]**

$\frac{72}{9} = 8$ **[A1]**

[Back to Question] *[Total Marks: 4]*

861 (a) Volume scale factor $= \left(\frac{15}{6}\right)^3 = \frac{125}{8}$ **[M1]**

$V_B = 48 \times \frac{125}{8} = 750$ **[A1]**

(b) Area scale factor $= \left(\frac{15}{6}\right)^2 = \frac{25}{4}$ **[M1]**

$A_A = 200 \div \frac{25}{4} = 32$ **[A1]**

[Back to Question] *[Total Marks: 4]*

862 (a) Length scale factor $= \frac{9}{3} = 3$ **[M1]**

15 **[A1]**

(b) Area scale factor $= 3^2 = 9$ **[M1]**

6 **[A1]**

[Back to Question] *[Total Marks: 4]*

863 **(a)** Scale factor $= \frac{4+2}{4} = 1.5$ (or $\frac{6}{4}$ or $\frac{3}{2}$) **[M1]**

7.5 **[A1]**

(b) Area scale factor $= 1.5^2 = 2.25$ **[M1]**

Area of triangle $ABC = 8 \times 2.25 = 18$ **[M1]**

10 **[A1]**

[Back to Question] *[Total Marks: 5]*

864 **(a)** Scale factor $= \frac{4+2}{4}$ or 1.5 **[M1]**

9 **[A1]**

(b) Area scale factor $= 1.5^2$ or 2.25 **[M1]**

Area of triangle $ABC = 32 \times 2.25$ or 72 **[M1]**

40 **[A1]**

[Back to Question] *[Total Marks: 5]*

865 **(a)** Identifies the scale factor is $\frac{AD+DB}{AD} = \frac{6}{4}$ or 1.5 **[M1]**

9 **[A1]**

(b) Uses the area scale factor: 1.5^2 or $\frac{9}{4}$ or 2.25 **[M1]**

Calculates the area of triangle ABC: $12 \times 2.25 = 27$ **[M1]**

15 **[A1]**

[Back to Question] *[Total Marks: 5]*

866 **(a)** Finds the linear scale factor: $\frac{15}{10} = 1.5$ (or $\frac{3}{2}$) **[M1]**

Finds the area scale factor: $1.5^2 = 2.25$ (or $\frac{9}{4}$) **[M1]**

$120 \times 2.25 = 270$ **[A1]**

(b) Finds the volume scale factor: $1.5^3 = 3.375$ (or $\frac{27}{8}$) **[M1]**

Sets up the correct division: $540 \div 3.375$ (or $540 \times \frac{8}{27}$) **[M1]**

160 **[A1]**

[Back to Question] *[Total Marks: 6]*

867 **(a)** $\frac{6}{9} = \frac{8}{QR}$ or scale factor $= \frac{9}{6}$ (or 1.5) **[M1]**

12 **[A1]**

(b) $\frac{6}{9} = \frac{PT}{12}$ or $PT = 12 \div 1.5$ **[M1]**

8 **[A1]**

(c) Area scale factor $= \left(\frac{9}{6}\right)^2$ or 1.5^2 or 2.25 **[M1]**

Area of triangle $PQR = 20 \times 2.25 = 45$ **[M1]**

25 **[A1]**

[Back to Question] *[Total Marks: 7]*

868 **(a)** Area scale factor $= \frac{45}{20}$ or $\frac{9}{4}$ **[M1]**

Linear scale factor $= \sqrt{\frac{9}{4}} = \frac{3}{2}$ **[M1]**

Volume $B = 16 \times \left(\frac{3}{2}\right)^3 = 16 \times \frac{27}{8} = 54$ **[A1]**

(b) Height $A = 12 \div \frac{3}{2}$ or $12 \times \frac{2}{3}$ **[M1]**

8 **[A1]**

[Back to Question] *[Total Marks: 5]*

869 **(a)** Volume scale factor $= \frac{1350}{400} = \frac{27}{8}$ (or 3.375) **[M1]**

Linear scale factor $= \sqrt[3]{\frac{27}{8}} = \frac{3}{2}$ (or 1.5) **[M1]**

18 **[A1]**

(b) Area scale factor $= \left(\frac{3}{2}\right)^2 = \frac{9}{4}$ (or 2.25) **[M1]**

$\frac{360}{2.25} = 160$ **[A1]**

[Back to Question] *[Total Marks: 5]*

870 **(a)** Angle A (or $\angle DAE$ and $\angle CAB$) is common **[B1]**

Angle ADE = Angle ACB (given) **[B1]**

Therefore the third angles are equal ($\angle AED = \angle ABC$) and the triangles are similar (AAA) **[B1]**

(b) Sets up correct ratio: $\frac{AD}{AC} = \frac{AE}{AB}$ or $\frac{x}{4+5} = \frac{4}{x+(x+1)}$ **[M1]**

$\frac{x}{9} = \frac{4}{2x+1}$ **[M1]**

$2x^2 + x - 36 = 0$ **[M1]**

$(2x+9)(x-4) = 0$ implies $x = 4$ (ignoring negative root) **[A1]**

(c) Area scale factor $= \left(\frac{4}{9}\right)^2 = \frac{16}{81}$ **[M1]**

$16 : 65$ **[A1]**

[Back to Question] *[Total Marks: 9]*

871 **(a)** Linear scale factor $= 14 \div 4$ or 3.5 **[M1]**

21 **[A1]**

(b) Area scale factor $= 3.5^2$ or 12.25 **[M1]**

104.125 **[A1]**

[Back to Question] *[Total Marks: 4]*

872 **(a)** Volume scale factor $= \frac{1200}{150} = 8$ **[M1]**

Length scale factor $= \sqrt[3]{8} = 2$ **[M1]**

24 **[A1]**

(b) Area scale factor $= 2^2 = 4$ **[M1]**

110 **[A1]**

[Back to Question] *[Total Marks: 5]*

873 **(a)** Volume scale factor $= \frac{1080}{320}$ or $\frac{27}{8}$ or 3.375 **[M1]**

Linear scale factor $= \sqrt[3]{\frac{27}{8}}$ or 1.5 **[M1]**

27 **[A1]**

(b) Area scale factor $= 1.5^2$ or 2.25 or $\frac{9}{4}$ **[M1]**

Surface area of $A = 630 \div 2.25$ or $630 \times \frac{4}{9}$ **[M1]**

280 **[A1]**

[Back to Question] *[Total Marks: 6]*

874 **(a)** Volume scale factor $= \left(\frac{30}{12}\right)^3$ or 2.5^3 or 15.625 **[M1]**

5000 **[A1]**

(b) Area scale factor $= \left(\frac{30}{12}\right)^2$ or 2.5^2 or 6.25 **[M1]**

$1125 \div 6.25 = 180$ **[A1]**

(c) $4.80 \times \left(\frac{30}{12}\right)^3$ or 4.80×15.625 **[M1]**

75 or 75.00 **[A1]**

[Back to Question] *[Total Marks: 6]*

875 **(a)** Volume scale factor $= \frac{8640}{320} = 27$ **[M1]**

Length scale factor $= \sqrt[3]{27} = 3$ **[M1]**

36 **[A1]**

(b) Area scale factor $= 3^2 = 9$ **[M1]**

250 **[A1]**

(c) Mass $= 8.5 \times 320 = 2720\,\text{g}$ **[M1]**

2.72 **[A1]**

[Back to Question] *[Total Marks: 7]*

876 **(a)** Angle A is common to both triangles **[B1]**

Angle ABC = Angle ADE (corresponding angles as BC is parallel to DE) **[B1]**

(b) $\frac{DE}{7.5} = \frac{6+4}{6}$ **[M1]**

12.5 **[A1]**

(c) Area scale factor $= \left(\frac{10}{6}\right)^2 = \frac{25}{9}$ **[M1]**

Area of $ADE = 18 \times \frac{25}{9} = 50$ **[M1]**

Area of $BCED = 50 - 18 = 32$ **[A1]**

[Back to Question] *[Total Marks: 7]*

877 **(a)** States that angle A is common to both triangles. **[B1]**

States that angle ADE = angle ABC (and/or angle AED = angle ACB) because they are corresponding angles (due to parallel lines). **[B1]**

(b) Calculates the full length of AB: $6 + 4 = 10$ **[M1]**

Finds the linear scale factor: $\frac{10}{6} = \frac{5}{3}$ and sets up equation $BC = 9 \times \frac{5}{3}$ **[M1]**

15 **[A1]**

(c) Finds the area scale factor: $\left(\frac{5}{3}\right)^2 = \frac{25}{9}$ **[M1]**

Calculates the area of triangle ABC: $27 \times \frac{25}{9} = 75$ **[M1]**

Calculates the area of the trapezium: $75 - 27 = 48$ **[A1]**

[Back to Question] *[Total Marks: 8]*

878 **(a)** Linear scale factor $= \frac{45}{15} = 3$ **[M1]**

Mass (Volume) scale factor $= 3^3 = 27$ **[M1]**

13.5 **[A1]**

(b) Mass scale factor from P to $R = \frac{4000}{500} = 8$ **[M1]**

Linear scale factor $= \sqrt[3]{8} = 2$ **[M1]**

30 **[A1]**

(c) Area scale factor from P to $Q = 3^2 = 9$ **[M1]**

270 **[A1]**

[Back to Question] *[Total Marks: 8]*

879 **(a)** Calculates the volume ratio: $\frac{1080}{320} = \frac{27}{8}$ or 3.375 **[M1]**

Calculates the length scale factor: $\sqrt[3]{\frac{27}{8}} = \frac{3}{2}$ or 1.5 **[M1]**

18 **[A1]**

(b) Uses the area scale factor: $450 \div 1.5^2$ or $450 \times \left(\frac{2}{3}\right)^2$ **[M1]**

200 **[A1]**

(c) Calculates the volume of cone R: $320 + 1080 = 1400$ **[B1]**

Finds the length scale factor from P to R: $\sqrt[3]{\frac{1400}{320}}$ or $\sqrt[3]{4.375}$ **[M1]**

Calculates the height of R: $12 \times \sqrt[3]{4.375} = 19.626\ldots$ and rounds to 19.6 **[A1]**

[Back to Question] *[Total Marks: 8]*

880 **(a)** Angle AXB = Angle CXD (vertically opposite) **[B1]**

Angle ABX = Angle CDX (alternate angles, AB parallel to DC) **[B1]**

(Accept Angle BAX = Angle DCX for the second B1 mark)

(b) Length scale factor $= \frac{15}{9} = \frac{5}{3}$ **[M1]**

$DX = 6 \times \frac{5}{3} = 10$ **[M1]**

$BD = 6 + 10 = 16$ **[A1]**

(c) Area scale factor $= \left(\frac{15}{9}\right)^2 = \frac{25}{9}$ **[M1]**

Area $= 27 \times \frac{25}{9} = 75$ **[A1]**

(d) Ratio of bases $\frac{CX}{AX} = \frac{15}{9} = \frac{5}{3}$ **[M1]**

Area $= 27 \times \frac{5}{3} = 45$ **[A1]**

[Back to Question] *[Total Marks: 9]*

881 **(a)** $\frac{4}{4+x} = \frac{5}{7+x}$ OR $\frac{4}{x} = \frac{5}{x+2}$ **[M1]**

$4(x+7) = 5(x+4)$ OR $4(x+2) = 5x$ **[M1]**

$x = 8$ **[A1]**

(b) Linear scale factor $= \frac{4+8}{4} = 3$ **[M1]**

$BC = 3.5 \times 3 = 10.5$ **[A1]**

(c) Area of $\triangle ABC = 3^2 \times A = 9A$ **[M1]**

Area of trapezium $= 9A - A = 8A$ **[A1]**

[Back to Question] *[Total Marks: 7]*

882 **(a)** Volume ratio $A : B = \frac{108}{500} = \frac{27}{125}$ **[M1]**

Area ratio $= \left(\sqrt[3]{\frac{27}{125}}\right)^2 = \left(\frac{3}{5}\right)^2 = \frac{9}{25}$ **[M1]**

Let surface areas be $9x$ and $25x$. $9x + 25x = 340 \implies 34x = 340 \implies x = 10$ **[M1]**

$25 \times 10 = 250$ **[A1]**

(b) Volume of small cone $= 1 - \frac{7}{8} = \frac{1}{8}$ of Cone B **[M1]**

Ratio of heights $= \sqrt[3]{\frac{1}{8}} = \frac{1}{2}$ **[M1]**

Height of small cone $= 20 \times \frac{1}{2} = 10$ **[A1]**

[Back to Question] *[Total Marks: 7]*

883 **(a)** $\angle AXB = \angle DXC$ (vertically opposite angles) **[B1]**

$\angle BAX = \angle CDX$ OR $\angle ABX = \angle DCX$ (angles in the same segment) **[B1]**

Third angle in triangle OR states the other pair of angles in the same segment, concluding triangles are similar (AAA) **[B1]**

(b) Linear scale factor $= \frac{7.5}{5} = 1.5$ OR $\frac{DX}{4} = \frac{7.5}{5}$ **[M1]**

$DX = 6$ **[A1]**

$DC = 9$ **[A1]**

(c) Area scale factor $= 1.5^2 = 2.25$ **[M1]**

26.55 **[A1]**

[Back to Question] *[Total Marks: 8]*

884 **(a)** $\dfrac{x+2}{(x+2)+(x-1)} = \dfrac{8}{8+x}$ or $\dfrac{x+2}{x-1} = \dfrac{8}{x}$ **[M1]**

$(x+2)(x+8) = 8(2x+1)$ or $x(x+2) = 8(x-1)$ **[M1]**

$x^2 + 10x + 16 = 16x + 8$ or $x^2 + 2x = 8x - 8$ leading to $x^2 - 6x + 8 = 0$ **[A1]**

(b) $(x-2)(x-4) = 0$ **[M1]**

2 and 4 **[A1]**

(c) When $x = 4$, length ratio $\dfrac{AD}{AB} = \dfrac{6}{9} = \dfrac{2}{3}$ **[M1]**

Area of $ABC = 24 \div \left(\dfrac{2}{3}\right)^2 = 54$ **[M1]**

30 **[A1]**

[Back to Question] *[Total Marks: 8]*

885 **(a)** Angle A is common to both triangles **[B1]**

Angle AED = Angle ABC (given in the question) **[B1]**

Therefore, Angle ADE = Angle ACB (sum of angles in a triangle is 180°), so the triangles are similar (AAA) **[B1]**

(b) $AB = 6 + 9 = 15$ cm **[M1]**

$\dfrac{AC}{AD} = \dfrac{AB}{AE} \implies \dfrac{AC}{6} = \dfrac{15}{5}$ **[M1]**

$AC = 18$, so $EC = 18 - 5 = 13$ **[A1]**

(c) Area scale factor for the triangles $= \left(\dfrac{5}{15}\right)^2 = \dfrac{1}{9}$ **[M1]**

1 : 8 **[A1]**

[Back to Question] *[Total Marks: 8]*

886 **(a)** States that angle BAX = angle DCX OR angle ABX = angle CDX due to alternate angles (between parallel lines) **[B1]**

States a second valid pair of angles are equal (e.g. vertically opposite angles at X or the other pair of alternate angles) concluding the triangles are similar (AAA) **[B1]**

(b) Identifies the length scale factor: $\dfrac{10.5}{7}$ or 1.5 **[M1]**

Calculates $CX = 4 \times 1.5 = 6$ **[M1]**

10 **[A1]**

(c) Calculates the area of triangle CDX: $16 \times 1.5^2 = 36$ **[M1]**

Recognises the ratio of areas of triangle BCX to triangle ABX is equal to the ratio of bases $CX : AX = 1.5 : 1$ (as they share the same height from B) **[M1]**

Calculates the area of triangle BCX (or ADX): $16 \times 1.5 = 24$ **[M1]**

Total area $= 16 + 36 + 24 + 24 = 100$ **[A1]**

[Back to Question] *[Total Marks: 9]*

887 **(a)** Angle A is common to both triangles **[B1]**

Angle ABC = Angle AED (given), so the third angles must be equal, hence AA similarity **[B1]**

(b) 2.5 or $\frac{5}{2}$ **[B1]**

(c) $\frac{AB}{AE} = 2.5 \implies \frac{2x + 2.5}{x} = 2.5$ (or equivalent valid ratio) **[M1]**

$2x + 2.5 = 2.5x$ **[M1]**

$x = 5$ **[A1]**

(d) Area scale factor $= 2.5^2 = 6.25$ **[M1]**

Formulates correct equation for areas, e.g., $6.25A - A = 42$ or $5.25A = 42$ **[M1]**

8 **[A1]**

[Back to Question] *[Total Marks: 9]*

888 **(a)** $\angle AXB = \angle CXD$ because they are vertically opposite angles **[B1]**

$\angle BAX = \angle DCX$ (or $\angle ABX = \angle CDX$) because they are alternate angles, $AB \parallel DC$ **[B1]**

All three pairs of corresponding angles are equal, so the triangles are similar (AAA) **[B1]**

(b) $\frac{2x - 3}{x} = \frac{6}{4}$ or equivalent **[M1]**

$4(2x - 3) = 6x$ or $2x - 3 = 1.5x$ **[M1]**

$0.5x = 3$ leading to $x = 6$ **[A1]**

(c) Area of triangle $CDX = 24 \times 1.5^2$ or 54 **[M1]**

Recognises triangle BCX shares altitude with ABX and base ratio is $6 : 4$ **[M1]**

Area of triangle BCX (and triangle ADX) $= 24 \times 1.5$ or 36 **[M1]**

Total area $= 24 + 54 + 36 + 36 = 150$ **[A1]**

[Back to Question] *[Total Marks: 10]*

889 **(a)** Finds the volume scale factor: $\frac{2700}{800} = 3.375$ (or $\frac{27}{8}$) **[M1]**

Finds the linear scale factor by cube rooting: $\sqrt[3]{3.375} = 1.5$ **[M1]**

$12 \times 1.5 = 18$ **[A1]**

(b) Finds the area scale factor from the linear scale factor: $1.5^2 = 2.25$ **[M1]**

Sets up the multiplication for the cost: 18×2.25 **[M1]**

40.50 **[A1]**

(c) Identifies the area scale factor between C and A is 1.44 **[B1]**

Finds the linear scale factor: $\sqrt{1.44} = 1.2$ **[M1]**

Finds the volume scale factor: $1.2^3 = 1.728$ **[M1]**

Calculates the volume of C: $800 \times 1.728 = 1382.4$ **[A1]**

[Back to Question] *[Total Marks: 10]*

890 **(a)** Slant height $l = \sqrt{10^2 + 24^2} = 26$ **[M1]**

Surface Area $= \pi(10)^2 + \pi(10)(26)$ **[M1]**

360π **[A1]**

(b) $V_A = \frac{1}{3}\pi(10)^2(24) = 800\pi$ **[M1]**

Volume Ratio $= \frac{100\pi}{800\pi} = \frac{1}{8}$ **[M1]**

Area Ratio $= \left(\frac{1}{8}\right)^{\frac{2}{3}} = \frac{1}{4}$ **[M1]**

Surface Area of $B = 360\pi \times \frac{1}{4} = 90\pi$ **[A1]**

(c) Volume of frustum $= 800\pi - 100\pi = 700\pi$ **[M1]**

$\frac{4}{3}\pi R^3 = 700\pi$ **[M1]**

$R^3 = 700 \times \frac{3}{4} = 525$ **[M1]**

$R = \sqrt[3]{525} = 8.07$ (accept $8.066\ldots$) **[A1]**

[Back to Question] *[Total Marks: 11]*

891 **(a)** 9 **[B1]**

(b) 2 **[B1]**

(c) Isosceles triangle **[B1]**

[Back to Question] *[Total Marks: 3]*

892 **(a)** 10 **[B1]**

(b) Rhombus OR Rectangle **[B1]**

(c) 72 **[B1]**

[Back to Question] *[Total Marks: 3]*

893 **(a)** 4 **[B1]**

(b) 0 **[B1]**

(c) Correct reflection of at least two 'petals' (e.g. vertices at $(1, 4)$ and $(4, 3)$) **[M1]**

Fully correct shape drawn **[A1]**

[Back to Question] *[Total Marks: 4]*

894 **(a)** **(i)** 15 **[B1]**

(ii) 15 **[B1]**

(b) Rectangle OR Rhombus **[B1]**

(c) 5 **[B1]**

[Back to Question] *[Total Marks: 4]*

895 Parallelogram: 0 lines, order 2 **[B2, B1 for each correct value]**

Regular Pentagon: 5 lines, order 5 **[B2, B1 for each correct value]**

Isosceles Trapezium: 1 line, order 1 **[B2, B1 for each correct value]**

[Back to Question] *[Total Marks: 6]*

896 **(a)** Identifies 6 vertical planes of symmetry or 1 horizontal plane of symmetry **[M1]**

7 **[A1]**

(b) 6 **[B1]**

(c) 4 **[B1]**

[Back to Question] *[Total Marks: 4]*

897 **(a)** 4 **[B1]**

(b) 6 **[B2]**

Strictly 5 (vertical) or 1 (horizontal) planes identified **[B1]**

(c) A straight line passing through the apex (top vertex) **[B1]**

and the centre of the circular base. **[B1]**

[Back to Question] *[Total Marks: 5]*

898 **(a)** Recognises that order 4 symmetry implies point symmetry (order 2) about M, so M is the midpoint of AC **[M1]**

$(5, -2)$ **[A1]**

(b) Gradient of $AC = \frac{-2-6}{5-1}$ or $\frac{-8}{4} = -2$ **[M1]**

Substitute $(1, 6)$ or $(5, -2)$ or $(3, 2)$ into $y = -2x + c$ OR $y - 6 = -2(x - 1)$ **[M1]**

$-2x + 8$ **[A1]**

(c) Deduces that $ABCD$ must be a square because it has order 4 rotational symmetry (area is $|AB|^2$) **[M1]**

$|AB|^2 = (7-1)^2 + (4-6)^2$ **[M1]**

40 **[A1]**

[Back to Question] *[Total Marks: 8]*

899 **(a)** 4 **[B2]**

If 0 scored, allow [B1] for identifying either 3 (vertical) OR 1 (horizontal/mid-plane).

(b) 3 **[B1]**

(c) 7 **[B2]**

If 0 scored, allow [B1] for 6.

[Back to Question] *[Total Marks: 5]*

900 **(a)** **(i)** $\frac{360}{15}$ **[M1]**

24 **[A1]**

(ii) 24 **[B1]**

(b) Kite OR Isosceles Trapezium (accept Isosceles Trapezoid) **[B1]**

(c) Infinite (accept an infinite number) **[B1]**

[Back to Question] *[Total Marks: 5]*

901 **(a)** **(i)** 4 **[B1]**

(ii) Infinite (or infinitely many) **[B1]**

(b) Uses volume formula: $\frac{1}{3} \times 6^2 \times h = 48$ **[M1]**

Solves for height: $h = 4$ **[A1]**

Calculates distance from centre of base to a corner $= \sqrt{3^2 + 3^2} = \sqrt{18}$ (or $3\sqrt{2}$ or 4.24...) **[M1]**

Uses Pythagoras' theorem for sloping edge $l = \sqrt{4^2 + (\sqrt{18})^2}$ **[M1]**

5.83 (or 5.830...) **[A1]**

[Back to Question] *[Total Marks: 7]*

902 **(a)** Understanding that diagonals of a rhombus intersect at 90°, hence gradient $m = -\frac{1}{3}$ **[M1]**

$-\frac{1}{3}x$ (accept equivalent forms) **[A1]**

(b) 2 **[B1]**

(c) Exterior angle $= 180° - 165° = 15°$ **[M1]**

$n = \frac{360}{15}$ **[M1]**

$n = 24$ **[A1]**

[Back to Question] *[Total Marks: 6]*

903 **(a)** **(i)** 8 **[B1]**

(ii) 9 **[B2]**

If B0, score B1 for 8 (identifying only the vertical planes)

(b) **(i)** 12 **[B1]**

(ii) $(12 - 2) \times 180$ **[M1]**

1800 **[A1]**

(c) $(-2, 3)$ **[B2]**

If B0, score M1 for point plotted correctly on grid OR one coordinate correct

[Back to Question] *[Total Marks: 8]*

904 **(a)** Calculates length $AB = \sqrt{(5-1)^2 + (4-2)^2} = \sqrt{20}$ **[M1]**

Calculates length $BC = \sqrt{(3-5)^2 + (8-4)^2} = \sqrt{20}$ **[M1]**

States $AB = BC$ therefore the triangle is isosceles **[A1]**

(b) Identifies midpoint of $AC = (\frac{1+3}{2}, \frac{2+8}{2}) = (2, 5)$ **[M1]**

Calculates gradient of line of symmetry $= \frac{4-5}{5-2} = -\frac{1}{3}$ (or uses perpendicular to AC) **[M1]**

Substitutes $(5, 4)$ or $(2, 5)$ into $y = -\frac{1}{3}x + c$ to find c **[M1]**

$-\frac{1}{3}x + \frac{17}{3}$ **[A1]**

(c) Identifies $n = 15$ and uses $\frac{(15-2) \times 180}{15}$ OR $180 - (360 \div 15)$ **[M1]**

156 **[A1]**

[Back to Question] *[Total Marks: 9]*

905 **(a)** States that a right prism with a regular n-gon base has n vertical planes of symmetry and 1 horizontal plane of symmetry (halfway along its length) **[B1]**

Sets up the equation $n + 1 = 7$, yielding $n = 6$ **[B1]**

(b) **(i)** Recognises the side length of the hexagon $= 12 \div 2 = 6$ cm **[B1]**

Calculates area of the hexagonal cross-section $= 6 \times \left(\frac{1}{2} \times 6 \times 6 \times \sin(60°)\right) = 54\sqrt{3}$ (or 93.53...) **[M1]**

Uses Volume = Area × Length to find Length $= \frac{450}{54\sqrt{3}}$ **[M1]**

4.81 (or 4.811...) **[A1]**

(ii) Method for total area = 2 × Base Area + 6 × Lateral Rectangular Area **[M1]**

Substitutes values: $2 \times 54\sqrt{3} + 6 \times (6 \times 4.811...)$ **[M1]**

Simplifies to $108\sqrt{3} + 100\sqrt{3}$ (or 187.06... + 173.2...) **[M1]**

360 (or 360.2... to 360.3...) **[A1]**

[Back to Question] *[Total Marks: 10]*

906 **(a)** 48 **[B1]**

(b) 56 **[B1]**

(c) $180 - 48 - 56 = 76$ **[B1]**

[Back to Question] *[Total Marks: 3]*

907 **(a)** 125 **[B1]**

(b) Corresponding angles (are equal) **[B1]**

(c) 144 **[B1]**

[Back to Question] *[Total Marks: 3]*

908 **(a)** Identifies alternate angles by drawing a parallel line through E OR forms a triangle by extending a line **[M1]**

85 **[A1]**

(b) $\frac{360}{24}$ **[M1]**

15 **[A1]**

[Back to Question] *[Total Marks: 4]*

909 **(a)** 52 **[B1]**

(b) Angles on a straight line sum to 180°: $180 - (52 + 61)$ **[M1]**

67 **[A1]**

(c) Identifies angle $TVU = 61°$ (alternate angle to QTV) or angle $VTW + TVW$ exterior relationship **[M1]**

Angle $TVW = 180 - 61 = 119$ **[A1]**

[Back to Question] *[Total Marks: 5]*

910 **(a)** $360 \div 24$ **[M1]**

15 **[A1]**

(b) $(15 - 2) \times 180$ **[M1]**

2340 **[A1]**

(c) Exterior angle $= 180 - 165 = 15°$ **[M1]**

$360 \div 15 = 24$ **[A1]**

[Back to Question] *[Total Marks: 6]*

911 **(a)** $180 - (55 + 70)$ **[M1]**

55 **[A1]**

(b) 55 **[B1]**

(c) Angle $DAE = 55°$ (from part a) AND Angle $ADE = 55°$ **[B1]**

Two angles are equal, so the triangle is isosceles **[B1]**

[Back to Question] *[Total Marks: 5]*

912 **(a)** Equates sum of angles to formula: $3 \times 140 + 165(n - 3) = 180(n - 2)$ **[M1]**

Expands brackets correctly: $420 + 165n - 495 = 180n - 360$ **[M1]**

Rearranges to isolate n: $15n = 285$ (or equivalent) **[M1]**

19 **[A1]**

(b) States $a + b + c = 180$ **[B1]**

Substitutes $a + b = c$ to get $c + c = 180 \implies 2c = 180 \implies c = 90$ completing the proof **[B1]**

(c) **(i)** Calculates exterior angle of $P = \dfrac{360}{12} = 30°$ so exterior angle of $Q = 15°$ **[M1]**

24 **[A1]**

(ii) Calculates interior angles of P and Q: $180 - 30 = 150°$ and $180 - 15 = 165°$ **[M1]**

$360 - (150 + 165) = 45$ **[A1]**

[Back to Question] *[Total Marks: 10]*

913 **(a)** Sum of interior angles of a pentagon $= (5 - 2) \times 180° = 540°$ **[M1]**

$x + (x + 20) + (2x - 10) + (2x + 30) + 140 = 540$ **[M1]**

$6x + 180 = 540 \implies 6x = 360 \implies x = 60$ **[A1]**

(b) Substituting $x = 60$ into $(2x + 30)$ or checking all angle sizes $(60°, 80°, 110°, 150°, 140°)$ **[M1]**

150 **[A1]**

[Back to Question] *[Total Marks: 5]*

914 **(a)** Recognises the sum of the interior angles of a pentagon is $540°$ **[B1]**

Sets up a correct equation: $(2x + 10) + 100 + 80 + 3x + (x + 50) = 540$ **[M1]**

50 **[A1]**

(b) Sets up a relationship for the angles, e.g., $4E + E = 180$ where E is the exterior angle **[M1]**

$E = 36°$ **[M1]**

10 **[A1]**

[Back to Question] *[Total Marks: 6]*

915 **(a)** $5x + 10 = 7x - 30$ **[M1]**

$x = 20$ **[A1]**

Corresponding angles (are equal) **[B1]**

(b) Angle $GHD = 7(20) - 30 = 110°$ **[B1]**

Angle $GHK = 180 - 110 = 70°$ (Angles on a straight line) **[B1]**

Angle $GKH = 180 - (70 + 35) = 75°$ (Angles in a triangle) **[B1]**

[Back to Question] *[Total Marks: 6]*

916 **(a)** **(i)** Equates the sum of the angles to the interior angle sum of a hexagon ($720°$):
$x + 2x + x + 30 + 2x + 10 + 3x - 20 + 160 = 720$ **[M1]**

Simplifies the equation to $9x + 180 = 720$ (or $9x = 540$) **[M1]**

60 **[A1]**

(ii) Sets up an equation for the exterior angle e:
$6.5e + e = 180°$ **[M1]**

$7.5e = 180° \implies e = 24°$ **[M1]**

Number of sides $n = \dfrac{360}{24} = 15$ **[A1]**

(iii) Let exterior angles be E_m and E_n. Sets up the first equation:
$E_m + E_n = 54$ **[B1]**

Relates interior difference to exterior difference:
$(180 - E_m) - (180 - E_n) = 18 \implies E_n - E_m = 18$ **[M1]**

Solves the simultaneous equations by addition (or substitution):
$2E_n = 72 \implies E_n = 36°$ **[M1]**

Calculates n: $n = \dfrac{360}{36} = 10$ **[A1]**

Calculates m: $E_m = 18° \implies m = \dfrac{360}{18} = 20$ **[A1]**

[Back to Question] *[Total Marks: 11]*

917 **(a)** Recognises that $\angle PXQ = \angle BPX + \angle DQX$ (e.g. by drawing a parallel line through X to form alternate angles) **[M1]**

$(3x + 15) + (2x + 10) = 75$ **[M1]**

$5x + 25 = 75 \implies 5x = 50 \implies x = 10$ **[A1]**

(b) Expression for exterior or interior angle of P: $\dfrac{360}{n}$ or $180 - \dfrac{360}{n}$ **[B1]**

Expression for exterior or interior angle of Q: $\dfrac{360}{3n}$ or $180 - \dfrac{120}{n}$ **[B1]**

Forms a correct equation: $\left(180 - \dfrac{120}{n}\right) - \left(180 - \dfrac{360}{n}\right) = 16$ OR $\dfrac{360}{n} - \dfrac{120}{n} = 16$ **[M1]**

$\dfrac{240}{n} = 16 \implies n = 15$ **[A1]**

[Back to Question] *[Total Marks: 7]*

918 **(a)** $\dfrac{(5-2) \times 180}{5}$ OR $180 - \dfrac{360}{5}$ **[M1]**

108 **[A1]**

(b) Interior angle of hexagon $= 120°$ **[B1]**

Angle $AEI = 360 - 108 - 120 = 132$ **[A1]**

(c) $AE = EI$ (sides of regular polygons sharing an edge are equal) **[M1]**

Triangle AEI is isosceles **[M1]**

Angle $EAI = \dfrac{180 - 132}{2} = 24$ **[A1]**

[Back to Question] *[Total Marks: 7]*

919 **(a)** $(5 - 2) \times 180 \div 5$ OR $180 - (360 \div 5)$ **[M1]**

108 **[A1]**

(b) Interior angle of square $= 90°$ **[B1]**

Interior angle of n-gon $= 360 - 90 - 108 = 162°$ **[M1]**

$n = 360 \div (180 - 162) = 20$ **[A1]**

(c) Recognises triangle PFA is isosceles (since $PA = PF$ from shared sides of regular polygons) **[M1]**

Angle $PFA = (180 - 162) \div 2 = 9$ **[A1]**

[Back to Question] *[Total Marks: 7]*

920 **(a)** Finds an expression for the interior angle ABC using the sum of angles at a point:
Interior angle $ABC = 360° - (300 - 2x)° = (60 + 2x)°$ **[M1]**

Draws a parallel line through B and uses alternate angles to form an equation:
Interior angle $ABC = (3x + 10)° + (2x + 20)°$ **[M1]**

Equates the two expressions for the interior angle ABC:

$5x + 30 = 60 + 2x$ **[M1]**

Solves the equation correctly to show the result:
$3x = 30 \implies x = 10$ **[A1]**

(b) Calculates the interior angle ABC using $x = 10$:
Angle $ABC = 60 + 2(10) = 80°$ **[M1]**

Recognises triangle ABC is isosceles and sets up the base angle calculation:
$\frac{180 - 80}{2}$ **[M1]**

50 **[A1]**

[Back to Question] *[Total Marks: 7]*

921 **(a)** 65 **[B1]**

(b) Uses isosceles triangle OAB, e.g. $\frac{180 - 130}{2}$ **[M1]**

25 **[A1]**

[Back to Question] *[Total Marks: 3]*

922 **(a)** 116 **[B1]**

(b) $\frac{180 - 116}{2}$ **[M1]**

32 **[A1]**

[Back to Question] *[Total Marks: 3]*

923 **(a)** 34 **[B1]**

(b) Recognises angle at centre is twice angle at circumference $\implies 2 \times 34$ **[M1]**

68 **[A1]**

[Back to Question] *[Total Marks: 3]*

924 **(a)** 55 **[B1]**

(b) 55 **[B1]**

(c) Angles in the same segment (are equal) **[B1]**

[Back to Question] *[Total Marks: 3]*

925 **(a)** 112 **[B1]**

(b) 124 **[B1]**

(c) Opposite angles of a cyclic quadrilateral sum to 180° **[B1]**

[Back to Question] *[Total Marks: 3]*

926 **(a)** **(i)** 48 **[B1]**

(ii) Angles in the same segment **[B1]**

(b) 96 **[B1]**

[Back to Question] *[Total Marks: 3]*

927 **(a)** 65 **[B1]**

(b) $180 - 65$ **[M1]**

115 **[A1]**

[Back to Question] *[Total Marks: 3]*

928 **(a)** 42 **[B1]**

(b) $180 - (42 + 65) = 73$ **[B1]**

(c) Angle BAD + Angle $BCD = 180$ (cyclic quadrilateral) **[M1]**

$180 - 73 = 107$ **[A1]**

[Back to Question] *[Total Marks: 4]*

929 **(a)** **(i)** 58 **[B1]**

(ii) Alternate segment theorem **[B1]**

(b) Angle $AOB = 2 \times 58 = 116°$ **[M1]**

Angle $OBA = \dfrac{180 - 116}{2} = 32°$ **[A1]**

[Back to Question] *[Total Marks: 4]*

930 **(a)** 55 **[B1]**

(b) Angle at the centre is twice the angle at the circumference. **[B1]**

(c) 30 **[B1]**

(d) $(180 - 110) \div 2$ **[M1]**

35 **[A1]**

[Back to Question] *[Total Marks: 5]*

931 **(a)** Angle at centre is twice the angle at the circumference **[M1]**

62 **[A1]**

(b) Opposite angles in a cyclic quadrilateral sum to 180° **[M1]**

$180 - 62 = 118$ **[A1]**

(c) Triangle OAC is isosceles ($OA = OC$) **[M1]**

$(180 - 124) \div 2 = 28$ **[A1]**

[Back to Question] *[Total Marks: 6]*

932 **(a)** **(i)** 42 **[B1]**

(ii) 55 **[B1]**

(b) $90 - 42$ **[M1]**

48 **[A1]**

(c) $180 - (55 + 42)$ **[M1]**

83 **[A1]**

[Back to Question] *[Total Marks: 6]*

933 **(a)** 55 **[B1]**

(b) Uses opposite angles of a cyclic quadrilateral (e.g., $180 - 100$) **[M1]**

80 **[A1]**

(c) Uses angles in a triangle sum to 180° (e.g., $180 - (55 + 80)$) **[M1]**

45 **[A1]**

[Back to Question] *[Total Marks: 5]*

934 **(a)** 42 **[B1]**

(b) 75 **[B1]**

(c) Uses sum of angles in triangle PRS equal to 180° ($180 - 75 - 48$) **[M1]**

57 **[A1]**

(d) 57 **[B1]**

[Back to Question] *[Total Marks: 5]*

935 **(a)** **(i)** 90 **[B1]**

(ii) Angle in a semicircle is 90° **[B1]**

(b) 54 **[B1]**

(c) $180 - 54$ **[M1]**

126 **[A1]**

[Back to Question] *[Total Marks: 5]*

936 **(a)** Angle at centre = 2× angle at circumference **[M1]**

$118 \times 2 = 236$ **[A1]**

(b) $360 - 236 = 124$ **[B1]**

(c) Triangle POR is isosceles, so angle OPR = angle $ORP = \dfrac{180 - 124}{2} = 28°$ **[M1]**

Angle $ORS = 90°$ (tangent-radius), so angle $PRS = 90 - 28 = 62$ **[A1]**

[Back to Question] *[Total Marks: 5]*

937 **(a)** Angle $OAT = 90°$ and Angle $OBT = 90°$ **[B1]**

$360 - 90 - 90 - 44 = 136$ **[A1]**

(b) 68 **[B1]**

(c) $180 - 68$ **[M1]**

112 **[A1]**

[Back to Question] *[Total Marks: 5]*

938 **(a)** **(i)** 62 **[B1]**

(ii) Angle in a semicircle (is 90°) **[B1]**

(b) 28 **[B1]**

(c) Angle $ABC = 62°$ OR Angle $ADB = 90°$ **[M1]**

118 **[A1]**

[Back to Question] *[Total Marks: 5]*

939 **(a)** **(i)** 58 **[B1]**

(ii) Alternate segment theorem **[B1]**

(b) Angle in a semicircle is 90° (Angle $ACB = 90°$) **[M1]**

$180 - 90 - 58$ **[M1]**

32 **[A1]**

(c) $180 - 32$ **[M1]**

148 **[A1]**

[Back to Question] *[Total Marks: 7]*

940 **(a)** Angle $PAD = 90°$ OR Angle $DAB = 90 - 58 = 32°$ **[M1]**

58 **[A1]**

(b) Angle $ADC = 58 + 22 = 80°$ **[M1]**

$180 - 80 = 100$ **[A1]**

(c) Angle $BOC = 2 \times 22 = 44°$ **[M1]**

$\frac{180-44}{2}$ **[M1]**

68 **[A1]**

[Back to Question] *[Total Marks: 7]*

941 **(a)** Forms correct equation using angle at centre theorem: $4x - 20 = 2(x + 32)$ **[M1]**

$4x - 20 = 2x + 64 \implies 2x = 84$ **[M1]**

$x = 42$ **[A1]**

(b) Recognises angle CDE = angle ABC (exterior angle of cyclic quadrilateral) OR finds angle $ADC = 180 - (42 + 32) = 106°$ **[M1]**

74 **[A1]**

[Back to Question] *[Total Marks: 5]*

942 **(a)** Recognition that tangent meets radius at 90°, so $\angle OAT = \angle OCT = 90°$ **[M1]**

$180 - 48 = 132$ **[A1]**

(b) 66 **[B1]**

(c) Use of isosceles triangle OAC ($OA = OC$ radii): $(180 - 132) \div 2$ **[M1]**

OR $\angle TAC = (180 - 48) \div 2 = 66°$ leading to $\angle OAC = 90 - 66$ **[M1]**

24 **[A1]**

[Back to Question] *[Total Marks: 5]*

943 **(a)** Angle $PQR = 90°$ **[M1]**

$180 - 90 - 32 = 58$ **[A1]**

(b) Angle $TPR = 90°$ **[M1]**

$90 - 32 = 58$ **[A1]**

(c) Angle $PRS = 180 - 90 - 41 = 49°$ **[M1]**

49 **[A1]**

[Back to Question] *[Total Marks: 6]*

944 **(a)** Angle $ACD = 90°$ (angle in a semicircle) **[M1]**

Angle $ADC = 180 - 90 - 24 = 66°$ **[A1]**

(b) $180 - 66 = 114°$ (opposite angles of cyclic quadrilateral) **[B1]**

(c) 24° (angles in same segment as angle CAD) **[B1]**

(d) Angle $ABD = 90°$ (angle in a semicircle) **[M1]**

In triangle ABX, angle $AXB = 180 - 90 - 32 = 58°$ **[A1]**

[Back to Question] *[Total Marks: 6]*

945 **(a)** Angle $ACB = 62$ **[B1]**

Alternate segment theorem **[B1]**

(b) $180 - 62$ **[M1]**

118 **[A1]**

(c) Angle $OAB = 90 - 62 = 28°$ **[M1]**

28 (since triangle OAB is isosceles) **[A1]**

[Back to Question] *[Total Marks: 6]*

946 **(a)** Angle $BAD = 180° - 115° = 65°$ **[B1]**

$(2x + 15) + 65 + (x + 10) = 180$ **[M1]**

$3x + 90 = 180$ **[M1]**

30 **[A1]**

(b) Angle $ABD = (x + 10)°$ OR Angle $ADB = (2x + 15)°$ **[M1]**

40 **[A1]**

[Back to Question] *[Total Marks: 6]*

947 **(a)** **(i)** 58 **[B1]**

(ii) Alternate segment theorem **[B1]**

(b) Angle $AQB = 180 - 58 - 58 = 64°$ or Angle $BAQ = 58°$ **[M1]**

128 **[A1]**

(c) Angle $ACB = 180 - 64$ (using cyclic quadrilateral $ACBQ$) **[M1]**

116 **[A1]**

[Back to Question] *[Total Marks: 6]*

948 **(a)** States Angle CAB = Angle BCQ because of the alternate segment theorem **[B1]**

$2x - 5 = x + 17$ **[M1]**

$x = 22$ **[A1]**

(b) Identifies angle $ACB = 90°$ (angle in a semicircle) or calculates angle $CAB = 39°$ **[M1]**

51 **[A1]**

(c) Angle $ADC = 180 - 51$ (Opposite angles of cyclic quadrilateral sum to 180°) **[M1]**

129 **[A1]**

[Back to Question] *[Total Marks: 7]*

949 **(a)** Identifies angle ABC = angle CAQ (alternate segment theorem) OR angle ACB = angle PAB **[M1]**

Sets up the equation: $3x - 5 = x + 25$ **[M1]**

$x = 15$ **[A1]**

(b) Calculates angle $ABC = 40°$ and angle $ACB = 40°$ (or uses angles on straight line PQ) **[M1]**

100 **[A1]**

(c) Uses opposite angles of cyclic quadrilateral $ABDC$ sum to 180° $(180 - 100)$ **[M1]**

80 **[A1]**

[Back to Question] *[Total Marks: 7]*

950 **(a)** Angle TAB = angle ACB (alternate segment theorem) **[M1]**

$2y - 10 = 3y - 40$ **[M1]**

30 **[A1]**

(b) $x + 10 = 180 - 2(50)$ **[M1]**

70 **[A1]**

(c) Angle $ADB = 180 - 50$ (opposite angles of cyclic quadrilateral) **[M1]**

130 **[A1]**

[Back to Question] *[Total Marks: 7]*

951 **(a)** **(i)** Angle ACT = Angle $CAT = \dfrac{180 - 48}{2} = 66°$ **[M1]**

Angle ABC = Angle $ACT = 66$ (alternate segment theorem) **[A1]**

(ii) Cyclic quadrilateral $ABCD$, so angle $ADC = 180 - 66$ **[M1]**

114 **[A1]**

(iii) $180 - (114 + 31) = 35$ **[B1]**

(iv) Angle at centre = 2× angle at circumference **[M1]**

$2 \times 66 = 132$ **[A1]**

[Back to Question] *[Total Marks: 7]*

952 **(a)** $180 - 2x$ **[B1]**

(b) $\dfrac{180 - 2x}{2} = 90 - x$ **[B1]**

(c) Angle $ADB = 180 - (90 - x)$ (opposite angles of cyclic quadrilateral) **[M1]**

$90 + x$ **[A1]**

(d) $90 + x = 3(90 - x)$ **[M1]**

$90 + x = 270 - 3x \implies 4x = 180$ **[M1]**

$x = 45$ **[A1]**

[Back to Question] *[Total Marks: 7]*

953 **(a)** **(i)** $2x + 3y = x + 2y + 10$ (from Alternate Segment Theorem) **[B1]**

$3x - y = x + y + 8$ (from Angles in Same Segment) **[B1]**

(ii) Rearranges to linear equations: $x + y = 10$ and $2x - 2y = 8 \implies x - y = 4$ **[M1]**

Valid method to solve simultaneous equations (e.g. adds equations to get $2x = 14$) **[M1]**

$x = 7$ and $y = 3$ **[A1]**

(b) Uses opposite angles of cyclic quadrilateral: angle $ADC = 180° - 95° = 85°$ **[B1]**

Calculates angle $BDC = (7) + (3) + 8 = 18°$ OR angle $BAC = 18°$ **[B1]**

Finds angle $ADB = 85° - 18° = 67°$ and uses alternate segment theorem $\implies$ angle $QAB = 67°$ **[B1]**

[Back to Question] *[Total Marks: 8]*

954 **(a)** Sets up equation for x: $2x + 25 = 4x - 5$ **[M1]**

$x = 15$ **[A1]**

Sets up equation for y: $y + 12 = 3y$ **[M1]**

$y = 6$ **[A1]**

(b) Evaluates angles on straight line: $180 - (4(15) - 5) - 3(6)$ OR evaluates angles in triangle $180 - (2(15) + 25) - (6 + 12)$ **[M1]**

107 **[A1]**

(c) Recognises $ABTC$ is a cyclic quadrilateral and uses $180 - \angle ABT$: $180 - (2(15) + 25)$ **[M1]**

125 **[A1]**

[Back to Question] *[Total Marks: 8]*

955 **(a)** **(i)** Angle $OAT = 90°$ and angle $OBT = 90°$ **[B1]**

Angle $AOB = 180 - x$ **[M1]**

$180 - x = 2(2x - 30)$ **[M1]**

$180 - x = 4x - 60 \implies 5x = 240 \implies x = 48$ **[A1]**

(ii) Angle $AOB = 180 - 48 = 132°$ **[M1]**

Angle $OAB = (180 - 132) \div 2 = 24$ **[A1]**

(b) Angle $ACB = 2(48) - 30 = 66°$ **[M1]**

Angle $ADB = 180 - 66 = 114$ (opposite angles of cyclic quad) **[A1]**

[Back to Question] *[Total Marks: 8]*

956 **(a)** Triangle PAB is isosceles ($PA = PB$) **[M1]**

$\frac{180 - 4x}{2}$ **[M1]**

$90 - 2x$ **[A1]**

(b) Angle ACB = Angle PAB (Alternate segment theorem) **[M1]**

$3x - 10 = 90 - 2x$ **[M1]**

$5x = 100 \implies x = 20$ **[A1]**

(c) $x = 20 \implies$ Angle $APB = 80°$ **[M1]**

Angles in quadrilateral $PAOB$ sum to 360°, with $OAP = OBP = 90°$ **[M1]**

$360 - 90 - 90 - 80 = 100$ **[A1]**

(d) Angle $ACB = 3(20) - 10 = 50°$ **[M1]**

$180 - 50$ (opposite angles of cyclic quad sum to 180°) **[M1]**

130 **[A1]**

[Back to Question] *[Total Marks: 12]*

957 **(a)** Recognises that tangents from an external point are equal in length, so triangle PAB is isosceles: $180 - 44 = 136$ **[M1]**

$136 \div 2 = 68$ **[A1]**

(b) Recognises angle between tangent and radius is 90°, setting up sum of angles in quadrilateral $OAPB$: $360 - 90 - 90 - 44$ **[M1]**

136 **[A1]**

[Back to Question] *[Total Marks: 4]*

958 **(a)** Tangents from an external point are equal in length **[B1]**

(b) 70 **[B1]**

(c) Angle $OAP = 90°$ OR Angle $OBP = 90°$ **[M1]**

140 **[A1]**

[Back to Question] *[Total Marks: 4]*

959 **(a)** Angle $PTO = 56 \div 2 = 28°$ **[M1]**

$180 - 90 - 28 = 62$ **[A1]**

(b) Tangents from an external point are equal in length (so $TP = TQ$) **[B1]**

(c) 62 **[B1]**

[Back to Question] *[Total Marks: 4]*

960 **(a)** Uses angle sum of quadrilateral $OATB$ with angle $OAT = 90°$ and angle $OBT = 90°$: $360° - 90° - 90° - 44°$ OR $180° - 44°$ **[M1]**

136 **[A1]**

(b) Recognises triangle OAB is isosceles ($OA = OB$) and uses $(180° - 136°) \div 2$ **[M1]**

22 **[A1]**

(c) Tangents from an external point are equal in length **[B1]**

[Back to Question] *[Total Marks: 5]*

961 **(a)** 24 **[B1]**

(b) Identifies $\angle OAP = 90°$ (tangent perpendicular to radius) **[M1]**

$OP^2 = 7^2 + 24^2 = 625 \implies OP = 25$ **[A1]**

(c) Let M be the intersection of OP and AB. Identifies $\triangle OAM \sim \triangle OPA$ OR uses Area of $\triangle OAP$ OR uses $\cos(\angle AOP)$ **[M1]**

$\cos(\angle AOP) = \dfrac{7}{25}$ and $OM = 7 \times \cos(\angle AOP)$ **[M1]**

$OM = \dfrac{49}{25}$ or 1.96 **[A1]**

[Back to Question] *[Total Marks: 6]*

962 **(a)** $x + 3 = 3x - 5$ **[M1]**

$x = 4$ **[A1]**

(b) Perpendicular distance = 7 (can be implied) **[B1]**

$\sqrt{25^2 - 7^2}$ OR 24 **[M1]**

48 **[A1]**

[Back to Question] *[Total Marks: 5]*

963 **(a)** Angle $OMB = 90°$ (line from centre to midpoint of chord is perpendicular to chord) **[B1]**

$MB = 7$ **[B1]**

$OM = \sqrt{8.5^2 - 7^2}$ **[M1]**

4.77 or 4.769... **[A1]**

(b) Half chord $= \sqrt{8.5^2 - 3^2}$ **[M1]**

$2 \times \sqrt{63.25} = 15.9$ or 15.90... **[A1]**

[Back to Question] *[Total Marks: 5]*

964 **(a)** Identifies right-angled triangle OAM (or OBM) and states $AM = 24 \div 2 = 12$ **[B1]**

Applies Pythagoras' theorem: $OM^2 + 12^2 = 13^2$ **[M1]**

$OM = \sqrt{169 - 144} = \sqrt{25} = 5$ **[A1]**

(b) 24 (using property that chords equidistant from the centre are equal in length) **[B1]**

(c) $\frac{1}{2} \times 24 \times 5$ **[M1]**

60 **[A1]**

[Back to Question] *[Total Marks: 6]*

965 **(a)** States or uses $OM \perp AB$ (perpendicular bisector of chord passes through centre), so $AM = 7$ **[B1]**

Uses Pythagoras' Theorem: $OM = \sqrt{25^2 - 7^2} = \sqrt{625 - 49} = \sqrt{576} = 24$ **[B1]**

(b) States $ON = OM = 24$ (equal chords are equidistant from centre) **[B1]**

Uses Cosine rule: $MN^2 = 24^2 + 24^2 - 2(24)(24)\cos 100°$ OR uses $MN = 2 \times 24 \times \sin 50°$ **[M1]**

36.8 (accept 36.77...) **[A1]**

(c) $\frac{1}{2} \times 24 \times 24 \times \sin 100°$ **[M1]**

284 (accept 283.6...) **[A1]**

[Back to Question] *[Total Marks: 7]*

966 **(a)** Forms a right-angled triangle with hypotenuse 13 and base $\frac{24}{2} = 12$ **[M1]**

Uses Pythagoras' theorem: $\sqrt{13^2 - 12^2}$ **[M1]**

5 **[A1]**

(b) 5 **[B1]**

(c) Uses Pythagoras' theorem: $\sqrt{13^2 - 12^2}$ **[M1]**

Half of chord $EF = 5$ **[A1]**

10 **[A1]**

(d) The perpendicular from the centre to a chord bisects the chord (accept "perpendicular bisector of a chord passes through the centre") **[B1]**

[Back to Question] *[Total Marks: 8]*

967 **(a)** Angle $OTA = 90°$ (tangent perpendicular to radius) **[B1]**

$AT = 12$ (perpendicular from centre to chord bisects chord) **[B1]**

In triangle OTA, $r^2 + 12^2 = R^2$ (Pythagoras theorem) **[M1]**

$R^2 - r^2 = 144$ **[A1]**

(b) Area $= \pi R^2 - \pi r^2 = \pi(R^2 - r^2)$ **[M1]**

$k = 144$ **[A1]**

(c) Distance $= r$ (or same distance as chord AB) **[B1]**

Equal chords are equidistant from the centre **[B1]**

[Back to Question] *[Total Marks: 7]*

968 **(a)** The perpendicular bisector of a chord passes through the centre of the circle **[B1]**

(b) $MX = 16 \div 2 = 8$ **[B1]**

$OX = \sqrt{10^2 - 8^2} = 6$ **[M1]**

$10 - 6 = 4$ **[A1]**

(c) Distance of PQ from $O = 11 - 6 = 5$ **[B1]**

Half length of $PQ = \sqrt{10^2 - 5^2}$ **[M1]**

$\sqrt{75}$ or 8.66... **[A1]**

$2 \times \sqrt{75} = 17.3$ (17.32...) **[A1]**

[Back to Question] *[Total Marks: 8]*

969 **(a)** Identifies $\angle OAT = 90°$ and $\angle AOT = 90° - x$ **[M1]**

Uses $AM = r\sin(90° - x) = r\cos x$ and perpendicular bisector property $AB = 2AM$ to get $2r\cos x$ **[A1]**

(b) $r\sin x$ (equal chords are equidistant from centre, distance $= OM = r\cos(90° - x)$) **[B1]**

(c) Area of $OAB = \dfrac{1}{2} \times AB \times OM = \dfrac{1}{2}(2r\cos x)(r\sin x) = r^2 \sin x \cos x$ **[M1]**

$TM = \dfrac{AM}{\tan x} = \dfrac{r\cos x}{\tan x} = \dfrac{r\cos^2 x}{\sin x}$ **[M1]**

Area of $ATB = \dfrac{1}{2} \times AB \times TM = \dfrac{1}{2}(2r\cos x)\left(\dfrac{r\cos^2 x}{\sin x}\right) = \dfrac{r^2\cos^3 x}{\sin x}$ **[M1]**

Equates areas: $r^2 \sin x \cos x = \dfrac{r^2\cos^3 x}{\sin x} \implies \sin^2 x \cos x = \cos^3 x \implies \tan^2 x = 1$ **[M1]**

45 **[A1]**

[Back to Question] *[Total Marks: 8]*

970 **(a)** Tangents from an external point are equal in length **[B1]**

(b) Recognises $BP = BQ$, $CR = CQ$, and $DR = DS$ **[M1]**

$AB + CD = AP + PB + CR + RD$ **[M1]**

Substitutes equal tangent segments:
$AS + BQ + CQ + DS = (AS + DS) + (BQ + CQ) = AD + BC$ **[A1]**

(c) Substitutes expressions into the relationship from part (b):
$(2x + 5) + (x + 8) = (2x + 1) + (3x - 2)$ **[M1]**

Correctly expands and simplifies to a linear equation, e.g., $3x + 13 = 5x - 1$ **[M1]**

7 **[A1]**

(d) Substitutes $x = 7$ into the lengths of the sides (at least two) OR uses $2(3x + 13)$ **[M1]**

68 **[A1]**

[Back to Question] *[Total Marks: 9]*

971 **(a)** Recognises angle $OPT = 90°$ and sets up Pythagoras: $TP^2 + 7^2 = 25^2$ **[M1]**

$TP = \sqrt{625 - 49} = \sqrt{576} = 24$ **[A1]**

(b) Uses trigonometry in triangle OPT: $\sin(\text{angle } PTO) = \frac{7}{25}$ or $\tan(\text{angle } PTO) = \frac{7}{24}$ **[M1]**

Angle $PTO = 16.26°$ **[A1]**

Angle $PTQ = 2 \times 16.26° = 32.5°$ (accept $32.52°$) **[A1]**

(c) Calculates total area of kite $OPTQ = 2 \times (\frac{1}{2} \times 7 \times 24) = 168$ **[M1]**

Calculates angle $POQ = 360 - 90 - 90 - 32.52 = 147.48°$ (or uses $180 - 32.52$) **[M1]**

Calculates area of sector $OPQ = \frac{147.48}{360} \times \pi \times 7^2 = 63.07\ldots$ **[M1]**

Shaded area $= 168 - 63.07 = 104.93$ (accept 104.9 to 105.0) **[A1]**

[Back to Question] *[Total Marks: 9]*

972 **(a)** 5.2 **[B1]**

(b) 4700 **[B1]**

(c) 850 **[B1]**

[Back to Question] *[Total Marks: 3]*

973 **(a)** $80 \times 45 = 3600$ **[M1]**

0.36 **[A1]**

(b) 3200 **[B1]**

[Back to Question] *[Total Marks: 3]*

974 **(a)** 35000 **[B1]**

(b) $60 \times 45 \times 30$ OR 2700×30 OR 81000 **[M1]**

their $81000 \div 1000$ **[M1]**

81 **[A1]**

[Back to Question] *[Total Marks: 4]*

975 **(a)** Converts units correctly to cm: 250 mm = 25 cm AND 0.2 m = 20 cm **[B1]**

20000 **[A1]**

(b) 20 **[B1]**

(c) Uses Mass = Density × Volume: 0.6×20000 (= 12000 g) **[M1]**

12 **[A1]**

[Back to Question] *[Total Marks: 5]*

976 **(a)** Converts all dimensions to cm: 250, 140, 0.8 **[M1]**

$250 \times 140 \times 0.8 = 28000$ **[A1]**

(b) Multiplies volume by 7.8: $28000 \times 7.8 = 218400$ **[M1]**

218.4 **[A1]**

[Back to Question] *[Total Marks: 4]*

977 **(a)** Converts dimensions to consistent units, e.g. 150 cm, 80 cm, 60 cm OR 1.5 m, 0.8 m, 0.6 m **[M1]**

$150 \times 80 \times 60 = 720000$ OR $1.5 \times 0.8 \times 0.6 = 0.72$ **[M1]**

720 **[A1]**

(b) $0.024 \times 1000000 = 24000$ **[M1]**

Mass $= 24000 \times 7.8 = 187200$ **[M1]**

187.2 **[A1]**

[Back to Question] *[Total Marks: 6]*

978 **(a)** Conversion of units: 15 $mm = 0.015$ m OR 15 $m = 1500$ cm etc. **[M1]**

Volume $= 15 \times 12 \times 0.015$ (or eq) **[M1]**

2700 **[A1]**

(b) Use of consistent units for tank volume equation (e.g. $r = 0.6$ m and $V = 2.7$ m^3 OR $V = 2700000$ cm^3 and $r = 60$ cm) **[M1]**

$\pi \times 0.6^2 \times h = 2.7$ OR $\pi \times 60^2 \times h = 2700000$ **[M1]**

$h = \dfrac{2.7}{0.36\pi}$ (m) OR $h = \dfrac{2700000}{3600\pi}$ (cm) **[M1]**

238.7 **[A1]**

[Back to Question] *[Total Marks: 7]*

979 **(a)** $\dfrac{4}{3} \times \pi \times 15^3$ OR 4500π OR 14137 **[M1]**

their 14137 $\div 1000000$ OR $\dfrac{4}{3} \times \pi \times 0.15^3$ **[M1]**

1.41×10^{-2} **[A1]**

(b) 125 kg $= 125000$ g **[B1]**

$\dfrac{125000}{4500\pi}$ OR $\dfrac{125000}{14137}$ **[M1]**

8.84 (8.841 . . .) **[A1]**

(c) **(i)** 24 mm $= 2.4$ cm **[B1]**

$\dfrac{1}{3} \times \pi \times 2.4^2 \times 5.5$ **[M1]**

33.2 (33.17 . . .) OR 10.56π **[A1]**

(ii) $\dfrac{4500\pi}{\text{their } 33.17}$ OR $426.1\ldots$ **[M1]**

426 **[A1]**

[Back to Question] *[Total Marks: 11]*

980 **(a)** Identifies consistent units for radius and length (e.g. $r = 1.5$ cm and length $= 80$ cm) **[M1]**

Volume in one second $= \pi \times 1.5^2 \times 80 = 180\pi\ cm^3$ $(= 565.48\ldots\ cm^3)$ **[M1]**

Converts to litres $(180\pi \div 1000)$ and confirms $0.56548\cdots \approx 0.565$ **[A1]**

(b) Volume of pool required $= 4 \times 2.5 \times 1.2 = 12\ m^3$ OR 12000 litres **[M1]**

Time in seconds $= 12000 \div 0.56548\ldots$ OR $12000 \div \dfrac{180\pi}{1000}$ **[M1]**

$= 21220.6\ldots$ seconds $(353.6\ldots$ minutes$)$ **[A1]**

5 hours 54 minutes **[A1]**

[Back to Question] *[Total Marks: 7]*

981 **(a)** Converts speed to cm/min: $5.4 \times 1000 \times 100 \div 60 = 9000$ **[M1]**

Calculates volume in 1 minute: $\pi \times 2.5^2 \times 9000 = 56250\pi\ cm^3$ **[M1]**

Converts cm^3 to litres and evaluates: $56250\pi \div 1000 = 176.714\ldots$ **[A1]**

(b) Calculates total volume in 45 minutes: $56250\pi \times 45 = 2531250\pi\ cm^3$ (or uses litres: $176.714\cdots \times 45 = 7952.1\ldots$ litres) **[M1]**

Converts tank base dimensions to cm and finds area: $300 \times 200 = 60000\ cm^2$ **[M1]**

Sets up equation for depth: Depth $= \dfrac{2531250\pi}{60000}$ **[M1]**

133 (or 132.5 to 132.6) **[A1]**

[Back to Question] *[Total Marks: 7]*

982 **(a)** Cross-sectional area $= \pi \times 2.5^2$ $(= 6.25\pi)$ **[M1]**

Speed conversion: $18\ km/h = \dfrac{18 \times 1000 \times 100}{60}\ cm/min$ $(= 30000\ cm/min)$ **[M1]**

Volume per minute $= 6.25\pi \times 30000$ $(= 187500\pi\ cm^3)$ **[M1]**

Convert to litres: $\dfrac{187500\pi}{1000}$ **[M1]**

589 or 589.0 to 589.1 (187.5π) **[A1]**

(b) Capacity of reservoir $= 50 \times 30 \times 4 = 6000\ m^3$ OR 6000000 litres **[M1]**

Time in minutes $= \dfrac{6000000}{187.5\pi}$ (or $6000000 \div$ their (a)) $(= 10185.9...)$ **[M1]**

Time in hours $= \dfrac{10185.9...}{60}$ $(= 169.76...)$ **[M1]**

$169.76... \div 24 = 7.07...$ days so 7 full days **[M1]**

$0.07... \times 24 = 1.76...$ hours (or $169.76... - 7 \times 24$) **[M1]**

7 **[A1]**

2 **[A1]**

[Back to Question] *[Total Marks: 12]*

983 **(a)** $\pi \times 2.5^2 \times 160$ OR $\pi \times 0.025^2 \times 1.6$ **[M1]**

$\times 60$ **[M1]**

$\div 1000$ (if using cm) OR $\times 1000$ (if using m) **[M1]**

188 (188.4 . . .) **[A1]**

(b) Volume of tank $= 4.2 \times 3.5 \times 2.4 = 35.28\ m^3$ **[M1]**

$35.28 \times 1000 = 35280$ litres **[M1]**

Time $= \dfrac{35280}{\text{their } 188.49}$ OR $\dfrac{35280}{60\pi}$ **[M1]**

187 minutes (187.1 . . .) **[A1]**

3 hours 7 minutes **[A1]**

(c) Calculates lateral area correctly: $2 \times 4.2 \times 2.4 + 2 \times 3.5 \times 2.4$ OR $36.96\ m^2$ [M2]
(If incorrect, M1 for one correct face area e.g., 4.2×2.4)

$36.96 \times 10000 = 369600\ cm^2$ OR $45000 \div 10000 = 4.5\ m^2$ **[M1]**

Litres of paint required $= \dfrac{369600}{45000}$ OR $\dfrac{36.96}{4.5}$ **[M1]**

Number of tins $= \dfrac{8.21\ldots}{2.5}$ $(= 3.28\ldots)$ **[M1]**

4 **[A1]**

[Back to Question] *[Total Marks: 15]*

984 **(a)** Calculates area of triangle: $\dfrac{1}{2} \times 12 \times 8 = 48$ **[M1]**

$4x = 48$

x = 12[*A*1]

(b) Uses Pythagoras to find slanted side: $\sqrt{6^2 + 8^2}$ **[M1]**

Slanted side = 10 **[A1]**

Perimeter $= 12 + 10 + 10 = 32$ **[A1]**

[Back to Question] *[Total Marks: 5]*

985 **(a)** $15 \times h = 120$ **[M1]**

8 **[A1]**

(b) $2(15 + AD) = 54$ OR $15 + AD = 27$ **[M1]**

12 **[A1]**

(c) 60 **[B1]**

[Back to Question] *[Total Marks: 5]*

986 **(a)** $2(2x+3)+2(x-1)=34$ OR $2x+3+x-1=17$ **[M1]**

$6x+4=34$ OR $3x+2=17$ **[M1]**

$x=5$ **[A1]**

(b) Substitution of $x=5$: $(2(5)+3)\times(5-1)$ OR 13×4 **[M1]**

52 **[A1]**

[Back to Question] *[Total Marks: 5]*

987 **(a)** $\frac{1}{2}(10+16)\times 8$ **[M1]**

104 **[A1]**

(b) Base of right-angled triangle $=16-10=6$ **[M1]**

Slanted edge $=\sqrt{6^2+8^2}=10$ **[M1]**

$8+10+16+10=44$ **[A1]**

[Back to Question] *[Total Marks: 5]*

988 **(a)** Uses Area $=\frac{1}{2}\times$ base $\times$ height $=\frac{1}{2}\times 14\times 9$ **[M1]**

63 **[A1]**

(b) Finds $AE=\sqrt{9^2+9^2}=\sqrt{162}$ or $12.7\ldots$ **[M1]**

Finds $BE=\sqrt{5^2+9^2}=\sqrt{106}$ or $10.2\ldots$ **[M1]**

Perimeter $=14+\sqrt{162}+\sqrt{106}$ **[M1]**

37.0 (or $37.02\ldots$) **[A1]**

[Back to Question] *[Total Marks: 6]*

989 **(a)** $\frac{1}{2}\times(15+24)\times 8$ **[M1]**

156 **[A1]**

(b) Horizontal distance from D to the altitude from $A=\sqrt{10^2-8^2}=6$ **[M1]**

Horizontal distance for the right-angled triangle under $B=24-15-6=3$ **[M1]**

$BC=\sqrt{8^2+3^2}$ $(=\sqrt{73}\approx 8.544)$ **[M1]**

57.5 OR 57.54 to 57.55 **[A1]**

[Back to Question] *[Total Marks: 6]*

990 **(a)** Area of rectangle $=4(2x+3)$ OR $8x+12$ **[M1]**

Area of triangle $= \frac{1}{2} \times 6 \times (3x - 1)$ OR $9x - 3$ **[M1]**

$8x + 12 = 9x - 3$ **[M1]**

$x = 15$ **[A1]**

(b) Length $= 2(15) + 3 = 33$ **[M1]**

Perimeter $= 2(4 + 33) = 74$ **[A1]**

[Back to Question] *[Total Marks: 6]*

991 **(a)** $\frac{1}{2} \times 8 \times ((x + 4) + (3x - 2)) = 120$ **[M1]**

$4(4x + 2) = 120$ OR $4x + 2 = 30$ **[M1]**

7 **[A1]**

(b) Evaluates parallel sides: 11 and 19 **[B1]**

Uses Pythagoras' theorem to find the slanted side: $\sqrt{8^2 + \left(\frac{19 - 11}{2}\right)^2}$ OR $\sqrt{8^2 + 4^2}$ [M2]

(M1 for finding the horizontal base of 4)

47.9 (47.88...) **[A1]**

[Back to Question] *[Total Marks: 7]*

992 **(a)** **(i)** Sets up equation for area: $\frac{1}{2}((x + 4) + (3x - 2))(x - 1) = 20$ **[M1]**

Simplifies expression to $\frac{1}{2}(4x + 2)(x - 1) = 20$ or $(2x + 1)(x - 1) = 20$ **[M1]**

Expands to $2x^2 - 2x + x - 1 = 20$ leading to $2x^2 - x - 21 = 0$ with no errors seen **[A1]**

(ii) Factorises $(2x - 7)(x + 3) = 0$ or uses quadratic formula to find $x = 3.5$ **[M1]**

Calculates straight sides: $AD = 7.5$, $BC = 8.5$, $AB = 2.5$ **[M1]**

Sets up Pythagoras for the slanted side CD: $\sqrt{(8.5 - 7.5)^2 + 2.5^2}$ **[M1]**

Calculates $CD = \sqrt{1^2 + 2.5^2} = \sqrt{7.25}$ (or 2.69...) **[A1]**

Perimeter $= 7.5 + 8.5 + 2.5 + 2.69\cdots = 21.2$ (or 21.19...) **[A1]**

[Back to Question] *[Total Marks: 8]*

993 **(a)** $\frac{1}{2}(x + x + 4)(x - 1) = 40$ **[M1]**

$\frac{1}{2}(2x + 4)(x - 1) = 40$ OR $(x + 2)(x - 1) = 40$ **[M1]**

$x^2 + 2x - x - 2 = 40 \implies x^2 + x - 42 = 0$ **[A1]**

(b) $(x + 7)(x - 6) = 0$ **[M1]**

6 **[A1]**

(c) Identifies lengths are Top = 6, Bottom = 10, Height = 5 **[M1]**

Horizontal difference $= 4$, so slanted side $= \sqrt{5^2 + 4^2}$ OR $\sqrt{41}$ **[M1]**

Perimeter $= 5 + 6 + 10 + \sqrt{41}$ **[M1]**

27.4 (or 27.40...) **[A1]**

[Back to Question] *[Total Marks: 9]*

994 **(a)** Area of rectangle $= (2x + 1)(x + 3) = 2x^2 + 6x + x + 3$ **[M1]**

Area of triangle $= \frac{1}{2}(3x + 2)(2x - 1) = \frac{1}{2}(6x^2 - 3x + 4x - 2)$ **[M1]**

Equates areas: $2x^2 + 7x + 3 = \frac{1}{2}(6x^2 + x - 2)$ **[M1]**

Multiplies by 2 and simplifies correctly to $2x^2 - 13x - 8 = 0$ with no errors **[A1]**

(b) $\frac{-(-13) \pm \sqrt{(-13)^2 - 4(2)(-8)}}{2(2)}$ [M2]

(M1 for correct substitution into quadratic formula)

$\frac{13 \pm \sqrt{233}}{4}$ **[M1]**

7.07 and -0.57 **[A1]**

(c) Substitutes the positive value of x into $2(2x + 1 + x + 3)$ OR calculates sides 15.14 and 10.07 **[M1]**

50.4 (accept 50.39 to 50.43) **[A1]**

[Back to Question] *[Total Marks: 10]*

995 **(a)** Perimeter of rectangle $= 6y$ OR Perimeter of square $= 60 - 6y$ **[M1]**

$\frac{60 - 6y}{4}$ OR $15 - 1.5y$ **[A1]**

(b) $(15 - 1.5y)^2 + y(2y) = 113$ **[M1]**

$225 - 45y + 2.25y^2 + 2y^2 = 113$ **[M1]**

$4.25y^2 - 45y + 112 = 0$ leading to $17y^2 - 180y + 448 = 0$ with no errors seen **[A1]**

(c) $\frac{-(-180) \pm \sqrt{(-180)^2 - 4(17)(448)}}{2(17)}$ **[M1]**

$\frac{180 \pm \sqrt{1936}}{34}$ OR $\frac{180 \pm 44}{34}$ **[M1]**

$y = 4$ and $y = 6.59$ (or 6.588... or $\frac{112}{17}$) **[A1]**

(d) Identifies $y = 4$ as the valid solution (since $15 - 1.5(6.59) < 6.59$) leading to side length $= 9$ **[M1]**

81 **[A1]**

[Back to Question] *[Total Marks: 10]*

996 **(a)** $\frac{45}{360} \times 2 \times \pi \times 12$ **[M1]**

3π **[A1]**

(b) $\frac{45}{360} \times \pi \times 12^2$ **[M1]**

18π **[A1]**

[Back to Question] *[Total Marks: 4]*

997 **(a)** $\frac{60}{360} \times 2 \times \pi \times 12$ **[M1]**

4π **[A1]**

(b) $\frac{60}{360} \times \pi \times 12^2$ **[M1]**

24π **[A1]**

[Back to Question] *[Total Marks: 4]*

998 **(a)** $\frac{45}{360} \times \pi \times 12^2$ **[M1]**

18π **[A1]**

(b) $\frac{45}{360} \times 2 \times \pi \times 12$ **[M1]**

3π **[A1]**

$3\pi + 24$ **[A1]**

[Back to Question] *[Total Marks: 5]*

999 **(a)** $\frac{150}{360} \times \pi \times 6^2$ **[M1]**

15π **[A1]**

(b) $\frac{150}{360} \times 2 \times \pi \times 6$ **[M1]**

5π **[A1]**

$5\pi + 12$ **[A1]**

[Back to Question] *[Total Marks: 5]*

1000 **(a)** $\frac{\theta}{360} \times \pi r^2 = 10\pi$ **[M1]**

$\frac{\theta}{360} \times 2\pi r = 2\pi$ **[M1]**

Valid method to solve simultaneously (e.g. dividing the area equation by the arc length equation to get $\frac{r}{2} = 5$) **[M1]**

$r = 10$ **[A1]**

$\theta = 36$ **[A1]**

(b) $\frac{x+20}{360} \times 2\pi x = \frac{16\pi}{5}$ **[M1]**

Simplifying to form a 3-term quadratic equation, e.g. $x^2 + 20x - 576 = 0$ **[M1]**

Valid method to solve the quadratic (e.g. $(x+36)(x-16) = 0$) **[M1]**

$x = 16$ (with -36 rejected or simply 16 as the final answer) **[A1]**

[Back to Question] *[Total Marks: 9]*

1001 **(a)** $\frac{72}{360} \times \pi \times r^2 = 85$ **[M1]**

$r^2 = \frac{85 \times 5}{\pi}$ or $135.28\ldots$ **[M1]**

11.6 or $11.63\ldots$ **[A1]**

(b) Arc length $= \frac{72}{360} \times 2 \times \pi \times 11.63$ or $14.6\ldots$ **[M1]**

Perimeter $= 2 \times 11.63 + 14.61$ **[M1]**

37.9 or 37.86 to 37.87 **[A1]**

[Back to Question] *[Total Marks: 6]*

1002 **(a)** $\frac{250}{360} \times 2 \times \pi \times r = 35$ **[M1]**

$r = \frac{35 \times 360}{250 \times 2 \times \pi}$ **[M1]**

8.02 or 8.021... **[A1]**

(b) $\frac{250}{360} \times \pi \times (\text{their } 8.02)^2$ **[M1]**

OR $\frac{1}{2} \times \text{their } 8.02 \times 35$ **[M1]**

140 or 140.3 to 140.4 **[A1]**

[Back to Question] *[Total Marks: 6]*

1003 **(a)** $\frac{120}{360} \times \pi \times 9^2$ OR $\frac{120}{360} \times \pi \times 5^2$ **[M1]**

$\frac{120}{360} \times \pi \times 9^2 - \frac{120}{360} \times \pi \times 5^2$ **[M1]**

58.6 $(58.64\ldots)$ OR $\frac{56\pi}{3}$ **[A1]**

(b) $\frac{120}{360} \times 2 \times \pi \times 9$ OR $\frac{120}{360} \times 2 \times \pi \times 5$ **[M1]**

$\frac{120}{360} \times 2 \times \pi \times 9 + \frac{120}{360} \times 2 \times \pi \times 5 + 4 + 4$ **[M1]**

37.3 $(37.32\ldots)$ OR $\frac{28\pi}{3} + 8$ **[A1]**

(c) $\pi r^2 = 58.64\ldots$ (or their area from (a)) **[M1]**

4.32 (4.320...) **[A1]**

[Back to Question] *[Total Marks: 8]*

1004 **(a)** $\frac{110}{360} \times \pi \times 8.5^2$ **[M1]**

69.4 (or 69.35...) **[A1]**

(b) Area of triangle $OAB = \frac{1}{2} \times 8.5^2 \times \sin(110°)$ (=33.94...) **[M1]**

Area of segment = $69.35\cdots - 33.94...$ **[M1]**

35.4 (or 35.40...) **[A1]**

(c) Arc length $AB = \frac{110}{360} \times 2 \times \pi \times 8.5$ (=16.31...) **[M1]**

Chord $AB = \sqrt{8.5^2 + 8.5^2 - 2(8.5)(8.5)\cos(110°)}$ OR $2 \times 8.5\sin(55°)$ **[M1]**

Chord $AB = 13.92...$ **[A1]**

Perimeter = $16.31\cdots + 13.92\cdots = 30.2$ (or 30.24...) **[A1]**

[Back to Question] *[Total Marks: 9]*

1005 **(a)** Area of sector $= \frac{110}{360} \times \pi \times 8^2$ (= 61.43...) **[M1]**

Area of triangle $= \frac{1}{2} \times 8 \times 8 \times \sin(110°)$ (= 30.07...) **[M1]**

Area of segment = Area of sector − Area of triangle **[M1]**

31.4 or 31.36... **[A1]**

(b) Arc length $= \frac{110}{360} \times 2 \times \pi \times 8$ (= 15.35...) **[M1]**

Chord $AB = \sqrt{8^2 + 8^2 - 2 \times 8 \times 8 \times \cos(110°)}$ OR $2 \times 8 \times \sin(55°)$ (= 13.11...) **[M1]**

28.5 or 28.46... **[A1]**

[Back to Question] *[Total Marks: 7]*

1006 **(a)** **(i)** $\frac{x}{360} \times 2\pi r = 6\pi$ OR $\frac{x}{360} \times \pi r^2 = 27\pi$ **[M1]**

Valid method to eliminate x, e.g. dividing the two equations: $\frac{\frac{x}{360}\pi r^2}{\frac{x}{360}2\pi r} = \frac{27\pi}{6\pi}$ leading to $\frac{r}{2} = 4.5$ **[M1]**

9 **[A1]**

(ii) Substitutes their r back into an equation, e.g. $\frac{x}{360} \times 2\pi(9) = 6\pi$ **[M1]**

120 **[A1]**

(b) Perimeter of sector $= \frac{y}{360} \times 2\pi R + 2R$ **[M1]**

Equates to circumference: $\frac{y}{360} \times 2\pi R + 2R = 2\pi R$ **[M1]**

Rearranges to isolate terms with y: $\frac{y}{360} \times 2\pi R = 2\pi R - 2R$ **[M1]**

$\frac{360(\pi - 1)}{\pi}$ OR $360 - \frac{360}{\pi}$ **[A1]**

[Back to Question] *[Total Marks: 9]*

1007 **(a)** Uses cosine rule: $\cos(AOB) = \frac{15^2 + 15^2 - 24^2}{2 \times 15 \times 15}$ or $24^2 = 15^2 + 15^2 - 2(15)(15)\cos(AOB)$ **[M1]**

$\cos(AOB) = -0.28$ **[M1]**

$106.26\ldots$ which rounds to $106.3°$ with no errors seen **[A1]**

(Alternative: $\sin(\frac{AOB}{2}) = \frac{12}{15}$ [M1], $\frac{AOB}{2} = 53.13\ldots°$ [M1], $AOB = 106.3°$ [A1])

(b) **(i)** Arc length $= \frac{106.3}{360} \times 2 \times \pi \times 15$ $(= 27.8\ldots)$ **[M1]**

Perimeter $= 24 + 27.8\ldots$ **[M1]**

51.8 (or 51.81 to 51.83) **[A1]**

(ii) Area of sector $= \frac{106.3}{360} \times \pi \times 15^2$ $(= 208.6\ldots$ to $208.7\ldots)$ **[M1]**

Area of triangle $= \frac{1}{2} \times 15 \times 15 \times \sin(106.3°)$ $(= 108.0\ldots)$ **[M1]**

Subtracts triangle area from sector area: $208.7 - 108.0$ **[M1]**

101 or 100.6 to 100.8 **[A1]**

[Back to Question] *[Total Marks: 10]*

1008 **(a)** $\frac{4}{3} \times \pi \times 3^3$ **[M1]**

36π **[A1]**

(b) $4 \times \pi \times 3^2$ **[M1]**

36π **[A1]**

[Back to Question] *[Total Marks: 4]*

1009 **(a)** Volume of cylinder $= \pi \times 3^2 \times 10$ $(= 90\pi)$ **[M1]**

Volume of hemisphere $= \frac{2}{3} \times \pi \times 3^3$ $(= 18\pi)$ **[M1]**

108π **[A1]**

(b) Curved surface area of cylinder $= 2 \times \pi \times 3 \times 10$ $(= 60\pi)$ **[M1]**

Area of circular base $= \pi \times 3^2$ $(= 9\pi)$ AND Curved surface area of hemisphere $= 2 \times \pi \times 3^2$ $(= 18\pi)$ **[M1]**

87π **[A1]**

[Back to Question] *[Total Marks: 6]*

1010 **(a)** Area of base $= \pi \times 6^2 = 36\pi$ **[M1]**

Curved surface area $= \pi \times 6 \times l = 6\pi l$ **[M1]**

$36\pi + 6\pi l = 96\pi$ leading to $6l = 60$ so $l = 10$ **[A1]**

(b) $h^2 + 6^2 = 10^2$ **[M1]**

$h = \sqrt{100 - 36} = \sqrt{64} = 8$ **[A1]**

Volume $= \frac{1}{3} \times \pi \times 6^2 \times 8$ **[M1]**

96π **[A1]**

[Back to Question] *[Total Marks: 7]*

1011 **(a)** Use of Pythagoras to find the height of the cone: $\sqrt{5^2 - 3^2}$ $(= 4)$ **[M1]**

Volume of cone $= \frac{1}{3} \times \pi \times 3^2 \times 4$ $(= 12\pi)$ **[M1]**

Volume of hemisphere $= \frac{1}{2} \times \frac{4}{3} \times \pi \times 3^3$ $(= 18\pi)$ **[M1]**

30π **[A1]**

(b) Curved surface area of cone $= \pi \times 3 \times 5$ $(= 15\pi)$ **[M1]**

Surface area of hemisphere $= 2 \times \pi \times 3^2$ $(= 18\pi)$ **[M1]**

33π **[A1]**

[Back to Question] *[Total Marks: 7]*

1012 **(a)** $\frac{1}{2} \times 6 \times 8 \times 15$ **[M1]**

360 **[A1]**

(b) Area of two triangular ends $= \frac{1}{2} \times 6 \times 8 \times 2$ $(= 48)$ **[M1]**

Area of three rectangular faces $= 15 \times (6 + 8 + 10)$ or $(15 \times 6) + (15 \times 8) + (15 \times 10)$ $(= 360)$ **[M1]**

408 **[A1]**

[Back to Question] *[Total Marks: 5]*

1013 **(a)** Curved surface area $= 2 \times \pi \times 5 \times 12$ **[M1]**

Total surface area $= 2 \times \pi \times 5 \times 12 + 2 \times \pi \times 5^2$ **[M1]**

534 or 534.07... or 170π **[A1]**

(b) Volume of cylinder $= \pi \times 5^2 \times 12 = 300\pi$ (942...) **[M1]**

$\frac{4}{3}\pi R^3 = 300\pi$ **[M1]**

$R = \sqrt[3]{225} = 6.08$ or 6.082... **[A1]**

[Back to Question] *[Total Marks: 6]*

1014 **(a)** $\frac{1}{3} \times 10^2 \times h = 400$ **[M1]**

12 **[A1]**

(b) Distance from centre of base to midpoint of side = 5 cm **[B1]**

Slant height of a triangular face = $\sqrt{12^2 + 5^2}$ (= 13) **[M1]**

Area of one triangular face = $\frac{1}{2} \times 10 \times 13$ (= 65) **[M1]**

360 **[A1]**

[Back to Question] *[Total Marks: 6]*

1015 **(a)** Height of cone = 24 − 12 = 12 cm **[B1]**

Volume of cylinder = $\pi \times 5^2 \times 12$ (=300π or 942.4...) **[M1]**

Volume of cone = $\frac{1}{3} \times \pi \times 5^2 \times 12$ (=100π or 314.1...) **[M1]**

1260 (or 1256 to 1257 or 400π) **[A1]**

(b) Slant height of cone (l) = $\sqrt{5^2 + 12^2} = 13$ **[M1]**

Curved surface area of cone = $\pi \times 5 \times 13$ (=65π or 204.2...) **[M1]**

Curved surface area of cylinder = $2 \times \pi \times 5 \times 12$ (=120π or 376.9...) **[M1]**

Area of circular base = $\pi \times 5^2$ (=25π or 78.5...) **[M1]**

660 (or 659.7... or 210π) **[A1]**

[Back to Question] *[Total Marks: 9]*

1016 **(a)** $\pi \times (x)^2 \times 3x$ **[M1]**

$3\pi x^3$ **[A1]**

(b) $\frac{4}{3} \times \pi \times \left(\frac{x}{2}\right)^3$ **[M1]**

$\frac{\pi x^3}{6}$ **[A1]**

(c) 18 **[B1]**

(d) Total surface area = $2\pi x^2 + 2\pi x(3x)$ **[M1]**

$8\pi x^2 = 140$ **[M1]**

$x = 2.36$ (or 2.360...) **[A1]**

[Back to Question] *[Total Marks: 8]*

1017 **(a)** Area of trapezium = $\frac{1}{2} \times (15 + 25) \times 18$ **[M1]**

360 **[A1]**

Volume = $360 \times 120 = 43200$ cm^3 **[M1]**

43.2 **[A1]**

(b) Volume of water after 30 seconds $= 0.8 \times 30 = 24$ litres $= 24000$ cm^3 **[M1]**

Let depth be h and the width of water surface be w.

$w = 15 + \frac{10}{18}h = 15 + \frac{5}{9}h$ **[M2]**

Area of cross-section of water $= \frac{1}{2} \times (15 + 15 + \frac{5}{9}h) \times h = 15h + \frac{5}{18}h^2$ **[M1]**

Volume $= (15h + \frac{5}{18}h^2) \times 120 = 24000$

$1800h + \frac{100}{3}h^2 = 24000$

$h^2 + 54h - 720 = 0$ **[M1]**

$(h - 11.23...)(h + 65.23...) = 0$

$h = 11.2$ or $11.23...$ **[A1]**

[Back to Question] *[Total Marks: 9]*

1018 **(a)** Volume of large cone $= \frac{1}{3} \times \pi \times 10^2 \times 24$ $(= 800\pi)$ **[M1]**

Volume of small cone $= \frac{1}{3} \times \pi \times 5^2 \times 12$ $(= 100\pi)$ **[M1]**

2200 or 2199.1 to 2199.2 **[A1]**

(b) Slant height of large cone $= \sqrt{10^2 + 24^2}$ $(= 26)$ AND Slant height of small cone $= \sqrt{5^2 + 12^2}$ $(= 13)$ **[M1]**

Curved surface area of frustum $= \pi \times 10 \times 26 - \pi \times 5 \times 13$ $(= 195\pi)$ **[M1]**

Total surface area $= 195\pi + \pi \times 10^2 + \pi \times 5^2$ **[M1]**

1010 or 1005.3 . . . **[A1]**

(c) Volume of one sphere $= \frac{4}{3} \times \pi \times 3^3$ $(= 36\pi$ or $113.1...)$ **[M1]**

$\frac{700\pi}{36\pi}$ or $\frac{2199.1}{113.1}$ **[M1]**

19 **[A1]**

[Back to Question] *[Total Marks: 10]*

1019 **(a)** $\frac{1}{3} \times 12^2 \times 10$ **[M1]**

480 **[A1]**

(b) Use of Pythagoras to find slant height: $\sqrt{10^2 + 6^2}$ **[M1]**

$\sqrt{136}$ or 11.6... or 11.7 **[A1]**

Area of four triangular faces $= 4 \times \frac{1}{2} \times 12 \times$ their $\sqrt{136}$ $(= 279.8...)$ **[M1]**

424 or 423.8... to 423.9 **[A1]**

(c) Volume of one sphere $= \frac{4}{3} \times \pi \times 2.5^3$ $(= 65.4...)$ **[M1]**

$\frac{480}{\text{their } 65.4...}$ **[M1]**

7 **[A1]**

(d) $\frac{1620}{480}$ or 3.375 **[M1]**

$\sqrt[3]{3.375}$ or 1.5 **[M1]**

15 **[A1]**

[Back to Question] *[Total Marks: 12]*

1020 Identifies the three straight edges as $10 + 6 + 10 = 26$ **[M1]**

Calculates the arc length of the semicircle as $\frac{1}{2} \times \pi \times 6$ (or 3π) **[M1]**

$26 + 3\pi$ **[A1]**

[Back to Question] *[Total Marks: 3]*

1021 **(a)** Arc length $BD = \frac{1}{4} \times 2 \times \pi \times 10 = 5\pi$ **[M1]**

Perimeter $= 10 + 10 + 5\pi = 20 + 5\pi$ **[A1]**

(b) Area of quarter circle $= \frac{1}{4} \times \pi \times 10^2 = 25\pi$ **[M1]**

Area of shaded region $= 100 - 25\pi$ **[A1]**

[Back to Question] *[Total Marks: 4]*

1022 **(a)** Curved edge of semi-circle $= \frac{1}{2} \times \pi \times 4 = 2\pi$ **[M1]**

$20 + 2\pi$ **[A1]**

(b) $(8 \times 4) + \left(\frac{1}{2} \times \pi \times 2^2\right)$ **[M1]**

$32 + 2\pi$ **[A1]**

[Back to Question] *[Total Marks: 4]*

1023 **(a)** Perimeter $= 12 + 12 + 8 + \frac{1}{2} \times \pi \times 8$ **[M1]**

$32 + 4\pi$ **[A1]**

(b) Area $= 12 \times 8 + \frac{1}{2} \times \pi \times 4^2$ **[M1]**

$96 + 8\pi$ **[A1]**

[Back to Question] *[Total Marks: 4]*

1024 **(a)** Arc length $= \frac{1}{2} \times \pi \times 8 = 4\pi$ **[M1]**

$32 + 4\pi$ **[A1]**

(b) Area of rectangle $= 12 \times 8 = 96$ **[M1]**

Area of semicircle $= \frac{1}{2} \times \pi \times 4^2 = 8\pi$ **[M1]**

$96 + 8\pi$ **[A1]**

[Back to Question] *[Total Marks: 5]*

1025 **(a)** Area of rectangle $= 10 \times 8 = 80$ **[M1]**

Area of semicircle $= \frac{1}{2} \times \pi \times 4^2 = 8\pi$ **[M1]**

$80 + 8\pi$ **[A1]**

(b) Identifies the three straight edges: $10 + 8 + 10 = 28$ **[M1]**

Perimeter of semicircle arc $= \frac{1}{2} \times \pi \times 8 = 4\pi$ **[M1]**

$28 + 4\pi$ **[A1]**

[Back to Question] *[Total Marks: 6]*

1026 **(a)** Sum of three straight edges: $10 + 6 + 10 = 26$ **[M1]**

Arc length of semicircle $= \frac{1}{2} \times \pi \times 6$ $(= 3\pi)$ **[M1]**

$26 + 3\pi$ **[A1]**

(b) Area of rectangle $= 10 \times 6$ $(= 60)$ **[M1]**

Area of removed semicircle $= \frac{1}{2} \times \pi \times 3^2$ $(= 4.5\pi)$ **[M1]**

$60 - 4.5\pi$ **[A1]**

[Back to Question] *[Total Marks: 6]*

1027 **(a)** Area of square $= 4 \times 4 = 16$ **[B1]**

Area of quarter circle $= \frac{1}{4} \times \pi \times 4^2 = 4\pi$ **[M1]**

$16 + 4\pi$ **[A1]**

(b) Arc length $= \frac{1}{4} \times 2 \times \pi \times 4 = 2\pi$ **[M1]**

Perimeter $= 4 + 4 + 4 + 2\pi$ **[M1]**

$12 + 2\pi$ **[A1]**

[Back to Question] *[Total Marks: 6]*

1028 **(a)** Arc length $= \frac{1}{2} \times \pi \times 10$ $(= 5\pi)$ **[M1]**

Perimeter $= 12 + 12 + 10 + 5\pi$ **[M1]**

$34 + 5\pi$ **[A1]**

(b) Area of rectangle $= 12 \times 10$ $(= 120)$ **[M1]**

Area of semicircle $= \frac{1}{2} \times \pi \times 5^2$ $(= 12.5\pi)$ **[M1]**

$120 + 12.5\pi$ **[A1]**

[Back to Question] *[Total Marks: 6]*

1029 **(a)** Calculates the area of the rectangle: $10 \times 6 = 60$ **[M1]**

Calculates the area of the semicircle: $\frac{1}{2} \times \pi \times 3^2$ $(= 4.5\pi)$ **[M1]**

$60 - 4.5\pi$ **[A1]**

(b) Calculates the sum of the three straight edges: $10 + 6 + 10 = 26$ **[M1]**

Calculates the arc length of the semicircle: $\frac{1}{2} \times \pi \times 6$ $(= 3\pi)$ **[M1]**

$26 + 3\pi$ **[A1]**

[Back to Question] *[Total Marks: 6]*

1030 **(a)** Volume of cylinder $= \pi \times 5^2 \times 12 = 300\pi$ **[M1]**

Volume of hemisphere $= \frac{2}{3} \times \pi \times 5^3 = \frac{250\pi}{3}$ **[M1]**

1200 (or 1204 or 1204.2 . . .) **[A1]**

(b) Curved surface of cylinder $= 2 \times \pi \times 5 \times 12 = 120\pi$ **[M1]**

Curved surface of hemisphere + Base of cylinder $= 2 \times \pi \times 5^2 + \pi \times 5^2 = 75\pi$ **[M1]**

613 (or 612.6 . . .) **[A1]**

[Back to Question] *[Total Marks: 6]*

1031 **(a)** Height of cylinder $= 25 - 5 = 20$ **[B1]**

Total Volume $= \pi \times 5^2 \times 20 + \frac{2}{3} \times \pi \times 5^3$ **[M1]**

1830 or 1832.5 . . . **[A1]**

(b) Curved surface area of cylinder $= 2 \times \pi \times 5 \times 20$ $(= 200\pi)$ **[M1]**

Area of circular base $= \pi \times 5^2$ $(= 25\pi)$ **[M1]**

Curved surface area of hemisphere $= 2 \times \pi \times 5^2$ $(= 50\pi)$ **[M1]**

864 or 863.9 . . . **[A1]**

[Back to Question] *[Total Marks: 7]*

1032 **(a)** Volume of cylinder $= \pi \times 5^2 \times 12$ $(= 300\pi$ or 942.47...) **[M1]**

Volume of cone $= \frac{1}{3} \times \pi \times 5^2 \times 6$ $(= 50\pi$ or 157.07...) **[M1]**

1100 or 1099.5... to 1099.6 **[A1]**

(b) Slant height of cone $= \sqrt{5^2 + 6^2}$ $(= \sqrt{61}$ or 7.81...) **[M1]**

Curved surface area of cone $= \pi \times 5 \times$ their $\sqrt{61}$ $(= 122.68...)$ **[M1]**

Curved surface area of cylinder + base $= (2 \times \pi \times 5 \times 12) + (\pi \times 5^2)$ $(= 145\pi$ or 455.53...) **[M1]**

578 or 578.2... **[A1]**

[Back to Question] *[Total Marks: 7]*

1033 **(a)** Volume of cylinder $= \pi \times 6^2 \times 14$ **[M1]**

Volume of hemisphere $= \frac{2}{3} \times \pi \times 6^3$ **[M1]**

2040 or 2035.7... **[A1]**

(b) Curved surface area of cylinder + base area $= 2 \times \pi \times 6 \times 14 + \pi \times 6^2$ (or $168\pi + 36\pi$) **[M1]**

Curved surface area of hemisphere $= 2 \times \pi \times 6^2$ (or 72π) **[M1]**

Correct addition of all three parts **[M1]**

867 or 867.0... to 867.1 **[A1]**

[Back to Question] *[Total Marks: 7]*

1034 **(a)** Volume of cylinder $= \pi \times 6^2 \times 15 = 540\pi$ **[M1]**

Volume of cone $= \frac{1}{3} \times \pi \times 6^2 \times 8 = 96\pi$ **[M1]**

1998 or 2000 (to 3 sf) **[A1]**

(b) Slant height of cone $l = \sqrt{6^2 + 8^2} = 10$ cm **[M1]**

Curved surface area of cone $= \pi \times 6 \times 10 = 60\pi$ **[M1]**

Curved surface area of cylinder + base area $=$
$2 \times \pi \times 6 \times 15 + \pi \times 6^2 = 180\pi + 36\pi = 216\pi$ **[M1]**

867 (to 3 sf) **[A1]**

[Back to Question] *[Total Marks: 7]*

1035 **(a)** Volume of cylinder $= \pi \times 3^2 \times 10 = 90\pi$ $(282.7...)$ **[M1]**

Volume of hemisphere $= \frac{2}{3} \times \pi \times 3^3 = 18\pi$ $(56.5...)$ **[M1]**

108π or 339 or 339.2 to 339.3 **[A1]**

(b) Area of circular base $= \pi \times 3^2 = 9\pi$ $(28.2...)$ **[B1]**

Curved area of cylinder $= 2 \times \pi \times 3 \times 10 = 60\pi$ $(188.4...)$ **[M1]**

Curved area of hemisphere $= 2 \times \pi \times 3^2 = 18\pi$ $(56.5...)$ **[M1]**

87π or 273 or 273.3 to 273.4 **[A1]**

[Back to Question] *[Total Marks: 7]*

1036 **(a)** $\pi \times 4^2 \times 12$ (or 192π) **[M1]**

$\frac{2}{3} \times \pi \times 4^3$ (or $\frac{128\pi}{3}$) **[M1]**

737 or 737.2... or $\frac{704\pi}{3}$ **[A1]**

(b) Calculates curved surface area of cylinder: $2 \times \pi \times 4 \times 12$ (or 96π) **[M1]**

Calculates curved surface area of hemisphere: $2 \times \pi \times 4^2$ (or 32π) **[M1]**

Adds all three external faces including the flat circular base: $96\pi + 32\pi + \pi(4^2)$ **[M1]**

452 or 452.3... to 452.4 or 144π **[A1]**

[Back to Question] *[Total Marks: 7]*

1037 **(a)** Volume of large cone $= \frac{1}{3} \times \pi \times 8^2 \times 24$ ($=512\pi$) **[M1]**

Volume of small cone $= \frac{1}{3} \times \pi \times 5^2 \times 15$ ($=125\pi$) **[M1]**

1215.8 (or 1220 or 387π) **[A1]**

(b) Slant height of large cone $= \sqrt{8^2 + 24^2}$ ($=8\sqrt{10}$ or $25.298\ldots$) **[M1]**

Slant height of small cone $= \sqrt{5^2 + 15^2}$ ($=5\sqrt{10}$ or $15.811\ldots$) **[M1]**

Curved SA of frustum $= \pi \times 8 \times 8\sqrt{10} - \pi \times 5 \times 5\sqrt{10}$ ($=387.4\ldots$) **[M1]**

Total SA = Curved SA $+ \pi(8^2) + \pi(5^2)$ **[M1]**

667 (or $667.0\ldots$) **[A1]**

[Back to Question] *[Total Marks: 8]*

1038 **(a)** Calculates the height of the cylinder: $18 - 5 = 13\,\text{cm}$ **[B1]**

Calculates the volume of the cylinder: $\pi \times 5^2 \times 13$ ($= 325\pi$) **[M1]**

Calculates the volume of the hemisphere: $\frac{2}{3} \times \pi \times 5^3$ ($= \frac{250\pi}{3}$) **[M1]**

1280 (accept 1280 to 1284) **[A1]**

(b) Calculates the curved surface area of the cylinder: $2 \times \pi \times 5 \times 13$ ($= 130\pi$) **[M1]**

Calculates the curved surface area of the hemisphere: $2 \times \pi \times 5^2$ ($= 50\pi$) **[M1]**

Calculates the area of the circular base: $\pi \times 5^2$ ($= 25\pi$) **[M1]**

644 (accept 643 to 645) **[A1]**

[Back to Question] *[Total Marks: 8]*

1039 **(a)** Height of cone $= 17 - 5 = 12$ cm **[M1]**

Volume of cone $= \frac{1}{3} \times \pi \times 5^2 \times 12$ (100π or 314.1...) AND Volume of hemisphere $= \frac{2}{3} \times \pi \times 5^3$ ($\frac{250}{3}\pi$ or 261.7...) **[M1]**

576 (accept 575.9 to 576.0) **[A1]**

(b) Slant height of cone $l = \sqrt{5^2 + 12^2} = 13$ **[M1]**

Curved SA of cone $= \pi \times 5 \times 13$ (65π) AND Curved SA of hemisphere $= 2 \times \pi \times 5^2$ (50π) **[M1]**

361 (accept 361.2 to 361.3) **[A1]**

(c) $\pi \times 4^2 \times h = 575.9...$ (or $\frac{550}{3}\pi$) **[M1]**

$16h = 183.3...$ (or $\frac{550}{3}$) **[M1]**

11.5 (accept 11.45 to 11.46) **[A1]**

[Back to Question] *[Total Marks: 9]*

1040 **(a)** Uses similar triangles: $\frac{h}{24} = \frac{4}{10}$ **[M1]**

$h = \frac{24 \times 4}{10} = 9.6$ cm **[A1]**

(b) Volume of large cone $= \frac{1}{3} \times \pi \times 10^2 \times 24 = 800\pi$ **[M1]**

Volume of small cone $= \frac{1}{3} \times \pi \times 4^2 \times 9.6 = 51.2\pi$ **[M1]**

2350 (or 2352.4...) **[A1]**

(c) Sets up equation: $\frac{4}{3} \times \pi \times r^3 = 748.8\pi$ (or 2352.4) **[M1]**

$r^3 = 561.6$ **[M1]**

8.25 (or 8.250...) **[A1]**

[Back to Question] *[Total Marks: 8]*

1041 **(a)** Total height of large cone $= 20 + 10 = 30$ cm **[B1]**

Volume of large cone $= \frac{1}{3} \times \pi \times 12^2 \times 30$ $(= 1440\pi$ or 4523.89...) **[M1]**

Volume of small cone $= \frac{1}{3} \times \pi \times 4^2 \times 10$ $(= 53.33\pi$ or 167.55...) **[M1]**

4360 or 4356... **[A1]**

(b) Slant height of large cone, $L = \sqrt{12^2 + 30^2}$ $(= \sqrt{1044}$ or 32.31...) AND slant height of small cone, $l = \sqrt{4^2 + 10^2}$ $(= \sqrt{116}$ or 10.77...) **[M1]**

Curved surface area of frustum $= \pi \times 12 \times \sqrt{1044} - \pi \times 4 \times \sqrt{116}$ $(= 1082.7...)$ **[M1]**

Total surface area $= 1082.7... + \pi \times 12^2 + \pi \times 4^2$ **[M1]**

1585 or 1585.3... to 1585.4 **[A1]**

[Back to Question] *[Total Marks: 8]*

1042 **(a)** Uses similar triangles: $\frac{r}{15} = \frac{16}{40}$ **[M1]**

$r = 15 \times \dfrac{16}{40} = 6$ leading correctly to given answer **[A1]**

(b) Volume of large cone $= \dfrac{1}{3} \times \pi \times 15^2 \times 40 = 3000\pi$ (9424.7...) **[M1]**

Volume of small cone $= \dfrac{1}{3} \times \pi \times 6^2 \times 16 = 192\pi$ (603.1...) **[M1]**

2808π or 8820 or 8821.5 to 8822 **[A1]**

(c) Equating volumes: $\dfrac{4}{3} \times \pi \times R^3 = 2808\pi$ **[M1]**

$R^3 = 2106$ **[M1]**

$R = \sqrt[3]{2106} = 12.8$ or 12.81 to 12.82 **[A1]**

[Back to Question] *[Total Marks: 8]*

1043 **(a)** Let x be the height of the removed cone. $\dfrac{x}{4} = \dfrac{x+15}{8}$ **[M1]**

$8x = 4x + 60 \implies 4x = 60 \implies x = 15$. Total height $= 15 + 15 = 30$ cm. **[A1]**

(b) Volume of large cone $= \dfrac{1}{3} \times \pi \times 8^2 \times 30 = 640\pi$ **[M1]**

Volume of small cone $= \dfrac{1}{3} \times \pi \times 4^2 \times 15 = 80\pi$ **[M1]**

1760 (to 3 sf) **[A1]**

(c) Slant height of large cone $L = \sqrt{8^2 + 30^2} \approx 31.05$ cm and small cone $l = \sqrt{4^2 + 15^2} \approx 15.52$ cm **[M1]**

Curved surface area of frustum $= \pi \times 8 \times \sqrt{964} - \pi \times 4 \times \sqrt{241} \approx 585.25$ **[M1]**

Area of top and bottom bases $= \pi \times 4^2 + \pi \times 8^2 = 80\pi \approx 251.33$ **[M1]**

837 (to 3 sf) **[A1]**

[Back to Question] *[Total Marks: 9]*

1044 **(a)** Identifies that the centers form an equilateral triangle of side $2r$ **[M1]**

Area of triangle $= \dfrac{1}{2} \times 2r \times 2r \times \sin(60°) = r^2\sqrt{3}$ **[M1]**

Identifies 3 sectors inside the triangle, each with angle 60°: Total sector area $= 3 \times \dfrac{60}{360} \times \pi r^2 = \dfrac{\pi}{2} r^2$ **[M1]**

Area $= r^2\sqrt{3} - \dfrac{\pi}{2} r^2 = r^2\left(\sqrt{3} - \dfrac{\pi}{2}\right)$ with no errors seen **[A1]**

(b) $r^2\left(\sqrt{3} - \dfrac{\pi}{2}\right) = 15$ **[M1]**

$r^2 = \dfrac{15}{\sqrt{3} - \pi/2}$ (≈ 93.02) **[M1]**

9.64 (or 9.644...) **[A1]**

(c) Perimeter consists of 3 arcs, each with angle 60° **[M1]**

Total perimeter $= 3 \times \dfrac{60}{360} \times 2\pi(9.644\ldots) = \pi(9.644\ldots)$ **[M1]**

30.3 (or 30.29 to 30.30) **[A1]**

[Back to Question] *[Total Marks: 10]*

1045 **(a)** $\frac{r}{10} = \frac{24-15}{24}$ or $\frac{r}{10} = \frac{9}{24}$ **[M1]**

3.75 **[A1]**

(b) Volume of large cone $= \frac{1}{3} \times \pi \times 10^2 \times 24$ (or 800π) **[M1]**

Volume of small cone $= \frac{1}{3} \times \pi \times 3.75^2 \times 9$ (or 42.1875π) **[M1]**

2380 or 2381 (accept 2380.7...) **[A1]**

(c) Slant height of large cone $L = \sqrt{10^2 + 24^2} = 26$ **[M1]**

Slant height of small cone $l = \sqrt{3.75^2 + 9^2} = 9.75$ **[M1]**

Curved surface area of frustum $= \pi \times 10 \times 26 - \pi \times 3.75 \times 9.75$ **[M1]**

Adding base and top areas: $+\pi \times 10^2 + \pi \times 3.75^2$ **[M1]**

1060 or 1060.2... to 1060.3 **[A1]**

[Back to Question] *[Total Marks: 10]*

1046 **(a)** Uses similar triangles: $\frac{h}{24} = \frac{3}{8}$ **[M1]**

$h = \frac{3 \times 24}{8} = 9$ **[A1]**

(b) Volume of large cone $= \frac{1}{3} \times \pi \times 8^2 \times 24$ (or 512π) **[M1]**

Volume of small cone $= \frac{1}{3} \times \pi \times 3^2 \times 9$ (or 27π) **[M1]**

1520 or 1524 or 485π **[A1]**

(c) Calculates slant height of large cone $L = \sqrt{8^2 + 24^2}$ (or $\sqrt{640}$) and small cone $l = \sqrt{3^2 + 9^2}$ (or $\sqrt{90}$) **[M1]**

Curved surface area of frustum $= \pi(8)(\sqrt{640}) - \pi(3)(\sqrt{90})$ **[M1]**

Calculates area of both flat circular bases: $\pi(8^2) + \pi(3^2)$ (or 73π) **[M1]**

Adds curved surface area and both base areas together **[M1]**

776 (or 775.5... to 775.6) **[A1]**

[Back to Question] *[Total Marks: 10]*

1047 **(a)** Let h be the height of the small cone. $\frac{h}{4} = \frac{h+15}{12}$ **[M1]**

$12h = 4h + 60$ **[M1]**

$h = 7.5$ **[A1]**

(b) Volume of large cone $= \frac{1}{3} \times \pi \times 12^2 \times 22.5$ (1080π or 3392.9...) **[M1]**

Volume of small cone $= \frac{1}{3} \times \pi \times 4^2 \times 7.5$ (40π or 125.6...) **[M1]**

3270 (accept 3267 to 3268) **[A1]**

(c) Slant height of large cone $= \sqrt{12^2 + 22.5^2} = 25.5$ **[M1]**

Slant height of small cone $= \sqrt{4^2 + 7.5^2} = 8.5$ **[M1]**

Curved surface area of frustum $= \pi \times 12 \times 25.5 - \pi \times 4 \times 8.5$ ($306\pi - 34\pi = 272\pi$) **[M1]**

Area of both circular bases $= \pi \times 12^2 + \pi \times 4^2$ ($144\pi + 16\pi = 160\pi$) **[M1]**

1360 (accept 1357 to 1358) **[A1]**

[Back to Question] *[Total Marks: 11]*

1048 **(a)** Let H be the height of the large cone. Height of small removed cone $= H - 8$ **[M1]**

Uses similar triangles: $\frac{H}{12} = \frac{H-8}{6}$ (or equivalent ratio) **[M1]**

$6H = 12H - 96$

6H = 96

H = 16*withnoerrorsseen*[*A*1]

(b) Volume of frustum $= \frac{1}{3} \times \pi \times 12^2 \times 16 - \frac{1}{3} \times \pi \times 6^2 \times 8$ ($= 768\pi - 96\pi = 672\pi$)

[M2, M1 for one correct cone volume]

Volume of hemisphere $= \frac{2}{3} \times \pi \times 6^3$ ($= 144\pi$) **[M1]**

Total volume $= 672\pi + 144\pi = 816\pi = 2560$ or $2563.5\ldots$ **[A1]**

(c) Slant height of large cone $= \sqrt{12^2 + 16^2} = 20$ AND Slant height of small cone $= \sqrt{6^2 + 8^2} = 10$ **[M1]**

Curved surface area of frustum $= \pi \times 12 \times 20 - \pi \times 6 \times 10$ ($= 180\pi$) **[M1]**

Total surface area $= 180\pi$ (frustum curved) $+\pi \times 12^2$ (base) $+2 \times \pi \times 6^2$ (hemisphere) **[M1]**

1240 or 1244 to 1244.2 **[A1]**

[Back to Question] *[Total Marks: 11]*

1049 **(a)** Sets up an equation using similar triangles (e.g. let x be the height of the smaller cone): $\frac{x}{4} = \frac{x+10}{6}$ (or equivalent) **[M1]**

Solves for the height of the smaller cone: $6x = 4x + 40 \implies 2x = 40 \implies x = 20$ **[M1]**

Calculates the height of the larger cone: $20 + 10 = 30$ **[A1]**

(b) Calculates the volume of the larger cone: $\frac{1}{3} \times \pi \times 6^2 \times 30$ ($= 360\pi$) **[M1]**

Calculates the volume of the smaller cone: $\frac{1}{3} \times \pi \times 4^2 \times 20$ ($= \frac{320\pi}{3}$) **[M1]**

796 (accept 795 to 797) **[A1]**

(c) Calculates the slant height of the larger cone: $\sqrt{6^2 + 30^2} = \sqrt{936}$ ($= 30.59...$) **[M1]**

Calculates the slant height of the smaller cone: $\sqrt{4^2 + 20^2} = \sqrt{416}$ $(= 20.39...)$ **[M1]**

Calculates the curved surface area of the frustum: $\pi \times 6 \times \sqrt{936} - \pi \times 4 \times \sqrt{416}$ $(= 320.38...)$ **[M1]**

Calculates the sum of the areas of the circular top and base: $\pi \times 6^2 + \pi \times 4^2$ $(= 52\pi$ or $163.36...)$ **[M1]**

484 (accept 483.7 to 484.5) **[A1]**

[Back to Question] *[Total Marks: 11]*

1050

Half of the chord length $= 24 \div 2 = 12$ **[M1]**

Uses Pythagoras' theorem: $r^2 = 5^2 + 12^2$ **[M1]**

13 **[A1]**

[Back to Question] *[Total Marks: 3]*

1051

Uses Pythagoras' theorem: $(8 - -4)^2 + (1 - 6)^2$ or $12^2 + (-5)^2$ **[M1]**

$144 + 25 = 169$ **[M1]**

13 **[A1]**

[Back to Question] *[Total Marks: 3]*

1052

$\Delta x = 4 - (-2) = 6$ and $\Delta y = 9 - 1 = 8$ **[M1]**

$\sqrt{6^2 + 8^2}$ **[M1]**

10 **[A1]**

[Back to Question] *[Total Marks: 3]*

1053

Use of Pythagoras' theorem: $(k - 1)^2 + (19 - 4)^2 = 17^2$ **[M1]**

$(k - 1)^2 + 15^2 = 289$ or $(k - 1)^2 = 64$ **[M1]**

$k - 1 = 8$ and $k - 1 = -8$ (or $k^2 - 2k - 63 = 0$

(k-9)(k+7) = 0)$[M1]$

9 and -7 **[A1]**

[Back to Question] *[Total Marks: 4]*

1054 **(a)**

$(x - 7)^2 + x^2 = (x + 1)^2$ **[M1]**

$x^2 - 14x + 49 + x^2 = x^2 + 2x + 1$ **[M1]**

$2x^2 - 14x + 49 = x^2 + 2x + 1$

$x^2 - 16x + 48 = 0$ **[A1]**

(b)

$(x - 4)(x - 12) = 0$ **[M1]**

$x = 4$ or $x = 12$ **[A1]**

Rejects $x = 4$ because $x - 7$ would be -3 (length must be positive), so $x = 12$ **[B1]**

Perimeter $= 5 + 12 + 13 = 30$ **[A1]**

[Back to Question] *[Total Marks: 7]*

1055 **(a)** $AM = 16 \div 2 = 8$ **[M1]**

$PM = \sqrt{10^2 - 8^2} = \sqrt{36} = 6$ with no errors seen **[A1]**

(b) In triangle AMQ, $QM^2 = 17^2 - 8^2$ or $289 - 64$ **[M1]**

$QM = \sqrt{225} = 15$ **[A1]**

$PQ = 6 + 15 = 21$ **[A1]**

[Back to Question] *[Total Marks: 5]*

1056 **(a)** $(2x + 1)^2 = (x - 1)^2 + (x + 6)^2$ **[M1]**

$4x^2 + 4x + 1 = (x^2 - 2x + 1) + (x^2 + 12x + 36)$ **[M1]**

$4x^2 + 4x + 1 = 2x^2 + 10x + 37$ **[M1]**

$2x^2 - 6x - 36 = 0$

$x^2 - 3x - 18 = 0$ with no errors seen [A1]

(b) $(x - 6)(x + 3) = 0$ **[M1]**

$x = 6$ or $x = -3$ **[A1]**

Selects $x = 6$ and rejects $x = -3$ because lengths must be strictly positive **[B1]**

(c) Base $= 6 - 1 = 5$ and Height $= 6 + 6 = 12$ **[M1]**

Area $= \frac{1}{2} \times 5 \times 12$ **[M1]**

30 **[A1]**

[Back to Question] *[Total Marks: 10]*

1057 Let M be the midpoint of AB. $AM^2 = 8.5^2 - 4^2$ **[M1]**

$AM = \sqrt{72.25 - 16} = \sqrt{56.25} = 7.5$ **[M1]**

Chord $AB = 2 \times 7.5 = 15$ **[A1]**

[Back to Question] *[Total Marks: 3]*

1058 **(a)** $AM = 15 \div 2 = 7.5$ **[B1]**

$d^2 + 7.5^2 = 8.5^2$ **[M1]**

$d = \sqrt{72.25 - 56.25} = \sqrt{16} = 4$ **[A1]**

(b) Let half-chord be x. $x^2 + 7.5^2 = 8.5^2$ **[M1]**

$x = \sqrt{72.25 - 56.25} = 4$ **[A1]**

Length of chord $= 2 \times 4 = 8$ **[A1]**

[Back to Question] *[Total Marks: 6]*

1059 **(a)** Let half the chord length be x. Uses $12.4^2 = x^2 + 5.2^2$ **[M1]**

$x = \sqrt{12.4^2 - 5.2^2}$ or $x = 11.257\ldots$ **[M1]**

22.5 (or 22.51 to 22.52) **[A1]**

(b) Let perpendicular distance from O to CD be y. Uses $12.4^2 = y^2 + 9^2$ **[M1]**

$y = \sqrt{12.4^2 - 9^2}$ or $y = 8.5299\ldots$ **[A1]**

Total distance $= 5.2 + 8.5299\ldots$ **[M1]**

13.7 (or 13.72 to 13.73) **[A1]**

[Back to Question] *[Total Marks: 7]*

1060 **(a)** Uses distance formula: $\sqrt{(6-(-2))^2 + (y-(-1))^2} = 10$ **[M1]**

$64 + (y+1)^2 = 100$ **[M1]**

$(y+1)^2 = 36$

$y + 1 = 6$ (since $y > 0$)

5 **[A1]**

(b) Let C be $(x, 0)$.

$AC^2 = (x-(-2))^2 + (0-(-1))^2 = (x+2)^2 + 1$ **[M1]**

$BC^2 = (x-6)^2 + (0-5)^2 = (x-6)^2 + 25$ **[M1]**

Equating distances squared: $x^2 + 4x + 5 = x^2 - 12x + 61$ **[M1]**

$16x = 56$

3.5 **[A1]**

[Back to Question] *[Total Marks: 7]*

1061 **(a)** $x^2 + (x+7)^2 = (x+8)^2$ **[M1]**

$x^2 + x^2 + 14x + 49 = x^2 + 16x + 64$ **[M1]**

$x^2 - 2x - 15 = 0$ **[A1]**

(b) $(x-5)(x+3) = 0$ or correct use of the quadratic formula **[M1]**

$x = 5$ (accept value 5 only, ignoring -3 if appropriately discarded) **[A1]**

(c) Base $= 5 + 7 = 12$ and Height $= 5$ **[M1]**

Area $= \frac{1}{2} \times 12 \times 5 = 30$ **[A1]**

[Back to Question] *[Total Marks: 7]*

1062 Identifies cosine ratio: $\cos(\theta) = \frac{2.1}{6.5}$ **[M1]**

$\theta = \cos^{-1}\left(\frac{2.1}{6.5}\right)$ **[M1]**

71.2 **[A1]**

[Back to Question] *[Total Marks: 3]*

1063 $\cos(58°) = \frac{9.4}{XZ}$ **[M1]**

$XZ = \frac{9.4}{\cos(58°)}$ **[M1]**

17.7 or 17.73... **[A1]**

[Back to Question] *[Total Marks: 3]*

1064 **(a)** $\tan(38°) = \frac{TB}{45}$ **[M1]**

35.2 (or 35.15 to 35.16) **[A1]**

(b) $\cos(38°) = \frac{45}{PT}$ or $PT^2 = 45^2 + 35.15^2$ **[M1]**

57.1 (or 57.10 to 57.11) **[A1]**

(c) $\tan(XPB) = \frac{20}{45}$ **[M1]**

Angle $= \tan^{-1}\left(\frac{20}{45}\right)$ **[M1]**

24.0 (or 23.96 . . .) **[A1]**

[Back to Question] *[Total Marks: 7]*

1065 **(a)** Uses $\sin(52°) = \frac{\text{distance}}{18}$ or $18 \times \sin(52°)$ **[M1]**

14.18... or 14.2 **[A1]**

(b) Distance to closest point $= 18 \times \cos(52°) = 11.082...$ km **[M1]**

Time $= \frac{11.082...}{14}$ $(= 0.7915...$ hours) **[M1]**

$0.7915... \times 60 = 47.49... = 47$ **[A1]**

[Back to Question] *[Total Marks: 5]*

1066 **(a)** Calculates $\angle PAB = 360° - (180° - 40°) - 130° = 90°$ (or similar valid reasoning to show right angle) **[M1]**

Uses Pythagoras' theorem: $PB^2 = 80^2 + 60^2$ **[M1]**

100 **[A1]**

(b) Uses trigonometry to find $\angle APB$: $\tan(APB) = \frac{60}{80}$ **[M1]**

$\angle APB = \tan^{-1}(0.75) = 36.869...°$ **[A1]**

Bearing $= 040° + 36.869...° = 076.9$ (accept 077) **[A1]**

(c) Recognises the shortest distance d is the perpendicular height of $\triangle PAB$ from A to PB. **[M1]**

Uses area equation: $\frac{1}{2} \times 100 \times d = \frac{1}{2} \times 80 \times 60$ OR uses trigonometry $80\sin(53.13°)$ or $60\sin(36.87°)$ **[M1]**

48 **[A1]**

[Back to Question] *[Total Marks: 9]*

1067 **(a)** Interior angle between parallel North lines $= 180° - 52° = 128°$ **[M1]**

Angles around point $B = 360° - 128° - 142° = 90°$ with no errors seen **[A1]**

(b) $AC = \sqrt{15^2 + 20^2}$ **[M1]**

$AC = \sqrt{625}$ **[M1]**

25 **[A1]**

(c) $\tan(BAC) = \frac{20}{15}$ **[M1]**

Angle $BAC = 53.13°$ **[A1]**

Bearing $= 052° + 53.13°$ **[M1]**

105.1 (or 105.13 . . .) **[A1]**

[Back to Question] *[Total Marks: 9]*

1068 **(a)** $PR = \sqrt{8.5^2 + 11.2^2}$ or $\sqrt{197.69}$ **[M1]**

14.06... or 14.1 **[A1]**

(b) $\sin(41°) = \frac{14.06...}{PS}$ or $PS = \frac{14.06...}{\sin(41°)}$ **[M1]**

21.43... or 21.4 **[A1]**

(c) Angle $RPS = 180° - 90° - 41° = 49°$ AND distance $= 14.06... \times \sin(49°)$
OR area method: $0.5 \times 21.43... \times h = 0.5 \times 14.06... \times (14.06.../\tan(41°))$ **[M1]**

10.60... or 10.6 **[A1]**

[Back to Question] *[Total Marks: 6]*

1069 **(a)** Sets up equation for $\triangle BCD$: $\tan(42°) = \frac{CD}{BC}$

BC $= \frac{CD}{\tan(42°)}$ [M1]

Sets up equation for $\triangle ACD$: $\tan(25°) = \frac{CD}{BC + 50}$

BC + 50 $= \frac{CD}{\tan(25°)}$ [M1]

Equates or substitutes to eliminate BC: $50 = \dfrac{CD}{\tan(25°)} - \dfrac{CD}{\tan(42°)}$ **[M1]**

Factorises to solve for CD: $CD\left(\dfrac{1}{\tan(25°)} - \dfrac{1}{\tan(42°)}\right) = 50$ **[M1]**

48.4 (accept 48.36...) **[A1]**

(b) Recognises the shortest distance d is the perpendicular from B to AD, forming a right-angled triangle with hypotenuse $AB = 50$. **[M1]**

Sets up trigonometric ratio: $\sin(25°) = \dfrac{d}{50}$ **[M1]**

21.1 (accept 21.13...) **[A1]**

[Back to Question] *[Total Marks: 8]*

1070 **(a)** $AC = \sqrt{8^2 + 15^2}$ **[M1]**

17 **[A1]**

(b) $AD^2 + 12^2 = 17^2$ (or their AC^2) **[M1]**

$AD = \sqrt{289 - 144} = \sqrt{145}$ **[M1]**

12.0 (or 12.04...) **[A1]**

(c) Area of $\triangle ADC = \dfrac{1}{2} \times 12 \times \sqrt{145} = 72.25\ldots$ **[M1]**

$\dfrac{1}{2} \times 17 \times h = 72.25\ldots$ **[M1]**

8.50 (or 8.499...) **[A1]**

(d) $\tan(BAC) = \dfrac{15}{8} \implies$ Angle $BAC = 61.93°$ **[M1]**

$\tan(CAD) = \dfrac{12}{\sqrt{145}} \implies$ Angle $CAD = 44.90°$ **[M1]**

Angle $DAB = 61.93° + 44.90°$ **[M1]**

106.8 (or 106.82 to 106.83) **[A1]**

[Back to Question] *[Total Marks: 12]*

1071 **(a)** $\dfrac{\sqrt{3}}{2}$ **[B1]**

(b) $\sin 30° = \dfrac{1}{2}$ and $\tan 60° = \sqrt{3}$ **[M1]**

$\dfrac{1 - 1/2}{(\sqrt{3})^2} = \dfrac{1/2}{3} = \dfrac{1}{6}$ with no errors seen **[A1]**

[Back to Question] *[Total Marks: 3]*

1072 **(a)** 2 **[B1]**

(b) $\dfrac{1}{2}$ **[B1]**

(c) $\dfrac{\sqrt{3}}{\sqrt{3}/2}$ **[M1]**

2 **[A1]**

[Back to Question] *[Total Marks: 4]*

1073 **(a)** Substitutes $\sin 60° = \dfrac{\sqrt{3}}{2}$ and $\tan 30° = \dfrac{1}{\sqrt{3}}$ (or $\dfrac{\sqrt{3}}{3}$) **[M1]**

1 **[A1]**

(b) Substitutes $\tan 45° = 1$ leading to $\cos x = \dfrac{1}{\sqrt{2}}$ **[M1]**

45 **[A1]**

[Back to Question] *[Total Marks: 4]*

1074 **(a)** $\sin 45° = \dfrac{6}{PR}$ (or $\cos 45° = \dfrac{6}{PR}$) **[M1]**

$PR = 6 \div \dfrac{\sqrt{2}}{2} = 6\sqrt{2}$ **[A1]**

(b) $\sin 60° = \dfrac{6\sqrt{2}}{PS}$ **[M1]**

$PS = 6\sqrt{2} \div \dfrac{\sqrt{3}}{2}$ **[M1]**

$PS = \dfrac{12\sqrt{2}}{\sqrt{3}} = \dfrac{12\sqrt{6}}{3} = 4\sqrt{6}$ **[A1]**

[Back to Question] *[Total Marks: 5]*

1075 **(a)** $\sin 45° = \dfrac{PQ}{8\sqrt{2}}$ **[M1]**

$PQ = 8\sqrt{2} \times \dfrac{1}{\sqrt{2}} = 8$ **[A1]**

(b) $\cos 30° = \dfrac{8\sqrt{2}}{PS}$ or $PS = \dfrac{8\sqrt{2}}{\cos 30°}$ **[M1]**

$PS = \dfrac{8\sqrt{2}}{\sqrt{3}/2} = \dfrac{16\sqrt{2}}{\sqrt{3}}$ **[M1]**

Rationalises the denominator to get $\dfrac{16\sqrt{6}}{3}$ **[A1]**

[Back to Question] *[Total Marks: 5]*

1076 In $\triangle ABD$, $AD = 8\sqrt{2}\cos 45° = 8$ **[M1]**

$BD = 8\sqrt{2}\sin 45° = 8$ **[A1]**

In $\triangle BDC$, $BC = \dfrac{BD}{\sin 30°} = \dfrac{8}{0.5} = 16$ **[M1]**

$DC = \dfrac{BD}{\tan 30°} = 8\sqrt{3}$ **[M1]**

Perimeter $= AB + BC + AD + DC = 8\sqrt{2} + 16 + 8 + 8\sqrt{3}$ **[M1]**

$24 + 8\sqrt{2} + 8\sqrt{3}$ **[A1]**

[Back to Question] *[Total Marks: 6]*

1077 **(a)** Substitutes $\tan 30° = \dfrac{1}{\sqrt{3}}$ to get $\tan^2 30° = \dfrac{1}{3}$ **[B1]**

Evaluates numerator as $1 - \dfrac{1}{3} = \dfrac{2}{3}$ AND denominator as $1 + \dfrac{1}{3} = \dfrac{4}{3}$ **[M1]**

Simplifies $\dfrac{2/3}{4/3} = \dfrac{1}{2}$ which is equal to $\cos 60°$ **[A1]**

(b) Substitutes exact values: $x\left(\dfrac{\sqrt{3}}{2}\right) = \sqrt{3} - x\left(\dfrac{1}{2}\right)$ **[M1]**

Rearranges to collect x terms: $x\left(\dfrac{\sqrt{3}}{2} + \dfrac{1}{2}\right) = \sqrt{3}$ **[M1]**

Isolates x: $x = \dfrac{2\sqrt{3}}{\sqrt{3}+1}$ **[M1]**

Rationalises denominator: $\dfrac{2\sqrt{3}(\sqrt{3}-1)}{(\sqrt{3}+1)(\sqrt{3}-1)} = \dfrac{6-2\sqrt{3}}{2} = 3 - \sqrt{3}$ **[A1]**

[Back to Question] *[Total Marks: 7]*

1078 **(a)** $\cos 30° = \dfrac{AD}{12}$ **[M1]**

$AD = 12 \times \dfrac{\sqrt{3}}{2} = 6\sqrt{3}$ with no errors seen **[A1]**

(b) Uses Area $= AB \times AD = 60\sqrt{3}$ to find $AB = 10$ **[B1]**

$DE = 12\sin 30° = 6$ **[M1]**

$CE = CD - DE = 10 - 6 = 4$ **[A1]**

(c) Identifies $BC = 6\sqrt{3}$ and $CE = 4$ **[M1]**

$\tan(\angle EBC) = \dfrac{CE}{BC} = \dfrac{4}{6\sqrt{3}}$ **[M1]**

$\dfrac{2}{3\sqrt{3}} = \dfrac{2\sqrt{3}}{9}$ **[A1]**

[Back to Question] *[Total Marks: 8]*

1079 **(a)** Uses cosine rule: $7^2 = x^2 + (x+3)^2 - 2(x)(x+3)\cos 60°$ **[M1]**

Substitutes $\cos 60° = 0.5$: $49 = x^2 + x^2 + 6x + 9 - x(x+3)$ **[M1]**

Expands and simplifies: $49 = 2x^2 + 6x + 9 - x^2 - 3x$ **[M1]**

$x^2 + 3x - 40 = 0$ **[A1]**

(b) $(x + 8)(x - 5) = 0$ **[M1]**

$x = 5$ (rejects $x = -8$) **[A1]**

Perimeter $= 5 + (5 + 3) + 7 = 20$ **[A1]**

(c) Uses sine area formula: Area $= \frac{1}{2} \times 5 \times 8 \times \sin 60°$ **[M1]**

Substitutes $\sin 60° = \frac{\sqrt{3}}{2}$ **[M1]**

$10\sqrt{3}$ **[A1]**

[Back to Question] *[Total Marks: 10]*

1080 $\sin x = \frac{1}{\sqrt{2}}$ **[M1]**

$x = 45°$ **[B1]**

$x = 135°$ **[B1]**

[Back to Question] *[Total Marks: 3]*

1081 **(a)** $(180, -1)$ **[B1]**

(b) $60°$ **[B1]**

$300°$ **[B1]**

[Back to Question] *[Total Marks: 3]*

1082 **(a)** Smooth sine wave shape passing through $(0, 0)$, $(180, 0)$ and $(360, 0)$ **[B1]**

Maximum turning point at $(90, 1)$ and minimum turning point at $(270, -1)$ **[B1]**

(b) $x = 30$ **[B1]**

$x = 150$ **[B1]**

[Back to Question] *[Total Marks: 4]*

1083 **(a)** $\cos x = 0.5$ **[M1]**

$x = 60°$ **[B1]**

$x = 300°$ **[B1]**

(b) 0 (since $\sin x = \frac{5}{3} > 1$) **[B1]**

[Back to Question] *[Total Marks: 4]*

1084 **(a)** Smooth U-shaped cosine curve starting at $(0, 1)$ and ending at $(4, 1)$ representing 360° **[B1]**

Curve passes through $(1, 0)$, $(3, 0)$ and has a minimum at $(2, -1)$ **[B1]**

(b) $(180°, -1)$ **[B1]**

(c) $\cos x = -0.5$ (or $-\frac{1}{2}$) **[M1]**

$x = 120°$ **[B1]**

$x = 240°$ **[B1]**

[Back to Question] *[Total Marks: 6]*

1085 **(a)** Smooth sine curve passing through $(0, 0), (90°, 1), (180°, 0), (270°, -1), (360°, 0)$ **[B1]**

Smooth cosine curve passing through $(0, 1), (90°, 0), (180°, -1), (270°, 0), (360°, 1)$ **[B1]**

Intersection points shown approximately accurately at $x = 45°$ and $x = 225°$ **[B1]**

(b) $\tan x = 1$ **[M1]**

$x = 45°, 225°$ (Both required) **[A1]**

(c) Identifies the region between the two intersection points **[M1]**

$45° < x < 225°$ **[A1]**

[Back to Question] *[Total Marks: 7]*

1086 **(a)** $\sin x = -\frac{\sqrt{3}}{2}$ **[M1]**

$240°$ **[A1]**

$300°$ **[A1]**

(b) $\cos^2 x = \frac{3}{4}$ **[M1]**

$\cos x = \pm\frac{\sqrt{3}}{2}$ **[M1]**

$30°$ and $150°$ **[A1]**

$210°$ and $330°$ **[A1]**

(Award maximum of 3 marks if $\pm$ is omitted, resulting in only two solutions)

[Back to Question] *[Total Marks: 7]*

1087 **(a)** $\sin x = -0.75$ **[M1]**

Reference angle $= 48.6°$ (or $48.59...°$) **[M1]**

$x = 228.6°, 311.4°$ (Both required for the final mark) **[A1]**

(b) x-coordinate is $270°$ **[B1]**

y-coordinate is -1 **[B1]**

[Back to Question] *[Total Marks: 5]*

1088 **(a)** Correct basic shape with multiple branches passing through $(0, 0)$, $(180, 0)$, and $(360, 0)$

[B1]

Asymptotes at $x = 90$ and $x = 270$ clearly indicated or graph showing asymptotic behaviour [B1]

(b) $\tan x = -\frac{5}{3}$ or $-1.66\ldots$ [M1]

$121.0°$ [A1]

$301.0°$ [A1]

[Back to Question] *[Total Marks: 5]*

1089 **(a)** $\cos x = -0.25$ [M1]

$x = 104.5$ [A1]

$x = 360 - 104.47... = 255.5$ [A1]

(b) $\tan x = 2.5$ [M1]

$x = 68.2$ [A1]

$x = 68.19... + 180 = 248.2$ [A1]

[Back to Question] *[Total Marks: 6]*

1090 **(a)** $\sin 30° = \frac{1}{2}$ leading to $a\left(\frac{1}{2}\right) - b = 0$ or $\frac{b}{a} = \frac{1}{2}$ [M1]

$a = 2$ and $b = 1$ [A1]

(b) 150 [B1]

(c) $k = \cos(242°) = -0.469$ [B1]

Uses symmetry of cosine curve: $360° - 242°$ [M1]

$x = 118°$ [A1]

[Back to Question] *[Total Marks: 6]*

1091 **(a)** $\tan x = -2.8$ [M1]

$x = 109.7°$ (accept $109.6°$ to $109.7°$) [B1]

$x = 289.7°$ (accept $289.6°$ to $289.7°$) [B1]

(b) Factorises to $\sin x(2\sin x - 1) = 0$ [M1]

Identifies $\sin x = 0$ or $\sin x = 0.5$ (or $\frac{1}{2}$) [M1]

$x = 0°, 180°, 360°$ (all three correct for the mark) [A1]

$x = 30°, 150°$ (both correct for the mark) [A1]

[Back to Question] *[Total Marks: 7]*

1092 **(a)** $\frac{1}{2} \times 8 \times 5 \times \sin 60°$ [M1]

$10\sqrt{3}$ **[A1]**

(b) $QR^2 = 8^2 + 5^2 - 2(8)(5)\cos 60°$ **[M1]**

$QR^2 = 64 + 25 - 80\left(\frac{1}{2}\right)$ **[M1]**

7 **[A1]**

[Back to Question] *[Total Marks: 5]*

1093 **(a)** $7^2 = 5^2 + x^2 - 2(5)(x)\cos 60°$ **[M1]**

$\cos 60° = \frac{1}{2}$ or 0.5 seen **[B1]**

$49 = 25 + x^2 - 5x \implies x^2 - 5x - 24 = 0$ (with no errors seen) **[A1]**

(b) $(x - 8)(x + 3) = 0$ **[M1]**

$x = 8$ **[A1]**

(c) Area $= \frac{1}{2} \times 5 \times 8 \times \sin 60°$ **[M1]**

$10\sqrt{3}$ **[A1]**

[Back to Question] *[Total Marks: 7]*

1094 **(a)** $0.5 \times 8 \times 12 \times \sin(42°)$ **[M1]**

32.1 (or 32.11 to 32.12) **[A1]**

(b) $BC^2 = 8^2 + 12^2 - 2(8)(12)\cos(42°)$ **[M1]**

$BC^2 = 65.3...$ **[M1]**

8.08 (or 8.081 to 8.082) **[A1]**

[Back to Question] *[Total Marks: 5]*

1095 **(a)** $\frac{1}{2} \times 8 \times 12 \times \sin 40°$ **[M1]**

30.9 (accept 30.85 . . .) **[A1]**

(b) $8^2 + 12^2 - 2 \times 8 \times 12 \times \cos 40°$ **[M1]**

60.9 . . . **[A1]**

7.80 or 7.81 **[A1]**

[Back to Question] *[Total Marks: 5]*

1096 **(a)** $0.5 \times 8.5 \times 11.2 \times \sin(54°)$ **[M1]**

38.5 (38.50 to 38.51) **[A1]**

(b) $BC^2 = 8.5^2 + 11.2^2 - 2(8.5)(11.2)\cos(54°)$ [M2]
(or [M1] for correct cosine rule formula with one substitution error)

$BC^2 = 85.77...$ **[A1]**

9.26 (9.261...) **[A1]**

[Back to Question] *[Total Marks: 5]*

1097 **(a)** $\frac{1}{2} \times 8 \times 12 \times \sin(50°)$ **[M1]**

36.77... so 36.8 **[A1]**

(b) $BC^2 = 8^2 + 12^2 - 2(8)(12)\cos(50°)$ **[M1]**

$BC^2 = 208 - 123.415... = 84.58...$ **[M1]**

9.197... so 9.20 **[A1]**

[Back to Question] *[Total Marks: 5]*

1098 **(a)** $\frac{1}{2} \times 7.2 \times 9.5 \times \sin 64°$ **[M1]**

30.7 or 30.73... **[A1]**

(b) $7.2^2 + 9.5^2 - 2(7.2)(9.5)\cos 64°$ **[M2]**

(or $M1$ for correct implicit cosine rule equation)

9.06 or 9.062... **[A1]**

[Back to Question] *[Total Marks: 5]*

1099 **(a)** Uses cosine rule: $BC^2 = 8^2 + 12^2 - 2(8)(12)\cos 65°$ **[M1]**

$BC^2 = 126.85...$ **[M1]**

11.3 **[A1]**

(b) Uses area formula: $\frac{1}{2} \times 8 \times 12 \times \sin 65°$ **[M1]**

43.5 **[A1]**

[Back to Question] *[Total Marks: 5]*

1100 **(a)** $\frac{1}{2} \times 8 \times 11 \times \sin(65°)$ **[M1]**

39.9 **[A1]**

(b) $BC^2 = 8^2 + 11^2 - 2(8)(11)\cos(65°)$ **[M2]**

($M1$ for correct substitution into implicit cosine rule)

10.5 **[A1]**

[Back to Question] *[Total Marks: 5]*

1101 **(a)** $\frac{BC}{\sin 45°} = \frac{12}{\sin 60°}$ **[M1]**

$BC = \dfrac{12\sin 45°}{\sin 60°}$ **[M1]**

9.80 **[A1]**

(b) Angle $ACB = 180° - (45° + 60°) = 75°$ **[B1]**

Area $= \dfrac{1}{2} \times 12 \times 9.798 \times \sin 75°$ **[M1]**

56.8 **[A1]**

[Back to Question] *[Total Marks: 6]*

1102 **(a)** $\dfrac{BC}{\sin 40°} = \dfrac{12}{\sin 60°}$ **[M1]**

$BC = \dfrac{12\sin 40°}{\sin 60°}$ **[M1]**

8.91 (accept 8.907 . . .) **[A1]**

(b) Angle $BCA = 180° - (40° + 60°) = 80°$ **[B1]**

$\dfrac{1}{2} \times 12 \times 8.91 \times \sin 80°$ **[M1]**

52.6 (accept 52.62 . . . to 52.65 . . .) **[A1]**

[Back to Question] *[Total Marks: 6]*

1103 **(a)** $\cos Q = \dfrac{7^2 + 8^2 - 12^2}{2 \times 7 \times 8}$ [M2] (M1 for correct implicit cosine rule equation e.g. $12^2 = 7^2 + 8^2 - 2(7)(8)\cos Q$)

106.1 (accept 106.07 . . .) **[A1]**

(b) $\dfrac{1}{2} \times 7 \times 8 \times \sin 106.1°$ **[M1]**

26.9 (accept 26.89 . . .) **[A1]**

[Back to Question] *[Total Marks: 5]*

1104 **(a)** $\dfrac{\sin P}{12} = \dfrac{\sin 35°}{8}$ **[M1]**

$\sin P = \dfrac{12\sin 35°}{8} = 0.8603...$ **[M1]**

Acute angle $= 59.35...°$ **[A1]**

Obtuse angle $P = 180° - 59.35...° = 120.64...°$ so $120.6°$ **[A1]**

(b) Angle $R = 180° - 35° - 120.64...° = 24.36...°$ **[M1]**

Area $= \dfrac{1}{2} \times 8 \times 12 \times \sin(24.36...°) = 19.79...$ so 19.8 **[A1]**

[Back to Question] *[Total Marks: 6]*

1105 **(a)** $\dfrac{\sin R}{15} = \dfrac{\sin 42°}{12}$ **[M1]**

$\sin R = \dfrac{15\sin 42°}{12}$ (evaluates to $0.836\ldots$) **[M1]**

123.2 (accept 123.2 to 123.3) **[A1]**

(b) Angle $P = 180° - 42° - 123.24° = 14.76°\ldots$ **[M1]**

$\dfrac{QR}{\sin 14.76°} = \dfrac{12}{\sin 42°}$ (or equivalent application of sine/cosine rule) **[M1]**

4.57 (accept 4.56 to 4.58) **[A1]**

[Back to Question] *[Total Marks: 6]*

1106 **(a)** $\dfrac{\sin R}{14} = \dfrac{\sin 35°}{10}$ **[M1]**

$\sin^{-1}\left(\dfrac{14\sin 35°}{10}\right)$ leading to 53.4° (53.41...) **[M1]**

126.6 (126.58...) **[A1]**

(b) Angle $PQR = 180° - 35° - 126.58° = 18.4°$ **[M1]**

Area $= \dfrac{1}{2} \times 14 \times 10 \times \sin(18.4°)$ **[M1]**

22.1 (22.11...) **[A1]**

[Back to Question] *[Total Marks: 6]*

1107 **(a)** $\dfrac{\sin Q}{14} = \dfrac{\sin 32°}{9}$ **[M1]**

$\sin Q = \dfrac{14\sin 32°}{9}$ or 0.824... **[M1]**

Acute angle $= 55.5°$ (55.51...) **[A1]**

$180 - 55.5... = 124.5°$ (124.48 to 124.5) **[A1]**

(b) Angle $PRQ = 180 - 32 - 124.48 = 23.52°$ **[B1]**

$0.5 \times 14 \times 9 \times \sin(23.52°)$ **[M1]**

25.1 (25.12 to 25.14) **[A1]**

[Back to Question] *[Total Marks: 7]*

1108 **(a)** $\cos Q = \dfrac{15^2 + 12^2 - 18^2}{2 \times 15 \times 12}$ **[M1]**

$\cos Q = \dfrac{45}{360}$ OR 0.125 **[M1]**

$Q = \cos^{-1}(0.125) = 82.819...°$ **[A1]**

(b) $0.5 \times 15 \times 12 \times \sin(82.8°)$ **[M1]**

89.3 (or 89.28 to 89.30) **[A1]**

(c) Area $= 0.5 \times 15 \times h = 89.3$ OR $h = 12 \times \sin(82.8°)$ **[M1]**

11.9 (or 11.90 to 11.91) **[A1]**

[Back to Question] *[Total Marks: 7]*

1109 **(a)** $\dfrac{\sin C}{12} = \dfrac{\sin 35°}{9}$ **[M1]**

$\sin C = \dfrac{12\sin 35°}{9}$ $(= 0.764...)$ **[M1]**

$49.9°$ (or $49.88...$) **[A1]**

$130.1°$ (or $130.11...$) **[A1]**

(b) Angle $ABC = 180 - (35 + 130.1) = 14.9°$ **[M1]**

Area $= \dfrac{1}{2} \times 12 \times 9 \times \sin(14.9°)$ **[M1]**

13.9 (accept $13.87...$ to $13.89...$) **[A1]**

[Back to Question] *[Total Marks: 7]*

1110 **(a)** Uses sine rule: $\dfrac{\sin R}{15.2} = \dfrac{\sin 72°}{18.5}$ **[M1]**

$\sin R = 0.781...$ **[M1]**

51.4 **[A1]**

(b) Angle $P = 180° - 72° - 51.38° = 56.62°$ **[B1]**

$\dfrac{QR}{\sin 56.62°} = \dfrac{18.5}{\sin 72°}$ **[M1]**

16.2 **[A1]**

(c) Shortest distance $= 15.2 \times \sin 72°$ **[M1]**

14.5 **[A1]**

[Back to Question] *[Total Marks: 8]*

1111 **(a)** $BD^2 = 8^2 + 10^2 - 2(8)(10)\cos 110°$ **[M1]**

$BD^2 = 164 - 160(-0.342) = 218.72$ **[M1]**

14.8 **[A1]**

(b) $\cos C = \dfrac{14^2 + 15^2 - 14.79^2}{2(14)(15)}$ **[M1]**

$\cos C = \dfrac{202.26}{420} = 0.4816$ **[M1]**

61.2 **[A1]**

(c) Area $ABD = \dfrac{1}{2}(8)(10)\sin 110° = 37.59$ **[M1]**

Area $BCD = \dfrac{1}{2}(14)(15)\sin 61.2° = 92.01$ **[M1]**

130 (or 129.6) **[A1]**

[Back to Question] *[Total Marks: 9]*

1112 **(a)** $AC^2 = 12.4^2 + 9.8^2 - 2(12.4)(9.8)\cos 105°$ **[M1]**

$AC^2 = 153.76 + 96.04 - (-62.90)$ or 312.7 **[M1]**

17.7 (accept $17.68\ldots$) **[A1]**

(b) $\dfrac{CD}{\sin 41°} = \dfrac{17.68}{\sin 68°}$ **[M1]**

$CD = \dfrac{17.68 \times \sin 41°}{\sin 68°}$ **[M1]**

12.5 (accept $12.51\ldots$) **[A1]**

(c) Area $ABC = \dfrac{1}{2} \times 12.4 \times 9.8 \times \sin 105° = 58.68\ldots$ **[B1]**

Area $ACD = \dfrac{1}{2} \times 17.68 \times 12.51 \times \sin(180° - 41° - 68°) = 104.6\ldots$ (accept correct alternative using $AD = 18.0$) **[M1]**

163 (accept 163.2 to 163.3) **[A1]**

[Back to Question] *[Total Marks: 9]*

1113 **(a)** $\dfrac{\sin C}{15} = \dfrac{\sin 48°}{12}$ **[M1]**

$\sin C = 0.9289\ldots$ **[A1]**

68.3 and 111.7 **[A1]**

(b) Calculates either possible angle B: $180° - 48° - 68.27° = 63.73°$ OR $180° - 48° - 111.73° = 20.27°$ **[M1]**

Area $1 = \dfrac{1}{2} \times 15 \times 12 \times \sin 63.73° = 80.7\ldots$ **[M1]**

Area $2 = \dfrac{1}{2} \times 15 \times 12 \times \sin 20.27° = 31.2\ldots$ **[M1]**

49.5 (accept 49.5 to 49.6) **[A1]**

[Back to Question] *[Total Marks: 7]*

1114 **(a)** $\dfrac{\sin C}{10.5} = \dfrac{\sin 35°}{8.2}$ **[M1]**

$\sin C = \dfrac{10.5 \times \sin 35°}{8.2}$ OR $0.734...$ **[M1]**

$C = 180° - 47.3° = 132.7°$ (or 132.73 to 132.74) **[A1]**

(b) Angle $B = 180° - 35° - 132.7° = 12.3°$ **[M1]**

$\dfrac{AC}{\sin 12.3°} = \dfrac{8.2}{\sin 35°}$ OR $AC^2 = 10.5^2 + 8.2^2 - 2(10.5)(8.2)\cos 12.3°$ **[M1]**

3.04 (or 3.035 to 3.045) **[A1]**

(c) $0.5 \times 3.04 \times 6.4 \times \sin(ACD) = 8.5$ **[M1]**

$\sin(ACD) = \dfrac{8.5}{9.728}$ OR Angle $ACD = \sin^{-1}\left(\dfrac{8.5}{9.728}\right) = 60.9°$ (or $60.8...$ to $61.1°$) **[A1]**

[Back to Question] *[Total Marks: 8]*

1115 **(a)** $\dfrac{\sin F}{10.5} = \dfrac{\sin 42^\circ}{8.2}$ **[M1]**

$\sin F = 0.8568\ldots$ yielding acute angle $58.96\ldots^\circ$ **[A1]**

$180^\circ - 58.96^\circ = 121.0$ (accept $121.04\ldots$) **[A1]**

(b) Angle $EDF = 180^\circ - 42^\circ - 121.04^\circ = 16.96^\circ$ **[B1]**

$\dfrac{EF}{\sin 16.96^\circ} = \dfrac{8.2}{\sin 42^\circ}$ **[M1]**

3.57 (accept $3.56\ldots$ to $3.58\ldots$) **[A1]**

(c) Shortest distance $= 10.5 \times \sin 42^\circ$ **[M1]**

7.03 (accept $7.025\ldots$ to $7.027\ldots$) **[A1]**

[Back to Question] *[Total Marks: 8]*

1116 **(a)** $BD^2 = 6^2 + 8^2 - 2(6)(8)\cos(110^\circ)$ **[M1]**

$BD^2 = 100 - 96(-0.342...) = 132.83...$ **[M1]**

$BD = 11.52...$ so 11.5 **[A1]**

(b) $\cos C = \dfrac{9^2 + 5^2 - (11.525...)^2}{2(9)(5)}$ **[M1]**

$\cos C = \dfrac{106 - 132.83...}{90} = -0.2981...$ **[M1]**

$C = 107.34...^\circ$ so 107.3° **[A1]**

(c) Area of triangle $ABD = \dfrac{1}{2} \times 6 \times 8 \times \sin(110^\circ) = 22.55...$ **[M1]**

Area of triangle $BCD = \dfrac{1}{2} \times 9 \times 5 \times \sin(107.34...^\circ) = 21.47...$ **[M1]**

Total Area $= 22.55... + 21.47... = 44.02...$ so 44.0 **[A1]**

[Back to Question] *[Total Marks: 9]*

1117 **(a)** $7^2 + 9^2 - 2 \times 7 \times 9 \times \cos 110^\circ$ **[M1]**

$173.09\ldots$ **[A1]**

13.2 (accept 13.15 to 13.2) **[A1]**

(b) $13.156^2 = 10^2 + 14^2 - 2 \times 10 \times 14 \times \cos(\angle BCD)$ **[M1]**

$\cos(\angle BCD) = \dfrac{10^2 + 14^2 - 173.09}{2 \times 10 \times 14}$ **[M1]**

64.0 (accept $63.9\ldots$ to 64.0) **[A1]**

(c) $\dfrac{1}{2} \times 7 \times 9 \times \sin 110^\circ$ (evaluates to $29.6\ldots$) **[M1]**

$\dfrac{1}{2} \times 10 \times 14 \times \sin 64.0^\circ$ (evaluates to $62.9\ldots$) **[M1]**

92.5 (accept 92.4 to 92.6) **[A1]**

[Back to Question] *[Total Marks: 9]*

1118 **(a)** $AC^2 = 7.4^2 + 9.1^2 - 2(7.4)(9.1)\cos(112°)$ [M2]
(or [M1] for correct cosine rule formula with one error)
$AC = 13.7$ (13.71 to 13.72) **[A1]**

(b) $\dfrac{\sin D}{13.712} = \dfrac{\sin 43°}{10.5}$ **[M1]**
$\sin D = \dfrac{13.712 \sin 43°}{10.5}$ or 0.890... **[M1]**
63.0 (62.95 to 63.0) **[A1]**

(c) Angle $ACD = 180 - 43 - 62.95 = 74.05°$ **[M1]**
Shortest distance $= 10.5 \times \sin(74.05°)$ **[M1]**
10.1 (10.09 to 10.1) **[A1]**

[Back to Question] *[Total Marks: 9]*

1119 **(a)** $BD^2 = 7.5^2 + 9.2^2 - 2(7.5)(9.2)\cos(105°)$ **[M2]**
($M1$ for correct substitution into implicit cosine rule)
13.3 (13.28...) **[A1]**

(b) $\cos C = \dfrac{13^2 + 10.5^2 - 13.289^2}{2(13)(10.5)}$ **[M2]**
($M1$ for correct substitution into implicit cosine rule)
67.9 (67.91...) **[A1]**

(c) Area $ABD = \dfrac{1}{2} \times 7.5 \times 9.2 \times \sin(105°)$ (33.32...) **[M1]**
Area $BCD = \dfrac{1}{2} \times 13 \times 10.5 \times \sin(67.9°)$ (63.24...) **[M1]**
96.6 (96.5... to 96.6) **[A1]**

[Back to Question] *[Total Marks: 9]*

1120 **(a)** Uses cosine rule: $BD^2 = 8.5^2 + 11.2^2 - 2(8.5)(11.2)\cos 105°$ **[M1]**
$BD^2 = 246.96...$ **[A1]**
15.7 **[A1]**

(b) Uses sine rule: $\dfrac{\sin C}{15.715} = \dfrac{\sin 38°}{11.0}$ **[M1]**
Acute angle $C = 61.59...°$ **[M1]**
Obtuse angle $BCD = 180° - 61.59...° = 118.4°$ **[A1]**

(c) Area of triangle $ABD = \dfrac{1}{2} \times 8.5 \times 11.2 \times \sin 105°$ $(= 45.97...)$ **[M1]**
Angle $DBC = 180° - 38° - 118.41° = 23.59°$ **[B1]**
Area of triangle $BCD = \dfrac{1}{2} \times 11.0 \times 15.715 \times \sin 23.59°$ $(= 34.59...)$ **[M1]**
80.6 **[A1]**

[Back to Question] *[Total Marks: 10]*

1121 **(a)** Angle $BAL = 085° - 050° = 35°$ **[B1]**

$\dfrac{\sin(ALB)}{40} = \dfrac{\sin 35°}{35}$ **[M1]**

$\sin(ALB) = \dfrac{40 \sin 35°}{35} = 0.6555\ldots$ **[M1]**

Acute angle $= 40.96°$, Obtuse angle $ALB = 180° - 40.96° = 139.0$ **[A1]**

(b) Angle $ABL = 180° - 35° - 139.04° = 5.96°$ **[B1]**

$\dfrac{AL}{\sin 5.96°} = \dfrac{35}{\sin 35°}$ **[M1]**

6.33 **[A1]**

(c) Bearing of A from $B = 180° + 050° = 230°$ **[M1]**

Bearing of L from $B = 230° -$ Angle $ABL = 230° - 5.96°$ **[M1]**

224.0 **[A1]**

[Back to Question] *[Total Marks: 10]*

1122 **(a)** $AC^2 = 8^2 + 6^2$ **[M1]**

$AC = 10$ **[A1]**

(b) $AG^2 = 10^2 + 5^2$ (or $AG^2 = 8^2 + 6^2 + 5^2$) **[M1]**

$AG = \sqrt{125} = 11.18...$ so 11.2 **[A1]**

(c) $\tan\theta = \dfrac{5}{10}$ (or $\sin\theta = \dfrac{5}{\sqrt{125}}$, etc.) **[M1]**

$\theta = 26.56...$ so $26.6°$ **[A1]**

[Back to Question] *[Total Marks: 6]*

1123 **(a)** Uses Pythagoras to find half diagonal AM: $AM = \dfrac{1}{2}\sqrt{10^2 + 10^2} = \sqrt{50}$ (or 7.07...) **[M1]**

$VA^2 = 12^2 + (\sqrt{50})^2 = 144 + 50 = 194$ **[M1]**

$VA = \sqrt{194} = 13.92...$ so 13.9 **[A1]**

(b) $\tan(\angle VAM) = \dfrac{12}{\sqrt{50}}$ **[M1]**

$\angle VAM = 59.49...$ so $59.5°$ **[A1]**

[Back to Question] *[Total Marks: 5]*

1124 **(a)** Uses Pythagoras to find AC: $AC = \sqrt{12^2 + 5^2} = 13$ **[M1]**

$AF^2 = 13^2 + 8^2 = 169 + 64 = 233$ **[M1]**

$AF = \sqrt{233} = 15.26...$ so 15.3 **[A1]**

(b) Identifies required angle as $\angle FAC$ and uses trig, e.g. $\tan(\angle FAC) = \frac{8}{13}$ **[M1]**

$\angle FAC = 31.60...$ so $31.6°$ **[A1]**

(c) Recognises the required angle is $\angle FAE$ because FE is perpendicular to plane $ABED$ **[B1]**

Uses trig in triangle FAE (where $\angle FEA = 90°$): $\sin(\angle FAE) = \frac{5}{\sqrt{233}}$ OR finds $AE = \sqrt{12^2 + 8^2} = \sqrt{208}$ and uses $\tan(\angle FAE) = \frac{5}{\sqrt{208}}$ **[M1]**

$\angle FAE = 19.11...$ so $19.1°$ **[A1]**

[Back to Question] *[Total Marks: 8]*

1125 **(a)** Correct triangle B with vertices at $(-3, 3), (-1, 3), (-1, 6)$ **[B2]**

Translation by vector $\begin{pmatrix} -4 \\ k \end{pmatrix}$ or $\begin{pmatrix} k \\ 2 \end{pmatrix}$ **[B1]**

(b) Correct triangle C with vertices at $(-3, 1), (-5, 1), (-5, 4)$ **[B2]**

Reflection in $y = -1$ or correct orientation but wrong position **[B1]**

[Back to Question] *[Total Marks: 4]*

1126 **(a)** Correct translation of Shape A **[B2]**

If incorrect, translation of $\begin{pmatrix} -4 \\ k \end{pmatrix}$ or $\begin{pmatrix} k \\ 2 \end{pmatrix}$ where $k \neq 0$ **[B1]**

(b) Correct reflection of Shape A to form Shape C with vertices at $(1, -3)$, $(3, -3)$, and $(1, -6)$ **[B2]**

If incorrect, reflection in $x = -1$ or $y = 1$ **[B1]**

[Back to Question] *[Total Marks: 4]*

1127 **(a)** Correct triangle with vertices at $(-3, 3), (-1, 3)$ and $(-1, 6)$ **[B2]**

Translation by correct vector but incorrect shape/orientation OR correct size, shape and orientation but wrong translation **[B1]**

(b) Reflection **[B1]**

in the y-axis OR $x = 0$ **[B1]**

[Back to Question] *[Total Marks: 4]*

1128 **(a)** Vertices of image plotted correctly at $(-1, -4)$, $(1, -4)$ and $(1, -1)$ **[B2]**

If not B2, award B1 for correct size and orientation, or 2 vertices correct **[B1]**

(b) Reflection **[B1]**

in the y-axis (or $x = 0$) **[B1]**

[Back to Question] *[Total Marks: 4]*

1129 **(a)** Image B drawn correctly at $(-4, 3), (-2, 3), (-2, 6)$ **[B2]**

If incorrect, 1 mark for translation by $\begin{pmatrix} -5 \\ k \end{pmatrix}$ or $\begin{pmatrix} k \\ 2 \end{pmatrix}$ **[M1]**

(b) Image C drawn correctly at $(1, -1), (3, -1), (3, -4)$ **[B2]**

If incorrect, 1 mark for reflection in y-axis or in a line $y = k$ **[M1]**

(c) Image D drawn correctly at $(-1, -1), (-3, -1), (-3, -4)$ **[B2]**

If incorrect, 1 mark for rotation of 180° about the wrong centre, or 90° about $(0, 0)$ **[M1]**

[Back to Question] *[Total Marks: 6]*

1130 **(a)** Correct shape with vertices at $(-2, -4), (-6, -4), (-6, -6), (-4, -6), (-4, -8)$ and $(-2, -8)$ **[B2]**

Enlargement, scale factor 2 centre $(0, 0)$ OR correct scale factor and orientation but wrong centre **[B1]**

(b) Correct shape with vertices at $(4, -1), (4, -3), (5, -3), (5, -2), (6, -2)$ and $(6, -1)$ **[B2]**

Rotation of 90° clockwise about the wrong centre OR rotation of 90° anti-clockwise about $(1, -1)$ **[B1]**

[Back to Question] *[Total Marks: 4]*

1131 **(a)** Rotation **[B1]**

90° clockwise OR 270° anti-clockwise **[B1]**

Centre $(0, 0)$ **[B1]**

(b) Correct triangle drawn with vertices at $(0, 2)$, $(1.5, 2)$, and $(1.5, 3)$ **[B2]**

If incorrect, correct size and orientation but wrong position OR correct centre of enlargement but scale factor 2 **[B1]**

[Back to Question] *[Total Marks: 5]*

1132 **(a)** Image B drawn correctly at $(2, -4), (2, -2), (6, -2)$ **[B2]**

If incorrect, 1 mark for reflection in $y = -x$ or correct orientation but wrong position **[M1]**

(b) Image C drawn correctly at $(-4, 2), (-2, 2), (-2, -2)$ **[B2]**

If incorrect, 1 mark for rotation 90° anti-clockwise about $(2, 2)$ or 90° clockwise about the wrong centre **[M1]**

(c) Reflection **[B1]**

in the line $y = 2$ **[B1]**

[Back to Question] *[Total Marks: 6]*

1133 **(a)** Vertices of image plotted correctly at $(-1, 2)$, $(-2, 2)$ and $(-2, 0.5)$ **[B2]**

If not B2, award B1 for enlargement SF $-\frac{1}{2}$ from wrong centre, or correct centre but SF $\frac{1}{2}$ **[B1]**

(b) Rotation **[B1]**

$180°$ (direction not required) **[B1]**

centre $(1, 2)$ **[B1]**

(c) Calculates area of $T = \frac{1}{2} \times 2 \times 3 = 3$ **[M1]**

Sets up $3 \times k^2 = 48 \implies k^2 = 16$ **[M1]**

$k = 4$ **[A1]**

[Back to Question] *[Total Marks: 7]*

1134 **(a)** Enlargement **[B1]**

Scale factor -2 **[B1]**

Centre $(0, 1)$ **[B1]**

(b) Enlargement scale factor $-\frac{1}{2}$ **[B1]**

Centre $(0, 1)$ **[B1]**

(c) Translating Shape T gives at least two correct coordinates of V: $(-2, 3)$, $(-1, 3)$, $(-1, 5)$ **[M1]**

Recognising reflection in $y = x$ swaps coordinates (x, y)

(y, x)$[M1]$

Final coordinates: $(3, -2)$, $(3, -1)$, $(5, -1)$ **[A1]**

[Back to Question] *[Total Marks: 8]*

1135 **(a)** Rotation **[B1]**

$90°$ anticlockwise OR $270°$ clockwise **[B1]**

Centre $(0, 0)$ **[B1]**

(b) Correct shape R with vertices at $(4, 4), (5, 4), (5, 5), (4.5, 5), (4.5, 4.5), (4, 4.5)$ **[B2]**

Any enlargement scale factor $\frac{1}{2}$ OR correct centre $(6, 6)$ with incorrect scale factor **[B1]**

[Back to Question] *[Total Marks: 5]*

1136 **(a)** Vertices of image plotted correctly at $(1, 1)$, $(2, 1)$, $(2, 1.5)$, $(1.5, 1.5)$, $(1.5, 2)$ and $(1, 2)$ **[B2]**

If not B2, award B1 for correct size and orientation, or correct enlargement from the wrong centre **[B1]**

(b) Rotation **[B1]**

$90°$ clockwise (or $-90°$) **[B1]**

centre $(0, 0)$ (or origin) **[B1]**

[Back to Question] *[Total Marks: 5]*

1137 **(a)** Enlargement **[B1]**

Centre $(0, 0)$ or origin **[B1]**

Scale factor $-\frac{1}{2}$ **[B1]**

(b) Image R drawn correctly at $(1, 3), (3, 3), (3, 4)$ **[B3]**

If incorrect, 2 marks for correct size and orientation but wrong position **[M2]**

1 mark for correct centre used but wrong scale factor **[M1]**

(c) Rotation **[B1]**

$180°$ **[B1]**

Centre $(0, 1)$ **[B1]**

OR Enlargement [B1], Scale factor -1 [B1], Centre $(0, 1)$ **[B1]**

[Back to Question] *[Total Marks: 9]*

1138 **(a)** Enlargement **[B1]**

Scale factor $-\frac{1}{2}$ **[B1]**

Centre $(0, 0)$ **[B1]**

(b) Reflection **[B1]**

in the x-axis OR $y = 0$ **[B1]**

[Back to Question] *[Total Marks: 5]*

1139 **(a)** Correct triangle U with vertices at $(1, 2), (3, 2)$ and $(3, 0)$ **[B2]**

Enlargement scale factor $\frac{1}{2}$ from wrong centre OR ray lines drawn from $(0, 4)$ but scale factor incorrect **[B1]**

(b) Correct triangle V with vertices at $(2, 1), (2, 3)$ and $(0, 3)$ **[B2]**

Reflection of U in the line $y = -x$ OR reflection of T in the line $y = x$ **[B1]**

(c) Enlargement **[B1]**

Scale factor 2 **[B1]**

Centre $(0, 4)$ **[B1]**

[Back to Question] *[Total Marks: 7]*

1140 **(a)** $\begin{pmatrix} 8 \\ -6 \end{pmatrix} + \begin{pmatrix} -2 \\ 5 \end{pmatrix}$ **[M1]**

$\begin{pmatrix} 6 \\ -1 \end{pmatrix}$ **[A1]**

(b) $(5, 1)$ **[B1]**

[Back to Question] *[Total Marks: 3]*

1141 **(a)** $\begin{pmatrix} 4 \\ -3 \end{pmatrix} + \begin{pmatrix} -4 \\ 10 \end{pmatrix}$ **[M1]**

$\begin{pmatrix} 0 \\ 7 \end{pmatrix}$ **[A1]**

(b) $\begin{pmatrix} 3 \\ -1 \end{pmatrix} + \begin{pmatrix} 4 \\ -3 \end{pmatrix} = \begin{pmatrix} 7 \\ -4 \end{pmatrix}$ [M1]

$(7, -4)$ **[A1]**

[Back to Question] *[Total Marks: 4]*

1142 **(a)** $3 \times \begin{pmatrix} 4 \\ -3 \end{pmatrix}$ **[M1]**

$\begin{pmatrix} 12 \\ -9 \end{pmatrix}$ **[A1]**

(b) $\begin{pmatrix} 4 \\ -3 \end{pmatrix} - \begin{pmatrix} -4 \\ 10 \end{pmatrix}$ or $\begin{pmatrix} 4 - 2(-2) \\ -3 - 2(5) \end{pmatrix}$ **[M1]**

$\begin{pmatrix} 8 \\ -13 \end{pmatrix}$ **[A1]**

(c) $\begin{pmatrix} x \\ y \end{pmatrix} + \begin{pmatrix} -2 \\ 5 \end{pmatrix} = \begin{pmatrix} 6 \\ -1 \end{pmatrix}$ or $(6 - (-2), -1 - 5)$ **[M1]**

$(8, -6)$ **[A1]**

[Back to Question] *[Total Marks: 6]*

1143 **(a)** **(i)** $\vec{OA} + \frac{1}{2}\vec{AB} = \mathbf{a} + \frac{1}{2}\mathbf{c}$ **[M1]**

$\mathbf{a} + \frac{1}{2}\mathbf{c}$ **[A1]**

(ii) $\vec{MB} + \vec{BN} = \frac{1}{2}\mathbf{c} - \frac{2}{3}\mathbf{a}$ **[M1]**

$\frac{1}{2}\mathbf{c} - \frac{2}{3}\mathbf{a}$ **[A1]**

(b) $\vec{ON} = \vec{OC} + \vec{CN} = \mathbf{c} + \frac{1}{3}\mathbf{a}$ **[M1]**

$\frac{1}{3}$ **[A1]**

[Back to Question] *[Total Marks: 6]*

1144 **(a)** $\begin{pmatrix} 8 \\ -1 \end{pmatrix}$ **[B1]**

(b) Deduces correct column vectors for **a** and **b**: $\mathbf{a} = \begin{pmatrix}2\\1\end{pmatrix}$ and $\mathbf{b} = \begin{pmatrix}-1\\2\end{pmatrix}$ **[B1]**

Sets up correct simultaneous equations from components: $2p - q = 8$ and $p + 2q = -1$ **[M1]**

Valid method to solve the simultaneous equations (e.g. multiplying first by 2 and adding: $4p - 2q = 16$

5p = 15)$[M1]$

$p = 3$ **[A1]**

$q = -2$ **[A1]**

[Back to Question] *[Total Marks: 5]*

1145 (a) $\begin{pmatrix}12\\-9\end{pmatrix}$ or $\begin{pmatrix}-4\\10\end{pmatrix}$ seen **[M1]**

$\begin{pmatrix}16\\-19\end{pmatrix}$ **[A1]**

(b) $\vec{AB} = \begin{pmatrix}2\\2\end{pmatrix}$ **[M1]**

$(5, 1)$ **[A1]**

[Back to Question] *[Total Marks: 4]*

1146 (a) $\vec{AB} = \mathbf{b} - \mathbf{a}$ or $\vec{BA} = \mathbf{a} - \mathbf{b}$ **[B1]**

$\vec{OP} = \mathbf{a} + \frac{3}{4}(\mathbf{b} - \mathbf{a})$ or $\vec{OP} = \mathbf{b} + \frac{1}{4}(\mathbf{a} - \mathbf{b})$ **[M1]**

$\frac{1}{4}\mathbf{a} + \frac{3}{4}\mathbf{b}$ **[A1]**

(b) $\vec{MP} = -\frac{1}{2}\mathbf{a} + \vec{OP}$ (oe) **[M1]**

$-\frac{1}{4}\mathbf{a} + \frac{3}{4}\mathbf{b}$ **[A1]**

[Back to Question] *[Total Marks: 5]*

1147 (a) Forms a correct equation for the horizontal components: $3(4) - k = 10$ **[M1]**

$k = 2$ **[A1]**

Forms a correct equation for the vertical components: $3(-3) + 2y = 6$

-9 + 2y = 6$[M1]$

$y = 7.5$ **[A1]**

(b) (i) $\begin{pmatrix}3\\6\end{pmatrix}$ **[B1]**

(ii) Finds vector $\vec{QR} = \begin{pmatrix}6\\12\end{pmatrix}$ OR $\vec{PR} = \begin{pmatrix}9\\18\end{pmatrix}$ **[M1]**

States that $\vec{QR} = 2\vec{PQ}$ (or equivalent scalar multiplier), showing the vectors are parallel, AND mentions that they share a common point Q (or P), concluding they are collinear **[A1]**

[Back to Question] *[Total Marks: 6]*

1148 **(a)** $\begin{pmatrix}7\\10\end{pmatrix} - \begin{pmatrix}1\\-2\end{pmatrix}$ **[M1]**

$\begin{pmatrix}6\\12\end{pmatrix}$ **[A1]**

(b) $\vec{AC} = \begin{pmatrix}4-1\\4-(-2)\end{pmatrix} = \begin{pmatrix}3\\6\end{pmatrix}$ **[M1]**

$\frac{1}{2}\vec{AB} = \frac{1}{2}\begin{pmatrix}6\\12\end{pmatrix} = \begin{pmatrix}3\\6\end{pmatrix}$, hence $\vec{AC} = \frac{1}{2}\vec{AB}$ **[A1]**

(c) $(4-5, 4+2)$ **[M1]**

$(-1, 6)$ **[A1]**

(d) $\begin{pmatrix}-1\\6\end{pmatrix} - \begin{pmatrix}7\\10\end{pmatrix}$ or equivalent path method $-\vec{AC} + \vec{CD}$ **[M1]**

$\begin{pmatrix}-8\\-4\end{pmatrix}$ **[A1]**

[Back to Question] *[Total Marks: 8]*

1149 **(a)** $\vec{AX} = \frac{1}{3}\mathbf{c}$ **[M1]**

$\mathbf{a} + \frac{1}{3}\mathbf{c}$ **[A1]**

(b) $\vec{BY} = 2\mathbf{a}$ or $\vec{CY} = 3\mathbf{a}$ **[M1]**

$3\mathbf{a} + \mathbf{c}$ **[A1]**

(c) $\vec{OY} = 3\mathbf{a} + \mathbf{c} = 3(\mathbf{a} + \frac{1}{3}\mathbf{c}) = 3\vec{OX}$ **[M1]**

Since $\vec{OY}$ is a scalar multiple of $\vec{OX}$ and they share the common point O, they are collinear. **[A1]**

(d) $1 : 2$ **[B1]**

[Back to Question] *[Total Marks: 7]*

1150 **(a)** $\begin{pmatrix}6\\7\end{pmatrix}$ - $\begin{pmatrix}2\\-1\end{pmatrix}$ $[M1]$

$\begin{pmatrix}4\\8\end{pmatrix}$ **[A1]**

(b) $\vec{AC} = \begin{pmatrix}k-1\\3k-1\end{pmatrix}$ **[B1]**

Gradient or ratio condition: $\dfrac{3k-1}{k-1} = \dfrac{8}{4}$ or $\vec{AC} = m\vec{AB}$ **[M1]**

$3k - 1 = 2(k-1) \implies k = -1$ **[A1]**

(c) $\vec{AC} = \begin{pmatrix} -2 \\ -4 \end{pmatrix}$ **[M1]**

$-\dfrac{1}{2}\vec{AB}$ **[A1]**

[Back to Question] *[Total Marks: 7]*

1151 **(a)** $\vec{PQ} = 2\begin{pmatrix} 3 \\ -2 \end{pmatrix} - \begin{pmatrix} -1 \\ 4 \end{pmatrix} = \begin{pmatrix} 7 \\ -8 \end{pmatrix}$ **[M1]**

$\begin{pmatrix} x \\ y \end{pmatrix} - \begin{pmatrix} 5 \\ -1 \end{pmatrix} = \begin{pmatrix} 7 \\ -8 \end{pmatrix}$ or $(5+7, -1-8)$ **[M1]**

$(12, -9)$ **[A1]**

(b) $\vec{PR} = \begin{pmatrix} 14-5 \\ 3-(-1) \end{pmatrix} = \begin{pmatrix} 9 \\ 4 \end{pmatrix}$ **[B1]**

$3h - k = 9$ and $-2h + 4k = 4$ **[M1]**

Rearranging to get $k = 3h - 9$ and substituting: $-2h + 4(3h-9) = 4$ or equivalent valid method to solve simultaneous equations **[M1]**

$10h = 40 \implies h = 4$ **[A1]**

$k = 3(4) - 9 = 3$ **[A1]**

[Back to Question] *[Total Marks: 8]*

1152 $\sqrt{5^2 + (-12)^2}$ or $\sqrt{25 + 144}$ **[M1]**

13 **[A1]**

[Back to Question] *[Total Marks: 2]*

1153 **(a)** Determines column vector $\mathbf{a} = \begin{pmatrix} -6 \\ 8 \end{pmatrix}$ **[M1]**

10 **[A1]**

(b) Determines column vector $\mathbf{b} = \begin{pmatrix} 3 \\ -4 \end{pmatrix}$ **[B1]**

Calculates resultant vector $2\begin{pmatrix} -6 \\ 8 \end{pmatrix} + 3\begin{pmatrix} 3 \\ -4 \end{pmatrix} = \begin{pmatrix} -12 \\ 16 \end{pmatrix} + \begin{pmatrix} 9 \\ -12 \end{pmatrix} = \begin{pmatrix} -3 \\ 4 \end{pmatrix}$ **[M1]**

5 **[A1]**

[Back to Question] *[Total Marks: 5]*

1154 **(a)** $\sqrt{(-8)^2 + 15^2}$ or $\sqrt{64 + 225}$ **[M1]**

17 **[A1]**

(b) $\mathbf{p} + 2\mathbf{q} = \begin{pmatrix} -8 \\ 15 \end{pmatrix} + \begin{pmatrix} 4 \\ -8 \end{pmatrix} = \begin{pmatrix} -4 \\ 7 \end{pmatrix}$ **[M1]**

$\sqrt{(-4)^2 + 7^2}$ or $\sqrt{16 + 49}$ **[M1]**

$\sqrt{65}$ **[A1]**

[Back to Question] *[Total Marks: 5]*

1155 **(a)** Sets up equation: $p^2 + p^2 = 18$ **[M1]**

3 **[A1]**

(b) Finds expression for $\mathbf{w}$: $\begin{pmatrix} 4 - t \\ 1 + 2t \end{pmatrix}$ **[M1]**

Forms magnitude squared equation: $(4 - t)^2 + (1 + 2t)^2 = 29$ **[M1]**

Expands brackets correctly: $16 - 8t + t^2 + 1 + 4t + 4t^2 = 29$ **[M1]**

Rearranges to a 3-term quadratic: $5t^2 - 4t - 12 = 0$ **[M1]**

$t = 2$, or $t = -1.2$ (or $-\frac{6}{5}$) **[A1]**

[Back to Question] *[Total Marks: 7]*

1156 **(a)** $\sqrt{x^2 + (x + 2)^2} = 10$ or $x^2 + (x + 2)^2 = 10^2$ **[M1]**

$x^2 + x^2 + 4x + 4 = 100$ **[M1]**

$2x^2 + 4x - 96 = 0$ leading to $x^2 + 2x - 48 = 0$ with no errors seen **[A1]**

(b) $(x + 8)(x - 6) = 0$ or correct substitution into quadratic formula **[M1]**

$x = -8, x = 6$ **[A1]**

(c) $\mathbf{u} = \begin{pmatrix} -8 \\ -6 \end{pmatrix}$ **[B1]**

$\mathbf{u} = \begin{pmatrix} 6 \\ 8 \end{pmatrix}$ **[B1]**

[Back to Question] *[Total Marks: 7]*

1157 **(a)** $\vec{AC} = \begin{pmatrix} 7 - 3 \\ 1 - (-2) \end{pmatrix}$ or $\begin{pmatrix} 4 \\ 3 \end{pmatrix}$ **[M1]**

5 **[A1]**

(b) $\vec{AB} = \begin{pmatrix} -8 \\ y + 2 \end{pmatrix}$ **[B1]**

$\sqrt{(-8)^2 + (y + 2)^2} = 17$ **[M1]**

$64 + (y + 2)^2 = 289$ or $(y + 2)^2 = 225$ **[M1]**

$y + 2 = 15$ or $y + 2 = -15$ **[M1]**

$y = 13$, $y = -17$ **[A1]**

[Back to Question] *[Total Marks: 7]*

1158 **(a)** Expresses vector $\vec{PQ}$ as $\begin{pmatrix} k-2 \\ 5-(-3) \end{pmatrix} = \begin{pmatrix} k-2 \\ 8 \end{pmatrix}$ **[M1]**

Sets up magnitude equation: $(k-2)^2 + 8^2 = 10^2$ **[M1]**

Simplifies to solve for k: $(k-2)^2 = 36$ **[M1]**

$k = 8$, or $k = -4$ **[A1]**

(b) Finds $\vec{QR}$ using $Q(8,5)$ and $R(4,1)$: $\vec{QR} = \begin{pmatrix} 4-8 \\ 1-5 \end{pmatrix} = \begin{pmatrix} -4 \\ -4 \end{pmatrix}$ **[M1]**

Calculates magnitude: $\sqrt{(-4)^2 + (-4)^2} = \sqrt{32}$ **[M1]**

5.66 **[A1]**

[Back to Question] *[Total Marks: 7]*

1159 **(a)** $\begin{pmatrix} 9-2 \\ 4-(-3) \end{pmatrix}$ **[M1]**

$\begin{pmatrix} 7 \\ 7 \end{pmatrix}$ **[A1]**

(b) $\sqrt{7^2 + 7^2}$ or $\sqrt{98}$ **[M1]**

9.90 **[A1]**

(c) $\sqrt{k^2 + 4^2} = 5$ or $k^2 + 4^2 = 5^2$ **[M1]**

$k^2 = 25 - 16$ or $k^2 = 9$ **[M1]**

$k = 3$, $k = -3$ **[A1]**

[Back to Question] *[Total Marks: 7]*

1160 **(a)** $\mathbf{u} + \mathbf{v} = \begin{pmatrix} 3k+1 \\ k-1 \end{pmatrix}$ **[M1]**

$(3k+1)^2 + (k-1)^2 = (\sqrt{50})^2$ **[M1]**

$9k^2 + 6k + 1 + k^2 - 2k + 1 = 50$ **[M1]**

$10k^2 + 4k + 2 = 50 \implies 5k^2 + 2k - 24 = 0$ **[A1]**

(b) $(5k + a)(k + b) = 0$ where $ab = -24$ or $5b + a = 2$, or correct substitution into quadratic formula **[M1]**

$(5k + 12)(k - 2) = 0$ **[M1]**

$k = 2$ **[A1]**

$k = -2.4$ or $-\frac{12}{5}$ **[A1]**

(c) Substituting $k = 2$ and $k = -2.4$ into $\begin{pmatrix} 2k \\ k-3 \end{pmatrix}$ **[M1]**

$\begin{pmatrix} 4 \\ -1 \end{pmatrix}$ and $\begin{pmatrix} -4.8 \\ -5.4 \end{pmatrix}$ **[A1]**

[Back to Question] *[Total Marks: 10]*

1161 **(a)** 0.17 **[B1]**

(b) The train does not arrive on time (or equivalent) **[B1]**

[Back to Question] *[Total Marks: 2]*

1162 **(a)** $0.15 + 2x + 0.31 + x + 0.24 = 1$ or $3x + 0.7 = 1$ **[M1]**

0.1 **[A1]**

(b) $P(2) = 2(0.1) = 0.2$ **[M1]**

0.8 **[A1]**

[Back to Question] *[Total Marks: 4]*

1163 **(a)** Sum of given probabilities: $0.15 + 0.3 + 0.45 = 0.9$ **[M1]**

0.1 **[A1]**

(b) 0.45 **[B1]**

(c) Uses $1 - P(G)$ formula: $1 - 0.45$ **[M1]**

0.55 **[A1]**

[Back to Question] *[Total Marks: 5]*

1164 **(a)** $\frac{1}{5} + 0.35 + x + 2x = 1$ or $0.55 + 3x = 1$ **[M1]**

0.15 **[A1]**

(b) $1 - 2(0.15)$ or $\frac{1}{5} + 0.35 + 0.15$ **[M1]**

0.7 or $\frac{7}{10}$ **[A1]**

[Back to Question] *[Total Marks: 4]*

1165 **(a)** Places $2x$ in the intersection **[B1]**

Places $38 - 2x$ in T only region and $24 - 2x$ in C only region **[B1]**

Places x in the region outside both circles **[B1]**

(b) Forms correct equation: $(38 - 2x) + 2x + (24 - 2x) + x = 60$ or $62 - x = 60$ **[M1]**

2 **[A1]**

(c) Finds the number of people in $T \cap C'$: $38 - 2(2) = 34$ **[M1]**

$\frac{34}{60}$ or $\frac{17}{30}$ **[A1]**

[Back to Question] *[Total Marks: 7]*

1166 **(a)** $x^2 + 3x + 2$ **[B1]**

(b) $\dfrac{x^2}{x^2+3x+2} = \dfrac{8}{15}$ **[M1]**

$15x^2 = 8(x^2+3x+2) \implies 15x^2 = 8x^2+24x+16$ **[M1]**

(c) $(7x+4)(x-4)=0$ or correct substitution into the quadratic formula [M2]
(Award M1 for $(7x+a)(x+b)=0$ where $ab=-16$ or $7b+a=-24$)

4 **[A1]**

(d) Total counters $= 4^2+3(4)+2 = 30$ and number of blue counters $= 3(4) = 12$ **[M1]**

$\dfrac{18}{30} = \dfrac{3}{5}$ **[A1]**

[Back to Question] *[Total Marks: 8]*

1167 **(a)** $1-\left(\dfrac{2}{5}+\dfrac{1}{3}\right)$ or $1-\dfrac{11}{15}$ **[M1]**

$\dfrac{4}{15}$ **[A1]**

(b) Equates their fraction to the number of green marbles: $\dfrac{4}{15}$ of total $= 12$ **[M1]**

Total $= 12 \times \dfrac{15}{4}$ **[M1]**

45 **[A1]**

(c) $\dfrac{2}{3}$ **[B1]**

[Back to Question] *[Total Marks: 6]*

1168 **(a)** 21 in the intersection $C \cap D$ **[B1]**

34 in C only and 47 in D only **[B1]**

18 outside both circles **[B1]**

(b) $\dfrac{21}{120}$ or $\dfrac{7}{40}$ **[B1]**

(c) $1-\dfrac{68}{120}$ or $\dfrac{34+18}{120}$ **[M1]**

$\dfrac{52}{120}$ or $\dfrac{13}{30}$ **[A1]**

(d) $\dfrac{34+47}{120}$ or $\dfrac{120-21-18}{120}$ **[M1]**

$\dfrac{81}{120}$ or $\dfrac{27}{40}$ **[A1]**

[Back to Question] *[Total Marks: 8]*

1169 **(a)** $(x+4)+x+(2x-1)+(3x+5) = 50$ or $7x+8=50$ **[M1]**

6 **[A1]**

(b) $(2(6)-1)+(3(6)+5)$ or $50-(6+4+6)$ or $\dfrac{11+23}{50}$ **[M1]**

$\dfrac{34}{50}$ or $\dfrac{17}{25}$ or 0.68 **[A1]**

(c) $(6+4)+(2(6)-1)$ or $10+11$ or $\dfrac{21}{50}$ **[M1]**

$\dfrac{21}{50}$ or 0.42 **[A1]**

[Back to Question] *[Total Marks: 6]*

1170 **(a)** $0.15+0.25+x+0.2+3x=1$ **[M1]**

$x=0.1$ **[A1]**

(b) $P(>3)=P(4)+P(5)=0.2+3(0.1)=0.5$ **[M1]**

Expected number $=250\times 0.5=125$ **[A1]**

[Back to Question] *[Total Marks: 4]*

1171 **(a)** $\dfrac{70}{200}$ or $\dfrac{7}{20}$ or 0.35 **[B1]**

(b) $1000\times\dfrac{45}{200}$ **[M1]**

225 **[A1]**

(c) No, because the relative frequencies are not all close to 0.25 (or 50 out of 200) **[B1]**

(d) A larger number of trials leads to a relative frequency that is closer to the theoretical probability **[B1]**

[Back to Question] *[Total Marks: 5]*

1172 **(a)** $x+2x+5+3x-2+x+8=60$ **[M1]**

$7x+11=60 \implies 7x=49$

$x=7$ **[A1]**

(b) Frequency of $3=3(7)-2=19$

$\dfrac{19}{60}$ **[B1]**

(c) Frequency of $2=2(7)+5=19$

Relative frequency of $2=\dfrac{19}{60}$

$1200\times\dfrac{19}{60}$ **[M1]**

380 **[A1]**

[Back to Question] *[Total Marks: 5]*

1173 **(a)** $x = 1 - (0.15 + 0.45 + 0.10)$

$x = 0.30$ **[B1]**

(b) 300×0.45 **[M1]**

135 **[A1]**

(c) Biased, because the relative frequencies are not all close to 0.25 (or specifically Blue is 0.45, which is much higher than 0.25) **[B1]**

[Back to Question] *[Total Marks: 4]*

1174 **(a)** $\frac{x}{300} = 0.22 \implies x = 300 \times 0.22$ **[M1]**

$y = 300 - (84 + 105 + 66)$ **[M1]**

$x = 66$, $y = 45$ **[A1]**

(b) Relative frequency of Yellow $= \frac{45}{300}$ or 0.15 **[M1]**

Total beads $= \frac{15}{0.15} = 100$ **[A1]**

(c) Probability of Blue or Green $= \frac{105 + 66}{300} = \frac{171}{300}$ or 0.57 **[M1]**

$800 \times 0.57 = 456$ **[A1]**

[Back to Question] *[Total Marks: 7]*

1175 **(a)** $\frac{260}{800}$ **[M1]**

$\frac{13}{40}$ **[A1]**

(b) $260 + 180 + w + w + 40 = 800 \implies 480 + 2w = 800$ **[M1]**

$w = 160$ **[A1]**

(c) $\frac{160}{800}$ or $\frac{1}{5}$ or 0.2 **[B1]**

(d) $P(\text{Black}) = \frac{180}{800} = \frac{9}{40}$ **[M1]**

Expected number $= 12000 \times \frac{9}{40} = 2700$ **[A1]**

[Back to Question] *[Total Marks: 7]*

1176 **(a)** $0.15 + 2x + x + 0.05 + 0.2 + x = 1$ **[M1]**

$4x + 0.4 = 1 \implies 4x = 0.6 \implies x = 0.15$ **[A1]**

(b) $P(2, 3, 5) = 2(0.15) + (0.15 + 0.05) + 0.15 = 0.3 + 0.2 + 0.15 = 0.65$ **[M1]**

$600 \times 0.65 = 390$ **[A1]**

(c) Sets up both equations: $N \times p = 54$ and $N \times (1 - 3p) = 36$ **[M1]**

Expands the second equation: $N - 3Np = 36$ **[M1]**

Substitutes $Np = 54$ into the expanded equation: $N - 3(54) = 36$ **[M1]**

$N = 198$, $p = \dfrac{54}{198} = \dfrac{3}{11}$ (or 0.272...) **[A1]**

[Back to Question] *[Total Marks: 8]*

1177 **(a)** Sum of probabilities: $3x + 2y = 1 \implies y = \dfrac{1 - 3x}{2}$ **[B1]**

$E_1 = 400x$ and $E_3 = 400y \implies 400x \times 400y = 5600 \implies xy = \dfrac{7}{200}$ **[M1]**

Substitute for y: $x\left(\dfrac{1 - 3x}{2}\right) = \dfrac{7}{200} \implies 100x(1 - 3x) = 7 \implies 300x^2 - 100x + 7 = 0$

[A1]

(b) Factorising $(30x - 7)(10x - 1) = 0$ or correct substitution into the quadratic formula **[M1]**

$x = \dfrac{7}{30}$ and $x = \dfrac{1}{10}$ (or 0.1) **[A1]**

When $x = \dfrac{7}{30}$, $y = 0.15$ (reject as $y < x$). When $x = 0.1$, $y = 0.35$ (accept as $y > x$) **[M1]**

$x = 0.1, y = 0.35$ **[A1]**

(c) $P(\text{even}) = 2x + y = 2(0.1) + 0.35 = 0.55$ **[M1]**

Expected number $= 400 \times 0.55 = 220$ **[A1]**

[Back to Question] *[Total Marks: 9]*

1178 **(a)** $300p + 200(p + 0.15) = 255$ **[M1]**

$300p + 200p + 30 = 255$ **[M1]**

$500p = 225 \implies p = \dfrac{225}{500}$

$p = 0.45$ **[A1]**

(b) Probability Coin B is Heads $= 0.45 + 0.15 = 0.60$

Probability of different outcomes = Probability (H, T) + Probability (T, H)

$(0.45 \times 0.40) + (0.55 \times 0.60)$ **[M1]**

$0.18 + 0.33 = 0.51$ **[M1]**

$1000 \times 0.51 = 510$ **[A1]**

(c) **(i)** $\dfrac{72}{N} = 0.48$ **[M1]**

$N = \dfrac{72}{0.48}$

$N = 150$ **[A1]**

(ii) Probability of Tails $= 1 - 0.45 = 0.55$

150×0.55 **[M1]**

82.5 **[A1]**

[Back to Question] *[Total Marks: 10]*

1179 **(a)** $\frac{3}{8}$ correctly placed on the first branch for Blue **[B1]**

$\frac{5}{8}$ and $\frac{3}{8}$ correctly placed on all second branches **[B1]**

(b) $\frac{5}{8} \times \frac{5}{8}$ **[M1]**

$\frac{25}{64}$ **[A1]**

[Back to Question] *[Total Marks: 4]*

1180 **(a)** $\frac{3}{8}$ correctly placed for top branch (Red given Red) **[B1]**

$\frac{4}{8}$ correctly placed for bottom branches (Red given Blue, and Blue given Blue) **[B1]**

(b) $\frac{4}{9} \times \frac{5}{8}$ or $\frac{5}{9} \times \frac{4}{8}$ **[M1]**

$\frac{4}{9} \times \frac{5}{8} + \frac{5}{9} \times \frac{4}{8}$ **[M1]**

$\frac{5}{9}$ or equivalent (e.g. $\frac{40}{72}$) **[A1]**

[Back to Question] *[Total Marks: 5]*

1181 **(a)** $\frac{6}{9}$ on the Green branch from Green **[B1]**

$\frac{2}{9}$ and $\frac{7}{9}$ on the branches from Red **[B1]**

(b) $\frac{7}{10} \times \frac{6}{9}$ or $\frac{3}{10} \times \frac{2}{9}$ **[M1]**

$\frac{7}{10} \times \frac{6}{9} + \frac{3}{10} \times \frac{2}{9}$ **[M1]**

$\frac{48}{90}$ or $\frac{8}{15}$ **[A1]**

[Back to Question] *[Total Marks: 5]*

1182 **(a)** Intersection region = 6 **[B1]**

French only = 11, Spanish only = 8, outside = 5 **[B1]**

(b) **(i)** $\frac{6}{30}$ or $\frac{1}{5}$ **[B1]**

(ii) $\frac{11}{30}$ **[B1]**

(iii) $\frac{25}{30}$ or $\frac{5}{6}$ **[B1]**

[Back to Question] *[Total Marks: 5]*

1183 **(a)** Number of students studying both B and C: $(18 + 15) - (30 - 4) = 7$ **[M1]**

Fully correct Venn diagram: Only $B = 11$, $B \cap C = 7$, Only $C = 8$, Neither $= 4$ **[A1]**

(b) **(i)** $\frac{8}{30}$ or $\frac{4}{15}$ **[B1]**

(ii) $\frac{26}{30}$ or $\frac{13}{15}$ **[B1]**

(iii) $\frac{7}{18}$ **[B1]**

[Back to Question] *[Total Marks: 5]*

1184 **(a)** $\frac{3}{8}$ correctly placed for 1st ball Blue **[B1]**

$\frac{3}{8}$, $\frac{5}{8}$, and $\frac{3}{8}$ correctly placed on all three remaining 2nd ball branches **[B1]**

(b) $\frac{25}{64}$ **[B1]**

(c) $\left(\frac{5}{8} \times \frac{3}{8}\right) + \left(\frac{3}{8} \times \frac{5}{8}\right)$ **[M1]**

$\frac{30}{64}$ or $\frac{15}{32}$ **[A1]**

[Back to Question] *[Total Marks: 5]*

1185 **(a)** First counter branches: Red $\frac{5}{8}$, Blue $\frac{3}{8}$ **[B1]**

Second counter branches: Red $\frac{5}{8}$, Blue $\frac{3}{8}$ on both sets of branches **[B1]**

(b) $\frac{5}{8} \times \frac{5}{8} = \frac{25}{64}$ **[B1]**

(c) $\left(\frac{5}{8} \times \frac{3}{8}\right) + \left(\frac{3}{8} \times \frac{5}{8}\right)$ **[M1]**

$= \frac{30}{64}$ (or $\frac{15}{32}$) **[A1]**

[Back to Question] *[Total Marks: 5]*

1186 **(a)** $\frac{6}{9}$ and $\frac{3}{9}$ correctly placed on the top branches **[B1]**

$\frac{7}{9}$ and $\frac{2}{9}$ correctly placed on the bottom branches **[B1]**

(b) $\frac{7}{10} \times \frac{6}{9}$ or $\frac{3}{10} \times \frac{2}{9}$ **[M1]**

$\left(\frac{7}{10} \times \frac{6}{9}\right) + \left(\frac{3}{10} \times \frac{2}{9}\right)$ **[M1]**

$\frac{48}{90}$ or $\frac{8}{15}$ **[A1]**

[Back to Question] *[Total Marks: 5]*

1187 **(a)** Number studying both $= (18 + 15) - (30 - 5) = 8$ **[M1]**

Fully correct Venn diagram: B only $= 10$, $B \cap C = 8$, C only $= 7$, Outside $= 5$ **[A1]**

(b) **(i)** $\frac{10}{30}$ or $\frac{1}{3}$ **[B1]**

(ii) $\frac{25}{30}$ or $\frac{5}{6}$ **[B1]**

(c) Identifies total number of Chemistry students as denominator: 15 **[M1]**

$\frac{8}{15}$ **[A1]**

[Back to Question] *[Total Marks: 6]*

1188 **(a)** 8 in the intersection **[B1]**

16 in F only, 7 in C only, and 9 outside the circles **[B1]**

(b) Identifies 9 as the number of students playing neither **[M1]**

$\frac{9}{40}$ **[A1]**

(c) $16 + 8 + 7$ or $40 - 9$ **[M1]**

31 **[A1]**

[Back to Question] *[Total Marks: 6]*

1189 **(a)** First stage branches correctly labeled: $\frac{7}{12}$ for R and $\frac{5}{12}$ for B **[B1]**

Second stage branches from R correctly labeled: $\frac{6}{11}$ for R and $\frac{5}{11}$ for B **[B1]**

Second stage branches from B correctly labeled: $\frac{7}{11}$ for R and $\frac{4}{11}$ for B **[B1]**

(b) Multiplies probabilities along the R, R path: $\frac{7}{12} \times \frac{6}{11}$ **[M1]**

$\frac{42}{132}$ or $\frac{7}{22}$ **[A1]**

(c) Identifies and sums both correct combinations: $\left(\frac{7}{12} \times \frac{5}{11}\right) + \left(\frac{5}{12} \times \frac{7}{11}\right)$ or $2 \times \left(\frac{35}{132}\right)$ **[M1]**

$\frac{70}{132}$ or $\frac{35}{66}$ **[A1]**

[Back to Question] *[Total Marks: 7]*

1190 **(a)** $\frac{3}{8}$ placed correctly on the lower first branch **[B1]**

$\frac{5}{8}, \frac{3}{8}$ and $\frac{5}{8}, \frac{3}{8}$ placed correctly on the second branches **[B1]**

(b) $\frac{5}{8} \times \frac{5}{8}$ **[M1]**

$\frac{25}{64}$ **[A1]**

(c) $\frac{5}{8} \times \frac{3}{8}$ or $\frac{15}{64}$ **[M1]**

$2 \times \left(\frac{5}{8} \times \frac{3}{8}\right)$ **[M1]**

$\frac{15}{32}$ **[A1]**

[Back to Question] *[Total Marks: 7]*

1191 **(a)** $\frac{x}{x+4} \times \frac{x-1}{x+3}$ **[M1]**

$= \frac{x(x-1)}{(x+4)(x+3)} = \frac{x(x-1)}{x^2+7x+12}$ **[A1]**

(b) **(i)** $\frac{x(x-1)}{x^2+7x+12} = \frac{1}{3}$ **[M1]**

$3(x^2-x) = x^2+7x+12$ or $3x^2-3x = x^2+7x+12$ **[M1]**

$2x^2-10x-12 = 0 \implies x^2-5x-6 = 0$ **[A1]**

(ii) $(x-6)(x+1) = 0$ **[M1]**

$x = 6$ or $x = -1$ **[A1]**

Rejects $x = -1$ to conclude there are 6 blue counters **[B1]**

[Back to Question] *[Total Marks: 8]*

1192 **(a)** $\frac{5}{x+5} \times \frac{4}{x+4} = \frac{2}{15}$ **[M1]**

$20 \times 15 = 2(x+5)(x+4)$ or $300 = 2(x^2+9x+20)$ **[M1]**

$150 = x^2+9x+20$

$x^2+9x-130 = 0$ (No errors seen) [A1]

(b) $(x+13)(x-10) = 0$ **[M1]**

$x = 10$ **[A1]**

(c) Total counters $= 15$ **[B1]**

$\left(\frac{10}{15} \times \frac{5}{14}\right) + \left(\frac{5}{15} \times \frac{10}{14}\right)$ **[M1]**

$\frac{100}{210}$ or $\frac{10}{21}$ **[A1]**

[Back to Question] *[Total Marks: 8]*

1193 **(a)** Sets up correct equation for probability of two red balls: $\frac{x}{x+4} \times \frac{x-1}{x+3} = \frac{1}{3}$ **[M1]**

Cross-multiplies correctly to clear fractions: $3x(x-1) = (x+4)(x+3)$ **[M1]**

Expands and rearranges correctly with no errors:
$3x^2 - 3x = x^2 + 7x + 12 \implies 2x^2 - 10x - 12 = 0 \implies x^2 - 5x - 6 = 0$ **[A1]**

(b) Factorises or solves the quadratic equation: $(x-6)(x+1) = 0$ **[M1]**

6 (and explicitly or implicitly rejects $x = -1$) **[A1]**

(c) Uses $x = 6$ to determine probabilities for one red and one blue ball: $\frac{6}{10} \times \frac{4}{9}$ **[M1]**

Accounts for both (R, B) and (B, R) combinations: $2 \times \left(\frac{6}{10} \times \frac{4}{9}\right)$ or $\frac{24}{90} + \frac{24}{90}$ **[M1]**

$\frac{48}{90}$ or $\frac{8}{15}$ **[A1]**

[Back to Question] *[Total Marks: 8]*

1194 **(a)** $\frac{3}{n} \times \frac{2}{n-1}$ **[M1]**

$\frac{6}{n(n-1)} = \frac{1}{15}$ **[M1]**

$n^2 - n = 90$

n$^2 - n - 90 = 0$ with no errors seen [A1]

(b) $(n-10)(n+9) = 0$ leading to $n = 10$ **[M1]**

7 **[A1]**

(c) $\frac{3}{10} \times \frac{7}{9}$ or $\frac{7}{10} \times \frac{3}{9}$ **[M1]**

$2 \times \left(\frac{3}{10} \times \frac{7}{9}\right)$ **[M1]**

$\frac{7}{15}$ or $\frac{42}{90}$ **[A1]**

[Back to Question] *[Total Marks: 8]*

1195 **(a)** $\frac{3}{n} \times \frac{2}{n-1}$ **[M1]**

$\frac{6}{n(n-1)} = \frac{1}{15}$ **[M1]**

$90 = n^2 - n$

n$^2 - n - 90 = 0$ [A1]

(b) $(n-10)(n+9) = 0$ **[M1]**

10 **[A1]**

(c) $n = 10$, giving 3 Red and 7 Blue balls **[B1]**

$2 \times \left(\frac{3}{10} \times \frac{7}{9}\right)$ **[M1]**

$\frac{42}{90}$ or $\frac{7}{15}$ **[A1]**

[Back to Question] *[Total Marks: 8]*

1196 **(a)** $\frac{3}{n} \times \frac{2}{n-1} = \frac{1}{15}$ **[M1]**

$\frac{6}{n(n-1)} = \frac{1}{15}$ **[M1]**

$n^2 - n = 90 \implies n^2 - n - 90 = 0$ **[A1]**

(b) $(n-10)(n+9) = 0$ **[M1]**

$n = 10$ **[A1]**

(c) Number of Blue counters $= 7$ **[B1]**

$\left(\frac{3}{10} \times \frac{7}{9}\right) + \left(\frac{7}{10} \times \frac{3}{9}\right)$ **[M1]**

$\frac{42}{90}$ or $\frac{7}{15}$ **[A1]**

[Back to Question] *[Total Marks: 8]*

1197 **(a)** $\frac{2}{5}$ correctly placed on all three Tails branches **[B1]**

$\frac{3}{5}$ correctly placed on the remaining Heads branch **[B1]**

(b) $\frac{3}{5} \times \frac{2}{5} + \frac{2}{5} \times \frac{3}{5}$ or $2 \times \left(\frac{3}{5} \times \frac{2}{5}\right)$ **[M1]**

$\frac{12}{25}$ or 0.48 **[A1]**

[Back to Question] *[Total Marks: 4]*

1198 **(a)** 9 in the intersection of H and G **[B1]**

14, 10, and 7 in the correct regions **[B1]**

(b) $\frac{14}{40}$ or $\frac{7}{20}$ **[B1]**

(c) $\frac{9}{23} \times \frac{8}{22}$ **[M1]**

$\frac{72}{506}$ or $\frac{36}{253}$ **[A1]**

[Back to Question] *[Total Marks: 5]*

1199 **(a)** 6 placed correctly in the intersection **[B1]**

12 in F only, 8 in T only, and 4 outside the circles **[B1]**

(b) $\frac{6}{30}$ or $\frac{1}{5}$ or 0.2 **[B1]**

(c) $\frac{12}{30}$ seen as the probability for the first student **[M1]**

$\frac{12}{30} \times \frac{11}{29}$ **[M1]**

$\frac{132}{870}$ or $\frac{22}{145}$ **[A1]**

[Back to Question] *[Total Marks: 6]*

1200 **(a)** $2x + x + (x + 4) + 8 = 40$ **[M1]**

$4x + 12 = 40 \implies x = 7$ **[A1]**

(b) $\frac{32}{40}$ or $\frac{4}{5}$ **[B1]**

(c) Number of students playing Football only = 14 **[B1]**

$\frac{14}{40} \times \frac{13}{39}$ **[M1]**

$\frac{182}{1560}$ or $\frac{7}{60}$ **[A1]**

[Back to Question] *[Total Marks: 6]*

1201 **(a)** 10 in the intersection **[B1]**

14 in F only, 9 in T only, and 7 outside the circles **[B1]**

(b) $\frac{10}{40}$ or $\frac{1}{4}$ **[B1]**

(c) $\frac{10}{24} \times \frac{14}{23}$ or $\frac{14}{24} \times \frac{10}{23}$ **[M1]**

$\left(\frac{10}{24} \times \frac{14}{23}\right) + \left(\frac{14}{24} \times \frac{10}{23}\right)$ **[M1]**

$\frac{35}{69}$ or $\frac{280}{552}$ **[A1]**

[Back to Question] *[Total Marks: 6]*

1202 **(a)** Places 6 outside both circles **[B1]**

Calculates intersection: $32 + 21 + 6 - 50 = 9$ and places 9 in $H \cap G$ **[B1]**

Places 23 in H only and 12 in G only **[B1]**

(b) **(i)** $\frac{9}{50}$ **[B1]**

(ii) $\frac{38}{50}$ or $\frac{19}{25}$ **[B1]**

(c) Recognises the reduced sample space and uses probability without replacement: $\frac{9}{32} \times \frac{8}{31}$ **[M1]**

$\frac{72}{992}$ or $\frac{9}{124}$ **[A1]**

[Back to Question] *[Total Marks: 7]*

1203 **(a)** Intersection $n(G \cap H) = 8$ correctly placed **[B1]**

Remaining regions: $n(G \cap H') = 16$, $n(H \cap G') = 10$, outside region $= 6$ **[B1]**

(b) **(i)** $\frac{10}{40}$ (or $\frac{1}{4}$ or 0.25) **[B1]**

(ii) Identifies regions $16 + 8 + 6 = 30$ or calculates $1 - \frac{10}{40}$ **[M1]**

$\frac{30}{40}$ (or $\frac{3}{4}$ or 0.75) **[A1]**

(c) $\frac{24}{40} \times \frac{23}{39}$ **[M1]**

$= \frac{552}{1560}$ (or $\frac{23}{65}$ or 0.354) **[A1]**

[Back to Question] *[Total Marks: 7]*

1204 **(a)** 7 correctly placed in the intersection of T and G **[B1]**

17, 11, and 5 correctly placed in the remaining regions **[B1]**

(b) Identifies 17 students play tennis but not golf **[B1]**

$\frac{17}{40} \times \frac{16}{39}$ **[M1]**

$\frac{34}{195}$ or 0.174 (0.1743...) **[A1]**

(c) Denominator 18 seen **[M1]**

$\frac{7}{18}$ or 0.389 (0.3888...) **[A1]**

[Back to Question] *[Total Marks: 7]*

1205 **(a)** First branch $B = \frac{3}{10}$ **[B1]**

Top branch after R: $B = \frac{3}{9}$ **[B1]**

Bottom branches after B: $R = \frac{7}{9}$ and $B = \frac{2}{9}$ **[B1]**

(b) $\frac{7}{10} \times \frac{6}{9}$ **[M1]**

$\frac{42}{90}$ or $\frac{7}{15}$ or 0.467 **[A1]**

(c) $\frac{7}{10} \times \frac{3}{9} + \frac{3}{10} \times \frac{7}{9}$ **[M1]**

$\frac{42}{90}$ or $\frac{7}{15}$ or 0.467 **[A1]**

[Back to Question] *[Total Marks: 7]*

1206 **(a)** First pick: $\mathrm{P}(R) = \frac{5}{9}$ and $\mathrm{P}(Y) = \frac{4}{9}$ **[B1]**

Second pick: After R, $\mathrm{P}(R) = \frac{4}{8}$, $\mathrm{P}(Y) = \frac{4}{8}$; After Y, $\mathrm{P}(R) = \frac{5}{8}$, $\mathrm{P}(Y) = \frac{3}{8}$ **[B1]**

(b) $\mathrm{P}(RR) + \mathrm{P}(YY) = \left(\frac{5}{9} \times \frac{4}{8}\right) + \left(\frac{4}{9} \times \frac{3}{8}\right)$ **[M1]**

$\frac{20}{72} + \frac{12}{72} = \frac{32}{72} = \frac{4}{9}$ **[A1]**

(c) Method 1: $1 - \mathrm{P}(RR)$ or Method 2: $\mathrm{P}(RY) + \mathrm{P}(YR) + \mathrm{P}(YY)$ **[M1]**

$1 - \left(\frac{5}{9} \times \frac{4}{8}\right) = 1 - \frac{20}{72}$ **[M1]**

$\frac{52}{72} = \frac{13}{18}$ **[A1]**

[Back to Question] *[Total Marks: 7]*

1207 **(a)** First branches correct: $\mathrm{P}(R_1) = \frac{5}{8}$ and $\mathrm{P}(B_1) = \frac{3}{8}$ **[B1]**

Second branches after R_1 correct: $\mathrm{P}(R_2) = \frac{4}{7}$ and $\mathrm{P}(B_2) = \frac{3}{7}$ **[B1]**

Second branches after B_1 correct: $\mathrm{P}(R_2) = \frac{5}{7}$ and $\mathrm{P}(B_2) = \frac{2}{7}$ **[B1]**

(b) $\mathrm{P}(RR) + \mathrm{P}(BB) = \left(\frac{5}{8} \times \frac{4}{7}\right) + \left(\frac{3}{8} \times \frac{2}{7}\right)$ **[M1]**

$\frac{26}{56}$ or $\frac{13}{28}$ (or 0.464 or 0.4642...) **[A1]**

(c) $1 - \mathrm{P}(BB) = 1 - \left(\frac{3}{8} \times \frac{2}{7}\right)$ or $\left(\frac{5}{8} \times \frac{4}{7}\right) + \left(\frac{5}{8} \times \frac{3}{7}\right) + \left(\frac{3}{8} \times \frac{5}{7}\right)$ **[M1]**

$\frac{50}{56}$ or $\frac{25}{28}$ (or 0.893 or 0.8928...) **[A1]**

[Back to Question] *[Total Marks: 7]*

1208 **(a)** 3 placed correctly outside the circles **[B1]**

x, $24 - x$, and $18 - x$ placed correctly in the regions **[B1]**

(b) $(24 - x) + x + (18 - x) + 3 = 40$ or equivalent **[M1]**

5 **[A1]**

(c) $\frac{19}{40} \times \frac{13}{39}$ **[M1]**

$2 \times \left(\frac{19}{40} \times \frac{13}{39}\right)$ **[M1]**

$\frac{19}{60}$ or $\frac{494}{1560}$ or 0.316... **[A1]**

[Back to Question] *[Total Marks: 7]*

1209 **(a)** $18 + 14 + 3 - x = 30$ or $35 - x = 30$ **[M1]**

5 **[A1]**

(b) 5 placed correctly in the intersection **[B1]**

13 in F only, 9 in S only, and 3 outside the circles **[B1]**

(c) Identifies number of students studying only French is 13 **[M1]**

$\frac{13}{30} \times \frac{12}{29}$ **[M1]**

$\frac{156}{870}$ or $\frac{26}{145}$ **[A1]**

[Back to Question] *[Total Marks: 7]*

1210 **(a)** $\frac{3}{9}$ on the Red

Bluebranch[B1]

$\frac{7}{9}$ on the Blue

Redbranch[B1]

$\frac{2}{9}$ on the Blue

Bluebranch[B1]

(b) $\frac{7}{10} \times \frac{6}{9}$ **[M1]**

$\frac{42}{90}$ or $\frac{7}{15}$ **[A1]**

(c) $\left(\frac{7}{10} \times \frac{3}{9}\right) + \left(\frac{3}{10} \times \frac{7}{9}\right)$ **[M1]**

$\frac{42}{90}$ or $\frac{7}{15}$ **[A1]**

[Back to Question] *[Total Marks: 7]*

1211 **(a)** $\frac{5}{n} \times \frac{4}{n-1} = \frac{2}{21}$ [M2]

(Award M1 for $\frac{5}{n} \times \frac{4}{n}$ or $\frac{5}{n} + \frac{4}{n-1}$)

$2(n^2 - n) = 420$ or $n(n-1) = 210$ **[M1]**

$n^2 - n - 210 = 0$ reached with no errors seen **[A1]**

(b) $(n-15)(n+14) = 0$ or correct substitution into quadratic formula **[M1]**

$n = 15$ **[A1]**

(c) $\frac{5}{15} \times \frac{10}{14}$ or $\frac{10}{15} \times \frac{5}{14}$ seen **[M1]**

Multiplies their probability by 2 (i.e. $2 \times \frac{50}{210}$) **[M1]**

$\frac{10}{21}$ or 0.476 (0.4761...) **[A1]**

[Back to Question] *[Total Marks: 8]*

1212 **(a)** $\text{P(Blue from A)} = \frac{x}{x+3}$ and $\text{P(Red from A)} = \frac{3}{x+3}$ **[B1]**

If Blue transferred to B, Box B has 5 Blue and 2 Red (total 7), so $\text{P(Blue from B)} = \frac{5}{7}$ **[B1]**

If Red transferred to B, Box B has 4 Blue and 3 Red (total 7), so $\text{P(Blue from B)} = \frac{4}{7}$ **[B1]**

$\text{Total probability} = \left(\frac{x}{x+3} \times \frac{5}{7}\right) + \left(\frac{3}{x+3} \times \frac{4}{7}\right) = \frac{5x+12}{7(x+3)}$ **[A1]**

(b) **(i)** $\frac{5x+12}{7(x+3)} = \frac{3}{4}$ **[M1]**

$4(5x+12) = 21(x+3) \implies 20x + 48 = 21x + 63$ **[M1]**

$x = 15$ **[A1]**

(ii) $\frac{\text{P(Red from A and Blue from B)}}{\text{P(Blue from B)}} = \frac{\frac{3}{18} \times \frac{4}{7}}{\frac{3}{4}} = \frac{\frac{1}{14}}{\frac{3}{4}} = \frac{2}{21}$ **[B1]**

[Back to Question] *[Total Marks: 8]*

1213 **(a)** $\frac{4}{x} \times \frac{3}{x-1}$ **[M1]**

$\frac{12}{x(x-1)} = \frac{1}{11}$ or $12 \times 11 = x(x-1)$ **[M1]**

$132 = x^2 - x \implies x^2 - x - 132 = 0$ **[A1]**

(b) $(x-12)(x+11) = 0$ or correct substitution into the quadratic formula **[M1]**

$x = 12$ **[A1]**

Number of blue discs $= 12 - 4 = 8$ **[A1]**

(c) $\frac{4}{12} \times \frac{8}{11}$ or $\frac{32}{132}$ **[M1]**

$\left(\frac{4}{12} \times \frac{8}{11}\right) \times 2$ **[M1]**

$\frac{64}{132}$ or $\frac{16}{33}$ **[A1]**

[Back to Question] *[Total Marks: 9]*

1214 **(a)** $\frac{4}{n} \times \frac{3}{n-1} = \frac{1}{14}$ **[M1]**

$\frac{12}{n^2-n} = \frac{1}{14}$ **[M1]**

$168 = n^2 - n$ leading to $n^2 - n - 168 = 0$ with no errors seen **[A1]**

(b) $(n-14)(n+12) = 0$ or correct use of quadratic formula **[M1]**

$n = 14$ (rejecting $n = -12$) **[A1]**

Number of dark chocolates $= 14 - 4 = 10$ **[A1]**

(c) $\frac{4}{14} \times \frac{10}{13}$ or $\frac{10}{14} \times \frac{4}{13}$ **[M1]**

$2 \times \left(\frac{4}{14} \times \frac{10}{13}\right)$ **[M1]**

$\frac{80}{182}$ or $\frac{40}{91}$ **[A1]**

[Back to Question] *[Total Marks: 9]*

1215 **(a)** **(i)** Sets up equation $\frac{6}{n} \times \frac{5}{n-1} = \frac{1}{3}$ **[M1]**

Simplifies to $\frac{30}{n(n-1)} = \frac{1}{3}$ or $30 \times 3 = n(n-1)$ **[M1]**

$n^2 - n = 90$

n$^2 - n - 90 = 0$ with no steps omitted [A1]

(ii) Factorises $(n-10)(n+9) = 0$ or uses quadratic formula correctly **[M1]**

Obtains $n = 10$ and $n = -9$ **[A1]**

Rejects negative root and states total number of sweets $= 10$ **[B1]**

(b) Calculates $P(\text{all 3 lemon}) = \frac{6}{10} \times \frac{5}{9} \times \frac{4}{8}$ **[M1]**

Identifies $P(\text{at least 1 orange}) = 1 - P(\text{all 3 lemon})$ **[M1]**

$= 1 - \frac{120}{720} = 1 - \frac{1}{6} = \frac{5}{6}$ (or 0.833) **[A1]**

[Back to Question] *[Total Marks: 9]*

1216 **(a)** Sets up correct initial probability equation: $\frac{4}{n} \times \frac{3}{n-1} = \frac{2}{15}$ **[M1]**

Simplifies equation: $\frac{12}{n(n-1)} = \frac{2}{15} \implies 2n(n-1) = 180$ **[M1]**

Expands and rearranges correctly to establish $n^2 - n - 90 = 0$ with no missing algebraic steps **[A1]**

(b) Factorises $(n-10)(n+9) = 0$ or uses quadratic formula correctly **[M1]**

Rejects negative root $n = -9$ and states $n = 10$ **[A1]**

(c) Identifies total number of chocolates $n = 10$, dark $= 4$, milk $= 6$ **[B1]**

Calculates probability of one correct arrangement (e.g., DD M): $\frac{4}{10} \times \frac{3}{9} \times \frac{6}{8} = \frac{72}{720}$ or $\frac{1}{10}$ **[M1]**

Multiplies single selection probability by 3 (accounting for DD M, D MD, M DD) **[M1]**

$\frac{216}{720}$ or $\frac{3}{10}$ (or 0.3) **[A1]**

[Back to Question] *[Total Marks: 9]*

1217 **(a)** $\frac{x}{x+4} \times \frac{x-1}{x+3} = \frac{5}{18}$ **[M1]**

$18x(x-1) = 5(x+4)(x+3)$ **[M1]**

$18x^2 - 18x = 5(x^2 + 7x + 12)$ **[M1]**

$13x^2 - 53x - 60 = 0$ obtained with no errors **[A1]**

(b) $(13x + 12)(x - 5) = 0$ or correct use of the quadratic formula [M2] (allow [M1] for one error in formula)

$x = 5$ or $x = -\frac{12}{13}$ **[A1]**

Selects $x = 5$ as the only valid solution for the number of apples **[B1]**

(c) Total apples = 9, green = 5, red = 4. Evaluates one branch: $\frac{5}{9} \times \frac{4}{8}$ or $\frac{4}{9} \times \frac{5}{8}$ **[M1]**

$\left(\frac{5}{9} \times \frac{4}{8}\right) + \left(\frac{4}{9} \times \frac{5}{8}\right)$ or multiplies one branch by 2 **[M1]**

$\frac{40}{72}$ or $\frac{5}{9}$ **[A1]**

[Back to Question] *[Total Marks: 10]*

1218 **(a)** Boys Total = 40 and Girls Football = 8 **[B1]**

Girls Basketball = 8 and Total Tennis = 27 **[B1]**

Boys Tennis = 13 **[B1]**

(b) $\frac{13}{70}$ **[B1]**

[Back to Question] *[Total Marks: 4]*

1219 **(a)** Tennis Total = 15, Girls Total = 22 **[B1]**

Boys Football = 12, Boys Swimming = 9 **[B1]**

Girls Football = 8 **[B1]**

(b) $\frac{12}{50}$ or $\frac{6}{25}$ **[B1]**

[Back to Question] *[Total Marks: 4]*

1220 **(a)** Boys Basketball = 8 **[B1]**

Girls Football = 6 AND Girls Total = 22 **[B1]**

Total Basketball = 18 AND Total Tennis = 11 **[B1]**

(b) Denominator of 11 seen or fraction $\frac{5}{n}$ where $n > 5$ **[M1]**

$\frac{5}{11}$ **[A1]**

[Back to Question] *[Total Marks: 5]*

1221 **(a)** 6 or more correct missing entries **[B3]**

4 or 5 correct missing entries **[B2]**

2 or 3 correct missing entries **[B1]**

(Correct Boys row: 24, 10, 11, 45)

(Correct Girls row: 14, 12, 9, 35)

(Correct Total row: 38, 22, 20, 80)

(b) $\frac{10}{80}$ or $\frac{1}{8}$ **[B1]**

(c) Denominator of 35 seen **[M1]**

$\frac{9}{35}$ **[A1]**

[Back to Question] *[Total Marks: 6]*

1222 **(a)** $x + (2x + 10) + (x + 5) + (3x - 11)$ **[M1]**

$7x + 4$ **[A1]**

(b) $7x + 4 = 130$ **[M1]**

$x = 18$ **[A1]**

(c) Correct Male row: 18, 46, 64 **[B1]**

Correct Female row: 23, 43, 66 **[B1]**

Correct Total row: 41, 89, 130 **[B1]**

(d) Denominator of 41 seen **[M1]**

$\frac{23}{41}$ **[A1]**

[Back to Question] *[Total Marks: 9]*

1223 **(a)** Total Women = 55, Total Tea = 70, Overall Total = 120 placed correctly **[B1]**

Men Coffee = 25 and Total Coffee = 50 or Men Total = 65 placed correctly **[B1]**

Men Tea = 40, Women Tea = 30, Women Coffee = 25 **[B1]**

(b) $\frac{30}{120} \times 100$ **[M1]**

25 **[A1]**

[Back to Question] *[Total Marks: 5]*

1224 **(a)** Total column correct: 75 (Men), 45 (Women), 120 (Total) **[B1]**

Bus column correct: 15 (Men), 15 (Women), 30 (Total) **[B1]**

Train column correct: 24 (Men), 14 (Women), 38 (Total) **[B1]**

Car column correct: 36 (Men), 16 (Women), 52 (Total) **[B1]**

(b) Denominator of 52 seen or fraction $\frac{36}{n}$ **[M1]**

$\frac{36}{52}$ or $\frac{9}{13}$ **[A1]**

[Back to Question] *[Total Marks: 6]*

1225 **(a)** Total Juniors = 240, Total Seniors = 160 **[B1]**

Total Bus = 180, Total Car = 110, Total Walk = 110 **[B1]**

Juniors: Bus = 100, Car = 80, Walk = 60 AND Seniors: Bus = 80, Car = 30, Walk = 50 **[B1]**

(b) $\frac{60}{400}$ or $\frac{3}{20}$ **[B1]**

(c) $\frac{80}{160} \times \frac{79}{159}$ **[M1]**

$\frac{79}{318}$ **[A1]**

[Back to Question] *[Total Marks: 6]*

1226 **(a)** $x + 2x + (x + 5) + (3x - 2) = 80$ **[M1]**

$7x + 3 = 80$ **[M1]**

$x = 11$ **[A1]**

(b) Boys Bus = 11, Boys Walk = 22, Boys Total = 33 **[B1]**

Girls Bus = 16, Girls Walk = 31, Girls Total = 47 and Column totals (Bus = 27, Walk = 53) correct **[B1]**

(c) Denominator of 53 seen **[M1]**

$\frac{31}{53}$ **[A1]**

[Back to Question] *[Total Marks: 7]*

1227 **(a)** Correct Managers row: 48, 12, 60 **[B1]**

Correct Office Workers row: 90, 60, 150 **[B1]**

Correct Technicians row: 78, 12, 90 **[B1]**

Correct Total row: 216, 84, 300 **[B1]**

(b) Denominator of 84 seen **[M1]**

$\frac{60}{84}$ or $\frac{5}{7}$ **[A1]**

(c) $\frac{12}{60} \times \frac{48}{59}$ **[M1]**

$\frac{12}{60} \times \frac{48}{59} + \frac{48}{60} \times \frac{12}{59}$ or $2 \times \left(\frac{12}{60} \times \frac{48}{59}\right)$ **[M1]**

$\frac{96}{295}$ or 0.325 (0.3254...) **[A1]**

[Back to Question] *[Total Marks: 9]*

1228 **(a)** $\frac{3}{4}x$ or $0.75x$ **[B1]**

(b) Forms correct equation: $\frac{3}{4}x + \left(\frac{1}{4}x + 30\right) = 110$ **[M1]**

Simplifies to $x + 30 = 110$ leading to $x = 80$ with no errors **[A1]**

(c) Male row correct: 20 (Golf), 60 (Tennis), 80 (Total) **[B1]**

Female row correct: 70 (Golf), 50 (Tennis), 120 (Total) **[B1]**

Total row correct: 90 (Golf), 110 (Tennis), 200 (Total) **[B1]**

(d) Identifies correct fraction for first female: $\frac{70}{90}$ **[M1]**

Multiplies by correct fraction for second female: $\frac{70}{90} \times \frac{69}{89}$ **[M1]**

$\frac{4830}{8010}$ or $\frac{161}{267}$ **[A1]**

[Back to Question] *[Total Marks: 9]*

1229 **(a)** Total number of left-handed students $= 3x$ **[B1]**

$\frac{3x}{120} \times \frac{3x-1}{119} = \frac{3}{34}$ **[M1]**

$\frac{9x^2 - 3x}{14280} = \frac{3}{34}$ or $3x(3x-1) = 1260$ **[M1]**

$9x^2 - 3x = 1260 \implies 3x^2 - x - 420 = 0$ (with no errors seen) **[A1]**

(b) $(3x - 36)(x + 11.66...) = 0$ or correct substitution into quadratic formula **[M1]**

$x = 12$ **[A1]**

Total students equation: $x + y + 2x + y - 6 = 120$ **[M1]**

$3(12) + 2y - 6 = 120 \implies 2y = 90 \implies y = 45$ **[A1]**

(c) Total right-handed $= 2y - 6 = 2(45) - 6 = 84$ OR Right-handed girls $= 45 - 6 = 39$ **[M1]**

$\frac{39}{84}$ or $\frac{13}{28}$ **[A1]**

[Back to Question] *[Total Marks: 10]*

1230 **(a)** Class B, because the mean (or median) is higher. **[B1]**

(b) Class B, because the range (or interquartile range) is smaller. **[B1]**

(c) Identifies that the table only gives summary statistics / does not show individual marks **[B1]**

Mentions overlapping ranges / a specific example (e.g. the lowest mark in Class B could be lower than the highest mark in Class A) **[B1]**

[Back to Question] *[Total Marks: 4]*

1231 **(a)** Median $= 2$ **[B1]**

Range $= 4$ **[B1]**

(b) Team A generally scored more goals because their median is higher $(2 > 1.5)$ **[B1]**

Team A was more consistent because their range is smaller $(4 < 5)$ **[B1]**

[Back to Question] *[Total Marks: 4]*

1232 **(a)** B **[B1]**

(b) 3 **[B1]**

(c) Total students $= 5+10+3+2+7+8+3+2 = 40$ AND Total Grade A $= 5+7 = 12$ **[M1]**

30 **[A1]**

[Back to Question] *[Total Marks: 4]*

1233 **(a)** 27 **[B1]**

(b) 31 **[B1]**

(c) 36 and 20 identified as quartiles **[M1]**

16 **[A1]**

(d) $\frac{6}{15}$ or $\frac{2}{5}$ **[B1]**

[Back to Question] *[Total Marks: 5]*

1234 **(a)** Identifies the middle value is 7 **[M1]**

Identifies that 3 must appear at least twice as the lowest values: $3, 3$ **[M1]**

Uses the range to find the maximum value: $3 + 11 = 14$ **[M1]**

Uses the mean to find the missing value: $40 - (3 + 3 + 7 + 14) = 13$, yielding $3, 3, 7, 13, 14$ **[A1]**

(b) New sum $= 6 \times 9 = 54$ AND Old sum $= 5 \times 8 = 40$ (or $54 - 40$) **[M1]**

14 **[A1]**

(c) No, because standard deviation measures the spread of all data points around the mean, not just the difference between the extreme values **[B1]**

[Back to Question] *[Total Marks: 7]*

1235 **(a)** 40 **[B1]**

(b) 25 (from $50 - 25$) **[B1]**

(c) Group Y was slower on average / had a higher median time ($40 > 35$) **[B1]**

Group Y's times were more consistent / had a smaller IQR ($15 < 25$) or smaller range ($40 < 60$) **[B1]**

(d) Identifies that exactly 25% of Group Y took less than 30 minutes (since lower quartile is 30) **[M1]**

States Group X had a larger proportion, because 30 is between their LQ (25) and median (35), meaning more than 25% took less than 30 minutes **[A1]**

[Back to Question] *[Total Marks: 6]*

1236 **(a)** 45 **[B1]**

(b) 10 **[B1]**

(c) Group A students were faster on average because their median time (45 min) is less than Group B (50 min). **[B1]**

Group B times are more consistent / less spread out because their interquartile range (10 min) is smaller than Group A (20 min). **[B1]**

(d) Recognises that 55 is the upper quartile, representing 25% or $\frac{1}{4}$ of the students **[M1]**

20 **[A1]**

[Back to Question] *[Total Marks: 6]*

1237 **(a)** 65 **[B1]**

(b) $75 - 40$ **[M1]**

35 **[A1]**

(c) Class X has a higher average score (or median is higher: $65 > 55$) **[B1]**

Class X scores are less spread out / more consistent (or IQR is lower: $30 < 35$, or Range is lower: $60 < 65$) **[B1]**

(d) No, because the ranges overlap (e.g., the maximum score in Class Y is 90, which is higher than the lower quartile of Class X) **[B1]**

[Back to Question] *[Total Marks: 6]*

1238 **(a)** $120 < m \leq 140$ **[B1]**

(b) Use of correct midpoints: $110, 130, 150, 170, 190$ **[M1]**

$110 \times 10 + 130 \times 20 + 150 \times 15 + 170 \times 10 + 190 \times 5$ **[M1]**

$= 1100 + 2600 + 2250 + 1700 + 950 = 8600$ **[M1]**

$\frac{8600}{60} = 143.33\ldots$ so 143 (to 3 sf) **[A1]**

(c) The exact masses of the individual mangoes are unknown because the data is grouped **[B1]**

The 15 mangoes could have any combination of masses between 140 g and 160 g, not necessarily exactly at the midpoint **[B1]**

[Back to Question] *[Total Marks: 7]*

1239 **(a)** Upper Quartile = 44 and Lower Quartile = 26 **[M1]**

$44 - 26 = 18$ **[A1]**

(b) The Adults generally completed the puzzle faster because their median time is lower ($32 < 36$) **[B1]**

The Students' times were more consistent (less spread out) because their interquartile range (12) is smaller than the Adults' interquartile range (18) **[B1]**

(c) The reasoning is invalid because range only measures spread, not the actual values or maximum times **[B1]**

However, the conclusion happens to be true because the maximum time for Students (48) is less than the maximum time for Adults (55) **[B1]**

[Back to Question] *[Total Marks: 6]*

1240 **(a)** Forms a correct equation: $x + 2x + 120 + x = 360$ **[M1]**

Solves to find $4x = 240$

x = 60[$M1$]

Calculates number of students: $\frac{60}{360} \times 120 = 20$ **[A1]**

(b) Calculates the percentage for Class P: $\frac{120^\circ}{360^\circ} \times 100 = 33.3\%$ (or $\frac{1}{3}$) **[M1]**

Identifies total ratio parts for Class Q $= 1 + 3 + 2 + 2 = 8$ **[M1]**

Calculates percentage for Class Q: $\frac{3}{8} \times 100 = 37.5\%$ and concludes $37.5\% > 33.3\%$ **[A1]**

(c) The two classes have different total numbers of students (so proportions/percentages must be used for a fair comparison). **[B1]**

[Back to Question] *[Total Marks: 7]*

1241 **(a)** 112 **[B1]**

(b) 120 and 104 seen as quartiles **[M1]**

16 **[A1]**

(c) Cumulative frequency of 110 seen, leading to $120 - 110 = 10$ apples **[M1]**

8.33 (8.333...) **[A1]**

(d) Farm X apples are heavier on average as their median (112 g) is greater than Farm Y (105 g). **[B1]**

Farm X apples are more consistent in mass / less spread out as their interquartile range

(16 g) is smaller than Farm Y (22 g). **[B1]**

[Back to Question] *[Total Marks: 7]*

1242 **(a)** Continuous **[B1]**

(b) Discrete **[B1]**

(c) Continuous **[B1]**

[Back to Question] *[Total Marks: 3]*

1243 **(a)** **(i)** Discrete **[B1]**

(ii) Continuous **[B1]**

(iii) Continuous **[B1]**

(iv) Discrete **[B1]**

[Back to Question] *[Total Marks: 4]*

1244 **(a)** Discrete **[B1]**

(b) Continuous **[B1]**

(c) Continuous **[B1]**

(d) Continuous **[B1]**

[Back to Question] *[Total Marks: 4]*

1245 **(a)** Discrete **[B1]**

(b) Continuous **[B1]**

(c) Continuous **[B1]**

(d) Discrete **[B1]**

[Back to Question] *[Total Marks: 4]*

1246 **(a)** Data Set A (time) is continuous, so it can take any decimal value and inequalities are needed to define exact boundary limits without gaps. **[B1]**

Data Set B (number of push-ups) is discrete, so it only takes integer values and a hyphen clearly defines the inclusive start and end integers. **[B1]**

(b) **(i)** 19.5 and 29.5 **[B1]**

(ii) 10 **[B1]**

(c) Age (usually given as an integer in years) OR Money/Currency (rounded to nearest cent/pence) OR Shoe size **[B1]**

[Back to Question] *[Total Marks: 5]*

1247 **(a)** Continuous **[B1]**

(b) Height is continuous / can take decimal values. **[B1]**

A plant with a height between the intervals (e.g. 19.5 cm) would not fit into any class. **[B1]**

(c) Discrete **[B1]**

[Back to Question] *[Total Marks: 4]*

1248 **(a)** Discrete data can only take specific or exact values (often counted) **[B1]**

Continuous data can take any value within a given range (often measured) **[B1]**

(b) **(i)** Continuous **[B1]**

(ii) Rounding limits the precision of the recording, but the actual true length could still be any value between the rounded boundaries **[B1]**

[Back to Question] *[Total Marks: 4]*

1249 **(a)** Continuous **[B1]**

(b) Uses midpoints: 90, 110, 130, 150 **[M1]**

$\Sigma fx = 90 \times 12 + 110 \times 26 + 130 \times 15 + 150 \times 7 = 6940$ **[M1]**

$\frac{6940}{60} = 115.66... = 116$ or 115.7 **[A1]**

(c) The exact masses of the apples in each class are not known (or similar reference to data being grouped) **[B1]**

[Back to Question] *[Total Marks: 5]*

1250 **(a)** Length **[B1]**

(b) **(i)** 14.45 **[B1]**

(ii) 14.55 **[B1]**

(c) Identifies that the discrete values include $5, 6, 7, 8, 9$ OR uses continuous boundaries $9.5 - 4.5$ **[M1]**

5 **[A1]**

[Back to Question] *[Total Marks: 5]*

1251 **(a)** **(i)** Discrete **[B1]**

(ii) Continuous **[B1]**

(b) The variable can still only take specific, fixed intervals/values (e.g., jumping by 0.5) **[B1]**

It cannot take any value within a range (e.g., a shoe size of 8.23 does not exist) **[B1]**

[Back to Question] *[Total Marks: 4]*

1252 **(a)** Continuous data can take any value within a given range (measured). **[B1]**

Discrete data can only take specific, exact values (usually counted). **[B1]**

(b) **(i)** Continuous **[B1]**

(ii) Shoe size can only take specific, set values (there are gaps between sizes like 7 and 7.5). **[B1]**

It cannot take every possible value (e.g. you cannot have a shoe size of 7.23), so it must be discrete. **[B1]**

[Back to Question] *[Total Marks: 5]*

1253 **(a)** **(i)** Discrete **[B1]**

(ii) Continuous **[B1]**

(b) Upper bound of mass of one book = 855 g **[M1]**

$24 \times 855 = 20520$ g $\implies$ 20.52 **[A1]**

(c) Lower bound of empty box = 1.15 kg **[B1]**

Lower bound of mass of one book = 845 g = 0.845 kg seen **[B1]**

Total lower bound $= 1.15 + (24 \times 0.845) = 21.43$ **[A1]**

[Back to Question] *[Total Marks: 7]*

1254 **(a)** 1 **[B1]**

(b) Ordered list: $0, 0, 1, 1, 1, 1, 2, 2, 3, 4$ **[M1]**

1 **[A1]**

(c) $\dfrac{2+0+1+4+1+0+3+1+2+1}{10} = \dfrac{15}{10}$ **[M1]**

1.5 **[A1]**

[Back to Question] *[Total Marks: 5]*

1255 **(a)** 16 **[B1]**

(b) 21 **[B1]**

(c) Ordered list seen: $5, 8, 10, 13, 14, 16, 18, 20, 22, 24, 26$ **[M1]**

12 **[A1]**

(d) $(14 + 20 + 8 + 24 + 16 + 5 + 26 + 13 + 22 + 10 + 18) \div 11$ or $176 \div 11$ **[M1]**

16 **[A1]**

[Back to Question] *[Total Marks: 6]*

1256 **(a)** 7 **[B1]**

(b) $12 - 7 = 5$ **[B1]**

(c) 8 **[B1]**

(d) $(7 + 7 + 7 + 8 + 10 + 12 + 12) \div 7$ or $\frac{63}{7}$ **[M1]**

9 **[A1]**

(e) $\frac{63 + y}{8} = 9.5$ or 9.5×8 **[M1]**

$76 - 63 = 13$ **[A1]**

[Back to Question] *[Total Marks: 7]*

1257 **(a)** $30 < h \leq 40$ **[B1]**

(b) Uses midpoints $15, 25, 35, 45, 55$ **[M1]**

$\Sigma fx = (15 \times 12) + (25 \times 25) + (35 \times 28) + (45 \times 11) + (55 \times 4)$ or $180 + 625 + 980 + 495 + 220 = 2500$ **[M1]**

$\frac{2500}{80}$ **[M1]**

31.25 **[A1]**

[Back to Question] *[Total Marks: 5]*

1258 **(a)** $100 < m \leq 110$ **[B1]**

(b) Midpoints used: $85, 95, 105, 115, 125$ **[M1]**

$\Sigma fx = 85 \times 8 + 95 \times 22 + 105 \times 31 + 115 \times 14 + 125 \times 5$ **[M1]**

$\Sigma fx = 680 + 2090 + 3255 + 1610 + 625 = 8260$ **[M1]**

Mean $= \frac{8260}{80} = 103.25$ **[A1]**

(c) The exact masses of individual apples are not known **[B1]**

The calculation uses the midpoints of the class intervals, assuming the data is evenly distributed within each class **[B1]**

[Back to Question] *[Total Marks: 7]*

1259 **(a)** $30 < t \leq 40$ **[B1]**

(b) Midpoints: $15, 25, 35, 45, 55$ **[M1]**

$\Sigma fx = 15(14) + 25(28) + 35(32) + 45(18) + 55(8)$ **[M1]**

$= 210 + 700 + 1120 + 810 + 440 = 3280$ **[M1]**

$\frac{3280}{100} = 32.8$ **[A1]**

(c) $18 + 8 = 26$ **[M1]**

26 **[A1]**

(d) $40 < t \leq 50$ **[B1]**

[Back to Question] *[Total Marks: 8]*

1260 **(a)** $p + q = 25$ **[B1]**

$\frac{10p + 420 + 800 + 70q + 450}{60} = 48$ or $10p + 70q = 1210$ [B2] (Allow B1 for unsimplified $\Sigma fx = 10p + 420 + 800 + 70q + 450$ seen)

Correctly sets up and solves simultaneous equations to eliminate one variable **[M1]**

$p = 9$, $q = 16$ **[A1]**

(b) Identifies median is between 30th and 31st value and shows cumulative frequencies 23 and 39 (or $9 + 14 = 23$ and $23 + 16 = 39$) **[M1]**

$40 < m \leq 60$ **[A1]**

[Back to Question] *[Total Marks: 7]*

1261 **(a)** Since the mode is 8, there must be at least two 8s. **[M1]**

Since the median is 11, the middle number when ordered is 11. The first three numbers are $8, 8, 11$. **[M1]**

The mean is 12, so the sum of the five numbers is $5 \times 12 = 60$. **[M1]**

The range is 15, so the largest number is $8 + 15 = 23$. **[M1]**

Let the fourth number be x. Then $8 + 8 + 11 + x + 23 = 60 \implies 50 + x = 60 \implies x = 10$. **[A1]**

(b) New sum $= 6 \times 13 = 78$ **[M1]**

$k = 78 - 60 = 18$ **[A1]**

[Back to Question] *[Total Marks: 7]*

1262 **(a)** $15 + x + y + 8 + 2 = 50$ leading to $x + y = 25$ **[B1]**

(b) Midpoints: $1, 3, 5, 7, 9$ **[M1]**

$\frac{1(15) + 3(x) + 5(y) + 7(8) + 9(2)}{50} = 3.8$ **[M1]**

$15 + 3x + 5y + 56 + 18 = 190$ **[M1]**

$3x + 5y + 89 = 190$ leading to $3x + 5y = 101$ **[A1]**

(c) $3x + 3y = 75$ or substitution method $3(25 - y) + 5y = 101$ **[M1]**

$2y = 26 \implies y = 13$ **[A1]**

$x = 12$ **[A1]**

(d) $2 < m \leq 4$ **[B1]**

[Back to Question] *[Total Marks: 9]*

1263 **(a)** 17.5 (or $17\frac{7}{15}$ or 17.46...) **[B1]**

(b) 17 **[B1]**

(c) Upper quartile = 21 and Lower quartile = 15 **[M1]**

6 **[A1]**

[Back to Question] *[Total Marks: 4]*

1264 **(a)** **(i)** 3.0476... **[M1]**

3.05 **[A1]**

(ii) 3 **[B1]**

(iii) Upper quartile = 4 and Lower quartile = 2 identified **[M1]**

2 **[A1]**

[Back to Question] *[Total Marks: 5]*

1265 **(a)** 16 **[B2]**

If B0, M1 for $\sum x = 240$ seen

(b) 15 **[B1]**

(c) 6 **[B2]**

If B0, M1 for $Q_1 = 13$ and $Q_3 = 19$ seen

[Back to Question] *[Total Marks: 5]*

1266 **(a)** $20 < t \leq 30$ **[B1]**

(b) Use of correct midpoints: $5, 15, 25, 35, 50$ **[M1]**

$\Sigma fx = 14(5) + 28(15) + 35(25) + 18(35) + 5(50)$ or 2245 seen **[M1]**

22.45 **[A1]**

(c) The exact times are not known so the midpoints of the intervals are used. **[B1]**

[Back to Question] *[Total Marks: 5]*

1267 **(a)** Use of correct mid-interval values $(5, 15, 25, 35, 50)$ **[M1]**

$\Sigma ft = 5(12) + 15(25) + 25(28) + 35(11) + 50(4)$ or 1720 **[M1]**

21.5 **[A1]**

(b) The exact times are not known / the data is grouped so mid-interval values are used. **[B1]**

(c) The mean will decrease because 12 is less than the current mean (21.5). **[B1]**

[Back to Question] *[Total Marks: 5]*

1268 **(a)** 21.25 **[B2]**

If B0, M1 for $\sum fx = 1700$ or midpoints $5, 15, 25, 35, 45$ seen

(b) The exact times are not known / the midpoints of each class are used as an estimate. **[B1]**

(c) $120 \times 24.5 = 2940$ **[M1]**

$(1700 + 2940) \div 200$ **[M1]**

23.2 **[A1]**

[Back to Question] *[Total Marks: 6]*

1269 **(a)** $3 + x + y + 6 + 4 + 1 = 35$ leading to $x + y = 21$ **[M1]**

Expression for total goals: $0(3) + 1(x) + 2(y) + 3(6) + 4(4) + 5(1)$ or $x + 2y + 39$ **[M1]**

$\dfrac{x + 2y + 39}{35} = 2.2$ leading to $x + 2y = 38$ **[M1]**

$x = 4$ and $y = 17$ **[A1]**

(b) 2 **[B1]**

[Back to Question] *[Total Marks: 5]*

1270 **(a)** $\dfrac{0 \times 6 + 1 \times x + 2 \times 8 + 3 \times 4 + 4 \times 2}{6 + x + 8 + 4 + 2} = 1.5$ **[M1]**

$\dfrac{x + 36}{x + 20} = 1.5$ **[M1]**

$x + 36 = 1.5x + 30 \implies 0.5x = 6 \implies x = 12$ **[A1]**

(b) 1 **[B1]**

(c) 1 **[B2]**

If B0, M1 for $Q_1 = 1$ and $Q_3 = 2$ seen

[Back to Question] *[Total Marks: 6]*

1271 **(a)** Total frequency $= 10 + k + 2k + 3 + 7 + 5 = 3k + 25$ **[M1]**

Total score $= 10(1) + k(2) + (2k + 3)(3) + 7(4) + 5(5) = 8k + 72$ **[M1]**

$\dfrac{8k + 72}{3k + 25} = 2.8$ **[M1]**

$8k + 72 = 8.4k + 70 \implies 0.4k = 2 \implies k = 5$ **[A1]**

(b) **(i)** 3 **[B1]**

(ii) 1.5 **[B1]**

(iii) 4 **[B1]**

[Back to Question] *[Total Marks: 7]*

1272 **(a)** 21 **[B1]**

(b) 25 **[B1]**

(c) 33 **[B1]**

(d) 6 (parcels identified) **[M1]**

$\frac{6}{15}$ or $\frac{2}{5}$ **[A1]**

[Back to Question] *[Total Marks: 5]*

1273 **(a)** Correct stems used **[B1]**

Unordered leaves all correct **[B1]**

Fully correct ordered leaves:
Row 1: 2 5
Row 2: 3 7 7
Row 3: 1 4 8
Row 4: 0 2 2 5 9
Row 5: 1 3 **[B1]**

(b) 34 **[B1]**

(c) 41 **[B1]**

[Back to Question] *[Total Marks: 5]*

1274 **(a)** $360 - (110 + 80 + 70 + 40) = 60$ **[B1]**

(b) $\frac{80}{360} \times 180$ **[M1]**

40 **[A1]**

(c) $\frac{110}{360}$ **[M1]**

$\frac{11}{36}$ **[A1]**

[Back to Question] *[Total Marks: 5]*

1275 **(a)** Correct stems $1, 2, 3, 4$ **[B1]**

Correct ordered leaves for at least two stems **[B1]**

Fully correct ordered diagram **[B1]**

(b) 2.9 **[B1]**

(c) $\frac{4}{15}$ **[B1]**

[Back to Question] *[Total Marks: 5]*

1276 **(a)** $\frac{150}{360} \times 120$ **[M1]**

50 **[A1]**

(b) $\frac{25}{120} \times 360$ **[M1]**

75 **[A1]**

(c) Angle for Bicycle $= 360 - 150 - 90 - 75 = 45°$ (or finding 15 students) **[M1]**

$\frac{45}{360} \times 100$ or $\frac{15}{120} \times 100$ **[M1]**

12.5 **[A1]**

[Back to Question] *[Total Marks: 7]*

1277 **(a)** $\frac{35}{120} \times 360$ **[M1]**

105 **[A1]**

(b) Sector angle for Tennis $= \frac{25}{120} \times 360 = 75°$ **[B1]**

$\frac{75}{360} \times \pi \times 6^2$ **[M1]**

23.6 (accept 23.56 to 23.57) **[A1]**

(c) $\frac{54}{108} \times 360$ or $54 \div \frac{108}{360}$ **[M1]**

180 **[A1]**

[Back to Question] *[Total Marks: 7]*

1278 **(a)** $2x + 30 + x + x + 50 + 2x - 20 = 360$ **[M1]**

$6x + 60 = 360$ leading to $x = 50$ **[A1]**

(b) Angle for Tennis $= 100°$ **[M1]**

Total $= 10 \times \frac{360}{100} = 36$ **[A1]**

(c) Number of students choosing Football $= 13$ and Basketball $= 5$ **[B1]**

$\frac{13}{36} \times \frac{5}{35} \times 2$ **[M1]**

$\frac{13}{126}$ **[A1]**

[Back to Question] *[Total Marks: 7]*

1279 **(a)** $x + 2x + 15 + 3x - 10 + x + 40 = 360$ **[M1]**

$7x + 45 = 360 \implies 7x = 315$ **[M1]**

$x = 45$ **[A1]**

(b) $\frac{45}{360} \times$ Total $= 24$ or $24 \times \frac{360}{45}$ **[M1]**

192 **[A1]**

(c) Angle for Grade C $= 3(45) - 10 = 125°$ **[M1]**

Percentage $= \dfrac{125}{360} \times 100$ **[M1]**

34.7 (accept 34.72 or 34.722 . . .) **[A1]**

[Back to Question] *[Total Marks: 8]*

1280 **(a)** Total Laptops $= 50 + 60 + 70 = 180$ **[B1]**

Total Tablets (Jan + Feb) $= 40 + 60 = 100$ **[B1]**

$180 = 1.2 \times (100 + x)$ or $\dfrac{180}{1.2} = 100 + x$ **[M1]**

$x = 50$ **[A1]**

(b) Total items (Jan to Mar) $= 180 + 150 = 330$ **[B1]**

Total items (April) $= 330 \times \dfrac{1}{3} = 110$ **[M1]**

Laptops (April) $= \dfrac{6}{6+5} \times 110$ **[M1]**

60 **[A1]**

(c) Tablet revenue $= 40 \times 200 = 8000$ and Laptop revenue $= 50 \times 450 = 22500$ **[M1]**

$\dfrac{8000}{8000 + 22500} \times 100$ **[M1]**

26.2 (accept 26.22 to 26.23) **[A1]**

[Back to Question] *[Total Marks: 11]*

1281 **(a)** Positive **[B1]**

(b) Single straight ruled line passing within the bounds of the data points, with roughly equal numbers of points on either side. **[B1]**

(c) 64 (Accept values between 62 and 67 strictly based on the student's line of best fit) **[B1]**

[Back to Question] *[Total Marks: 3]*

1282 **(a)** Plotting the remaining 6 points correctly. **[P2]**

(Allow P1 for 4 or 5 points correct).

(b) Negative **[B1]**

(c) A ruled straight line of best fit drawn by eye, with a roughly equal number of points on each side. **[B1]**

[Back to Question] *[Total Marks: 4]*

1283 **(a)** All 8 points plotted correctly **[B2]**

(5, 6, or 7 points plotted correctly [B1])

(b) Positive **[B1]**

(c) Ruled line passing through the mean point $(45, 45)$ and fitting the trend of the data **[B1]**

(d) Value read correctly from their line of best fit (typically $42 - 48$) **[B1]**

[Back to Question] *[Total Marks: 5]*

1284 **(a)** $y = 5.96x + 34.7$ (or $m = 5.96$, $c = 34.7$) **[B2]**

(Allow B1 for either $m = 5.96$ or $c = 34.7$ correct to 3 sf)

(b) Substitution of $x = 7$ into their equation from (a): $5.96(7) + 34.7$ **[M1]**

76.4 (or 76) **[A1ft]**

(c) The value $x = 15$ is outside the range of the given data (extrapolation). **[B1]**

[Back to Question] *[Total Marks: 5]*

1285 **(a)** Points $(7, 5)$ and $(9, 1)$ correctly plotted **[B1]**

(b) Mean age $= 4.5$ **[B1]**

Mean price $= 11$ **[B1]**

(c) Mean point $(4.5, 11)$ plotted accurately **[B1]**

Ruled line of best fit passing through $(4.5, 11)$ **[B1]**

(d) Negative (or strong negative) **[B1]**

(e) Value read correctly from their line of best fit (typically $2 - 4$) **[B1]**

[Back to Question] *[Total Marks: 7]*

1286 **(a)** All three points $(6, 81)$, $(7, 88)$, and $(8, 100)$ plotted correctly **[B2]**

(Allow B1 if only 1 or 2 points are plotted correctly)

(b) $\bar{x} = \dfrac{35}{7} = 5$ **[B1]**

$\bar{y} = \dfrac{490}{7} = 70$ **[B1]**

(c) Mean point $(5, 70)$ correctly plotted (allow a distinct mark e.g. a circle around the cross) **[B1]**

Single straight ruled line passing exactly through their mean point, following the trend of the data **[B1]**

(d) 75 (Accept values between 73 and 77 strictly based on the student's line of best fit) **[B1]**

[Back to Question] *[Total Marks: 7]*

1287 **(a)** $\bar{x} = \dfrac{2100}{12} = 175$ **[B1]**

$\bar{y} = \dfrac{924}{12} = 77$ **[B1]**

(b) Substitution of $(\bar{x}, \bar{y})$ into the equation: $77 = 0.85(175) - c$ **[M1]**

$c = 148.75 - 77 = 71.75$ **[A1]**

(c) Predicted mass $= 0.85(180) - 71.75 = 81.25$ **[M1]**

New mean $= \dfrac{924 + 81.25}{13} = 77.3$ (or 77.32...) **[A1]**

[Back to Question] *[Total Marks: 6]*

1288 **(a)** **(i)** $m = -5.82$ (accept -5.81 to -5.82) **[B1]**

$c = 27.0$ (accept 27.0 to 27.1) **[B1]**

(ii) The decrease in fuel consumption (in km/l) for every 1 litre increase in engine size. **[B1]**

(b) Substitution of $x = 1.6$ into their equation: $-5.816(1.6) + 27.0$ **[M1]**

17.7 (accept 17.6 to 17.8 depending on rounding of m and c) **[A1]**

(c) Because 5.0 is outside the range of the given data (extrapolation). **[B1]**

[Back to Question] *[Total Marks: 6]*

1289 **(a)** Mean temperature $= 21.25$ **[B1]**

Mean ice creams sold $= 80.25$ **[B1]**

(b) $m = 4.28$ **[B1]**

$c = -10.6$ **[B1]**

(c) $y = 4.28 \times 24 - 10.6$ **[M1]**

92 (or 92.12) **[A1]**

(d) Unreliable because 42°C is outside the range of the given data (extrapolation) **[B1]**

[Back to Question] *[Total Marks: 7]*

1290 **(a)** 32 to 34 **[B1]**

(b) Reading Upper Quartile as 40 and Lower Quartile as 23 to 24 **[M1]**

16 to 17.5 **[A1]**

(c) $120 - 100$ (or 120 – their reading at $t = 45$) **[M1]**

18 to 22 **[A1]**

[Back to Question] *[Total Marks: 5]*

1291 **(a)** **(i)** 24 **[B1]**

(ii) Reading Lower Quartile = 15 and Upper Quartile = 32.5 (Accept 32 to 33) **[M1]**

17.5 (Accept 17 to 18) **[A1]**

(b) Reading Cumulative Frequency at 45 km = 56 (Accept 55 to 57) **[M1]**

$60 - 56$ **[M1]**

4 (Accept 3 to 5) **[A1]**

[Back to Question] *[Total Marks: 6]*

1292 **(a)** 6 points plotted correctly at the upper boundaries **[P2]**

(Allow P1 for 4 or 5 points plotted correctly)

A smooth curve drawn passing through all their points **[C1]**

(b) Reading from cumulative frequency 40 **[M1]**

$26 \leq$ answer ≤ 28 **[A1]**

[Back to Question] *[Total Marks: 5]*

1293 **(a)** $90, 160, 188, 200$ **[B2]**

(B1 for 2 or 3 correct values)

(b) 5 points plotted correctly from their table **[B2]**

(B1 for 3 or 4 points plotted correctly)

Smooth curve drawn through their points **[B1]**

(c) Reading from curve at cumulative frequency $= 140$ **[M1]**

46 to 48 **[A1]**

[Back to Question] *[Total Marks: 7]*

1294 **(a)** Reading from cumulative frequency 60 **[M1]**

$22 \leq$ answer ≤ 24 **[A1]**

(b) Attempt to read from cumulative frequencies 90 (UQ) and 30 (LQ) **[M1]**

Lower quartile ≈ 15.5 and Upper quartile ≈ 29.5 seen **[M1]**

$12 \leq$ answer ≤ 16 **[A1]**

(c) Attempt to read at $d = 35$ (approx 105) **[M1]**

120 – their reading **[M1]**

$13 \leq$ answer ≤ 17 **[A1]**

[Back to Question] *[Total Marks: 8]*

1295 **(a)** 46, 92, 112 **[B2]**

(Allow [B1] for 2 correct values)

(b) Points correctly positioned horizontally and vertically starting at $(50, 0)$ **[B1]**

All 5 points plotted correctly at class upper boundaries (Tolerance $\pm$ half small square) **[B1]**

Smooth curve drawn through all 6 points **[B1]**

(c) Finding target Cumulative Frequency = 60% of 120 = 72 **[M1]**

178 (Accept 176 to 180) **[A1]**

(d) Reading Cumulative Frequency at $m = 220$ is 100 **[M1]**

$120 - 100$ **[M1]**

20 (Accept 18 to 22) **[A1]**

[Back to Question] *[Total Marks: 10]*

1296 **(a)** $0.8 \times 200 = 160$ **[M1]**

$28 \leq$ answer ≤ 29 **[A1]**

(b) Frequencies 30, 60, 80, 30 correct **[B2]**

(Allow B1 for 2 or 3 correct frequencies)

(c) Using midpoints 5, 15, 25, 35 **[M1]**

$\Sigma fx = 30 \times 5 + 60 \times 15 + 80 \times 25 + 30 \times 35$ (using their frequencies) **[M1]**

their $4100 \div 200$ **[M1]**

20.5 **[A1]**

[Back to Question] *[Total Marks: 8]*

1297 **(a)** Reading from curve at $m = 220$: 81 to 85 **[M1]**

100 − their reading **[M1]**

15 to 19 **[A1]**

(b) $p = 10$ **[B1]**

$q = 26$ **[B1]**

$r = 18$ **[B1]**

(c) Use of correct midpoints: $75, 125, 175, 225, 275$ **[M1]**

$\sum f \times m = 10 \times 75 + 26 \times 125 + 40 \times 175 + 18 \times 225 + 6 \times 275 = 16700$ **[M1]**

$16700 \div 100 = 167$ **[A1]**

[Back to Question] *[Total Marks: 9]*

1298 **(a)** Reading Lower Quartile = 38 (Accept 37 to 39) and Upper Quartile = 64 (Accept 63 to 66) **[M1]**

26 (Accept 24 to 29) **[A1]**

(b) Finding Cumulative Frequency = 15% of 200 = 30 **[M1]**

30 (Accept 29 to 31) **[A1]**

(c) Finding target Cumulative Frequency = 200 − 30 = 170 **[M1]**

71 (Accept 70 to 72) **[A1]**

(d) Reading Cumulative Frequency at 45 marks = 74 (Accept 72 to 76) **[M1]**

Number of candidates NOT taking test = 200 − 74 = 126 **[M1]**

$\frac{126}{200} \times 100\%$ **[M1]**

63 (Accept 62 to 64) **[A1]**

[Back to Question] *[Total Marks: 10]*

1299 **(a)** 090
090 **[B1]**

(b) 054
+
180
054
+180

[M1]

234
234 **[A1]**

(c) Angle
=
90
54
=
36
BAC=90
54
=36
and angle
=
180
36
2
=
72
ABC=
2
180
36
=72

[M1]

Bearing
=
234
72
=
162
=234
72

=162 **[A1]**

[Back to Question] *[Total Marks: 5]*

1300 **(a)** 065
065 (accept
063
063 to
067
067) **[B1]**

(b) Measured length
=
6
PQ=6 cm (accept
5.8
5.8 cm to
6.2
6.2 cm) **[B1]**

30
30 (accept
29
29 to
31
31) **[A1]**

(c) Line drawn from
Q on bearing
140
140
(accept
138
138
to
142
142
) **[B1]**

Point
R marked at distance
5
5 cm from
Q (accept
4.8
4.8 cm to
5.2
5.2 cm) and labelled **[B1]**

(d) 065
+
180
065
+180

[M1]

245
245 **[A1]**

[Back to Question] *[Total Marks: 7]*

1301 **(a)** 270
270 **[B1]**

(b) 018.6
+
180
018.6
+180

[M1]

198.6
198.6 **[A1]**

(c) **(i)** 198.6
48.4
198.6
48.4

[M1]

150.2
150.2 **[A1]**

(ii) Angle
=
90
18.6
=
71.4
ADC=90
18.6
=71.4

[B1]

Angle
=
322.8
270
=
52.8
BCD=322.8
270
=52.8

[B1]

Angle
=
360
(
71.4
+
150.2

+
52.8
)
=
85.6
ABC=360
(71.4
+150.2
+52.8
)=85.6 **[A1]**

(d) Bearing of
A from
=
048.4
+
180
=
228.4
B=048.4
+180
=228.4

[M1]

Bearing of
C from
=
228.4
85.6
=
142.8
B=228.4
85.6
=142.8 **[A1]**

[Back to Question] *[Total Marks: 10]*