IGCSE 0607 Practice Questions — E3.1 — Combined Questions & Markscheme


Graphs of functions

Total Questions: 39 — **Total Marks:** 215 Marks

Table of Contents & Quick Navigation

- **E3.1** Graphs of functions
  → **Paper 2 (Non-Calculator)** *(20 Qs)* · **Paper 4 (Calculator)** *(19 Qs)*

1 The diagram shows the graph of $y = a\sin(bx)$ for $0° \leq x \leq 360°$.




(a) Find the value of a.

$a =$ ..... **[1]**

(b) Find the value of b.

$b =$ ..... **[1]**

2 The diagram shows a sketch of the graph of $y = a\cos(bx)$ for $0° \leq x \leq 360°$.




(a) Write down the value of a.

$a =$ ..... **[1]**

(b) Find the value of b.

$b =$ ..... **[2]**

3 The diagram shows a sketch of the graph of the function $f(x) = ax^2 + c$.
The graph passes through the points $(0, -5)$ and $(2, 7)$.




(a) Write down the value of c.

$c =$ ..... **[1]**

(b) Find the value of a.

$a =$ ..... **[2]**

4 The diagram shows the sketch of the graph of $y = a\cos(bx)$ for $0° \leq x \leq 120°$.




(a) Write down the amplitude of the graph.

..... **[1]**

(b) Find the value of a and the value of b.

$a =$ ..... , $b =$ ..... **[2]**

5 The diagram shows the graph of the trigonometric function $f(x) = a\cos(bx)$ for $0° \leq x \leq 360°$.




(a) Write down the value of a and the value of b.

$a =$ ..... , $b =$ ..... **[2]**

(b) Write down the coordinates of the first minimum point where $x > 0$.

(..... , ...) **[1]**

6 The diagram shows the graph of $y = a\sin(bx)$ for $0° \leq x \leq 360°$.




(a) Find the value of a and the value of b.

$a =$ ..... , $b =$ ..... **[2]**

(b) Write down the period of the function.

..... ° **[1]**

7 Four graphs are shown below, labelled A, B, C, and D.



(a) Write down the letter of the graph that represents each function.

(i)

$$y = x^2 - 1$$

..... **[1]**

(ii)

$$y = 2^x$$

..... **[1]**

(iii)

$$y = \frac{1}{x}$$


..... **[1]**

(iv)

$$y = 2x - 1$$

..... **[1]**

8 The diagram shows the graph of $y = a\cos(bx)$ for $0° \leq x \leq 360°$.



(a) Write down the amplitude of the graph.

..... **[1]**


(b) Write down the period of the graph.

..... ° **[1]**

(c) Find the value of a and the value of b.

$a =$ ..... , $b =$ ..... **[2]**

9 The diagram shows the graph of $y = a\sin(bx)$ for $0° \leq x \leq 180°$.



(a) State the amplitude.

..... **[1]**


(b) State the period of the function.

..... ° **[1]**

(c) Write down the value of a and the value of b.

$a =$ ..... , $b =$ ..... **[2]**

10 The diagram shows the graph of $y = a\sin(bx)$ for $0° \leq x \leq 360°$.




(a) Find the value of a and the value of b.

$a =$ ..... , $b =$ ..... **[2]**

(b) The graph of $y = kx^3$ passes through the point $(2, -24)$.
Find the value of k.

$k =$ ..... **[2]**

11 The diagram shows the graph of $y = a\sin(bx)$ for $0° \leq x \leq 360°$.
The values of a and b are positive integers.



(a) Find the value of a and the value of b.

$a =$ ..... , $b =$ ..... **[2]**


(b) Write down the period of the function.

..... ° **[1]**

(c) State the number of solutions to the equation $a\sin(bx) = 2$ in the interval $0° \leq x \leq 360°$.

..... **[1]**

12 The diagram shows the graph of the function $y = x^2 + bx + c$.



(a) Write down the value of c.

$c =$ ..... **[1]**

(b) Find the value of b.

$b =$ ..... **[2]**


(c) Write down the coordinates of the minimum point of the graph.

(..... , ...) **[2]**

13 Six equations are shown below.

$$y = 2^x \qquad y = x^2 \qquad y = -x^2 \qquad y = x^3 \qquad y = \frac{1}{x} \qquad y = \cos x$$

The diagrams show four of these graphs, labelled A, B, C and D.




Write down the correct equation for each graph.

$A =$ ..... , $B =$ ..... , $C =$ ..... , $D =$ .....

[4]

14 The diagram shows a sketch of the graph of $y = x^2 + bx + c$.
The minimum point on the graph is $M(3, -4)$.




(a) Work out the value of b and the value of c.

$b =$ ..... , $c =$ ..... **[3]**

(b) Find the coordinates of the points where the graph crosses the x-axis.

..... **[2]**

15 The diagram shows a sketch of the graph of $f(x) = a\sin(bx)$ for $0° \leq x \leq 360°$.



(a) Write down the value of a.

$a =$ ..... **[1]**

(b) Write down the value of b.

$b =$ ..... **[1]**

(c) Solve the equation $f(x) = 1.5$ for $0° \leq x \leq 180°$.


$x =$ ..... **[3]**

16 The graph of the quadratic function $f(x) = ax^2 + bx + c$ has a line of symmetry at $x = 2$.
The graph passes through the points $(0, -5)$ and $(2, 3)$.

Find the values of a, b and c.

$a =$ ..... , $b =$ ..... , $c =$ ..... **[5]**

17 The diagram shows a sketch of the curve with equation $y = x^3 + ax^2 + bx$.
The curve passes through the points $Q(-1, 3)$ and $R(2, 0)$.



(a) Use the coordinates of Q and R to form two simultaneous equations in terms of a and b.

.....

..... **[2]**


(b) Solve your equations to find the value of a and the value of b.

$a =$ ..... , $b =$ ..... **[2]**

(c) Find the coordinates of the other two points where the curve crosses the x-axis.

..... **[3]**

18 The diagram shows the graph of $y = f(x)$ where $f(x) = ax^3 + bx^2$.
The graph passes through the origin $(0, 0)$, the point $(3, 0)$ and has a local minimum point at $(2, -4)$.



(a) Show that $3a + b = 0$.

[1]


(b) Find the value of a and the value of b.

$a =$ ..... , $b =$ ..... **[3]**

(c) Write down the range of values of k for which the equation $f(x) = k$ has exactly one solution.

..... **[2]**

19 The diagram shows a sketch of the graph of $y = f(x)$, where $f(x) = ax^2 + bx + c$.
The curve passes through the origin O and intersects the x-axis at the point A.
The point $M(3, 18)$ is the maximum point of the curve.



(a) Write down the value of c.

$c =$ ..... **[1]**

(b) Write down the coordinates of A.

(..... , ...) **[1]**


(c) Find the value of a and the value of b.

$a =$ ..... , $b =$ ..... **[3]**

(d) Find the range of values of k for which the equation $f(x) = k$ has two distinct positive real roots.

..... **[2]**

20 The diagram shows the graph of the cubic function $y = ax^3 + bx^2 + cx + d$.



(a) Write down the value of d.

$d =$ ..... **[1]**

(b) The graph intersects the x-axis at $x = -3$ and $x = 3$.
Show that $b = 0$ and $9a + c = 0$.

[3]

(c) The graph passes through the point $(1, -16)$.
Find the value of a and the value of c.


$a =$ ..... , $c =$ ..... **[3]**

(d) A different function $g(x) = px^2 + qx + r$ has a minimum point at $(0, -5)$ and passes through the point $(2, 7)$.

Find the values of p, q and r.

$p =$ ..... , $q =$ ..... , $r =$ ..... **[3]**

21 The diagram shows the graph of $y = a\sin(bx)$ for $0° \leq x \leq 360°$.



(a) Write down the value of a.

$a =$ ..... **[1]**


(b) Find the value of b.

$b =$ ..... **[1]**

(c) Calculate the value of y when $x = 30°$.
Give your answer correct to 3 significant figures.

$y =$ ..... **[2]**

22 The diagram shows the graph of $y = a\sin(bx)$ for $0° \leq x \leq 360°$.



(a) Find the value of a and the value of b.

$a =$ ..... , $b =$ ..... **[2]**


(b) The graph of $y = p^x$ passes through the point $(3, 125)$.
Find the value of p.

$p =$ ..... **[1]**

(c) The graph of $y = \dfrac{k}{x}$ passes through the point $(0.5, 8)$.
Find the value of k.

$k =$ ..... **[1]**

23 The diagram shows the graph of $y = \frac{a}{x} + b$.
The equation of the horizontal asymptote is $y = 2$.
The graph passes through the point $(1, 6)$.



(a) Write down the value of b.

$b =$ ..... **[1]**


(b) Find the value of a.

$a =$ ..... **[2]**

(c) The graph of $y = k^x$, where $k > 0$, also passes through the point $(2, 4)$, which lies on the curve $y = \frac{a}{x} + b$.
Find the value of k.

$k =$ ..... **[2]**

24 The diagram shows a sketch of the graph of $f(x) = \frac{a}{x} + b$, where $xe0$.
The dashed line is a horizontal asymptote.
The graph passes through the point $(2, 7)$.



(a) Write down the value of b.

$b =$ ..... **[1]**


(b) Find the value of a.

$a =$ ..... **[2]**

(c) Find the coordinates of the point where the graph crosses the x-axis.

(..... , ...) **[2]**

25 The diagram shows a sketch of the graph of $f(x) = a\cos(bx)$ for $0° \leq x \leq 180°$.



(a) Write down the value of a.

$a =$ ..... **[1]**


(b) Find the value of b.

$b =$ ..... **[2]**

(c) Determine the number of solutions to the equation $a\cos(bx) = 1.5$ in the interval $0° \leq x \leq 360°$.

..... **[2]**

26 The diagram shows the graph of a quadratic function $f(x) = ax^2 + bx + c$.
The graph crosses the y-axis at $(0, -4)$ and the x-axis at $(-1, 0)$ and $(4, 0)$.



(a) Write down the value of c.

$c =$ ..... **[1]**


(b) Determine the value of a and the value of b.

$a =$ ..... , $b =$ ..... **[3]**

(c) Calculate the exact coordinates of the minimum point of the graph.

(..... , ...) **[2]**

27 The diagram shows a sketch of the graphs of $y = a^x$ and $y = \dfrac{b}{x}$ for $x > 0$.
The graphs intersect at the point $P(3, 8)$.



(a) Find the value of a and the value of b.

$a =$ ..... , $b =$ ..... **[3]**


(b) The graph of $y = a^x$ intersects the y-axis at the point Q.
Write down the coordinates of Q.

(..... , ...) **[1]**

(c) Calculate the gradient of the straight line joining Q and P.


Gradient = ... **[2]**

28 Four graphs A, B, C and D are shown below.



Graph A

Graph B



Graph C

Graph D

(a) Write down the letter of the graph which could represent the function

(i) $f(x) = \dfrac{a}{x}$ where $a > 0$

..... **[1]**

(ii) $f(x) = a^x$ where $a > 1$

..... **[1]**

(iii) $f(x) = -ax^2$ where $a > 0$

..... **[1]**

(b) The equation of Graph B is $y = p^x$.
The graph passes through the point $(3, 125)$.
Find the value of p.

$p =$ ..... **[2]**


(c) The equation of Graph C is $y = \dfrac{q}{x}$.

The graph passes through the point $(0.5, 8)$.

Find the value of q.

$q =$ ..... **[2]**

29 The diagram shows the graph of $y = ax^2 + bx + c$.
The graph intersects the x-axis at $(-1, 0)$ and $(5, 0)$, and the y-axis at $(0, -5)$.



(a) Find the values of a, b and c.

$a =$ ..... , $b =$ ..... , $c =$ ..... **[3]**


(b) Write down the coordinates of the minimum point of the graph.

(..... , ...) **[2]**

(c) The graph of $y = p \times q^x$ passes through the points $(0, 2)$ and $(2, 18)$, where $q > 0$.
Find the value of p and the value of q.

$p =$ ..... , $q =$ ..... **[3]**

30 The diagram shows a sketch of the curve $f(x) = x^3 + ax^2 + bx + c$.
The curve crosses the y-axis at $(0, 4)$.
It crosses the x-axis at $(-1, 0)$ and touches the x-axis at $(2, 0)$.



(a) Write down the value of c.

$c =$ ..... **[1]**


(b) By using the x-intercepts, find the value of a and the value of b.

$a =$ ..... , $b =$ ..... **[4]**

(c) Use the graph to state the number of solutions to the equation $f(x) = 3$.

..... **[1]**

31 The diagram shows a sketch of the graph of $f(x) = x^3 + bx^2 + cx + d$.
The curve intersects the y-axis at 4 and the x-axis at -1 and 2.
The point $(2, 0)$ is a local minimum.



(a) Write down the value of d.

$d =$ ..... **[1]**


(b) Find the values of b and c.

$b =$ ..... , $c =$ ..... **[4]**

(c) Write down the number of solutions to the equation $f(x) = -3$.

Number of solutions = ... **[1]**

32 The diagram shows the graph of $y = x^3 + bx^2 + cx + d$.
The graph crosses the axes at $(-2, 0)$, $(2, 0)$, $(p, 0)$ and $(0, 12)$.



(a) Write down the value of d.

$d =$ ..... **[1]**


(b) Find the values of b and c.

$b =$ ..... , $c =$ ..... **[4]**

(c) Find the value of p.

$p =$ ..... **[2]**

33 The diagram shows a sketch of the graph of $f(x) = ax^3 + bx$.
The curve passes through the points $P(2, -16)$ and $Q(-3, 9)$.



(a) Find the value of a and the value of b.

$a =$ ..... , $b =$ ..... **[3]**


(b) Find the coordinates of the local maximum point of the graph.

(..... , ...) **[2]**

(c) The equation $f(x) = k$ has exactly one solution.
Write down the range of values for k.

..... **[2]**

34 The diagram shows a sketch of the curve $y = ax^3 + bx$.
The curve passes through the origin O, and the points $(2, 10)$ and $(-1, 4)$.
The curve crosses the x-axis at the points A, O, and B.




(a) Find the value of a and the value of b.

$a =$ ..... , $b =$ ..... **[4]**

(b) Calculate the exact length of the line segment AB.

..... **[3]**

35 The diagram shows the graph of the exponential function $f(x) = a \cdot b^x + c$.
The horizontal line $y = -3$ is an asymptote to the graph.
The graph passes through the points $(0, -1)$ and $(1, 3)$.



(a) State the value of c.

$c =$ ..... **[1]**

(b) Work out the value of a and the value of b.


$a =$ ..... , $b =$ ..... **[4]**

(c) The graph of $g(x) = \dfrac{k}{x}$ intersects the graph of $f(x)$ at the point where $x = 1$.

Find the value of k.

$k =$ ..... **[2]**

36 The diagram shows the graph of a cubic function $f(x) = ax^3 + bx^2 + cx + d$.
The graph crosses the x-axis at $(-3, 0)$, $(1, 0)$ and $(4, 0)$.
It crosses the y-axis at $(0, -24)$.



(a) Find the values of a, b, c and d.


$a =$ ..... , $b =$ ..... , $c =$ ..... , $d =$ ..... **[6]**

(b) The graph of $y = f(x)$ is translated by the vector $\begin{pmatrix} 0 \\ 10 \end{pmatrix}$ to give the graph of $y = g(x)$.

Find the coordinates of the y-intercept of the graph of $y = g(x)$.

(..... , ...) **[2]**

37 The diagram shows part of the graph of $y = \frac{a}{x} + b$ for $x > 0$.
The graph passes through the points $(1, 8)$ and $(4, 3.5)$.



(a) Find the value of a and the value of b.

$a =$ ..... , $b =$ ..... **[3]**

(b) Write down the equation of the horizontal asymptote to the graph.

..... **[1]**

(c) The line $y = x - 3$ intersects the graph of $y = \frac{6}{x} + 2$ at the point P.


(i) Show that the x-coordinate of P satisfies the equation $x^2 - 5x - 6 = 0$.

[2]

(ii) Find the coordinates of P.

(..... , ...) **[2]**

38 The diagram shows a sketch of the graph of $f(x) = x^3 + bx^2 + cx + 4$.
The graph crosses the x-axis at $(-1, 0)$ and touches the x-axis at $(2, 0)$.
The graph crosses the y-axis at $(0, 4)$.



(a) **(i)** Use the point $(-1, 0)$ to show that $b - c = -3$.

[2]


(ii) Find the values of b and c.

$b =$ ..... , $c =$ ..... **[4]**

(b) The graph of $g(x) = p \cdot q^x$ passes through the points $(0, 2)$ and $(3, 54)$.
Given that $q > 0$, work out the values of p and q.

$p =$ ..... , $q =$ ..... **[3]**

39 The diagram shows the graphs of $y = f(x)$ and $y = g(x)$ for $x \geq 0$.
It is given that $f(x) = px^2 + qx + r$.
The graph of $y = f(x)$ has a maximum point at $M(2, 9)$ and intersects the y-axis at $P(0, 5)$.



(a) Find the values of p, q, and r.

$p =$ ..... , $q =$ ..... , $r =$ ..... **[4]**

(b) The graph of $y = f(x)$ intersects the positive x-axis at the point R.
Find the coordinates of R.

(..... , ...) **[2]**

(c) It is given that $g(x) = a\sin(bx^\circ)$, where $a > 0$ and $b > 0$.
The graph of $y = g(x)$ has its first positive x-intercept at the point R.
The amplitude of $g(x)$ is equal to the y-coordinate of P.

Find the value of a and the value of b.

$a =$ ..... , $b =$ ..... **[3]**

1 **(a)** 4 **[B1]**

(b) 2 **[B1]**

[Back to Question] *[Total Marks: 2]*

2 **(a)** 4 **[B1]**

(b) Identifies period is $180°$ OR uses $360 \div 180$ **[M1]**

2 **[A1]**

[Back to Question] *[Total Marks: 3]*

3 **(a)** -5 **[B1]**

(b) M1 for substituting $(2, 7)$ and their c into the equation, e.g. $7 = a(2)^2 - 5$

3 **[A1]**

[Back to Question] *[Total Marks: 3]*

4 **(a)** 4 **[B1]**

(b) $a = 4$ **[B1]**

$b = 3$ **[B1]**

[Back to Question] *[Total Marks: 3]*

5 **(a)** $a = 3$ **[B1]**

$b = 2$ **[B1]**

(b) $(90, -3)$ **[B1]**

[Back to Question] *[Total Marks: 3]*

6 **(a)** $a = 4$ **[B1]**

$b = 2$ **[B1]**

(b) 180 **[B1]**

[Back to Question] *[Total Marks: 3]*

7 **(a)** **(i)** B **[B1]**

(ii) A **[B1]**

(iii) D **[B1]**

(iv) C **[B1]**

[Back to Question] *[Total Marks: 4]*

8 **(a)** 5 **[B1]**

(b) 180 **[B1]**

(c) $a = 5$ **[B1]**

$b = 2$ **[B1]**

[Back to Question] *[Total Marks: 4]*

9 **(a)** 4 **[B1]**

(b) 180 **[B1]**

(c) $a = 4$ **[B1]**

$b = 2$ **[B1]**

[Back to Question] *[Total Marks: 4]*

10 **(a)** $a = 5$ **[B1]**

$b = 3$ **[B1]**

(b) $-24 = k(2^3)$ or $8k = -24$ **[M1]**

-3 **[A1]**

[Back to Question] *[Total Marks: 4]*

11 **(a)** $a = 4$ **[B1]**

$b = 3$ **[B1]**

(b) 120 **[B1]**

(c) 6 **[B1]**

[Back to Question] *[Total Marks: 4]*

12 **(a)** -8 **[B1]**

(b) Substitutes an x-intercept from the graph, e.g. $4^2 + 4b - 8 = 0$ **[M1]**

-2 **[A1]**

(c) $(1, -9)$ [B2]
(If incorrect, B1 for $x = 1$ or $y = -9$)

[Back to Question] *[Total Marks: 5]*

13 A: $y = 2^x$ **[B1]**

B: $y = \frac{1}{x}$ **[B1]**

C: $y = -x^2$ **[B1]**

D: $y = x^3$ **[B1]**

[Back to Question] *[Total Marks: 4]*

14 (a) x-coordinate of minimum is 3 $\implies -\frac{b}{2} = 3$ **[M1]**

$b = -6$ **[A1]**

Substitute $x = 3, y = -4, b = -6$ to find c: $-4 = 3^2 - 6(3) + c \implies c = 5$ **[A1]**

(b) $x^2 - 6x + 5 = 0$ **[M1]**

$(1, 0)$ and $(5, 0)$ **[A1]**

[Back to Question] *[Total Marks: 5]*

15 (a) 3 **[B1]**

(b) 2 **[B1]**

(c) M1 for setting up $3\sin(2x) = 1.5$ leading to $\sin(2x) = 0.5$

B1 for one correct value, e.g. 15° or 75°

15° and 75° **[A1]**

[Back to Question] *[Total Marks: 5]*

16 $c = -5$ **[B1]**

Uses line of symmetry $x = 2$ to form equation $-\frac{b}{2a} = 2 \implies b = -4a$ **[M1]**

Substitutes $(2, 3)$ and $c = -5$ into equation: $4a + 2b - 5 = 3$ **[M1]**

Solves simultaneous equations for a: $4a + 2(-4a) = 8 \implies -4a = 8 \implies a = -2$ **[M1]**

$b = -4(-2) = 8$ **[A1]**

[Back to Question] *[Total Marks: 5]*

17 (a) Substitute $(-1, 3)$: $(-1)^3 + a(-1)^2 + b(-1) = 3$

a - b = 4[$B1$]

Substitute $(2, 0)$: $(2)^3 + a(2)^2 + b(2) = 0$

2a + b = -4[$B1$]

(b) Valid method to solve simultaneous equations (e.g. addition: $3a = 0$) **[M1]**

$a = 0$ and $b = -4$ **[A1]**

(c) $x^3 - 4x = 0$ **[M1]**

$x(x^2 - 4) = 0$

x(x-2)(x+2) = 0[$M1$]

$(0, 0)$ and $(-2, 0)$ **[A1]**

[Back to Question] *[Total Marks: 7]*

18 **(a)** Substitutes $(3,0)$ into $f(x)$ to get $a(3)^3 + b(3)^2 = 0 \implies 27a + 9b = 0$, leading to $3a + b = 0$. **[B1]**

(b) Substitutes $(2,-4)$ to form a second equation: $8a + 4b = -4$ (or $2a + b = -1$) **[M1]**

Solves simultaneous equations (e.g. substitutes $b = -3a$ into $2a + b = -1$) **[M1]**

$a = 1, b = -3$ **[A1]**

(c) Identifies the turning points at $y = 0$ and $y = -4$ as critical values **[M1]**

$k < -4$ or $k > 0$ **[A1]**

[Back to Question] *[Total Marks: 6]*

19 **(a)** 0 **[B1]**

(b) $(6,0)$ **[B1]**

(c) Uses the vertex form $y = a(x-3)^2 + 18$ OR substitutes $(3,18)$ into $18 = 9a + 3b$ (knowing $c = 0$) **[M1]**

Uses a second point e.g., $(6,0)$ to set up $0 = 36a + 6b$ OR substitutes $(0,0)$ into vertex form: $0 = a(-3)^2 + 18$ **[M1]**

$a = -2, b = 12$ **[A1]**

(d) Identifies that $y = k$ must intersect the curve strictly between $y = 0$ and $y = 18$ on the positive x-axis **[M1]**

$0 < k < 18$ **[A1]**

[Back to Question] *[Total Marks: 7]*

20 **(a)** 0 **[B1]**

(b) Sets up equation for $x = 3$: $27a + 9b + 3c = 0$ (or $9a + 3b + c = 0$) **[M1]**

Sets up equation for $x = -3$: $-27a + 9b - 3c = 0$ (or $-9a + 3b - c = 0$) **[M1]**

Adds or subtracts equations correctly to deduce $18b = 0 \implies b = 0$, and substitutes to conclude $9a + c = 0$ **[A1]**

(c) Substitutes $x = 1, y = -16$ into the equation to get $a + c = -16$ **[M1]**

Solves $9a + c = 0$ and $a + c = -16$ simultaneously **[M1]**

$a = 2$, $c = -18$ **[A1]**

(d) $r = -5$ **[B1]**

$q = 0$ **[B1]**

Sets up $4p - 5 = 7$ to find $p = 3$ **[B1]**

[Back to Question] *[Total Marks: 10]*

21 **(a)** 4 **[B1]**

(b) 2 **[B1]**

(c) $4\sin(2 \times 30°)$ or $4\sin(60°)$ **[M1]**

3.46 **[A1]**

[Back to Question] *[Total Marks: 4]*

22 **(a)** $a = 4$ **[B1]**

$b = 2$ **[B1]**

(b) 5 **[B1]**

(c) 4 **[B1]**

[Back to Question] *[Total Marks: 4]*

23 **(a)** 2 **[B1]**

(b) Substitutes $(1, 6)$ into equation: $6 = \frac{a}{1} + 2$ **[M1]**

4 **[A1]**

(c) Substitutes $(2, 4)$ into $y = k^x$: $4 = k^2$ **[M1]**

2 **[A1]**

[Back to Question] *[Total Marks: 5]*

24 **(a)** 3 **[B1]**

(b) $7 = \frac{a}{2} + 3$ **[M1]**

$a = 8$ **[A1]**

(c) $\frac{8}{x} + 3 = 0$ **[M1]**

$\left(-\frac{8}{3}, 0\right)$ or $(-2.67, 0)$ **[A1]**

[Back to Question] *[Total Marks: 5]*

25 **(a)** 4 **[B1]**

(b) Recognises the period is 120° or sets $b \times 120 = 360$ **[M1]**

3 **[A1]**

(c) Recognises there are 2 solutions per period (or sketches horizontal line) **[M1]**

6 **[A1]**

[Back to Question] *[Total Marks: 5]*

26 **(a)** -4 **[B1]**

(b) Uses x-intercepts to set up equation: $a(-1)^2 + b(-1) - 4 = 0$ or $a(x+1)(x-4)$ **[M1]**

Substitute y-intercept $(0, -4)$ into factored form: $a(1)(-4) = -4 \implies a = 1$ **[M1]**

$b = -3$ **[A1]**

(c) Identifies x-coordinate of minimum as $x = \dfrac{-1+4}{2} = 1.5$ (or uses $x = \dfrac{-b}{2a}$) **[M1]**

$(1.5, -6.25)$ **[A1]**

[Back to Question] *[Total Marks: 6]*

27 **(a)** Uses $P(3, 8)$ in $y = a^x$: $a^3 = 8$ **[M1]**

$a = 2$ **[A1]**

Uses $P(3, 8)$ in $y = \dfrac{b}{x}$: $8 = \dfrac{b}{3} \implies b = 24$ **[B1]**

(b) $(0, 1)$ **[B1]**

(c) Gradient $= \dfrac{8-1}{3-0}$ **[M1]**

$\dfrac{7}{3}$ (or 2.33 or $2\dfrac{1}{3}$) **[A1]**

[Back to Question] *[Total Marks: 6]*

28 **(a)** **(i)** Graph C (or C) **[B1]**

(ii) Graph B (or B) **[B1]**

(iii) Graph D (or D) **[B1]**

(b) $p^3 = 125$ **[M1]**

$p = 5$ **[A1]**

(c) $\dfrac{q}{0.5} = 8$ **[M1]**

$q = 4$ **[A1]**

[Back to Question] *[Total Marks: 7]*

29 **(a)** $c = -5$ **[B1]**

Uses the roots to write $y = k(x+1)(x-5)$ **[M1]**

Expands to $x^2 - 4x - 5$ to find $a = 1$ and $b = -4$ **[A1]**

(b) $x = 2$ **[B1]**

$(2, -9)$ **[B1]**

(c) $2 = p \times q^0 \implies p = 2$ **[B1]**

$18 = 2 \times q^2 \implies q^2 = 9$ **[M1]**

$q = 3$ **[A1]**

[Back to Question] *[Total Marks: 8]*

30 **(a)** 4 **[B1]**

(b) Sets up equation using $(-1, 0)$: $(-1)^3 + a(-1)^2 + b(-1) + 4 = 0 \implies a - b = -3$ **[M1]**

Sets up equation using $(2, 0)$: $(2)^3 + a(2)^2 + b(2) + 4 = 0 \implies 4a + 2b = -12$ OR $2a + b = -6$ **[M1]**

Correct method to solve simultaneous equations (e.g. adding the equations) **[M1]**

$a = -3$ and $b = 0$ **[A1]**

(c) 3 **[B1]**

[Back to Question] *[Total Marks: 6]*

31 **(a)** 4 **[B1]**

(b) M1 for substituting $(-1, 0)$ to form an equation: $-1 + b - c + 4 = 0$ (or $b - c = -3$)

M1 for substituting $(2, 0)$ to form a second equation: $8 + 4b + 2c + 4 = 0$ (or $2b + c = -6$)

M1 for a valid method to solve their simultaneous equations

OR M2 for expanding $(x + 1)(x - 2)^2$

$b = -3$ **[A1]**

$c = 0$ **[A1]**

(c) 1 **[B1]**

[Back to Question] *[Total Marks: 6]*

32 **(a)** 12 **[B1]**

(b) Substitution of $(-2, 0)$ or $(2, 0)$ into the equation: $0 = -8 + 4b - 2c + 12$ or $0 = 8 + 4b + 2c + 12$ **[M1]**

Forming two correct simultaneous equations, e.g., $2b - c = -2$ and $2b + c = -10$ **[M1]**

$b = -3$ **[A1]**

$c = -4$ **[A1]**

(c) Setting up an equation for the roots, e.g., $(x - 2)(x + 2)(x - p) = x^3 - 3x^2 - 4x + 12$ OR $p^3 - 3p^2 - 4p + 12 = 0$ **[M1]**

$p = 3$ **[A1]**

[Back to Question] *[Total Marks: 7]*

33 **(a)** $8a + 2b = -16$ or $4a + b = -8$ **[M1]**

$-27a - 3b = 9$ or $-9a - b = 3$ **[M1]**

$a = 1, b = -12$ **[A1]**

(b) -2 **[B1]**

16 **[B1]**

(c) $k > 16$ **[B1]**

$k < -16$ **[B1]**

[Back to Question] *[Total Marks: 7]*

34 **(a)** $10 = a(2)^3 + b(2) \implies 8a + 2b = 10$ or $4a + b = 5$ **[M1]**

$4 = a(-1)^3 + b(-1) \implies -a - b = 4$ **[M1]**

Attempt to solve simultaneous equations (e.g., adding to eliminate b) **[M1]**

$a = 3$, $b = -7$ **[A1]**

(b) $3x^3 - 7x = 0$ **[M1]**

$x(3x^2 - 7) = 0 \implies x = \pm\sqrt{\frac{7}{3}}$ **[M1]**

Length $AB = 2 \times \sqrt{\frac{7}{3}} = \frac{2\sqrt{21}}{3}$ (or equivalent exact form) **[A1]**

[Back to Question] *[Total Marks: 7]*

35 **(a)** -3 **[B1]**

(b) Substitutes $(0, -1)$ into the equation: $a \cdot b^0 - 3 = -1$ **[M1]**

$a = 2$ **[A1]**

Substitutes $(1, 3)$ into the updated equation: $2 \cdot b^1 - 3 = 3$ **[M1]**

$b = 3$ **[A1]**

(c) Identifies that $g(1) = f(1)$, so $g(1) = 3$ **[M1]**

$\frac{k}{1} = 3 \implies k = 3$ **[A1]**

[Back to Question] *[Total Marks: 7]*

36 **(a)** Forms equation using roots: $f(x) = k(x + 3)(x - 1)(x - 4)$ **[M1]**

Substitutes $(0, -24)$ to find k: $k(3)(-1)(-4) = -24$ **[M1]**

$12k = -24 \implies k = -2$ **[A1]**

Expands brackets: $-2(x^2 + 2x - 3)(x - 4) = -2(x^3 - 4x^2 + 2x^2 - 8x - 3x + 12)$ **[M1]**

Simplifies to: $-2(x^3 - 2x^2 - 11x + 12) = -2x^3 + 4x^2 + 22x - 24$ **[M1]**

$a = -2, b = 4, c = 22, d = -24$ **[A1]**

(b) Recognises translation moves the y-intercept up by 10 (calculates $-24 + 10$) **[M1]**

$(0, -14)$ **[A1]**

[Back to Question] *[Total Marks: 8]*

37 **(a)** Sets up simultaneous equations: $8 = a + b$ and $3.5 = \frac{a}{4} + b$ **[M1]**

Subtracts to eliminate b: $4.5 = \frac{3a}{4} \implies a = 6$ **[A1]**

$b = 2$ **[A1]**

(b) $y = 2$ **[B1]**

(c) **(i)** $x - 3 = \frac{6}{x} + 2$ **[M1]**

$x - 5 = \frac{6}{x} \implies x^2 - 5x = 6 \implies x^2 - 5x - 6 = 0$ **[A1]**

(ii) Solves $(x - 6)(x + 1) = 0$ to get $x = 6$ (rejects $x = -1$ as $x > 0$) **[M1]**

$(6, 3)$ **[A1]**

[Back to Question] *[Total Marks: 8]*

38 **(a)** **(i)** $f(-1) = (-1)^3 + b(-1)^2 + c(-1) + 4 = 0$ **[M1]**

$-1 + b - c + 4 = 0$ leading to $b - c = -3$ **[A1]**

(ii) Use point $(2, 0)$ to form equation: $2^3 + b(2)^2 + c(2) + 4 = 0$ **[M1]**

$8 + 4b + 2c + 4 = 0$ leading to $2b + c = -6$ **[A1]**

Attempt to solve simultaneous equations $b - c = -3$ and $2b + c = -6$ **[M1]**

$b = -3$, $c = 0$ **[A1]**

(b) $p \cdot q^0 = 2 \implies p = 2$ **[B1]**

$2 \cdot q^3 = 54 \implies q^3 = 27$ **[M1]**

$q = 3$ **[A1]**

[Back to Question] *[Total Marks: 9]*

39 **(a)** $r = 5$ **[B1]**

Uses vertex $(2, 9)$ to form an equation, e.g. $9 = p(2)^2 + q(2) + 5$ or uses vertex form $y = p(x - 2)^2 + 9$ **[M1]**

Forms a second equation using symmetry, e.g.

$$\frac{-q}{2p} = 2$$

q = -4p, *or substitutes* (0,5) *into vertex form* 5 = p(0-2)^2 + 9 [M1]

$p = -1$ and $q = 4$ **[A1]**

(b) Equating $f(x)$ to zero: $-x^2 + 4x + 5 = 0$

$x^2 - 4x - 5 = 0$ [M1]

$(5, 0)$ **[A1]**

(c) $a = 5$ **[B1]**

First positive intercept occurs when $5b = 180$ **[M1]**

$b = 36$ **[A1]**

[Back to Question] *[Total Marks: 9]*