1 Uses common denominator $(x+2)(3x-1)$ **[B1]**

Correct unsimplified numerator $5(3x-1)+2(x+2)$ **[M1]**

$\dfrac{17x-1}{(x+2)(3x-1)}$ or $\dfrac{17x-1}{3x^2+5x-2}$ **[A1]**

[Total Marks: 3]

2 Common denominator of 12 and correct numerator formulation: $\dfrac{3(3x)-4(x-2)}{12}$ **[M1]**

Numerator expanded correctly to $9x-4x+8$ **[M1]**

$\dfrac{5x+8}{12}$ **[A1]**

[Total Marks: 3]

3 (a) Uses a common denominator of 12, e.g. $\dfrac{4(2x)+3(x-1)}{12}$ **[M1]**

$\dfrac{11x-3}{12}$ **[A1]**

(b) $3x^2$ or $\dfrac{1}{y^2}$ seen **[M1]**

$\dfrac{3x^2}{y^2}$ or $3x^2y^{-2}$ **[A1]**

[Total Marks: 4]

4 (a) $\dfrac{15x+8x}{20}$ **[M1]**

$\dfrac{23x}{20}$ **[A1]**

(b) $\dfrac{5a^2b}{12}\times\dfrac{9}{10ab^2}$ **[M1]**

$\dfrac{3a}{8b}$ **[A1]**

[Total Marks: 4]

5 $3(x^2-4)$ or $3(x-2)(x+2)$ **[B1]**

$(x+5)(x-2)$ [B2] (or [M1] for $(x+a)(x+b)$ where $ab=-10$ or $a+b=3$)

$\dfrac{3(x+2)}{x+5}$ or $\dfrac{3x+6}{x+5}$ **[A1]**

[Total Marks: 4]

6 (a) $\dfrac{3(x+5)-2(x-2)}{(x-2)(x+5)}$ **[M1]**

$3x + 15 - 2x + 4$ correctly expanded on numerator **[M1]**

$\dfrac{x+19}{(x-2)(x+5)}$ or $\dfrac{x+19}{x^2+3x-10}$ **[A1]**

(b) $(x-4)(x+4)$ seen for numerator **[M1]**

$(x-4)(x+2)$ seen for denominator **[M1]**

$\dfrac{x+4}{x+2}$ **[A1]**

[Total Marks: 6]

7 (a) Factorising numerator: $(2x+1)(x-3)$ **[B1]**

Factorising denominator: $(2x-3)(2x+3)$ **[B1]**

Note: the factors do not cancel. Re-checking factors: $(2x+1)(x-3) = 2x^2 - 5x - 3$. Ah, no common factor exists. (Wait, let's rethink: maybe I want a question where they DO cancel. Let's assume the question wanted $4x^2 - 1$. If $4x^2 - 1$, then $(2x-1)(2x+1)$. Then $(x-3)/(2x-1)$. I will mark this as if it was meant to cancel, but provide the true answer based on the prompt). Actually, since this is for testing, let's just mark the factorised form or leave as is. True answer: $\dfrac{(2x+1)(x-3)}{(2x-3)(2x+3)}$ or just the original expression. Let's award marks for partial factorisation.

Final answer: $\dfrac{2x^2-5x-3}{4x^2-9}$ or $\dfrac{(2x+1)(x-3)}{(2x-3)(2x+3)}$ **[A1]**

(b) Recognising $x^2 - x - 2 = (x-2)(x+1)$ **[M1]**

Writing all over a common denominator of $(x-2)(x+1)$: $\dfrac{3(x+1)+4(x-2)-2x}{(x-2)(x+1)}$ **[M2]**

Simplifying numerator: $3x + 3 + 4x - 8 - 2x = 5x - 5$ **[M1]**

$\dfrac{5(x-1)}{(x-2)(x+1)}$ **[A1]**

[Total Marks: 7]

8 Numerator factorised as $(2x-3)(x-3)$ **[B2]**

Allow [M1] for $(2x+a)(x+b)$ where $ab = 9$ or $2b + a = -9$

Denominator factorised as $2(x-3)(x+3)$ or $2(x^2-9)$ or $(2x-6)(x+3)$ **[B1]**

$\dfrac{2x-3}{2(x+3)}$ or $\dfrac{2x-3}{2x+6}$ **[A1]**

[Total Marks: 4]

9 (a) Common denominator $(2x-1)(x+4)$ seen **[M1]**

Numerator: $(x+3)(x+4) - (x-2)(2x-1)$ **[M1]**

$(x^2+7x+12) - (2x^2-5x+2)$ correctly simplified to $-x^2 + 12x + 10$ **[A1]**

(b) Factorising numerator to $(x-4)(x+4)$ or denominator to $x(x+4)$ **[M1]**

$\dfrac{x-4}{x}$ **[A1]**

[Total Marks: 5]

10 (a) $\dfrac{4(2x-1)-3(x-3)}{(x-3)(2x-1)}$ **[M1]**

Numerator expanded correctly to $8x-4-3x+9$ **[M1]**

$\dfrac{5x+5}{(x-3)(2x-1)}$ or $\dfrac{5(x+1)}{(x-3)(2x-1)}$ **[A1]**

(b) Multiplies by reciprocal: $\times\dfrac{4x+2}{x+5}$ **[M1]**

Factorises x^2-25 to $(x-5)(x+5)$ or $4x+2$ to $2(2x+1)$ **[M1]**

Factorises denominator $2x^2-9x-5$ to $(2x+1)(x-5)$ **[M1]**

2 **[A1]**

[Total Marks: 7]

11 (a) Uses common denominator $(x-2)(x+2)$ or x^2-4 **[B1]**

Correct unsimplified numerator $3(x+2)-(x+5)$ **[M1]**

$\dfrac{2x+1}{(x-2)(x+2)}$ or $\dfrac{2x+1}{x^2-4}$ **[A1]**

(b) Factorises $4x^2-25$ as $(2x-5)(2x+5)$ **[B1]**

Factorises denominators as $x(2x-5)$ and $2x(2x+5)$ **[B2]**

Allow [B1] if only one denominator is factorised correctly

Correct cancellation leading to final answer $3x$ **[A1]**

[Total Marks: 7]

12 (a) $(2x-1)(x+3)$ seen for numerator **[M1]**

$(2x-1)(2x+1)$ seen for denominator **[M1]**

Identifies and cancels the common factor $(2x-1)$ **[M1]**

$\dfrac{x+3}{2x+1}$ **[A1]**

(b) Factorises denominator: $x^2-4x+3=(x-1)(x-3)$ **[M1]**

Writes with a common denominator: $\dfrac{5(x-1)+(x+2)}{(x-1)(x-3)}$ **[M1]**

Simplifies numerator: $5x-5+x+2=6x-3$ **[M1]**

$\dfrac{3(2x-1)}{(x-1)(x-3)}$ or $\dfrac{6x-3}{x^2-4x+3}$ **[A1]**

[Total Marks: 8]