IGCSE 0607 Practice Questions — E2.5 — Combined Questions & Markscheme

Equations

Total Questions: 30 — **Total Marks:** 205 Marks

Table of Contents & Quick Navigation

- **E2.5** Equations
  → **Paper 2 (Non-Calculator)** *(17 Qs)* · **Paper 4 (Calculator)** *(13 Qs)*

1 Solve the equation.

$$\frac{3x-1}{4} = \frac{x+5}{3}$$

$x =$ ..... **[3]**

2 Solve the simultaneous equations.
You must show all your working.

$4x + 3y = 5$

$2x - 5y = 22$

$x =$ ..... , $y =$ ..... **[3]**

3 **(a)** Solve the equation.

$$4(x-3) = 2x + 10$$

$x =$ ..... **[2]**

(b) Rearrange the formula to make x the subject.

$$y = \frac{2x-1}{5}$$

$x =$ ..... **[2]**

4 **(a)** Solve the equation.

$3(2x - 1) = 5x + 8$

$x =$ ..... **[2]**

(b) Rearrange the formula to make x the subject.

$y = 4x - 7$

$x =$ ..... **[2]**

5 **(a)** Solve the equation.

$3(2x - 1) = 4x + 11$

$x =$ ..... **[2]**

(b) Solve the simultaneous equations.

$3x + 2y = 16$

$x - y = 2$

$x =$ ..... , $y =$ ..... **[3]**

6 **(a)** Solve

$$5x - 3 = 2(x + 6)$$

$x =$ ..... **[2]**

(b) Solve

$$\frac{x+2}{3} + \frac{2x-1}{4} = 3$$

$x =$ ..... **[3]**

7 **(a)** Solve the equation.

$$3(2x - 5) = 4x + 7$$

$x =$ ..... **[3]**

(b) Rearrange the formula to make x the subject.

$$y = \frac{2x + 3}{5}$$

$x =$ ..... **[2]**

8 **(a)** Solve the equation.

$$4(2x - 1) = 3x + 11$$

$x =$ ..... **[3]**

(b) Make x the subject of the formula.

$$y = \frac{3x - 5}{2}$$

$x =$ ..... **[3]**

9 Consider the following algebraic tasks.

(a) Solve the equation.

$$5(x-2) = 3x + 14$$

$x =$ ..... **[3]**

(b) Rearrange the formula to make b the subject.

$$A = \frac{h(a+b)}{2}$$

$b =$ ..... **[3]**

10 **(a)** Solve.

$4(x - 3) = 2x + 7$

$x =$ ..... **[3]**

(b) Solve the simultaneous equations.
You must show all your working.

$5x - 2y = 16$

$3x + 4y = 7$

$x =$ ..... , $y =$ ..... **[3]**

11 Rearrange the formula to make x the subject.

$$y = \frac{3x - 2}{x + 4}$$

$x =$ ..... **[4]**

12 **(a)** Solve the simultaneous equations.
You must show all your working.

$3x + 2y = 12$

$5x - 3y = 1$

$x =$ ..... , $y =$ ..... **[3]**

(b) Solve the equation by factorisation.

$2x^2 - 5x - 12 = 0$

$x =$ ..... , $x =$ ..... **[3]**

13 **(a)** Solve the simultaneous equations.
You must show all your working.

$3x + 2y = 12$

$5x - 3y = 1$

$x =$ ..... , $y =$ ..... **[4]**

(b) Solve the equation by factorisation.

$2x^2 - 7x - 15 = 0$

$x =$ ..... , or $x =$ ..... **[3]**

14 **(a)** Solve the simultaneous equations.
You must show all your working.

$3x - 2y = 19$

$4x + y = 18$

$x =$ ..... , $y =$ ..... **[3]**

(b) Rearrange the formula to make p the subject.

$$T = \frac{3p + 4}{p - 2}$$

$p =$ ..... **[4]**

15 **(a)** Solve the simultaneous equations.
You must show all your working.

$2x - y = 9$ $3x + 2y = 10$

$x =$ ..... , $y =$ ..... **[4]**

(b) Solve the equation.

$$\frac{2x}{3} - \frac{x-4}{4} = \frac{5}{2}$$

$x =$ ..... **[4]**

16 Solve the following equations.

(a) Solve the simultaneous equations.
You must show all your working.

$4x + 3y = 5$

$2x - 5y = 9$

$x =$ ..... , $y =$ ..... **[4]**

(b) Solve the equation by factorisation.

$3x^2 - 10x - 8 = 0$

$x =$ ..... or $x =$ **[4]**

17 Solve the equation.

$$\frac{2}{x-1} + \frac{1}{x-2} = 1$$

Give your answers in the form $a \pm \sqrt{b}$, where a and b are integers.
You must show all your working.

$x =$ ..... **[6]**

18 **(a)** Rearrange the formula to make x the subject.

$$p = \frac{3x - 1}{x + 4}$$

$x =$ ..... **[4]**

(b) Solve the equation by factorising.

$$2x^2 - 5x - 12 = 0$$

$x =$ ..... , $x =$ ..... **[3]**

19 **(a)** A rectangle has length $(3x + 2)$ cm and width $(x - 1)$ cm.
The perimeter of the rectangle is 42 cm.

Work out the value of x.

$x =$ ..... **[3]**

(b) Make x the subject of the formula.

$$p = \sqrt{\frac{x-3}{x}}$$

$x =$ ..... **[4]**

20 **(a)** Make x the subject of the formula.

$$y = \frac{2x + 5}{x - 3}$$


$x =$ ..... **[3]**

(b) Solve the equation.
Show all your working and give your answers correct to 2 decimal places.

$$\frac{x + 1}{3} - \frac{2}{x - 2} = 1$$

$x =$ ..... , $x =$ ..... **[6]**

21 The diagram shows a rectangle with length $(2x + 3)$ cm and width $(x - 1)$ cm.
The area of the rectangle is $45\,\text{cm}^2$.



NOT TO SCALE

(a) Show that $2x^2 + x - 48 = 0$.

[2]

(b) Solve the equation $2x^2 + x - 48 = 0$.
You must show all your working and give your answers correct to 2 decimal places.

$x =$ ..... , $x =$ ..... **[4]**

(c) Make x the subject of the formula.

$$y = \frac{ax + 5}{x - 3}$$

$x =$ ..... **[4]**

22 Solve the equation.
Show all your working and give your answers correct to 2 decimal places.

$$\frac{3}{x-2} + \frac{4}{2x+1} = 2$$

$x =$ ..... , $x =$ ..... **[6]**

23 Solve the equation.
Show all your working and give your answers correct to 2 decimal places.

$$\frac{5}{x-2} - \frac{3}{2x+1} = 2$$

$x =$ ..... , $x =$ ..... **[7]**

24 **(a)** Show that $\frac{3}{x-1} - \frac{2}{x+3} = 1$ simplifies to $x^2 + x - 14 = 0$.

[4]

(b) Solve the equation $x^2 + x - 14 = 0$.
Show all your working and give your answers correct to 2 decimal places.

$x =$ ..... , $x =$ ..... **[3]**

25 **(a)** A train travels 120 km at an average speed of x km/h.
It then travels a further 150 km at an average speed of $(x + 10)$ km/h.
The total time for the journey is 4 hours.
Write down an equation in x and show that it simplifies to $2x^2 - 115x - 600 = 0$.

[4]

(b) Solve the equation $2x^2 - 115x - 600 = 0$ to find the value of x.
Give your answer correct to 1 decimal place.

$x =$ ..... **[2]**

(c) Use your graphic display calculator to solve the equation.

$$2^x - 3 = \frac{5}{x}$$

Give your answers correct to 3 significant figures.

$x =$ ..... , $x =$ ..... **[2]**

26 **(a)** Solve the equation.

Give your answers in the exact form $\frac{p \pm \sqrt{q}}{2}$.

$$\frac{x}{x+2} = \frac{3}{x-6}$$

$x =$ ..... **[5]**

(b) Rearrange the formula to make x the subject.

$$y = \frac{2x-5}{3x+1}$$

$x =$ ..... **[4]**

27 A bus travels 100 km at an average speed of x km/h.
A car travels 100 km at an average speed of $(x + 15)$ km/h.
The car takes 30 minutes less than the bus.

(a) Write down an equation in x and show that it simplifies to $x^2 + 15x - 3000 = 0$.

[3]

(b) Solve the equation $x^2 + 15x - 3000 = 0$.
You must show all your working and give your answers correct to 2 decimal places.

$x =$ ..... , $x =$ ..... **[4]**

(c) Work out the time taken by the car to travel the 100 km.
Give your answer in hours and minutes, correct to the nearest minute.

hours, minutes **[3]**

28 **(a)** Make x the subject of the formula.

$$y = \frac{3x - 2}{x + 4}$$

$x =$ ..... **[4]**

(b) Solve the equation.
Show all your working and give your answers correct to 2 decimal places.

$$\frac{5}{x - 2} - \frac{3}{2x + 1} = 2$$

$x =$ ..... , or $x =$ ..... **[6]**

29 Consider the following algebraic equations and formulas.

(a) Rearrange the formula to make x the subject.

$$y = \frac{3x - 1}{x + 2}$$

$x =$ ..... **[4]**

(b) Solve the equation.
Give your answers correct to 2 decimal places.
Show all your working.

$$\frac{2}{x - 1} + \frac{3}{x + 2} = 1$$

$x =$ ..... or $x =$ **[6]**

30 A train travels 150 km at an average speed of x km/h.

(a) Write down an expression, in terms of x, for the time taken in hours.

..... hours **[1]**

(b) A bus travels the same distance of 150 km at an average speed of $(x - 15)$ km/h.
The bus takes 30 minutes longer than the train.
Show that x satisfies the equation $x^2 - 15x - 4500 = 0$.

[4]

(c) Solve the equation $x^2 - 15x - 4500 = 0$ to find the average speed of the train.

..... km/h **[3]**

(d) Rearrange the formula to make q the subject.

$$P = \frac{2q + 3}{q - 1}$$

$q =$ ..... **[4]**

1 $3(3x-1)=4(x+5)$ or $9x-3=4x+20$ **[M1]**

$5x=23$ **[M1]**

4.6 or $\frac{23}{5}$ **[A1]**

[Back to Question] *[Total Marks: 3]*

2 Multiplies one or both equations to equate coefficients of x or y (e.g. $4x-10y=44$) **[M1]**

Adds or subtracts equations to eliminate one variable (e.g. $13y=-39$) **[M1]**

$x=3.5$, $y=-3$ **[A1]**

[Back to Question] *[Total Marks: 3]*

3 (a) $4x-12=2x+10$ **[M1]**

$x=11$ **[A1]**

(b) $5y=2x-1$ **[M1]**

$\frac{5y+1}{2}$ **[A1]**

[Back to Question] *[Total Marks: 4]*

4 (a) $6x-3=5x+8$ **[M1]**

11 **[A1]**

(b) $y+7=4x$ or $\frac{y}{4}=x-\frac{7}{4}$ **[M1]**

$\frac{y+7}{4}$ or $\frac{y}{4}+\frac{7}{4}$ **[A1]**

[Back to Question] *[Total Marks: 4]*

5 (a) Expands brackets correctly to $6x-3=4x+11$ **[M1]**

7 **[A1]**

(b) Multiplies second equation by 2 to get $2x-2y=4$ OR substitutes $x=y+2$ into first equation **[M1]**

Correctly finds $x=4$ **[A1]**

Correctly finds $y=2$ **[A1]**

[Back to Question] *[Total Marks: 5]*

6 (a) $5x-3=2x+12$ **[M1]**

5 **[A1]**

(b) Multiply by 12 to get $4(x+2)+3(2x-1)=36$ (or equivalent common denominator method) **[M1]**

$4x + 8 + 6x - 3 = 36$ **[M1]**

$\frac{31}{10}$ or 3.1 **[A1]**

[Back to Question] *[Total Marks: 5]*

7 **(a)** Expands bracket to obtain $6x - 15$ **[M1]**

Rearranges terms to isolate x, e.g., $6x - 4x = 7 + 15$ or $2x = 22$ **[M1]**

11 **[A1]**

(b) Multiplies by 5 to obtain $5y = 2x + 3$ **[M1]**

$\frac{5y - 3}{2}$ **[A1]**

[Back to Question] *[Total Marks: 5]*

8 **(a)** $8x - 4 = 3x + 11$ **[M1]**

$5x = 15$ **[M1]**

3 **[A1]**

(b) $2y = 3x - 5$ **[M1]**

$2y + 5 = 3x$ **[M1]**

$\frac{2y + 5}{3}$ or $\frac{2y}{3} + \frac{5}{3}$ **[A1]**

[Back to Question] *[Total Marks: 6]*

9 **(a)** $5x - 10 = 3x + 14$ **[M1]**

$2x = 24$ **[M1]**

12 **[A1]**

(b) $2A = h(a + b)$ or $\frac{A}{h} = \frac{a + b}{2}$ **[M1]**

$\frac{2A}{h} = a + b$ or $2A = ha + hb$ **[M1]**

$\frac{2A}{h} - a$ or $\frac{2A - ah}{h}$ **[A1]**

[Back to Question] *[Total Marks: 6]*

10 **(a)** Expands brackets: $4x - 12 = 2x + 7$ **[M1]**

Rearranges to collect terms: $2x = 19$ **[M1]**

9.5 or $\frac{19}{2}$ **[A1]**

(b) Multiplies first equation by 2 to equate coefficients of y: $10x - 4y = 32$ (or equivalent valid method) **[M1]**

Adds equations to eliminate y: $13x = 39 \implies x = 3$ **[A1]**

Substitutes $x = 3$ into either equation to find y: $15 - 2y = 16 \implies y = -0.5$ **[A1]**

[Back to Question] *[Total Marks: 6]*

11 $y(x + 4) = 3x - 2$ **[M1]**

$xy - 3x = -4y - 2$ or $3x - xy = 4y + 2$ **[M1]**

$x(y - 3) = -4y - 2$ or $x(3 - y) = 4y + 2$ **[M1]**

$\frac{4y + 2}{3 - y}$ or $\frac{-4y - 2}{y - 3}$ **[A1]**

[Back to Question] *[Total Marks: 4]*

12 (a) Multiplies equations to equate coefficients of x or y, e.g., $9x + 6y = 36$ and $10x - 6y = 2$ **[M1]**

Adds or subtracts equations to eliminate a variable **[M1]**

$x = 2$ and $y = 3$ **[A1]**

(b) $(2x \pm a)(x \pm b) = 0$ where $ab = 12$ **[M1]**

$(2x + 3)(x - 4) = 0$ **[M1]**

$x = 4$ and $x = -1.5$ (or $-\frac{3}{2}$) **[A1]**

[Back to Question] *[Total Marks: 6]*

13 (a) Multiplies equations to equate coefficients, e.g., $9x + 6y = 36$ and $10x - 6y = 2$ **[M1]**

Adds or subtracts equations to eliminate one variable, e.g., $19x = 38$ **[M1]**

$x = 2$ **[A1]**

$y = 3$ **[A1]**

(b) Finds correct product and sum components, e.g., $-10x$ and $3x$ **[M1]**

Correctly factorises to $(2x + 3)(x - 5) = 0$ **[M1]**

$x = -1.5, x = 5$ **[A1]**

[Back to Question] *[Total Marks: 7]*

14 (a) Multiply the second equation by 2 to get $8x + 2y = 36$ (or correct substitution method) **[M1]**

$x = 5$ **[A1]**

$y = -2$ **[A1]**

(b) $T(p - 2) = 3p + 4$ **[M1]**

$Tp - 2T = 3p + 4$ **[M1]**

$Tp - 3p = 2T + 4$ or $p(T - 3) = 2T + 4$ **[M1]**

$\frac{2T+4}{T-3}$ or $\frac{-2T-4}{3-T}$ **[A1]**

[Back to Question] *[Total Marks: 7]*

15 (a) Multiplies the first equation by 2 to give $4x - 2y = 18$ (or rearranges to $y = 2x - 9$) **[M1]**

Adds the equations to eliminate y, yielding $7x = 28$ (or correctly substitutes to form an equation in one variable) **[M1]**

$x = 4$ **[A1]**

$y = -1$ **[A1]**

(b) Multiplies throughout by a common denominator (e.g., 12) to obtain $4(2x) - 3(x-4) = 30$ **[M1]**

Expands brackets correctly: $8x - 3x + 12 = 30$ **[M1]**

$5x = 18$ **[M1]**

3.6 or $\frac{18}{5}$ **[A1]**

[Back to Question] *[Total Marks: 8]*

16 (a) Multiply second equation by 2: $4x - 10y = 18$ **[M1]**

Subtract equations to eliminate x: $13y = -13$ **[M1]**

$y = -1$ **[A1]**

$x = 2$ **[A1]**

(b) $(3x+2)(x-4) = 0$ **[M2]**

If M0, M1 for $(3x \pm a)(x \pm b)$ where $ab = 8$ or $a \pm 3b = -10$

$x = -\frac{2}{3}$ **[A1]**

$x = 4$ **[A1]**

[Back to Question] *[Total Marks: 8]*

17 Forms common denominator or multiplies through: $2(x-2) + 1(x-1) = (x-1)(x-2)$ **[M1]**

Expands brackets correctly: $2x - 4 + x - 1 = x^2 - 3x + 2$ **[M1]**

Rearranges to a 3-term quadratic: $x^2 - 6x + 7 = 0$ **[A1]**

Applies quadratic formula: $\frac{-(-6) \pm \sqrt{(-6)^2 - 4(1)(7)}}{2(1)}$ or completes the square: $(x-3)^2 - 9 + 7 = 0$ **[M1]**

Correct unsimplified surd form: $\frac{6 \pm \sqrt{8}}{2}$ or $(x-3)^2 = 2$ **[M1]**

$3 \pm \sqrt{2}$ **[A1]**

[Back to Question] *[Total Marks: 6]*

18 **(a)** $p(x+4) = 3x - 1$ **[M1]**

$px + 4p = 3x - 1$ **[M1]**

$3x - px = 4p + 1$ (or equivalent step isolating x terms on one side) **[M1]**

$\frac{4p+1}{3-p}$ (or $\frac{-4p-1}{p-3}$) **[A1]**

(b) $(2x \pm a)(x \pm b) = 0$ where $ab = -12$ or $2b \pm a = -5$ **[M1]**

$(2x+3)(x-4) = 0$ **[M1]**

$x = -1.5$, $x = 4$ **[A1]**

[Back to Question] *[Total Marks: 7]*

19 **(a)** $2(3x + 2 + x - 1) = 42$ or $3x + 2 + x - 1 = 21$ **[M1]**

$4x + 1 = 21$ **[M1]**

5 **[A1]**

(b) $p^2 = \frac{x-3}{x}$ **[M1]**

$p^2x = x - 3$ **[M1]**

$p^2x - x = -3$ or $x - p^2x = 3$ or $x(p^2 - 1) = -3$ **[M1]**

$\frac{3}{1-p^2}$ or $\frac{-3}{p^2-1}$ **[A1]**

[Back to Question] *[Total Marks: 7]*

20 **(a)** Multiplies by denominator: $y(x-3) = 2x + 5$ **[M1]**

Expands and collects x terms on one side: $xy - 2x = 3y + 5$ **[M1]**

$\frac{3y+5}{y-2}$ or $\frac{-3y-5}{2-y}$ **[A1]**

(b) Multiplies throughout by common denominator $3(x-2)$ to reach $(x+1)(x-2) - 6 = 3(x-2)$ **[M1]**

Expands brackets correctly: $x^2 - 2x + x - 2 - 6 = 3x - 6$ **[M1]**

Rearranges to form a correct 3-term quadratic equation: $x^2 - 4x - 2 = 0$ **[A1]**

Uses the quadratic formula $\frac{-(-4) \pm \sqrt{(-4)^2 - 4(1)(-2)}}{2(1)}$ **[M1]**

4.45 **[A1]**

-0.45 **[A1]**

[Back to Question] *[Total Marks: 9]*

21 **(a)** Sets up equation for area: $(2x+3)(x-1) = 45$ **[M1]**

Expands and simplifies to given result: $2x^2 - 2x + 3x - 3 = 45 \implies 2x^2 + x - 48 = 0$ **[A1]**

(b) Uses quadratic formula with correct values: $\frac{-1 \pm \sqrt{1^2 - 4(2)(-48)}}{2(2)}$ **[M1]**

Correct discriminant: $\frac{-1 \pm \sqrt{385}}{4}$ **[M1]**

4.655... and $-5.155...$ **[A1]**

4.66 and -5.16 **[A1]**

(c) Multiplies by denominator: $y(x-3) = ax+5$ **[M1]**

Expands brackets: $xy - 3y = ax + 5$ **[M1]**

Collects terms in x on one side: $xy - ax = 3y + 5$ **[M1]**

Factorises x and divides: $\frac{3y+5}{y-a}$ (or equivalent) **[A1]**

[Back to Question] *[Total Marks: 10]*

22 Multiplies by the common denominator to get $3(2x+1)+4(x-2) = 2(x-2)(2x+1)$ [M2]
(allow M1 for $\frac{3(2x+1)+4(x-2)}{(x-2)(2x+1)} = 2$)

Expands brackets to get $10x - 5 = 2(2x^2 - 3x - 2)$ or $10x - 5 = 4x^2 - 6x - 4$ **[M1]**

Rearranges to form a correct 3-term quadratic equation equal to zero: $4x^2 - 16x + 1 = 0$ **[A1]**

Substitutes correctly into the quadratic formula: $x = \frac{-(-16) \pm \sqrt{(-16)^2 - 4(4)(1)}}{2(4)}$ **[M1]**

$x = 3.94$ and $x = 0.06$ **[A1]**

[Back to Question] *[Total Marks: 6]*

23 Multiplies by the common denominator to get $5(2x+1) - 3(x-2) = 2(x-2)(2x+1)$ **[M2]**

If not M2, M1 for a correct single fraction $\frac{5(2x+1)-3(x-2)}{(x-2)(2x+1)}$ seen

$4x^2 - 13x - 15 = 0$ **[A2]**

If not A2, A1 for correctly expanding sides to $7x + 11$ or $4x^2 - 6x - 4$

$x = \frac{-(-13) \pm \sqrt{(-13)^2 - 4(4)(-15)}}{2(4)}$ **[M1]**

4.15 **[A1]**

-0.90 **[A1]**

[Back to Question] *[Total Marks: 7]*

24 (a) $\frac{3(x+3)-2(x-1)}{(x-1)(x+3)} = 1$ **[M1]**

$3x + 9 - 2x + 2 = (x-1)(x+3)$ **[M1]**

$x + 11 = x^2 + 2x - 3$ **[M1]**

$x^2 + x - 14 = 0$ established with no errors seen **[A1]**

(b) $\frac{-1 \pm \sqrt{1^2 - 4(1)(-14)}}{2(1)}$ **[M1]**

$\frac{-1 \pm \sqrt{57}}{2}$ **[M1]**

$3.27, -4.27$ **[A1]**

[Back to Question] *[Total Marks: 7]*

25 **(a)** Constructs initial time equation: $\frac{120}{x} + \frac{150}{x+10} = 4$ **[M1]**

Multiplies by common denominator $x(x+10)$ to reach $120(x+10) + 150x = 4x(x+10)$ **[M1]**

Expands brackets correctly: $120x + 1200 + 150x = 4x^2 + 40x$ **[M1]**

Rearranges to $4x^2 - 230x - 1200 = 0$ and divides by 2 to yield $2x^2 - 115x - 600 = 0$ with no errors seen **[A1]**

(b) Substitutes values into the quadratic formula $\frac{115 \pm \sqrt{(-115)^2 - 4(2)(-600)}}{4}$ **[M1]**

62.3 **[A1]**

(c) -1.84 **[B1]**

2.36 **[B1]**

[Back to Question] *[Total Marks: 8]*

26 **(a)** $x(x-6) = 3(x+2)$ **[M1]**

$x^2 - 6x = 3x + 6$ **[M1]**

$x^2 - 9x - 6 = 0$ **[A1]**

Uses the quadratic formula, e.g., $x = \frac{-(-9) \pm \sqrt{(-9)^2 - 4(1)(-6)}}{2(1)}$ **[M1]**

$\frac{9 \pm \sqrt{105}}{2}$ **[A1]**

(b) $y(3x+1) = 2x - 5$ **[M1]**

$3xy + y = 2x - 5$ **[M1]**

Isolates terms containing x on one side, e.g., $3xy - 2x = -y - 5$ or $2x - 3xy = y + 5$ **[M1]**

$\frac{-y-5}{3y-2}$ or $\frac{y+5}{2-3y}$ **[A1]**

[Back to Question] *[Total Marks: 9]*

27 **(a)** $\frac{100}{x} - \frac{100}{x+15} = \frac{1}{2}$ (or equivalent e.g., $\frac{100}{x} = \frac{100}{x+15} + 0.5$) **[M1]**

$200(x+15) - 200x = x(x+15)$ **[M1]**

$200x + 3000 - 200x = x^2 + 15x$ leading to $x^2 + 15x - 3000 = 0$ **[A1]**

(b) $\frac{-15 \pm \sqrt{15^2 - 4(1)(-3000)}}{2(1)}$ **[M2]**

Allow [M1] for correctly substituting values into the quadratic formula or evaluating the

discriminant $\sqrt{12225}$

47.78 **[A1]**

-62.78 **[A1]**

(c) Speed of car $= 47.78 + 15 = 62.78$ km/h **[M1]**

Time $= \dfrac{100}{62.78} = 1.592...$ hours **[M1]**

1 hour 36 minutes **[A1]**

[Back to Question] *[Total Marks: 10]*

28 (a) Multiplies by the denominator to obtain $y(x+4) = 3x - 2$ **[M1]**

Expands the bracket to obtain $xy + 4y = 3x - 2$ **[M1]**

Collects terms in x on one side, e.g., $xy - 3x = -4y - 2$ or $3x - xy = 4y + 2$ **[M1]**

Factorises x out and divides: $\dfrac{-4y-2}{y-3}$ or $\dfrac{4y+2}{3-y}$ **[A1]**

(b) Writes with common denominator or multiplies through to eliminate fractions:
$5(2x+1) - 3(x-2) = 2(x-2)(2x+1)$ **[M1]**

Correctly expands LHS: $10x + 5 - 3x + 6$ **[M1]**

Correctly expands RHS: $2(2x^2 - 3x - 2)$ or $4x^2 - 6x - 4$ **[M1]**

Rearranges to form a correct 3-term quadratic equation, e.g., $4x^2 - 13x - 15 = 0$ **[A1]**

Correctly substitutes into the quadratic formula: $\dfrac{-(-13) \pm \sqrt{(-13)^2 - 4(4)(-15)}}{2(4)}$ **[M1]**

$x = 4.15$ and $x = -0.90$ **[A1]**

[Back to Question] *[Total Marks: 10]*

29 (a) $y(x+2) = 3x - 1$ **[M1]**

$yx + 2y = 3x - 1$ **[M1]**

$2y + 1 = 3x - yx$ or $yx - 3x = -2y - 1$ **[M1]**

$x = \dfrac{2y+1}{3-y}$ or $x = \dfrac{-2y-1}{y-3}$ **[A1]**

(b) Multiply throughout by $(x-1)(x+2)$: $2(x+2) + 3(x-1) = (x-1)(x+2)$ **[M1]**

Expand terms: $2x + 4 + 3x - 3 = x^2 + x - 2$ **[M1]**

Collect to form a quadratic equation: $x^2 - 4x - 3 = 0$ **[A1]**

Use the quadratic formula: $x = \dfrac{-(-4) \pm \sqrt{(-4)^2 - 4(1)(-3)}}{2(1)}$ **[M1]**

$x = \dfrac{4 \pm \sqrt{28}}{2}$ **[M1]**

$x = 4.65$ and $x = -0.65$ **[A1]**

[Back to Question] *[Total Marks: 10]*

30 **(a)** $\dfrac{150}{x}$ **[B1]**

(b) $\dfrac{150}{x-15} - \dfrac{150}{x} = 0.5$ (or $\dfrac{30}{60}$ or $\dfrac{1}{2}$) **[M1]**

$150x - 150(x-15) = 0.5x(x-15)$ **[M1]**

$150x - 150x + 2250 = 0.5x^2 - 7.5x$ **[M1]**

$4500 = x^2 - 15x \implies x^2 - 15x - 4500 = 0$ with no errors seen **[A1]**

(c) $(x-75)(x+60) = 0$ or $\dfrac{-(-15) \pm \sqrt{(-15)^2 - 4(1)(-4500)}}{2(1)}$ **[M2]**

(or $M1$ for correct factors $(x+a)(x+b)$ where $ab = -4500$ or $a+b = -15$, or for correct substitution into quadratic formula)

75 (rejects negative solution -60) **[A1]**

(d) $P(q-1) = 2q+3$ **[M1]**

$Pq - P = 2q + 3$ **[M1]**

$Pq - 2q = P + 3$ **[M1]**

$\dfrac{P+3}{P-2}$ or $\dfrac{-P-3}{2-P}$ **[A1]**

[Back to Question] *[Total Marks: 12]*