IGCSE 0607 Practice Questions — E2.7 — Combined Questions & Markscheme

Sequences

Total Questions: 12 — **Total Marks:** 72 Marks

Table of Contents & Quick Navigation

- **E2.7** Sequences
  → **Paper 2 (Non-Calculator)** *(5 Qs)* · **Paper 4 (Calculator)** *(7 Qs)*

---

1 Here are the first four terms of a sequence.

5 12 19 26

(a) Write down the next term in the sequence.

..... **[1]**

(b) Find an expression for the nth term of the sequence.

..... **[2]**

(c) Explain why 150 is not a term in this sequence.

..... **[1]**

2 Here are the first four terms of a sequence.

5 11 17 23

(a) Find the next term in this sequence.

..... **[1]**

(b) Find an expression for the nth term of this sequence.

..... **[2]**

(c) Explain why 150 is not a term in this sequence.

..... **[1]**

3 The first four terms of a sequence are

7 11 15 19

(a) Write down the next two terms of the sequence.

..... **[1]**

(b) Find an expression for the nth term of this sequence.

nth term = ... **[2]**

(c) Explain why 102 is not a term in this sequence.

..... **[2]**

4 **(a)** Sequence A is a linear sequence.

$$5 \quad 9 \quad 13 \quad 17 \quad \ldots$$

(i) Find the next two terms of sequence A.

..... **[1]**

(ii) Find the nth term of sequence A.

..... **[2]**

(b) Sequence B is a geometric sequence.

$$2 \quad 6 \quad 18 \quad 54 \quad \ldots$$

Find the nth term of sequence B.

..... **[2]**

5 The first five terms of a sequence, T_n, are given below.

$$4 \qquad 14 \qquad 34 \qquad 68 \qquad 120$$

(a) By finding the first, second and third differences, show that the nth term of the sequence is a cubic expression in n.

[2]

(b) Find an expression for T_n, the nth term of the sequence.

$T_n =$ ..... **[5]**

6 **(a)** The nth term of a sequence is $n^2 - 3n + 5$.
Work out the first three terms of this sequence.

1st term = ... , 2nd term = ... , 3rd term = ...
[2]

(b) The first four terms of another sequence are

2 9 20 35

Find an expression for the nth term of this sequence.

..... **[3]**

7 The first four terms of a sequence, T, are

$$2 \quad 9 \quad 22 \quad 41$$

(a) Find an expression for the nth term, T_n.

$T_n =$ ..... **[3]**

(b) A second sequence is given by the formula $U_n = 2^n + 3$.
Work out the value of U_5.

$U_5 =$ ..... **[1]**

(c) Find the value of n for which $U_n = 131$.

$n =$ ..... **[2]**

8 The first four terms of a sequence are

$$-1 \quad 2 \quad 9 \quad 20$$

(a) Find the nth term of this sequence.

..... **[3]**

(b) The nth term of another sequence is given by $3^n - 2$.
Find the value of the term in this sequence that is closest to 1000.

..... **[3]**

9 The first five terms of a sequence are

2 9 20 35 54 ...

(a) Find the next term.

..... **[1]**

(b) Find the nth term of the sequence.

..... **[3]**

(c) The kth term of the sequence is 464.
Find the value of k.

$k =$ ..... **[3]**

10 Consider the following three sequences, P, Q and R.
Sequence P:

$$2 \quad 9 \quad 28 \quad 65 \quad 126 \quad \ldots$$

Sequence Q:

$$3 \quad 8 \quad 15 \quad 24 \quad 35 \quad \ldots$$

Sequence R:

$$\frac{2}{3} \quad \frac{9}{8} \quad \frac{28}{15} \quad \frac{65}{24} \quad \frac{126}{35} \quad \ldots$$

(a) Find the nth term of sequence P.

..... **[2]**

(b) Find the nth term of sequence Q.

..... **[2]**

(c) Write down the nth term of sequence R.

..... **[1]**

(d) Find the 12th term of sequence R, giving your answer as a fraction in its simplest form.

..... **[2]**

11 **(a)** The first five terms of a sequence are

$$-1 \quad 8 \quad 33 \quad 80 \quad 155$$

Use the method of differences to find an expression for the nth term of this sequence.

..... **[4]**

(b) An exponential sequence has nth term $P_n = 5 \times 3^{n-1}$.
The kth term of this sequence is 3645.
Work out the value of k.

$k =$ ..... **[2]**

(c) The nth term of sequence T is $T_n = \dfrac{2n-1}{3n+2}$.
Write down the value that T_n approaches as n becomes very large.

..... **[1]**

12 **(a)** A sequence has the following first four terms:

$$1 \quad 15 \quad 53 \quad 127$$

Use the method of differences to find an expression for the nth term.

nth term = ... **[4]**

(b) Another sequence, V, has its nth term defined by

$$V_n = \frac{n^2 + 1}{2^n}$$

(i) Find the values of V_1, V_2 and V_3.
Give each fraction in its simplest form.

V_1 = ... , V_2 = ... , V_3 = ... **[2]**

(ii) Work out the exact value of $V_5 - V_4$.

..... **[3]**

1 **(a)** 33 **[B1]**

(b) $7n + k$ for any integer k **[M1]**

$7n - 2$ **[A1]**

(c) $7n - 2 = 150$

$7\text{n} = 152 whichdoesnotgiveanintegervaluefor\text{n}.[B1]$

[Back to Question] *[Total Marks: 4]*

2 **(a)** 29 **[B1]**

(b) $6n + c$ or finding common difference of 6 **[M1]**

$6n - 1$ **[A1]**

(c) Sets $6n - 1 = 150 \implies 6n = 151$ and states that 151 is not divisible by 6 (or n is not an integer) **[B1]**

[Back to Question] *[Total Marks: 4]*

3 **(a)** $23, 27$ **[B1]**

(b) $4n + k$ (for any k) **[M1]**

$4n + 3$ **[A1]**

(c) Equates $4n + 3 = 102$ and attempts to solve for n **[M1]**

$n = 24.75$, which is not a whole number/integer **[A1]**

[Back to Question] *[Total Marks: 5]*

4 **(a)** **(i)** 21 and 25 **[B1]**

(ii) Identifies common difference of 4 leading to $4n$ **[M1]**

$4n + 1$ **[A1]**

(b) Identifies common ratio of 3 **[M1]**

$2 \times 3^{n-1}$ **[A1]**

[Back to Question] *[Total Marks: 5]*

5 **(a)** First differences: $10, 20, 34, 52$ **[M1]**

Second differences: $10, 14, 18$
Third differences: $4, 4$
Since the third difference is a non-zero constant, the sequence is cubic. **[A1]**

(b) Uses $6a = 4 \implies a = \dfrac{2}{3}$ **[M1]**

Uses $12a + 2b = 10 \implies 12(\dfrac{2}{3}) + 2b = 10 \implies 8 + 2b = 10 \implies b = 1$ **[M1]**

Uses $7a+3b+c=10 \implies 7(\frac{2}{3})+3(1)+c=10 \implies c=10-3-\frac{14}{3}=\frac{7}{3}$ **[M1]**

Uses $a+b+c+d=4 \implies \frac{2}{3}+1+\frac{7}{3}+d=4 \implies 4+d=4 \implies d=0$ **[M1]**

$\frac{2}{3}n^3+n^2+\frac{7}{3}n$ **[A1]**

[Back to Question] *[Total Marks: 7]*

6 **(a)** One correct term evaluated **[M1]**

$3, 3, 5$ **[A1]**

(b) Second differences are 4 **[M1]**

Identifies $2n^2$ as part of the expression **[M1]**

$2n^2+n-1$ **[A1]**

[Back to Question] *[Total Marks: 5]*

7 **(a)** Finds first differences of $7, 13, 19$ and constant second differences of 6 **[M1]**

Deduces the quadratic term is $3n^2$ **[M1]**

$3n^2-2n+1$ **[A1]**

(b) 35 **[B1]**

(c) $2^n+3=131$ leading to $2^n=128$ **[M1]**

7 **[A1]**

[Back to Question] *[Total Marks: 6]*

8 **(a)** Finds first differences: $3, 7, 11$ and second differences: $4, 4$ **[M1]**

Deduces the term involves $2n^2$ **[M1]**

$2n^2-3$ **[A1]**

(b) $3^n-2 \approx 1000 \implies 3^n \approx 1002$ **[M1]**

Uses logarithms or trial and improvement: $n=6 \implies 3^6-2=729-2=727$;
$n=7 \implies 3^7-2=2187-2=2185$ **[M1]**

727 **[A1]**

[Back to Question] *[Total Marks: 6]*

9 **(a)** 77 **[B1]**

(b) Finds first differences $(7, 11, 15, 19)$ and constant second difference of 4 **[M1]**

Deduces the coefficient of n^2 is 2 **[A1]**

Subtracts $2n^2$ from the sequence terms to identify the linear sequence $n-1$ **[M1]**

$2n^2+n-1$ **[A1]**

(c) Sets up the equation $2k^2 + k - 1 = 464$ **[M1]**

Rearranges to form a quadratic equal to zero: $2k^2 + k - 465 = 0$ **[M1]**

15 **[A1]**

[Back to Question] *[Total Marks: 7]*

10 (a) Recognises the relationship to cube numbers $(1, 8, 27, 64, 125)$ **[M1]**

$n^3 + 1$ **[A1]**

(b) Identifies second difference is 2, or relates to square numbers **[M1]**

$n^2 + 2n$ or $(n+1)^2 - 1$ **[A1]**

(c) $\dfrac{n^3 + 1}{n^2 + 2n}$ **[B1]**

(d) Substitutes $n = 12$ into their expression for sequence R, yielding $\dfrac{1728 + 1}{144 + 24}$ or $\dfrac{1729}{168}$ **[M1]**

Simplifies the fraction correctly to $\dfrac{247}{24}$ **[A1]**

[Back to Question] *[Total Marks: 7]*

11 (a) Finds third differences are 6 (first diffs: $9, 25, 47, 75$; second diffs: $16, 22, 28$; third diffs: 6) **[M1]**

Identifies n^3 as the cubic term **[B1]**

Subtracts n^3 from the sequence to get a quadratic sequence $(-2, 0, 6, 16, 30)$ and finds its second differences are 4 to give $2n^2$ **[M1]**

$n^3 + 2n^2 - 4n$ **[A1]**

(b) $5 \times 3^{k-1} = 3645$

$3^{k-1} = 729$ [M1]

7 **[A1]**

(c) $\dfrac{2}{3}$ or 0.667 **[B1]**

[Back to Question] *[Total Marks: 7]*

12 (a) Finds first differences $14, 38, 74$ and second differences $24, 36$ **[M1]**

Finds third difference of 12 leading to a cubic term of $2n^3$ **[M1]**

Evaluates the sequence of residuals $T_n - 2n^3$ to find a constant difference of -1 **[M1]**

$2n^3 - 1$ **[A1]**

(b) (i) $V_1 = 1$, $V_2 = \dfrac{5}{4}$, $V_3 = \dfrac{5}{4}$ **[B2]**

Award B1 if any two are correct **[B1]**

(ii) Substitutes $n = 5$ to find $V_5 = \dfrac{25 + 1}{32} = \dfrac{13}{16}$ **[B1]**

Substitutes $n = 4$ to find $V_4 = \dfrac{16+1}{16} = \dfrac{17}{16}$ **[B1]**

Evaluates $\dfrac{13}{16} - \dfrac{17}{16} = -\dfrac{1}{4}$ (accept -0.25) **[A1]**

[Back to Question] *[Total Marks: 9]*