IGCSE 0607 Practice Questions — E3.4 — Combined Questions & Markscheme

Finding a quadratic function using given information

Total Questions: 39 — **Total Marks:** 210 Marks

Table of Contents & Quick Navigation

- **E3.4** Finding a quadratic function using given information
  → **Paper 2 (Non-Calculator)** *(21 Qs)* · **Paper 4 (Calculator)** *(18 Qs)*

---

1 A quadratic function has x-intercepts at $x = -3$ and $x = 4$.
The coefficient of x^2 is 1.

Write down the equation of this function in the form $y = ax^2 + bx + c$.

$y =$ ..... **[2]**

2 A quadratic function has x-intercepts at $x = 4$ and $x = -2$.
The coefficient of x^2 is 1.

Find the equation of the quadratic function in the form $y = x^2 + bx + c$.

$y =$ ..... **[2]**

3 A quadratic function $f(x)$ has x-intercepts at $x = -4$ and $x = 3$.
The coefficient of x^2 is 1.

Find the equation of the function in the form $f(x) = x^2 + bx + c$.

$f(x) =$ ..... **[2]**

4 A quadratic function has x-intercepts at $x = -4$ and $x = 7$.
The coefficient of x^2 is 1.
Write down the equation of the function.
Give your answer in the form $y = x^2 + bx + c$.

$y =$ ..... **[2]**

5 A quadratic function has the equation $y = x^2 + bx + c$.
The vertex of the graph of this function is $(4, -3)$.

Find the value of b and the value of c.

$b =$ ..... , $c =$ ..... **[3]**

6 A quadratic function is given by

$$y = x^2 + bx + c$$

The vertex of the graph of this function is $(3, -4)$.

Find the value of b and the value of c.

$b =$ ..... , $c =$ ..... **[3]**

7 A quadratic function has a minimum point at $(3, -4)$ and passes through the point $(5, 0)$.
The coefficient of x^2 is 1.

Find the equation of the function in the form $y = x^2 + bx + c$.

$y =$ ..... **[3]**

8 The graph of a quadratic function, $y = x^2 + bx + c$, crosses the x-axis at $(-4, 0)$ and $(2, 0)$.

(a) Work out the value of b and the value of c.

$b =$ ..... , $c =$ ..... **[2]**

(b) Find the coordinates of the vertex of the graph.

(..... , ...) **[2]**

9 The graph of a quadratic function $f(x) = x^2 + bx + c$ crosses the x-axis at $(-3, 0)$ and $(5, 0)$.

(a) Find the values of b and c.

$b =$ ..... , $c =$ ..... **[3]**

(b) Write down the coordinates of the y-intercept of the graph of $y = f(x)$.

(..... , ...) **[1]**

10 A quadratic function $y = f(x)$ has its vertex at $(3, -4)$.
The coefficient of x^2 is 1.

(a) Find the equation of the function in the form $y = x^2 + bx + c$.

$y =$ ..... **[2]**

(b) Find the coordinates of the points where the graph of $y = f(x)$ crosses the x-axis.

..... **[2]**

11 A quadratic function is given by the equation $y = (x - h)^2 + k$.
The vertex of the graph of this function is $(3, -2)$.


(a) Write down the value of h and the value of k.

$h =$ ..... , $k =$ ..... **[2]**

(b) The graph passes through the point $(5, p)$.
Work out the value of p.

$p =$ ..... **[2]**

12 The diagram shows a sketch of the quadratic curve $y = x^2 + bx + c$.
The vertex of the curve is at $(3, -4)$.



(a) Write down the equation of the curve in the form $y = (x - h)^2 + k$.

$y =$ ..... **[2]**


(b) Find the values of b and c.

$b =$ ..... , $c =$ ..... **[2]**

(c) Write down the coordinates of the point where the curve crosses the y-axis.

(..... , ...) **[1]**

13 The diagram shows the graph of a quadratic function $y = x^2 + px + q$.
The graph has a minimum point (vertex) at $M(2, -9)$.



(a) Find the value of p and the value of q.

$p =$ ..... , $q =$ ..... **[3]**

(b) The graph crosses the x-axis at the points A and B.
Find the coordinates of A and the coordinates of B.


$A =$ ..... , $B =$ ..... **[3]**

14 The graph of a quadratic function $y = f(x)$ intersects the x-axis at $(-2, 0)$ and $(5, 0)$.
The graph intersects the y-axis at $(0, -20)$.

Work out the function $f(x)$, giving your answer in the form $ax^2 + bx + c$.

$f(x) =$ ..... **[4]**


15 The diagram shows a sketch of the quadratic graph $y = f(x)$.
The graph crosses the x-axis at $(-2, 0)$ and $(6, 0)$.
It crosses the y-axis at $(0, -24)$.



Find the expression for $f(x)$ in the form $ax^2 + bx + c$.

$f(x) =$ ..... **[4]**

16 The diagram shows the graph of a quadratic function $g(x)$.
The vertex of the parabola is at $(2, -9)$ and the curve passes through the point $(5, 9)$.



(a) Find the equation of the quadratic function in the form $g(x) = a(x - h)^2 + k$.

$g(x) =$ ..... **[3]**


(b) Express $g(x)$ in the form $ax^2 + bx + c$.

$g(x) =$ ..... **[2]**

(c) Write down the equation of the line of symmetry of the graph of $y = g(x)$.

..... **[1]**

17 The diagram shows a sketch of the quadratic function $y = f(x)$.
The graph crosses the x-axis at $(-1, 0)$ and $(4, 0)$.
The graph crosses the y-axis at $(0, -8)$.




(a) Find the equation of the quadratic function in the form $y = ax^2 + bx + c$.

$y =$ ..... **[3]**

(b) Find the coordinates of the minimum point of the graph.

(..... , ...) **[3]**

18 The diagram shows the graph of a quadratic function $y = ax^2 + bx + c$.
The curve crosses the x-axis at $(-2, 0)$ and $(4, 0)$, and crosses the y-axis at $(0, 16)$.



(a) Find the equation of the curve in the form $y = ax^2 + bx + c$.

$y =$ ..... **[4]**

(b) Find the coordinates of the turning point of the curve.

(..... , ...) **[2]**

(c) Write down the equation of the line of symmetry of the curve.

..... **[1]**

19 A quadratic function is given by $f(x) = a(x-h)^2 + k$.
The graph of $y = f(x)$ has a maximum point at $(-1, 8)$ and passes through the point $(2, -10)$.

(a) Write down the value of h and the value of k.

$h =$ ..... , $k =$ ..... **[2]**


(b) Find the value of a.

$a =$ ..... **[3]**

(c) Write $f(x)$ in the form $ax^2 + bx + c$.

$f(x) =$ ..... **[3]**


20 The diagram shows the graph of a quadratic function $y = f(x)$.
The graph crosses the x-axis at $x = -1$ and $x = 4$.
The graph crosses the y-axis at $(0, -8)$.



Find the equation of the quadratic function in the form $f(x) = ax^2 + bx + c$.

$f(x) =$ ..... **[4]**

21 The diagram shows a sketch of the curve $y = f(x)$ where $f(x)$ is a quadratic function.
The maximum point on the curve is $M(-2, 8)$ and the curve crosses the y-axis at $(0, -4)$.



(a) Find the equation of the curve.
Give your answer in the form $f(x) = ax^2 + bx + c$.

$f(x) =$ ..... **[4]**

(b) Solve $f(x) = 0$ to find the exact x-coordinates of the points where the curve crosses the x-axis.

Give your answers in the form $\frac{p \pm q\sqrt{6}}{r}$, where p, q and r are integers.

$x =$ ..... **[3]**

22 A quadratic curve has its vertex at $(3, -5)$ and passes through the point $(1, 7)$.
Find the equation of the curve in the form $y = a(x - h)^2 + k$.

$y =$ ..... **[3]**

23 The graph of a quadratic function has its vertex at $(-2, 5)$ and passes through the point $(1, -13)$. Find the equation of the function in the form $y = a(x - h)^2 + k$.

$y =$ ..... **[3]**

24 The graph of a quadratic function has its vertex at $(2, 7)$ and passes through the point $(4, -5)$.

(a) Find the equation of the function in the form $y = a(x - h)^2 + k$.

$y =$ ..... **[3]**

(b) Find the y-intercept of the graph.

..... **[1]**

25 A quadratic curve crosses the x-axis at $(-1, 0)$ and $(5, 0)$.

The curve also passes through the point $(2, -18)$.

Find the equation of the curve in the form $y = ax^2 + bx + c$.


$y =$ ..... **[4]**

26 A quadratic curve has a vertex at the point $(2, -5)$ and passes through the point $(4, 7)$.

Find the equation of the curve.
Give your answer in the form $y = ax^2 + bx + c$.

$y =$ ..... **[4]**

27 The diagram shows a sketch of the graph of a quadratic function $y = a(x - h)^2 + k$.
The vertex of the graph is $V(2, 5)$.
The graph passes through the point $P(4, -7)$.



(a) Write down the value of h and the value of k.

$h =$ ..... , $k =$ ..... **[2]**

(b) Work out the value of a.

$a =$ ..... **[2]**

(c) Find the coordinates of the y-intercept of the graph.

(..... , ...) **[2]**

28 The graph of a quadratic function $f(x)$ intersects the x-axis at $x = -2$ and $x = 5$.
The graph passes through the point $(3, 10)$.


(a) Find an expression for $f(x)$ in the form $ax^2 + bx + c$.

$f(x) =$ ..... **[4]**

(b) Find the coordinates of the vertex of the graph of $y = f(x)$.

(..... , ...) **[2]**


29 The diagram shows a sketch of the graph of $y = ax^2 + bx + c$.
The vertex of the graph is $V(-1, 8)$ and it passes through the point $P(2, -10)$.



Work out the values of a, b and c.

$a =$ ..... , $b =$ ..... , $c =$ ..... **[5]**

30 The diagram shows the sketch of a quadratic function.
The curve crosses the x-axis at -3 and 2, and intersects the y-axis at 12.




(a) Find the equation of the quadratic function.
Give your answer in the form $y = ax^2 + bx + c$.

$y =$ ..... **[4]**

(b) Calculate the coordinates of the vertex of this curve.

((..... , ...)) **[2]**

31 The diagram shows a sketch of the quadratic function $y = f(x)$.
The curve crosses the x-axis at $x = -1.5$ and $x = 4$.
It passes through the point $(2, -14)$.



(a) Find the equation of the quadratic function in the form $f(x) = ax^2 + bx + c$.

$f(x) =$ ..... **[4]**

(b) Find the exact coordinates of the turning point of the graph.

(..... , ...) **[3]**

32 A quadratic function is given by $f(x) = a(x - h)^2 + k$.
The maximum point of the graph of $y = f(x)$ is $(-4, 20)$.
The graph intersects the x-axis at $x = -2$ and $x = p$.

(a) **(i)** Write down the value of h and the value of k.

$h =$ ..... , $k =$ ..... **[2]**

(ii) Find the value of a.

$a =$ ..... **[2]**

(iii) Find the value of p.

$p =$ ..... **[1]**

(iv) Express $f(x)$ in the form $ax^2 + bx + c$.

$f(x) =$ ..... **[2]**

33 A quadratic function $h(x) = ax^2 + bx + c$ has x-intercepts at $(-2, 0)$ and $(4, 0)$, and passes through the point $(1, 13.5)$.

(a) Find the values of a, b, and c.

$a =$ ..... , $b =$ ..... , $c =$ ..... **[4]**


(b) Hence find the coordinates of the vertex of the graph of $y = h(x)$.

(..... , ...) **[2]**

(c) State the range of $h(x)$.

..... **[1]**

34 The graph of a quadratic function $y = f(x)$ intersects the x-axis at the points $A(-1, 0)$ and $B(5, 0)$. The graph intersects the y-axis at the point $C(0, 10)$.



(a) Find the equation of the function $f(x)$.
Give your answer in the form $f(x) = ax^2 + bx + c$.

$f(x) =$ ..... **[4]**


(b) Find the coordinates of the turning point (vertex) of the graph.

(..... , ...) **[2]**

(c) The graph of $y = f(x)$ is translated by the vector $\begin{pmatrix} 2 \\ -3 \end{pmatrix}$ to give the graph of $y = g(x)$.
Find the equation of $g(x)$ in the form $g(x) = a(x - h)^2 + k$.

$g(x) =$ ..... **[2]**

35 The diagram shows the graph of a quadratic function $y = f(x)$.
The vertex of the graph is at $V(-2, 8)$ and it intersects the y-axis at $(0, 5)$.



(a) Find the equation of the function $f(x)$ in the form $f(x) = a(x - h)^2 + k$.

$f(x) =$ ..... **[3]**


(b) Write $f(x)$ in the form $ax^2 + bx + c$.

$f(x) =$ ..... **[2]**

(c) Calculate the exact x-coordinates of the points where the graph intersects the x-axis.
Give your answers correct to 3 significant figures.

$x =$ ..... , $x =$ ..... **[3]**

36 The diagram shows a sketch of the quadratic curve $y = ax^2 + bx + c$.
The curve crosses the x-axis at $A(-2, 0)$ and $B(5, 0)$, and crosses the y-axis at $C(0, 15)$.



(a) Find the values of a, b, and c.

$a =$ ..... , $b =$ ..... , $c =$ ..... **[5]**


(b) Find the coordinates of the vertex of the curve.

(..... , ...) **[3]**

(c) The equation $ax^2 + bx + c = k$ has no real roots.
Write down the inequality for k.

..... **[1]**

37 The diagram shows a sketch of the graph of a quadratic function $y = f(x)$.
The maximum point of the graph is $V(-2, 18)$.
The graph crosses the y-axis at $(0, 10)$.



(a) Find the equation of the quadratic function in the form $y = ax^2 + bx + c$.

$y =$ ..... **[3]**

(b) The graph crosses the x-axis at A and B.
Find the length of the line segment AB.

$AB =$ ..... **[3]**

(c) The line $y = -2x + k$ is a tangent to the curve $y = f(x)$.

Find the value of k.

$k =$ ..... **[4]**

38 A quadratic function $f(x) = a(x-h)^2 + k$ has its vertex at $(4, -5)$ and passes through the point $(1, 8.5)$.

(a) Find the values of a, h, and k.

$a =$ ..... , $h =$ ..... , $k =$ ..... **[3]**

(b) Solve the equation $f(x) = 0$, giving your answers correct to 2 decimal places.

$x =$ ..... , $x =$ ..... **[3]**

(c) The line $y = 2x - 7$ intersects the curve $y = f(x)$ at two points, P and Q.
Find the x-coordinates of P and Q, giving your answers correct to 2 decimal places.

$x =$ ..... , $x =$ ..... **[3]**

(d) The graph of $y = f(x)$ is translated by the vector $\begin{pmatrix} -3 \\ 9 \end{pmatrix}$ to give the graph of $y = g(x)$.
Write down the expression for $g(x)$ in the form $p(x-q)^2 + r$.

$g(x) =$ ..... **[2]**

39 The quadratic function $f(x) = ax^2 + bx + c$ has a minimum turning point at $(1, -8)$.
The graph of $y = f(x)$ crosses the x-axis at $x = -1$ and $x = 3$.

(a) Show that $f(x) = 2x^2 - 4x - 6$.

[3]

(b) Solve the equation.

$f(x) = 10$

$x =$ ..... , $x =$ ..... **[3]**

(c) A straight line has equation $y = mx - 8$.
The line is a tangent to the graph of $y = f(x)$.

Find the possible values of m.

$m =$ ..... , $m =$ ..... **[4]**

(d) The graph of $y = f(x)$ is translated by the vector $\begin{pmatrix} 0 \\ k \end{pmatrix}$.

The new graph has no x-intercepts.

Write down the inequality for k.

..... **[2]**

1 $y = (x+3)(x-4)$ **[M1]**

$x^2 - x - 12$ **[A1]**

[Back to Question] *[Total Marks: 2]*

2 $(x-4)(x+2)$ **[M1]**

$x^2 - 2x - 8$ **[A1]**

[Back to Question] *[Total Marks: 2]*

3 Uses $(x-p)(x-q)$ with $p = -4$ and $q = 3$: $(x+4)(x-3)$ **[M1]**

$x^2 + x - 12$ **[A1]**

[Back to Question] *[Total Marks: 2]*

4 Uses the intercepts to form $y = (x+4)(x-7)$ **[M1]**

$x^2 - 3x - 28$ **[A1]**

[Back to Question] *[Total Marks: 2]*

5 Uses the vertex form $y = (x-4)^2 - 3$ **[M1]**

Expands to $x^2 - 8x + 16 - 3$ **[M1]**

$b = -8$ and $c = 13$ **[A1]**

[Back to Question] *[Total Marks: 3]*

6 Uses the vertex form $y = (x-3)^2 - 4$ OR uses $-\dfrac{b}{2(1)} = 3$ **[M1]**

$b = -6$ **[A1]**

$c = 5$ **[A1]**

[Back to Question] *[Total Marks: 3]*

7 $y = (x-3)^2 - 4$ **[M1]**

$y = x^2 - 6x + 9 - 4$ **[M1]**

$x^2 - 6x + 5$ **[A1]**

[Back to Question] *[Total Marks: 3]*

8 (a) $x^2 + bx + c = (x+4)(x-2)$ **[M1]**

$b = 2$, $c = -8$ **[A1]**

(b) Finds x-coordinate of vertex: $x = \dfrac{-4+2}{2} = -1$ **[M1]**

$(-1, -9)$ **[A1]**

[Back to Question] *[Total Marks: 4]*

9 **(a)** Writes $f(x) = (x+3)(x-5)$ **[M1]**

Expands correctly: $x^2 - 2x - 15$ **[M1]**

$b = -2$, $c = -15$ **[A1]**

(b) $(0, -15)$ **[B1]**

[Back to Question] *[Total Marks: 4]*

10 **(a)** Uses vertex form: $y = (x-3)^2 - 4$ **[M1]**

$x^2 - 6x + 5$ **[A1]**

(b) Sets $x^2 - 6x + 5 = 0 \implies (x-1)(x-5) = 0$ **[M1]**

$(1, 0)$ and $(5, 0)$ **[A1]**

[Back to Question] *[Total Marks: 4]*

11 **(a)** $h = 3$ **[B1]**

$k = -2$ **[B1]**

(b) $p = (5-3)^2 - 2$ **[M1]**

$p = 2$ **[A1]**

[Back to Question] *[Total Marks: 4]*

12 **(a)** $(x-3)^2$ seen **[B1]**

$(x-3)^2 - 4$ **[B1]**

(b) $y = x^2 - 6x + 9 - 4$ **[M1]**

$b = -6$ and $c = 5$ **[A1]**

(c) $(0, 5)$ **[B1]**

[Back to Question] *[Total Marks: 5]*

13 **(a)** Uses vertex form: $y = (x-2)^2 - 9$ or x-coordinate of vertex $\dfrac{-p}{2} = 2$ **[M1]**

Expands to find p and q, or solves $2^2 + 2p + q = -9$ **[M1]**

$p = -4$ and $q = -5$ **[A1]**

(b) $x^2 - 4x - 5 = 0$ **[M1]**

$(x-5)(x+1) = 0$ **[M1]**

$A = (-1, 0)$ and $B = (5, 0)$ **[A1]**

[Back to Question] *[Total Marks: 6]*

14 $f(x) = a(x + 2)(x - 5)$ **[M1]**

$a(0 + 2)(0 - 5) = -20$ or $-10a = -20$ **[M1]**

$a = 2$ **[A1]**

$2x^2 - 6x - 20$ **[A1]**

[Back to Question] *[Total Marks: 4]*

15 $f(x) = a(x + 2)(x - 6)$ **[M1]**

$-24 = a(0 + 2)(0 - 6)$ **[M1]**

$a = 2$ **[A1]**

$2x^2 - 8x - 24$ **[A1]**

[Back to Question] *[Total Marks: 4]*

16 (a) Writes $g(x) = a(x - 2)^2 - 9$ **[M1]**

Substitutes $(5, 9)$: $9 = a(5 - 2)^2 - 9$ or $9a = 18$ **[M1]**

$2(x - 2)^2 - 9$ **[A1]**

(b) $2(x^2 - 4x + 4) - 9$ **[M1]**

$2x^2 - 8x - 1$ **[A1]**

(c) $x = 2$ **[B1]**

[Back to Question] *[Total Marks: 6]*

17 (a) Uses intercept form: $y = a(x + 1)(x - 4)$ **[M1]**

Substitutes $(0, -8)$ to find a: $-8 = a(0 + 1)(0 - 4) \implies -4a = -8 \implies a = 2$ **[M1]**

$2x^2 - 6x - 8$ **[A1]**

(b) Finds the x-coordinate of the vertex by symmetry: $x = \dfrac{-1 + 4}{2} = 1.5$ (or uses $x = -\dfrac{b}{2a}$) **[M1]**

Substitutes $x = 1.5$ into the equation: $y = 2(1.5)^2 - 6(1.5) - 8$ **[M1]**

$(1.5, -12.5)$ (or $(\dfrac{3}{2}, -\dfrac{25}{2})$) **[A1]**

[Back to Question] *[Total Marks: 6]*

18 (a) $y = a(x + 2)(x - 4)$ **[M1]**

$16 = a(0 + 2)(0 - 4)$ **[M1]**

$a = -2$ **[A1]**

$y = -2x^2 + 4x + 16$ **[A1]**

(b) $x = \dfrac{-2+4}{2} = 1$ **[M1]**

$(1, 18)$ **[A1]**

(c) $x = 1$ **[B1]**

[Back to Question] *[Total Marks: 7]*

19 **(a)** $h = -1$ **[B1]**

$k = 8$ **[B1]**

(b) $-10 = a(2 - (-1))^2 + 8$ **[M1]**

$9a = -18$ **[M1]**

-2 **[A1]**

(c) $-2(x+1)^2 + 8$ **[M1]**

$-2(x^2 + 2x + 1) + 8$ **[M1]**

$-2x^2 - 4x + 6$ **[A1]**

[Back to Question] *[Total Marks: 8]*

20 $f(x) = a(x+1)(x-4)$ **[M1]**

$-8 = a(0+1)(0-4)$ **[M1]**

$a = 2$ **[A1]**

$2x^2 - 6x - 8$ **[A1]**

[Back to Question] *[Total Marks: 4]*

21 **(a)** Forms the equation $f(x) = a(x+2)^2 + 8$ **[M1]**

Substitutes $(0, -4)$ into their equation:

$-4 = a(0+2)^2 + 8$ **[M1]**

$4a = -12 \implies a = -3$ **[A1]**

$-3x^2 - 12x - 4$ **[A1]**

(b) Sets $-3(x+2)^2 + 8 = 0$ or $-3x^2 - 12x - 4 = 0$ **[M1]**

Rearranges to $(x+2)^2 = \dfrac{8}{3}$ OR correctly substitutes into the quadratic formula:

$x = \dfrac{-(-12) \pm \sqrt{(-12)^2 - 4(-3)(-4)}}{2(-3)}$ **[M1]**

$\dfrac{-6 \pm 2\sqrt{6}}{3}$ **[A1]**

[Back to Question] *[Total Marks: 7]*

22 $y = a(x-3)^2 - 5$ **[M1]**

$7 = a(1-3)^2 - 5 \implies 12 = 4a \implies a = 3$ **[M1]**

$3(x-3)^2 - 5$ **[A1]**

[Back to Question] *[Total Marks: 3]*

23 $y = a(x+2)^2 + 5$ **[M1]**

$-13 = a(1+2)^2 + 5$ **[M1]**

$-18 = 9a$

a = -2$[M1]$

$-2(x+2)^2 + 5$ **[A1]**

[Back to Question] *[Total Marks: 3]*

24 **(a)** Uses vertex form: $y = a(x-2)^2 + 7$ **[M1]**

Substitutes $(4, -5)$ into their equation: $-5 = a(4-2)^2 + 7$ **[M1]**

$-3(x-2)^2 + 7$ **[A1]**

(b) -5 **[B1]**

[Back to Question] *[Total Marks: 4]*

25 Writes the function in the factored form $y = a(x+1)(x-5)$ **[M1]**

Substitutes $(2, -18)$ into their equation:

$-18 = a(2+1)(2-5)$ **[M1]**

$a = 2$ **[A1]**

$2x^2 - 8x - 10$ **[A1]**

[Back to Question] *[Total Marks: 4]*

26 Uses vertex form $y = a(x-2)^2 - 5$ **[M1]**

Substitutes $(4, 7)$ into their equation: $7 = a(4-2)^2 - 5$ **[M1]**

$4a = 12$

a = 3$[A1]$

$3x^2 - 12x + 7$ **[A1]**

[Back to Question] *[Total Marks: 4]*

27 **(a)** $h = 2$ **[B1]**

$k = 5$ **[B1]**

(b) Substitutes $(4, -7)$ into equation: $-7 = a(4-2)^2 + 5$ **[M1]**

$a = -3$ **[A1]**

(c) Substitutes $x = 0$ into $y = -3(x-2)^2 + 5$ **[M1]**

$(0, -7)$ **[A1]**

[Back to Question] *[Total Marks: 6]*

28 **(a)** Sets up the factored form $f(x) = a(x+2)(x-5)$ **[M1]**

Substitutes the point $(3, 10)$: $10 = a(3+2)(3-5)$ **[M1]**

Solves for a: $10 = a(5)(-2) \implies a = -1$ **[A1]**

$f(x) = -x^2 + 3x + 10$ **[A1]**

(b) Finds the x-coordinate of the vertex (e.g., midpoint of intercepts $\frac{-2+5}{2} = 1.5$ or $x = \frac{-b}{2a}$) **[M1]**

$(1.5, 12.25)$ **[A1]**

[Back to Question] *[Total Marks: 6]*

29 $y = a(x+1)^2 + 8$ **[M1]**

$-10 = a(2+1)^2 + 8$ **[M1]**

$9a = -18 \implies a = -2$ **[A1]**

$-2(x^2 + 2x + 1) + 8$ **[M1]**

$a = -2$, $b = -4$, $c = 6$ **[A1]**

[Back to Question] *[Total Marks: 5]*

30 **(a)** $y = a(x+3)(x-2)$ **[M1]**

$12 = a(0+3)(0-2)$ **[M1]**

$12 = -6a$

a = -2[$M1$]

$-2x^2 - 2x + 12$ **[A1]**

(b) $x = \frac{-3+2}{2}$ or $x = \frac{-(-2)}{2(-2)} = -0.5$ **[M1]**

$y = -2(-0.5)^2 - 2(-0.5) + 12 = 12.5$ **[M1]**

$(-0.5, 12.5)$ **[A1]**

[Back to Question] *[Total Marks: 6]*

31 **(a)** Sets up equation using x-intercepts: $f(x) = a(x+1.5)(x-4)$ OR $f(x) = a(2x+3)(x-4)$ **[M1]**

Substitutes $(2, -14)$ to find a: $-14 = a(2+1.5)(2-4)$ **[M1]**

Solves for a: $a = 2$ **[A1]**

$2x^2 - 5x - 12$ **[A1]**

(b) Identifies x-coordinate of vertex as midpoint of roots: $x = \dfrac{-1.5 + 4}{2} = 1.25$ OR uses $x = \dfrac{-b}{2a}$ **[M1]**

Substitutes $x = 1.25$ into their function: $y = 2(1.25)^2 - 5(1.25) - 12$ **[M1]**

$(1.25, -15.125)$ OR $\left(\dfrac{5}{4}, -\dfrac{121}{8}\right)$ **[A1]**

[Back to Question] *[Total Marks: 7]*

32 **(a)** **(i)** $h = -4$ **[B1]**

$k = 20$ **[B1]**

(ii) $0 = a(-2 + 4)^2 + 20$ **[M1]**

$a = -5$ **[A1]**

(iii) -6 **[B1]**

(iv) $f(x) = -5(x + 4)^2 + 20 = -5(x^2 + 8x + 16) + 20$ **[M1]**

$-5x^2 - 40x - 60$ **[A1]**

[Back to Question] *[Total Marks: 7]*

33 **(a)** Writes $h(x) = a(x + 2)(x - 4)$ **[M1]**

Substitutes $(1, 13.5)$: $13.5 = a(3)(-3) \implies -9a = 13.5$ **[M1]**

$a = -1.5$ (or $-\dfrac{3}{2}$) **[A1]**

Expands $-1.5(x^2 - 2x - 8)$ to obtain $b = 3$ and $c = 12$ **[A1]**

(b) x-coordinate of vertex $= \dfrac{-2 + 4}{2} = 1$ **[M1]**

$(1, 13.5)$ **[A1]**

(c) $h(x) \leq 13.5$ or $y \leq 13.5$ **[B1]**

[Back to Question] *[Total Marks: 7]*

34 **(a)** Sets up the equation in factored form: $f(x) = a(x + 1)(x - 5)$ **[M1]**

Substitutes $(0, 10)$ to find a: $10 = a(0 + 1)(0 - 5)$ **[M1]**

$a = -2$ **[A1]**

$-2x^2 + 8x + 10$ **[A1]**

(b) Finds the x-coordinate of the vertex using symmetry $x = \dfrac{-1 + 5}{2} = 2$ OR uses $x = -\dfrac{b}{2a}$ **[M1]**

$(2, 18)$ **[A1]**

(c) Translates the vertex $(2, 18)$

(4, 15)$ORappliestransformation$-2((x-2)-2)$^2 + 18 - 3$ [M1]

$-2(x-4)^2 + 15$ **[A1]**

[Back to Question] *[Total Marks: 8]*

35 **(a)** Uses the vertex coordinates to write $f(x) = a(x+2)^2 + 8$ **[M1]**

Substitutes the y-intercept $(0, 5)$: $5 = a(0+2)^2 + 8$ **[M1]**

$f(x) = -0.75(x+2)^2 + 8$ or equivalent using fractions like $-\frac{3}{4}$ **[A1]**

(b) Expands $(x+2)^2$ as $x^2 + 4x + 4$ **[M1]**

$f(x) = -0.75x^2 - 3x + 5$ **[A1]**

(c) Equates function to zero: $-0.75(x+2)^2 + 8 = 0$ or $-0.75x^2 - 3x + 5 = 0$ **[M1]**

Rearranges to $(x+2)^2 = \frac{32}{3}$ or correctly applies the quadratic formula **[M1]**

$x = 1.27$ and $x = -5.27$ **[A1]**

[Back to Question] *[Total Marks: 8]*

36 **(a)** $c = 15$ **[B1]**

$y = a(x+2)(x-5)$ **[M1]**

$15 = a(0+2)(0-5)$ **[M1]**

$a = -1.5$ **[A1]**

$b = 4.5$ **[A1]**

(b) $x = \frac{-2+5}{2} = 1.5$ **[M1]**

$y = -1.5(1.5)^2 + 4.5(1.5) + 15$ **[M1]**

$(1.5, 18.375)$ or $\left(\frac{3}{2}, \frac{147}{8}\right)$ **[A1]**

(c) $k > 18.375$ **[B1]**

[Back to Question] *[Total Marks: 9]*

37 **(a)** Uses vertex form: $y = a(x+2)^2 + 18$ **[M1]**

Substitutes $(0, 10)$ to find a: $10 = a(0+2)^2 + 18 \implies 4a = -8 \implies a = -2$ **[M1]**

$-2x^2 - 8x + 10$ **[A1]**

(b) Sets $f(x) = 0$: $-2x^2 - 8x + 10 = 0$ **[M1]**

$x^2 + 4x - 5 = 0 \implies (x+5)(x-1) = 0 \implies x = -5$ or $x = 1$ **[M1]**

6 **[A1]**

(c) Equates line and curve: $-2x^2 - 8x + 10 = -2x + k$ **[M1]**

Rearranges to form a quadratic equation: $2x^2 + 6x + (k-10) = 0$ (or equivalent) **[M1]**

Sets discriminant $\Delta = 0$ for a tangent: $6^2 - 4(2)(k-10) = 0$ **[M1]**

14.5 (or $\frac{29}{2}$) **[A1]**

[Back to Question] *[Total Marks: 10]*

38 **(a)** $h = 4$ and $k = -5$ **[B1]**

$8.5 = a(1-4)^2 - 5$ **[M1]**

$a = 1.5$ **[A1]**

(b) $1.5(x-4)^2 - 5 = 0$ or $(x-4)^2 = \frac{10}{3}$ **[M1]**

$x = 4 \pm \sqrt{\frac{10}{3}}$ **[M1]**

$x = 2.17$ and $x = 5.83$ **[A1]**

(c) $1.5(x-4)^2 - 5 = 2x - 7$ **[M1]**

$3x^2 - 28x + 52 = 0$ **[A1]**

$x = 2.56$ and $x = 6.77$ **[A1]**

(d) New vertex at $(1, 4)$ **[M1]**

$1.5(x-1)^2 + 4$ **[A1]**

[Back to Question] *[Total Marks: 11]*

39 **(a)** Forms equation using x-intercepts: $f(x) = a(x+1)(x-3)$ OR using vertex $f(x) = a(x-1)^2 - 8$ **[M1]**

Substitutes appropriate point to find a: $a(1+1)(1-3) = -8 \implies -4a = -8 \implies a = 2$ **[M1]**

Expands to show final form: $2(x^2 - 2x - 3) = 2x^2 - 4x - 6$ **[A1]**

(b) $2x^2 - 4x - 6 = 10 \implies 2x^2 - 4x - 16 = 0$ **[M1]**

Factors correctly to $(x-4)(x+2) = 0$ OR uses quadratic formula correctly **[M1]**

$x = 4$ and $x = -2$ **[A1]**

(c) Sets equations equal: $2x^2 - 4x - 6 = mx - 8$ leading to $2x^2 - (4+m)x + 2 = 0$ **[M1]**

States condition for tangent (discriminant $= 0$): $(-(4+m))^2 - 4(2)(2) = 0$ **[M1]**

Simplifies to $(4+m)^2 = 16$ or $m^2 + 8m = 0$ **[M1]**

$m = 0$, $m = -8$ **[A1]**

(d) Recognises minimum y-value must be strictly greater than 0: $-8 + k > 0$ **[M1]**

$k > 8$ **[A1]**

[Back to Question] *[Total Marks: 12]*