IGCSE 0607 Practice Questions — E2.4 — Combined Questions & Markscheme

Indices II

Total Questions: 9 — **Total Marks:** 50 Marks

Table of Contents & Quick Navigation

- **E2.4** Indices II
  → **Paper 2 (Non-Calculator)** *(8 Qs)* · **Paper 4 (Calculator)** *(1 Q)*

---

1 **(a)** Simplify.

$$4x^2 \times 3x^{-5}$$

..... **[1]**

(b) Simplify.

$$\left(\frac{8x^9}{27}\right)^{\frac{1}{3}}$$

..... **[2]**

2 **(a)** Simplify.

$w^4 \times w^7$

..... **[1]**

(b) Simplify.

$(3x^3)^2$

..... **[2]**

(c) Simplify.

$$\frac{15y^8}{5y^{-2}}$$

..... **[2]**

3 **(a)** Simplify.

$p^4 \times p^7$

..... **[1]**

(b) Simplify.

$(3q^3)^2$

..... **[2]**

(c) Work out the value of $27^{\frac{2}{3}}$.

..... **[2]**

4 **(a)** Work out the exact value of

$$32^{-\frac{2}{5}}$$

..... **[2]**

(b) Solve the equation.

$$5^{x+2} = 125^{x}$$

$x =$ ..... **[3]**

5 **(a)** Simplify fully.

$$4x^{-3} \times \frac{3}{2}x^{\frac{1}{2}}$$

..... **[2]**

(b) Solve the equation.

$$3^{2x+1} = 27$$

$x =$ ..... **[2]**

(c) Simplify fully.

$$\left(\frac{8a^6}{b^3}\right)^{\frac{1}{3}}$$

..... **[2]**

6 **(a)** Simplify fully.

$$\left(\frac{16x^{12}}{y^8}\right)^{\frac{3}{4}}$$

..... **[3]**

(b) Solve the equation.

$$2^{x+3} = 8^x$$

$x =$ ..... **[3]**

7 **(a)** Solve the equation.

$$16^{2x-1} = 8^{x+2}$$

$x =$ ..... **[3]**

(b) Simplify fully.

$$\left(\frac{81x^{12}}{16y^{8}}\right)^{-\frac{3}{4}}$$

..... **[3]**

8 **(a)** Simplify fully.

$$\frac{4x^{-\frac{1}{2}} \times 3x^2}{6x^{\frac{1}{3}}}$$

..... **[3]**

(b) Solve the equation.

$$9^{2x-1} \times 3^{x+1} = 27^x$$

$x =$ ..... **[4]**

9 **(a)** Simplify fully.

$$\frac{12x^{\frac{3}{2}}y^{-2}}{4x^{-\frac{1}{2}}y^{4}}$$

..... **[3]**

(b) Solve the equation.

$$4^{2x-1} \times 8^{x+1} = 32^{x}$$

$x =$ ..... **[4]**

1 **(a)** $12x^{-3}$ or $\frac{12}{x^3}$ **[B1]**

(b) $\frac{2x^3}{3}$ or $\frac{2}{3}x^3$ **[B2]**

Correct cube root of coefficient $\frac{2}{3}$ OR correct index x^3 seen **[M1]**

[Back to Question] *[Total Marks: 3]*

2 **(a)** w^{11} **[B1]**

(b) 3^2 or 9 seen, or x^6 seen **[M1]**

$9x^6$ **[A1]**

(c) $\frac{15}{5} = 3$ or $y^{8-(-2)}$ **[M1]**

$3y^{10}$ **[A1]**

[Back to Question] *[Total Marks: 5]*

3 **(a)** p^{11} **[B1]**

(b) $9q^6$ **[B2]**

Partial correct answer $9q^k$ or cq^6 **[B1]**

(c) $(\sqrt[3]{27})^2$ or 3^2 or $\sqrt[3]{729}$ **[M1]**

9 **[A1]**

[Back to Question] *[Total Marks: 5]*

4 **(a)** $\frac{1}{4}$ or 0.25 **[B2]**

Correct intermediate step e.g., 2^{-2} or $\left(\frac{1}{32}\right)^{\frac{2}{5}}$ or $32^{\frac{1}{5}} = 2$ **[M1]**

(b) Writes 125 as 5^3 **[M1]**

Equates indices: $x + 2 = 3x$ **[M1]**

1 **[A1]**

[Back to Question] *[Total Marks: 5]*

5 **(a)** $4 \times \frac{3}{2} = 6$ or $x^{-3+\frac{1}{2}}$ **[M1]**

$6x^{-\frac{5}{2}}$ or $\frac{6}{x^{\frac{5}{2}}}$ **[A1]**

(b) $3^{2x+1} = 3^3$ or $2x + 1 = 3$ **[M1]**

1 **[A1]**

(c) $8^{\frac{1}{3}} = 2$ or $(a^6)^{\frac{1}{3}} = a^2$ or $(b^3)^{\frac{1}{3}} = b$ **[M1]**

$\dfrac{2a^2}{b}$ **[A1]**

[Back to Question] *[Total Marks: 6]*

6 (a) 8 **[B1]**

x^9 **[B1]**

y^6 as denominator or y^{-6} **[B1]**

(b) $8 = 2^3$ seen or implied by $8^x = 2^{3x}$ **[M1]**

$x + 3 = 3x$ **[M1]**

1.5 or $\dfrac{3}{2}$ **[A1]**

[Back to Question] *[Total Marks: 6]*

7 (a) Expressing both sides with base 2: $(2^4)^{2x-1} = (2^3)^{x+2}$ **[M1]**

Equating powers: $4(2x - 1) = 3(x + 2)$ or $8x - 4 = 3x + 6$ **[M1]**

2 **[A1]**

(b) Taking the reciprocal to remove negative index: $\left(\dfrac{16y^8}{81x^{12}}\right)^{\frac{3}{4}}$ **[M1]**

Evaluating the fractional power of the numbers: $16^{\frac{3}{4}} = 8$ and $81^{\frac{3}{4}} = 27$ **[M1]**

$\dfrac{8y^6}{27x^9}$ **[A1]**

[Back to Question] *[Total Marks: 6]*

8 (a) $12x^{\frac{3}{2}}$ seen as numerator **[M1]**

$2x^k$ for any k or $cx^{\frac{7}{6}}$ for any c **[M1]**

$2x^{\frac{7}{6}}$ **[A1]**

(b) Converting all terms to base 3: $3^{2(2x-1)} \times 3^{x+1} = 3^{3x}$ **[M1]**

Applying exponent multiplication law: $3^{4x-2+x+1}$ or 3^{5x-1} **[M1]**

Equating indices: $5x - 1 = 3x$ **[M1]**

0.5 or $\dfrac{1}{2}$ **[A1]**

[Back to Question] *[Total Marks: 7]*

9 (a) $3x^2y^{-6}$ or $\dfrac{3x^2}{y^6}$ **[B3]**

Coefficient 3 seen **[M1]**

Correct subtraction of indices $x^{\frac{3}{2}-\left(-\frac{1}{2}\right)}$ leading to x^2 OR y^{-2-4} leading to y^{-6} **[M1]**

(b) Expresses all terms as powers of 2: $(2^2)^{2x-1} \times (2^3)^{x+1} = (2^5)^x$ **[M1]**

Correctly adds powers on the LHS: $2(2x-1)+3(x+1)$ or $4x-2+3x+3$ **[M1]**

Forms linear equation equating LHS and RHS powers: $7x+1=5x$ **[M1]**

$-\frac{1}{2}$ or -0.5 **[A1]**

[Back to Question] *[Total Marks: 7]*