1 **(a)** $x = 4$ **[B1]**

(b) $y = 3$ **[B1]**

[Total Marks: 2]

2 **(a)** $x = 2$ **[B1]**

(b) $y = 1$ **[B1]**

(c) $x = -2$ **[B1]**

[Total Marks: 3]

3 **(a)** Vertical asymptote: $x = -2$ **[B1]**

Horizontal asymptote: $y = 1$ **[B1]**

(b) Substitute $y = 0$ to find x-intercept: $0 = \dfrac{3}{x+2} + 1 \implies x = -5$ **[M1]**

x-axis at $(-5, 0)$ and y-axis at $(0, 2.5)$ **[A1]**

(c) Two correct branches of the hyperbola sketched **[B1]**

Graph approaches $x = -2$ and $y = 1$ correctly without touching or crossing them **[B1]**

Graph passes through $(-5, 0)$ and $(0, 2.5)$ approximately **[B1]**

[Total Marks: 7]

4 **(a)** $x = -2$ **[B1]**

(b) $y = -4$ **[B1]**

(c) Sets $x = 0$ to find the y-intercept: $y = \dfrac{3}{0+2} - 4$ **[M1]**

y-intercept is $(0, -2.5)$ **[A1]**

Sets $y = 0$ to find the x-intercept: $0 = \dfrac{3}{x+2} - 4$ **[M1]**

$4 = \dfrac{3}{x+2} \implies 4(x+2) = 3 \implies 4x + 8 = 3$ **[M1]**

x-intercept is $(-1.25, 0)$ **[A1]**

(d) $x = 90°$ **[B1]**

$x = 270°$ **[B1]**

[Total Marks: 9]

5 **(a)** $b = 2$ **[B1]**

$c = -1$ **[B1]**

(b) Substitutes $(5, 0)$ into the equation: $\dfrac{a}{5-2} - 1 = 0$ **[M1]**

$a = 3$ **[A1]**

(c) Sets up the equation $\frac{3}{x-2} - 1 = -x + 5$ or uses GDC to find intersection of $y = f(x)$ and $y = -x + 5$ **[M1]**

$x = 3$ and $x = 5$ **[A1]**

[Total Marks: 6]

6 (a) $q = 5$ **[B1]**

$r = -3$ **[B1]**

(b) Substitutes $x = 7, y = -2$ and their values of q and r into the equation: $-2 = \frac{p}{7-5} - 3$ **[M1]**

$-2 = \frac{p}{2} - 3 \implies \frac{p}{2} = 1$ **[M1]**

$p = 2$ **[A1]**

(c) Sets $y = \frac{2}{x-5} - 3$ and attempts to make x the subject, or swaps x and y **[M1]**

$y + 3 = \frac{2}{x-5} \implies x - 5 = \frac{2}{y+3}$ **[M1]**

$x = \frac{2}{y+3} + 5$ **[M1]**

$f^{-1}(x) = \frac{2}{x+3} + 5$ or equivalent **[A1]**

(d) $x = -3$ **[B1]**

$y = 5$ **[B1]**

(Strict follow through from their answers in part (a): $x = r, y = q$)

[Total Marks: 11]