IGCSE 0607 Practice Questions — E4.3 — Combined Questions & Markscheme

Length and midpoint


Total Questions: 9 — **Total Marks:** 60 Marks

Table of Contents & Quick Navigation

- **E4.3** Length and midpoint
  → **Paper 2 (Non-Calculator)** *(4 Qs)* · **Paper 4 (Calculator)** *(5 Qs)*

---

1 The points $A(-2, 5)$ and $B(4, -3)$ are shown on the coordinate grid.



(a) Find the coordinates of the midpoint of the line segment AB.

((..... , ...)) **[2]**

(b) Calculate the exact length of the line segment AB.

..... **[3]**

2 Point A has coordinates $(1, 4)$ and point B has coordinates $(7, -4)$.

(a) Calculate the coordinates of the midpoint of the line segment AB.

(..... , ...) **[2]**

(b) Calculate the length of the line segment AB.

..... **[3]**

3 A circle has its centre at the origin, $O(0, 0)$.
The point $P(-3, 4)$ lies on the circumference of this circle.

(a) Work out the radius of the circle.

..... **[2]**

(b) The line segment PQ is a diameter of the circle.
Find the coordinates of Q.

(..... , ...) **[2]**

(c) The point R has coordinates $(k, \sqrt{11})$.
R also lies on the circumference of the circle, and k is a negative number.
Work out the exact value of k.

$k =$ ..... **[3]**

4 A circle has diameter AB, where A is the point $(-1, 2)$ and B is the point $(5, 8)$.

(a) Find the coordinates of the centre of the circle, C.

(..... , ...) **[2]**

(b) Find the exact radius of the circle.
Give your answer in the form $a\sqrt{b}$, where a and b are positive integers and b is as small as possible.

..... **[3]**

(c) Point D is such that the midpoint of CD is $M\left(\frac{5}{2}, \frac{3}{2}\right)$.

Find the exact length of AD, giving your answer in the form $c\sqrt{d}$, where c and d are integers and d is as small as possible.

..... **[5]**

5 The points A and B have coordinates $(-2, 5)$ and $(4, -3)$ respectively.

(a) Calculate the exact length of the line segment AB.

..... **[3]**

(b) Find the coordinates of the midpoint of AB.

(..... , ...) **[2]**

6 Point P has coordinates $(3, k)$ and point Q has coordinates $(7, 5)$.
The length of the line segment PQ is 5 units.

(a) Find the two possible values of k.

$k =$ ..... , $k =$ ..... **[3]**

(b) M is the midpoint of PQ.
For the case where $k < 5$, find the coordinates of M.

((..... , ...)) **[2]**

7 Point P has coordinates $(-3, 5)$ and point Q has coordinates $(k, -1)$.
The length of the line segment PQ is 10 units.


(a) Find the two possible values of k.

$k =$ ..... , $k =$ ..... **[4]**

(b) Given that $k > 0$, point M is the midpoint of PQ.
Find the coordinates of M.

(..... , ...) **[2]**

8 The points $A(-3, 2)$ and $B(5, 8)$ are the endpoints of a diameter of a circle.



(a) Find the coordinates of the centre of the circle, C.

((..... , ...)) **[2]**

(b) Calculate the area of the circle.
Give your answer correct to 1 decimal place.

..... **[3]**

(c) The point $D(k, 8)$ lies on the circumference of the circle.
Find the possible values of k.

$k =$ ..... , $k =$ ..... **[3]**

9 The points $A(2p, p)$ and $B(3p, 4p)$ are joined by a straight line, where $p > 0$.

(a) The length of the line segment AB is $5\sqrt{10}$.
Find the value of p.

$p =$ ..... **[4]**

(b) Point C lies on the line segment AB such that the ratio $AC : CB = 2 : 3$.
Using your value of p from part (a), find the coordinates of C.

(..... , ...) **[3]**

(c) The point $D(x, y)$ is such that C is the midpoint of BD.
Work out the coordinates of D.

(..... , ...) **[2]**

1 **(a)** $\left(\frac{-2+4}{2}, \frac{5+(-3)}{2}\right)$ **[M1]**

$(1, 1)$ **[A1]**

(b) $(4-(-2))^2 + (-3-5)^2$ or $6^2 + (-8)^2$ **[M1]**

$\sqrt{36+64}$ or $\sqrt{100}$ **[M1]**

10 **[A1]**

[Back to Question] *[Total Marks: 5]*

2 **(a)** $\left(\frac{1+7}{2}, \frac{4+(-4)}{2}\right)$ **[M1]**

$(4, 0)$ **[A1]**

(b) $(7-1)^2 + (-4-4)^2$ oe **[M1]**

$\sqrt{36+64}$ or $\sqrt{100}$ **[M1]**

10 **[A1]**

[Back to Question] *[Total Marks: 5]*

3 **(a)** Calculates length OP: $\sqrt{(-3)^2+4^2}$ or $\sqrt{9+16}$ **[M1]**

5 **[A1]**

(b) Recognises $O(0,0)$ is the midpoint of PQ, so sets up $\frac{-3+x}{2} = 0$ and $\frac{4+y}{2} = 0$ OR uses translation from P to O **[M1]**

$(3, -4)$ **[A1]**

(c) Sets up equation for length OR: $k^2 + (\sqrt{11})^2 = 5^2$ **[M1]**

$k^2 + 11 = 25$

$k^2 = 14$ [M1]

$-\sqrt{14}$ **[A1]**

[Back to Question] *[Total Marks: 7]*

4 **(a)** $\left(\frac{-1+5}{2}, \frac{2+8}{2}\right)$ **[M1]**

$(2, 5)$ **[A1]**

(b) $(2-(-1))^2 + (5-2)^2$ or $\frac{1}{2}\sqrt{(5-(-1))^2 + (8-2)^2}$ oe **[M1]**

$\sqrt{9+9}$ or $\sqrt{18}$ **[M1]**

$3\sqrt{2}$ **[A1]**

(c) $\frac{2+x}{2} = \frac{5}{2}$ and $\frac{5+y}{2} = \frac{3}{2}$ oe **[M1]**

$x = 3$ and $y = -2$ to give $D(3, -2)$ **[A1]**

Length $AD = \sqrt{(3-(-1))^2 + (-2-2)^2}$ oe **[M1]**

$\sqrt{16+16}$ or $\sqrt{32}$ **[M1]**

$4\sqrt{2}$ **[A1]**

[Back to Question] *[Total Marks: 10]*

5 **(a)** Uses distance formula: $\sqrt{(4-(-2))^2+(-3-5)^2}$ **[M1]**

Simplifies to $\sqrt{6^2+(-8)^2}$ or $\sqrt{36+64}$ **[M1]**

10 **[A1]**

(b) Uses midpoint formula: $\left(\frac{-2+4}{2}, \frac{5+(-3)}{2}\right)$ **[M1]**

$(1, 1)$ **[A1]**

[Back to Question] *[Total Marks: 5]*

6 **(a)** $(7-3)^2+(5-k)^2=5^2$ or equivalent **[M1]**

$16+(5-k)^2=25$ or $(5-k)^2=9$ **[M1]**

$k=2$ and $k=8$ **[A1]**

(b) Identifies $P(3,2)$ to use in midpoint calculation: $\left(\frac{3+7}{2}, \frac{2+5}{2}\right)$ **[M1]**

$(5, 3.5)$ **[A1]**

[Back to Question] *[Total Marks: 5]*

7 **(a)** $(k-(-3))^2+(-1-5)^2=10^2$ oe **[M1]**

$(k+3)^2+36=100$ or $k^2+6k+9+36=100$ **[M1]**

$(k+3)^2=64$ or $k^2+6k-55=0$ **[M1]**

$k=5$ or $k=-11$ **[A1]**

(b) Chooses $k=5$ to establish coordinates of Q as $(5,-1)$ **[M1]**

$(1, 2)$ **[A1]**

[Back to Question] *[Total Marks: 6]*

8 **(a)** $\left(\frac{-3+5}{2}, \frac{2+8}{2}\right)$ **[M1]**

$(1, 5)$ **[A1]**

(b) Radius $r=\sqrt{(1-(-3))^2+(5-2)^2}$ or $\sqrt{4^2+3^2}=5$ **[M1]**

Area $=\pi\times 5^2$ **[M1]**

78.5 **[A1]**

(c) Distance $CD=5 \implies (k-1)^2+(8-5)^2=5^2$ **[M1]**

$(k-1)^2+9=25 \implies (k-1)^2=16$ **[M1]**

$k = 5$ and $k = -3$ **[A1]**

[Back to Question] *[Total Marks: 8]*

9 **(a)** Sets up the distance formula: $\sqrt{(3p-2p)^2+(4p-p)^2} = 5\sqrt{10}$ **[M1]**

Simplifies the terms inside the root: $\sqrt{p^2+(3p)^2} = 5\sqrt{10}$ **[M1]**

Squares both sides and simplifies: $10p^2 = 250$ **[M1]**

$p = 5$ (rejects -5 as $p > 0$) **[A1]**

(b) Evaluates coordinates of $A(10, 5)$ and $B(15, 20)$ **[B1]**

Sets up a valid expression for the coordinates of C, e.g., x-coordinate $= 10 + \frac{2}{5}(15-10)$ or y-coordinate $= 5 + \frac{2}{5}(20-5)$ **[M1]**

$(12, 11)$ **[A1]**

(c) Uses the midpoint formula, e.g. $\frac{15+x}{2} = 12$ and $\frac{20+y}{2} = 11$, or $D = 2C - B$ **[M1]**

$(9, 2)$ **[A1]**

[Back to Question] *[Total Marks: 9]*