IGCSE 0607 Practice Questions — E1.7 — Combined Questions & Markscheme

Indices I

Total Questions: 6 — **Total Marks:** 36 Marks

Table of Contents & Quick Navigation

- **E1.7** Indices I
  → **Paper 2 (Non-Calculator)** *(5 Qs)* · **Paper 4 (Calculator)** *(1 Q)*

---

1 **(a)** Work out the value of

$125^{\frac{1}{3}}$

..... **[1]**

(b) Work out the value of

6^{-2}

..... **[1]**

(c) Simplify.

$t^3 \times t^5$

..... **[1]**

2 **(a)** Find the exact value of

$16^{-\frac{3}{4}}$

..... **[2]**

(b) Simplify fully.

$(5x^3y^{-2})^2$

..... **[2]**

(c) Solve the equation.

$2^{3x-1} \times 4^x = 32$

$x =$ ..... **[3]**

3 **(a)** Simplify fully.

$$\left(\frac{16x^8}{y^4}\right)^{-\frac{1}{2}}$$

..... **[3]**

(b) Solve the equation.

$$4^x - 2^{2x+1} + 8 = 0$$

$x =$ ..... **[4]**

4 **(a)** Find the value of 7^0.

..... **[1]**

(b) Find the value of $81^{\frac{1}{2}}$.

..... **[1]**

(c) Write as a single power of x.

$x^5 \times x^{-2}$

..... **[1]**

5 **(a)** Find the exact value of each of the following.

(i) $64^{\frac{2}{3}}$

..... **[2]**

(ii) 2^{-4}

..... **[1]**

(b) **(i)** Simplify fully.

$$\frac{15a^6b^3}{3a^2b^7}$$

..... **[2]**

(ii) Simplify fully.

$$(4m^6n^{-2})^{\frac{1}{2}}$$

..... **[2]**

6 **(a)** Find the value of k when $5^k = \dfrac{1}{\sqrt{125}}$.

$k =$ ..... **[3]**

(b) Simplify fully.

$$\left(\frac{16x^{12}}{81y^8}\right)^{-\frac{3}{4}}$$

..... **[3]**

(c) Solve the equation.

$$2^{2x-1} \times 4^x = 32$$

$x =$ ..... **[3]**

1 **(a)** 5 **[B1]**

(b) $\dfrac{1}{36}$ **[B1]**

(c) t^8 **[B1]**

[Back to Question] *[Total Marks: 3]*

2 **(a)** $16^{\frac{1}{4}} = 2$ or equivalent correct first step **[M1]**

$\dfrac{1}{8}$ **[A1]**

(b) 25 **[B1]**

x^6y^{-4} or $\dfrac{x^6}{y^4}$ **[B1]**

(c) Expresses terms with a common base, e.g. $4^x = 2^{2x}$ or $32 = 2^5$ **[M1]**

Sets up a correct linear equation from indices, e.g. $3x - 1 + 2x = 5$ **[M1]**

$x = 1.2$ or $x = \dfrac{6}{5}$ **[A1]**

[Back to Question] *[Total Marks: 7]*

3 **(a)** Inverts fraction to remove negative power or correctly applies negative power to all terms, e.g. $\left(\dfrac{y^4}{16x^8}\right)^{\frac{1}{2}}$ or $\dfrac{16^{-\frac{1}{2}}x^{-4}}{y^{-2}}$ **[M1]**

Correctly evaluates the coefficient $16^{-\frac{1}{2}} = \dfrac{1}{4}$ or correctly halves powers **[M1]**

$\dfrac{y^2}{4x^4}$ or $0.25x^{-4}y^2$ **[A1]**

(b) Recognises that $4^x = 2^{2x}$ or substitutes $u = 4^x$ **[M1]**

Expresses 2^{2x+1} as 2×2^{2x} or $2(4^x)$ **[M1]**

Simplifies to a single exponential term, e.g. $-2^{2x} = -8$ or $u - 2u = -8$ **[M1]**

$x = 1.5$ or $x = \dfrac{3}{2}$ **[A1]**

[Back to Question] *[Total Marks: 7]*

4 **(a)** 1 **[B1]**

(b) 9 **[B1]**

(c) x^3 **[B1]**

[Back to Question] *[Total Marks: 3]*

5 **(a)** **(i)** 16 **[B2]**

If 0 scored, M1 for $(\sqrt[3]{64})^2$ or 4^2 or $4096^{\frac{1}{3}}$ seen **[M1]**

(ii) $\frac{1}{16}$ or 0.0625 **[B1]**

(b) **(i)** $\frac{5a^4}{b^4}$ or $5a^4b^{-4}$ **[B2]**

If 0 scored, M1 for $\frac{15}{3} = 5$ or a^{6-2} or b^{3-7} **[M1]**

(ii) $2m^3n^{-1}$ or $\frac{2m^3}{n}$ **[B2]**

If 0 scored, M1 for $4^{\frac{1}{2}} = 2$ or $m^{6\times\frac{1}{2}}$ or $n^{-2\times\frac{1}{2}}$ **[M1]**

[Back to Question] *[Total Marks: 7]*

6 **(a)** -1.5 or $-\frac{3}{2}$ **[B3]**

If 0 scored, M2 for $5^k = (5^3)^{-\frac{1}{2}}$ or $5^k = 5^{-\frac{3}{2}}$ **[M2]**

If 0 or M2 not scored, M1 for $\sqrt{125} = 125^{\frac{1}{2}}$ or $\frac{1}{125^{0.5}}$ or $125 = 5^3$ **[M1]**

(b) $\frac{27y^6}{8x^9}$ or $3.375y^6x^{-9}$ **[B3]**

If 0 scored, M2 for $\left(\frac{81y^8}{16x^{12}}\right)^{\frac{3}{4}}$ or $\frac{16^{-\frac{3}{4}}x^{-9}}{81^{-\frac{3}{4}}y^{-6}}$ **[M2]**

If 0 or M2 not scored, M1 for finding the 4th root: $\frac{2}{3}$, x^3, or y^2 correctly applied to at least one term **[M1]**

(c) 1.5 or $\frac{3}{2}$ **[B3]**

If 0 scored, M2 for $2x - 1 + 2x = 5$ or $4x - 1 = 5$ **[M2]**

If 0 or M2 not scored, M1 for writing with common base 2: $4^x = 2^{2x}$ or $32 = 2^5$ **[M1]**

[Back to Question] *[Total Marks: 9]*