1 **(a)** Median = 2 **[B1]**

Range = 4 **[B1]**

(b) Team A generally scored more goals because their median is higher ($2 > 1.5$) **[B1]**

Team A was more consistent because their range is smaller ($4 < 5$) **[B1]**

[Total Marks: 4]

2 **(a)** B **[B1]**

(b) 3 **[B1]**

(c) Total students $= 5+10+3+2+7+8+3+2 = 40$ AND Total Grade A $= 5+7 = 12$ **[M1]**

30 **[A1]**

[Total Marks: 4]

3 **(a)** 27 **[B1]**

(b) 31 **[B1]**

(c) 36 and 20 identified as quartiles **[M1]**

16 **[A1]**

(d) $\frac{6}{15}$ or $\frac{2}{5}$ **[B1]**

[Total Marks: 5]

4 **(a)** Identifies the middle value is 7 **[M1]**

Identifies that 3 must appear at least twice as the lowest values: $3, 3$ **[M1]**

Uses the range to find the maximum value: $3 + 11 = 14$ **[M1]**

Uses the mean to find the missing value: $40 - (3 + 3 + 7 + 14) = 13$, yielding $3, 3, 7, 13, 14$ **[A1]**

(b) New sum $= 6 \times 9 = 54$ AND Old sum $= 5 \times 8 = 40$ (or $54 - 40$) **[M1]**

14 **[A1]**

(c) No, because standard deviation measures the spread of all data points around the mean, not just the difference between the extreme values **[B1]**

[Total Marks: 7]

5 **(a)** 45 **[B1]**

(b) 10 **[B1]**

(c) Group A students were faster on average because their median time (45 min) is less than Group B (50 min). **[B1]**

Group B times are more consistent / less spread out because their interquartile range (10

min) is smaller than Group A (20 min). **[B1]**

(d) Recognises that 55 is the upper quartile, representing 25% or $\frac{1}{4}$ of the students **[M1]**

20 **[A1]**

[Total Marks: 6]

6 (a) 65 **[B1]**

(b) $75 - 40$ **[M1]**

35 **[A1]**

(c) Class X has a higher average score (or median is higher: $65 > 55$) **[B1]**

Class X scores are less spread out / more consistent (or IQR is lower: $30 < 35$, or Range is lower: $60 < 65$) **[B1]**

(d) No, because the ranges overlap (e.g., the maximum score in Class Y is 90, which is higher than the lower quartile of Class X) **[B1]**

[Total Marks: 6]

7 (a) $120 < m \leq 140$ **[B1]**

(b) Use of correct midpoints: $110, 130, 150, 170, 190$ **[M1]**

$110 \times 10 + 130 \times 20 + 150 \times 15 + 170 \times 10 + 190 \times 5$ **[M1]**

$= 1100 + 2600 + 2250 + 1700 + 950 = 8600$ **[M1]**

$\frac{8600}{60} = 143.33\ldots$ so 143 (to 3 sf) **[A1]**

(c) The exact masses of the individual mangoes are unknown because the data is grouped **[B1]**

The 15 mangoes could have any combination of masses between 140 g and 160 g, not necessarily exactly at the midpoint **[B1]**

[Total Marks: 7]

8 (a) Upper Quartile = 44 and Lower Quartile = 26 **[M1]**

$44 - 26 = 18$ **[A1]**

(b) The Adults generally completed the puzzle faster because their median time is lower ($32 < 36$) **[B1]**

The Students' times were more consistent (less spread out) because their interquartile range (12) is smaller than the Adults' interquartile range (18) **[B1]**

(c) The reasoning is invalid because range only measures spread, not the actual values or maximum times **[B1]**

However, the conclusion happens to be true because the maximum time for Students (48) is less than the maximum time for Adults (55) **[B1]**

[Total Marks: 6]

9 **(a)** 112 **[B1]**

(b) 120 and 104 seen as quartiles **[M1]**

16 **[A1]**

(c) Cumulative frequency of 110 seen, leading to $120 - 110 = 10$ apples **[M1]**

8.33 (8.333...) **[A1]**

(d) Farm X apples are heavier on average as their median (112 g) is greater than Farm Y (105 g). **[B1]**

Farm X apples are more consistent in mass / less spread out as their interquartile range (16 g) is smaller than Farm Y (22 g). **[B1]**

[Total Marks: 7]

10 **(a)** Class B, because the mean (or median) is higher. **[B1]**

(b) Class B, because the range (or interquartile range) is smaller. **[B1]**

(c) Identifies that the table only gives summary statistics / does not show individual marks **[B1]**

Mentions overlapping ranges / a specific example (e.g. the lowest mark in Class B could be lower than the highest mark in Class A) **[B1]**

[Total Marks: 4]

11 **(a)** 40 **[B1]**

(b) 25 (from $50 - 25$) **[B1]**

(c) Group Y was slower on average / had a higher median time ($40 > 35$) **[B1]**

Group Y's times were more consistent / had a smaller IQR ($15 < 25$) or smaller range ($40 < 60$) **[B1]**

(d) Identifies that exactly 25% of Group Y took less than 30 minutes (since lower quartile is 30) **[M1]**

States Group X had a larger proportion, because 30 is between their LQ (25) and median (35), meaning more than 25% took less than 30 minutes **[A1]**

[Total Marks: 6]

12 **(a)** Forms a correct equation: $x + 2x + 120 + x = 360$ **[M1]**

Solves to find $4x = 240$

x = 60[$M1$]

Calculates number of students: $\frac{60}{360} \times 120 = 20$ **[A1]**

(b) Calculates the percentage for Class P: $\frac{120^\circ}{360^\circ} \times 100 = 33.3\%$ (or $\frac{1}{3}$) **[M1]**

Identifies total ratio parts for Class Q $= 1 + 3 + 2 + 2 = 8$ **[M1]**

Calculates percentage for Class Q: $\frac{3}{8} \times 100 = 37.5\%$ and concludes $37.5\% > 33.3\%$ **[A1]**

(c) The two classes have different total numbers of students (so proportions/percentages must be used for a fair comparison). **[B1]**

[Total Marks: 7]