1 $\sqrt{5^2 + (-12)^2}$ or $\sqrt{25 + 144}$ **[M1]**

13 **[A1]**

[Total Marks: 2]

2 **(a)** $\sqrt{(-8)^2 + 15^2}$ or $\sqrt{64 + 225}$ **[M1]**

17 **[A1]**

(b) $\mathbf{p} + 2\mathbf{q} = \begin{pmatrix} -8 \\ 15 \end{pmatrix} + \begin{pmatrix} 4 \\ -8 \end{pmatrix} = \begin{pmatrix} -4 \\ 7 \end{pmatrix}$ **[M1]**

$\sqrt{(-4)^2 + 7^2}$ or $\sqrt{16 + 49}$ **[M1]**

$\sqrt{65}$ **[A1]**

[Total Marks: 5]

3 **(a)** $\sqrt{x^2 + (x+2)^2} = 10$ or $x^2 + (x+2)^2 = 10^2$ **[M1]**

$x^2 + x^2 + 4x + 4 = 100$ **[M1]**

$2x^2 + 4x - 96 = 0$ leading to $x^2 + 2x - 48 = 0$ with no errors seen **[A1]**

(b) $(x + 8)(x - 6) = 0$ or correct substitution into quadratic formula **[M1]**

$x = -8, x = 6$ **[A1]**

(c) $\mathbf{u} = \begin{pmatrix} -8 \\ -6 \end{pmatrix}$ **[B1]**

$\mathbf{u} = \begin{pmatrix} 6 \\ 8 \end{pmatrix}$ **[B1]**

[Total Marks: 7]

4 **(a)** $\vec{AC} = \begin{pmatrix} 7 - 3 \\ 1 - (-2) \end{pmatrix}$ or $\begin{pmatrix} 4 \\ 3 \end{pmatrix}$ **[M1]**

5 **[A1]**

(b) $\vec{AB} = \begin{pmatrix} -8 \\ y + 2 \end{pmatrix}$ **[B1]**

$\sqrt{(-8)^2 + (y+2)^2} = 17$ **[M1]**

$64 + (y+2)^2 = 289$ or $(y+2)^2 = 225$ **[M1]**

$y + 2 = 15$ or $y + 2 = -15$ **[M1]**

$y = 13$, $y = -17$ **[A1]**

[Total Marks: 7]

5 **(a)** $\begin{pmatrix} 9 - 2 \\ 4 - (-3) \end{pmatrix}$ **[M1]**

$\begin{pmatrix}7\\7\end{pmatrix}$ **[A1]**

(b) $\sqrt{7^2+7^2}$ or $\sqrt{98}$ **[M1]**

9.90 **[A1]**

(c) $\sqrt{k^2+4^2}=5$ or $k^2+4^2=5^2$ **[M1]**

$k^2=25-16$ or $k^2=9$ **[M1]**

$k=3$, $k=-3$ **[A1]**

[Total Marks: 7]

6 (a) $\mathbf{u}+\mathbf{v}=\begin{pmatrix}3k+1\\k-1\end{pmatrix}$ **[M1]**

$(3k+1)^2+(k-1)^2=(\sqrt{50})^2$ **[M1]**

$9k^2+6k+1+k^2-2k+1=50$ **[M1]**

$10k^2+4k+2=50 \implies 5k^2+2k-24=0$ **[A1]**

(b) $(5k+a)(k+b)=0$ where $ab=-24$ or $5b+a=2$, or correct substitution into quadratic formula **[M1]**

$(5k+12)(k-2)=0$ **[M1]**

$k=2$ **[A1]**

$k=-2.4$ or $-\dfrac{12}{5}$ **[A1]**

(c) Substituting $k=2$ and $k=-2.4$ into $\begin{pmatrix}2k\\k-3\end{pmatrix}$ **[M1]**

$\begin{pmatrix}4\\-1\end{pmatrix}$ and $\begin{pmatrix}-4.8\\-5.4\end{pmatrix}$ **[A1]**

[Total Marks: 10]

7 (a) Determines column vector $\mathbf{a}=\begin{pmatrix}-6\\8\end{pmatrix}$ **[M1]**

10 **[A1]**

(b) Determines column vector $\mathbf{b}=\begin{pmatrix}3\\-4\end{pmatrix}$ **[B1]**

Calculates resultant vector $2\begin{pmatrix}-6\\8\end{pmatrix}+3\begin{pmatrix}3\\-4\end{pmatrix}=\begin{pmatrix}-12\\16\end{pmatrix}+\begin{pmatrix}9\\-12\end{pmatrix}=\begin{pmatrix}-3\\4\end{pmatrix}$ **[M1]**

5 **[A1]**

[Total Marks: 5]

8 (a) Expresses vector $\vec{PQ}$ as $\begin{pmatrix}k-2\\5-(-3)\end{pmatrix}=\begin{pmatrix}k-2\\8\end{pmatrix}$ **[M1]**

Sets up magnitude equation: $(k-2)^2+8^2=10^2$ **[M1]**

Simplifies to solve for k: $(k-2)^2 = 36$ **[M1]**

$k = 8$, or $k = -4$ **[A1]**

(b) Finds $\vec{QR}$ using $Q(8,5)$ and $R(4,1)$: $\vec{QR} = \begin{pmatrix} 4-8 \\ 1-5 \end{pmatrix} = \begin{pmatrix} -4 \\ -4 \end{pmatrix}$ **[M1]**

Calculates magnitude: $\sqrt{(-4)^2 + (-4)^2} = \sqrt{32}$ **[M1]**

5.66 **[A1]**

[Total Marks: 7]

9 (a) Sets up equation: $p^2 + p^2 = 18$ **[M1]**

3 **[A1]**

(b) Finds expression for $\mathbf{w}$: $\begin{pmatrix} 4-t \\ 1+2t \end{pmatrix}$ **[M1]**

Forms magnitude squared equation: $(4-t)^2 + (1+2t)^2 = 29$ **[M1]**

Expands brackets correctly: $16 - 8t + t^2 + 1 + 4t + 4t^2 = 29$ **[M1]**

Rearranges to a 3-term quadratic: $5t^2 - 4t - 12 = 0$ **[M1]**

$t = 2$, or $t = -1.2$ (or $-\frac{6}{5}$) **[A1]**

[Total Marks: 7]