

Contents

- [Features](#)
- [Prerequisites](#)
 - [Installation of the AstroBinUpload.py script](#)
 - [config.ini generation](#)
 - [config.ini contents and editing](#)
 - [\[defaults\]](#)
 - [\[filters\]](#)
 - [\[secrets\]](#)
 - [\[sites\]](#)
 - [Editing the initial config.ini](#)
- [Running the Script](#)
 - [Initialization](#)
 - [Single directory or symbolic link passed to script](#)
 - [Multiple directory paths or symbolic links are passed to the script](#)
 - [Debug output](#)
- [Example calls and outputs](#)
 - [Single site, non-mosaic, no Masters. Data resides in structured single directory, symbolic links used for calibration data](#)
 - [Single site, 2 panel mosaic, symbolic links to calibration data, use of MASTERFLATS](#)
 - [Dual site, structured directory, 2 panel mosaic, use of MASTERCALS](#)
 - [WBPP two-panel mosaic](#)
- [References](#)
 - [AstroBin's Acquisition CSV File Format](#)
 - [AstroBin's Long Exposure Acquisition Fields](#)
 - [Astrobin Filter-Code mappings](#)
 - [Finding the AstroBin's Numeric ID for Filters](#)
 - [Accessing sky quality data](#)
 - [Reverse Geocoding](#)
 - [FWHM values](#)
 - [Data Sources](#)
- [Contributing to AstroBinUpload.py Processing Script](#)
- [Contact](#)
- [Licence](#)

Features

When run this Python script creates a detailed observation session summary and an acquisition.csv file suitable for upload using AstroBin's import CSV dialogue.

Data is obtained by extracting FITS (Flexible Image Transport System) or XISF (Extensible Image Serialization Format) headers from image and calibration files associated with the given astronomical target.

Key features include:

- **The ability to pass multiple directories via the command line:** Multiple directories can be passed to the script via the command line. All images results contained within the directories will be accumulated as part of the target.
- **Structured and unstructured directories:** Image files, including calibration files, can be collected into a single directory, the root directory. The root directory structure can be flat or contain subdirectories.
- **Symbolic links to directories:** Symbolic links can be used within the root directory or passed directly via the command line, this is useful when reusing calibration directories. The first directory passed should be the root directory.
- **MASTER calibration files:** If MASTER calibration files are found, these will be used. If the non-MASTER versions of the MASTER files are also found, the non-MASTER versions will be ignored.
- **Processing of PixInsight's Weighted Batch Pre-processing (WBPP) output:** When the target is a WBPP directory the script will use the calibrated LIGHT frames as well as any MASTER calibration files found in the directory. MASTERLIGHT or processed image files are ignored.
- **Multiple panel mosaic imaging sessions:** Mosaic imaging sessions are detected from the OBJECT entry in the FITS headers. LIGHT frames are processed on a per-panel basis, whilst calibration data is processed per target. For this to work correctly the image names (OBJECT in FITS header) must have the format

```
"target name Panel x"
```

where x is the panel number. N.I.N.A does this automatically but in Sequence Generator Pro the user will have to edit the directory name in Target Settings before starting the sequence.

- **Multiple site support:** Multi-site collaborative target acquisition or remote observatory image capture is supported. Site location data is reverse-geocoded from HEADER location data. Data from multiple sites is reported with summary outputs that correctly identify the site contribution, for instance equipment, LIGHT, and calibration data. All data is, however, aggregated in the AstroBin.csv file for the image target.
- **Support for multiple file formats:** Extracts headers for all FITS/FIT/FTS/XISF files in specified directories. Directories can have a mix of files.

- **Accepts files generated by N.I.N.A, SGPro and PixInsight**
- **Sky Quality Retrieval:** Recovers SQM and Bortle scale classification based on the observation location coordinates.
- **Auxiliary Parameter Calculation:** Calculates additional parameters like Image Scale (IMSCALE), and Full-width Half Maximum (FWHM) from measured/estimated HFR values for each image.
- **AstroBin Compatibility:** Formats aggregated data for upload to AstroBin's import CSV file dialogue.
- **Target summary:** Creates a detailed summary text file for a target acquisition session. Caters for single or multi-site data as well as single or mosaic imaging data.

Pre-requisites

Before using this script, ensure you have Python 3.x installed. [Python Installation Instructions](#).

Installation of the AstroBinUpload.py script

To install this script, follow these steps:

1. Create a directory to hold the script and associated files
2. Clone or download the repository from my [GitHub repository](#)
3. Ensure the following files are in the new directory:
 - AstroBinUpload.py
 - utils.py
 - requirements.txt
4. To install the required python libraries, navigate to the new directory and run the following command:

```
pip install -r requirements.txt
```

Initialization and Config.ini generation

The script is first run with no arguments. This will create a default config.ini file in the local directory. The config.ini file defines parameters the script needs to run correctly. This file can be personalized by the user. A description of the config.ini file parameters is given later. Then you run the script for the first time or if your config.ini file has been deleted, you will see the following:

```
'A new config.ini file was created. Please edit this before re-running the script:'
```

The script will halt, allowing you to edit the config.ini file using a text editor. You can then re-run the script.

If you run the script when the config.ini file exists and pass no arguments, an error will be flagged, and the script will exit.

Config.ini contents and editing

The config.ini file contains the following sections:

[defaults]

The [defaults] section holds:

1. Default FITS keywords that should be present in the header files and that are needed in the production of the astrobin.csv output file
2. Default site location and auxiliary information, used in the production of the summary.txt output file

The data in the [defaults] section can be modified by the user. If the script cannot find the information in the header files it will take it from the [defaults] section of the config.ini file

[filters]

The filter section holds the filter name to AstroBin code mappings. The filter names and codes can be modified here.

[secrets]

The secret section holds:

1. The sky quality API keys and API endpoint required by the script to be able to obtain values of Bortle and SQM for the site location. Only the API key is to be edited. If there is no valid API key the values of Bortle and SQM are taken from [defaults][BORTLE] and [defaults][SQM] in the config.ini file
2. User email address. This is used as part of an information string sent to the reverse geocoding API, which is used to recover the site address. Unique site latitude and longitude values extracted from the headers are passed to the API to generate the site address. Your email address is passed to the API as courtesy, so the provider can see who is using their API. If the API request fails the site location information is taken from [defaults][SITE], [defaults][SITELAT] and [defaults][SITELONG] in the config.ini file.

[sites]

The [sites] section holds historic site information found by the script. When a script is run it first looks here to collect site information. Only if a site found in the headers does not exist does it access the external API's. The script automatically updates this section if a new site is found. The user does not usually have to edit this section, but remote site information can be added here if the script cannot access the API.

It is advisable to back up your config.ini file regularly.

Editing the config.ini

A config.ini file with an explanation of the sections is given below:

```
[defaults]
    IMAGETYP = LIGHT
    EXPOSURE = 0.0
    DATE-OBS = 2023-01-01
    XBINNING = 1
```

These are place-holders and are fall backs. These parameters should be created by the capture software.

```
GAIN = -1
EGAIN = -1
```

If you have a CCD camera, leave these values as they are. If you have a CMOS camera these gain values can be set to the typical values of your camera. The script will, however, collect the correct results for both CCD and CMOS cameras from the headers processed.

```
INSTRUME = None
TELESCOP = None
FOCNAME = None
FWHEEL = None
ROTATOR = None
XPIXSZ = 1
CCD-TEMP = -10
FOCALLEN = 540
FOCRATIO = 5.4
```

This is where default the equipment configuration. Again the script should be able to populate these parameters from the header information. XPIXSZ is the X-pixel size in um and is used to represent the sensor pixel size in the script.

```
SITE = Papworth Everard
SITELAT = 52.2484
SITELONG = -0.1231
BORTLE = 4
SQM = 20.5
```

You should modify these parameters to reflect your own site. If any API call fails the script will fall back to these parameters

```
FILTER = No Filter
```

If you use a color camera and don't report your filters automatically you should enter your filter name here as there may be no filter information in the header.

```
OBJECT = No target
FOCTEMP = 20
```

These are place-holders and should not be required as they should be populated by the capture software

```
HFR = 1.6
```

You should set this value to the typical value for your imaging train. If you use N.I.N.A you can add the measured HFR for the image to the file name. The script looks for HFR=X.XX in the image file name and if present uses the value found, if HFR is not in the image file name the script falls back to this value.

```
SWCREATE = Unknown package
```

This is a place-holder and should not be required, it should be created by the capture software

```
[filters]
#Filter      code
Ha           = 4663
SII          = 4844
OIII         = 4752
Red          = 4649
Green        = 4643
Blue         = 4637
Lum          = 2906
```

Modify the [filters] section to reflect your imaging set up, see [Astrobin Filter-Code mappings](#) for information on how to populate this table. If you use a color camera and don't report your filters automatically you should enter your filter name and the corresponding code here. Delete filters you don't require.

```
[secret]
#API key      API endpoint
xxxxxxxxxxxxx = https://www.lightpollutionmap.info/QueryRaster/
EMAIL_ADDRESS = id@provider.com
```

If you wish to automatically generate an address and sky quality information for the observation site enter the [Sky Quality API key](#) and your [email address](#) here. If you don't wish to do this or if the API calls fail the script falls back to site parameters found in the [defaults] section

```
[sites]
```

If the script is run and a new site identified the script will update the [sites] section with the new site information. For example after one run my home location was added to the [sites] as shown below:

```
[sites]
[["Norton Close, Papworth Everard, South Cambridgeshire,
Cambridgeshire, Cambridgeshire and Peterborough, England, CB23 3XT, United
Kingdom"]]
    latitude = 52.2484
    longitude = -0.1231
    bortle = 4
    sqm = 20.52
```

When the script processes [SITELAT] and [SITELONG] header entries it looks here first to see if a site has been seen before. If it has the script uses the site information found, if not it calls the external APIs to retrieve the information. If the external API call fails the script falls back to site parameters found in the [defaults] section of the configs.ini file. New sites can also be manually added in the [sites] section following the format shown above.

Running the Script

The script is called from the command line. There are three calling methods:

Initialization, no arguments are passed:

```
Linux/MACOS example: python3 AstroBinUpload.py
Windows example:     python  AstroBinUpload.py
```

When called with no argument the script will create a new default config.ini file in the local directory and then exit. The user can then edit the config.ini file, using a text editor, before running the script again. If an existing config.ini file is lost and the script run, a new default config.ini file will be created, and the code will exit. Once you have personalized your config.ini file make a backup.

A single directory path or symbolic link argument is passed to the script

Note: only Linux calling examples are used going forward.

```
python3 AstroBinUpload.py "dir 1"
```

The script expects to find all data contained in the directory passed to it. Symbolic links can be used as the argument passed to script and can also be present in the directory. The directory leaf or child directory name must be the target name if the output files are to be named correctly. From the processing perspective the only condition required to ensure data is associated with a given target is that all data and links must reside in the one directory.

Multiple directory paths or symbolic links are passed to the script

```
python3 AstroBinUpload.py "dir 1" "dir 2" ....
```

All directory arguments are assumed to belong to one target. Again the first directory leaf, or child directory name should contain the target name for the output files to be named correctly.

Debug output

```
python3 AstroBinUpload.py "dir 1" "dir 2" ... --debug
```


This will dump four processed header files to the current directory as .csv files. The files are

```
Target_name_basic_headers.csv  
Target_name_headers.csv  
Target_name_modified.csv  
Target_name_aggregated.csv
```

These files can be used to assist the debugging of any issues

Example calls and outputs

Example 1: Single site, non-mosaic, no Masters, data resides in structured single directory, symbolic links used for calibration data



Example 1: Directory structure

```
python3 AstroBinUpload.py "/mnt/HDD_8TB/Preselected/Sadr Region"
```

Example 1: Script calling syntax

The output files being named:

- Sadr_Region_session_summary.txt
- Sadr_Region_aquisition.csv

```
Observation session summary
Generated 2024-01-31 09:34:42

Target: Gamma Cygni Nebula

Site: Norton Close, Papworth Everard, South Cambridgeshire, Cambridgeshire, Cambridgeshire and Peterborough, England, CB23 3XT, United Kingdom
Latitude: 52.2485°, Longitude: -0.1232°
Bottle scale: 4.0, SQM: 20.52 mag/arcsec²

Equipment used:
Telescope      : NP101is
Camera         : ZWO ASI6200MM PRO
Filterwheel    : Starlight Xpress Fil
Focuser       : FocusLynx Focuser 1
Capture software : N.I.N.A. 2.3.0.2001
               : N.I.N.A. 3.0.0.1028

Observation period:
Start date      : 2023-07-06
End date       : 2023-08-11
Number of days  : 36
Observation sessions : 6
Min temperature : 7.8°C
Max temperature : 18.9°C
Mean temperature : 13.9°C

LIGHTS:
Target: Gamma Cygni Nebula

Filter  Gain      Egain      Mean FWHM  Frames  Sensor Temperature  Mean Temperature  Exposure  Total Exposure
Blue   0.0 dB    0.78 e/ADU  4.74 arcsec  17      -10.0°C            16.9°C            60.00 secs  17 mins 0 secs
Green  0.0 dB    0.78 e/ADU  4.68 arcsec  18      -10.0°C            16.8°C            60.00 secs  18 mins 0 secs
Ha     10.0 dB   0.25 e/ADU  4.78 arcsec  51      -10.0°C            13.5°C            600.00 secs  8 hrs 30 mins 0 secs
OIII   10.0 dB   0.25 e/ADU  4.74 arcsec  54      -10.0°C            13.2°C            600.00 secs  9 hrs 0 mins 0 secs
Red    0.0 dB    0.78 e/ADU  4.81 arcsec  30      -10.0°C            16.9°C            60.00 secs  30 mins 0 secs
SII    10.0 dB   0.25 e/ADU  4.85 arcsec  51      -10.0°C            12.7°C            600.00 secs  8 hrs 30 mins 0 secs

Exposure time for Gamma Cygni Nebula: 27 hrs 5 mins 0 secs
Total LIGHT Exposure Time: 27 hrs 5 mins 0 secs

FLATS:

Filter  Frames  Gain      Egain      Exposure  Total Exposure
Blue   100     0.0 dB    0.78 e/ADU  0.22 secs  22 secs
Green  100     0.0 dB    0.78 e/ADU  0.17 secs  17 secs
Ha     100     10.0 dB   0.25 e/ADU  0.47 secs  47 secs
Lum    100     0.0 dB    0.78 e/ADU  0.05 secs  5 secs
OIII   100     10.0 dB   0.25 e/ADU  0.55 secs  55 secs
Red    100     0.0 dB    0.78 e/ADU  0.10 secs  10 secs
SII    100     10.0 dB   0.25 e/ADU  0.58 secs  57 secs

Total FLAT Exposure Time: 0 hrs 3 mins 34 secs

BIASES:

Filter  Frames  Gain      Egain      Exposure  Total Exposure
Ha     99      0.0 dB    0.78 e/ADU  0.00 secs  0 secs
Ha    100     10.0 dB   0.25 e/ADU  0.00 secs  0 secs

Total BIAS Exposure Time: 0 hrs 0 mins 0 secs

DARKS:

Filter  Frames  Gain      Egain      Sensor Temperature  Exposure  Total Exposure
Ha     49      0.0 dB    0.78 e/ADU  -10.0°C            60.00 secs  49 mins 0 secs
Ha     50      10.0 dB   0.25 e/ADU  -10.0°C            600.00 secs  8 hrs 20 mins 0 secs

Total DARK Exposure Time: 9 hrs 9 mins 0 secs

Total number of images processed: 1219
```

Example 1: Summary output

```
Sadr_Region_acquisition.csv
```

date	filter	number	duration	binning	gain	sensorCooling	fNumber	darks	flats	flatDarks	bias	bortle	meanSqm	meanFwhm	temperature
2023-07-06	4663	20	600.0	1	100	-10	5.4	50	100	0	100	4.0	20.52	4.76	10.27
2023-07-07	4663	24	600.0	1	100	-10	5.4	50	100	0	100	4.0	20.52	4.75	13.08
2023-07-08	4663	7	600.0	1	100	-10	5.4	50	100	0	100	4.0	20.52	4.82	17.16
2023-07-08	4844	5	600.0	1	100	-10	5.4	50	100	0	100	4.0	20.52	4.91	16.82
2023-07-29	4844	2	600.0	1	100	-10	5.4	50	100	0	100	4.0	20.52	4.80	13.40
2023-07-30	4844	1	600.0	1	100	-10	5.4	50	100	0	100	4.0	20.52	4.80	12.80
2023-08-06	4844	12	600.0	1	100	-10	5.4	50	100	0	100	4.0	20.52	4.83	11.29
2023-08-07	4844	18	600.0	1	100	-10	5.4	50	100	0	100	4.0	20.52	4.87	9.19
2023-08-08	4752	12	600.0	1	100	-10	5.4	50	100	0	100	4.0	20.52	4.78	9.92
2023-08-08	4844	5	600.0	1	100	-10	5.4	50	100	0	100	4.0	20.52	4.90	9.86
2023-08-09	4752	15	600.0	1	100	-10	5.4	50	100	0	100	4.0	20.52	4.89	10.58
2023-08-09	4844	8	600.0	1	100	-10	5.4	50	100	0	100	4.0	20.52	4.86	15.78
2023-08-10	4752	25	600.0	1	100	-10	5.4	50	100	0	100	4.0	20.52	4.68	15.76
2023-08-10	4649	15	60.0	1	0	-10	5.4	49	100	0	99	4.0	20.52	4.82	17.08
2023-08-11	4637	17	60.0	1	0	-10	5.4	49	100	0	99	4.0	20.52	4.74	16.86
2023-08-11	4643	18	60.0	1	0	-10	5.4	49	100	0	99	4.0	20.52	4.68	16.77
2023-08-11	4752	2	600.0	1	100	-10	5.4	50	100	0	100	4.0	20.52	4.61	16.65
2023-08-11	4649	15	60.0	1	0	-10	5.4	49	100	0	99	4.0	20.52	4.80	16.80


```

Processing summary exported to Sadr_Region_session_summary.txt
AstroBin data exported to Sadr_Region_acquisition.csv
Processing completed.

```

Example 1: AstroBin.csv output

Example 2: Single site, 2 panel mosaic, symbolic links to calibration data, use of MASTERFLATS



Example 2: Directory structure

```
python3 AstroBinUpload.py '/mnt/HDD_8TB/Preselected/NGC 1499 Mosaic'
```

or using a symbolic link:

```
python3 AstroBinUpload.py '/home/steve/Desktop/AstroData/Link to NGC 1499 Mosaic'
```

Example 2: Script calling syntax

```
Observation session summary
Generated 2024-01-31 09:36:32

Target: NGC 1499 2 Panel Mosaic

Site: Norton Close, Papworth Everard, South Cambridgeshire, Cambridgeshire, Cambridgeshire and Peterborough, England, CB23 3XT, United Kingdom
Latitude: 52.2486°, Longitude: -0.123°
Bortle scale: 4.0, SQM: 20.52 mag/arcsec²

Equipment used:
Telescope      : NP101is
Camera         : ZWO ASI6200MM PRO
Filterwheel    : Starlight Xpress Filter Wheels
Focuser       : FocusLynx Focuser 1
Capture software : N.I.N.A. 2.3.2.9001 (x64)
               : N.I.N.A. 3.0.0.2003 (x64)

Observation period:
Start date      : 2023-11-14
End date       : 2024-01-10
Number of days  : 57
Observation sessions : 10
Min temperature : -3.9°C
Max temperature : 10.5°C
Mean temperature : 1.2°C

LIGHTS:
Target: NGC 1499 Panel 1

Filter  Gain    Egain      Mean FWHM   Frames   Sensor Temperature  Mean Temperature  Exposure  Total Exposure
Blue   0.0 dB    0.78 e/ADU   4.60 arcsec  30      -10.0°C            -3.7°C            60.00 secs  30 mins 0 secs
Green  0.0 dB    0.78 e/ADU   4.66 arcsec  30      -10.0°C            -3.7°C            60.00 secs  30 mins 0 secs
Ha     10.0 dB   0.25 e/ADU   4.74 arcsec  66      -10.0°C            2.4°C            600.00 secs  11 hrs 0 mins 0 secs
OIII   10.0 dB   0.25 e/ADU   5.52 arcsec  84      -10.0°C            2.1°C            600.00 secs  14 hrs 0 mins 0 secs
Red    0.0 dB    0.78 e/ADU   4.79 arcsec  25      -10.0°C            -3.5°C            60.00 secs   25 mins 0 secs
SII    10.0 dB   0.25 e/ADU   4.94 arcsec  67      -10.0°C            3.7°C            600.00 secs  11 hrs 10 mins 0 secs

Exposure time for NGC 1499 Panel 1: 37 hrs 35 mins 0 secs

Target: NGC 1499 Panel 2

Filter  Gain    Egain      Mean FWHM   Frames   Sensor Temperature  Mean Temperature  Exposure  Total Exposure
Blue   0.0 dB    0.78 e/ADU   5.17 arcsec  30      -10.0°C            -3.2°C            60.00 secs  30 mins 0 secs
Green  0.0 dB    0.78 e/ADU   4.93 arcsec  30      -10.0°C            -3.2°C            60.00 secs  30 mins 0 secs
Ha     10.0 dB   0.25 e/ADU   4.88 arcsec  66      -10.0°C            -1.0°C            600.00 secs  11 hrs 0 mins 0 secs
OIII   10.0 dB   0.25 e/ADU   4.80 arcsec  65      -10.0°C            1.7°C            600.00 secs  10 hrs 50 mins 0 secs
Red    0.0 dB    0.78 e/ADU   4.97 arcsec  30      -10.0°C            -3.3°C            60.00 secs   30 mins 0 secs
SII    10.0 dB   0.25 e/ADU   5.11 arcsec  57      -10.0°C            5.2°C            600.00 secs  9 hrs 30 mins 0 secs

Exposure time for NGC 1499 Panel 2: 32 hrs 50 mins 0 secs

Total LIGHT Exposure Time: 70 hrs 25 mins 0 secs

Total number of images processed: 580
```

Example 2: Summary Output

NGC_1499_Mosaic_acquisition.csv

date	filter	number	duration	binning	gain	sensorCooling	fNumber	darks	flats	flatDarks	bias	bottle	meanSqm	meanFwhm	temperature
2023-11-14	4663	18	600.0	1	100	-10	5.4	0	0	0	0	4.0	20.52	4.76	5.86
2023-11-15	4663	41	600.0	1	100	-10	5.4	0	0	0	0	4.0	20.52	4.78	5.22
2023-11-15	4844	10	600.0	1	100	-10	5.4	0	0	0	0	4.0	20.52	4.96	3.31
2023-11-16	4844	3	600.0	1	100	-10	5.4	0	0	0	0	4.0	20.52	4.89	3.17
2023-11-18	4844	6	600.0	1	100	-10	5.4	0	0	0	0	4.0	20.52	5.09	9.35
2023-11-19	4844	24	600.0	1	100	-10	5.4	0	0	0	0	4.0	20.52	4.91	8.70
2023-11-24	4663	11	600.0	1	100	-10	5.4	0	0	0	0	4.0	20.52	4.76	-0.34
2023-11-24	4844	13	600.0	1	100	-10	5.4	0	0	0	0	4.0	20.52	4.97	0.17
2023-11-25	4663	35	600.0	1	100	-10	5.4	0	0	0	0	4.0	20.52	4.86	-1.59
2023-11-30	4637	30	60.0	1	0	-10	5.4	0	0	0	0	4.0	20.52	4.60	-3.71
2023-11-30	4643	30	60.0	1	0	-10	5.4	0	0	0	0	4.0	20.52	4.66	-3.73
2023-11-30	4663	7	600.0	1	100	-10	5.4	0	0	0	0	4.0	20.52	4.69	-3.76
2023-11-30	4663	4	600.0	1	100	-10	5.4	0	0	0	0	4.0	20.52	4.75	-3.38
2023-11-30	4649	25	60.0	1	0	-10	5.4	0	0	0	0	4.0	20.52	4.79	-3.54
2023-11-30	4844	11	600.0	1	100	-10	5.4	0	0	0	0	4.0	20.52	4.83	-2.75
2023-12-01	4663	5	600.0	1	100	-10	5.4	0	0	0	0	4.0	20.52	4.75	-3.78
2023-12-05	4752	4	600.0	1	100	-10	5.4	0	0	0	0	4.0	20.52	4.88	1.38
2023-12-06	4752	31	600.0	1	100	-10	5.4	0	0	0	0	4.0	20.52	4.83	-1.98
2023-12-09	4752	15	600.0	1	100	-10	5.4	0	0	0	0	4.0	20.52	4.97	6.80
2023-12-10	4663	10	600.0	1	100	-10	5.4	0	0	0	0	4.0	20.52	5.38	5.74
2023-12-10	4752	2	600.0	1	100	-10	5.4	0	0	0	0	4.0	20.52	5.00	6.50
2023-12-10	4844	30	600.0	1	100	-10	5.4	0	0	0	0	4.0	20.52	5.11	5.87
2023-12-11	4844	8	600.0	1	100	-10	5.4	0	0	0	0	4.0	20.52	5.07	7.09
2023-12-19	4752	2	600.0	1	100	-10	5.4	0	0	0	0	4.0	20.52	4.89	2.50
2023-12-19	4844	19	600.0	1	100	-10	5.4	0	0	0	0	4.0	20.52	5.15	2.51
2023-12-20	4752	16	600.0	1	100	-10	5.4	0	0	0	0	4.0	20.52	5.01	2.21
2023-12-29	4752	21	600.0	1	100	-10	5.4	0	0	0	0	4.0	20.52	4.67	2.83
2024-01-02	4752	6	600.0	1	100	-10	5.4	0	0	0	0	4.0	20.52	4.66	5.62
2024-01-03	4752	17	600.0	1	100	-10	5.4	0	0	0	0	4.0	20.52	4.64	5.62
2024-01-09	4663	1	600.0	1	100	-10	5.4	0	0	0	0	4.0	20.52	4.80	-2.40
2024-01-09	4752	11	600.0	1	100	-10	5.4	0	0	0	0	4.0	20.52	8.74	-2.64
2024-01-09	4752	14	600.0	1	100	-10	5.4	0	0	0	0	4.0	20.52	4.75	-2.59
2024-01-10	4637	30	60.0	1	0	-10	5.4	0	0	0	0	4.0	20.52	5.17	-3.22
2024-01-10	4643	30	60.0	1	0	-10	5.4	0	0	0	0	4.0	20.52	4.93	-3.20
2024-01-10	4752	10	600.0	1	100	-10	5.4	0	0	0	0	4.0	20.52	4.83	-2.97
2024-01-10	4649	30	60.0	1	0	-10	5.4	0	0	0	0	4.0	20.52	4.97	-3.27

Processing summary exported to NGC_1499_Mosaic_session_summary.txt


AstroBin data exported to NGC_1499_Mosaic_acquisition.csv

Processing completed.


Example 2: AstroBin.csv output

Example 3: Dual site, structured directory, 2 panel mosaic, use of MASTERCALS

Note: although data is reported on a per-site basis, data is aggregated from all sites to create the AstroBin.csv output. Symbolic links can also be used. A non-structured directory can also be used as long as all files reside under the main target directory. Mosaics can be generated per site, however, summary files can be quite large.



Example 3: Directory structure



Example 3: Script calling syntax

```
python3 AstroBinUpload.py "/mnt/HDD_8TB/AstroBinTest/M51"
```

Example 3: Summary Output

Site 1

```
Observation session summary
Generated 2024-01-31 09:40:13

Target: M51 2 Panel Mosaic

Site: Church Street, Belmont, St. George's, Saint George, Grenada
Latitude: 12.0529°, Longitude: -61.7523°
Bottle scale: 4.0, SQM: 20.69 mag/arcsec²

Equipment used:
Telescope      : SkyWatcher 80ED
Camera         : ZWO ASI2600M PRO
Filterwheel    : QHYCFW3-S
Focuser       : PrimaluceLab Sesto Senso 2
Rotator       : Pegasus Falcon Rotator
Capture software : N.I.N.A. 2.3.2.9001 (x64)
               : PixInsight 1.8.9-2

Observation period:
Start date      : 2023-09-11
End date       : 2023-09-23
Number of days  : 12
Observation sessions : 4
Min temperature : 10.0°C
Max temperature : 10.0°C
Mean temperature : 10.0°C

LIGHTS:
Target: M51 Panel 1

Filter  Gain  Egain      Mean FWHM  Frames  Sensor Temperature  Mean Temperature  Exposure  Total Exposure
Blue   0.0 dB  0.78 e/ADU  7.56 arcsec  1      -10.0°C            10.0°C            60.00 secs  1 mins 0 secs
Green  0.0 dB  0.78 e/ADU  9.40 arcsec  1      -10.0°C            10.0°C            60.00 secs  1 mins 0 secs
Ha     10.0 dB 0.25 e/ADU  11.72 arcsec 1      -10.0°C            10.0°C            600.00 secs 10 mins 0 secs
Lum    0.0 dB  0.78 e/ADU  11.81 arcsec 1      -10.0°C            10.0°C            60.00 secs  1 mins 0 secs
OIII   10.0 dB 0.25 e/ADU  11.11 arcsec 1      -10.0°C            10.0°C            600.00 secs 10 mins 0 secs
Red    0.0 dB  0.78 e/ADU  10.08 arcsec 1      -10.0°C            10.0°C            60.00 secs  1 mins 0 secs
SII    10.0 dB 0.25 e/ADU  9.44 arcsec  1      -10.0°C            10.0°C            600.00 secs 10 mins 0 secs

Exposure time for M51 Panel 1: 0 hrs 34 mins 0 secs

Target: M51 Panel 2

Filter  Gain  Egain      Mean FWHM  Frames  Sensor Temperature  Mean Temperature  Exposure  Total Exposure
Blue   0.0 dB  0.78 e/ADU  8.86 arcsec  1      -10.0°C            10.0°C            60.00 secs  1 mins 0 secs
Green  0.0 dB  0.78 e/ADU  11.48 arcsec 1      -10.0°C            10.0°C            60.00 secs  1 mins 0 secs
Ha     10.0 dB 0.25 e/ADU  10.67 arcsec 1      -10.0°C            10.0°C            600.00 secs 10 mins 0 secs
Lum    0.0 dB  0.78 e/ADU  12.51 arcsec 1      -10.0°C            10.0°C            60.00 secs  1 mins 0 secs
OIII   10.0 dB 0.25 e/ADU  7.44 arcsec  1      -10.0°C            10.0°C            600.00 secs 10 mins 0 secs
Red    0.0 dB  0.78 e/ADU  7.95 arcsec  1      -10.0°C            10.0°C            60.00 secs  1 mins 0 secs
SII    10.0 dB 0.25 e/ADU  8.76 arcsec  1      -10.0°C            10.0°C            600.00 secs 10 mins 0 secs

Exposure time for M51 Panel 2: 0 hrs 34 mins 0 secs

Total LIGHT Exposure Time: 1 hrs 8 mins 0 secs

MASTERFLATS:

Filter  Frames  Gain  Egain      Exposure  Total Exposure
Blue   100     0.0 dB  0.78 e/ADU  1.65 secs  2 mins 45 secs
Green  100     0.0 dB  0.78 e/ADU  1.65 secs  2 mins 45 secs
Ha     100    10.0 dB  0.25 e/ADU  1.65 secs  2 mins 45 secs
Lum    100     0.0 dB  0.78 e/ADU  1.65 secs  2 mins 45 secs
OIII   100    10.0 dB  0.25 e/ADU  1.65 secs  2 mins 45 secs
Red    100     0.0 dB  0.78 e/ADU  1.65 secs  2 mins 45 secs
SII    100    10.0 dB  0.25 e/ADU  1.65 secs  2 mins 45 secs

Total MASTERFLAT Exposure Time: 0 hrs 19 mins 15 secs

MASTERBIASES:

Filter  Frames  Gain  Egain      Exposure  Total Exposure
No filter 100     0.0 dB  0.78 e/ADU  1.65 secs  2 mins 45 secs
No filter 100    10.0 dB  0.25 e/ADU  1.65 secs  2 mins 45 secs

Total MASTERBIAS Exposure Time: 0 hrs 5 mins 30 secs

MASTERDARKS:

Filter  Frames  Gain  Egain      Sensor Temperature  Exposure  Total Exposure
No filter 100     0.0 dB  0.78 e/ADU  -10.0°C            1.65 secs  2 mins 45 secs
No filter 100    10.0 dB  0.25 e/ADU  -10.0°C            1.65 secs  2 mins 45 secs

Total MASTERDARK Exposure Time: 0 hrs 5 mins 30 secs

Total number of images processed: 68
```

Site 2

Target: M51 2 Panel Mosaic

Site: 천주교학산교회, 54-41, 서산로, 학산면, 영동군, 충청북도, 29166, 대한민국

Latitude: 36.0976°, Longitude: 127.6844°

Bortle scale: 4.0, SQM: 21.12 mag/arcsec²

Equipment used:

Telescope : Tele Vue-85

Camera : ZWO ASI1600GT

Filterwheel : QHYCFW3-L

Focuser : Starlight Instruments FocusLynx Focuser

Rotator : Primaluce ARCO 2

Capture software : N.I.N.A. 2.3.2.9001 (x64)

: PixInsight 1.8.9-2

Observation period:

Start date : 2023-09-11

End date : 2023-09-22

Number of days : 11

Observation sessions : 6

Min temperature : 10.0°C

Max temperature : 10.0°C

Mean temperature : 10.0°C

LIGHTS:

Target: M51 Panel 1

Filter	Gain	Egain	Mean FWHM	Frames	Sensor Temperature	Mean Temperature	Exposure	Total Exposure
Blue	0.0 dB	0.78 e/ADU	10.27 arcsec	1	-10.0°C	10.0°C	60.00 secs	1 mins 0 secs
Green	0.0 dB	0.78 e/ADU	12.71 arcsec	1	-10.0°C	10.0°C	60.00 secs	1 mins 0 secs
Ha	10.0 dB	0.25 e/ADU	8.40 arcsec	7	-10.0°C	10.0°C	600.00 secs	1 hrs 10 mins 0 secs
Lum	0.0 dB	0.78 e/ADU	6.50 arcsec	1	-10.0°C	10.0°C	60.00 secs	1 mins 0 secs
OIII	10.0 dB	0.25 e/ADU	8.45 arcsec	1	-10.0°C	10.0°C	600.00 secs	10 mins 0 secs
Red	0.0 dB	0.78 e/ADU	7.92 arcsec	1	-10.0°C	10.0°C	60.00 secs	1 mins 0 secs
SII	10.0 dB	0.25 e/ADU	12.61 arcsec	1	-10.0°C	10.0°C	600.00 secs	10 mins 0 secs

Exposure time for M51 Panel 1: 1 hrs 34 mins 0 secs

Target: M51 Panel 2

Filter	Gain	Egain	Mean FWHM	Frames	Sensor Temperature	Mean Temperature	Exposure	Total Exposure
Blue	0.0 dB	0.78 e/ADU	10.66 arcsec	1	-10.0°C	10.0°C	60.00 secs	1 mins 0 secs
Green	0.0 dB	0.78 e/ADU	11.34 arcsec	1	-10.0°C	10.0°C	60.00 secs	1 mins 0 secs
Ha	10.0 dB	0.25 e/ADU	7.94 arcsec	7	-10.0°C	10.0°C	600.00 secs	1 hrs 10 mins 0 secs
Lum	0.0 dB	0.78 e/ADU	6.59 arcsec	1	-10.0°C	10.0°C	60.00 secs	1 mins 0 secs
OIII	10.0 dB	0.25 e/ADU	5.18 arcsec	1	-10.0°C	10.0°C	600.00 secs	10 mins 0 secs
Red	0.0 dB	0.78 e/ADU	5.21 arcsec	1	-10.0°C	10.0°C	60.00 secs	1 mins 0 secs
SII	10.0 dB	0.25 e/ADU	11.09 arcsec	1	-10.0°C	10.0°C	600.00 secs	10 mins 0 secs

Exposure time for M51 Panel 2: 1 hrs 34 mins 0 secs

Total LIGHT Exposure Time: 3 hrs 8 mins 0 secs

MASTERFLATS:

Filter	Frames	Gain	Egain	Exposure	Total Exposure
Blue	100	0.0 dB	0.78 e/ADU	1.65 secs	2 mins 45 secs
Green	100	0.0 dB	0.78 e/ADU	1.65 secs	2 mins 45 secs
Ha	700	10.0 dB	0.25 e/ADU	1.65 secs	19 mins 15 secs
Lum	100	0.0 dB	0.78 e/ADU	1.65 secs	2 mins 45 secs
OIII	100	10.0 dB	0.25 e/ADU	1.65 secs	2 mins 45 secs
Red	100	0.0 dB	0.78 e/ADU	1.65 secs	2 mins 45 secs
SII	100	10.0 dB	0.25 e/ADU	1.65 secs	2 mins 45 secs

Total MASTERFLAT Exposure Time: 0 hrs 35 mins 45 secs

MASTERBIASES:

Filter	Frames	Gain	Egain	Exposure	Total Exposure
No filter	100	0.0 dB	0.78 e/ADU	1.65 secs	2 mins 45 secs
No filter	100	10.0 dB	0.25 e/ADU	1.65 secs	2 mins 45 secs

Total MASTERBIAS Exposure Time: 0 hrs 5 mins 30 secs

MASTERDARKS:

Filter	Frames	Gain	Egain	Sensor Temperature	Exposure	Total Exposure
No filter	100	0.0 dB	0.78 e/ADU	-10.0°C	1.65 secs	2 mins 45 secs
No filter	100	10.0 dB	0.25 e/ADU	-10.0°C	1.65 secs	2 mins 45 secs

Total MASTERDARK Exposure Time: 0 hrs 5 mins 30 secs

Total number of images processed: 68

M51_acquisition.csv

date	filter	number	duration	binning	gain	sensorTemperature	fNumber	darks	flats	flatDarks	bias	bortle	meanSqm	meanFwhm	temperature
2023-09-11	4637	1	60.0	1	0	-10	7.5	100	100	0	100	4.0	20.69	8.86	10.0
2023-09-11	4844	1	600.0	1	100	-10	7.5	100	100	0	100	4.0	20.69	8.76	10.0
2023-09-13	4649	1	60.0	1	0	-10	7.5	100	100	0	100	4.0	20.69	10.08	10.0
2023-09-15	2906	1	60.0	1	0	-10	7.5	100	100	0	100	4.0	20.69	12.51	10.0
2023-09-16	4643	1	60.0	1	0	-10	7.5	100	100	0	100	4.0	20.69	11.48	10.0
2023-09-16	2906	1	60.0	1	0	-10	7.5	100	100	0	100	4.0	20.69	11.81	10.0
2023-09-16	4844	1	600.0	1	100	-10	7.5	100	100	0	100	4.0	20.69	9.44	10.0
2023-09-17	4752	1	600.0	1	100	-10	7.5	100	100	0	100	4.0	20.69	11.11	10.0
2023-09-19	4663	1	600.0	1	100	-10	7.5	100	100	0	100	4.0	20.69	11.72	10.0
2023-09-19	4649	1	60.0	1	0	-10	7.5	100	100	0	100	4.0	20.69	7.95	10.0
2023-09-21	4752	1	600.0	1	100	-10	7.5	100	100	0	100	4.0	20.69	7.44	10.0
2023-09-22	4643	1	60.0	1	0	-10	7.5	100	100	0	100	4.0	20.69	9.40	10.0
2023-09-23	4637	1	60.0	1	0	-10	7.5	100	100	0	100	4.0	20.69	7.56	10.0
2023-09-23	4663	1	600.0	1	100	-10	7.5	100	100	0	100	4.0	20.69	10.67	10.0
2023-09-11	4637	1	60.0	1	0	-10	7.0	100	100	0	100	4.0	21.12	10.66	10.0
2023-09-11	2906	1	60.0	1	0	-10	7.0	100	100	0	100	4.0	21.12	6.50	10.0
2023-09-11	4649	1	60.0	1	0	-10	7.0	100	100	0	100	4.0	21.12	5.21	10.0
2023-09-12	4663	1	600.0	1	100	-10	7.0	100	100	0	100	4.0	21.12	6.78	10.0
2023-09-13	4663	2	600.0	1	100	-10	7.0	100	100	0	100	4.0	21.12	6.92	10.0
2023-09-13	4752	1	600.0	1	100	-10	7.0	100	100	0	100	4.0	21.12	8.45	10.0
2023-09-13	4844	1	600.0	1	100	-10	7.0	100	100	0	100	4.0	21.12	12.61	10.0
2023-09-14	4663	2	600.0	1	100	-10	7.0	100	100	0	100	4.0	21.12	8.08	10.0
2023-09-15	4663	1	600.0	1	100	-10	7.0	100	100	0	100	4.0	21.12	9.53	10.0
2023-09-16	4663	3	600.0	1	100	-10	7.0	100	100	0	100	4.0	21.12	9.42	10.0
2023-09-17	4643	1	60.0	1	0	-10	7.0	100	100	0	100	4.0	21.12	11.34	10.0
2023-09-18	4643	1	60.0	1	0	-10	7.0	100	100	0	100	4.0	21.12	12.71	10.0
2023-09-18	4663	1	600.0	1	100	-10	7.0	100	100	0	100	4.0	21.12	10.63	10.0
2023-09-19	4663	2	600.0	1	100	-10	7.0	100	100	0	100	4.0	21.12	7.32	10.0
2023-09-20	4637	1	60.0	1	0	-10	7.0	100	100	0	100	4.0	21.12	10.27	10.0
2023-09-20	2906	1	60.0	1	0	-10	7.0	100	100	0	100	4.0	21.12	6.59	10.0
2023-09-20	4649	1	60.0	1	0	-10	7.0	100	100	0	100	4.0	21.12	7.92	10.0
2023-09-20	4844	1	600.0	1	100	-10	7.0	100	100	0	100	4.0	21.12	11.09	10.0
2023-09-21	4752	1	600.0	1	100	-10	7.0	100	100	0	100	4.0	21.12	5.18	10.0
2023-09-22	4663	1	600.0	1	100	-10	7.0	100	100	0	100	4.0	21.12	9.18	10.0
2023-09-22	4663	1	600.0	1	100	-10	7.0	100	100	0	100	4.0	21.12	5.89	10.0

Example 3: AstroBin.csv output

Example 4: WBPP two-panel mosaic

California Nebula (NGC1499)

File Edit View Go Bookmarks Help

< > ^

/home/steve/Desktop/Current PixInsight Projects/California Nebula (NGC1499)

⌂ 🔍 ⌵ ⌶ ⌷

My Computer

Home

Desktop

Documents

Music

Pictures

calibrated

cosmetized

fastIntegration

logs

master

5 items, Free space: 932.2 GB

calibrated

File Edit View Go Bookmarks Help

< > ^

/home/steve/Desktop/Current PixInsight Projects/California Nebula (NGC1499)/calibrated

⌂ 🔍 ⌵ ⌶ ⌷

My Computer

Home

Desktop

Documents

Music

Pictures

Videos

Downloads


Recent

File System

Rubbish Bin

Name	Size	Type	Date Modified
Light_BIN-1_9576x6388_EXPOSURE-60.00s_FILTER-Blue_mono_GAIN-0_PANEL-1	30 items	Folder	Wed 10 Jan 2024 08:49:33 GMT
Light_BIN-1_9576x6388_EXPOSURE-60.00s_FILTER-Blue_mono_GAIN-0_PANEL-2	30 items	Folder	Wed 10 Jan 2024 08:49:55 GMT
Light_BIN-1_9576x6388_EXPOSURE-60.00s_FILTER-Green_mono_GAIN-0_PANEL-1	30 items	Folder	Wed 10 Jan 2024 08:50:41 GMT
Light_BIN-1_9576x6388_EXPOSURE-60.00s_FILTER-Green_mono_GAIN-0_PANEL-2	30 items	Folder	Wed 10 Jan 2024 08:51:04 GMT
Light_BIN-1_9576x6388_EXPOSURE-60.00s_FILTER-Red_mono_GAIN-0_PANEL-1	25 items	Folder	Wed 10 Jan 2024 08:51:47 GMT
Light_BIN-1_9576x6388_EXPOSURE-60.00s_FILTER-Red_mono_GAIN-0_PANEL-2	30 items	Folder	Wed 10 Jan 2024 08:52:17 GMT
Light_BIN-1_9576x6388_EXPOSURE-600.00s_FILTER-Ha_mono_GAIN-100_PANEL-1	66 items	Folder	Wed 10 Jan 2024 08:43:53 GMT
Light_BIN-1_9576x6388_EXPOSURE-600.00s_FILTER-Ha_mono_GAIN-100_PANEL-2	66 items	Folder	Wed 10 Jan 2024 08:42:49 GMT
Light_BIN-1_9576x6388_EXPOSURE-600.00s_FILTER-OIII_mono_GAIN-100_PANEL-1	84 items	Folder	Wed 10 Jan 2024 08:45:27 GMT
Light_BIN-1_9576x6388_EXPOSURE-600.00s_FILTER-OIII_mono_GAIN-100_PANEL-2	65 items	Folder	Wed 10 Jan 2024 08:46:44 GMT
Light_BIN-1_9576x6388_EXPOSURE-600.00s_FILTER-SII_mono_GAIN-100_PANEL-1	67 items	Folder	Wed 10 Jan 2024 08:47:49 GMT
Light_BIN-1_9576x6388_EXPOSURE-600.00s_FILTER-SII_mono_GAIN-100_PANEL-2	57 items	Folder	Wed 10 Jan 2024 08:48:48 GMT

12 items, Free space: 932.2 GB



Example 4: Directory structure

```
python3 AstroBinUpload.py "/home/steve/Desktop/Current PixInsight
Projects/California Nebula (NGC1499)"
```

Example 4: Script calling syntax

```
Observation session summary
Generated 2024-01-31 09:42:21

Target: NGC 1499 2 Panel Mosaic

Site: Norton Close, Papworth Everard, South Cambridgeshire, Cambridgeshire, Cambridgeshire and Peterborough, England, CB23 3XT, United Kingdom
Latitude: 52.2485°, Longitude: -0.1231°
Bortle scale: 4.0, SQM: 20.52 mag/arcsec²

Equipment used:
Telescope      : NP101is
Camera         : ZWO ASI6200MM PRO
Filterwheel    : Starlight Xpress Filter Wheels
Focuser       : FocusLynx Focuser 1
Capture software : N.I.N.A. 3.0.0.2003 (x64)
               : PixInsight 1.8.9-2
               : N.I.N.A. 2.3.2.9001 (x64)

Observation period:
Start date      : 2023-11-14
End date       : 2024-01-10
Number of days  : 57
Observation sessions : 10
Min temperature : -3.9°C
Max temperature : 10.5°C
Mean temperature : 1.2°C

LIGHTS:
Target: NGC 1499 Panel 1

Filter  Gain    Egain    Mean FWHM  Frames  Sensor Temperature  Mean Temperature  Exposure  Total Exposure
Blue   0.0 dB    0.78 e/ADU  4.60 arcsec  30      -10.0°C            -3.7°C            60.00 secs  30 mins 0 secs
Green  0.0 dB    0.78 e/ADU  4.66 arcsec  30      -10.0°C            -3.7°C            60.00 secs  30 mins 0 secs
Ha     10.0 dB   0.25 e/ADU  4.74 arcsec  66      -10.0°C            2.4°C            600.00 secs  11 hrs 0 mins 0 secs
OIII   10.0 dB   0.25 e/ADU  5.52 arcsec  84      -10.0°C            2.1°C            600.00 secs  14 hrs 0 mins 0 secs
Red    0.0 dB    0.78 e/ADU  4.79 arcsec  25      -10.0°C            -3.5°C            60.00 secs  25 mins 0 secs
SII    10.0 dB   0.25 e/ADU  4.94 arcsec  67      -10.0°C            3.7°C            600.00 secs  11 hrs 10 mins 0 secs

Exposure time for NGC 1499 Panel 1: 37 hrs 35 mins 0 secs

Target: NGC 1499 Panel 2

Filter  Gain    Egain    Mean FWHM  Frames  Sensor Temperature  Mean Temperature  Exposure  Total Exposure
Blue   0.0 dB    0.78 e/ADU  5.17 arcsec  30      -10.0°C            -3.2°C            60.00 secs  30 mins 0 secs
Green  0.0 dB    0.78 e/ADU  4.93 arcsec  30      -10.0°C            -3.2°C            60.00 secs  30 mins 0 secs
Ha     10.0 dB   0.25 e/ADU  4.88 arcsec  66      -10.0°C            -1.0°C            600.00 secs  11 hrs 0 mins 0 secs
OIII   10.0 dB   0.25 e/ADU  4.80 arcsec  65      -10.0°C            1.7°C            600.00 secs  10 hrs 50 mins 0 secs
Red    0.0 dB    0.78 e/ADU  4.97 arcsec  30      -10.0°C            -3.3°C            60.00 secs  30 mins 0 secs
SII    10.0 dB   0.25 e/ADU  5.11 arcsec  57      -10.0°C            5.2°C            600.00 secs  9 hrs 30 mins 0 secs

Exposure time for NGC 1499 Panel 2: 32 hrs 50 mins 0 secs

Total LIGHT Exposure Time: 70 hrs 25 mins 0 secs

MASTERFLATS:

Filter  Frames  Gain    Egain    Exposure  Total Exposure
Blue   100     0.0 dB  0.78 e/ADU  0.22 secs  22 secs
Green  100     0.0 dB  0.78 e/ADU  0.17 secs  17 secs
Ha     100     10.0 dB 0.25 e/ADU  0.47 secs  47 secs
Lum    100     0.0 dB  0.78 e/ADU  0.05 secs  5 secs
OIII   100     10.0 dB 0.25 e/ADU  0.55 secs  55 secs
Red    100     0.0 dB  0.78 e/ADU  0.10 secs  10 secs
SII    100     10.0 dB 0.25 e/ADU  0.58 secs  57 secs

Total MASTERFLAT Exposure Time: 0 hrs 3 mins 34 secs

MASTERBIASES:

Filter  Frames  Gain    Egain    Exposure  Total Exposure
No Filter 100     0.0 dB  0.78 e/ADU  0.00 secs  0 secs
No Filter 100     10.0 dB 0.25 e/ADU  0.00 secs  0 secs

Total MASTERBIAS Exposure Time: 0 hrs 0 mins 0 secs

MASTERDARKS:

Filter  Frames  Gain    Egain    Sensor Temperature  Exposure  Total Exposure
No Filter 50     0.0 dB  0.78 e/ADU  -10.0°C            60.00 secs  50 mins 0 secs
No Filter 50     10.0 dB 0.25 e/ADU  -10.0°C            600.00 secs  8 hrs 20 mins 0 secs

Total MASTERDARK Exposure Time: 9 hrs 10 mins 0 secs

Total number of images processed: 660
```

Example 4: Summary Output

```
California_Nebula_(NGC1499)_acquisition.csv

date filter number duration binning gain sensorCooling fNumber darks flats flatDarks bias bortle meanSqm meanFwhm temperature
2023-11-14 4663 18 600.0 1 100 -10 5.4 50 100 0 100 4.0 20.52 4.76 5.86
2023-11-15 4663 41 600.0 1 100 -10 5.4 50 100 0 100 4.0 20.52 4.78 5.22
2023-11-15 4844 10 600.0 1 100 -10 5.4 50 100 0 100 4.0 20.52 4.96 3.31
2023-11-16 4844 3 600.0 1 100 -10 5.4 50 100 0 100 4.0 20.52 4.89 3.17
2023-11-18 4844 6 600.0 1 100 -10 5.4 50 100 0 100 4.0 20.52 5.09 9.35
2023-11-19 4844 24 600.0 1 100 -10 5.4 50 100 0 100 4.0 20.52 4.91 8.70
2023-11-24 4663 11 600.0 1 100 -10 5.4 50 100 0 100 4.0 20.52 4.76 -0.34
2023-11-24 4844 13 600.0 1 100 -10 5.4 50 100 0 100 4.0 20.52 4.97 0.17
2023-11-25 4663 35 600.0 1 100 -10 5.4 50 100 0 100 4.0 20.52 4.86 -1.59
2023-11-30 4637 30 60.0 1 0 -10 5.4 50 100 0 100 4.0 20.52 4.60 -3.71
2023-11-30 4643 30 60.0 1 0 -10 5.4 50 100 0 100 4.0 20.52 4.66 -3.73
2023-11-30 4663 7 600.0 1 100 -10 5.4 50 100 0 100 4.0 20.52 4.69 -3.76
2023-11-30 4663 4 600.0 1 100 -10 5.4 50 100 0 100 4.0 20.52 4.75 -3.38
2023-11-30 4649 25 60.0 1 0 -10 5.4 50 100 0 100 4.0 20.52 4.79 -3.54
2023-11-30 4844 11 600.0 1 100 -10 5.4 50 100 0 100 4.0 20.52 4.83 -2.75
2023-12-01 4663 5 600.0 1 100 -10 5.4 50 100 0 100 4.0 20.52 4.75 -3.78
2023-12-05 4752 4 600.0 1 100 -10 5.4 50 100 0 100 4.0 20.52 4.88 1.38
2023-12-06 4752 31 600.0 1 100 -10 5.4 50 100 0 100 4.0 20.52 4.83 -1.98
2023-12-09 4752 15 600.0 1 100 -10 5.4 50 100 0 100 4.0 20.52 4.97 6.00
2023-12-10 4663 10 600.0 1 100 -10 5.4 50 100 0 100 4.0 20.52 5.38 5.74
2023-12-10 4752 2 600.0 1 100 -10 5.4 50 100 0 100 4.0 20.52 5.00 6.50
2023-12-10 4844 30 600.0 1 100 -10 5.4 50 100 0 100 4.0 20.52 5.11 5.87
2023-12-11 4844 8 600.0 1 100 -10 5.4 50 100 0 100 4.0 20.52 5.07 7.09
2023-12-19 4752 2 600.0 1 100 -10 5.4 50 100 0 100 4.0 20.52 4.89 2.50
2023-12-19 4844 19 600.0 1 100 -10 5.4 50 100 0 100 4.0 20.52 5.15 2.51
2023-12-20 4752 16 600.0 1 100 -10 5.4 50 100 0 100 4.0 20.52 5.01 2.21
2023-12-29 4752 21 600.0 1 100 -10 5.4 50 100 0 100 4.0 20.52 4.67 2.83
2024-01-02 4752 6 600.0 1 100 -10 5.4 50 100 0 100 4.0 20.52 4.66 5.62
2024-01-03 4752 17 600.0 1 100 -10 5.4 50 100 0 100 4.0 20.52 4.64 5.62
2024-01-09 4663 1 600.0 1 100 -10 5.4 50 100 0 100 4.0 20.52 4.80 -2.40
2024-01-09 4752 11 600.0 1 100 -10 5.4 50 100 0 100 4.0 20.52 8.74 -2.64
2024-01-09 4752 14 600.0 1 100 -10 5.4 50 100 0 100 4.0 20.52 4.75 -2.59
2024-01-10 4637 30 60.0 1 0 -10 5.4 50 100 0 100 4.0 20.52 5.17 -3.22
2024-01-10 4643 30 60.0 1 0 -10 5.4 50 100 0 100 4.0 20.52 4.93 -3.20
2024-01-10 4752 10 600.0 1 100 -10 5.4 50 100 0 100 4.0 20.52 4.83 -2.97
2024-01-10 4649 30 60.0 1 0 -10 5.4 50 100 0 100 4.0 20.52 4.97 -3.27

Processing summary exported to California_Nebula_(NGC1499)_session_summary.txt
AstroBin data exported to California_Nebula_(NGC1499)_acquisition.csv
Processing completed.
```

Example 4: AstroBin.csv output

References

AstroBin's Acquisition CSV File Format

This section details the required data fields for AstroBin's `acquisition.csv` dialogue.

AstroBin Long Exposure Acquisition Fields

Field	Description	Validation
date	The date when the acquisition took place	YYYY-MM-DD format
filter	Filter used	Numeric ID of a valid filter (found in the URL of the filter's page in the equipment database)
number*	Number of frames	Whole number
duration*	Duration of each frame in seconds	Number, Max decimals: 4, Min value: 0.0001, Max value: 999999.9999
iso	ISO setting on the camera	Whole number
binning	Binning of pixels	One of [1, 2, 3, 4]
gain	Gain setting on the camera	Number, Max decimals: 2
sensorCooling	The temperature of the chip in Celsius degrees, e.g., -20	Whole number, Min value: -274, Max value: 100
fNumber	If a camera lens was used, specify the f-number used for this acquisition session	Number, Max decimals: 2, Min value: 0
darks	The number of dark frames	Whole number, Min value: 0
flats	The number of flat frames	Whole number, Min value: 0
flatDarks	The number of flat dark frames	Whole number, Min value: 0
bias	The number of bias/offset frames	Whole number, Min value: 0
bortle	Bortle dark-sky scale	Whole number, Min value: 1, Max value: 9
meanSqm	Mean SQM mag/arcsec^2 as measured by a Sky Quality Meter	Number, Max decimals: 2, Min value: 0
meanFwhm	Mean Full Width at Half Maximum in arc seconds, a measure of seeing	Number, Max decimals: 2, Min value: 0
temperature	Ambient temperature in Celsius degrees	Number, Max decimals: 2, Min value: -88, Max value: 58

Astrobin Filter-Code mappings

The [filters] section of the config.ini file defines the mapping from the filter name to AstroBin's filter code.

The contents of my [filters] section is given below. It shows the names my Astronomik 2 inch filters, as generated by N.I.N.A, and their corresponding AstroBin codes:

[filters]

Filter	Code
Ha	4663
SII	4844
OIII	4752
Red	4649
Green	4643
Blue	4637
Lum	2906
CLS	4061

This is the default filter table in the config.ini. You should this section so that it reflects the filters you use. The filter names should match the names the image capture software generates for your filters.

Finding AstroBin's Numeric ID for Filters

The numeric ID of a filter can be found by examining the URL of the filter's page in the [AstroBin equipment database](#).

For example, consider a [2-inch H-alpha CCD 6nm filter from Astronomik](#). By using [AstroBin's filter explorer](#) to navigate to this filter's page the URL is found to be :

<https://app.astrobin.com/equipment/explorer/filter/4663/astronomik-h-alpha-ccd-6nm-2>

From this URL, the AstroBin code for this Astronomik 2-inch H-alpha CCD 6nm filter is 4663.

Accessing sky quality data

The artificial_brightness of the sky at a given latitude and longitude is obtained from the excellent web resource <https://www.lightpollutionmap.info>. This can be by done by visiting the website and entering the latitude and longitude of the observation site and obtaining the parameters Bortle and SQM. These parameters can then be entered into the [defaults] section of the cconfig.ini file. It can also be done programmatically by the code. To do this you need to place an API_KEY and API_ENDPOINT for the service in the secret.csv file. The only API_ENDPOINT supported currently is <https://www.lightpollutionmap.info/QueryRaster/>. You will have to apply to Jurij Stare, the website owner, for an API key. Jurij's email address is starej@t-2.net. The approach I suggest is: donate a small amount in support of his website. He will send you a thank you e-mail and in response ask for an API key.

[secrets] section format relating to the sky quality API is shown below:

API Key	API Endpoint
*****	https://www.lightpollutionmap.info/QueryRaster/

Reverse Geocoding

The script process the latitude and longitudes found in the header files and uses the [Geopy Nomintim library](#), which in turn, uses the [Open Street Map](#) API. As a courtesy the script provides the email address of the user accessing the API. To enable this the [secrets][EMAIL_ADDRESS] is used and should be set to your email address. Reverse geocoding is required to be able to produce accurate summary information with multi-site data. Without it all data is aggregated under the default site. It does not affect the Astrobin.csv output as this is aggregated across sites for any target.

[secrets] section format relating reverse geocoding API is shown below:

API Key	API Endpoint
EMAIL_ADDRESS	id@provider.com

Set the EMAIL_ADDRESS to your email. If the API call fails then the [default] site information is used

FWHM Values

The AstroBin Long Exposure Acquisition Fields has an entry for meanFwhm. This is not directly available from the header file. But N.I.N.A allows for the mean HFR value of an image to be embedded in the image file name. An example of my file naming convention, with HFR embedded, is given below:

```
'NGC 7822_Panel_1_Date_2023-09-02_Time_21-09-01_Filter_Ha_Exposure_600.00s_HFR_1.64px_FrameNo_0002.fits'
```

The code will look for the keyword HFR in the image file name. If found it will extract the HFR value and assign it to a variable HFR. As HFR is given in pixels, the script calculates the FWHM from the telescope information held in the FITS header. In particular XPIXSZ the x pixel size in microns and FOCALLEN the telescope focal length in mm.

$$\text{IMSCALE} = \text{XPIXSZ} / \text{FOCALLEN} * 206.265$$

$$\text{FWHM} = 2 * \text{hfr} * \text{imscale}$$

The calculations above assume that all stars are circular making FWHM a scaled version of HFR. This is a reasonable approximation as the code averages HFR across all images taken on a particular date with a given filter and gain and then AstroBin further averages HFR across all entries in the uploaded CSV file. Where HFR is not available in the filename it is obtained from [defaults][HFR] in the config.ini file.

Data Sources

The script was developed to work with the following sources of image files

1. Night Time Imaging N' Astronomy ([N.I.N.A](#))

2. Sequence Generator Pro ([SGPro](#))
3. [PixInsight](#) for Master calibration frames

FIT, FITS and FTS file headers are accessed in the code using [Astropy's FITS header library](#). To access XISF headers functions were developed based upon the [Pixinsight XISF header specification](#).

Contributing to AstroBinUpload.py Processing Script

This script is intended for educational purposes in the field of astrophotography. It is part of an open-source project and contributions or suggestions for improvements are welcome.

To contribute to this project, follow these steps:

1. Fork this repository.
2. Create a branch: `git checkout -b <branch_name>`.
3. Make your changes and commit them: `git commit -m '<commit_message>'`.
4. Push to the original branch: `git push origin <project_name>/<location>`.
5. Create the pull request.

Alternatively, see the GitHub documentation on [creating a pull request](#).

Contact

If you want to contact me, you can reach me at sgreaves139@gmail.com.

License

This project uses the following licence: [GNU General Public Licence v3.0](#).