

# BASIC TRAINING IN CHEMISTRY

---

STEVEN HOENIG



# **Basic Training in Chemistry**

*This page intentionally left blank.*

# **Basic Training in Chemistry**

**Steven L. Hoenig**

*Ridgewood, New York*

**Kluwer Academic Publishers**  
New York • Boston • Dordrecht • London • Moscow

eBook ISBN: 0-306-46926-X

Print ISBN: 0-306-46546-9

©2002 Kluwer Academic Publishers  
New York, Boston, Dordrecht, London, Moscow

All rights reserved

No part of this eBook may be reproduced or transmitted in any form or by any means, electronic, mechanical, recording, or otherwise, without written consent from the Publisher

Created in the United States of America

Visit Kluwer Online at: <http://www.kluweronline.com>  
and Kluwer's eBookstore at: <http://www.ebooks.kluweronline.com>

**To Lena and Alan**

*This page intentionally left blank.*

## Preface

This book was written as a quick reference to the many different concepts and ideas encountered in chemistry. Most books these days go into a detailed explanation of one subject and go no further. This is simply an attempt to present briefly some of the various subjects that make up the whole of chemistry. The different subjects covered include general chemistry, inorganic chemistry, organic chemistry, and spectral analysis. The material is brief, but hopefully detailed enough to be of use. Keep in mind that the material is written for a reader who is familiar with the subject of chemistry. It has been the author's intention to present in one ready source several disciplines that are used and referred to often.

This book was written not to be a chemistry text unto itself, but rather as a supplement that can be used repeatedly throughout a course of study and thereafter. This does not preclude it from being used by others that would find it useful as a reference source as well.

Having kept this in mind during its preparation, the material is presented in a manner in which the reader should have some knowledge of the material. Only the basics are stated because a detailed explanation was not the goal but rather to present a number of chemical concepts in one source.

The first chapter deals with material that is commonly covered in almost every first year general chemistry course. The concepts are presented in, I hope, a clear and concise manner. No detailed explanation of the origin of the material or problems are presented. Only that which is needed to understand the concept is stated. If more detailed explanation is needed any general chemistry text would suffice. And if examples are of use, any review book could be used.

The second chapter covers inorganic chemistry. Those most commonly encountered concepts are presented, such as, coordination numbers, crystal systems, and ionic crystals. More detailed explanation of the coordination encountered in bonding of inorganic compounds requires a deeper explanation than this book was intended for.

Chapter three consists mostly of organic reactions listed according to their preparation and reactions. The mechanisms of the various reactions are not discussed since there are numerous texts which are devoted to the subject. A section is devoted to the concept of isomers since any treatment of organic chemistry must include an understanding it. A section on polymer structures is also presented since it is impossible these days not to come across some discussion of it.

The fourth chapter covers instrumental analysis. No attempt is made to explain the inner workings of the different instruments or the mechanisms by which various spectra is produced. The material listed is for the use by those that are familiar with the different type of spectra encountered in the instrumental analysis of chemical compounds. The tables and charts would be useful for the interpretation of various spectra generated in the course of analyzing a chemical substance. Listed are tables that would be useful as for the interpretation of ultra-violet (uv), infra-red (ir), nuclear magnetic resonance (nmr) and mass spectroscopy (ms) spectra.

Chapter five consists of physical constants and unit measurements that are commonly encountered throughout the application of chemistry.

Chapter six contains certain mathematical concepts that are useful to have when reviewing or working with certain concepts encountered in chemistry.

*Steven L. Hoenig*

## **Acknowledgements**

I wish to express my deepest gratitude to Richard Leff, without whose faith and belief in me this book would not have been possible. And I wish to express my thanks to the following Richard Kolodkin, Robert Glatt, Cindy and Harriet Cuccias, and Anthony Woll, who know what the meaning of friendship is.

*This page intentionally left blank.*

# Contents

Acknowledgements	ix
Preface	vii
1. General Chemistry	1
1.1 Atomic Structure and the Periodic Table	1
1.2 Chemical Bonding	18
1.3 Gases	21
1.4 Solutions	23
1.5 Acids and Bases	24
1.6 Thermodynamics	25
1.7 Equilibria	28
1.8 Kinetics	36
2. Inorganic Chemistry	41
2.1 Group IA Elements	42
2.2 Group IIA Elements	43
2.3 Group IIIA Elements	44
2.4 Group IVA Elements	45
2.5 Group VA Elements	45
2.6 Group VIA Elements	46
2.7 Group VIIA Elements	46
2.8 Group VIIIA Elements	47
2.9 Transition Elements	48

2.10 Ionic Solids	49
2.11 Coordination Numbers	49
2.12 Ionic Crystal Systems	50
2.13 Crystal Lattice Packing	51
2.14 Crystal Lattice Types	52
2.15 Crystal Lattice Energy	53
2.16 Complexes	56
3. Organic Chemistry	59
3.1 Classification of Organic Compounds	60
3.2 Alkanes	62
3.3 Alkenes	63
3.4 Dienes	66
3.5 Alkynes	68
3.6 Benzene	69
3.7 Alkylbenzenes	71
3.8 Alkenylbenzenes	74
3.9 Alkyl Halides	75
3.10 Aryl Halides	79
3.11 Alcohols	81
3.12 Phenols	84
3.13 Ethers	89
3.14 Epoxides	91
3.15 Aldehydes and Ketones	92
3.16 Carboxylic Acids	96
3.17 Acid Chlorides	98
3.18 Acid Anhydrides	100
3.19 Esters	102
3.20 Amides	104
3.21 Amines	107
3.22 Alicyclic Compounds	110
3.23 Heterocyclic Compounds	111
3.24 Isomers	115
3.25 Polymer Structures	123
4. Instrumental Analysis	133
4.1 The Electromagnetic Spectrum	133
4.2 Ultraviolet-Visible Spectroscopy	135
4.3 Infrared Spectroscopy	138
4.4 Nuclear Magnetic Resonance	151
4.5 Mass Spectroscopy	163

5.	Units and Measurements	169
5.1	Fundamental Physical Constants	169
5.2	Units	170
5.3	Prefixes	171
5.4	Conversion Factors	172
6.	Mathematical Concepts	173
6.1	Algebraic Formulas	173
6.2	Plane Figure Formulas	175
6.3	Solid Figure Formulas	177
	Index	179

*This page intentionally left blank.*

# Chapter 1

## General Chemistry

- 1.1 *Atomic Structure and the Periodic Table*
- 1.2 *Chemical Bonding*
- 1.3 *Gases*
- 1.4 *Solutions*
- 1.5 *Acids and Bases*
- 1.6 *Thermodynamics*
- 1.7 *Equilibria*
- 1.8 *Kinetics*

### 1.1 Atomic Structure and the Periodic Table

#### 1.1.1 Constituents of the Atom


The atom of any element consists of three basic types of particles ... the electron (a negatively charged particle), the proton (a positively charged particle), and the neutron (a neutrally charged particle). The protons and neutrons occupy the nucleus while the electrons are outside of the nucleus. The protons and neutrons contribute very little to the total volume but account for the majority of the atom's mass. However, the atoms volume is determined by the electrons, which contribute very little to the mass. Table 1.1 summarizes the properties of these three particles.

*Table 1.1. Properties of the Proton, Electron, and Neutron*


Particle	Mass	Electric Charge	Unit Charge
Proton	$1.672 \times 10^{-24}$ g	$+1.602 \times 10^{-19}$ coulomb	+1
Electron	$9.108 \times 10^{-28}$ g	$-1.602 \times 10^{-19}$ coulomb	-1
Neutron	$1.675 \times 10^{-24}$ g	0	0

The **atomic number** (Z) of an element is the number of protons within the nucleus of an atom of that element. In a neutral atom, the number of protons and electrons are equal and the atomic number also indicates the number of electrons.

The **mass number** (A) is the sum of the protons and neutrons present in the atom. The number of neutrons can be determined by (A - Z). The symbol for denoting the atomic number and mass number for an element X is as follows:



Atoms that have the same atomic number (equal number of protons) but different atomic masses (unequal number of neutrons) are referred to as isotopes. For example, carbon consists of two isotopes, carbon-12 and carbon-13:



The **atomic mass unit** (amu) is defined as 1/12 the mass of a carbon-12 isotope. The relative **atomic mass** of an element is the weighted average of the isotopes relative to 1/12 of the carbon-12 isotope. For example, the atomic mass of neon is 20.17 amu and is calculated from the following data: neon-19 (amu of 19.99245, natural abundance of 90.92%), neon-20 (amu of 20.99396, natural abundance of 0.260%) and neon-21 (amu of 21.99139, natural abundance of 8.82%):

$$\begin{aligned} \text{a.m.} \quad \text{neon} &= (19.99245 * 0.9092) + (20.99396 * 0.00260) + (21.99139 * 0.0882) \\ &= 20.17 \text{ amu} \end{aligned}$$

The relative **molecular mass** is the sum of the atomic masses for each atom in the molecule. For  $\text{H}_2\text{SO}_4 = (1 * 2) + 32 + (16 * 4) = 98$ .

The **mole** (mol) is simply a unit of quantity, it represents a certain amount of material, i.e. atoms or molecules. The numerical value of one mole is **6.023 x 10<sup>23</sup>** and is referred to as **Avogadro's number**. The mole is defined as the mass, in grams, equal to the atomic mass of an element or molecule. Therefore, 1 mole of carbon weighs 12 grams and contains **6.023 x 10<sup>23</sup>** carbon atoms. The following formula can be used to find the number of moles:

$$\text{moles} = \frac{\text{mass in grams}}{\text{atomic (or molecular) mass}}$$

### 1.1.2 Quantum Numbers

From quantum mechanics a set of equations called wave equations are obtained. A series of solutions to these equations, called wave functions, gives the four quantum numbers required to describe the placement of the electrons in the hydrogen atom or in other atoms.

The principal quantum number,  $n$ , determines the energy of an orbital and has a value of  $n = 1, 2, 3, 4, \dots$

The angular momentum quantum number,  $\lambda$ , determines the "shape" of the orbital and has a value of 0 to  $(n - 1)$  for every value of  $n$ .

The magnetic quantum number,  $m_\lambda$ , determines the orientation of the orbital in space and has a value of  $-\lambda$  to  $+\lambda$ .

The electron spin quantum number,  $m_s$ , determines the magnetic field generated by the electron and has a value of  $-\frac{1}{2}$  or  $+\frac{1}{2}$ .

Table 1.2. Quantum Numbers and Electron Distribution

Shell	Principal quantum number $n$	Angular momentum quantum number $\lambda$	Orbital designation*	Magnetic quantum number $m_\lambda$	Spin quantum number $m_s$	Total number of electrons per orbital
K	1	0	s	0	$-\frac{1}{2}, +\frac{1}{2}$	2
L	2	0	s	0	$-\frac{1}{2}, +\frac{1}{2}$	2
		1	$p_x$	-1	$-\frac{1}{2}, +\frac{1}{2}$	6
			$p_y$	0	$-\frac{1}{2}, +\frac{1}{2}$	
			$p_z$	+1	$-\frac{1}{2}, +\frac{1}{2}$	
M	3	0	s	0	$-\frac{1}{2}, +\frac{1}{2}$	2
			$p_x$	-1	$-\frac{1}{2}, +\frac{1}{2}$	6
		1	$p_y$	0	$-\frac{1}{2}, +\frac{1}{2}$	
			$p_z$	+1	$-\frac{1}{2}, +\frac{1}{2}$	
			$d_{xy}$	-2	$-\frac{1}{2}, +\frac{1}{2}$	10
			$d_{xz}$	-1	$-\frac{1}{2}, +\frac{1}{2}$	
		2	$d_{yz}$	0	$-\frac{1}{2}, +\frac{1}{2}$	
			$d_{z^2}$	+1	$-\frac{1}{2}, +\frac{1}{2}$	
			$d_{x^2-y^2}$	+2	$-\frac{1}{2}, +\frac{1}{2}$	

The following is a summary in which the quantum numbers are used to fill the atomic orbitals:

1. No two electrons can have the same four quantum numbers. This is the Pauli exclusion principle.
2. Orbitals are filled in the order of increasing energy.
3. Each orbital can only be occupied by a maximum of two electrons and must have different spin quantum numbers (opposite spins).
4. The most stable arrangement of electrons in orbitals is the one that has the greatest number of equal spin quantum numbers (parallel spins). This is Hund's rule.

Also note that the energy of an electron also depends on the angular momentum quantum number as well as the principal quantum number. Therefore the order that the orbitals get filled does not strictly follow the principal quantum number. The order in which orbitals are filled is given in figure 1.1.




Figure 1.1 Filling Order and Relative Energy Levels of Orbitals

### 1.1.3 Atomic Orbitals

The quantum numbers mentioned earlier were obtained as solutions to a set of wave equations. These wave equations cannot tell precisely where an electron is at any given moment or how fast it is moving. But rather it states the probability of finding the electron at a particular place. An orbital is a region of space where the electron is most likely to be found. An orbital has no definite boundary to it, but can be thought of as a cloud with a specific shape. Also, the orbital is not uniform throughout, but rather densest in the region where the probability of finding the electron is highest.

The shape of an orbital represents 90% of the probability of finding the electron within that space. As the quantum numbers change so do the shapes

and direction of the orbitals. Figure 1.2 show the shapes for principal quantum number  $n = 1, 2$ , and  $3$ .




Figure 1.2 Representation of Atomic Orbitals

Another type of orbital is one that originates from the mixing of the different atomic orbitals and is called a hybrid orbital. Hybridization (mixing) of atomic orbitals results in a new set of orbitals with different

shapes and orientations. The orbitals are designated according to which of the separate atomic orbitals have been mixed. For instance, an s orbital mixing with a single p orbital is designated  $sp$ . An s orbital mixing with two separate p orbitals ( $p_x + p_y$  or  $p_x + p_z$  or  $p_y + p_z$ ) is designated  $sp^2$  and a s orbital mixing with three separate p orbitals is designated  $sp^3$ . Combinations of other orbitals can occur as well. Table 1.3 lists some of the possible hybrid orbitals.

Table 1.3. Hybrid Orbitals

Hybrid orbital	Atomic orbitals hybridized	Number of hybrid orbitals	Angle of hybrid orbital	Geometry	Example
sp dp	s + p d + p	2 2	180° 180°	—	CO <sub>2</sub> HgCl <sub>2</sub>
sp <sup>2</sup>	s + $p_x + p_y$	3	120°		PCl <sub>3</sub> SO <sub>3</sub> H <sub>2</sub> CO
sp <sup>3</sup>	s + $p_x + p_y + p_z$	4	109°28'		CH <sub>4</sub> AlCl <sub>4</sub>
dsp <sup>2</sup> d <sup>2</sup> p <sup>2</sup>	1 + px + py + $d_{x^2-y^2}$ $d_{x^2-y^2} + d_{z^2} +$ px + py	4	90° 90°		XeF <sub>4</sub>
dsp <sup>3</sup>	s + $p_x + p_y + p_z +$ $d_{x^2-y^2}$	5	90° 120° 180°		PCl <sub>5</sub> PF <sub>5</sub>
d <sup>2</sup> sp <sup>3</sup>	s + $p_x + p_y + p_z +$ $d_{x^2-y^2} + d_{z^2}$	6	90° 180°		SF <sub>6</sub> XeF <sub>6</sub> MoF <sub>6</sub>

## 1.1.4 Electronic Configuration of the Elements

Table 1.4. Electronic Configuration of the Elements

	Shells	K	L		M			N				O	P	Q						
	Sub-Levels	1s	2s	2p	3s	3p	3d	4s	4p	4d	4f	5s	5p	5d	5f	6s	6p	6d	6f	7s
1	Hydrogen	1																		
2	Helium	2																		
3	Lithium	2		1																
4	Beryllium	2		2																
5	Boron	2		2	1															
6	Carbon	2		2	2															
7	Nitrogen	2		2	3															
8	Oxygen	2		2	4															
9	Fluorine	2		2	5															
10	Neon	2		2	6															
11	Sodium	2		2	6		1													
12	Magnesium	2		2	6		2													
13	Aluminum	2		2	6		2	1												
14	Silicon	2		2	6		2	2												
15	Phosphorus	2		2	6		2	3												
16	Sulfur	2		2	6		2	4												
17	Chlorine	2		2	6		2	5												
18	Argon	2		2	6		2	6												
19	Potassium	2		2	6		2	6			1									
20	Calcium	2		2	6		2	6			2									
21	Scandium	2		2	6		2	6	1		2									
22	Titanium	2		2	6		2	6	2		2									
23	Vanadium	2		2	6		2	6	3		2									
24	Chromium	2		2	6		2	6	5		1									
25	Manganese	2		2	6		2	6	5		2									
26	Iron	2		2	6		2	6	6		2									
27	Cobalt	2		2	6		2	6	7		2									
28	Nickel	2		2	6		2	6	8		2									
29	Copper	2		2	6		2	6	10		1									
30	Zinc	2		2	6		2	6	10		2									
31	Gallium	2		2	6		2	6	10	2	1									
32	Germanium	2		2	6		2	6	10	2	2									
33	Arsenic	2		2	6		2	6	10	2	3									
34	Selenium	2		2	6		2	6	10	2	4									
35	Bromine	2		2	6		2	6	10	2	5									
36	Krypton	2		2	6		2	6	10	2	6									
37	Rubidium	2		2	6		2	6	10	2	6			1						
38	Strontium	2		2	6		2	6	10	2	6			2						
39	Yttrium	2		2	6		2	6	10	2	6	1		2						
40	Zirconium	2		2	6		2	6	10	2	6	2		2						
41	Niobium	2		2	6		2	6	10	2	6	4		1						
42	Molybdenum	2		2	6		2	6	10	2	6	5		1						

Table 1.4. (Continued).

	Shells	K	L		M			N				O			P			Q		
	Sub-Levels	1s	2s	2p	3s	3p	3d	4s	4p	4d	4f	5s	5p	5d	5f	6s	6p	6d	6f	7s
43	Technetium	2	2	6	2	6	10	2	6	6	1									
44	Ruthenium	2	2	6	2	6	10	2	6	7	1									
45	Rhodium	2	2	6	2	6	10	2	6	8	1									
46	Palladium	2	2	6	2	6	10	2	6	10										
47	Silver	2	2	6	2	6	10	2	6	10	1									
48	Cadmium	2	2	6	2	6	10	2	6	10	2									
49	Indium	2	2	6	2	6	10	2	6	10	1	2								
50	Tin	2	2	6	2	6	10	2	6	10	2	2								
51	Antimony	2	2	6	2	6	10	2	6	10	2	3								
52	Tellurium	2	2	6	2	6	10	2	6	10	2	4								
53	Iodine	2	2	6	2	6	10	2	6	10	2	5								
54	Xenon	2	2	6	2	6	10	2	6	10	2	6								
55	Cesium	2	2	6	2	6	10	2	6	10	2	6								1
56	Barium	2	2	6	2	6	10	2	6	10	2	6								2
57	Lanthanum	2	2	6	2	6	10	2	6	10	2	6	1							2
58	Cerium	2	2	6	2	6	10	2	6	10	2	6								2
59	Praseodymium	2	2	6	2	6	10	2	6	10	3	2	6							2
60	Neodymium	2	2	6	2	6	10	2	6	10	4	2	6							2
61	Promethium	2	2	6	2	6	10	2	6	10	5	2	6							2
62	Samarium	2	2	6	2	6	10	2	6	10	6	2	6							2
63	Europium	2	2	6	2	6	10	2	6	10	7	2	6							2
64	Gadolinium	2	2	6	2	6	10	2	6	10	7	2	6	1						2
65	Terbium	2	2	6	2	6	10	2	6	10	9	2	6							2
66	Dysprosium	2	2	6	2	6	10	2	6	10	10	2	6							2
67	Holium	2	2	6	2	6	10	2	6	10	11	2	6							2
68	Erbium	2	2	6	2	6	10	2	6	10	12	2	6							2
69	Thulium	2	2	6	2	6	10	2	6	10	13	2	6							2
70	Ytterbium	2	2	6	2	6	10	2	6	10	14	2	6							2
71	Lutetium	2	2	6	2	6	10	2	6	10	14	2	6	1						2
72	Hafnium	2	2	6	2	6	10	2	6	10	14	2	6	2						2
73	Tantalum	2	2	6	2	6	10	2	6	10	14	2	6	3						2
74	Tungsten	2	2	6	2	6	10	2	6	10	14	2	6	4						2
75	Rhenium	2	2	6	2	6	10	2	6	10	14	2	6	5						2
76	Osmium	2	2	6	2	6	10	2	6	10	14	2	6	6						2
77	Iridium	2	2	6	2	6	10	2	6	10	14	2	6	9						1
78	Platinum	2	2	6	2	6	10	2	6	10	14	2	6	9	1					
79	Gold	2	2	6	2	6	10	2	6	10	14	2	6	10	1					
80	Mercury	2	2	6	2	6	10	2	6	10	14	2	6	10	2					
81	Thallium	2	2	6	2	6	10	2	6	10	14	2	6	10	2	1				
82	Lead	2	2	6	2	6	10	2	6	10	14	2	6	10	2	2				
83	Bismuth	2	2	6	2	6	10	2	6	10	14	2	6	10	2	3				
84	Polonium	2	2	6	2	6	10	2	6	10	14	2	6	10	2	4				
85	Astatine	2	2	6	2	6	10	2	6	10	14	2	6	10	2	5				
86	Radon	2	2	6	2	6	10	2	6	10	14	2	6	10	2	6				

Table 1.4. (Continued)

## 1.1.5 Periodic Table of Elements

**Table 1.5. Periodic Table of Elements**

IA		IIA		VIIIA		VIIIA		VIIIA		VIIIA		VIIIA		
1	H	3	Li	5	B	6	C	7	N	8	O	9	F	
1.008		6.939	9.012	10.811	12.011	14.0067	15.9994	18.998	20.179					
3	Li	11	12	13	14	15	16	17	18	Cl	Ar			
		Na	Mg	Al	Si	P	S							
		22.991	24.312	26.982	28.086	30.974	38.96	35.453	39.948					
19	K	20	Ca	21	Sc	22	Ti	23	V	24	Cr	25	Mn	
39.102	40.08	44.956	47.90	50.942	51.996	54.938	55.847	58.933	58.71	63.546	65.37	69.72	72.59	
37	Rb	38	Sr	39	Y	40	Zr	41	Nb	42	Mo	43	Tc	47
85.47	87.62	88.905	91.22	92.906	96.94	98.906	101.07	102.905	106.4	107.868	112.40	114.82	118.69	121.75
55	Cs	56	Ba	57	La	72	Hf	73	Ta	74	W	75	Re	76
132.905	137.34	138.91	138.91	178.49	180.948	183.85	186.2	190.2	192.2	195.09	196.967	200.59	204.37	207.19
87	Fr	88	Ra	89	Ac	104	Unq	105	Unp	106	Unh	107	Uns	
(223)	226.02	(227)	(227)	(261)	(261)	(262)	(262)	(261)	(263)	(261)				

58	59	60	61	62	63	64	65	66	67	68	69	70	71
Ce	Pr	Nd	Pm	Sm	Eu	Gd	Tb	Dy	Ho	Er	Tm	Yb	Lu
140.12	140.90	144.24	(145)	150.35	151.96	157.25	158.924	162.50	164.930	167.26	168.934	173.04	174.97
90	91	92	93	94	95	96	97	98	99	100	101	102	103
Th	Pa	U	Np	Pu	Am	Cm	Bk	Cr	Es	Fm	Md	No	Lr
232.038	231.03	238.03	237.04	(244)	(243)	(247)	(247)	(251)	(254)	(257)	(258)	(259)	(260)

### 1.1.6 Ionization Energy

Ionization energy is the minimum amount of energy needed to remove an electron from a gasous atom or ion, and is expressed in electron volts (eV).

Going across the periodic table the I.E. increases due to the fact that the principal energy level (principal quantum number) remains the same while the number of electrons increase, thereby enhancing the electrostatic attraction between the protons in the nucleus and the electrons. Going down the table the I.E. decreases because the outer electrons are now further from the nucleus and the protons.

Table 1.6 shows the first ionization energy for the elements.




Table 1.6. First Ionization Energy (in eV)

Z	Element	I.E.	Z	Element	I.E.	Z	Element	I.E.
1	Hydrogen	13.59	30	Zinc	9.39	59	Praseodymium	5.40
2	Helium	24.58	31	Gallium	6.00	60	Neodymium	5.49
3	Lithium	5.39	32	Germanium	7.88	61	Promethium	5.55
4	Beryllium	9.32	33	Arsenic	9.81	62	Samarium	5.61
5	Boron	8.30	34	Selenium	9.75	63	Europium	5.64
6	Carbon	11.26	35	Bromine	11.84	64	Gadolinium	6.26
7	Nitrogen	14.53	36	Krypton	14.00	65	Terbium	5.89
8	Oxygen	13.61	37	Rubidium	4.18	66	Dysprosium	5.82
9	Fluorine	17.42	38	Strontium	5.69	67	Holium	5.89
10	Neon	21.56	39	Yttrium	6.38	68	Erbium	5.95
11	Sodium	5.14	40	Zirconium	6.84	69	Thulium	6.03
12	Magnesium	7.64	41	Niobium	6.88	70	Ytterbium	6.04
13	Aluminum	5.98	42	Molybdenum	7.10	71	Lutetium	5.32
14	Silicon	8.15	43	Technetium	7.28	72	Hafnium	7.00
15	Phosphorus	10.48	44	Ruthenium	7.36	73	Tantalum	7.88
16	Sulfur	10.36	45	Rhodium	7.46	74	Tungsten	7.98
17	Chlorine	12.97	46	Palladium	8.33	75	Rhenium	7.87
18	Argon	15.76	47	Silver	7.57	76	Osmium	8.73
19	Potassium	4.34	48	Cadmium	8.99	77	Iridium	9.1
20	Calcium	6.11	49	Indium	5.79	78	Platinum	8.96
21	Scandium	6.54	50	Tin	7.34	79	Gold	9.22
22	Titanium	6.82	51	Antimony	8.64	80	Mercury	10.43
23	Vanadium	6.74	52	Tellurium	9.01	81	Thallium	6.11
24	Chromium	6.87	53	Iodine	10.45	82	Lead	7.42
25	Manganese	7.43	54	Xenon	12.13	83	Bismuth	7.29
26	Iron	7.87	55	Cesium	3.89	84	Polonium	8.43
27	Cobalt	7.86	56	Barium	5.21	85	Astatine	9.5
28	Nickel	7.63	57	Lanthanum	5.61	86	Radon	10.75
29	Copper	7.72	58	Cerium	6.54			




Figure 1.3. First ionization energy

### 1.1.7 Electronegativity

Electronegativity ( $X$ ) is the relative attraction of an atom for an electron in a covalent bond. But due to the complexity of a covalent bond it is not possible to define precise electronegativity values. Originally the element fluorine, whose atoms have the greatest attraction for an electron, was given an arbitrary value of 4.0. All other electronegativity values are based on this.

Note that the greater the difference in electronegativities the more ionic in nature is the bond, and the smaller the difference the more covalent is the bond.

Going across the periodic table the electronegativity increases because the principal energy level remain the same and the electrostatic attraction increases. The atoms also have a desire to have the most stable configuration which is that of the noble gas configuration. Going down the table the electronegativity decreases due to the increased distance from the nucleus.

Table 1.7 lists relative electronegativities for the elements.

Table 1.7. Relative Electronegativities

Z	Element	X	Z	Element	X	Z	Element	X
1	Hydrogen	2.2	27	Cobalt	1.9	53	Iodine	2.5
2	Helium	---	28	Nickel	1.9	54	Xenon	---
3	Lithium	1.0	29	Copper	1.9	55	Cesium	0.7
4	Beryllium	1.5	30	Zinc	1.6	56	Barium	0.9
5	Boron	2.0	31	Gallium	1.6			
6	Carbon	2.5	32	Germanium	1.8	72	Hafnium	1.3
7	Nitrogen	3.0	33	Arsenic	2.0	73	Tantalum	1.5
8	Oxygen	3.5	34	Selenium	2.4	74	Tungsten	1.7
9	Fluorine	4.0	35	Bromine	2.8	75	Rhenium	1.9
10	Neon	---	36	Krypton	---	76	Osmium	2.2
11	Sodium	0.9	37	Rubidium	0.8	77	Iridium	2.2
12	Magnesium	1.2	38	Strontium	1.0	78	Platinum	2.2
13	Aluminum	1.5	39	Yttrium	1.2	79	Gold	2.4
14	Silicon	2.8	40	Zirconium	1.4	80	Mercury	1.9
15	Phosphorus	2.1	41	Niobium	1.6	81	Thallium	1.8
16	Sulfur	2.5	42	Molybdenum	1.8	82	Lead	1.9
17	Chlorine	3.0	43	Technetium	1.9	83	Bismuth	1.9
18	Argon	---	44	Ruthenium	2.2	84	Polonium	2.0
19	Potassium	0.8	45	Rhodium	2.2	85	Astatine	2.2
20	Calcium	1.0	46	Palladium	2.2	86	Radon	---
21	Scandium	1.3	47	Silver	1.9	87	Francium	0.7
22	Titanium	1.5	48	Cadmium	1.7	88	Radium	0.9
23	Vanadium	1.6	49	Indium	1.7	89	Actinium	1.1
24	Chromium	1.6	50	Tin	1.8	90	Thorium	1.3
25	Manganese	1.5	51	Antimony	1.9	91	Protactinium	1.4
26	Iron	1.8	52	Tellurium	2.1	92	Uranium	1.4




Figure 1.4. Electronegativity

### 1.1.8 Radius of Atoms

The radius of an atom can be estimated by taking half the distance between the nucleus of two of the same atoms. For example, the distance between the nuclei of  $I_2$  is 2.66 Å, half that distance would be the radius of atomic iodine or 1.33 Å. Using this method the atomic radius of nearly all the elements can be estimated.

Note that going across the periodic table, the atomic radius decreases. This is due to the fact that the principal energy level (principal quantum number) remains the same, but the number of electrons increase. The increase in the number of electrons causes an increase in the electrostatic attraction which causes the radius to decrease. However, going down the periodic table the principal energy level increases and hence the atomic radius increases.

Table 1.8 lists the atomic radii of some of the elements.

Table 1.8. Atomic Radii (in Å)

Z	Element	X	Z	Element	X	Z	Element	X
1	Hydrogen	0.37	25	Manganese	1.29	49	Indium	1.62
2	Helium	----	26	Iron	1.26	50	Tin	1.40
3	Lithium	1.52	27	Cobalt	1.26	51	Antimony	1.41
4	Beryllium	1.12	28	Nickel	1.24	52	Tellurium	1.37
5	Boron	0.88	29	Copper	1.28	53	Iodine	1.33
6	Carbon	0.77	30	Zinc	1.33	54	Xenon	----
7	Nitrogen	0.70	31	Gallium	1.22	55	Cesium	2.62
8	Oxygen	0.66	32	Germanium	1.22	56	Barium	2.17
9	Fluorine	0.64	33	Arsenic	1.21			
10	Neon	----	34	Selenium	1.17	72	Hafnium	1.57
11	Sodium	1.86	35	Bromine	1.14	73	Tantalum	1.43
12	Magnesium	1.60	36	Krypton	----	74	Tungsten	1.37
13	Aluminum	1.43	37	Rubidium	2.41	75	Rhenium	1.37
14	Silicon	1.17	38	Strontium	2.15	76	Osmium	1.34
15	Phosphorus	1.10	39	Yttrium	1.80	77	Iridium	1.35
16	Sulfur	1.04	40	Zirconium	1.57	78	Platinum	1.38
17	Chlorine	0.99	41	Niobium	1.43	79	Gold	1.44
18	Argon	----	42	Molybdenum	1.36	80	Mercury	1.50
19	Potassium	2.31	43	Technetium	1.30	81	Thallium	1.71
20	Calcium	1.97	44	Ruthenium	1.33	82	Lead	1.75
21	Scandium	1.60	45	Rhodium	1.34	83	Bismuth	1.46
22	Titanium	1.46	46	Palladium	1.38	84	Polonium	1.40
23	Vanadium	1.31	47	Silver	1.44	85	Astatine	1.40
24	Chromium	1.25	48	Cadmium	1.49	86	Radon	----




Figure 1.5. Atomic Radii

## 1.1.9 Atomic Weights

Table 1.9. Atomic Weights

Name	Symbol	Atomic Wt	Name	Symbol	Atomic Wt
Actinium	Ac	227.0278	Neodymium	Nd	144.24
Aluminum	Al	26.9815	Neon	Ne	20.183
Americium	Am	(243.0614)	Neptunium	Np	237.0482
Antimony	Sb	121.75	Nickel	Ni	58.71
Argon	Ar	39.948	Niobium	Nb	92.906
Arsenic	As	74.9216	Nitrogen	N	14.0067
Astatine	At	(209.9871)	Nobelium	No	(259.1009)
Barium	Ba	137.34	Osmium	Os	190.2
Berkelium	Bk	(247.0703)	Oxygen	O	15.9994
Beryllium	Be	9.0122	Palladium	Pd	106.4
Bismuth	Bi	208.980	Phosphorus	P	30.9738
Boron	B	10.811	Platinum	Pt	195.09
Bromine	B	79.909	Plutonium	Pu	(244.0642)
Cadmium	Cd	12.401	Polonium	Po	(208.9824)
Calcium	Ca	40.08	Potassium	K	39.102
Californium	Cf	(251.0796)	Praseodymium	Pr	140.907
Carbon	C	12.01115	Promethium	Pm	(144.9127)
Cerium	Ce	140.12	Protactinium	Pa	231.0359
Cesium	Cs	132.905	Radium	Ra	226.0254
Chlorine	Cl	35.453	Radon	Rn	(222.0176)
Chromium	Cr	51.996	Rhenium	Re	186.2
Cobalt	Co	58.9332	Rhodium	Rh	102.905
Copper	Cu	3.546	Rubidium	Rb	85.47
Curium	Cm	(247.0703)	Ruthenium	Ru	101.07
Dysprosium	Dy	162.50	Samarium	Sm	150.35
Einsteinium	Es	(252.083)	Scandium	Sc	44.956
Erbium	Er	167.26	Selenium	Se	78.96
Europium	Eu	151.96	Silicon	Si	28.086
Fermium	Fm	(257.0951)	Silver	Ag	107.870
Fluorine	F	18.9984	Sodium	Na	22.9898
Francium	Fr	(223.0197)	Strontium	Sr	87.62
Gadolinium	Gd	157.25	Sulfur	S	32.064
Gallium	Ga	69.72	Tantalum	Ta	180.948
Germanium	Ge	72.59	Technetium	Tc	98.906
Gold	Au	196.967	Tellurium	Te	127.60
Hafnium	Hf	178.49	Terbium	Tb	158.924
Helium	He	4.0026	Thallium	Tl	204.37
Holmium	Ho	164.930	Thorium	Th	232.038
Hydrogen	H	1.00797	Thulium	Tm	168.934
Indium	In	114.82	Tin	Sn	118.69
Iodine	I	126.9044	Titanium	Ti	47.90
Iridium	Ir	192.2	Tungsten	W	183.85
Iron	Fe	55.847	Unnilquadium	Unq	(261.11)

Table 1.9. (Continued)

Krypton	Kr	83.80	Unnilpentium	Unp	(62.114)
Lanthanum	La	183.91	Unnilhexium	Unh	(263.118)
Lawrencium	Lr	(262.11)	Unnilseptium	Uns	(262.12)
Lead	Pb	207.19	Uranium	U	238.03
Lithium	Li	6.939	Vanadium	V	50.942
Lutetium	Lu	174.97	Xenon	Xe	131.30
Magnesium	Mg	24.312	Ytterbium	Yb	173.04
Manganese	Mn	54.9380	Yttrium	Y	88.905
Mendelevium	Md	(258.10)	Zinc	Zn	65.37
Mercury	Hg	200.59	Zirconium	Zr	91.22
Molybdenum	Mo	95.94			

## 1.2 Chemical Bonding


### 1.2.1 Covalent Bonding

A **covalent bond** is a bond in which a pair of electrons is shared between two atoms. Depending on the atoms electronegativity the bond is either polar or non-polar.

A pair of atoms with the same electronegativity would form a **non-polar covalent bond**, such as:




A **polar covalent bond** is one in which the atoms have different electronegativities, such as:




### 1.2.2 Coordinate Covalent Bond (Dative Bond)

A **coordinate covalent bond** is a bond in which both pairs of electrons are donated by one atom and are shared between the two, for example:



### 1.2.3 Ionic Bonding

An **ionic bond** is one in which one or more electrons are transferred from one atom's valence shell (becoming a positively charged ion, called a **cation**) to the other's valence shell (becoming a negatively charged ion, called a **anion**). The resulting electrostatic attraction between oppositely charged ions results in the formation of the ionic bond.



Not all compounds will be either purely covalent or purely ionic, most are somewhere in between. As a rule of thumb, if a compound has less than 50% ionic character it is considered covalent and more than 50%, ionic. The **ionic character** can be related to the difference in electronegativities of the bonded atoms. If the electronegativity difference is 1.7, the bond is about 50% ionic.

### 1.2.4 Dipole-Dipole Bonding

A **dipole-dipole bond** occurs between polar molecules and is a weak electrostatic attraction.




Figure 1.6. Dipole-dipole attraction

### 1.2.5 Ion-Dipole Bonding (Solvation)

An **ion-dipole bond** is another electrostatic attraction between an ion and several polar molecules. When an ionic substance is dissolved in a polar solvent, it is this kind of interaction that takes place. The negative ends of the solvent aligned themselves to the positive charge, and the positive ends aligned with the negative charge. This process is **solvation**. When the solvent is water the process is the same but called **hydration**.




Figure 1.7. Ion-dipole attraction

### 1.2.6 Hydrogen Bonding

When hydrogen is bonded covalently to a small electronegative atom the electron cloud around the hydrogen is drawn to the electronegative atom and a strong dipole is created. The positive end of the dipole approaches close to the negative end of the neighboring dipole and a uniquely strong dipole-dipole bond forms, this is referred to as a **hydrogen bond**.




Figure 1.8. Hydrogen bonding

### 1.2.7 van der Waals

When the electron cloud around an atom or molecule shifts (for whatever reason), a temporary dipole is created, this in turns creates an induced dipole in the next molecule. This induced dipole (**van der Waals**) induces another and so on. The induced dipoles now are electrostatically attracted to each other and a weak induced dipole attraction occurs.




Figure 1.9. van der Waals attraction

## 1.3 Gases

The following laws and equations are for ideal gases. An **ideal gas** is considered to be composed of small particles whose volume is negligible when compared to the whole volume, and the gas particles do not exert any force on one another. Unless otherwise noted:

P - pressure

V - volume

T - absolute temperature (Kelvin)

n - number of moles

R - gas constant

$$8.314 \text{ J / mole K}$$

### 1.3.1 Boyle's Law

For a fixed amount of gas, held at constant temperature, the volume is inversely proportional to the applied pressure.

$$PV = \text{constant}$$

### 1.3.2 Charles' Law

For a fixed amount of gas, held at constant pressure, the volume is directly proportional to the temperature.

$$V/T = \text{constant}$$

### 1.3.3 Gay-Lussac's Law

For a fixed amount of gas, held at constant volume, the pressure is directly proportional to the absolute temperature.

$$P/T = \text{constant}$$

### 1.3.4 Avogadro's Law

At constant temperature and pressure, equal volumes of gas contain equal amounts of molecules.

$$V/n = \text{constant}$$

### 1.3.5 Dalton's Law of Partial Pressures

The total pressure exerted by a mixture of several gases is equal to the sum of the gases individual pressure (partial pressure).

$$P_T = p_a + p_b + p_c + \dots$$

$P_T$  pressure of gas a + b + c

$p_a$  pressure of gas a

$p_b$  pressure of gas b

$p_c$  pressure of gas c

### 1.3.6 Ideal Gas Law

Combining the above relationships and Avogadro's principle (under constant pressure and temperature, equal volumes of gas contain the equal numbers of molecules) into one equation we obtain the Ideal Gas Law:

$$PV = nRT$$

### 1.3.7 Equation of State of Real Gases

An ideal gas has negligible volume and exerts no force. However, real gases do have volumes and do exert forces upon one another. When these factors are taken into consideration, the following equation can be obtained:

$$(P + n^2a/V^2) - (V - nb) = nRT$$

**a** - proportionality constant

**b** - covolume

Note that a and b are dependent on the individual gas, since molecular volumes and molecular attractions vary from gas to gas.

Table 1.10. van der Waals constants for real gases.

Gas	a (atm liter <sup>2</sup> / mole <sup>2</sup> )	b (liter / mole)
He	0.034	0.0237
O <sub>2</sub>	1.36	0.0318
NH <sub>3</sub>	4.17	0.0371
H <sub>2</sub> O	5.46	0.0305
CH <sub>4</sub>	2.25	0.0428

### 1.3.8 Changes of Pressure, Volume, or Temperature

By combining the equations for Boyle's law, Charles' law, and Gay-Lussac's law, a single equation can be obtained that is useful for many computations:

$$\frac{P_1V_1}{T_1} = \frac{P_2V_2}{T_2}$$

## 1.4 Solutions

### 1.4.1 Mass Percent

The **mass percent** of a solution is the mass of the solute divided by the total mass (solute + solvent) multiplied by 100.

$$\text{percent by mass of solute} = \frac{\text{mass of solute}}{\text{mass of solute} + \text{mass of solvent}} \times 100$$

### 1.4.2 Mole Fraction (*X*)

The **mole fraction** (*X*) is the number of moles of component A divided by the total number of moles of all components.

$$\text{mole fraction of A} = \frac{\text{moles of A}}{\text{moles of all components}}$$

### 1.4.3 Molarity (*M*)

The **molarity** (*M*) of a solution is the number of moles of solute dissolved in 1 liter of solvent.

$$\text{molarity} = \frac{\text{moles of solute}}{\text{liters of solvent}}$$

#### 1.4.4 Molality (*m*)

**Molality (*m*)** is the number of moles of solute dissolved in 1000 g (1 kg) of solvent.

$$\text{molality} = \frac{\text{moles of solute}}{\text{mass of solvent}}$$

#### 1.4.5 Dilutions

A handy and useful formula when calculating dilutions is:

$$M_{\text{initial}} V_{\text{initial}} = M_{\text{final}} V_{\text{final}}$$


### 1.5 Acids and Bases

#### 1.5.1 Arrhenius Concept

An acid is any species that increases the concentration of **hydronium ions ( $H_3O^+$ )**, in aqueous solution.

A base is any species that increases the concentration of **hydroxide ion, ( $OH^-$ )**, in aqueous solution.

For an acid:



For a base:



However, the drawback with the Arrhenius concept is that it only applies to aqueous solutions.


#### 1.5.2 Bronsted-Lowery Concept

An acid is a species which can donate a proton (i.e., a hydrogen ion,  $H^+$ ) to a proton acceptor.

A base is a species which can accept a proton from a proton donor.

Along with the Bronsted-Lowery concept of a proton donor (acid) and a proton acceptor (base), arises the concept of conjugate acid-base pairs. For

example, when the acid HCl reacts, it donates a proton thereby leaving  $\text{Cl}^-$  (which is now a proton acceptor, or the conjugate base of HCl). Using  $\text{NH}_3$  as the base and  $\text{H}_2\text{O}$  as the acid:



### 1.5.3 Lewis Concept

An acid is a species that can accept a pair of electrons.

A base is a species that can donate a pair of electrons.



## 1.6 Thermodynamics

### 1.6.1 First Law of Thermodynamics

The energy change of a system is equal to the heat absorbed by the system plus the work done by the system. The reason for the minus sign for work,  $w$ , is that any work done by the system results in a loss of energy for the system as a whole.

$$\Delta E = q - w$$

$E$  = internal energy of the system

$q$  = heat absorbed by the system

$w$  = work done by the system

Table 1.11. Thermodynamic Processes

Process	Sign
work done by system	-
work done on system	+
heat absorbed by system (endothermic)	+
heat absorbed by surroundings (exothermic)	-

Making a substitution for work, the equation can be expressed as:

$$\Delta E = q - P\Delta V$$

For constant volume, the equation becomes:

$$\Delta E = q_v$$

### 1.6.2 Enthalpy

**Enthalpy, H,** is the heat content of the system at constant pressure.

$$\Delta E = q_p - P\Delta V$$

$$q_p = \Delta E + P\Delta V$$

$$q_p = (E_2 - E_1) + P(V_2 - V_1)$$

$$= (E_2 + PV_2) - (E_1 + PV_1)$$

$$q_p = H_2 - H_1 = \Delta H$$

### 1.6.3 Entropy

**Entropy, S,** is the measure of the degree of randomness of a system.

$$\Delta S = \frac{q_{rev}}{T}$$

T = temperature in °K

### 1.6.4 Gibbs Free Energy

**Gibbs free energy, G,** is the amount of energy available to the system to do useful work.

$$\Delta G = \Delta H - T\Delta S$$

$\Delta G < 0$	spontaneous process from 1 $\rightarrow$ 2
$\Delta G > 0$	spontaneous process from 2 $\rightarrow$ 1
$\Delta G = 0$	equilibrium

### 1.6.5 Standard States

The **standard state** is the standard or normal condition of a species.

*Table 1.12. Standard States*

State of Matter	Standard State
Gas	1 atm pressure
Liquid	Pure liquid
Solid	Pure solid
Element	Free energy of formation = 0
Solution	1 molar concentration


Note also that  $\Delta H^\circ_f$  for an element in its natural state at 25°C and 1 atm is taken to be equal to zero.

### 1.6.6 Hess' Law of Heat Summation



The final value of  $\Delta H$  for the overall process is the sum of all the enthalpy changes.

$$\Delta H^\circ = \Sigma \Delta H^\circ_f (\text{products}) - \Sigma \Delta H^\circ_f (\text{reactants})$$




For example, to vaporize 1 mole of  $\text{H}_2\text{O}$  at 100°C and 1 atm, the process absorbs 41 kJ of heat,  $\Delta H = +41$  kJ.



If a different path to the formation of 1 mole of gasous  $\text{H}_2\text{O}$  is taken, the same amount of net heat will still be absorbed.



Reversing the first reaction, then adding the two reactions together and cancelling common terms, results in the original reaction, and the amount of heat absorbed by the system



Using the Hess' Law of Summation:

$$\Delta H^\circ = \sum \Delta H_f^\circ(\text{products}) - \sum \Delta H_f^\circ(\text{reactants})$$

$$\Delta H^\circ = (-242 \text{ kJ}) - (283 \text{ kJ}) = +41 \text{ kJ}$$




Figure 1.10. Enthalpy Diagram for  $\text{H}_2\text{O}(l) \rightarrow \text{H}_2\text{O}(g)$

## 1.7 Equilibria





Figure 1.11. Concentrations of reactants and products approaching equilibrium.

When equilibrium is reached in a chemical reaction the rate of the forward reaction is equal to the rate of the reverse reaction, and the concentrations of the reactants and products do not change over time.

### 1.7.1 Homogeneous Equilibrium

**Homogeneous equilibrium** occurs when all reacting species are in the same phase. For the general reaction,



the equation expressing the **law of mass action**, at equilibrium, is:

$$\frac{[C]^c[D]^d}{[A]^a[B]^b} = K_c$$

The quantity,  $K_c$ , is a constant, called the **equilibrium constant** (in this case it denotes the equilibrium constant for species in solution, expressed as moles per liter). The magnitude of  $K_c$  tells us to what extent the reaction proceeds. A large  $K_c$  indicates that the reaction proceeds to the right of the reaction. A low value indicates that the reaction proceeds to the left of the reaction.

For gas-phase equilibrium the expression becomes

$$K_p = \frac{P_C^c P_D^d}{P_A^a P_B^b}$$

where  $P$  is the partial pressures of the species in the reaction.  $K_p$  can be related to  $K_c$  by the following equation,

$$K_p = K_c (0.08206T)^{\Delta n}$$

**T = the absolute temperature**

**$\Delta n$  = moles of product - moles of reactants.**

### 1.7.2 Heterogeneous Equilibrium

**Heterogeneous equilibrium** involves reactants and products in different phases. For example, when calcium carbonate is heated in a closed vessel, the following equilibrium reaction occurs:



The reactant is a solid, while the products are in the solid and gas phase. The equilibrium expression is written as the following:

$$K_c' = \frac{[\text{CaO}][\text{CO}_2]}{[\text{CaCO}_3]}$$

In any reaction that includes a solid, the solid concentration will remain constant and therefore is not included in the equilibrium expression. The equilibrium expression now becomes:

$$K_c'' \frac{[\text{CaCO}_3]}{\text{CaO}} = [\text{CO}_2] = K_c$$

### 1.7.3 Le Chatelier's Principle

**Le Chatelier's Principle** states that *when a system is in equilibrium and there is a change in one of the factors which affect the equilibrium, the system reacts in such a way as to cancel out the change and restore equilibrium.*

An increase in temperature will shift the reaction in the direction of heat absorption.


An increase in the pressure will shift the reaction in the direction in which the number of moles is decreased.

An increase or decrease in pressure does not affect a reaction in which there is no variation in the number of moles.

An increase in the concentration of one of the components will cause the reaction to shift so as to decrease the added component.

### 1.7.4 Equilibrium of Water

The reaction for autoionization of water is:



The equilibrium expression is:

$$K = \frac{[\text{H}_3\text{O}^+][\text{OH}^-]}{[\text{H}_2\text{O}][\text{H}_2\text{O}]}$$

Since the concentration of water is a constant ( $\approx 55.55 \text{ M}$ ),  $[\text{H}_2\text{O}]^2$  can be included in the equilibrium constant, K. this new constant is now called K<sub>w</sub>.

$$K_w = K[\text{H}_2\text{O}]^2 = [\text{H}_3\text{O}^+] [\text{OH}^-]$$

K<sub>w</sub> is the **ion product constant** for water, also called the **ionization constant** or **dissociation constant** for water. The ionization constant for water at 25 °C has a value of  $1.0 \times 10^{-14}$ . The equilibrium expression now becomes:

$$K_w = 1.0 \times 10^{-14} = [\text{H}_3\text{O}^+] [\text{OH}^-]$$

$$\text{Since } [\text{H}_3\text{O}^+] = [\text{OH}^-]$$

$$[\text{H}_3\text{O}^+] = [\text{OH}^-] = 1.0 \times 10^{-7}$$

When the concentration of hydrogen ions equals the concentration of hydroxide ions the solution is said to be neutral.

### 1.7.5 pH

pH is the measure of how strong or weak an acid is, and is defined as the negative of the log of the hydrogen ion concentration, or

$$\text{pH} = -\log[\text{H}_3\text{O}^+]$$

Water has a pH of 7, this is calculated from the dissociation constant for water:

$$[\text{H}_3\text{O}^+] = 1.0 \times 10^{-7}$$

$$\text{pH} = -\log[\text{H}_3\text{O}^+] = -\log(1.0 \times 10^{-7})$$

$$\text{pH} = 7$$

The concept of pH can be applied to any system in which hydrogen ions are produced. An acidic solution would have an excess of hydrogen ions, a basic solution would have an excess of hydroxide ions, and a neutral solution the hydrogen ions would equal the hydroxide ions. Since pH is a measure of the hydrogen ion concentration, acidic and basic solutions can be distinguished on the basis of their pH.

acidic solutions:	$[\text{H}_3\text{O}^+] > 10^{-7} \text{ M}$ , $\text{pH} < 7$
basic solutions:	$[\text{H}_3\text{O}^+] < 10^{-7} \text{ M}$ , $\text{pH} > 7$
neutral solutions:	$[\text{H}_3\text{O}^+] = 10^{-7} \text{ M}$ , $\text{pH} = 7$

### 1.7.6 Ionic Equilibrium

For a monoprotic acid HA, the equilibrium reaction is:




and the equilibrium expression is:


$$K_a = \frac{[\text{H}_3\text{O}^+][\text{A}^-]}{[\text{HA}]}$$

The equilibrium constant,  $K_a$ , is called the **acid dissociation constant**.


Similarly for a polyprotic acid (i.e. phosphoric acid), the equilibrium reactions are:



$$K_a' = \frac{[\text{H}^+][\text{H}_3\text{PO}_4^-]}{\text{H}_3\text{PO}_4} = 7.5 \times 10^{-3}$$




$$K_a'' = \frac{[\text{H}^+][\text{HPO}_4^{2-}]}{[\text{H}_2\text{PO}_4^-]} = 6.2 \times 10^{-8}$$



$$K_a''' = \frac{[\text{H}^+][\text{PO}_4^{3-}]}{[\text{HPO}_4^{2-}]} = 4.8 \times 10^{-13}$$

For a base the equilibrium reaction is:




and the equilibrium expression is:


$$K_b = \frac{[\text{BH}^+][\text{OH}^-]}{[\text{B}]}$$

The equilibrium constant,  $K_b$ , is called the **base dissociation constant**.

### 1.7.7 Relationship between $K_a$ and $K_b$ Conjugate Pair



$$K_a = \frac{[\text{H}_3\text{O}^+][\text{A}^-]}{[\text{HA}]}$$



$$K_b = \frac{[\text{HA}][\text{OH}^-]}{[\text{A}^-]}$$

$$K_a K_b = [\text{H}_3\text{O}^+] [\text{OH}^-] = 10^{-14}$$

$$\text{p}K_a + \text{p}K_b = 14$$

### 1.7.8 Hydrolysis


Hydrolysis is the reaction between water and the ions of a salt.

#### 1.7.8.1 Salt of a strong acid - strong base.

Consider NaCl, the salt of a strong acid and a strong base. The hydrolysis of this salt would yield NaOH and HCl. Since both species would completely dissociate into their respective ions yielding equivalent amounts of  $\text{H}_3\text{O}^+$  and  $\text{OH}^-$ , the overall net effect would be that no hydrolysis takes place. Since  $[\text{H}_3\text{O}^+] = [\text{OH}^-]$ , the pH would be 7, a neutral solution.

#### 1.7.8.2 Salt of a strong acid - weak base.

Consider the hydrolysis of  $\text{NH}_4\text{Cl}$ :




$$K_h = \frac{[H_3O^+][NH_3]}{[NH_4^+]}$$

$$K_h = \frac{K_w}{K_b} = \frac{[H_3O^+][OH^-]}{[NH_4^+][OH^-]/[NH_3]} = \frac{[H_3O^+][NH_3]}{[NH_4^+]}$$

### 1.7.8.3 Salt of a weak acid - strong base.

Consider the hydrolysis of  $NaC_2H_3O_2$ :




$$K_h = \frac{[HC_2H_3O_2][OH^-]}{[C_2H_3O_2^-]}$$

$$K_h = \frac{K_w}{K_a} = \frac{[H_3O^+][OH^-]}{[H_3O^+][C_2H_3O_2^-]/[HC_2H_3O_2]} = \frac{[HC_2H_3O_2][OH^-]}{[C_2H_3O_2^-]}$$

### 1.7.9 Solubility Product

In the case for which a solid is being dissolved, the general chemical reaction becomes:



and the equilibrium expression is:

$$K = \frac{[A]^a[B]^b}{[A_aB_b]}$$

the denominator in the expression  $[A_aB_b]$  represents the concentration of the pure solid and is constant, therefore it can be incorporated into the equilibrium constant, K. The expression now becomes:

$$K_{sp} = [A]^a[B]^b$$

For example, a saturated solution of AgCl, would have the following equilibrium:



$$K_{\text{sp}} = [\text{Ag}^+] [\text{Cl}^-]$$

The value of the  $K_{\text{sp}}$  for AgCl is  $1.7 \times 10^{-10}$

$$1.7 \times 10^{-10} = [\text{Ag}^+] [\text{Cl}^-]$$

If the ion product is equal to or less than the  $K_{\text{sp}}$ , no precipitate will form. If the ion product is greater than the  $K_{\text{sp}}$  value, the material will precipitate out of solution so that the ion product will be equal to the  $K_{\text{sp}}$ .

*Table 1.13. Solubility Products*

Compound	$K_{\text{sp}}$
AgCl	$1.7 \times 10^{-10}$
AgBr	$5.0 \times 10^{-13}$
BaSO <sub>4</sub>	$1.5 \times 10^{-9}$
CuS	$8.5 \times 10^{-36}$
PbCl <sub>2</sub>	$1.6 \times 10^{-5}$
HgS	$1.6 \times 10^{-54}$

### 1.7.10 Common Ion Effect

The common ion is when an ion common to one of the salt ions is introduced to the solution. The introduction of a common ion produces an effect on the equilibrium of the solution and according to Le Chatelier's principle, i.e. the equilibrium is shifted so as to reduce the effect of the added ion. This is referred to as the **common ion effect**.

In the case of a solution of AgCl, if NaCl is added, the common ion being Cl<sup>-</sup>, the equilibrium would be shifted to the left so that the ion product will preserve the value of the  $K_{\text{sp}}$ .

## 1.8 Kinetics

**Kinetics** deals with the rate (how fast) that a chemical reaction proceeds with. The reaction rate can be determined by following the concentration of either the reactants or products. The rate is also dependent on the concentrations, temperature, catalysts, and nature of reactants and products.

### 1.8.1 Zero-Order Reactions


Zero-order reactions are independent of the concentrations of reactants.



$$\text{rate} = -\frac{\Delta[A]}{\Delta t} = k[A]^0 = k$$

### 1.8.2 First-Order Reactions


First-order reactions are dependent on the concentration of the reactant.



$$\text{rate} = k[A]^1 = k[A]$$

### 1.8.3 Second-Order Reactions

There are two type of second-order reactions. The first kind involves a single kind of reactant.



$$\text{rate} = k[A]^2$$


The second kind of reaction involves two different kind of reactants.



$$\text{rate} = k[A][B]$$

### 1.8.4 Collision Theory

Consider the decomposition of HI.



In order for the decomposition of HI to take place, two molecules of HI must collide with each other with the proper orientation as shown in Figure 1.12. If the molecules collide without the proper orientation then no decomposition takes place.




Figure 1.12. Effective and Ineffective collisions

Not all collisions with the proper orientation will react. Only those collisions with the proper orientation and sufficient energy to allow for the breaking and forming of bonds will react. The minimum energy available in a collision which will allow a reaction to occur is called the **activation energy**.

### 1.8.5 Transition State Theory

When a collision with the proper orientation and sufficient activation energy occurs, an intermediate state exists before the products are formed. This intermediate state, also called an **activated complex** or **transition state**, is neither the reactant or product, but rather a highly unstable combination of both, as represented in Figure 1.13 for the decomposition of HI.




Figure 1.13. Transition state or activated complex




Figure 1.14. Potential energy diagram for the decomposition of HI.

Figure 1.14 shows the potential energy diagram for the decomposition of HI. As can be seen, in order to reach the activated complex or transition state the proper orientation and sufficient collision energy must be achieved. Once these requirements are achieved the reaction continues on to completion and the products are formed.

### 1.8.6 Catalysts

A **catalyst** is a substance that affects the rate of a chemical reaction without itself being consumed or chemically altered. The catalyst takes part in the reaction by providing an alternative route to the production of products. The catalyzed reaction has a lower activation energy than that of the uncatalyzed reaction, as shown in Figure 1.15. By lowering the activation energy there are more molecules with sufficient energy that can react and thus the rate of the reaction is affected.

A **homogeneous catalyst** is in the same phase as the reactants. The catalyst and the reactants form a reactive intermediate.

A **heterogeneous catalyst** is not in the same phase as the reactants. The reactants are absorbed on the surface of the heterogeneous catalyst and the reaction then takes place.




Figure 1.15. Energy diagram for a reaction with and without a catalysts

*This page intentionally left blank.*

## Chapter 2

### **Inorganic Chemistry**

- 2.1 Group IA Elements*
- 2.2 Group IIA Elements*
- 2.3 Group IIIA Elements*
- 2.4 Group IVA Elements*
- 2.5 Group VA Elements*
- 2.6 Group VIA Elements*
- 2.7 Group VIIA Elements*
- 2.8 Group VIIIA Elements*
- 2.9 Transition Elements*
- 2.10 Ionic Solids*
- 2.11 Coordination Numbers*
- 2.12 Ionic Crystal Systems*
- 2.13 Crystal Lattice Packing*
- 2.14 Crystal Lattice Types*
- 2.15 Crystal Lattice Energy*
- 2.16 Complexes*

## 2.1 Group IA Elements

Alkali Metals -Li, Na, K, Rb, Cs, Fr

Table 2.1. Group IA Properties

Element	Li	Na	K	Rb	Cs	Fr
Electronic configuration	[He]2s	[Ne]3s	[Ar]4s	[Kr]5s	[Xe]6s	[Rn]7s
M.P. (°K)	453.7	371.0	336.35	312.64	301.55	300
B.P. (°K)	1615	1156	1032	961	944	950
Pauling's Electronegativity	0.98	0.93	0.82	0.82	0.79	0.7
Atomic radius (Å)	2.05	2.23	2.77	2.98	3.34	-
Covalent radius (Å)	1.23	1.54	2.03	2.16	2.35	-
Ionic radius (Å)(+1)	0.68	0.98	1.33	1.48	1.67	1.8
Ionization enthalpy (eV)	5.392	5.139	4.341	4.177	3.894	-
Crystal Structure	bcc	bcc	bcc	bcc	bcc	bcc

Table 2.2. Group IA Compounds

	Li	Na	K	Rb	Cs	Fr
H <sup>-</sup>	x	x	x	x	x	-
X <sup>-</sup>	x	x	x	x	x	-
CH <sub>3</sub> COO <sup>-</sup>	x	x	x	x	x	-
HCO <sub>3</sub> <sup>-</sup>	x	x	x	x	x	-
ClO <sup>-</sup>	-	x	x	-	-	-
ClO <sub>3</sub> <sup>-</sup>	x	x	x	x	x	-
ClO <sub>4</sub> <sup>-</sup>	x	x	x	x	x	-
OH <sup>-</sup>	x	x	x	x	x	-
NO <sub>3</sub> <sup>-</sup>	x	x	x	x	x	-
NO <sub>2</sub> <sup>-</sup>	x	x	x	-	x	-
H <sub>2</sub> PO <sub>4</sub> <sup>-</sup>	x	x	x	-	-	-
HSO <sub>4</sub> <sup>-</sup>	x	x	x	x	x	-
HSO <sub>3</sub> <sup>-</sup>	-	x	x	-	-	-
CO <sub>3</sub> <sup>-2</sup>	x	x	x	x	x	-
C <sub>2</sub> O <sub>4</sub> <sup>-2</sup>	x	x	x	-	x	-
HPO <sub>4</sub> <sup>-2</sup>	-	x	x	-	-	-
SO <sub>4</sub> <sup>-2</sup>	x	x	x	x	x	-
SO <sub>3</sub> <sup>-2</sup>	x	x	x	-	-	-
PO <sub>3</sub> <sup>-3</sup>	x	x	x	-	-	-
PO <sub>4</sub> <sup>-3</sup>	x	x	x	-	-	-
N <sup>-3</sup>	x	x	x	-	-	-

## 2.2 Group IIA Elements

Alkaline Earth Metals -Be, Mg, Ca, Sr, Ba, Ra

Table 2.3. Group IIA Properties

Element	Be	Mg	Ca	Sr	Ba	Ra
Electronic configuration	[He]2s <sup>2</sup>	[Ne]3s <sup>2</sup>	[Ar]4s <sup>2</sup>	[Kr]5s <sup>2</sup>	[Xe]6s <sup>2</sup>	[Rn]7s <sup>2</sup>
M.P. (°K)	1560	922	1112	1041	1002	973
B.P. (°K)	2745	1363	1757	1650	2171	1809
Pauling's Electronegativity	1.57	1.31	1.00	0.95	0.89	0.9
Atomic radius (Å)	1.40	1.72	2.223	2.45	2.78	-
Covalent radius (Å)	0.90	1.36	1.74	1.91	1.98	-
Ionic radius (Å)(+2)	0.35	0.66	1.18	1.112	1.34	1.43
Ionization enthalpy (eV)	9.322	7.646	6.113	5.695	5.212	5.279
Crystal Structure	hex	hex	fcc	fcc	bcc	bcc

Table 2.4. Group IIA Compounds

	Be	Mg	Ca	Sr	Ba	Ra
H <sup>+</sup>	x	x	x	x	x	-
X <sup>-</sup>	x	x	x	x	x	-
CH <sub>3</sub> COO <sup>-</sup>	x	x	x	x	x	-
HCO <sub>3</sub> <sup>-</sup>	-	-	-	-	-	-
ClO <sup>-</sup>	-	-	x	-	x	-
ClO <sub>3</sub> <sup>-</sup>	-	x	x	x	x	-
ClO <sub>4</sub> <sup>-</sup>	-	x	x	x	x	-
OH <sup>-</sup>	x	x	x	x	x	-
NO <sub>3</sub> <sup>-</sup>	x	x	x	x	x	-
NO <sub>2</sub> <sup>-</sup>	-	x	x	x	x	-
H <sub>2</sub> PO <sub>4</sub> <sup>-</sup>	-	-	x	-	x	-
HSO <sub>4</sub> <sup>-</sup>	-	-	-	x	-	-
HSO <sub>3</sub> <sup>-</sup>	-	-	x	-	-	-
CO <sub>3</sub> <sup>-2</sup>	x	x	x	x	x	-
C <sub>2</sub> O <sub>4</sub> <sup>-2</sup>	x	x	x	x	x	-
HPO <sub>4</sub> <sup>-2</sup>	-	x	x	x	x	-
SO <sub>4</sub> <sup>-2</sup>	x	x	x	x	x	-
SO <sub>3</sub> <sup>-2</sup>	-	x	x	x	x	-
PO <sub>3</sub> <sup>-3</sup>	-	-	x	-	x	-
PO <sub>4</sub> <sup>-3</sup>	x	x	x	-	x	-
N <sup>-3</sup>	x	x	x	x	x	-

## 2.3 Group IIIA Elements

Boron group - B, Al, Ga, In, Tl

Table 2.5. Group IIIA Properties

Element	B	Al	Ga	In	Tl
Electronic configuration	[He]2s <sup>2</sup> 2p	[Ne]3s <sup>2</sup> 3p	[Ar]3d <sup>10</sup> 4s <sup>2</sup> 4p	[Kr]4d <sup>10</sup> 5s <sup>2</sup> 5p	[Xe]4f <sup>14</sup> 5d <sup>10</sup> 6s <sup>2</sup> 6p
M.P. (°K)	2300	933.25	301.90	429.76	577
B.P. (°K)	4275	2793	2478	2346	1746
Pauling's Electronegativity	2.04	1.61	1.81	1.78	2.04
Atomic radius (Å)	1.17	1.82	1.81	2.00	2.08
Covalent radius (Å)	0.82	1.18	1.26	1.44	1.48
Ionic radius (Å)(+3)	0.23	0.51	0.81	0.81	0.95
Ionization enthalpy (eV)	8.298	5.986	5.999	5.786	6.108
Crystal Structure	rhom	fcc	orthorho	tetrag	hex

Table 2.6. Group IIIA Compounds

	B	Al	Ga	In	Tl
H <sup>-</sup>	x	-	x	-	-
X <sup>-</sup>	x	x	x	x	x
CH <sub>3</sub> COO <sup>-</sup>	-	x	x	-	x
HCO <sub>3</sub> <sup>-</sup>	-	-	-	-	-
ClO <sup>-</sup>	-	-	-	-	-
ClO <sub>3</sub> <sup>-</sup>	-	-	-	-	x
ClO <sub>4</sub> <sup>-</sup>	-	-	x	x	x
OH <sup>-</sup>	-	x	x	x	x
NO <sub>3</sub> <sup>-</sup>	-	x	x	x	x
NO <sub>2</sub> <sup>-</sup>	-	-	-	-	x
H <sub>2</sub> PO <sub>4</sub> <sup>-</sup>	-	-	-	-	x
HSO <sub>4</sub> <sup>-</sup>	-	-	-	-	x
HSO <sub>3</sub> <sup>-</sup>	-	-	-	-	x
CO <sub>3</sub> <sup>-2</sup>	-	-	-	-	x
C <sub>2</sub> O <sub>4</sub> <sup>-2</sup>	-	x	x	-	x
HPO <sub>4</sub> <sup>-2</sup>	-	-	-	-	-
SO <sub>4</sub> <sup>-2</sup>	-	x	x	x	x
SO <sub>3</sub> <sup>-2</sup>	-	-	-	-	x
PO <sub>3</sub> <sup>-3</sup>	-	x	-	-	-
PO <sub>4</sub> <sup>-3</sup>	-	x	-	-	x
N <sup>-3</sup>	x	x	x	-	-

## 2.4 Group IVA Elements

Carbon group - C, Si, Ge, Sn, Pb

Table 2.7. Group IVA Properties

Element	C	Si	Ge	Sn	Pb
Electronic configuration	[He]2s <sup>2</sup> 2p <sup>2</sup>	[Ne]3s <sup>2</sup> 3p <sup>2</sup>	[Ar]3d <sup>10</sup> 4s <sup>2</sup> 4p <sup>2</sup>	[Kr]4d <sup>10</sup> 5s <sup>2</sup> 5p <sup>2</sup>	[Xe]4f <sup>14</sup> 5d <sup>10</sup> 6s <sup>2</sup> 6p <sup>2</sup>
M.P. (°K)	4100	1685	1210.4	505.06	600.6
B.P. (°K)	4470	3540	3107	2876	2023
Pauling's Electronegativity	2.55	1.90	2.01	1.96	2.33
Atomic radius (Å)	0.91	1.46	1.52	1.72	1.81
Covalent radius (Å)	0.77	1.11	1.22	1.41	1.47
Ionic radius (Å)(xx)					
Ionization enthalpy (eV)	11.260	8.151	7.899	7.344	7.416
Crystal Structure	hex	fcc	orthorho	tetrag	fcc

## 2.5 Group VA Elements

Nitrogen group - N, P, As, Sb, Bi

Table 2.8. Group VA Properties

Element	N	P	As	Sb	Bi
Electronic configuration	[He]2s <sup>2</sup> 2p <sup>3</sup>	[Ne]3s <sup>2</sup> 3p <sup>3</sup>	[Ar]3d <sup>10</sup> 4s <sup>2</sup> 4p <sup>3</sup>	[Kr]4d <sup>10</sup> 5s <sup>2</sup> 5p <sup>3</sup>	[Xe]4f <sup>14</sup> 5d <sup>10</sup> 6s <sup>2</sup> 6p <sup>3</sup>
M.P. (°K)	63.14	317.3	1081	904	544.52
B.P. (°K)	77.35	550	876 (sub)	1860	1837
Pauling's Electronegativity	3.04	2.19	2.18	2.05	2.02
Atomic radius (Å)	0.75	1.23	1.33	1.53	1.63
Covalent radius (Å)	0.75	1.06	1.20	1.40	1.46
Ionic radius (Å)(xx)					
Ionization enthalpy (eV)	14.534	10.486	9.81	8.641	7.289
Crystal Structure	hex	monoclini	rhom	rhom	rhom

## 2.6 Group VIA Elements

Oxygen group - O, S, Se, Te, Po

Table 2.9. Group VIA Properties

Element	O	S	Se	Te	Po
Electronic configuration	[He]2s <sup>2</sup> 2p <sup>4</sup>	[Ne]3s <sup>2</sup> 3p <sup>4</sup>	[Ar] 3d <sup>10</sup> 4s <sup>2</sup> 4p <sup>4</sup>	[Kr]4d <sup>10</sup> 5s <sup>2</sup> 5p <sup>4</sup>	[Xe]4f <sup>14</sup> 5d <sup>10</sup> 6s <sup>2</sup> 6p <sup>4</sup>
M.P. (°K)	50.35	388.36	494	722.65	527
B.P. (°K)	90.18	717.75	958	1261	1235
Pauling's Electronegativity	3.44	2.58	2.55	2.1	2.0
Atomic radius (Å)	0.65	1.09	1.22	1.42	1.53
Covalent radius (Å)	0.73	1.02	1.16	1.36	1.46
Ionic radius (Å)(-2)	1.32	1.84	1.91	2.11	-
Ionization enthalpy (eV)	13.618	10.360	9.752	9.009	8.42
Crystal Structure	cubic	orthorho	hex	hex	monoclin

## 2.7 Group VIIA Elements

Halogens - F, Cl, Br, I, At

Table 2.10. Group VIIA Properties

Element	F	Cl	Br	I	At
Electronic configuration	[He]2s <sup>2</sup> 2p <sup>5</sup>	[Ne]3s <sup>2</sup> 3p <sup>5</sup>	[Ar] 3d <sup>10</sup> 4s <sup>2</sup> 4p <sup>5</sup>	[Kr]4d <sup>10</sup> 5s <sup>2</sup> 5p <sup>5</sup>	[Xe]4f <sup>14</sup> 5d <sup>10</sup> 6s <sup>2</sup> 6p <sup>5</sup>
M.P. (°K)	53.48	172.16	265.90	386.7	575
B.P. (°K)	84.95	239.1	332.25	458.4	610
Pauling's Electronegativity	3.98	3.16	2.96	2.66	2.2
Atomic radius (Å)	0.57	0.97	1.12	1.32	1.43
Covalent radius (Å)	0.72	0.99	1.14	1.33	1.45
Ionic radius (Å)(-1)	1.33	1.81	1.96	2.20	-
Ionization enthalpy (eV)	17.411	112.967	11.814	10.451	-
Crystal Structure	cubic	orthorho	orthorho	orthorho	-

## 2.8 Group VIIA Elements

Noble (Inert) Gases - He, Ne, Ar, Kr, Xe, Rn

Table 2.11. Group VIIA Properties

Element	He	Ne	Ar	Kr	Xe	Rn
Electronic configuration	$1s^2$	$[He]$ $2s^22p^6$	$[Ne]$ $3s^23p^6$	$[Ar]$ $3d^{10}4s^2$ $4p^6$	$[Kr]$ $4d^{10}5s^2$ $5p^6$	$[Xe]4f^{14}$ $5d^{10}6s^26p^6$
M.P. (°K)	0.95	24.553	83.81	115.78	165.03	202
B.P. (°K)	4.215	27.096	87.30	119.80	161.36	211
Pauling's Electronegativity	0	0	0	0	0	0
Atomic radius (Å)	0.49	0.51	0.88	1.03	1.24	1.34
Covalent radius (Å)	0.93	0.71	0.98	1.12	1.31	-
Ionic radius (Å)	-	-	-	-	-	-
Ionization enthalpy (eV)	24.58	21.56	15.75	13.99	12.13	10.74
Crystal Structure	hex	fcc	fcc	fcc	fcc	fcc

Table 2.12. Some Group VIIA Compounds

Oxidation State	Compound	Form	Mp (°C)	Structure
II	XeF <sub>2</sub>	Crystal	129	Linear
IV	XeF <sub>4</sub>	Crystal	117	Square
VI	XeF <sub>6</sub>	Crystal	49.6	
	Cs <sub>2</sub> XeF <sub>8</sub>	Solid		
	XeOF <sub>4</sub>	Liquid	-46	Sq. Pyramid
	XeO <sub>3</sub>	Crystal		Pyramidal
VIII	XeO <sub>4</sub>	Gas		Tetrahedral
	XeO <sub>6</sub> <sup>4-</sup>	Salts		Octahedral

## 2.9 Transition Metal Elements

Table 2.13. Transition Elements Properties

Element	Elec. conf.	M.P. (°C)	B.P. (°C)	Density (g/cm³)	Atomic Radius	Ionic Radius (Å)			
						+2	+3	+4	+X
Sc	3d <sup>1</sup> 4s <sup>2</sup>	1540	2730	3.0	1.61		0.81		
Y	4d <sup>1</sup> 5s <sup>2</sup>	1500	2927	4.472	1.8		0.92		
La	5d <sup>1</sup> 6s <sup>2</sup>	920	3470	6.162	1.86		1.14		
Ti	3d <sup>2</sup> 4s <sup>2</sup>	1670	3260	4.5	1.45	0.90	0.87	0.68	
Zr	4d <sup>2</sup> 5s <sup>2</sup>	1850	3580	6.4	1.60			0.79	
Hf	5d <sup>2</sup> 6s <sup>2</sup>	2000	5400	13.2	-			0.78	
									X=5
V	3d <sup>3</sup> 4s <sup>2</sup>	1900	3450	5.8	1.32	0.88	0.74		0.59
Nb	4d <sup>4</sup> 5s <sup>1</sup>	2420	4930	8.57	1.45				0.69
Ta	5d <sup>3</sup> 6s <sup>2</sup>	3000	-	-	1.45				0.68
									X=6
Cr	3d <sup>5</sup> 4s <sup>1</sup>	1900	2640	7.2	1.37	0.88	0.63		0.52
Mo	4d <sup>5</sup> 5s <sup>1</sup>	2610	5560	10.2	1.36				0.62
W	5d <sup>4</sup> 6s <sup>2</sup>	3410	5930	19.3	1.37				0.62
									X=7
Mn	3d <sup>5</sup> 4s <sup>2</sup>	1250	2100	7.4	1.37	0.80	0.66		0.46
Tc	4d <sup>5</sup> 5s <sup>2</sup>	2140	-	-	1.36				
Re	5d <sup>5</sup> 6s <sup>2</sup>	3180	-	21	1.37				0.56
Fe	3d <sup>6</sup> 4s <sup>2</sup>	1540	3000	7.9	1.24	0.76	0.64		
Ru	4d <sup>7</sup> 5s <sup>1</sup>	2300	3900	12.2	1.33			0.67	
Os	5d <sup>6</sup> 6s <sup>2</sup>	3000	5500	22.4	1.34			0.69	
Co	3d <sup>7</sup> 4s <sup>2</sup>	1490	2900	8.9	1.25	0.74	0.63		
Rh	4d <sup>8</sup> 5s <sup>1</sup>	1970	3730	12.4	1.34	0.86	0.68		
Ir	5d <sup>7</sup> 6s <sup>2</sup>	2450	4500	22.5	1.36			0.68	
Ni	3d <sup>8</sup> 4s <sup>2</sup>	1450	2730	8.9	1.25	0.72	0.62		
Pd	4d <sup>10</sup>	1550	3125	12.0	1.38	0.80		0.65	
Pt	5d <sup>9</sup> 6s <sup>1</sup>	1770	3825	21.4	1.39	0.80		0.65	
									X=1
Cu	3d <sup>10</sup> 4s <sup>1</sup>	1083	2600	8.9	1.28	0.69			0.96
Ag	4d <sup>10</sup> 5s <sup>1</sup>	961	2210	10.5	1.44	0.89			1.26
Au	5d <sup>10</sup> 6s <sup>1</sup>	1063	2970	19.3	1.44		0.85		1.37
Zn	3d <sup>10</sup> 4s <sup>2</sup>	419	906	7.3	1.33	0.74			
Cd	4d <sup>10</sup> 5s <sup>2</sup>	321	765	8.64	1.49	0.97			
Hg	5d <sup>10</sup> 6s <sup>2</sup>	-39	357	13.54	1.52	1.10			

Table 2.14. Oxidation States of Transition Elements

Sc	Ti	V	Cr	Mn	Fe	Co	Ni	Cu	Zn
+III	+III	+II	+II	+II	+II	+II	+II	+I	+II
+IV	+III	+III	+III	+III	+III	+III	+III	+II	
	+IV	+VI	+IV	+IV					
	+V		+VI	+VI	+VI				
				+VII					
Ce		Mo		Ru	Rh	Pd	Ag	Cd	
+III		+III		+II	+III	+II	+I	+II	
+IV		+IV		+III	+IV	+IV	+II		
		+V		+IV					
		+VI		+VI					
	W		Os	Ir	Pt	Au	Hg		
	+IV		+IV	+III	+II	+I	+I		
	+V		+VI	+IV	+IV	+III	+II		
	+VI		+VIII	+VI					

## 2.10 Ionic Solids

**Ionic solids** are mostly inorganic compounds, which are held together by ionic bonds (Chapter 1.2.3). The ions are treated as thought they were hard spheres with either positive (cations) or negative (anions) charges. Using this model the ionic solids or **crystals**, are found to be arranged in a very specific order called a **crystal lattice**. The crystal lattice consists of repeating units called unit cells (see below).

## 2.11 Coordination Number

For ionic compounds the **coordination number** is the number of anions that are arranged about the cation in an organized structure. For example, NaCl has a coordination number of 6. In otherwords, 6 Cl<sup>-</sup> atoms surround 1 Na<sup>+</sup> atom. The number of anions that can surround a cation is dependent (but not entirely) on the relative sizes of the ions involved. Table 2.15 illustrates the ratios of the radii of the ions and their coordination number.

Table 2.15. Radius Ratios and Coordination Number

Coordination Number	Geometry	Ratio (+/-)
2	Linear	0.000 - 0.155
3	Trigonal	0.155 - 0.225
4	Tetrahedral	0.225 - 0.414
4/6	Sq. Planar/Octahedral	0.414 - 0.732
8	Cubic	0.732 - 1.000
12	Dodecahedral	1.000 -

Taking NaCl, the radii of  $\text{Na}^+$  ion is  $0.95\text{\AA}$  and  $\text{Cl}^-$  is  $1.81\text{\AA}$ . Their ratio would be as follows:

$$\frac{r_{\text{cation}}}{r_{\text{anion}}} = \frac{r_{\text{Na}}}{r_{\text{Cl}}} = \frac{0.95}{1.81} = 0.52$$

forming the sodium chloride lattice with coordination number 6.

## 2.12 Ionic Crystal Systems

A crystal lattice can be broken down to a small repeating unit called a unit cell. There are only seven distinct unit cells possible, and are shown in Figure 2.1.

	All Edges Equal	Two Edges Equal	No Edges Equal
All 90° Angles	<b>cubic</b>  $a = b = c$ $\alpha = \beta = \gamma = 90^\circ$	<b>tetragonal</b>  $a = b \neq c$ $\alpha = \beta = \gamma = 90^\circ$	<b>orthorhombic</b>  $a \neq b \neq c$ $\alpha = \beta = \gamma = 90^\circ$
Two 90° angles		<b>hexagonal</b>  $a = b \neq c$ $\alpha = \beta = 90^\circ; \gamma = 120^\circ$	<b>monoclinic</b>  $a \neq b \neq c$ $\alpha = \gamma = 90^\circ; \beta \neq 90^\circ$

Figure 2.1. Unit Cells




Figure 2.1. Unit Cells

## 2.13 Crystal Lattice Packing

When a crystal lattice forms the ions are arranged in the most efficient way of packing spheres into the smallest possible space. Starting with a single layer as in figure 2.2a, a second layer can be placed on top of it in the hollows of the first Figure 2.2b. At this point a third layer can be placed. If the third layer is placed directly over the first, Figure 2.2c, the structure is **hexagonal close-packed (hcp)**. The layering in hexagonal close-packed is ABABAB. If the third layer is placed so it is not directly over the first a different arrangement is obtained. This structure is **cubic close-packed (ccp)**. The layering in cubic close-packed is ABCABC.




Figure 2.2. Close Packing of Atoms




Figure 2.2. Close Packing of Atoms

## 2.14 Crystal Lattice Types

Shown below are commonly encountered crystal lattice structures. The lattice type depends on the radius ratio favoring a particular coordination number for the structure type.




Figure 2.3. Common Lattice Types




Figure 2.3. Common Lattice Types

## 2.15 Crystal Lattice Energy

### 2.15.1 Born-Haber Cycle

An important property of an ionic crystal is the energy required to break the crystal apart into individual ions, this is the **crystal lattice energy**. It can be measured by a thermodynamic cycle, called the **Born-Haber** cycle.




Figure 2.4. Born-Haber Cycle

The Born-Haber cycle follows the Law of Conservation of Energy, that is when a system goes through a series of changes and is returned to its initial state the sum of the energy changes is equal to zero. Thus the equation:

$$0 = \Delta h_f + \Delta h_{vap} + \frac{1}{2} \Delta h_{diss} + \Delta H_{ion} + \Delta H_{EA} + U$$

From this the crystal lattice energy, U, can be calculated from the following enthalpies:

enthalpy of formation	$(\Delta h_f)$	-411
vaporization of sodium	$(\Delta h_{vap})$	-108
dissociation of Cl <sub>2</sub> (g) into gaseous atoms	$(\frac{1}{2} \Delta h_{diss})$	-121
ionization of Na(g) to Na <sup>+</sup> (g)	$(\Delta h_{ion})$	-502
<u>electron attachment to Cl(g) to give Cl<sup>-</sup>(g)</u>	$(\Delta H_{EA})$	354
crystal lattice energy	U	-788 kJ/mol

### 2.15.2 Madelung Constant

The crystal lattice energy can be estimated from a simple electrostatic model. When this model is applied to an ionic crystal only the electrostatic charges and the shortest anion-cation internuclear distance need be considered. The summation of all the geometrical interactions between the ions is called the **Madelung constant**. From this model an equation for the crystal lattice energy is derived:

$$U = -1389 \frac{M}{r} \left( 1 - \frac{1}{n} \right)$$

U = crystal lattice energy

M = Madelung constant

r = shortest internuclear distance

n = Born exponent

The Madelung constant is unique for each crystal structure and is defined only for those whose interatomic vectors are fixed by symmetry. The **Born exponent**, n, can be estimated for alkali halides by the noble-gas-like electron configuration of the ion. It can also be estimated from the compressibility of the crystal system. For NaCl, n equals 9.1.

Table 2.16. Madelung Constants

Structure Type	M
NaCl	1.74756
CsCl	1.76267
CaF <sub>2</sub>	5.03878
Zinc Blende	1.63805
Wurtzite	1.64132

Table 2.17. Born Exponents

Configuration	n
He	5
Ne	7
Ar	9
Kr	10
Xe	12

For NaCl, substituting in the appropriate values into the equation we obtain:

$$U = -1389 \frac{1.747}{2.82} \left( 1 - \frac{1}{9.1} \right)$$

$$U = -860 + 95 = -765 \text{ kJ mol}^{-1}$$

As can be seen the result is close (within 3%) of the value of U obtained from using the Born-Haber cycle. More accurate calculations can be obtained


if other factors are taken into account, such as, van der Waals repulsion, zero-point energy, etc....

## 2.16 Complexes


Transition and non-transition metal ions form a great many complex ions and molecules. Bonding is achieved by an ion or molecule donating a pair of electrons to the metal ion. This type of bond is an coordinate covalent bond (section 1.2.2), the resulting complexes are called **coordination complexes**. The species donating the electron pair is called a **ligand**. More than one type of ligand can bond to the same metal ion, i.e.  $\text{K}_2\text{PtCl}_6$ . In addition a ligand can bond to more than one site on the metal ion, a phenomenon called **chelation**.

The bonding involved in the formation of coordination complexes involve the d orbitals (section 1.1.3) of the metal ion. The electron pair being donated occupies the empty d orbitals and accounts for the geometry of the complex.


### 2.16.1 Unidentate Ligands




### 2.16.2 Bidentate Ligands




Oxalate Ion



Ethylenediamine




Diacetyl dioxime




Acetylacetone

## 2.16.3 Tridentate Ligands




Diethylenetriamine

## 2.16.4 Quadridentate Ligands





Tricarboxymethylamine


## 2.16.5 Pentadentate Ligands




Ethylenediaminetriacetic acid

## 2.16.6 Hexadentate Ligands

Ethylenediaminetetraacetic acid  
- EDTADiaminocyclohexanetetraacetic  
acid - CDTA



Diethylenetriaminepentaacetic  
acid - DTPA



Dioxaoctamethylenedinitriolo  
tetraacetic acid - PGTA

## Chapter 3

### **Organic Chemistry**

- 3.1 *Classification of Organic Compounds*
- 3.2 *Alkanes*
- 3.3 *Alkenes*
- 3.4 *Dienes*
- 3.5 *Alkynes*
- 3.6 *Benzene*
- 3.7 *Alkylbenzenes*
- 3.8 *Alkenylbenzenes*
- 3.9 *Alkyl Halides*
- 3.10 *Aryl Halides*
- 3.11 *Alcohols*
- 3.12 *Phenols*
- 3.13 *Ethers*
- 3.14 *Epoxides*
- 3.15 *Aldehydes and Ketones*
- 3.16 *Carboxylic Acids*
- 3.17 *Acid Chlorides*
- 3.18 *Esters*
- 3.19 *Acid Anhydrides*
- 3.20 *Amides*
- 3.21 *Amines*
- 3.22 *Alicyclic Compounds*
- 3.23 *Heterocyclic Compounds*
- 3.24 *Isomers*
- 3.25 *Polymer Structures*

### 3.1 Classification Of Organic Compounds

#### 3.1.1 General Classification














Figure 3.1. General Classification of Organic Compounds

#### 3.1.2 Classification by Functional Group

Table 3.1. Organic Functional Groups

Type	Functional Group	Example	Name
Alkane	R-H	CH <sub>3</sub> CH <sub>2</sub> CH <sub>3</sub>	propane
Alkene	R=R	CH <sub>2</sub> =CHCH <sub>3</sub>	propene
Diene	R=R-R=R	CH <sub>2</sub> =CH-CH=CH <sub>2</sub>	1,4-butene
Alkyne	R ≡ R	CH ≡ CH	ethyne
Halide	R-X	CH <sub>3</sub> CH <sub>2</sub> -Br	bromoethane
Alcohol	R-OH	CH <sub>3</sub> CH <sub>2</sub> -OH	ethanol
Ether	R-O-R	CH <sub>3</sub> CH <sub>2</sub> -O-CH <sub>2</sub> CH <sub>3</sub>	ethoxyether


Table 3.1. (Continued)

Epoxide			ethylene oxide
Aldehyde	R-CHO		ethanal
Ketone	R-CO-R		2-propanone
Carboxylic Acid	R-CO <sub>2</sub> H		ethanoic acid
Acid Chloride	R-CO-Cl		acetyl chloride
Acid Anhydride	(RCO) <sub>2</sub> O		acetic anhydride
Ester	R-CO <sub>2</sub> R		methyl ethanoate
Amide	R-CONH <sub>2</sub>		ethanamide
Amine 1°	R-NH <sub>2</sub>	CH <sub>3</sub> CH <sub>2</sub> -NH <sub>2</sub>	ethaneamine
Amine 2°	R-NH-R	CH <sub>3</sub> CH <sub>2</sub> -NH-CH <sub>2</sub> CH <sub>3</sub>	diethaneamine
Amine 3°	R <sub>3</sub> -N	(CH <sub>3</sub> CH <sub>2</sub> ) <sub>3</sub> -N	triethaneamine
Nitro Compound	R-NO <sub>2</sub>		nitromethane
Nitrile	R-C≡N	CH <sub>3</sub> C≡N	ethanenitrile
Thiol	R-SH	CH <sub>3</sub> CH <sub>2</sub> -SH	ethanethiol


## 3.2 Alkanes

### 3.2.1 Preparation of Alkanes



#### 3.2.1.1 Wurtz Reaction




#### 3.2.1.2 Grignard Reduction




#### 3.2.1.3 Reduction



#### 3.2.1.4 Kolbe Reaction




#### 3.2.1.5 Hydrogenation




## 3.2.2 Reactions of Alkanes

### 3.2.2.1 Combustion




### 3.2.2.2 Halogenation



Reactivity X:  $Cl_2 > Br_2$   
 H:  $3^\circ > 2^\circ > 1^\circ > CH_3-H$


### 3.2.2.3 Free Radical Substitution



## 3.3 Alkenes


### 3.3.1 Preparation of Alkenes

#### 3.3.1.1 Dehydrohalogenation of Alkyl Halides




Ease of dehydrohalogenation  $3^\circ > 2^\circ > 1^\circ >$



#### 3.3.1.2 Dehalogenation of Vicinal Dihalides



### 3.3.1.3 Dehydration of Alcohols




### 3.3.1.4 Reduction of Alkynes




## 3.3.2 Reactions of Alkenes

### 3.3.2.1 Hydrogenation





### 3.3.2.2 Halogenation




$\text{X}_2 = \text{Cl}_2, \text{Br}_2$

### 3.3.2.3 Addition of Hydrogen Halide




**Markovnikov's rule:** The hydrogen of the acid attaches itself to the carbon atom which already has the greatest number of hydrogens. In the presence of peroxide, HBr will undergo anti-Markovnikov addition.



### 3.3.2.4 Addition of Sulfuric Acid



### 3.3.2.5 Addition of Water




### 3.3.2.6 Halohydrin Formation




### 3.3.2.7 Oxymercuration-Demercuration




### 3.3.2.8 Hydroboration-Oxidation




### 3.3.2.9 Polymerization



### 3.3.2.10 Hydroxylation



### 3.3.2.11 Halogenation - Allylic Substitution




$$\text{X}_2 = \text{Cl}_2, \text{Br}_2$$

## 3.4 Dienes


Isolated dienes can be prepared following the methods for alkanes using difunctional starting materials.

### 3.4.1 Preparation of Conjugated Dienes

#### 3.4.1.1 Dehydration of 1,3-Diol





#### 3.4.1.2 Dehydrogenation




### 3.4.2 Reactions of Dienes

#### 3.4.2.1 1,4-Addition




#### 3.4.2.2 Polymerization




### 3.5 Alkynes

#### 3.5.1 Preparation of Alkynes

##### 3.5.1.1 Dehydrohalogenation of Alkyl Dihalides



##### 3.5.1.2 Dehalogenation of Tetrahalides





##### 3.5.1.3 Reaction of Water and Calcium Carbide




#### 3.5.2 Reactions of Alkynes

##### 3.5.2.1 Hydrogenation






### 3.5.2.2 Halogenation




$\text{X}_2 = \text{Cl}_2, \text{Br}_2$

### 3.5.2.3 Addition of Hydrogen Halide



$\text{X} = \text{Cl}, \text{Br}, \text{I}$


### 3.5.2.4 Addition of Water (Hydration)




## 3.6 Benzene

### 3.6.1 Preparation of Benzene


#### 3.6.1.1 Ring Formation



### 3.6.1.2 Cyclization




### 3.6.1.3 Elimination




## 3.6.2 Reactions of Benzenes


### 3.6.2.1 Nitration



### 3.6.2.2 Sulfonation




### 3.6.2.3 Halogenation




$\text{X} = \text{Cl, Br}$


### 3.6.2.4 Friedel-Crafts Alkylation




## 3.6.2.5 Friedel-Crafts Acylation




## 3.6.2.6 Hydrogenation



## 3.6.2.7 Bromination




## 3.6.2.8 Combustion



## 3.7 Alkylbenzenes

## 3.7.1 Preparation of Alkylbenzenes



## 3.7.1.1 Freidel-Crafts Alkylation




Lewis acid:  $\text{AlCl}_3, \text{BF}_3, \text{HF}$

Ar-H cannot be used in place of R-X


### 3.7.1.2 Side Chain Conversion



### 3.7.1.3 Electrophilic Aromatic Substitution




### 3.7.1.4 Hydrogenation




## 3.7.2 Reactions of Alkylbenzenes


### 3.7.2.1 Hydrogenation





### 3.7.2.2 Oxidation



## 3.7.2.3 Substitution in ring - electrophilic aromatic substitution




## 3.7.2.4 Substitution in the side chain




### 3.8 Alkenylbenzenes

#### 3.8.1 Preparation of Alkenylbenzenes


##### 3.8.1.1 Dehydrogenation



##### 3.8.1.2 Dehydrohalogenation




##### 3.8.1.3 Dehydration




#### 3.8.2 Reactions of Alkenylbenzenes


##### 3.8.2.1 Catalytic Hydrogenation



### 3.8.2.2 Oxidation




### 3.8.2.3 Ring Halogenation




## 3.9 Alkyl Halides

### 3.9.1 Preparation of Alkyl Halides


#### 3.9.1.1 From Alcohols




#### 3.9.1.2 Addition of Hydrogen Halide to Alkenes







#### 3.9.1.3 Halogenation of Alkanes



### 3.9.1.4 Halide Exchange




### 3.9.1.5 Halogenation of Alkenes and Alkynes



## 3.9.2 Reactions of Alkyl Halides

### 3.9.2.1 Addition of Hydroxide





### 3.9.2.2 Addition of Water




### 3.9.2.3 Addition of Alkoxide



**3.9.2.4     Addition of Carboxylate****3.9.2.5     Addition of Hydrosulfide****3.9.2.6     Addition of Thioalkoxide****3.9.2.7     Addition of Sulfide****3.9.2.8     Addition of Thiocyanide****3.9.2.9     Addition of Iodide****3.9.2.10    Addition of Amide****3.9.2.11    Addition of Ammonia**


### 3.9.2.12 Addition of 1° Amine




### 3.9.2.13 Addition of 2° Amine




### 3.9.2.14 Addition of 3° Amine



### 3.9.2.15 Addition of Azide



### 3.9.2.16 Addition of Nitrite



### 3.9.2.17 Addition of Phosphine





### 3.9.2.18 Addition of Cyanide



### 3.9.2.19 Addition of Alkynyl Anion





### 3.9.2.20 Addition of Carbanion



## 3.10 Aryl Halides

### 3.10.1 Preparation of Aryl Halides


#### 3.10.1.1 Halogenation by Substitution




#### 3.10.1.2 From Arylthallium Compounds



### 3.10.1.3 From Diazonium Salt





### 3.10.1.4 Halogenation by Addition




## 3.10.2 Reactions of Aryl Halides

### 3.10.2.1 Grignard Reagent Formation




### 3.10.2.2 Nucleophilic Aromatic Substitution



$\text{Z}^-$  = strong base

### 3.10.2.3 Electrophilic Aromatic Substitution



X deactivates and directs ortho, para in electrophilic aromatic substitution.



## 3.11 Alcohols

### 3.11.1 Preparation of Alcohols


#### 3.11.1.1 Addition of Hydroxide




#### 3.11.1.2 Grignard Synthesis



primary alcohol




secondary alcohol




tertiary alcohol

### 3.11.1.3 Reduction of Carbonyl Compounds




primary alcohol




secondary alcohol


### 3.11.1.4 Hydration of Alkenes




$\text{R} > \text{R}'$



### 3.11.1.5 Reaction of Amines with Nitrous Acid




### 3.11.1.6 Oxymercuration-Demercuration



Markovnikov addition


### 3.11.1.7 Hydroboration-Oxidation



Anti-Markovnikov addition

### 3.11.2 Reactions of Alcohols

#### 3.11.2.1 Reaction with Hydrogen Halides




Reactivity of HX: HI > HBr > HCl

Reactivity of ROH: allyl, benzyl > 3° > 2° > 1°


#### 3.11.2.2 Reaction with Phosphorus Trihalide




#### 3.11.2.3 Dehydration




#### 3.11.2.4 Ester Formation



### 3.11.2.5 Reaction with Active Metals




### 3.11.2.6 Oxidation




## 3.12 Phenols

### 3.12.1 Preparation of Phenols


#### 3.12.1.1 Nucleophilic Displacement of Halides




#### 3.12.1.2 Oxidation of Cumene



#### 3.12.1.3 Hydrolysis of Diazonium Salts



### 3.12.1.4 Oxidation of Arylthallium Compounds




### 3.12.2 Reactions of Phenols


#### 3.12.2.1 Salt Formation




#### 3.12.2.2 Ether Formation - Williamson Synthesis




#### 3.12.2.3 Ester Formation




### 3.12.2.4 Ring Substitution - Nitration





### 3.12.2.5 Ring Substitution - Sulfonation




### 3.12.2.6 Ring Substitution - Nitrosation




## 3.12.2.7 Ring Substitution - Halogenation




## 3.12.2.8 Ring Substitution - Friedel-Crafts Alkylation




## 3.12.2.9 Ring Substitution - Friedel-Crafts Acylation




### 3.12.2.10 Coupling with Diazonium Salts




### 3.12.2.11 Carbonation. Kolbe Reaction




### 3.12.2.12 Carboxylation. Kolbe Reaction




### 3.12.2.13 Aldehyde Formation. Reimer-Tiemann Reaction



### 3.12.2.14 Carboxylic Acid Formation. Reimer-Tiemann Reaction



### 3.12.2.15 Reaction with Formaldehyde



## 3.13 Ethers


### 3.13.1 Preparation of Ethers

#### 3.13.1.1 Williamson Synthesis




$R'$  = alkyl or aryl

#### 3.13.1.2 Oxymercuration-Demercuration






### 3.13.1.3 Dehydration of Alcohols






### 3.13.2 Reactions of Ethers


#### 3.13.2.1 Single Cleavage by Acids



#### 3.13.2.2 Double Cleavage by Acids




### 3.13.2.3 Substitution on the Hydrocarbon Chain




## 3.14 Epoxides

### 3.14.1 Preparation of Epoxides

#### 3.14.1.1 Halohydrin Reaction




#### 3.14.1.2 Peroxidation




### 3.14.2 Reactions of Epoxides


#### 3.14.2.1 Acid-Catalyzed Cleavage



### 3.14.2.2 Base-Catalyzed Cleavage





### 3.14.2.3 Grignard Reaction




## 3.15 Aldehydes And Ketones

### 3.15.1 Preparation of Aldehydes


#### 3.15.1.1 Oxidation



#### 3.15.1.2 Reduction




### 3.15.1.3 Reduction




### 3.15.2 Reactions Specific to Aldehydes

#### 3.15.2.1 Oxidation




#### 3.15.2.2 Cannizzaro Reaction




### 3.15.3 Preparation of Ketones

#### 3.15.3.1 Oxidation




#### 3.15.3.2 Friedel-Crafts Acylation



R' = aryl or alkyl

### 3.15.3.3 Grignard Reaction




### 3.15.4 Reactions Specific to Ketones

#### 3.15.4.1 Halogenation




#### 3.15.4.2 Oxidation




### 3.15.5 Reactions Common to Aldehydes And Ketones


#### 3.15.5.1 Reduction to Alcohol




#### 3.15.5.2 Reduction to Hydrocarbon




## 3.15.5.3 Grignard Reaction




## 3.15.5.4 Cyanohydrin Formation



## 3.15.5.5 Addition of Bisulfite




## 3.15.5.6 Addition of Ammonia Derivatives




G	Product
R	=NR
OH	=NOH
NH <sub>2</sub>	=NNH <sub>2</sub>
NHC <sub>6</sub> H <sub>5</sub>	=NNHC <sub>6</sub> H <sub>5</sub>
NHCONH <sub>2</sub>	=NNHCONH <sub>2</sub>

## 3.15.5.7 Aldol Condensation



### 3.15.5.8 Wittig Reaction



### 3.15.5.9 Acetal Formation




## 3.16 Carboxylic Acids

### 3.16.1 Preparation of Carboxylic Acids


#### 3.16.1.1 Oxidation of Primary Alcohols




#### 3.16.1.2 Oxidation of Alkylbenzenes



#### 3.16.1.3 Carbonation of Grignard Reagents





#### 3.16.1.4 Hydrolysis of Nitriles




### 3.16.2 Reactions of Carboxylic Acids

#### 3.16.2.1 Acidity Salt Formation




#### 3.16.2.2 Conversion to Acid Chloride




$\text{R}'$  = alkyl or aryl

#### 3.16.2.3 Conversion to Esters




$\text{R}$  = alkyl or aryl

#### 3.16.2.4 Conversion to Amides



$\text{R}'$  = alkyl or aryl

### 3.16.2.5 Reduction




$\text{R}'$  = alkyl or aryl

### 3.16.2.6 Alpha-Halogenation of Aliphatic Acids




### 3.16.2.7 Ring Substitution in Aromatic Acids



## 3.17 Acid Chlorides

### 3.17.1 Preparation of Acid Chlorides



$\text{R}'$  = alkyl or aryl


### 3.17.2 Reactions of Acid Chlorides

#### 3.17.2.1 Hydrolysis (Acid Formation)




$\text{R}'$  = alkyl or aryl

#### 3.17.2.2 Ammonolysis (Amide Formation)




$\text{R}'$  = alkyl or aryl

#### 3.17.2.3 Alcoholysis (Ester Formation)




$\text{R}'$  = alkyl or aryl

#### 3.17.2.4 Friedel-Crafts Acylation (Ketone Formation)




$\text{R}'$  = alkyl or aryl

### 3.17.2.5 Ketone Formation by Reaction with Organocadium Compounds




R' must be an acyl or primary alkyl alcohol


### 3.17.2.6 Aldehyde Formation by Reduction



### 3.17.2.7 Rosenmund Reduction




### 3.17.2.8 Reduction to Alcohols




### 3.18 Acid Anhydrides

### 3.18.1 Preparation of Acid Anhydrides

### 3.18.1.1 Ketene Reaction




### 3.18.1.2 Dehydration of Dicarboxylic Acids



$n = 2, 3, 4$

### 3.18.2 Reactions of Acid Anhydrides

#### 3.18.2.1 Hydrolysis (Acid Formation)




$\text{R}'$  = alkyl or aryl

#### 3.18.2.2 Ammonolysis (Amide Formation)




$\text{R}'$  = alkyl or aryl

#### 3.18.2.3 Alcoholsysis (Ester Formation)



$\text{R}'$  = alkyl or aryl

#### 3.18.2.4 Friedel-Crafts Acylation (Ketone Formation)



### 3.19 Esters


#### 3.19.1 Preparation of Esters

##### 3.19.1.1 From Acids




$R'$  = alkyl or aryl

##### 3.19.1.2 From Acid Chlorides




$R'$  = alkyl or aryl


##### 3.19.1.3 From Acid Anhydrides



##### 3.19.1.4 Transesterification




##### 3.19.1.5 From Ketene and Alcohols




### 3.19.2 Reactions of Esters

#### 3.19.2.1 Hydrolysis




$\text{R}'$  = alkyl or aryl

#### 3.19.2.2 Saponification




$\text{R}'$  = alkyl or aryl


#### 3.19.2.3 Ammonolysis




#### 3.19.2.4 Transesterification




#### 3.19.2.5 Grignard Reaction




### 3.19.2.6 Hydrogenolysis




### 3.19.2.7 Bouvauel - Blanc Method



### 3.19.2.8 Chemical Reduction



### 3.19.2.9 Claisen Condensation




## 3.20 Amides

### 3.20.1 Preparation of Amides


#### 3.20.1.1 From Acid Chlorides





$R'$  = alkyl or aryl


### 3.20.1.2 From Acid Anhydrides




### 3.20.1.3 From Esters by Ammonolysis




### 3.20.1.4 From Carboxylic Acids



### 3.20.1.5 From Nitriles




### 3.20.1.6 From Ketenes and Amines




### 3.20.2 Reactions of Amides

#### 3.20.2.1 Hydrolysis




$\text{R}'$  = alkyl or aryl


#### 3.20.2.2 Conversion to Imides




#### 3.20.2.3 Reaction with Nitrous Acid




#### 3.20.2.4 Dehydration



#### 3.20.2.5 Reduction





#### 3.20.2.6 Hoffman Degradation




### 3.21 Amines

#### 3.21.1 Preparation of Amines




##### 3.21.1.1 Reduction of Nitro Compounds




##### 3.21.1.2 Reaction of Ammonia with Halides



##### 3.21.1.3 Reductive Amination




### 3.21.1.4 Reduction of Nitriles



$\text{R}'$  = alkyl or aryl



### 3.21.1.5 Hofmann Degradation




$\text{R}'$  = alkyl or aryl

## 3.21.2 Reactions of Amines

### 3.21.2.1 Alkylation






### 3.21.2.2 Salt Formation





$\text{R}'$  = alkyl or aryl

### 3.21.2.3 Amide Formation






### 3.21.2.4 Reaction of Amines with Nitrous Acid



## 3.22 Alicyclic Compounds

### 3.22.1 Preparation of Alicyclic Compounds

#### 3.22.1.1 Cyclization




When  $n = 1$  cyclopropane


$n = 2$  cyclobutane

$n = 3$  cyclopentane

#### 3.22.1.2 Hydrogenation




#### 3.22.1.3 Cycloaddition




$\text{X} = \text{H, Cl, Br}$


### 3.22.2 Reactions of Alicyclic Compounds

#### 3.22.2.1 Free Radical Addition






### 3.22.2.2 Addition Reaction




## 3.23 Heterocyclic Compounds

### 3.23.1 Preparation of Pyrrole, Furan, and Thiophene


#### 3.23.1.1 Pyrrole



#### 3.23.1.2 Furan





#### 3.23.1.3 Thiophene






### 3.23.2 Reactions of Pyrrole, Furan, and Thiophene


#### 3.23.2.1 Pyrrole






#### 3.23.2.2 Furan





#### 3.23.2.3 Thiophene








### 3.23.3 Preparation of Pyridine, Quinoline, and Isoquinoline

#### 3.23.3.1 Pyridine




#### 3.23.3.2 Quinoline









### 3.23.3.3 Isoquinoine




## 3.23.4 Reactions of Pyridine, Quinoline, and Isoquinoline

### 3.23.4.1 Pyridine






### 3.23.4.2 Quinoline



### 3.23.4.3 Isoquinoline



## 3.24 Isomers

### 3.24.1 Isomers and Stereoisomers

Organic compounds that have the same chemical formula but are attached to one another in different ways are called **isomers**. Isomers that have the same chemical formula and are attached to one another in same way but whose orientation to one another differ are called **Stereoisomers**. There are several different types of isomers that are encountered in organic chemistry.

To represent three dimensional structures on paper the chiral center of a molecule is taken at the cross point of a cross and the groups are attached at

the ends. The horizontal line represents the bonds projecting out of the plane of the paper. The vertical line represents the bonds projecting into the plane of the paper.




Figure 3.2. Three Dimensional Representations

### 3.24.2 Optical Activity

In addition to having different arrangements of atoms, certain organic compounds exhibit a unique property of rotating **plane-polarized light** (light that has its amplitude in one plane). Compounds that rotate light are said to be **optically active**. Optically active compounds that rotate light to the right are called dextrorotatory and are symbolized by D or +. Compounds that rotate light to the left are called levorotatory and are symbolized by L or -.

### 3.24.3 Enantiomers

Enantiomers are stereoisomers that are non-superimposable mirror images of each other. Enantiomers have identical physical and chemical (except towards optically active reagents) properties except for the direction in which plane-polarized light is rotated. Enantiomers account for a compound's optical activity.




Figure 3.3. Enantiomers

### 3.24.4 Chirality

Molecules that are not superimposable on their mirror images are **chiral**. Chiral molecules exist as enantiomers but achiral molecules cannot exist as enantiomers. A carbon atom to which four different groups are attached is a **chiral center**. Not all molecules that contain a chiral center are chiral. Not all chiral molecules contain a chiral center.

### 3.24.5 Racemic Mixture

A mixture of equal parts of enantiomers is called a **racemic mixture**. A racemic mixture contains equal parts of D and L components and therefore, the mixture is optically inactive.

### 3.24.6 Diastereomers

**Diastereomers** are stereoisomers that are not mirror images of each other. Diastereomers have different physical properties. And they maybe dextrorotatory, levorotatory or inactive.




Figure 3.4 Enantiomers and Diastereomers

Structures 1 and 2 are enantiomers, structure 3 is a diastereomer of structures 1 and 2.

### 3.24.7 Meso Compounds

Meso compounds are superimposable mirror images of each other, even though they contain a chiral center.




Figure 3.5. Meso Compound

Meso compounds can be recognized by the fact that half the molecule is a mirror image of the other half.




Figure 3.6. Plane of symmetry of a meso compound

The upper half of the molecule is a non-superimposable mirror image of the lower half, making the top half an enantiomer of the lower half. However, since the two halves are in the same molecule the rotation of plane-polarized light by the upper half is cancelled by the lower half and the compound is optically inactive.

### 3.24.8 Positional Isomers

**Positional isomers** are compounds that have the same number and kind of atoms but are arranged (or bonded) in a different order. They also have different physical and chemical properties. Butane (figure 3.1) can have two different structures, n-butane and 2-methylpropane:




Figure 3.7. n-butane and 2-methylpropane

### 3.24.9 Geometric Isomers

**Geometric isomers** or cis-trans isomerism can exist in compounds that contain a double bond or a ring structure. In order for this type of isomerism to exist the groups coming off the same end of the double bond must be different. For example bromoethene does not have cis-trans isomerism.




Figure 3.8. Bromoethene structure

However, 1,2-dibromoethene can exist as cis-1,2-dibromoethene and trans-1,2-dibromoethene




Figure 3.9. Trans and cis butene

Ring structures confer restricted rotation around the bonds and thereby give rise to geometric isomer. In trans-1,3-dichlorocyclopentane one chlorine is above the plane of the ring and one is below. In cis-1,3-dichlorocyclopentane both chlorines are above the plane of the ring.




Figure 3.10. Trans and cis- 1,3-cyclopropane

### 3.24.10 Conformational Isomers

**Conformational isomers** deal with the orientations within a molecule. The free rotation around a single bond accounts for the different

conformations that can exist within a molecule. For example, n-butane can have the following conformations:




Figure 3.11. Conformational isomers of butane

Figure 1 has a *staggered* or *anti* conformation. Since in figure 1, the two methyl groups are farthest apart, this form is referred to as *anti*. Figures 3 and 5 have *staggered* or *gauche* conformations. Figures 2, 4, and 6 have an *eclipsed* conformation.

A different type of projection used to view isomers, called Newman projections, is sometimes used. The following figures are the same as above:




Figure 3.12. Newman projections




Figure 3.12. Newman projections

Although cyclohexane is a ring structure it does have free rotation around single bonds. Cyclohexane has two main conformations. The most stable form is called the chair form, the less stable is called the boat form:



Figure 3.13. Chair and boat conformations of cyclohexane

The bonds in cyclohexane occupy two kinds of position, six hydrogens lie in the plane and six hydrogens lie either above or below the plane. Those that are in the plane of the ring lie in the “equator” of it, and are called the **equatorial bonds**. Those bonds that are above or below are pointed along the axis perpendicular to the plane and are called **axial bonds**.




Figure 3.14. Equatorial and axial bonds of cyclohexane

### 3.24.11 Configurational Isomers

The arrangement of atoms that characterizes a certain stereoisomer is called its configuration. In general, optically active compounds can have more than one configuration. Determination of the Configuration can be determined by the following two steps:

Step 1. Following a set of sequence rule we assign a sequence of priority to the four atoms attached to the chiral center.

Step 2. The molecule is oriented so that the group of lowest priority is directed away from us. The arrangement of the remaining groups is then observed. If the sequence of highest priority to lowest priority is clockwise, the configuration is designated R. If the sequence is counterclockwise, the configuration is designated S.

From these steps a set of sequence rules can be formulated that will allow a configuration to be designated as either R or S.

Sequence 1. If the four atoms attached to the chiral center are all different, priority depends on the atomic number, with the atom of higher atomic number getting higher priority.

Sequence 2. If the relative priority of two groups cannot be decided by Sequence 1, it shall be determined by a similar comparison of the atoms attached to it.

For example, bromochloroiodomethane,  $\text{CHClBrI}$ , has two possible configurations as shown in figure 3.8. Using the sequence rules, the order of the atoms for the configuration is I, Br, Cl, H. However, the figure on the left has a different sequence than the one on the right. Hence, (R)-bromochloroiodomethane and (S)-bromochloroiodomethane





Figure 3.15. R and S configuration

### 3.25 Polymer Structures


The following listing of common polymers provides respective structure. The reader should note that the name of the polymer often provides the key to its representative structure. There are, however, names such as polycarbonate that can represent a variety of polymeric materials.

Acrylonitrile-butadiene-styrene terpolymer (ABS):




### Buna-N:

## Elastomeric copolymer of butadiene and acrylonitrile.




## Buna-S:


## Elastomeric copolymer of butadiene and styrene.




## Butyl rubber:




Cellulose:




Epoxy resins:



Ethylene-methacrylic acid copolymers (ionomers):




Ethylene-propylene elastomers:




Formaldehyde resins:


*Phenol-formaldehyde (PF):*




### Urea-formaldehyde (UF):




### *Melamine-formaldehyde (MF):*




Nitrile rubber (NBR):



## Polyacrylamide:




Polyacrylonitrile:




Polyamides (nylons):

*Nylon 6,6 and nylon 6:*




nylon 6,6




nylon 6

*Aromatic polyamides:*




Nomex




Kevlar

*Polyamide imides and polyimides:*




poly(amide imides)




polyimides


Polybutadiene (butadiene rubber, BR):



Polycarbonate (PC):




Polychloroprene:




Polyesters:


*Poly(ethylene terephthalate) (PET):*




*Poly(butylene terephthalate) (PBT):*




*Aromatic polyesters:*



Polyether (polyoxymethylene; polyacetal):



Polyethylene (PE):




*Low-density polyethylene (LDPE).*

*High-density polyethylene (HDPE).*


*Linear low-density polyethylene (LLDPE).*

*Ultrahigh molecular weight polyethylene (UHMWPE).*


Poly(ethylene glycol) (PEG):




Polyisobutylene (PIB):




Polyisoprene:



Poly(methyl methacrylate) (PMMA):




Poly(phenylene oxide) (PPO):




**poly(2,6-dimethyl-p-phenylene ether)**


Poly(phenylene sulfide) (PPS):




Polyphosphazenes:




Polypropylene (PP):




Polystyrene (PS):




Poly sulfone:




Polytetrafluoroethylene (PTFE):




Polyurethane:




Poly(vinyl acetate) (PVA):




Poly(vinyl alcohol) (PVAL):




Poly(vinyl butyral) (PVB):




Poly(vinyl carbazole):




Poly(vinyl chloride) (PVC):




Poly(vinyl formal) (PVF):




Poly(vinylidene chloride):




Poly(vinyl pyridine):




Poly(vinyl pyrrolidone):



Silicones (siloxanes):




Starch:




linear amylose

Styrene-acrylonitrile copolymer (SAN):



Styrene-butadiene rubber (SBR):



## Chapter 4

### Instrumental Analysis

- 4.1 *The Electromagnetic Spectrum*
- 4.2 *Ultraviolet-Visible Spectrum*
- 4.3 *Infrared Spectrum*
- 4.4 *Nuclear Magnetic Resonance*
- 4.5 *Mass Spectroscopy*

### 4.1 The Electromagnetic Spectrum

#### 4.1.1 The Electromagnetic Spectrum

The various regions of the electromagnetic spectrum are somewhat arbitrary and not very sharply defined. Each region overlaps at both ends the adjacent regions. The units defining the different regions can be in frequency (Hz,  $\text{cm}^{-1}$ ), energy (kcal/mole, eV) and wavelengths (m and  $\lambda$ , "official" or most common unit used for a given region). Some useful relationships and constants are the following: 1 nanometer (nm) = 1 millimicron ( $\text{m}\mu$ ) , $=10^{-9}$  m; 1  $\mu\text{m}$  (formerly micron) =  $10^{-6}$  m = 10 mm; 1 angstrom ( $\text{\AA}$ ) = 0.1 nm =  $10^{-8}$  cm; 1 eV = 23.06 kcal/mole = 8063  $\text{cm}^{-1}$ . Energy, E =  $h\nu$  [ $h$  = Planck's constant =  $9.534 \times 10^{-14}$  kcal-sec/mole =  $1.583 \times 10^{-34}$  cal-sec/molecule;  $\nu$  in Hz (cycles/sec)]. E in kcal/mole =  $2.8635/\lambda$  ( $\lambda$  in  $\mu\text{m}$ ); E in eV =  $12,3451$  ( $\lambda$  in  $\text{\AA}$  ).

Table 4.1 The Electromagnetic Spectrum

Region	Spectral Use	Hz	cm <sup>-1</sup>	kcal/mole	eV	m	$\lambda$
Cosmic rays		10 <sup>22</sup>	3x10 <sup>11</sup>	9.5x10 <sup>8</sup>	4.1x10 <sup>7</sup>	3x10 <sup>-14</sup>	3x10 <sup>-4</sup> Å
Gamma rays	Nuclear transitions	10 <sup>19</sup>	3x10 <sup>7</sup>	9.5x10 <sup>5</sup>	4.1x10 <sup>4</sup>	3x10 <sup>-10</sup>	3 Å
Soft X-rays	Inner electron transitions	10 <sup>17</sup>	1x10 <sup>5</sup>	9.5x10 <sup>3</sup>	4.1x10 <sup>2</sup>	3x10 <sup>-8</sup>	300 Å
Vacuum uv	Valance electron transitions	1.5x10 <sup>15</sup>	5x10 <sup>4</sup>	143	6.2	2x10 <sup>-7</sup>	200 nm
Quartz uv	(electronic spectra)	7.5x10 <sup>14</sup>	2.5x10 <sup>4</sup>	71.5	3.1	4x10 <sup>-7</sup>	400 nm
Visible		3.8x10 <sup>14</sup>	1.2x10 <sup>4</sup>	36.2	1.6	8x10 <sup>-7</sup>	800 nm
Near ir	Vibrational ir overtone and combination region	1.2x10 <sup>14</sup>	4x10 <sup>3</sup>	12	0.52	2.5x10 <sup>-6</sup>	2.5 $\mu$ m
Vibrational ir	Fundamental region	2.5x10 <sup>13</sup>	8x10 <sup>2</sup>	2.4	0.10	1.2x10 <sup>-5</sup>	12.5 $\mu$ m
Far ir	Sketal, ring torsional, solid state (lattice) deformations	10 <sup>12</sup>	33	9.5x10 <sup>-2</sup>	4.1x10 <sup>-3</sup>	3x10 <sup>-4</sup>	300 $\mu$ m
Microwave	Molecular rotations; bond torsions	10 <sup>9</sup>	3.3x10 <sup>-2</sup>	9.5x10 <sup>-5</sup>	4.1x10 <sup>-6</sup>	3x10 <sup>-1</sup>	300 mm
Short Radio	Spin orientations	1.5x10 <sup>6</sup>	5x10 <sup>-5</sup>	1.4x10 <sup>-7</sup>	6.1x10 <sup>-9</sup>	2x10 <sup>2</sup>	200 m
Broadcast	Radio, tv, etc.	5.5x10 <sup>5</sup>	1.8x10 <sup>-5</sup>	5.2x10 <sup>-8</sup>	2.2x10 <sup>-9</sup>	5.6x10 <sup>2</sup>	550 m
Long Radio	Induction heating; longwave communication	3x10 <sup>3</sup>	10 <sup>-7</sup>	2.9x10 <sup>-10</sup>	1.3x10 <sup>-11</sup>	10 <sup>5</sup>	105 m
Electric Power	Commercial power and light	3x10 <sup>-1</sup>	10 <sup>-11</sup>	2.9x10 <sup>-14</sup>	1.3x10 <sup>-14</sup>	10 <sup>9</sup>	10 <sup>9</sup> m

(From The Chemist's Companion, Gordon &amp; Ford, Copyright © 1972 John Wiley &amp; Sons, Inc. Reprinted with Permission of John Wiley &amp; Sons, Inc.)

## 4.2 Ultraviolet-Visible Spectroscopy

### 4.2.1 Solvents

Table 4.2. UV-Vis Solvents

Substance	Formula	BP (°C)	Cut-off (nm)	
			1 mm	10 mm
2-Methylbutane	C <sub>4</sub> H <sub>9</sub> CH <sub>3</sub>	28	---	210
Pentane	C <sub>5</sub> H <sub>12</sub>	36	---	200
Hexane	C <sub>6</sub> H <sub>14</sub>	69	190	200
Heptane	C <sub>7</sub> H <sub>16</sub>	96	---	195
<i>i</i> -Octane	C <sub>8</sub> H <sub>18</sub>	98	190	205
Cyclopentane	C <sub>5</sub> H <sub>10</sub>	49	---	210
Methylcyclopentane	C <sub>5</sub> H <sub>9</sub> CH <sub>3</sub>	72	---	200
Cyclohexane	C <sub>6</sub> H <sub>12</sub>	81	190	195
Methylcyclohexane	C <sub>6</sub> H <sub>11</sub> CH <sub>3</sub>	101		205
Benzene	C <sub>6</sub> H <sub>6</sub>	80	275	280
Toluene	C <sub>6</sub> H <sub>5</sub> CH <sub>3</sub>	110	280	285
<i>m</i> -Xylene	C <sub>6</sub> H <sub>4</sub> (CH <sub>3</sub> ) <sub>2</sub>	139	285	290
Decalin	C <sub>10</sub> H <sub>18</sub>	190	---	200
Water	HOH	100	187	191
Methanol	CH <sub>3</sub> OH	65	200	205
Ethanol	CH <sub>3</sub> CH <sub>2</sub> OH	78	200	205
2-Propanol	CH <sub>3</sub> CH(OH)CH <sub>3</sub>	81	200	210
Glycerol	(HOCH <sub>2</sub> ) <sub>2</sub> CH(OH)	290d	200	205
Sulfuric acid	96% H <sub>2</sub> SO <sub>4</sub>	300	---	210
Diethyl ether	Et <sub>2</sub> O	35	205	215
THF	CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> O	67	---	220
1,4-Dioxan	(OCH <sub>2</sub> CH <sub>2</sub> ) <sub>2</sub>	102	210	220
Dibutyl ether	(C <sub>4</sub> H <sub>9</sub> ) <sub>2</sub> O	142	---	210
Carbon disulfide	CS <sub>2</sub>	46	---	380
Chloroform	CHCl <sub>3</sub>	60	230	245
Carbon tetrachloride	CCl <sub>4</sub>	76	245	260
Methylene chloride	CH <sub>2</sub> Cl <sub>2</sub>	40	220	230
1,1-Dichloroethane	ClCH <sub>2</sub> CH <sub>2</sub> Cl	84	220	230
1,1,2,2-Dichloroethene	Cl <sub>2</sub> C=CCl <sub>2</sub>	120	280	290
Tribromomethane	CHBr <sub>3</sub>	150	315	330
Bromotrichloromethane	BrCCl <sub>3</sub>	105	320	340
Acetonitrile	CH <sub>3</sub> CN	81	190	195
Methylnitrite	CH <sub>3</sub> NO <sub>2</sub>	100	360	380
Pyridine	C <sub>5</sub> H <sub>5</sub> N	112	300	305
N,N-DMF	HCON(CH <sub>3</sub> ) <sub>2</sub>	152	---	270
DMSO	(CH <sub>3</sub> ) <sub>2</sub> SO	189d	250	265
Acetone	CH <sub>3</sub> COCH <sub>3</sub>	56	320	330

## 4.2.2 Woodward's Rules for Diene Absorption

Parent heteroannular diene	214
Parent homoannular diene	253
Add for each substituent:	
Double bond extending conjugation	30
Alkyl substituent, ring residue, etc.	5
Exocyclic double bond	5
N(alkyl) <sub>2</sub>	60
S(alkyl)	30
O(alkyl)	6
OAc	0
$\lambda_{\text{calc max}} = \text{TOTAL, nm}$	

## 4.2.3 Selected UV-Vis Tables

Table 4.3. Characteristics of Simple Chromophoric Groups

Chromophore	Example	$\lambda_{\text{max}}$ , m $\mu$	$\epsilon_{\text{max}}$	Solvent
C=C	Ethylene	171	15,530	Vapor
	1-Octene	177	12,600	Heptane
C≡C	2-Octyne	178	10,000	Heptane
		196	2,100	Heptane
C=O	Acetaldehyde	223	160	Heptane
		160	20,000	Vapor
	Acetone	180	10,000	Vapor
		290	17	Hexane
		166	16,000	Vapor
		189	900	Hexane
		279	15	Hexane
		208	32	Ethanol
-CO <sub>2</sub> H	Acetic acid	220	100	Hexane
-COCl	Acetyl chloride	178	9,500	Hexane
-CONH <sub>2</sub>	Acetamide	220	63	Water
-CO <sub>2</sub> R	Ethyl acetate	211	57	Ethanol
-NO <sub>2</sub>	Nitromethane	201	5,000	Methanol
		274	17	Methanol
-ONO <sub>2</sub>	Butyl nitrate	270	17	Ethanol
-ONO	Butyl nitrite	220	14,500	Hexane
		356	87	Hexane
-NO	Nitrosobutane	300	100	Ether
		665	20	Ether
C=N	<i>neo</i> -Pentylidene n-butylamine	235	100	Ethanol
-C≡N	Acetonitrile	167	weak	Vapor
-N <sub>3</sub>	Azidoacetic ester	285	20	Ethanol
=N <sub>2</sub>	Diazomethane	410	3	Vapor
-N=N-	Azomethane	338	4	Ethanol

Table 4.4. Characteristics of Simple Conjugated Chromophoric Groups

Chromophore	Example	$\lambda_{\max}$ , m $\mu$	$\epsilon_{\max}$	Solvent
C=C-C=C	Butadiene	217	20,900	Hexane
C=C-C≡C	Vinylacetylene	219	7,600	Hexane
		228	7,800	Hexane
C=C-C=O	Crotonaldehyde	218	18,000	Ethanol
		320	30	Ethanol
	3-Penten-2-one	224	9,750	Ethanol
		314	38	Ethanol
-C≡C-C=O	1-Hexyn-3-one	214	4,500	Ethanol
		308	20	Ethanol
C=C-CO <sub>2</sub> H	cis-Crotonic acid	206	13,500	Ethanol
		242	250	Ethanol
-C≡C-CO <sub>2</sub> H	n-Butylpropionic acid	210	6,000	Ethanol
C=C-C=N-	N-n-Butylcrotonaldimine	219	25,000	Hexane
C=C=C≡N	Methacrylonitrile	215	680	Ethanol
C=C=NO <sub>2</sub>	1-Nitro-1-propene	229	9,400	Ethanol
		235	9,800	Ethanol
HO <sub>2</sub> C-CO <sub>2</sub> H	Oxalic acid	185	4,000	Water

Table 4.5. Ultraviolet Absorption of Some Monosubstituted Benzenes (in Water)

C <sub>6</sub> H <sub>5</sub> X	Primary Band		Secondary Band	
	X=	$\lambda_{\max}$ , m $\mu$	$\epsilon_{\max}$	$\lambda_{\max}$ , m $\mu$
-H		203.5	7,400	254
-NH <sub>3</sub>		203	7,500	254
-CH <sub>3</sub>		206.5	7,000	261
-I		207	7,000	257
-Cl		209.5	7,400	263.5
-Br		210	7,900	261
-OH		210.5	6,200	270
-OCH <sub>3</sub>		217	6,400	269
-SO <sub>2</sub> NH <sub>2</sub>		217.5	9,700	264.5
-CN		224	13,000	271
-CO <sub>2</sub> <sup>-</sup>		224	8,700	268
-CO <sub>2</sub> H		230	11,600	273
-NH <sub>2</sub>		230	8,600	280
-O <sup>-</sup>		235	9,400	287
-NHCOCH <sub>3</sub>		238	10,500	---
-COCH <sub>3</sub>		245.5	9,800	---
-CHO		249.5	11,400	---
-NO <sub>2</sub>		268.5	7,800	---

## 4.3 Infrared Spectroscopy

### 4.3.1 Infrared Media

Table 4.6. IR media

Substance	Usable Regions of the Spectrum: $\text{cm}^{-1}$ ( $\mu\text{m}$ )		
	Near IR <sup>1</sup>	Mid IR	Far IR
KBr	10000 - 3333 (1-3)	5000 - 667 (2-15)	800 - 250 (12.5-40)
KCl	10000 - 3333 (1-3)	5000 - 667 (2-15)	800 - 526 (12.5-19)
CsBr	10000 - 3333 (1-3)	5000 - 667 (2-15)	800 - 250 (12.5-40)
CsI	10000 - 3333 (1-3)	5000 - 667 (2-15)	800 - 130 (12.5-77)
AgCl	10000 - 3333 (1-3)	5000 - 667 (2-15)	800 - 530 (12.5-19)
TlCl	10000 - 3333 (1-3)	5000 - 667 (2-15)	800 - 530 (12.5-19)
Polyethylene	10000 - 3333 (1-3) <sup>2</sup>	2500 - 1540 (4-6.5) 1250 - 741 (8-13.5)	625 - 278 (16-36)
Polystyrene			400 - 278 (25-36) <sup>3</sup>
Teflon		5000 - 1333 (2-7.5) <sup>4</sup> 1111 - 690 (9-14.5)	
Kel-F		5000 - 1333 (2-7.5) 870 - 690 (11.5-14.5)	
Nujol <sup>5</sup>		5000 - 3333 (2-3) 2500 - 1540 (4-6.5) 1250 - 667 (8-15)	667 - 286 (15-35)
Fluorolube <sup>6</sup>		5000 - 1430 (2-7)	
$\text{C}_4\text{Cl}_6$ <sup>7</sup>		5000 - 1667 (2-6) 1430 - 1250 (7-8)	

(From The Chemist's Companion, Gordon & Ford, Copyright © 1972 John Wiley & Sons, Inc. Reprinted with permission of John Wiley & Sons, Inc.)

1. Very little work has been reported for near ir spectra of pellets or mulls; most measurements are made in solution.
2. For a 0.1 mm sheet when compensated; otherwise, interfering bands occur at 1.9, 2.3, and 2.6 - 2.8  $\mu\text{m}$ .
3. For 0.025 mm sheet.
4. For 0.01 mm thick specimen; Teflon and Kel-F powders are very good for observing details in the OH stretch region.
5. Common brand of heavy mineral oil.
6. A saturated chlorofluorocarbon oil.
7. Hexachloro-1,3-butadiene.

## 4.3.2 Infrared Absorption Frequencies of Functional Groups

Table 4.7. Saturated Compounds sp = sharp, br = broad, (w) = weak, (s) = strong

Functional Group	Absorption Range (cm <sup>-1</sup> )	Example (cm <sup>-1</sup> )	Example Compound
<b>SATURATED COMPOUNDS</b>			
<b>Linear</b>			
CH <sub>3</sub> asymmetric	2970-2950	2967	<i>n</i> -Octane
CH <sub>3</sub> symmetric	2885-2865	2868	
CH <sub>3</sub> asymmetric	1465-1440	1466	
CH <sub>3</sub> symmetric	1380-1370	1380	
CH <sub>2</sub> asymmetric	2930-2915	2920	<i>n</i> -Octane
CH <sub>2</sub> symmetric	2860-2840	2854	
CH <sub>2</sub>	1480-1450	1470	
(CH <sub>2</sub> ) <sub>n</sub> , n > 4	723-720	723	<i>n</i> -Octane
	735-725	733	<i>n</i> -Pentane
	755-735	741	2-Methylpentane
	800-770	781	<i>n</i> -Propane
<b>Branched</b>			
CH	2890	2890	Triphenylmethane
	1340	1341	
(CH <sub>3</sub> ) <sub>2</sub> -CH-	1385-1380	1384	2-Methylheptane
	1372-1366	1366	
	1175-1165		
	1160-1140		
	922-917		
(CH <sub>3</sub> ) <sub>3</sub> -C-	1395-1380	1393	2,2-Dimethylhexane
	1375-1365	1366	
	1252-1245		
	1225-1195		
	930-925		
-C(CH <sub>3</sub> )-C(CH <sub>3</sub> )-	1165-1150	1160	3,4-Dimethylhexane
	1130-1120	1122	
	1080-1065	1071	
(CH <sub>3</sub> ) <sub>2</sub> -C-R <sub>2</sub>	1391-1381	1389	3,3-Dimethylhexane
	1220-1190	1192	
	1195-1185	1189	
(C <sub>2</sub> H <sub>5</sub> ) <sub>2</sub> -CH-R	1250	1250	3-Ethylhexane
	1150	1155	
	1130	1131	
C-C(CH <sub>3</sub> )-C	1160-1150		

Table 4.7. (Continued)

<b>Cyclic Compounds</b>			
Cyclopropane derivatives	3100-3072 3033-2995 1030-1000	3075 3028 1024	Cyclopropane
Cyclobutane derivatives	3000-2975 2924-2874 1000-960 or 930-890	2974 2896 901	Cyclobutane
Cyclopentane derivatives	2959-2952 2870-2853 1000-960 930-890	2951 2871 968 894	Cyclopentane
Cyclohexane derivatives	1055-1000 1015-950	1038 1014	Cyclohexane

Table 4.8. Unsaturated Compounds

Functional Group	Absorption Range (cm <sup>-1</sup> )	Example (cm <sup>-1</sup> )	Example Compound
<b>UNSATURATED COMPOUNDS</b>			
<b>Isolated -C=C- bonds</b>			
CH <sub>2</sub> =CH-	3095-3075 3030-2990 1648-1638 1420-1410 1000-980 915-905	3096 2994 1645 1420 994 912	1-Butene
CH <sub>2</sub> =C	3095-3075 1660-1640 1420-1410 895-885	3096 1661 1420 887	Methylpropene
-CH=C	3040-3010 1680-1665 1350-1340 840-805	3037 1675 1351 812	3-Methyl-2-pentene
-CH=CH- (cis)	3040-3010 1660-1640 1420-1395 730-675	3030 1661 1406 675	cis-2-Butene

Table 4.8. (Continued)

-CH=CH-( <i>trans</i> )	3040-3010 1700-1670 1310-1295 980-960	3021 1701 1302 964	<i>trans</i> -2-Butene
<b>Conjugated -C=C- bonds</b>			
-C=C-C=C-	1629-1590 1820-1790	1592 1821	1,3-Butadiene
<b>Allenic -C=C- bonds</b>			
-C=C=C-	1960-1940 1070-1060		
<b>-C≡C- bonds</b>			
-C≡C-	2270-2250	2268	2-Pentyne
<b>-C≡CH groups</b>			
CH (stretch)	3320-3300*	3320	1-Butyne
-C≡C-	2140-2100	2122	
CH (bend)	700-600		
* CCl <sub>4</sub> Solutions only			

Table 4.9. Aromatic Compounds

Functional Group	Absorption Range (cm <sup>-1</sup> )	Example (cm <sup>-1</sup> )	Example Compound
<b>AROMATIC COMPOUNDS</b>			
<b>General</b>			
CH	3060-3010		
CH substitution bands			
overtones	2000-1650 (w)		
C=C	1620-1590 sp 1590-1560 sp		
CH	1510-1480 sp 1450 sp		
<b>Mono-substitution</b>			
	1175-1125	1170	Toluene
	1110-1070	1088	
	1070-1000	1032	
	765-725	728 (s)	
	720-690	693 (s)	
<b>Di-substitution</b>			
<i>ortho</i>	1225-1175 1125-1090 1070-1000 765-735	1185 1121 1053 741 (s)	<i>o</i> -Xylene
<i>meta</i>	1175-1125 1110-1070 1070-1000 900-770 710-690	1171 1095 1039 769 (s) 690 (s)	<i>m</i> -Xylene

Table 4.9. (Continued)

<i>para</i>	1225-1175 1125-1090 1070-1000 855-790	1219 1120 1043 796 (s)	<i>p</i> -Xylene
<b>Tri-substitution</b>			
1,2,3-	1175-1125 1110-1070 1000-960 800-755 740-695	1162 1095 1009 765 (s) 710 (s)	1,2,3-Trimethylbenzene
1,2,4-	1225-1175 1130-1090 1000-960 900-865 855-800	1156 1130 1000 873 (s) 805 (s)	1,2,4-Trimethylbenzene
1,3,5-	1175-1125 1070-1000 860-810 705-685	1165 1039 836 (s) 690 (s)	1,3,5-Trimethylbenzene

Table 4.10. Alcohol Compounds

Functional Group	Absorption Range (cm <sup>-1</sup> )	Example (cm <sup>-1</sup> )	Example Compound
<b>ALCOHOLS</b>			
<b>General</b>			
OH unbridged group	3650-3590 sp		
OH inter- and intra- molecularly H - bonded	3570-3450		
OH intermolecularly H-bonded	3400-3200 br		
<b>Primary alcohols</b>			
	1350-1260	1339	
	1065-1020	1028	1-Pentanol
<b>Secondary alcohols</b>			
	1370-1260	1369	
	1120-1080	1111	2-Pentanol
<b>Tertiary alcohols</b>			
	1410-1310	1379	
	1170-1120	1124	2-Methylbutanol
<b>Aromatic ring hydroxy compounds</b>			
OH unbridged	3617-3599 sp		
OH dimer	3460-3322 br		
OH polymer	3370-3322 br		
	1410-1310	1350	
	1225-1175	1225	Phenol

Table 4.11. Peroxide Compounds

Functional Group	Absorption Range (cm <sup>-1</sup> )	Example (cm <sup>-1</sup> )	Example Compound
<b>PEROXIDES</b>			
<b>Aliphatic</b>			
	1820-1810		
	1800-1780		
	890-820		
<b>Aromatic</b>			
	1805-1780		
	1785-1755		
	1020-980		

Table 4.12. Ether Type Compounds

Functional Group	Absorption Range (cm <sup>-1</sup> )	Example (cm <sup>-1</sup> )	Example Compound
<b>ETHERS</b>			
<b>Aliphatic</b>			
O-CH <sub>3</sub>	2830-2815		
C-O-C	1150-1060	1140	
O-(CH <sub>2</sub> ) <sub>4</sub>	742-734		Diethyl ether
O-CH <sub>3</sub>	1455		
<b>Aromatic</b>			
=C-O-C	1275-1200	1247	
C-O-C	1075-1020	1038	Anisol
<b>Cyclic</b>			
C-O-C	1140-1070		
<b>Epoxides</b>			
general	1260-1240	1261	1,2-Epoxybutane
<i>trans</i> compounds	890		
<i>cis</i> compounds	830	826	1,2-Epoxybutane
<b>Tetrahydrofuran derivatives</b>			
	1098-1075	1076	Tetrahydrofuran
	915-913	912	
<b>Trioxans</b>			
	1175	1172	Trioxan
	958	957	
<b>Tetrahydropyran derivatives</b>			
	1120-1080		
	1100-900		
	825-805		
<b>Dioxan derivatives</b>			
	1125	1122	Dioxan
<b>KETALS, ACETALS</b>			
R <sub>2</sub> -C-(O-C) <sub>2</sub>	1190-1158		
	1143-1124		
	1098-1063		

Table 4.13. Ketone and Aldehyde Compounds

Functional Group	Absorption Range (cm <sup>-1</sup> )	Example (cm <sup>-1</sup> )	Example Compound
<b>KETONES</b>			
<b>Aliphatic</b>			
	1725-1705	1727	Butanone
	1325-1215	1269	
	1200	1215	
<b>Unsaturated</b>			
C=C	1650-1620	1618	Methyl vinyl ketone
C=O	1685-1665	1684	
<b>Aromatic</b>			
Aryl, alkyl	1700-1680	1694	Acetophenone
Aryl, aryl	1670-1660		
<b>Cyclic</b>			
4- & 5-membered rings	1775-1740	1739	Cyclopentanone
6- & 7-membered rings	1725-1700	1703	Cycloheptanone
<b>Diketones</b>			
α-Diketones	1730-1710	1721	Diacyl ketone
β-Diketones	1640-1540		
γ-Diketones	1725-1705		
<b>Halogen substituted</b>			
α,α-Dihalogen substitution	1765-1745		
α-Dihalogen substitution	1745-1725		
<b>ALDEHYDES</b>			
<b>General</b>			
CH	2900-2700 2 band 2720-2700 975-780		
<b>Aliphatic</b>			
C=O	1740-1720	1735	Butyraldehyde
CH	1440-1325	1390	
<b>Unsaturated</b>			
C=O	1650-1620	1637	Crotonaldehyde
C=O α,β unsaturated	1690-1650		

Table 4.14. Carboxylic Acid Compounds

Functional Group	Absorption Range (cm <sup>-1</sup> )	Example (cm <sup>-1</sup> )	Example Compound
<b>CARBOXYLIC ACIDS</b>			
<b>General</b>			
OH	3200-2500 br		
CH	1440-1396 1320-1210		
OH dimer	950-900 br		
C=O halogen substitution	1740-1720		
C=O aliphatic	1720-1700	1718	Acetic acid
C=O unsaturated	1710-1690	1698	Crotonic acid
C=O aromatic	1700-1680	1695	Benzoic acid

Table 4.14. (Continued)

C=C	1660-1620	1655	Crotonic acid
<b>Carboxylic Ions</b>			
C=O	1610-1560		
C=O	1420-1300		

Table 4.15. Ester Compounds

Functional Group	Absorption Range (cm <sup>-1</sup> )	Example (cm <sup>-1</sup> )	Example Compound
<b>ESTERS</b>			
C=O unsaturated, aryl	1800-1770		
C=C unsaturated, aryl	1730-1710	1718	
C-O acrylates, fumarate	1300-1200	1282	Ethyl acetate
C-O	1190-1130		
C=O electronegatively substituted	1770-1745		
C=O $\alpha, \gamma$ keto	1755-1740		
C=O saturated	1750-1735	1744	Methyl acetate
C=O $\beta$ keto	1660-1640		
C-O benzoates, phthalates	1310-1250	1277	Methyl benzoate
	1150-1100	1108	
C-O acetates	1250-1230	1246	Propyl acetate
	1060-1000	1047	
C-O phenolic acetates	1205		
C-O formate	1200-1180	1190	Propyl formate
<b>LACTONES</b>			
$\beta$ -Lactones	1840-1800		
$\gamma$ -Lactones	1780-1760	1776	Butyrolactone
$\delta$ -Lactones	1750-1730		
	1280-1150	1168	

Table 4.16. Anhydride Compounds

Functional Group	Absorption Range (cm <sup>-1</sup> )	Example (cm <sup>-1</sup> )	Example Compound
<b>ANHYDRIDES</b>			
<b>Aliphatic</b>			
C=O	1850-1800	1842	Acetic acid anhydride
C=O	1785-1760	1783	
C-O	1170-1050	1134	
<b>Aromatic</b>			
C=O	1880-1840	1866	Phthalic acid anhydride
C=O	1790-1770	1773	
C-O	1300-1200	1267	
<b>Cyclic</b>			
C=O	1870-1820	1818	Glutaric acid anhydride
C=O	1800-1750	1772	

Table 4.17. Amide Compounds

Functional Group	Absorption Range (cm <sup>-1</sup> )	Example (cm <sup>-1</sup> )	Example Compound
<b>AMIDES</b>			
<b>Primary</b>			
NH free	3500		
NH free	3400		
NH bridges	3350	3346	Butyramide
NH bridged	3190	3191	
C=O	1660-1640	1660	
	1430-1400	1430	
<b>Secondary</b>			
NH free <i>trans</i>	3460-3400		
NH free <i>cis</i>	3440-3420		
NH bridged <i>trans</i>	3320-3270	3280	N-Methylacetamide
NH bridged <i>cis</i>	3180-3140		
bridged <i>cis, trans</i>	3100-3070	3090	
C=O	1680-1630	1652	
NH	1570-1510	1564	
	720 br	725	
<b>Tertiary</b>			
C=O	1670-1630	1670	N,N-Dimethyl formamid

Table 4.18. Amino Acid Compounds

Functional Group	Absorption Range (cm <sup>-1</sup> )	Example (cm <sup>-1</sup> )	Example Compound
<b>AMINO ACIDS</b>			
NH	3130-3030 br 2760-2530 2140-2080		
C=O	1720-1680		
ionized form	1600-1560 1300		
C=O $\alpha$ -amino acids	1754-1720		
C=O $\beta, \gamma$ -amino acids	1730-1700		
Amino acid HCl	3030-2500		
NH amino acid HCl's	1660-1590		
NH amino acid HCl's	1550-1490		

Table 4.19. Amine Compounds

Functional Group	Absorption Range (cm <sup>-1</sup> )	Example (cm <sup>-1</sup> )	Example Compound
<b>AMINES</b>			
<b>General</b>			
N-CH <sub>3</sub>	2820-2730		
N-CH <sub>3</sub>	1426		
C-N	1410		

Table 4.19. (Continued)

<b>Aliphatic, primary</b>			
NH free	3500-3200 (2 bands)	3350 3210	Ethylamine
NH	1650-1590 1200-1150 1120-1030	1630 1100	
<b>Aliphatic, secondary</b>			
NH free	3500-3200 1 band	3230	Dipropylamine
NH	1650-1550		
C-N	1200-1120	1126	
C-N	1150-1080	1090	
<b>Aliphatic, tertiary</b>			
C-N	1230-1130	1175	Ethyldimethylamine
C-N	1130-1030	1070	
<b>Aromatic, primary</b>			
	3510-3450 3420-3380 1630-1600	3460 3413 1621	Aniline
<b>Aromatic, secondary</b>			
Free	3450-3430		
Bridged	3400-3300	3400	N-Methylaniline

Table 4.20. Unsaturated Nitrogen Compounds

Functional Group	Absorption Range (cm <sup>-1</sup> )	Example (cm <sup>-1</sup> )	Example Compound
<b>UNSATURATED NITROGEN COMPOUNDS</b>			
<b>Imines</b>			
NH	3400-3300		
C=N	1690-1640		
<b>Oximes</b>			
Liquid	3602-3590		
Solid	3250		
Solids	3115		
Aliphatic	1680-1665		
Aromatic	1650-1620 1300 900		

Table 4.21. Cyanide and Isocyanide Compounds

Functional Group	Absorption Range (cm <sup>-1</sup> )	Example (cm <sup>-1</sup> )	Example Compound
<b>CYANIDES, ISOCYANIDES</b>			
C≡N unconjugated	2265-2240	2256	Ethyl cyanide
C≡N conjugated or aromatic	2240-2220	2222	Benzyl cyanide
C≡N cyanide, thiocyanide complex	2200-2000		
N≡C alkyl isocyanide	2183-2150	2166	Methyl isocyanide
N≡C aryl isocyanide	2140-2080	2100	Phenyl isocyanide

*Table 4.22. Cyclic Nitrogen Compounds*

Functional Group	Absorption Range (cm <sup>-1</sup> )	Example (cm <sup>-1</sup> )	Example Compound
<b>CYCLIC NITROGEN COMPOUNDS</b>			
<b>Pyridines, quinolines</b>			
CH	3100-3000	3030	
C=C, C=N	1615-1590	1590	
	1585-1550		
	1520-1465	1490	
	1440-1410		
	920-690	707	
<b>Pyrimidines</b>			
CH	3060-3010		
C=C, C=N	1580-1520		
Ring	1000-900		

*Table 4.23. Unsaturated Nitrogen-Nitrogen Compounds*

Functional Group	Absorption Range (cm <sup>-1</sup> )	Example (cm <sup>-1</sup> )	Example Compound
<b>UNSATURATED NITROGEN-NITROGEN COMPOUNDS</b>			
<b>Azo compounds</b>			
	1630-1575		
N=N azides	2160-2120	2130	
N=N azides	1340-1180	1297	Phenylazide

*Table 4.24. Nitro Compounds*

Functional Group	Absorption Range (cm <sup>-1</sup> )	Example (cm <sup>-1</sup> )	Example Compound
<b>NITRO COMPOUNDS</b>			
<b>Aliphatic</b>			
	1570-1500	1546	
	1385-1365	1362	
	880	879	
<b>Aromatic</b>			
	1550-1510	1527	
	1370-1330	1351	
	849	853	Nitrobenzene

*Table 4.25. Phosphorus Compounds*

Functional Group	Absorption Range (cm <sup>-1</sup> )	Example (cm <sup>-1</sup> )	Example Compound
<b>PHOSPHORUS COMPOUNDS</b>			
<b>O-H phosphoric acids</b>			
P-H	2700-2560 br		
P=O	2440-2350 sp		
P=O	1350-1250		
P-O-C	1250-1150		
P-O-R	1240-1190		
P-O-C	1190		
	1170-1150		

Table 4.25. (Continued)

P-O-C	1050-990		
P-O-P	970-940		
P-F	885		
P=S	840-600		
O-P-H	865-840		
O-P-O	590-520		
O-P-O	460-440		
<b>PHOSPHORUS-CARBON COMPOUNDS</b>			
P-C aromatic	1450-1435		
P-C aliphatic	1320-1280	1298	Trimethylphosphine
P-C	750-650	707	
PO <sub>4</sub> <sup>-3</sup> aryl phosphates	1080-1040		
PO <sub>4</sub> <sup>-3</sup> alkyl phosphates	1180-1150		
PO <sub>4</sub> <sup>-3</sup> alkyl phosphates	1080		

Table 4.26. Deuterated Compounds

Functional Group	Absorption Range (cm <sup>-1</sup> )	Example (cm <sup>-1</sup> )	Example Compound
<b>DEUTERATED COMPOUNDS</b>			
O-D deuterated alcohols	2650-2400		
O-D deuterated carboxylic acids	675		

Table 4.27. Sulfur Compounds

Functional Group	Absorption Range (cm <sup>-1</sup> )	Example (cm <sup>-1</sup> )	Example Compound
<b>SULFUR COMPOUNDS</b>			
C=S	1400-1300	1357	Dithioacetic acid
S=S	1200-1050		
P=S	840-600		
SH mercaptans	2600-2550	2580	Ethyl mercaptan
C-S mercaptans	700-600	665	
C-S-C dialkyl sulfides	750-600	726	Methyl ethyl sulfide
	710-570	676	
	660-630	654	
Aliphatic sulphones	1410-1390	1407	Dimethylsulphone
	1350-1300	1316	
Sulphonic acids	1210-1150		
	1060-1030		
	650		

Table 4.28. Silicon Compounds

Functional Group	Absorption Range (cm <sup>-1</sup> )	Example (cm <sup>-1</sup> )	Example Compound
<b>SILICON COMPOUNDS</b>			
SiH alkylsilanes	2300-2100	2175	Dimethylsilane
Si(CH <sub>3</sub> ) <sub>2</sub>	1265-1258	1262	

Table 4.28. (Continued)

	814-800			
	800			
Si(CH <sub>3</sub> ) <sub>3</sub>	1260-1240	1259		Methoxytrimethylsilane
	850-830	844		
	760	763		
Si-C aromatic	1429			
	1130-1090			
Si-C	860-715			
Si-O siloxanes	1100-1000			
Si-O-C open-chain	1090-1020			
Si-O-Si open-chain	1097			
Si-O-Si cyclic	1080-1010			

Table 4.29. Halogen Compounds

Functional Group	Absorption Range (cm <sup>-1</sup> )	Example (cm <sup>-1</sup> )	Example Compound
<b>HALOGEN COMPOUNDS</b>			
<b>Iodine Compounds</b>			
	500		
<b>Bromine Compounds</b>	700-500		
<b>Chlorine Compounds</b>			
Monochloro	800-600		
	750-700		
Fully chlorinated compounds	780-710		
<b>Fluorine Compounds</b>			
	1400-1000		
	1100-1000		
Fully fluorinated compounds	745-730		

Table 4.30. Inorganic Compounds

Functional Group	Absorption Range (cm <sup>-1</sup> )	Example (cm <sup>-1</sup> )	Example Compound
<b>INORGANIC COMPOUNDS</b>			
<b>Sulfates</b>			
	1200-1140	1143	Potassium sulfate
	1130-1080	1117	
	680-610	617	
<b>Nitrates</b>			
	1380-1350	1370	Potassium nitrate
	840-815	825	
<b>Nitrites</b>			
	840-800		
	750		
<b>Water of Crystallization</b>			
	1630-1615		

Table 4.30. (Continued)

Halogen-Oxygen salts				
Chlorates		980-930	978	Potassium chlorate
		930-910	932	
Bromates		810-790	793	Potassium bromate
Iodates		785-730	756	Potassium iodate
Carbonates		1450-1410	1410	Calcium carbonate
		880-860	875	

## 4.4 Nuclear Magnetic Resonance Spectroscopy

### 4.4.1 Common NMR Solvents

Table 4.31. NMR Solvents

Compound	M.W.	d <sup>20/4</sup>	m.p. <sup>1</sup>	b.p. <sup>1</sup>	δ <sup>2</sup> H(mult) <sup>3</sup>	δ <sup>2</sup> C(mult) <sup>3</sup>
Acetic Acid-d <sub>4</sub>	64.078	1.12	17	118	11.53 (1) 2.03 (5)	178 (br) 20.0 (7)
Acetone-d <sub>6</sub>	64.117	0.87	-94	57	2.04 (5)	206.0 (13) 29.8 (7)
Acetonitrile-d <sub>3</sub>	44.071	0.84	-45	82	1.93 (5)	118.2 (br)
Benzene-d <sub>6</sub>	84.152	0.95	5	80	7.15 (br)	128.0 (3)
Chloroform-d	120.384	1.50	-64	62	7.24 (1)	77.0 (3)
Cyclohexane-d <sub>12</sub>	96.236	0.89	6	81	1.38 (br)	26.4 (5)
Deuterium Oxide	20.028	1.11	3.8	101.4	4.67 (TSP)	
1,2-Dichloroethane-d <sub>4</sub>	102.985	1.25	-40	84	3.72 (br)	43.6 (5)
Diethyl-d <sub>10</sub> Ether	84.185	0.82	-116	35	3.34 (m) 1.07 (m)	65.3 (5) 14.5 (7)
Dimethylformamide-d <sub>7</sub>	80.138	1.04	-61	153	8.01 (br) 2.91 (5) 2.74 (5)	162.7 (3) 35.2 (7) 30.2 (7)
Dimethyl-d <sub>6</sub> Sulphoxide	84.170	1.18	18	189	2.49 (5)	39.5 (7)
p-Dioxane-d <sub>8</sub>	96.156	1.13	122	101	3.53 (m)	66.5 (5)
Ethyl Alcohol-d <sub>6</sub>	52.106	0.91	<130	79	5.19 (1) 3.55 (br) 1.11 (m)	56.8 (5) 17.2 (7)

Table 4.31. (Continued)

Hexafluoroacetone	198.067	1.71	21		5.26 (1)	122.5 (4)
Deuterate						92.9 (7)
HMPT-d <sub>18</sub>	197.314	1.14	7	106	2.53 (2x5)	35.8 (7)
Methyl Alcohol-d <sub>4</sub>	36.067	0.89	-98	65	4.78 (1)	49.0 (7)
Methylene Chloride-d <sub>2</sub>	86.945	1.35	-95	40	5.32 (3)	53.8 (5)
Tetrahydrofuran-d <sub>8</sub>	80.157	0.99	-109	66	3.58 (br) 1.73 (br)	67.4 (5) 25.3 (br)
Trifluoroacetic Acid-d	115.030	1.50	-15	72	11.50 (1)	164.2 (4) 116.6 (4)

1. Melting and boiling points (in °C) are those of the corresponding light compounds. (except for **D<sub>2</sub>O**) and are intended only to indicate the useful liquid range of the materials.
2. Chemicals shifts in ppm relative to TMS.
3. The multiplicity br indicates a broad peak without resolvable fine structure, while m indicates one with fine structure.
4. Note that chemical shifts can be dependent on solute, concentration and temperature.

#### 4.4.2 Reference Standards for Proton NMR

Table 4.32. NMR Reference Standards

Compound (Abbrev.)	Formula	m.p. °C	b.p. °C	Hydrogen bands		
				Group <sup>1</sup>	δ	τ
Tetramethylsilane (TMS) <sup>2</sup>	(CH <sub>3</sub> ) <sub>4</sub> Si		26.5	CH <sub>3</sub> (s)	0.00	10.00
Hexamethyl siloxane (HMDS) <sup>3</sup>	[(CH <sub>3</sub> ) <sub>3</sub> Si] <sub>2</sub> O	-59	100.4	CH <sub>3</sub> (s)	0.04	9.96
Sodium-3-trimethylsilyl- 1-propane sulfonate (DSS) <sup>4</sup>	(CH <sub>3</sub> ) <sub>3</sub> Si- CH <sub>2</sub> -CH <sub>2</sub> - CH <sub>2</sub> -SO <sub>3</sub> -Na	Solid Salt		CH <sub>3</sub> (s) 1-CH <sub>2</sub> (m) 2-CH <sub>2</sub> (m) 3-CH <sub>2</sub> (m)	0.00 0.6 1.8 2.9	10.00 9.4 8.2 7.1
Sodium 3-trimethylsilyl propionate-d <sub>4</sub> (TSP) <sup>5</sup>	(CH <sub>3</sub> ) <sub>3</sub> Si- CD <sub>2</sub> -CD <sub>2</sub> - COO-Na	Solid Salt		CH <sub>3</sub> (s)	0.00	10.00

1. The functional group which produces the observed band. The multiplicity of the band is indicated in parenthesis; s = singlet; m = complex multiplet.
2. Primary reference standard for room temperature and below.
3. Can be used as reference up to 180°C.
4. Reference for water solutions. The CH<sub>2</sub> bands can interfere with weak sample bands.
5. Reference for water solutions.

## 4.4.3 NMR Proton Chemical Shift Chart




Table 4.33. Proton Chemical Shifts




Table 4.33. (Continued)

## 4.4.4 NMR Chemical Shift Tables

The data in all of the tables are for the compound in dilute carbon tetrachloride or deuteriochloroform relative to internal TMS where such data were available. It is important to remember that solvent effects, especially in the case of aromatics, can cause significant variation in the observed chemical shifts. All chemical shifts listed are in ppm or  $\delta$ .

Table 4.34.  $^1\text{H}$  Chemical Shifts of Paraffinic Compounds with a Single Functional Groups

Group X	$\text{CH}_3\text{-X}$	$\text{C-CH}_2\text{-X}$	$(\text{C})_2\text{-CH-X}$	$\text{CH}_3\text{-C-X}$	$\text{C-CH}_2\text{-C-X}$	$(\text{C})_2\text{CH-C-X}$
-H	0.233	0.86	1.33	0.86	1.33	1.56
- $\text{CH=CR}_2$	1.73	2.00	1.73	1.55	1.35	1.00
- $\text{C}\equiv\text{CR}$	1.75	2.15	2.7	1.15	1.50	1.80
- $\text{C}\equiv\text{N}$	1.98	2.35	2.8	1.30	1.6	2.00
-Ph	2.34	2.60	2.87	1.18	1.6	1.8
-CHO	2.17	2.4	2.4	1.13	1.65	---
-COR	2.10	2.4	2.55	1.05	1.5	1.7
-COPh	2.5	2.9	3.4	1.18	1.6	2.0
- $\text{CO}_2\text{R}$	2.1	2.2	2.5	1.15	1.7	1.8
-CONR <sub>2</sub>	2.05	2.23	2.4	1.1	1.6	1.8
-I	2.16	3.17	4.25	1.8	1.8	2.1
-Br	2.68	3.36	4.2	1.8	1.9	2.0
-Cl	3.05	3.44	4.1	1.5	1.8	2.0
-F	4.26	4.4	4.8	1.4	1.8	2.1
-OR	3.38	3.4	3.6	1.2	---	---
-OPh	3.82	3.95	4.6	1.3	1.5	1.7
-OCOR	3.65	4.1	5.0	1.25	1.6	1.8
-OCOPh	3.82	4.2	5.1	1.5	1.7	1.9
-OCOCF <sub>3</sub>	3.95	4.3	---	1.4	1.6	---
-ONO	---	4.75	---	1.4	---	---
-NR <sub>2</sub>	2.3	2.6	2.9	1.05	1.45	1.7
- $\text{N}^+\text{R}_3$	~3.2	~3.1	~3.6	1.4	1.7	2.0
-NRPh	~2.7	~3.1	~3.6	1.1	1.5	1.8
-NHCOR	2.8	3.3	3.8	1.1	1.5	1.9
-NO <sub>2</sub>	4.30	4.4	4.6	1.6	2.05	2.5
-N≡C	2.85	---	4.8	1.6	---	---
-SR	2.09	2.5	3.0	1.25	1.6	1.9
-SSR	2.30	2.7	---	1.3	1.7	---
-SOR	2.5	3.0	2.8	1.35	1.7	---
-SO <sub>2</sub> R	2.8	2.9	3.1	1.35	1.7	---

(From The Chemist's Companion, Gordon & Ford, Copyright © 1972 John Wiley & Sons, Inc.  
Reprinted with permission of John Wiley & Sons, Inc.)

Table 4.35.  $^1\text{H}$  Chemical Shifts of Paraffinic Compounds with Two Functional Groups

Y\X	CN	CF <sub>3</sub>	Ph	C≡C	C=C	CH <sub>3</sub>
CH <sub>3</sub>	2.31	(1.84)	2.63	2.14	1.97	1.34
C=C	3.15	(2.69)	3.30	3.39	2.73	
C≡C	(3.37)	(2.81)	(3.52)	(3.11)		
Ph	3.68	3.50	3.92			
CF <sub>3</sub>	(3.07)	(2.51)				
CN	4.13					

Y\X	COPh	CONR <sub>2</sub>	COOR	COR
COR	(3.77)	3.52	3.32	3.62
COOR	(3.62)	(4.95)	4.22	
CONR <sub>2</sub>	(3.66)	3.30		
COPh	(3.91)			

Y\X	OCOR	OPh	OR	OH
OH	(5.92)	(6.02)	(5.15)	(5.35)
OR	(5.72)	(5.82)	4.49	
OPh	(6.59)	(6.69)		
OCOR	(6.49)			

Y\X	I	Br	Cl
Cl	4.99	5.16	5.28
Br	(4.38)	4.94	
I	3.89		

Y\X	NHCOR	N <sub>3</sub>	NR <sub>2</sub>
NR <sub>2</sub>	(4.07)	(3.77)	3.10
N <sub>3</sub>	(4.47)	(4.17)	
NHCOR	(4.77)		

Values in parentheses were calculated by the empirical method of Shoolery. All values refer to methylene ( $\text{CH}_2$ ) protons in  $\text{X}-\text{CH}_2-\text{Y}$ .

Table 4.36.  $^1\text{H}$  Chemical Shifts of Olefinic Compounds


Compound	X=	a	b	c
$\text{H}_2\text{C}=\text{CH}_2$		5.33		
	R	5.70	4.88	4.96
	F	6.17	4.03	4.37
	Cl	6.30	5.44	5.52
	Br	6.49	5.88	6.03
	C <sub>6</sub> H <sub>5</sub>	6.69	5.21	5.71
	CCl <sub>3</sub>	6.41	5.30	5.78
	CN	5.53	6.05	5.78
	CH <sub>2</sub> OH	6.00	5.13	5.25
	OCH <sub>3</sub>	6.43	3.90	4.04

Table 4.36. (Continued)







$\text{H}_2\text{C}=\text{C}=\text{CH}_2$		4.55		
	Cl Br I	5.76 5.85 5.62	5.05 4.82 4.46	
		5.20	5.11	
		5.38		
		4.70		
		4.82		
		4.55		
$\text{CH}_3\text{-CHO}$		9.72		
$\text{Ph-CHO}$		9.96		
$\text{CH}_2=\text{CH-CHO}$		9.48		
$(\text{CH}_3)_2\text{N-CHO}$		7.84		
$\text{CH}_3\text{O-CHO}$		8.08		

Table 4.37.  $^1\text{H}$  Chemical Shifts of Acetylenic Compounds

Compound	$\delta$	Compound	$\delta$
$\text{H-C}\equiv\text{C-H}$	1.80	$\text{OHC-C}\equiv\text{C-H}$	1.89
$\text{CH}_3\text{-C}\equiv\text{C-H}$	1.80	$\text{HOCH}_2\text{-C}\equiv\text{C-H}$	2.33
$\text{CH}_3\text{CH}_2\text{-C}\equiv\text{C-H}$	1.76	$\text{Cl-CH}_2\text{-C}\equiv\text{C-H}$	2.40
$\text{Ph-C}\equiv\text{C-H}$	3.05	$\text{Br-CH}_2\text{-C}\equiv\text{C-H}$	2.33
$\text{CH}_2=\text{CH-C}\equiv\text{C-H}$	2.92	$\text{I-CH}_2\text{-C}\equiv\text{C-H}$	2.19
$\text{C}_2\text{H}_5\text{-C}\equiv\text{C-C}\equiv\text{C-H}$	1.95	$\text{CH}_3\text{O-C}\equiv\text{C-H}$	1.33
$\text{CH}_3(\text{C}\equiv\text{C})_2\text{C}\equiv\text{C-H}$	1.87	$\text{CH}_2=\text{CH-O-C}\equiv\text{C-H}$	1.89

(From The Chemist's Companion, Gordon & Ford, Copyright © 1972 John Wiley & Sons, Inc.  
Reprinted with permission of John Wiley & Sons, Inc.)

Table 4.38.  $^1\text{H}$  Chemical Shifts of Cycloparaffinic Compounds














Compound	$\delta$	Compound	$\delta$	Compound	$\delta$
Cyclopropane	0.22	Cyclobutane	1.96	Cyclopentane	1.51
Cyclohexane	1.44	Cycloheptane	1.54	Cyclooctane	1.54
Adamantane	1.78				
	0.02		~1.4		2.37
	a 1.51 b ~2.1		a 1.56 b 0.87 c 2.49 d 1.58		a 1.21 b 1.49 c 1.18 d 2.20
	1.65		a 3.03 b 1.96		a 2.06 b 2.02
	2.25		0.99		a 2.70 b 1.92
	a 2.70 b 1.92		1.5		

Table 4.39.  $^1\text{H}$  Chemical Shifts Of Cycloolefinic Compounds




Compound	$\delta$	Compound	$\delta$	Compound	$\delta$
	a 0.92 b 7.01		a 2.57 b 5.97		a 2.28 b 5.60

Table 4.39. (Continued)










	6.42		a 1.96 b 5.57		2.15
	a 1.95 b 3.53		5.95		6.66
	6.25		6.27		6.70

Table 4.40.  $^1\text{H}$  Chemical Shifts Of Monosubstituted Benzenes

Substituent	Ortho	Meta	Para
H	7.27	7.27	7.27
$\text{CH}_3$	7.07	7.07	7.07
$\text{CH}_2\text{CH}_3$	7.13	7.13	7.13
$\text{CH}_2\text{OH}$	7.28	7.28	7.28
$\text{CH}_2\text{Cl}$	7.32	7.32	7.32
$\text{CHCl}_2$	7.42	7.42	7.42
$\text{CCl}_3$	7.91	7.40	7.37
$\text{CH}=\text{CH}_2$	7.5	7.5	7.5
CHO	7.83	7.49	7.56
$\text{COCH}_3$	7.89	7.41	7.56
$\text{CO}_2\text{H}$	8.12	7.43	7.51
$\text{CO}_2\text{CH}_3$	7.98	7.38	7.48
COCl	8.11	7.49	7.63
COBr	8.07	7.48	7.64
CONH <sub>2</sub>	7.8	7.5	7.5
CN	7.63	7.45	7.55
F	6.99	7.24	7.08
Cl	7.30	7.25	7.18
Br	7.45	7.19	7.23
I	7.67	7.06	7.27
$\text{NH}_2$	6.52	7.02	6.62
$\text{NHCH}_3$	6.47	7.05	6.59
$\text{N}(\text{CH}_3)_2$	6.61	7.09	6.60
NHCOCH <sub>3</sub>	7.7	7.1	7.0

Table 4.40. (Continued)

$\text{NH}_3^+$	7.7	7.5	7.5
NO	7.81	7.55	7.61
$\text{NO}_2$	8.22	7.53	7.65
OH	6.68	7.15	6.82
$\text{OCH}_3$	7.77	7.15	7.37
$\text{OCOCH}_3$	6.79	7.18	6.83
$\text{SCH}_3$	7.4	7.2	7.1
$\text{SO}_2\text{Cl}$	8.04	7.62	7.72
$\text{SO}_3\text{CH}_3$	7.87	7.53	7.60

(From The Chemist's Companion, Gordon & Ford, Copyright © 1972 John Wiley & Sons, Inc. Reprinted with permission of John Wiley & Sons, Inc.)

Table 4.41.  $^1\text{H}$  Chemical Shifts Of Heteroaromatic Compounds











Compound	$\delta$	Compound	$\delta$
	a 7.38 b 6.30		a 7.19 b 7.04
	8.19		8.58
	a 6.62 b 6.05		a 8.88 b 7.41 c 7.98
	a 7.55 b 6.25 c 7.55		a 7.14 b 7.14 c 7.70
	a 8.50 b 7.06 c 7.46		a 6.57 b 7.26 c 6.15 d 7.13

Table 4.41. (Continued)





















	a 9.17 b 7.68		a 9.15 b 8.60 c 7.09
	8.5		9.18
	a 6.54 b 6.34		8.95
	a 8.03 b 7.77 c 7.12 d 7.34 e 7.58		a 8.63 b 9.16 c 8.95
	a 7.14 b 6.64 c 6.28 d 7.25 e 6.50 f 6.35 g 7.76 a 7.52 b 6.66 c 7.49 d 7.13 e 7.19 f 7.42		a 8.81 b 7.26 c 8.00 d 7.68 e 7.43 f 7.61 g 8.05 a 8.20 b 7.69 c 7.39 d 7.80 e 8.53

(From The Chemist's Companion, Gordon & Ford, Copyright © 1972 John Wiley & Sons, Inc.  
Reprinted with permission of John Wiley & Sons, Inc.)

Table 4.42.  $^1\text{H}$  Chemical Shifts Of Heterocyclic Compounds

Compound	$\delta$	Compound	$\delta$	Compound	$\delta$
	2.54		1.48		2.27

Table 4.42. (Continued)

	a 4.73 b 2.72		a 3.54 b 2.23		a 2.82 b 1.93
	a 3.63 b 1.79		a 2.74 b 1.62		a 2.82 b 1.93
	a 3.56 b 1.58		a 2.69 b 1.49		a 2.57 b 1.60
	a 3.48 b 4.22		a 2.31 b 2.08 c 4.28		a 2.27 b 1.62 c 4.06
	a 4.77 b 3.77		a 4.82 b 3.80 c 1.68		3.59
	5.00		3.69		4.18
	a 2.92 b 2.16		a 3.57 b 2.83		a 3.88 b 2.57

(From The Chemist's Companion, Gordon & Ford, Copyright © 1972 John Wiley & Sons, Inc.  
Reprinted with permission of John Wiley & Sons, Inc.)

Table 4.43.  $^1\text{H}$  Chemical Shifts of Hydrogen Bonded To Oxygen, Nitrogen, Sulfur

Compound	$\delta$	Compound	$\delta$
ROH (monomeric) <sup>a</sup>	0.5	Oximes	7-11
ROH (hydrogen bonded)	0.5-5.0	RNH <sub>2</sub>	1.1-1.8
ArOH (monomeric) <sup>a</sup>	4.5	R <sub>2</sub> NH	1.2-2.1
ArOH (hydrogen bonded)	4.5-9	ArNH <sub>2</sub>	3.3-4.0
Enols (intramolecular bonded)	15-19	ArNHR	3.1-3.8
Carboxylic acids	10-13	RCONH <sub>2</sub> , ArCONH <sub>2</sub>	5-6.5
Sulfonic acids	11-12	RCONHR, ArCONHR	6-8.2
RSH	1-2	RCONHAr, ArCONHAr	7.8-9.4
ArSH	3-4	R <sub>3</sub> N <sup>+</sup> H	7.1-7.7

(From The Chemist's Companion, Gordon & Ford, Copyright © 1972 John Wiley & Sons, Inc.  
Reprinted with permission of John Wiley & Sons, Inc.)

## 4.5 Mass Spectroscopy

### 4.5.1 Common Molecular Ions Table

Table 4.44. Common Elemental Compositions of Molecular Ions<sup>1</sup>

m/z	Composition
16	CH <sub>4</sub>
17	NH <sub>3</sub>
18	H <sub>2</sub> O
26	C <sub>2</sub> H <sub>2</sub>
27	CHN
28	C <sub>2</sub> H <sub>4</sub> , CO, N <sub>2</sub>
30	C <sub>2</sub> H <sub>6</sub> , CH <sub>2</sub> O, NO
31	CH <sub>5</sub> N
32	CH <sub>4</sub> O, N <sub>2</sub> H <sub>4</sub> , SiH <sub>4</sub> , O <sub>2</sub>
34	CH <sub>3</sub> F, PH <sub>3</sub> , H <sub>2</sub> S
36	HCl
40	C <sub>3</sub> H <sub>4</sub>
41	C <sub>2</sub> H <sub>3</sub> N
42	C <sub>3</sub> H <sub>6</sub> , C <sub>2</sub> H <sub>2</sub> O, CH <sub>2</sub> N <sub>2</sub>
43	C <sub>2</sub> H <sub>5</sub> N, N <sub>3</sub> H
44	C <sub>2</sub> H <sub>4</sub> O, C <sub>3</sub> H <sub>8</sub> , C <sub>2</sub> HF, CO <sub>2</sub> , N <sub>2</sub> O
45	C <sub>2</sub> H <sub>7</sub> N, CH <sub>3</sub> NO
46	C <sub>2</sub> H <sub>6</sub> O, C <sub>2</sub> H <sub>3</sub> F, CH <sub>6</sub> Si, CH <sub>2</sub> O <sub>2</sub> , NO <sub>2</sub>
48	C <sub>2</sub> H <sub>5</sub> F, CH <sub>4</sub> S, CH <sub>5</sub> P
50	C <sub>4</sub> H <sub>2</sub> , CH <sub>3</sub> Cl
52	C <sub>4</sub> H <sub>4</sub> , CH <sub>2</sub> F <sub>2</sub>
53	C <sub>3</sub> H <sub>3</sub> N, HF <sub>2</sub> N
54	C <sub>4</sub> H <sub>6</sub> , F <sub>2</sub> O
55	C <sub>3</sub> H <sub>5</sub> N
56	C <sub>4</sub> H <sub>8</sub> , C <sub>3</sub> H <sub>4</sub> O, C <sub>2</sub> H <sub>4</sub> N <sub>2</sub>
57	C <sub>3</sub> H <sub>7</sub> N, C <sub>2</sub> H <sub>3</sub> NO

Table 4.44. (Continued)

58	$\text{C}_3\text{H}_6\text{O}$ , $\text{C}_4\text{H}_{10}$ , $\text{C}_2\text{H}_2\text{O}_2$ , $\text{C}_2\text{H}_6\text{N}_2$
59	$\text{C}_3\text{H}_9\text{N}$ , $\text{C}_2\text{H}_5\text{NO}$ , $\text{CH}_5\text{N}_3$
60	$\text{C}_3\text{H}_8\text{O}$ , $\text{C}_3\text{H}_5\text{F}$ , $\text{C}_2\text{H}_8\text{N}_2$ , $\text{C}_2\text{H}_4\text{O}_2$ , $\text{C}_2\text{H}_8\text{Si}$ , $\text{C}_2\text{H}_4\text{S}$ , $\text{C}_2\text{HCl}$ , $\text{CH}_4\text{N}_2\text{O}$ , $\text{COS}$
61	$\text{C}_2\text{H}_7\text{NO}$ , $\text{CH}_3\text{NO}_2$ , $\text{CClN}$
62	$\text{C}_3\text{H}_7\text{F}$ , $\text{C}_2\text{H}_7\text{P}$ , $\text{C}_2\text{H}_6\text{O}_2$ , $\text{C}_2\text{H}_6\text{S}$ , $\text{C}_2\text{H}_3\text{Cl}$
64	$\text{C}_2\text{H}_2\text{F}_2$ , $\text{C}_2\text{H}_5\text{FO}$ , $\text{C}_2\text{H}_5\text{CL}$ , $\text{SO}_2$
66	$\text{C}_5\text{H}_6$ , $\text{C}_2\text{H}_4\text{F}_2$ , $\text{CF}_2\text{O}$ , $\text{F}_2\text{N}_2$
67	$\text{C}_4\text{H}_5\text{N}$ , $\text{CH}_3\text{F}_2\text{N}$ , $\text{ClO}_2$
68	$\text{C}_5\text{H}_8$ , $\text{C}_4\text{H}_4\text{O}$ , $\text{C}_3\text{H}_4\text{N}_2$ , $\text{C}_3\text{O}_2$ , $\text{CH}_2\text{ClF}$
69	$\text{C}_4\text{H}_7\text{N}$ , $\text{C}_3\text{H}_3\text{NO}$ , $\text{C}_2\text{H}_3\text{N}_3$
70	$\text{C}_5\text{H}_{10}$ , $\text{C}_4\text{H}_6\text{O}$ , $\text{C}_3\text{H}_6\text{N}_2$ , $\text{CH}_2\text{N}_4$ , $\text{CHF}_3$
71	$\text{C}_4\text{H}_9\text{N}$ , $\text{C}_3\text{H}_5\text{NO}$ , $\text{F}_3\text{N}$
72	$\text{C}_4\text{H}_8\text{O}$ , $\text{C}_5\text{H}_{12}$ , $\text{C}_3\text{H}_4\text{O}_2$ , $\text{C}_4\text{H}_5\text{F}$
73	$\text{C}_4\text{H}_{11}\text{N}$ , $\text{C}_3\text{H}_7\text{NO}$ , $\text{C}_2\text{H}_3\text{NS}$ , $\text{C}_2\text{H}_7\text{N}_3$
74	$\text{C}_4\text{H}_{10}\text{O}$ , $\text{C}_3\text{H}_6\text{O}_2$ , $\text{C}_3\text{H}_{10}\text{N}_2$ , $\text{C}_3\text{H}_6\text{S}$ , $\text{C}_2\text{H}_6\text{N}_2\text{O}$ , $\text{C}_3\text{H}_{10}\text{Si}$ , $\text{C}_3\text{H}_3\text{Cl}$ , $\text{CH}_6\text{N}_4$ , $\text{C}_2\text{H}_6\text{N}_2\text{O}$ , $\text{C}_2\text{H}_2\text{O}_3$ , $\text{C}_3\text{H}_9\text{NO}$ , $\text{C}_2\text{H}_5\text{NO}_2$ , $\text{C}_2\text{H}_2\text{ClN}$
76	$\text{C}_3\text{H}_8\text{O}_2$ , $\text{C}_3\text{H}_8\text{S}$ , $\text{C}_3\text{H}_5\text{Cl}$ , $\text{C}_4\text{H}_9\text{F}$ , $\text{C}_4\text{N}_2$ , $\text{C}_3\text{H}_9\text{P}$ , $\text{C}_3\text{H}_5\text{FO}$ , $\text{C}_2\text{H}_4\text{O}_3$ , $\text{C}_2\text{H}_4\text{OS}$ , $\text{CH}_8\text{Si}_2$ , $\text{CH}_4\text{N}_2\text{S}$ , $\text{CS}_2$
77	$\text{CH}_3\text{NO}_3$
78	$\text{C}_6\text{H}_6$ , $\text{C}_3\text{H}_7\text{Cl}$ , $\text{C}_4\text{HSN}_2$ , $\text{C}_2\text{H}_6\text{OS}$ , $\text{C}_2\text{H}_3\text{ClO}$ , $\text{C}_2\text{H}_3\text{FO}_2$ , $\text{CF}_2\text{N}_2$ , $\text{CH}_6\text{N}_2\text{O}_2$
79	$\text{C}_5\text{H}_5\text{N}$
80	$\text{C}_6\text{H}_8$ , $\text{C}_4\text{H}_4\text{N}_2$ , $\text{C}_3\text{H}_6\text{F}_2$ , $\text{C}_2\text{H}_5\text{ClO}$ , $\text{C}_2\text{H}_2\text{ClF}$ , $\text{CH}_4\text{O}_2\text{S}$ , $\text{HBr}$
81	$\text{C}_5\text{H}_7\text{N}$ , $\text{C}_3\text{H}_3\text{N}_3$ , $\text{C}_2\text{H}_5\text{F}_2\text{N}$
82	$\text{C}_6\text{H}_{10}$ , $\text{C}_4\text{H}_6\text{N}_2$ , $\text{C}_5\text{H}_6\text{O}$ , $\text{C}_2\text{H}_4\text{ClF}$ , $\text{C}_2\text{H}_2\text{N}_4$ , $\text{C}_2\text{HF}_3$ , $\text{CClFO}$
83	$\text{C}_5\text{H}_9\text{N}$ , $\text{C}_3\text{H}_5\text{N}_3$ , $\text{C}_4\text{H}_5\text{NO}$
84	$\text{C}_6\text{H}_{12}$ , $\text{C}_5\text{H}_8\text{O}$ , $\text{C}_4\text{H}_8\text{N}_2$ , $\text{C}_4\text{H}_4\text{O}_2$ , $\text{C}_2\text{H}_4\text{N}_4$ , $\text{C}_4\text{H}_4\text{S}$ , $\text{C}_2\text{H}_3\text{F}_3$ , $\text{CH}_2\text{Cl}_2$
85	$\text{C}_5\text{H}_{11}\text{N}$ , $\text{C}_4\text{H}_7\text{NO}$ , $\text{C}_3\text{H}_3\text{NS}$ , $\text{CH}_3\text{N}_5$
86	$\text{C}_5\text{H}_{10}\text{O}$ , $\text{C}_4\text{H}_6\text{O}_2$ , $\text{C}_6\text{H}_{14}$ , $\text{C}_4\text{H}_{10}\text{N}_2$ , $\text{C}_4\text{H}_6\text{S}$ , $\text{C}_4\text{H}_3\text{Cl}$ , $\text{C}_3\text{H}_6\text{N}_2\text{O}$ , $\text{C}_2\text{H}_2\text{N}_2\text{S}$ , $\text{CHClF}_2$ , $\text{HF}_2\text{O}$ , $\text{Cl}_2\text{O}$ , $\text{F}_2\text{OS}$
87	$\text{C}_5\text{H}_{13}\text{N}$ , $\text{C}_4\text{H}_9\text{NO}$ , $\text{C}_3\text{H}_9\text{N}_3$ , $\text{C}_3\text{H}_5\text{NS}$ , $\text{C}_3\text{H}_2\text{ClN}$ , $\text{ClF}_2\text{N}$ , $\text{F}_3\text{NO}$
88	$\text{C}_5\text{H}_{12}\text{O}$ , $\text{C}_4\text{H}_8\text{O}_2$ , $\text{C}_4\text{H}_{12}\text{N}_2$ , $\text{C}_4\text{H}_8\text{S}$ , $\text{C}_3\text{H}_8\text{N}_2\text{O}$ , $\text{C}_3\text{H}_4\text{O}_3$ , $\text{C}_4\text{H}_{12}\text{Si}$ , $\text{C}_4\text{H}_5\text{Cl}$ , $\text{CF}_4$
89	$\text{C}_4\text{H}_{11}\text{NO}$ , $\text{C}_3\text{H}_7\text{NO}_2$ , $\text{C}_3\text{H}_4\text{ClN}$
90	$\text{C}_4\text{H}_{10}\text{O}_2$ , $\text{C}_4\text{H}_{10}\text{S}$ , $\text{C}_4\text{H}_7\text{Cl}$ , $\text{C}_3\text{H}_6\text{O}_3$ , $\text{C}_3\text{H}_{10}\text{OSi}$ , $\text{C}_3\text{H}_6\text{OS}$ , $\text{C}_3\text{H}_3\text{ClO}$ , $\text{C}_2\text{H}_6\text{N}_2\text{O}_2$ , $\text{C}_2\text{H}_6\text{N}_2\text{S}$ , $\text{C}_2\text{H}_2\text{O}_4$ , $\text{C}_4\text{H}_{11}\text{P}$
91	$\text{C}_2\text{H}_5\text{NO}_3$ , $\text{CH}_5\text{N}_3\text{S}$ , $\text{C}_3\text{H}_6\text{ClN}$
92	$\text{C}_7\text{H}_8$ , $\text{C}_4\text{H}_9\text{Cl}$ , $\text{C}_3\text{H}_5\text{ClO}$ , $\text{C}_2\text{H}_4\text{O}_2\text{S}$ , $\text{C}_3\text{H}_8\text{O}_3$ , $\text{C}_3\text{H}_9\text{FSi}$ , $\text{C}_5\text{H}_4\text{N}_2$ , $\text{C}_6\text{H}_4\text{O}$
93	$\text{C}_6\text{H}_7\text{N}$ , $\text{C}_5\text{H}_3\text{NO}$ , $\text{C}_4\text{H}_3\text{N}_3$
94	$\text{C}_5\text{H}_6\text{N}_2$ , $\text{C}_7\text{H}_{10}$ , $\text{C}_6\text{H}_6\text{O}$ , $\text{C}_3\text{H}_7\text{ClO}$ , $\text{C}_2\text{H}_2\text{S}_2$ , $\text{C}_2\text{H}_6\text{O}_2\text{SC}_3\text{HF}_3$ , $\text{C}_2\text{H}_3\text{ClO}_2$ , $\text{C}_2\text{Cl}_2$ , $\text{CH}_3\text{Br}$
95	$\text{C}_5\text{H}_5\text{NO}$ , $\text{C}_6\text{H}_9\text{N}$ , $\text{C}_4\text{H}_5\text{N}_3$ , $\text{C}_2\text{F}_3\text{N}$
96	$\text{C}_7\text{H}_{12}$ , $\text{C}_6\text{H}_8\text{O}$ , $\text{C}_5\text{H}_8\text{N}_2$ , $\text{C}_5\text{H}_4\text{O}_2$ , $\text{C}_4\text{H}_4\text{N}_2\text{O}$ , $\text{C}_2\text{H}_2\text{Cl}_2$ , $\text{C}_3\text{H}_3\text{F}_3$ , $\text{C}_2\text{H}_6\text{F}_2\text{Si}$
97	$\text{C}_5\text{H}_7\text{NO}$ , $\text{C}_6\text{H}_{11}\text{N}$
98	$\text{C}_7\text{H}_{14}$ , $\text{C}_6\text{H}_{10}\text{O}$ , $\text{C}_5\text{H}_6\text{O}_2$ , $\text{C}_4\text{H}_6\text{O}_2$ , $\text{C}_4\text{H}_6\text{N}_2\text{O}$ , $\text{C}_3\text{H}_6\text{N}_4$ , $\text{C}_5\text{H}_{16}\text{N}_2$ , $\text{C}_5\text{H}_6\text{S}$ , $\text{C}_4\text{H}_2\text{O}_3$ , $\text{C}_2\text{H}_4\text{Cl}_2$ , $\text{C}_2\text{HClF}_2$ , $\text{CCl}_2\text{O}$
99	$\text{C}_6\text{H}_{13}\text{N}$ , $\text{C}_5\text{H}_9\text{NO}$ , $\text{C}_4\text{H}_5\text{NS}$ , $\text{C}_4\text{H}_5\text{NO}_2$ , $\text{CH}_3\text{F}_2\text{NS}$

Table 4.44. (Continued)

100	C <sub>6</sub> H <sub>12</sub> O, C <sub>5</sub> H <sub>8</sub> O <sub>2</sub> , C <sub>7</sub> H <sub>16</sub> , C <sub>4</sub> H <sub>4</sub> O <sub>3</sub> , C <sub>3</sub> H <sub>4</sub> N <sub>2</sub> S, C <sub>5</sub> H <sub>12</sub> N <sub>2</sub> , C <sub>6</sub> H <sub>9</sub> F, C <sub>5</sub> H <sub>12</sub> Si, C <sub>4</sub> H <sub>4</sub> OS, C <sub>3</sub> H <sub>4</sub> N <sub>2</sub> O <sub>2</sub> , C <sub>2</sub> H <sub>3</sub> F <sub>3</sub> O, C <sub>2</sub> F <sub>4</sub>
101	C <sub>6</sub> H <sub>15</sub> N, C <sub>5</sub> H <sub>11</sub> NO, C <sub>4</sub> H <sub>7</sub> NO <sub>2</sub> , C <sub>4</sub> H <sub>7</sub> NS, C <sub>2</sub> H <sub>3</sub> N <sub>3</sub> S
102	C <sub>6</sub> H <sub>14</sub> O, C <sub>5</sub> H <sub>10</sub> O <sub>2</sub> , C <sub>5</sub> H <sub>10</sub> S, C <sub>4</sub> H <sub>10</sub> N <sub>2</sub> O, C <sub>4</sub> H <sub>6</sub> O <sub>3</sub> , C <sub>2</sub> H <sub>2</sub> F <sub>4</sub> , C <sub>3</sub> H <sub>6</sub> N <sub>2</sub> S, C <sub>8</sub> H <sub>6</sub> , CHCl <sub>2</sub> F, CHF <sub>3</sub> S, HF <sub>2</sub> PS, Cl <sub>2</sub> S
103	C <sub>4</sub> H <sub>9</sub> NO <sub>2</sub> , C <sub>7</sub> H <sub>5</sub> N, C <sub>5</sub> H <sub>13</sub> NO, C <sub>4</sub> H <sub>13</sub> N <sub>3</sub> , C <sub>4</sub> H <sub>6</sub> CIN, C <sub>2</sub> H <sub>2</sub> CIN <sub>3</sub>
104	C <sub>5</sub> H <sub>12</sub> O <sub>2</sub> , C <sub>5</sub> H <sub>12</sub> S, C <sub>5</sub> H <sub>9</sub> Cl, C <sub>4</sub> H <sub>8</sub> O <sub>3</sub> , C <sub>4</sub> H <sub>8</sub> O <sub>3</sub> , C <sub>4</sub> H <sub>8</sub> OS, C <sub>8</sub> H <sub>8</sub> , C <sub>6</sub> H <sub>13</sub> F, C <sub>6</sub> H <sub>4</sub> N <sub>2</sub> , C <sub>4</sub> H <sub>12</sub> N <sub>2</sub> O, C <sub>4</sub> H <sub>12</sub> OSi, C <sub>4</sub> H <sub>5</sub> ClO, C <sub>3</sub> H <sub>8</sub> N <sub>2</sub> S, CClF <sub>3</sub> , SiF <sub>4</sub>
105	C <sub>7</sub> H <sub>7</sub> N, C <sub>3</sub> H <sub>7</sub> NO <sub>3</sub> , C <sub>4</sub> H <sub>11</sub> NO <sub>2</sub> , C <sub>4</sub> H <sub>8</sub> CIN, CBrN
106	C <sub>8</sub> H <sub>10</sub> , C <sub>5</sub> H <sub>11</sub> Cl, C <sub>6</sub> H <sub>6</sub> N <sub>2</sub> , C <sub>7</sub> H <sub>6</sub> O, C <sub>4</sub> H <sub>10</sub> O <sub>3</sub> , C <sub>4</sub> H <sub>7</sub> ClO, C <sub>4</sub> H <sub>10</sub> OS, C <sub>3</sub> H <sub>6</sub> O <sub>2</sub> S, C <sub>2</sub> H <sub>3</sub> Br
107	C <sub>7</sub> H <sub>9</sub> N, C <sub>6</sub> H <sub>5</sub> NO, C <sub>2</sub> H <sub>5</sub> NO <sub>4</sub>
108	C <sub>6</sub> H <sub>8</sub> N <sub>2</sub> , C <sub>8</sub> H <sub>12</sub> , C <sub>7</sub> H <sub>8</sub> O, C <sub>4</sub> H <sub>9</sub> ClO, C <sub>3</sub> H <sub>9</sub> ClSi, C <sub>3</sub> H <sub>3</sub> ClO <sub>2</sub> , C <sub>6</sub> H <sub>4</sub> O <sub>2</sub> , C <sub>3</sub> H <sub>8</sub> S <sub>2</sub> , C <sub>2</sub> H <sub>5</sub> Br, C <sub>2</sub> H <sub>4</sub> O <sub>3</sub> S, SF <sub>4</sub>
109	C <sub>6</sub> H <sub>7</sub> NO, C <sub>7</sub> H <sub>11</sub> N, C <sub>2</sub> H <sub>4</sub> CINO <sub>2</sub>
110	C <sub>8</sub> H <sub>14</sub> , C <sub>7</sub> H <sub>10</sub> O, C <sub>5</sub> H <sub>6</sub> N <sub>2</sub> O, C <sub>3</sub> H <sub>4</sub> Cl <sub>2</sub> , C <sub>7</sub> H <sub>7</sub> F, C <sub>6</sub> H <sub>6</sub> O <sub>2</sub> , C <sub>4</sub> H <sub>6</sub> N <sub>4</sub> , C <sub>2</sub> H <sub>6</sub> O <sub>3</sub> S, C <sub>2</sub> H <sub>7</sub> O <sub>3</sub> P, C <sub>3</sub> H <sub>7</sub> ClO <sub>2</sub> , C <sub>6</sub> H <sub>10</sub> N <sub>2</sub>
111	C <sub>7</sub> H <sub>13</sub> N, C <sub>6</sub> H <sub>9</sub> NO, C <sub>5</sub> H <sub>5</sub> NO <sub>2</sub> , C <sub>4</sub> H <sub>5</sub> N <sub>3</sub> O, C <sub>2</sub> ClF <sub>2</sub> N, C <sub>6</sub> H <sub>6</sub> FN, C <sub>5</sub> H <sub>9</sub> N <sub>3</sub> , C <sub>5</sub> H <sub>5</sub> NS
112	C <sub>8</sub> H <sub>16</sub> , C <sub>7</sub> H <sub>12</sub> O, C <sub>6</sub> H <sub>8</sub> O <sub>2</sub> , C <sub>6</sub> H <sub>12</sub> N <sub>2</sub> , C <sub>6</sub> H <sub>12</sub> N <sub>2</sub> , C <sub>6</sub> H <sub>8</sub> S, C <sub>5</sub> H <sub>8</sub> N <sub>2</sub> O, C <sub>5</sub> H <sub>4</sub> O <sub>3</sub> , C <sub>4</sub> H <sub>4</sub> N <sub>2</sub> O <sub>2</sub> , C <sub>3</sub> H <sub>6</sub> Cl <sub>2</sub> , C <sub>6</sub> H <sub>5</sub> FO, C <sub>6</sub> H <sub>5</sub> Cl, C <sub>5</sub> H <sub>4</sub> OS, C <sub>3</sub> F <sub>4</sub> , C <sub>3</sub> H <sub>3</sub> F <sub>3</sub> O, CH <sub>2</sub> BrF
113	C <sub>7</sub> H <sub>15</sub> N, C <sub>6</sub> H <sub>11</sub> NO, C <sub>5</sub> H <sub>7</sub> NO <sub>2</sub> , C <sub>5</sub> H <sub>7</sub> NS, C <sub>5</sub> H <sub>4</sub> CIN, C <sub>3</sub> H <sub>3</sub> NO <sub>2</sub> , C <sub>2</sub> H <sub>6</sub> F <sub>2</sub> NP
114	C <sub>7</sub> H <sub>14</sub> O, C <sub>6</sub> H <sub>10</sub> O <sub>2</sub> , C <sub>8</sub> H <sub>18</sub> , C <sub>6</sub> H <sub>14</sub> N <sub>2</sub> , C <sub>4</sub> H <sub>6</sub> N <sub>2</sub> O <sub>2</sub> , C <sub>5</sub> H <sub>6</sub> N <sub>2</sub> O <sub>2</sub> , C <sub>5</sub> H <sub>6</sub> OS, C <sub>6</sub> H <sub>4</sub> F <sub>2</sub> , C <sub>5</sub> H <sub>2</sub> O <sub>3</sub> , C <sub>6</sub> H <sub>7</sub> Cl, C <sub>4</sub> H <sub>6</sub> N <sub>2</sub> S, C <sub>2</sub> H <sub>4</sub> Cl <sub>2</sub> O, C <sub>2</sub> HCl <sub>2</sub> F, C <sub>2</sub> HClF <sub>2</sub> O, C <sub>2</sub> HF <sub>3</sub> O <sub>2</sub>
115	C <sub>6</sub> H <sub>13</sub> NO, C <sub>7</sub> H <sub>17</sub> N, C <sub>5</sub> H <sub>9</sub> NO <sub>2</sub> , C <sub>5</sub> H <sub>9</sub> NS, C <sub>4</sub> H <sub>5</sub> NOS, C <sub>3</sub> H <sub>5</sub> N <sub>3</sub> S, C <sub>5</sub> H <sub>13</sub> N <sub>3</sub>
116	C <sub>6</sub> H <sub>12</sub> O <sub>2</sub> , C <sub>7</sub> H <sub>16</sub> O, C <sub>6</sub> H <sub>12</sub> S, C <sub>9</sub> H <sub>8</sub> , C <sub>6</sub> H <sub>16</sub> N <sub>2</sub> , C <sub>6</sub> H <sub>16</sub> Si, C <sub>6</sub> H <sub>9</sub> Cl, C <sub>5</sub> H <sub>12</sub> N <sub>2</sub> O, C <sub>5</sub> H <sub>8</sub> O <sub>3</sub> , C <sub>4</sub> H <sub>8</sub> N <sub>2</sub> O <sub>2</sub> , C <sub>4</sub> H <sub>4</sub> O <sub>4</sub> , C <sub>4</sub> H <sub>8</sub> N <sub>2</sub> S, C <sub>4</sub> H <sub>4</sub> S <sub>2</sub> , C <sub>2</sub> H <sub>3</sub> Cl <sub>2</sub> F, C <sub>2</sub> ClF
117	C <sub>8</sub> H <sub>7</sub> N, C <sub>6</sub> H <sub>15</sub> NO, C <sub>5</sub> H <sub>11</sub> NO <sub>2</sub> , C <sub>5</sub> H <sub>6</sub> CIN, C <sub>4</sub> H <sub>7</sub> NO <sub>3</sub> , C <sub>3</sub> H <sub>4</sub> CIN
118	C <sub>6</sub> H <sub>14</sub> O <sub>2</sub> , C <sub>6</sub> H <sub>14</sub> S, C <sub>5</sub> H <sub>10</sub> O <sub>3</sub> , C <sub>9</sub> H <sub>10</sub> , C <sub>7</sub> H <sub>16</sub> N <sub>2</sub> , C <sub>5</sub> H <sub>10</sub> OS, C <sub>6</sub> H <sub>15</sub> P, C <sub>7</sub> H <sub>15</sub> F, C <sub>6</sub> H <sub>11</sub> Cl, C <sub>5</sub> H <sub>14</sub> OSi, C <sub>4</sub> H <sub>10</sub> N <sub>2</sub> O <sub>2</sub> , C <sub>4</sub> H <sub>6</sub> O <sub>4</sub> , C <sub>4</sub> H <sub>3</sub> CIS, C <sub>4</sub> H <sub>10</sub> N <sub>2</sub> S, C <sub>4</sub> H <sub>14</sub> Si <sub>2</sub> , C <sub>3</sub> H <sub>3</sub> Br, C <sub>2</sub> H <sub>2</sub> ClF <sub>3</sub> , CHCl <sub>3</sub>
119	C <sub>7</sub> H <sub>5</sub> NO, C <sub>6</sub> H <sub>4</sub> N <sub>3</sub> , C <sub>8</sub> H <sub>9</sub> N, C <sub>4</sub> H <sub>9</sub> NO <sub>3</sub> , C <sub>4</sub> H <sub>9</sub> NOS
120	C <sub>9</sub> H <sub>12</sub> , C <sub>8</sub> H <sub>8</sub> O, C <sub>4</sub> H <sub>8</sub> O <sub>2</sub> S, C <sub>4</sub> H <sub>8</sub> S <sub>2</sub> , C <sub>7</sub> H <sub>8</sub> S <sub>2</sub> , C <sub>7</sub> H <sub>8</sub> N <sub>2</sub> , C <sub>6</sub> H <sub>13</sub> Cl, C <sub>5</sub> H <sub>12</sub> O <sub>3</sub> , C <sub>6</sub> H <sub>4</sub> N <sub>2</sub> O, C <sub>5</sub> H <sub>4</sub> N <sub>4</sub> , C <sub>4</sub> H <sub>8</sub> O <sub>4</sub> , C <sub>5</sub> H <sub>9</sub> OCl, C <sub>4</sub> H <sub>12</sub> O <sub>2</sub> Si, C <sub>3</sub> H <sub>5</sub> Br, CCl <sub>2</sub> F <sub>2</sub>
121	C <sub>8</sub> H <sub>11</sub> N, C <sub>7</sub> H <sub>7</sub> NO, C <sub>6</sub> H <sub>7</sub> N <sub>3</sub> , C <sub>4</sub> H <sub>3</sub> N <sub>5</sub> , C <sub>7</sub> H <sub>3</sub> FN
122	C <sub>8</sub> H <sub>10</sub> O, C <sub>7</sub> H <sub>10</sub> N <sub>2</sub> , C <sub>7</sub> H <sub>6</sub> O <sub>2</sub> , C <sub>4</sub> H <sub>7</sub> ClO <sub>2</sub> , C <sub>4</sub> H <sub>10</sub> S <sub>2</sub> , C <sub>9</sub> H <sub>14</sub> , C <sub>3</sub> H <sub>7</sub> Br, C <sub>8</sub> H <sub>7</sub> F, C <sub>6</sub> H <sub>6</sub> N <sub>2</sub> O, C <sub>4</sub> H <sub>10</sub> O <sub>2</sub> S, C <sub>4</sub> H <sub>4</sub> Cl <sub>2</sub> , C <sub>3</sub> H <sub>6</sub> O <sub>3</sub> S, C <sub>2</sub> H <sub>6</sub> N <sub>2</sub> S <sub>2</sub> , C <sub>2</sub> H <sub>3</sub> BrO
123	C <sub>7</sub> H <sub>9</sub> NO, C <sub>6</sub> H <sub>5</sub> NO <sub>2</sub> , C <sub>8</sub> H <sub>13</sub> N, C <sub>6</sub> H <sub>9</sub> N <sub>3</sub> , C <sub>4</sub> H <sub>4</sub> F <sub>3</sub> N, C <sub>3</sub> H <sub>9</sub> NO <sub>2</sub> S
124	C <sub>3</sub> H <sub>8</sub> O <sub>3</sub> S, C <sub>8</sub> H <sub>9</sub> F, C <sub>7</sub> H <sub>12</sub> N <sub>2</sub> , C <sub>7</sub> H <sub>5</sub> FO, C <sub>5</sub> H <sub>4</sub> N <sub>2</sub> O <sub>2</sub> , C <sub>4</sub> H <sub>6</sub> F <sub>2</sub> O <sub>2</sub> , C <sub>3</sub> H <sub>5</sub> ClO <sub>3</sub> , C <sub>2</sub> H <sub>5</sub> BrO, C <sub>2</sub> H <sub>4</sub> S <sub>3</sub>
125	C <sub>8</sub> H <sub>15</sub> N, C <sub>6</sub> H <sub>11</sub> N <sub>3</sub> , C <sub>6</sub> H <sub>7</sub> NS, C <sub>7</sub> H <sub>11</sub> NO, C <sub>6</sub> H <sub>7</sub> NO <sub>2</sub> , C <sub>2</sub> H <sub>8</sub> NO <sub>3</sub> P, C <sub>2</sub> H <sub>4</sub> CINO <sub>3</sub>
126	C <sub>9</sub> H <sub>18</sub> , C <sub>8</sub> H <sub>14</sub> O, C <sub>4</sub> H <sub>8</sub> Cl <sub>2</sub> , C <sub>7</sub> H <sub>10</sub> O <sub>2</sub> , C <sub>5</sub> H <sub>6</sub> N <sub>2</sub> O <sub>2</sub> , C <sub>7</sub> H <sub>7</sub> Cl, C <sub>6</sub> H <sub>10</sub> N <sub>2</sub> O, C <sub>6</sub> H <sub>6</sub> O <sub>3</sub> , C <sub>7</sub> H <sub>10</sub> S, C <sub>3</sub> H <sub>4</sub> Cl <sub>2</sub> , C <sub>2</sub> H <sub>6</sub> S <sub>3</sub> , C <sub>7</sub> H <sub>14</sub> N <sub>2</sub> , C <sub>7</sub> H <sub>7</sub> FO, C <sub>6</sub> H <sub>6</sub> OS, C <sub>4</sub> H <sub>6</sub> N <sub>4</sub> O, C <sub>4</sub> H <sub>5</sub> F <sub>3</sub> O, C <sub>3</sub> H <sub>2</sub> N <sub>6</sub> , C <sub>2</sub> ClO <sub>2</sub>
127	C <sub>7</sub> H <sub>13</sub> NO, C <sub>8</sub> H <sub>17</sub> N, C <sub>6</sub> H <sub>2</sub> CIN, C <sub>5</sub> H <sub>9</sub> N <sub>3</sub> O, C <sub>5</sub> H <sub>5</sub> NO <sub>3</sub> , C <sub>6</sub> H <sub>9</sub> NO <sub>2</sub> , C <sub>4</sub> H <sub>5</sub> N <sub>3</sub> S, C <sub>2</sub> Cl <sub>2</sub> FN

Table 4.44. (Continued)

128	C <sub>8</sub> H <sub>16</sub> O, C <sub>9</sub> H <sub>20</sub> , C <sub>7</sub> H <sub>12</sub> O <sub>2</sub> , C <sub>7</sub> H <sub>12</sub> O <sub>2</sub> , C <sub>7</sub> H <sub>12</sub> S, C <sub>8</sub> H <sub>4</sub> N <sub>2</sub> , C <sub>6</sub> H <sub>8</sub> O <sub>3</sub> , C <sub>6</sub> H <sub>6</sub> OS, C <sub>6</sub> H <sub>5</sub> ClO, C <sub>4</sub> H <sub>4</sub> N <sub>2</sub> OS, C <sub>3</sub> H <sub>6</sub> Cl <sub>2</sub> O, C <sub>10</sub> H <sub>8</sub> , C <sub>2</sub> H <sub>6</sub> Cl <sub>2</sub> Si, C <sub>2</sub> H <sub>2</sub> Cl <sub>2</sub> O <sub>2</sub> , CH <sub>2</sub> BrCl, C <sub>2</sub> H <sub>5</sub> ClO <sub>2</sub> S, C <sub>8</sub> H <sub>13</sub> F, HI
129	C <sub>8</sub> H <sub>19</sub> N, C <sub>7</sub> H <sub>15</sub> NO, C <sub>6</sub> H <sub>11</sub> NO <sub>2</sub> , C <sub>9</sub> H <sub>7</sub> N, C <sub>7</sub> H <sub>3</sub> N <sub>3</sub> , C <sub>6</sub> H <sub>15</sub> N <sub>3</sub> , C <sub>5</sub> H <sub>7</sub> NO <sub>3</sub> , C <sub>5</sub> H <sub>7</sub> NOS, C <sub>4</sub> H <sub>7</sub> N <sub>3</sub> S, C <sub>4</sub> H <sub>4</sub> CIN <sub>3</sub> , C <sub>4</sub> H <sub>3</sub> NO <sub>2</sub> S
130	C <sub>7</sub> H <sub>14</sub> O <sub>2</sub> , C <sub>8</sub> H <sub>18</sub> O, C <sub>6</sub> H <sub>10</sub> O <sub>3</sub> , C <sub>6</sub> H <sub>6</sub> N <sub>2</sub> , C <sub>10</sub> H <sub>10</sub> , C <sub>9</sub> H <sub>6</sub> O, C <sub>7</sub> H <sub>14</sub> S, C <sub>6</sub> H <sub>14</sub> N <sub>2</sub> O, C <sub>6</sub> H <sub>4</sub> ClF, C <sub>5</sub> H <sub>6</sub> O <sub>4</sub> , C <sub>5</sub> H <sub>6</sub> S <sub>2</sub> , C <sub>3</sub> H <sub>5</sub> Cl <sub>2</sub> F, C <sub>5</sub> H <sub>10</sub> N <sub>2</sub> O <sub>2</sub> , C <sub>3</sub> H <sub>6</sub> N <sub>4</sub> S, C <sub>3</sub> H <sub>2</sub> ClF <sub>3</sub> , C <sub>3</sub> H <sub>2</sub> N <sub>2</sub> O <sub>2</sub> S, C <sub>2</sub> HCl <sub>3</sub> , CHBrF <sub>2</sub>
131	C <sub>9</sub> H <sub>9</sub> N, C <sub>7</sub> H <sub>17</sub> NO, C <sub>5</sub> H <sub>9</sub> NO <sub>3</sub> , C <sub>7</sub> H <sub>5</sub> N <sub>3</sub> , C <sub>6</sub> H <sub>17</sub> N <sub>3</sub> , C <sub>6</sub> H <sub>13</sub> NO <sub>2</sub> , C <sub>4</sub> H <sub>9</sub> N <sub>3</sub> S, CF <sub>3</sub> NOS
132	C <sub>7</sub> H <sub>16</sub> O <sub>2</sub> , C <sub>6</sub> H <sub>12</sub> O <sub>3</sub> , C <sub>10</sub> H <sub>12</sub> , C <sub>9</sub> H <sub>8</sub> O, C <sub>7</sub> H <sub>16</sub> S, C <sub>8</sub> H <sub>8</sub> N <sub>2</sub> , C <sub>6</sub> H <sub>16</sub> OSi, C <sub>5</sub> H <sub>8</sub> O <sub>4</sub> , C <sub>6</sub> H <sub>12</sub> OS, C <sub>6</sub> H <sub>9</sub> ClO, C <sub>5</sub> H <sub>12</sub> N <sub>2</sub> O <sub>2</sub> , C <sub>5</sub> H <sub>12</sub> N <sub>2</sub> S, C <sub>5</sub> H <sub>8</sub> O <sub>2</sub> S, C <sub>5</sub> H <sub>5</sub> ClO <sub>2</sub> , C <sub>4</sub> H <sub>4</sub> OS <sub>2</sub> , C <sub>3</sub> H <sub>12</sub> Si <sub>3</sub> , C <sub>3</sub> H <sub>4</sub> N <sub>2</sub> S <sub>2</sub> , C <sub>2</sub> H <sub>3</sub> Cl <sub>3</sub> , C <sub>2</sub> Cl <sub>2</sub> F <sub>2</sub> , C <sub>2</sub> F <sub>4</sub> O <sub>2</sub>
133	C <sub>9</sub> H <sub>11</sub> N, C <sub>8</sub> H <sub>7</sub> NO, C <sub>7</sub> H <sub>7</sub> N <sub>3</sub> , C <sub>5</sub> H <sub>11</sub> NO <sub>3</sub> , C <sub>4</sub> H <sub>7</sub> NO <sub>2</sub> S, C <sub>3</sub> H <sub>4</sub> BrN, C <sub>2</sub> H <sub>3</sub> ClF <sub>3</sub> N
134	C <sub>9</sub> H <sub>10</sub> O, C <sub>10</sub> H <sub>14</sub> , C <sub>6</sub> H <sub>14</sub> O <sub>3</sub> , C <sub>6</sub> H <sub>6</sub> N <sub>4</sub> , C <sub>5</sub> H <sub>10</sub> O <sub>2</sub> S, C <sub>8</sub> H <sub>6</sub> O <sub>2</sub> , C <sub>8</sub> H <sub>6</sub> O <sub>2</sub> , C <sub>8</sub> H <sub>10</sub> N <sub>2</sub> , C <sub>8</sub> H <sub>6</sub> S, C <sub>7</sub> H <sub>15</sub> Cl, C <sub>7</sub> H <sub>6</sub> N <sub>2</sub> O, C <sub>6</sub> H <sub>14</sub> OS, C <sub>6</sub> H <sub>11</sub> ClO, C <sub>5</sub> H <sub>11</sub> ClSi, C <sub>5</sub> H <sub>10</sub> S <sub>2</sub> , C <sub>5</sub> H <sub>7</sub> ClO <sub>2</sub> , C <sub>3</sub> H <sub>3</sub> F <sub>5</sub> , C <sub>3</sub> Cl <sub>2</sub> N <sub>2</sub> , C <sub>2</sub> H <sub>2</sub> Cl <sub>2</sub> F <sub>2</sub>
135	C <sub>9</sub> H <sub>13</sub> N, C <sub>8</sub> H <sub>9</sub> NO, C <sub>7</sub> H <sub>5</sub> NS, C <sub>5</sub> H <sub>5</sub> N <sub>5</sub> , C <sub>7</sub> H <sub>5</sub> NO <sub>2</sub> , C <sub>6</sub> H <sub>5</sub> N <sub>3</sub> O, C <sub>4</sub> H <sub>9</sub> NO <sub>2</sub> S, C <sub>3</sub> H <sub>6</sub> BrN, C <sub>3</sub> F <sub>3</sub> N <sub>3</sub>
136	C <sub>9</sub> H <sub>12</sub> O, C <sub>10</sub> H <sub>16</sub> , C <sub>8</sub> H <sub>8</sub> O <sub>2</sub> , C <sub>4</sub> H <sub>9</sub> Br, C <sub>8</sub> H <sub>12</sub> N <sub>2</sub> , C <sub>8</sub> H <sub>12</sub> Si, C <sub>8</sub> H <sub>8</sub> S, C <sub>8</sub> H <sub>5</sub> Cl, C <sub>7</sub> H <sub>8</sub> N <sub>2</sub> O, C <sub>7</sub> H <sub>4</sub> O <sub>3</sub> , C <sub>5</sub> H <sub>12</sub> S <sub>2</sub> , C <sub>6</sub> H <sub>4</sub> N <sub>2</sub> O <sub>2</sub> , C <sub>6</sub> H <sub>4</sub> N <sub>2</sub> O <sub>2</sub> , C <sub>6</sub> H <sub>4</sub> N <sub>2</sub> S, C <sub>5</sub> H <sub>12</sub> O <sub>4</sub> , C <sub>5</sub> H <sub>12</sub> OS, C <sub>5</sub> H <sub>9</sub> ClO <sub>2</sub> , C <sub>5</sub> H <sub>4</sub> N <sub>4</sub> O, C <sub>2</sub> HClF <sub>4</sub> , CCl <sub>3</sub> F
137	C <sub>8</sub> H <sub>11</sub> NO, C <sub>9</sub> H <sub>15</sub> N, C <sub>7</sub> H <sub>7</sub> NO <sub>2</sub> , C <sub>7</sub> H <sub>7</sub> NS, C <sub>7</sub> H <sub>4</sub> CIN, C <sub>6</sub> H <sub>7</sub> N <sub>3</sub> O, C <sub>5</sub> H <sub>6</sub> F <sub>3</sub> N, C <sub>5</sub> H <sub>3</sub> N <sub>3</sub> S, C <sub>3</sub> H <sub>7</sub> NO <sub>5</sub>
138	C <sub>10</sub> H <sub>18</sub> , C <sub>9</sub> H <sub>14</sub> O, C <sub>8</sub> H <sub>10</sub> O <sub>2</sub> , C <sub>8</sub> H <sub>10</sub> S, C <sub>7</sub> H <sub>6</sub> O <sub>3</sub> , C <sub>6</sub> H <sub>6</sub> N <sub>2</sub> O <sub>2</sub> , C <sub>8</sub> H <sub>7</sub> Cl, C <sub>7</sub> H <sub>10</sub> N <sub>2</sub> O, C <sub>7</sub> H <sub>6</sub> OS, C <sub>6</sub> H <sub>10</sub> N <sub>4</sub> , C <sub>5</sub> H <sub>6</sub> N <sub>4</sub> O, C <sub>4</sub> H <sub>11</sub> O <sub>3</sub> P, C <sub>4</sub> H <sub>10</sub> O <sub>3</sub> S, C <sub>3</sub> H <sub>7</sub> BrO, C <sub>3</sub> H <sub>6</sub> S <sub>3</sub> , C <sub>2</sub> H <sub>3</sub> BrO <sub>2</sub> , C <sub>2</sub> F <sub>6</sub>
139	C <sub>7</sub> H <sub>9</sub> NS, C <sub>9</sub> H <sub>17</sub> N, C <sub>7</sub> H <sub>13</sub> N <sub>3</sub> , C <sub>7</sub> H <sub>9</sub> NO <sub>2</sub> , C <sub>6</sub> H <sub>5</sub> NO <sub>3</sub> , C <sub>5</sub> H <sub>5</sub> N <sub>3</sub> O <sub>2</sub> , C <sub>6</sub> H <sub>5</sub> NOS
140	C <sub>9</sub> H <sub>16</sub> O, C <sub>10</sub> H <sub>20</sub> , C <sub>8</sub> H <sub>9</sub> Cl, C <sub>8</sub> H <sub>12</sub> O <sub>2</sub> , C <sub>8</sub> H <sub>12</sub> S, C <sub>7</sub> H <sub>8</sub> O <sub>3</sub> , C <sub>5</sub> H <sub>10</sub> Cl <sub>2</sub> , C <sub>7</sub> H <sub>8</sub> OS, C <sub>8</sub> H <sub>9</sub> FO, C <sub>8</sub> H <sub>16</sub> N <sub>2</sub> , C <sub>8</sub> H <sub>6</sub> F <sub>2</sub> , C <sub>7</sub> H <sub>5</sub> ClO, C <sub>6</sub> H <sub>8</sub> N <sub>2</sub> O <sub>2</sub> , C <sub>6</sub> H <sub>8</sub> N <sub>2</sub> S, C <sub>6</sub> H <sub>4</sub> O <sub>4</sub> , C <sub>6</sub> H <sub>4</sub> S <sub>2</sub> , C <sub>4</sub> H <sub>6</sub> Cl <sub>2</sub> O, C <sub>2</sub> H <sub>2</sub> BrCl
141	C <sub>8</sub> H <sub>15</sub> NO, C <sub>7</sub> H <sub>11</sub> NO <sub>2</sub> , C <sub>6</sub> H <sub>11</sub> N <sub>3</sub> O, C <sub>6</sub> H <sub>7</sub> NO <sub>3</sub> , C <sub>9</sub> H <sub>19</sub> N, C <sub>7</sub> H <sub>15</sub> N <sub>3</sub> , C <sub>5</sub> H <sub>7</sub> N <sub>3</sub> S, C <sub>4</sub> H <sub>6</sub> F <sub>3</sub> NO, C <sub>4</sub> H <sub>3</sub> F <sub>4</sub> N
142	C <sub>8</sub> H <sub>14</sub> O <sub>2</sub> , C <sub>9</sub> H <sub>18</sub> O, C <sub>10</sub> H <sub>22</sub> , C <sub>6</sub> H <sub>6</sub> O <sub>4</sub> , C <sub>8</sub> H <sub>14</sub> S, C <sub>3</sub> H <sub>4</sub> Cl <sub>2</sub> O <sub>2</sub> , C <sub>7</sub> H <sub>7</sub> ClO, C <sub>11</sub> H <sub>10</sub> , C <sub>9</sub> H <sub>15</sub> F, C <sub>8</sub> H <sub>18</sub> N <sub>2</sub> , C <sub>7</sub> H <sub>14</sub> N <sub>2</sub> O, C <sub>7</sub> H <sub>10</sub> O <sub>3</sub> , C <sub>6</sub> H <sub>10</sub> N <sub>2</sub> O <sub>2</sub> , C <sub>6</sub> H <sub>6</sub> O <sub>2</sub> S, C <sub>5</sub> H <sub>6</sub> N <sub>2</sub> OS, C <sub>4</sub> H <sub>8</sub> Cl <sub>2</sub> O, C <sub>4</sub> H <sub>5</sub> ClF <sub>2</sub> O, C <sub>2</sub> H <sub>4</sub> BrCl, C <sub>2</sub> HBrF <sub>2</sub> , CH <sub>3</sub> I
143	C <sub>10</sub> H <sub>9</sub> N, C <sub>8</sub> H <sub>17</sub> NO, C <sub>7</sub> H <sub>13</sub> NO <sub>2</sub> , C <sub>9</sub> H <sub>21</sub> N, C <sub>7</sub> H <sub>10</sub> CIN, C <sub>6</sub> H <sub>13</sub> N <sub>3</sub> O, C <sub>6</sub> H <sub>9</sub> NO <sub>3</sub> , C <sub>4</sub> H <sub>9</sub> NOS, C <sub>5</sub> H <sub>9</sub> N <sub>3</sub> S, C <sub>2</sub> Cl <sub>3</sub> N
144	C <sub>8</sub> H <sub>16</sub> O <sub>2</sub> , C <sub>9</sub> H <sub>20</sub> O, C <sub>9</sub> H <sub>8</sub> N <sub>2</sub> , C <sub>7</sub> H <sub>12</sub> O <sub>3</sub> , C <sub>6</sub> H <sub>8</sub> O <sub>4</sub> , C <sub>10</sub> H <sub>8</sub> O, C <sub>11</sub> H <sub>12</sub> , C <sub>8</sub> H <sub>16</sub> S, C <sub>8</sub> H <sub>13</sub> Cl, C <sub>7</sub> H <sub>16</sub> N <sub>2</sub> O, C <sub>7</sub> H <sub>12</sub> OS, C <sub>6</sub> H <sub>9</sub> CIN <sub>2</sub> , C <sub>5</sub> H <sub>8</sub> N <sub>2</sub> O <sub>3</sub> , C <sub>3</sub> H <sub>3</sub> Cl <sub>3</sub>
145	C <sub>10</sub> H <sub>11</sub> N, C <sub>9</sub> H <sub>7</sub> NO, C <sub>8</sub> H <sub>7</sub> N <sub>3</sub> , C <sub>7</sub> H <sub>15</sub> NO <sub>2</sub> , C <sub>6</sub> H <sub>11</sub> NO <sub>3</sub> , C <sub>5</sub> H <sub>11</sub> N <sub>3</sub> S, C <sub>3</sub> H <sub>3</sub> N <sub>3</sub> O <sub>2</sub> S
146	C <sub>8</sub> H <sub>18</sub> O <sub>2</sub> , C <sub>7</sub> H <sub>14</sub> O <sub>3</sub> , C <sub>8</sub> H <sub>18</sub> S, C <sub>6</sub> H <sub>10</sub> O <sub>4</sub> , C <sub>6</sub> H <sub>10</sub> O <sub>2</sub> S, C <sub>11</sub> H <sub>14</sub> , C <sub>10</sub> H <sub>10</sub> O, C <sub>9</sub> H <sub>10</sub> N <sub>2</sub> , C <sub>10</sub> H <sub>7</sub> F, C <sub>8</sub> H <sub>6</sub> N <sub>2</sub> O, C <sub>7</sub> H <sub>18</sub> OSi, C <sub>7</sub> H <sub>14</sub> OS, C <sub>7</sub> H <sub>6</sub> N <sub>4</sub> , C <sub>6</sub> H <sub>14</sub> N <sub>2</sub> O <sub>2</sub> , C <sub>6</sub> H <sub>4</sub> Cl <sub>2</sub> , C <sub>5</sub> H <sub>6</sub> OS <sub>2</sub> , C <sub>3</sub> H <sub>5</sub> Cl <sub>3</sub> , C <sub>3</sub> H <sub>3</sub> BrN <sub>2</sub> , C <sub>2</sub> HCl <sub>3</sub> O, CHBrClF
147	C <sub>10</sub> H <sub>13</sub> N, C <sub>9</sub> H <sub>9</sub> NO, C <sub>8</sub> H <sub>9</sub> N <sub>3</sub> , C <sub>8</sub> H <sub>5</sub> NO <sub>2</sub> , C <sub>7</sub> H <sub>5</sub> N <sub>3</sub> O, C <sub>6</sub> H <sub>13</sub> NO <sub>3</sub> , C <sub>5</sub> H <sub>9</sub> NO <sub>4</sub> , C <sub>2</sub> H <sub>2</sub> BrN <sub>3</sub>

Table 4.44. (Continued)

148	C <sub>11</sub> H <sub>16</sub> , C <sub>10</sub> H <sub>12</sub> O, C <sub>9</sub> H <sub>8</sub> O <sub>2</sub> , C <sub>7</sub> H <sub>8</sub> N <sub>4</sub> , C <sub>9</sub> H <sub>12</sub> N <sub>2</sub> , C <sub>6</sub> H <sub>12</sub> O <sub>2</sub> S, C <sub>8</sub> H <sub>17</sub> Cl, C <sub>8</sub> H <sub>8</sub> N <sub>2</sub> O, C <sub>8</sub> H <sub>4</sub> O <sub>3</sub> , C <sub>9</sub> H <sub>8</sub> S, C <sub>7</sub> H <sub>16</sub> O <sub>3</sub> , C <sub>7</sub> H <sub>16</sub> OS, C <sub>7</sub> H <sub>4</sub> N <sub>2</sub> O <sub>2</sub> , C <sub>6</sub> H <sub>12</sub> O <sub>4</sub> , C <sub>6</sub> H <sub>4</sub> N <sub>4</sub> O, C <sub>5</sub> H <sub>8</sub> O <sub>3</sub> S, C <sub>3</sub> HF <sub>5</sub> O, C <sub>2</sub> H <sub>3</sub> Cl <sub>3</sub> O, C <sub>2</sub> Cl <sub>3</sub> F, CBrF <sub>3</sub>
149	C <sub>10</sub> H <sub>15</sub> N, C <sub>9</sub> H <sub>11</sub> NO, C <sub>7</sub> H <sub>7</sub> N <sub>3</sub> O, C <sub>8</sub> H <sub>11</sub> N <sub>3</sub> , C <sub>8</sub> H <sub>7</sub> NO <sub>2</sub>
150	C <sub>10</sub> H <sub>14</sub> O, C <sub>9</sub> H <sub>10</sub> O <sub>2</sub> , C <sub>6</sub> H <sub>14</sub> S <sub>2</sub> , C <sub>8</sub> H <sub>10</sub> N <sub>2</sub> O, C <sub>6</sub> H <sub>11</sub> ClO <sub>2</sub> , C <sub>5</sub> H <sub>11</sub> Br, C <sub>11</sub> H <sub>18</sub> , C <sub>8</sub> H <sub>14</sub> N <sub>2</sub> , C <sub>9</sub> H <sub>10</sub> S, C <sub>8</sub> H <sub>6</sub> O <sub>3</sub> , C <sub>7</sub> H <sub>6</sub> N <sub>2</sub> S, C <sub>6</sub> H <sub>14</sub> O <sub>4</sub> , C <sub>6</sub> H <sub>14</sub> O <sub>2</sub> S, C <sub>6</sub> H <sub>6</sub> N <sub>4</sub> O, C <sub>6</sub> H <sub>2</sub> F <sub>4</sub> , C <sub>3</sub> F <sub>6</sub> , C <sub>2</sub> H <sub>2</sub> Cl <sub>3</sub> F
151 <sup>2</sup>	(see m/z 137) C <sub>7</sub> H <sub>5</sub> NO <sub>3</sub>
152	(see m/z 138) C <sub>8</sub> H <sub>5</sub> ClO, C <sub>6</sub> H <sub>10</sub> Cl <sub>2</sub> , CCl <sub>4</sub>
153	(see m/z 139) C <sub>9</sub> H <sub>15</sub> NO, C <sub>7</sub> H <sub>5</sub> ClNO
154	(see m/z 140; C <sub>10</sub> H <sub>8</sub> O has highest occurrence of data base) C <sub>8</sub> H <sub>7</sub> ClO, C <sub>2</sub> ClO, C <sub>2</sub> ClF <sub>5</sub> , C <sub>8</sub> H <sub>10</sub> OS
155	(see m/z 141) C <sub>8</sub> H <sub>13</sub> NS
156	(see m/z 142) C <sub>7</sub> H <sub>5</sub> ClO <sub>2</sub> , C <sub>6</sub> H <sub>5</sub> Br, C <sub>6</sub> H <sub>4</sub> O <sub>5</sub>
157	(see m/z 143) C <sub>5</sub> H <sub>4</sub> BrN
158	(see m/z 144) CF <sub>6</sub> S
159	(see m/z 145) C <sub>6</sub> H <sub>9</sub> NS <sub>2</sub> , C <sub>5</sub> H <sub>6</sub> ClN <sub>3</sub> O
160	(see m/z 146) C <sub>7</sub> H <sub>20</sub> Si <sub>2</sub> , C <sub>6</sub> H <sub>9</sub> Br, C <sub>5</sub> H <sub>16</sub> Si <sub>3</sub>
161	(see m/z 147) C <sub>6</sub> H <sub>5</sub> Cl <sub>2</sub> N
162	(see m/z 148) C <sub>10</sub> H <sub>7</sub> Cl, C <sub>8</sub> H <sub>6</sub> N <sub>2</sub> S, C <sub>6</sub> H <sub>11</sub> Br, C <sub>4</sub> F <sub>6</sub> , C <sub>6</sub> H <sub>4</sub> Cl <sub>2</sub> O, CHBrCl <sub>2</sub>
163	(see m/z 149) C <sub>7</sub> H <sub>9</sub> N <sub>5</sub> , C <sub>3</sub> H <sub>2</sub> BrNS, CCl <sub>3</sub> NO <sub>2</sub>
164	(see m/z 150) C <sub>9</sub> H <sub>8</sub> OS, C <sub>8</sub> H <sub>8</sub> N <sub>2</sub> O <sub>2</sub> , C <sub>6</sub> H <sub>5</sub> ClN <sub>2</sub> , C <sub>7</sub> H <sub>16</sub> O <sub>4</sub> , C <sub>6</sub> H <sub>4</sub> N <sub>4</sub> O <sub>2</sub> , C <sub>5</sub> H <sub>9</sub> BrO, C <sub>5</sub> H <sub>8</sub> O <sub>2</sub> S <sub>2</sub> , C <sub>2</sub> Cl <sub>4</sub> , CBrClF <sub>2</sub>
165	(see m/z 137, 151) C <sub>6</sub> H <sub>7</sub> N <sub>5</sub> O
166	(see m/z 138, 152) C <sub>13</sub> H <sub>10</sub> , C <sub>8</sub> H <sub>6</sub> O <sub>4</sub> , C <sub>8</sub> H <sub>6</sub> O <sub>2</sub> S, C <sub>8</sub> H <sub>6</sub> S <sub>2</sub> , C <sub>6</sub> H <sub>6</sub> N <sub>4</sub> O <sub>2</sub> , C <sub>6</sub> H <sub>6</sub> N <sub>4</sub> S, C <sub>3</sub> H <sub>6</sub> N <sub>2</sub> O <sub>6</sub> , C <sub>3</sub> ClF <sub>5</sub> , C <sub>3</sub> F <sub>6</sub> O
167	(see m/z 139, 153) C <sub>7</sub> H <sub>13</sub> N <sub>5</sub> , C <sub>7</sub> H <sub>5</sub> NO <sub>4</sub> , C <sub>7</sub> H <sub>5</sub> NS <sub>2</sub>
168	(see m/z 140, 154) C <sub>13</sub> H <sub>12</sub> , C <sub>9</sub> H <sub>16</sub> N <sub>2</sub> O, C <sub>8</sub> H <sub>8</sub> O <sub>4</sub> , C <sub>8</sub> H <sub>8</sub> O <sub>2</sub> S, C <sub>8</sub> H <sub>8</sub> S <sub>2</sub> , C <sub>6</sub> H <sub>4</sub> N <sub>2</sub> O <sub>4</sub> , C <sub>6</sub> HF <sub>5</sub> , C <sub>3</sub> H <sub>5</sub> I, C <sub>2</sub> HCl <sub>3</sub> F <sub>2</sub> , Cl <sub>4</sub> Si
169	(see m/z 141, 155) C <sub>8</sub> H <sub>8</sub> ClNO, C <sub>7</sub> H <sub>7</sub> NO <sub>2</sub> S
170	(see m/z 142, 156) C <sub>9</sub> H <sub>16</sub> NO <sub>2</sub> , C <sub>8</sub> H <sub>10</sub> S <sub>2</sub> , C <sub>3</sub> H <sub>6</sub> S <sub>4</sub> , C <sub>2</sub> Cl <sub>2</sub> F <sub>4</sub> , C <sub>2</sub> F <sub>6</sub> S
171	(see m/z 143, 157) C <sub>7</sub> H <sub>9</sub> NO <sub>2</sub> S, C <sub>6</sub> H <sub>9</sub> N <sub>3</sub> O <sub>3</sub> , C <sub>5</sub> H <sub>2</sub> ClN <sub>3</sub> S
172	(see m/z 144, 158) C <sub>12</sub> H <sub>9</sub> F, C <sub>8</sub> H <sub>6</sub> Cl <sub>2</sub> , C <sub>8</sub> H <sub>9</sub> ClO <sub>2</sub> , C <sub>7</sub> H <sub>8</sub> O <sub>3</sub> S, C <sub>6</sub> H <sub>8</sub> N <sub>2</sub> O <sub>2</sub> S, C <sub>6</sub> H <sub>5</sub> BrO, C <sub>6</sub> H <sub>4</sub> O <sub>6</sub> , CH <sub>2</sub> Br <sub>2</sub>
	(see m/z 145, 159) C <sub>6</sub> H <sub>4</sub> CINOS, C <sub>5</sub> H <sub>8</sub> CIN <sub>5</sub>
173	(see m/z 146, 160) C <sub>11</sub> H <sub>10</sub> S, C <sub>10</sub> H <sub>6</sub> O <sub>3</sub> , C <sub>10</sub> H <sub>6</sub> OS, C <sub>9</sub> H <sub>6</sub> N <sub>2</sub> O <sub>2</sub> , C <sub>6</sub> H <sub>4</sub> BrF, C <sub>5</sub> F <sub>6</sub>
174	(see m/z 147, 161) C <sub>8</sub> H <sub>5</sub> N <sub>3</sub> O <sub>2</sub> , C <sub>5</sub> H <sub>9</sub> N <sub>3</sub> S <sub>2</sub> , C <sub>7</sub> H <sub>10</sub> CINO <sub>2</sub>

1. Compositions are listed by m/z value, ranked in decreasing order of Occurrence probability for compounds in the *Registry of Mass Spectral Data* (Stenhamer *et al.* 1974). Only the more probable combinations of the elements H, C, N, O, F, Si, P, S, Cl, Br and I are included. Note that these are odd-electron ion compositions; many common even-electron fragment ions have compositions differing by  $\pm 1$  hydrogen atom, and can therefore be found  $\pm 1$  mass unit from those listed. The above table can also be used to suggest possible elemental compositions of fragment ions
2. For masses above 150 the only compositions included are those for which a corresponding composition differing by a CH<sub>2</sub> less unit.

*This page intentionally left blank.*

## Chapter 5

### Units and Measurements

#### 5.1 Fundamental Physical Constants

#### 5.2 Units

#### 5.3 Prefixes

#### 5.4 Conversion Factors

### 5.1 Fundamental Physical Constants

Table 5.1. Physical Constants

Symbol	Name	Value	Units
amu	Atomic Mass Unit	$1.6605655 \times 10^{-24}$	g
$N_0$	Avogadro's Number	$6.022045 \times 10^{23}$	units/mole
K	Boltzman's Constant	$1.380663 \times 10^{-16}$	erg/ $^{\circ}$ K
e	Electron Charge	$1.6021892 \times 10^{-19}$	coulomb
$m_e$	Electron Rest Mass	$9.109534 \times 10^{-28}$	g
		$5.485803 \times 10^{-4}$	amu
eV	Electron Volt	$1.60 \times 10^{-19}$	joule
F	Faraday's Constant	$9.648456 \times 10^4$	coulombs/equiv
		$2.8925342 \times 10$	cal/volt
R	Gas Constant	$8.2056 \times 10^{-2}$	1 atm/ $^{\circ}$ K-mole
		8.3144	joules/ $^{\circ}$ K-mole
		$8.3144 \times 10^7$	erg/ $^{\circ}$ K-mole
		1.9872	cal/ $^{\circ}$ K-mole
$m_n$	Neutron Rest Mass	$1.6749543 \times 10^{-24}$	g
		1.0086650	amu
h	Planck's Constant	$6.626176 \times 10^{-27}$	erg/sec
$m_p$	Proton Rest Mass	$1.6726485 \times 10^{-24}$	g
		1.0072674	amu
$R^*$	Rydberg Constant	$1.0973718 \times 10^5$	cm
c	Speed of Light (in vacuum)	$2.9979246 \times 10^{10}$	cm/sec
atm	Standard Pressure	101.3	kPa
		760	mmHg

## 5.2 Units

### 5.2.1 Base SI Units

Table 5.2. Base SI Units

Symbol	Physical Quantity	Name of Base Unit
m	Length	Meter
kg	Mass	Kilogram
s	Time	Second
A	Electric Current	Ampere
K	Thermodynamic Temperature	Kelvin
mol	Amount of Substance	Mole
cd	Luminous Intensity	Candela

### 5.2.2 Derived SI Units

Table 5.3. Common Derived SI Units

Symbol	Physical Quantity	Name of Unit	Definition of Unit
Å	length	Angstrom	$10^{-10} \text{ m}$
μ	length	Micron	$10^{-6} \text{ m}$
dyn	force	Dyne	$10^{-5} \text{ N}$
bar	pressure	Bar	$10^{-5} \text{ N} \cdot \text{m}^{-2}$
erg	energy	Erg	$10^{-7} \text{ J}$

Table 5.4. Derived SI Units with Special Names

Symbol	Physical Quantity	Name of Unit	Definition of Unit
Hz	frequency	Hertz	1/s
J	energy	Joule	$\text{N} \cdot \text{m}$
N	force	Newton	$\text{kg} \cdot \text{m/s}^2$
W	power	Watt	J/s
Pa	pressure	Pascal.	$\text{N/m}^2$
C	electric charge	Coulomb	$\text{A} \cdot \text{s}$
V	difference	Volt	W/A
Ohm	electrical resistance	Ohm	V/A

### 5.2.3 Non SI Units

Table 5.5. Non SI Units

Symbol	Physical Quantity	Name of Unit	Definition of Unit
in	length	inch	$2.54 \cdot 10^{-2} \text{ m}$
lb	mass	pound	$0.45359237 \text{ kg}$
kgf	force	kilogram-force	$9.80665 \text{ N}$
atm	pressure	atmosphere	$101.325 \text{ N} \cdot \text{m}^{-2}$
torr	pressure	torr	$(101.325/760)\text{N} \cdot \text{m}^{-2}$
BTU	energy	British Thermal Unit	$1055.056 \text{ J}$
kW	energy	kilowatt-hour	$3.6 \times 10^6 \text{ J}$
cal <sub>th</sub>	energy	thermochemical calorie	$4.184 \text{ J}$

Table 5.5. (Continued)

eV	energy	electron Volt	$1.60219 \cdot 10^{-19}$ J
amu	mass	atomic mass unit	$1.6605655 \cdot 10^{-27}$ kg
D	electric dipole moment	Debye	$3.3356 \cdot 10^{-30}$ A · m · s
F	charge per molecule	Faraday	$9.648456 \cdot 10^4$ C mol <sup>-1</sup>

## 5.3 Prefixes

### 5.3.1 SI Prefixes

Table 5.6. SI Prefixes

Factor	Prefix	Symbol
$10^{15}$	penta	P
$10^{12}$	tera	T
$10^9$	giga	G
$10^6$	mega	M
$10^3$	kilo	k
$10^2$	hecto	h
$10^1$	deka	da
$10^{-1}$	deci	d
$10^{-2}$	centi	c
$10^{-3}$	milli	m
$10^{-6}$	micro	$\mu$
$10^{-9}$	nano	n
$10^{-12}$	pico	p
$10^{-15}$	femto	f

### 5.3.2 Greek Prefixes

Table 5.7. Greek Prefixes

Value	Prefix
1	mono
2	di
3	tri
4	tetra
5	penta
6	hexa
7	hepta
8	octa
9	ennea
10	deca

## 5.4 Conversion Factors

### 5.4.1 Linear Conversion

1 inch = 2.5400 centimeters  
 1 foot = 0.3048 meter  
 1 yard = 0.9144 meter  
 1 mile = 1.6093 kilometers

1 centimeter = 0.3937 inch  
 1 meter = 3.281 feet  
 1 meter = 1.0936 yards  
 1 kilometer = 0.62137 miles

### 5.4.2 Area Conversion

1 sq. inch = 6.4516 sq. centimeters  
 1 sq. foot = 0.0929 sq. meter  
 1 sq. yard = 0.8361 sq. meter  
 1 sq. mile = 2.59 sq kilometers

1 sq. centimeter = 0.155 sq. inch  
 1 sq. meter = 10.764 sq. feet  
 1 sq. meter = 1.196 sq. yards  
 1 sq. kilometers = 0.3861 sq. mile

### 5.4.3 Cubic Conversion

1 cu. inch = 16.3872 cu. centimeters  
 1 cu. foot = 28.317 cu. centimeters  
 1 cu. yard = 0.7645 cu. meter

1 cu. centimeter = 0.0610 cu. inch  
 1 cu. decimeter = 0.0353 cu. foot  
 1 cu. meter = 1.3079 cu. yards

### 5.4.4 Capacity Conversion

1 fluid ounce = 29.5730 milliliters  
 1 liquid pint = 0.4732 liter  
 1 liquid quart = 0.9463 liter  
 1 gallon = 3.7853 liters  
 1 dry quart = 1.1012 liters

1 milliliter = 0.0338 fluid ounce  
 1 liter = 2.1134 fluid pints  
 1 liter = 1.0567 liquid quarts  
 1 liter = 0.2642 gallon  
 1 liter = 0.9081 dry quart

### 5.4.5 Weight Conversion

1 ounce = 28.350 grams  
 1 pound = 0.4536 kilograms

1 gram = 0.0353 ounce  
 1 kilogram = 2.2046 pounds

### 5.4.6 Temperature Conversion

Temperature given in	To Convert to		
	°C	K	°F
°C	°C	°C + 273.15	1.8 °C + 32
K	K - 273.15	K	1.8K - 459.4
°F	0.556 °F - 17.8	0.556 °F + 255.3	°F

## Chapter 6

### Mathematical Concepts

- 6.1 *Algebraic Formulas*
- 6.2 *Plane Figure Formulas*
- 6.3 *Solid Figure Formulas*

#### 6.1 Algebraic Formulas

##### 6.1.1 Laws of Exponents

$$x^m \cdot x^n = x^{m+n}$$

$$x^m \div x^n = x^{m-n}$$

$$(x^m)^n = x^{m \cdot n}$$

$$(x \cdot y)^m = x^m \cdot y^m$$

$$(x/y)^m = x^m/y^m$$

$$x^0 = 1$$

$$x^{1/n} = \sqrt[n]{x}$$

$$x^{-m} = 1/x^m$$

##### 6.1.2 Laws of Logarithms

$$\log (A \cdot B) = \log A + \log B$$

$$\log \frac{A}{B} = \log A - \log B$$

$$\log A^b = b \log A$$

$$\log \sqrt[b]{A} = \frac{\log A}{b} = \frac{1}{b} \log A$$

##### 6.1.3 Quadratic Equation

$$ax^2 + bx + c = 0 \text{ (Where } a \neq 0 \text{ and } a, b, \text{ and } c \text{ are real numbers)}$$

If the roots of  $ax^2 + bx + c = 0$  are represented by  $r_1$  and  $r_2$ , then:

$$1. r_1 = \frac{-b + \sqrt{b^2 - 4ac}}{2a} \quad \text{and} \quad r_2 = \frac{-b - \sqrt{b^2 - 4ac}}{2a}$$

$$2. r_1 + r_2 = -b/a$$

$$3. r_1 r_2 = c/a$$

$$4. x^2 - (r_1 + r_2)x + r_1 r_2 = 0$$

Using the discriminant to determine the nature of the roots of  $ax^2 + bx + c = 0$ :

5. If  $b^2 - 4ac$  is zero or positive, the roots are real.

6. If  $b^2 - 4ac$  is negative, the roots are imaginary.

7. If  $b^2 - 4ac$  is zero, the roots are equal.

8. If  $b^2 - 4ac$  is not zero, the roots are unequal.

9. If  $b^2 - 4ac$  is a perfect square, the roots are rational numbers.

10. If  $b^2 - 4ac$  is positive and not a perfect square, the roots are irrational numbers.

#### 6.1.4 Graphs (a, b, c, m, and r are real numbers)

1. The slope of a line that passes through two points  $P_1(x_1, y_1)$  and  $P_2(x_2, y_2)$ ,  $x_1 \neq x_2$ :


$$m = \text{slope of } P_1 P_2 = \frac{y_2 - y_1}{x_2 - x_1} = \frac{\Delta y}{\Delta x}$$

2. Equation of a straight line:  $y = mx + b$  where  $m$  = slope of the line and  $b$  = y intercept.

3. Equation of a parabola:  $y = ax^2 + bx + c$  or  $x = ay^2 + by + c$ ,  $a \neq 0$ .

4. Equation of a circle:  $(x-h)^2 + (y-k)^2 = r^2$  where the center is  $(h, k)$  and the radius is  $r$ ,  $r > 0$ .


5. Equation of a circle:  $x^2 + y^2 = r^2$  where the center is at the origin and the radius is  $r$ ,  $r > 0$ .



6. Equation of an ellipse:  $ax^2 + by^2 = c$  where the center is at the origin; a, b, and c are positive;  $c \neq 0$ .
7. Equation of a hyperbola:  $ax^2 - by^2 = c$ ,  $ay^2 - bx^2 = c$  where the center is at the origin; a and b are positive. Also  $xy = k$ , k is a constant.
8. At the turning point of the parabola  $y = ax^2 + bx + c$ ,  $x = -b/2a$
9. The graph of the parabola  $y = ax^2 + bx + c$  opens upward and has a minimum turning point when a is positive,  $a > 0$ .
10. The graph of the parabola  $y = ax^2 + bx + c$  opens downward and has a maximum turning point when a is negative,  $a < 0$ .

## 6.2 Plane Figure Formulas

### 6.2.1 Rectangle




$$\text{Area } A = ab$$

$$\text{Perimeter } P = 2(a + b)$$

$$\text{Diagonal } d = \sqrt{a^2 + b^2}$$

If  $a = b$  then it is a square

### 6.2.2 Parallelogram



$$\text{Area } A = ah_a = bh_b = ab \sin \theta$$


$$\text{Perimeter } P = 2(a + b)$$

Diagonal:

$$d1 = \sqrt{a^2 + b^2 - 2ab \cos \theta}$$

$$d2 = \sqrt{a^2 + b^2 + 2ab \cos \theta}$$

## 6.2.3 Trapezoid



$$\text{Area } A = \frac{(a+b)h}{2}$$

$$\text{Perimeter } P = a + b + c + d$$


$$\text{Diagonal } d_1 = \sqrt{ab + \frac{ac^2 - bd^2}{a-b}}$$

$$d_2 = \sqrt{ab + \frac{ad^2 - bc^2}{a-b}}$$

$$\text{Height } h = \frac{2}{a-b} \cdot \sqrt{s(s-a+b)(s-c)(s-d)}$$

$$\text{where } s = \frac{1}{2}(a+b+c+d)$$

## 6.2.4 Equilateral Triangle



$a$  = all sides equal

$$\text{Area } A = a^2 \frac{\sqrt{3}}{4} = 0.433 a^2$$

$$\text{Perimeter } P = 3a$$

$$h = \frac{\sqrt{3}}{2} a = 0.866 a$$

## 6.2.5 Circle

$$\text{Area of circle } A = \pi r^2 = 3.14159 r^2$$

$$= \pi \frac{d^2}{4} = 0.78539 d$$

$$\text{Area of sector } AsB0A = \frac{\pi \theta r^2}{360}$$


$$\text{Area of segment } AsB1A = \frac{\pi r^2 \theta}{360} - \frac{r^2 \sin \theta}{2}$$

$$\text{Circumference} = 2 \pi r = 6.28318 r$$


$$= \pi d = 3.14159 d$$

$$\text{Length of chord } AlB = 2r \sin \frac{\theta}{2}$$

$$\text{Length of arc } AsB = \frac{\pi r \theta}{180}$$




## 6.2.6 Ellipse



$$\text{Area } A = \pi ab = 3.14159 ab$$

$$\text{Circumference} \approx 2\pi\sqrt{\frac{a^2+b^2}{2}}$$

## 6.2.7 Parabola




$$\text{Area ABCA} = \frac{2}{3} ab$$

$$\text{Length of arc ABC} = b \left\{ \frac{1}{2} (1 + 16 (a/b)^2)^{1/2} + \frac{1}{8 (\frac{a}{b})} \ln [4 n + (1 + 16 (a/b)^2)^{1/2}] \right\}$$

## 6.3 Solid Figure Formulas

## 6.3.1 Parallelpiped



$$\text{Surface area } A = 2(ab + bc + ca)$$

$$\text{Volume } V = abc$$

$$\text{Diagonal } d = \sqrt{a^2 + b^2 + c^2}$$


In the case of a cube,  $a = b = c$  then;

$$\text{Surface} = 6 a^2$$

$$\text{Volume } V = a^3$$

$$\text{Diagonal } d = a\sqrt{3}$$

### 6.3.2 Right Cylinder




$$\begin{aligned}\text{Surface area of convex surface } A_c &= 2 \pi r h \\ &= 6.283 r h\end{aligned}$$

$$\text{Total surface area } A = 2 \pi r (r + h)$$

$$\text{Volume } V = \pi r^2 h$$


### 6.3.3 Right Cone



$$\begin{aligned}\text{Surface area of convex surface } A_c &= \pi r l \\ (l &= \text{slant height} = \sqrt{r^2 + h^2})\end{aligned}$$

$$\text{Total surface area } A = \pi r (r + l)$$

$$\text{Volume } V = 1/3 \pi r^2 h$$



### 6.3.4 Sphere

$$\text{Surface area } A = 4 \pi r^2 = \pi d^2$$

$$\text{Volume } V = 4/3 \pi r^3 = 1/6 \pi d^3$$

## **Index**

*This page intentionally left blank.*

—A—

Acid anhydrides, 100  
preparation, 100  
reactions, 101

Acid chlorides, 98  
preparation, 98  
reactions, 99

Acid dissociation constant, 32

Acids, 24  
Arrhenius, 24  
Bronsted-Lowery, 25  
Lewis acid, 25

Activated complex, 38

Activation energy, 38

Alcohols, 81  
preparation, 81  
reactions, 83

Aldehydes, 92  
preparation, 92  
reactions, 93 - 94

Algebraic formulas, 173

Alicyclic compounds, 104  
preparation, 104  
reactions, 104

Alkali metals, 42

Alkaline earth metals, 43

Alkanes, 62  
preparation, 62  
reactions, 62

Alkenes, 63  
preparation, 63  
reactions, 64

Alkenylbenzenes, 74  
preparation, 74  
reactions, 74

Alkyl halides, 75  
preparation, 75  
reactions, 76

Alkylbenzenes, 71  
preparation, 71  
reactions, 72

Alkynes, 68  
preparation, 68  
reactions, 68

Amides, 104  
preparation, 104  
reactions, 106

Amines, 107  
preparation, 107  
reactions, 108

Angular momentum quantum number, 3  
Anti-Markovnikov addition, 65  
Arrhenius acid, 24  
Aryl halides, 79  
preparation, 79  
reactions, 80

Atomic mass unit, 2  
Atomic number, 2  
Atomic orbitals, 4  
Atomic radius, 15  
Atomic weights, 17  
Avogadro's law, 22  
Avogadro's number, 3

—B—

Base dissociation constant, 33  
Base SI units, 170  
Bases, 24  
Benzene, 69  
preparation, 69  
reactions, 70

Bidentate ligands, 56

Born exponent, 55

Born-Haber cycle, 54

Boron group, 44

Boyle's law, 21  
Bronsted-Lowery acid, 25

—C—

Carbon group, 45  
Carboxylic acids, 96  
preparation, 96  
reactions, 97

Catalysis, 39

Charles law, 21

Chelation, 56

Chirality, 117

Chromophoric groups, 137

Circle, 176

Classification of organic compounds, 60

Collision theory, 37

Common ion effect, 36

Common molecular ions table, 163

Common nmr solvents, 151

Configurational isomers, 122  
 Conformational isomers, 119  
 Conjugate acid-base pairs, 25  
 Conjugated chromophoric groups, 137  
 Conversion factors, 172  
 Coordinate covalent bond, 18  
 Coordination complexes, 56  
 Coordination number, 49  
 Covalent bond, 18  
 Covalent bonding, 18  
 Crystal lattice, 49  
 Crystal lattice energy, 53  
 Crystal lattice packing, 51  
 Crystal lattice types, 52  
 Cubic close-packed, 51

## —D—

Dalton's law of partial pressures, 22  
 Derived SI units, 170  
 Diastereomers, 117  
 Dienes, 66  
     preparation, 67  
     reactions, 67  
 Dipole-dipole bond, 19

## —E—

Electron, 2  
 Electron spin quantum number, 3  
 Electronegativity, 13  
 Electronic configuration of elements, 8  
 Ellipse, 177  
 Enantiomers, 116  
 Enthalpy, 26  
 Entropy, 26  
 Epoxides, 91  
     preparation, 91  
     reactions, 91  
 Equation of state, 22  
 Equilateral triangle, 176  
 Equilibrium, 28  
 Equilibrium constant, 29  
 Esters, 102  
     preparation, 102  
     reactions, 103  
 Ethers, 89  
     preparation, 89  
     reactions, 90

## —F—

First law of thermodynamics, 25  
 First-order reactions, 36  
 Functional group, 60  
 Fundamental physical constants, 169

## —G—

Gases, 21  
 Gay-Lussac's law, 22  
 Geometric isomers, 119  
 Gibbs free energy, 27  
 Graphs, 174  
 Greek prefixes, 171  
 Group IA, 42  
 Group IIA, 43  
 Group IIIA, 44  
 Group IVA, 45  
 Group VA, 45  
 Group VIA, 46  
 Group VIIA, 46  
 Group VIIIA, 47

## —H—

Halogens, 46  
 Hess' law, 27  
 Heterocyclic compounds, 111  
     preparation of pyridine, quinoline, and isoquinoline, 113  
     preparation of pyrrole, furan, and thiophene, 111  
     reactions of pyridine, quinoline, and isoquinoline, 114  
     reactions of pyrrole, furan, and thiophene, 112  
 Heterogeneous catalysis, 39  
 Heterogeneous equilibrium, 30  
 Hexadentate ligands, 57  
 Hexagonal close-packed, 51  
 Homogeneous catalysis, 39  
 Homogeneous equilibrium, 29  
 Hund's rule, 4  
 Hybrid orbital, 6  
 Hybridization, 6  
 Hydration, 19  
 Hydrogen bond, 20

Hydrolysis, 33  
Hydronium ions, 24  
Hydroxide ion, 24

**—I—**

Ideal gas, 21  
Infrared absorption frequencies, 139  
Infrared media, 138  
Infrared spectroscopy, 138  
Ion-&pole bond, 19  
Ionic bonding, 19  
Ionic character, 19  
Ionic crystal systems, 50  
Ionic equilibrium, 32  
Ionic solids, 49  
Ionization energy, 11  
Isomers, 115

configurational isomers, 122  
conformational isomers, 119  
geometric isomers, 119  
positional isomers, 118

Molarity, 24  
Mole, 2  
Mole fraction, 23  
Molecular mass, 2

**—N—**

Neutron, 2  
Nitrogen group, 45  
NMR chemical shift chart, 153  
NMR chemical shift tables, 155  
NMR spectroscopy, 151  
NMR reference standards, 152  
Noble (inert) gases, 47  
Non SI units, 170

**—O—**

Optically active compounds, 115  
Oxygen group, 46

**—P—**

Parabola, 177  
Parallelogram, 175  
Parallelepiped, 177  
Partial pressures, 22  
Pauli exclusion principle, 3  
Pentadentate ligands, 57  
Periodic table, 10  
Phenols, 84  
preparation, 84  
reactions, 85  
Plane figure formulas, 175  
Plane-polarized light, 116  
Polar covalent bond, 18  
Polymer structures, 123  
Positional isomers, 118  
Principal quantum number, 3  
Proton, 1

**—L—**

Laws of exponents, 173  
Laws of logarithms, 173  
Le Chatelier's principle, 30  
Lewis acid, 25  
Ligand, 56

**—M—**

Madelung constant, 55  
Magnetic quantum number, 3  
Markovnikov addition, 65  
Mass number, 2  
Mass percent, 23  
Mass spectroscopy, 163  
Meso compounds, 117  
Molarity, 24

**—Q—**

Quadratic equation, 174  
Quadridentate ligands, 57  
Quantum numbers, 3

## —R—

Racemic mixture, 117  
 Rectangle, 175  
 Right cone, 178  
 Right cylinder, 178

## —S—

Second-order reactions, 36  
 SI prefixes, 171  
 Solid figure formulas, 177  
 Solubility product, 34  
 Solutions, 23  
 Solvation, 19  
 Sphere, 178  
 Standard state, 27  
 Stereoisomers, 115

## —W—

Woodward's rules, 136

## —Z—

Zero-order reactions, 36

## —T—

Thermodynamics, 25  
 Transition metal, 48  
 Transition state, 38  
 Transition state theory, 38  
 Trapezoid, 176  
 Tridentate ligands, 57

## —U—

Uv spectroscopy, 135  
 Unidentate ligands, 56  
 Unit cells, 50  
     cubic, 50  
     hexagonal, 50  
     monoclinic, 50  
     orthorhombic, 50  
     rhombohedral, 51  
     tetragonal, 50  
     triclinic, 51  
 Uv-vis solvents, 135

## —V—

van der Waals, 20




**KLUWER ACADEMIC /  
PLENUM PUBLISHERS**

233 Spring Street, New York, New York 10013-1578

PRINTED IN U.S.A.

ISBN 0-306-46546-9

90000



9 780306 465468