Difficult transition for sugar maple in Boreal forest under climate change? Impact of alternative stable states on Sugar maple migration.

Research proposal

Master in Wildlife management

Ву

Steve Vissault

For

Richard Cloutier, Pr. Director of the program committee

Dominique Arsenault, Pr. President of the jury

Matt Talluto, PhD Research Co-director

Dominique Gravel, Pr. Research Director

Université du Québec à Rimouski January 17, 2014

1 Introduction.

Context. The boreal region is warming twice as fast as the global average and will inevitably alter species composition in boreal forest [10, 5]. Sugar maple is one of those species expected to migrate northward towards it's northern limits [7, 1]. Predict shifts in the repartition of sugar maple under climate change is an important challenge whereas this species is highly coveted by wood and maple syrup producers, two main economic sectors in Quebec. Indeed, Sugar maple is a widespread and abundant tree in north-eastern North America and one of the most representative species of northern temperate forests [3, 8, 6]. This northward migration will result in increasing the surface of the ecotone between the boreal and temperate forest of Quebec. Many ecotone studies and modeling efforts on transition between forest to non-forest ecosystems [10, 9, 4] but little attention has been given to evaluate the transitionnal dynamics of forest-forest ecotone [2, 3]. This project aim to develop a transitionnal model between the boreal and temperate nordic forest in order to investigate the migration rate of sugar maple under climate change.

Theorical framework.

2 Objectives.

This project aims to determine whether alternative stable states are present in the temperate-boreal forest ecotone and if so, look at the impact of plant-soil and disturbances feedback on the alternatives stables states. To assess this main objective, we will (O1) generate a transitionnal model between the temperate and the boreal forest; (O2) study the equilibrium states based on the model; (O3) investigate the spatial structure of the transitionnal zone; and finally (O4) run simulations based on different climate change scenarios.

3 Methods.

References

- [1] D. GOLDBLUM AND L. RIGG, Tree growth response to climate change at the deciduous boreal forest ecotone, Ontario, Canada, Canadian Journal of Forest Research, 2718 (2005), pp. 2709–2718.
- [2] D. GOLDBLUM AND L. S. RIGG, *The Deciduous Forest Boreal Forest Ecotone*, Geography Compass, 4 (2010), pp. 701–717.
- [3] N. Graignic, F. Tremblay, and Y. Bergeron, Geographical variation in reproductive capacity of sugar maple (Acer saccharum Marshall) northern peripheral populations, Journal of Biogeography, (2013), pp. n/a-n/a.
- [4] M. HIROTA, M. HOLMGREN, E. V. NES, AND M. SCHEFFER, Global Resilience of Tropical Forest and Savanna to Critical Transitions, Science, (2011), pp. 232–235.
- [5] L. Hughes, Biological consequences of global warming: is the signal already apparent?, Trends in Ecology & Evolution, 15 (2000), pp. 56–61.
- [6] M. Kellman, Sugar maple (Acer saccharum Marsh.) establishment in boreal forest: results of a transplantation experiment, Journal of Biogeography, 31 (2004), pp. 1515–1522.
- [7] D. W. Mc Kenney, J. H. Pedlar, K. Lawrence, K. Campbell, and M. F. Hutchinson, Potential Impacts of Climate Change on the Distribution of North American Trees, BioScience, 57 (2007), p. 939.
- [8] Y. MESSAOUD, Y. BERGERON, AND A. LEDUC, Ecological factors explaining the location of the boundary between the mixedwood and coniferous bioclimatic zones in the boreal biome of eastern North America, Global Ecology and Biogeography, 16 (2007), pp. 90–102.
- [9] M. Scheffer, S. Carpenter, J. A. Foley, C. Folke, and B. Walker, *Catastrophic shifts in ecosystems.*, Nature, 413 (2001), pp. 591–6.
- [10] M. SCHEFFER, M. HIROTA, M. HOLMGREN, E. H. VAN NES, AND F. S. CHAPIN, Thresholds for boreal biome transitions., Proceedings of the National Academy of Sciences of the United States of America, 109 (2012), pp. 21384–9.